1 **Determinants of target prioritization and regulatory hierarchy for the bacterial**
2 **small RNA SgrS**
3

4 **Maksym Bobrovskyy[1,2] Jane Frandsen[3,4], Jichuan Zhang[5], Anustup Poddar[5],**
5 **Muhammad S. Azam[1], Tina M. Henkin[3], Taekjip Ha[5,6] and Carin K. Vanderpool[1]***
6

[1]Department of Microbiology, University of Illinois at Urbana-Champaign, 601 S.
Goodwin Ave., Urbana, Illinois 61801
[2]Present address: Department of Microbiology, The University of Chicago, 920 E. 58[th]
St., Chicago, Illinois 60637
[3]Department of Microbiology and Center for RNA Biology, The Ohio State University,
Columbus, Ohio 43210
[4]Biochemistry Program, The Ohio State University
[5]Department of Biophysics and Biophysical Chemistry, Johns Hopkins University,
Baltimore, Maryland, USA 21205
[6]Howard Hughes Medical Institute, Baltimore, Maryland, USA 21205
17

*Corresponding author
Carin K. Vanderpool, Ph.D.
Department of Microbiology
University of Illinois at Urbana-Champaign
C213 CLSL, MC-110
601 S. Goodwin Ave.
Urbana, IL 61801
(t) 217-333-7033
(f) 217-244-6697
cvanderp@life.illinois.edu
28

29 Running title: Target prioritization by small RNAs

30 Abstract word count: 241

31 Text word count: 6,824

ABSTRACT

The mechanisms by which small RNA (sRNA) regulators select and prioritize target mRNAs remain poorly understood, but serve to promote efficient responses to environmental cues and stresses. We sought to uncover mechanisms establishing regulatory hierarchy for a model sRNA, SgrS, found in enteric bacteria and produced under conditions of metabolic stress when sugar transport and metabolism are unbalanced. SgrS post-transcriptionally controls a nine-gene regulon to restore growth and homeostasis under stress conditions. An in vivo reporter system was used to quantify SgrS-dependent regulation of target genes and established that SgrS exhibits a clear preference for certain targets, and regulates those targets efficiently even at low SgrS levels. Higher SgrS concentrations are required to regulate other targets. The position of particular targets in the regulatory hierarchy is not well-correlated with the predicted thermodynamic stability of SgrS-mRNA interactions or the SgrS-mRNA binding affinity as measured in vitro. Detailed analyses of SgrS interaction with *asd* mRNA demonstrate that SgrS binds cooperatively to two sites and remodels *asd* mRNA secondary structure. SgrS binding at both sites increases the efficiency of *asd* mRNA regulation compared to mutants that have only a single SgrS binding site. Our results suggest that sRNA selection of target mRNAs and regulatory hierarchy are influenced by several molecular features. The sRNA-mRNA interaction, including the number and position of sRNA binding sites on the mRNA and cofactors like the RNA chaperone Hfq seem to tune the efficiency of regulation of specific mRNA targets.

IMPORTANCE

To survive, bacteria must respond rapidly to stress and simultaneously maintain metabolic homeostasis. The small RNA (sRNA) SgrS mediates the response to stress arising from imbalanced sugar transport and metabolism. To coordinate the stress response, SgrS regulates genes involved in sugar uptake and metabolism. Intrinsic properties of sRNAs such as SgrS allow them to regulate extensive networks of genes. To date, sRNA regulation of targets has largely been studied in the context of "one sRNA-one target", and little is known about coordination of multi-gene regulons and sRNA regulatory network structure. Here, we explore the molecular basis for regulatory hierarchy in sRNA regulons. Our results reveal a complex interplay of factors that influence the outcome of sRNA regulation. The number and location of sRNA binding sites on mRNA targets and the participation of an RNA chaperone dictate prioritized regulation of targets to promote an efficient response to stress.

INTRODUCTION

Bacteria live in diverse niches, often encountering rapidly changing and stressful environments. Bacterial stress responses can mitigate the negative effects of stress on cell structure and function. Usually stress responses are coordinated by molecules—either RNAs or proteins, that alter expression of a regulon comprised of multiple genes. Coordinated control of the regulon prepares the cell to survive or adapt to the stress (1, 2). Proteins control expression of target regulons by binding to DNA sequences and modulating the frequency of transcription initiation, but RNAs often modulate gene expression post-transcriptionally. A prevalent type of RNA regulator in bacteria is referred to simply as small RNA (sRNA). These sRNAs are often produced in response to a particular stress, and regulate target mRNAs through base pairing interactions that modify mRNA translation or stability (3, 4). In diverse bacteria, hundreds of sRNAs have been identified (5-7). While the majority of sRNAs have not been characterized, many studies suggest that sRNA regulatory networks are as extensive and complex as those controlled by proteins (8, 9).

A large body of work has illuminated base pairing-dependent molecular mechanisms of post-transcriptional regulation by sRNAs (10, 11). The sRNA SgrS (sugar-phosphate stress sRNA) has been an important model for discovery of both negative and positive mechanisms of target mRNA regulation. SgrS is induced in response to metabolic stress associated with disruption of glycolytic flux and intracellular accumulation of sugar phosphates (also referred to as glucose-phosphate stress) (12, 13). SgrS regulates at least 9 genes and promotes recovery from glucose-phosphate stress. SgrS-dependent repression of mRNAs encoding sugar transporters

(*ptsG, manXYZ*) reduces uptake of sugars to prevent further sugar-phosphate

accumulation (Fig. 1) (12, 14, 15). Activation of a sugar phosphatase (*yigL*) mRNA

promotes dephosphorylation and efflux of accumulated sugars (16), and repression of

other mRNAs is hypothesized to reroute metabolism to promote recovery from stress

(Fig. 1) (17). Each target of SgrS is regulated by a distinct molecular mechanism. How

different mechanisms of regulation yield effects of variable magnitude with respect to

mRNA stability and translation is an open question.

Temporally-ordered and hierarchical patterns of gene regulation carried out by

protein transcription factors have been characterized in many systems (18-21). These

regulatory patterns allow cells to efficiently respond to environmental signals by

prioritizing induction or repression of products needed to respond to those signals.

Protein regulators establish a hierarchy of regulation based on their affinities for binding

sites in the operator regions of different target genes. As the concentration of active

regulator increases, genes are sequentially regulated based on binding site affinity (22).

There is growing evidence that sRNAs also regulate their target genes hierarchically

(23, 24). However, the mechanisms involved in establishing and maintaining prioritized

regulation of sRNA targets are not known.

We hypothesize that the temporal progression of target regulation by SgrS is

specifically optimized to promote efficient recovery from glucose-phosphate stress (Fig.

1). To test this hypothesis, first defined the efficiency of SgrS regulation of each target

and found that SgrS indeed prioritizes regulation of some targets over others. We

examined the factors that determine regulatory efficiency, including the the arrangement

and strength of SgrS target binding sites and the roles of other factors like RNase E and

Hfq. Detailed characterization of a specific SgrS-mRNA target interaction revealed

cooperative binding of SgrS to two binding sites and a requirement for both binding sites

for maximal SgrS-dependent regulation. Collectively, our results upheld the hypothesis

that sRNAs regulate expression of genes in their target regulons hierarchically.

Features of each sRNA-mRNA pair and molecular mechanisms of regulation precisely

determine the regulatory priority for each target.

RESULTS

SgrS differentially regulates targets at the level of translation. Previous studies

suggested the possibility of a hierarchy of regulatory effects carried out by the small

RNA SgrS, which controls translation of a diverse set of mRNA targets (11, 12, 15, 16,

25). To study this, we used a two-plasmid system to control expression of SgrS and

target translational fusions (Fig. 2A). All target transcript fragments fused to *gfp* contain

experimentally confirmed SgrS binding sites. Regulation of target translation by SgrS

was measured as described previously (24).

To quantify translational regulation by SgrS and facilitate comparisons of

regulatory efficiency among targets, we analyzed the data as described by Levine, et al.

(24). Activity of reporter fusions was measured by monitoring GFP fluorescence over

time. By plotting the GFP fluorescence (RFU) as a function of growth (OD_{600}) for target-

gfp fusions in the <u>absence</u> of SgrS, we defined "basal activity" at different inducer

concentrations (example in Fig. S1A). This method of quantifying translational fusion

activity accounts for the fact that fluorescence levels are not directly proportional to

inducer concentrations ((24) and Fig. S1A). While the absolute values for basal activity

differ among different target fusions, all fusions responded to induction in a dose-

dependent manner (Fig. S2A). Similar plots (RFU/OD$_{600}$) were generated for each

fusion induced in the presence of SgrS (examples Fig. S1B-F). We define "regulated

activity" as the slope of the curve (RFU/OD$_{600}$) under conditions where both the fusion

and SgrS are induced (example in Fig. S1B). As levels of SgrS increase, clear patterns

of repression or induction are seen for all target fusions (Figs S1B-F and S2B-F).

To define the efficiency of regulation of each target we plotted regulated activity

as a function of basal activity for *ptsG*, *manX, asdI, purR,* and *yigL*. When there is no

SgrS-mediated regulation, a line with a slope of 1 is seen for all targets (Fig. 2B-F).

Slopes less than 1 indicate that the fusion is repressed by SgrS. This is true for *ptsG*,

manX, *asdI* and *purR* reporter fusions (Figure 2B-E). Slopes greater than 1 are

indicative of activation by SgrS, which is true only for *yigL* (Fig. 2F). Importantly, the

magnitude of regulation was responsive to SgrS levels. As concentrations of SgrS

inducer (aTc) increased, slopes of lines for repressed targets were correspondingly

reduced (Fig. 2B-E). This was not the case for *yigL,* the only positively regulated target

of SgrS (Fig. 2F). The magnitude of activation did not increase beyond a maximal level

obtained at 20 ng/mL of inducer. While the basis of this difference is unclear, it likely

reflects the inherently different molecular mechanisms of regulation: mRNA stabilization

for *yigL* and translational repression for all other targets.

We then compared regulatory efficiency among different targets at different

levels of SgrS induction. At the two lowest levels of SgrS induction (10ng/mL and 20

ng/mL aTc), only *ptsG* and *yigL* showed substantial repression and activation,

respectively (Fig. 3A, B). In contrast, *manX*, *asdI* and *purR* fusions yielded curves

whose slopes remained at ~1, indicating no regulation at these lower levels of SgrS.

Our interpretation of these results is that *ptsG* and *yigL* are the high-priority or

"strongest" targets of SgrS, since they are regulated preferentially when SgrS levels are

low. With increasing SgrS levels (20-50 ng/ml aTc), regulation of "weaker" targets

manX, asdI and *purR* became apparent (Fig. 3C, D, E). As SgrS levels increased, *ptsG*

repression became more efficient up to a maximal repression at 40 ng/mL of aTc, and it

remained the most strongly repressed target at all levels of SgrS. Collectively, these

data suggest that SgrS targets are preferentially regulated in the following order: 1/2)

ptsG and *yigL*, 3) *manX*, 4) *asdI*, and 5) *purR* (Fig. 3A-E). We hypothesize that the

position of each target within the regulatory hierarchy is determined by characteristics of

SgrS-target mRNA interactions and the mechanism of SgrS-dependent regulation.

Differences in in vitro binding affinity are not correlated with regulation efficiency.

One of the initial steps in sRNA-mediated regulation is formation of base-pairing

interactions with the target mRNA. Binding of the sRNA with its target mRNA is

dependent on sequence complementarity and RNA secondary structure. We examined

the characteristics of SgrS-target mRNA binding in vitro to determine whether the

strength of binding is correlated with the target hierarchy observed at the level of

translation.

Electrophoretic mobility shift assays (EMSAs) were performed to measuring

binding of SgrS to its target mRNAs *ptsG*, *manX*, *purR*, *yigL* and *asd*. Binding of SgrS to

ptsG had a K_D of 0.11 ± 0.01 µM (Fig. 4A, B), which was lower than K_Ds for SgrS

binding to most of the other targets (Fig. 4A-E). SgrS-*manX* mRNA binding had a K_D of

19.7 ± 2.78 µM (Fig. 4A, C) which is weaker than the interaction with *ptsG* (Fig. 4B), but

stronger compared to *purR* and *yigL* (Fig. 4A). Three different fragments of *asd* mRNA

were tested, because previous work demonstrated that SgrS pairs at two distinct sites

on *asd* mRNA (17). The first site, *asdI*, is adjacent to the ribosome binding site and is

sufficient for modest SgrS-dependent translational repression. The second site, *asdII,* is

in the coding sequence of *asd,* 60-nt downstream of the start codon. When both sites

are present, *i.e.,* on *asdI-II,* stronger SgrS-dependent translational repression is

observed (17). Surprisingly, while *asdI* (containing only the upstream SgrS binding site)

regulation is less efficient compared to *manX* (Fig. 3A-E), *in vitro* it binds SgrS more

strongly with a K_D of 0.15 ± 0.04 µM (Fig. 4A, D), which is comparable to SgrS-*ptsG*

binding (Fig. 4A, B). SgrS interaction with *asdII* was very weak (Fig. 4A). We could not

determine K_D values for SgrS interaction with *asdII*, *purR* and *yigL*, due to limitations in

obtaining high enough concentrations of RNA, but it is apparent that SgrS binding to

these targets is much weaker compared to *ptsG*, *manX* and *asdI* (Fig. 4A).

Results of EMSAs with SgrS and *asdI-II* (containing both SgrS binding sites)

revealed apparent binding cooperativity. SgrS binding to *asdI-II* has a K_D of 0.07 ± 0.01

µM (Fig. 4E, F), even slightly lower than that of SgrS-*ptsG* mRNA binding. Moreover, we

observed two shifted species that correspond to one or two SgrS sRNAs pairing with a

single *asdI-II* transcript (Fig. 4E).

Structural analyses of SgrS-*asd* mRNA interactions. Our data thus far indicate that

SgrS regulates mRNA targets in a hierarchical fashion (Figs. 2, 3). SgrS-mRNA binding

affinities alone do not explain the target hierarchy, as SgrS-*ptsG* mRNA and SgrS-*asd*

mRNA interactions have very similar K_Ds, but *ptsG* is much more efficiently regulated

than *asd* at all concentrations of SgrS (Fig. 3). To further understand the features that

influence the efficiency of target regulation, we performed more detailed analyses of

SgrS-*asd* mRNA interactions.

Previous work demonstrated that SgrS binding site I encompasses nt +31 to +49

and site II from nt +110 to +127 ((17), Fig. 5A) We used IntaRNA (26, 27) to predict the

free energy (ΔG) for SgrS interactions with *asd* mRNA segments containing both sites,

or each site individually (Fig. 5B). IntaRNA accounts for the energy necessary to open

double-stranded regions of RNA secondary structure, to make them accessible for

pairing. We first analyzed SgrS interactions with *asdI-II* mRNA (encompassing nt +1 to

+240), which we denote as "structured" (Fig. 5B). Interaction of SgrS with *asd* site I has

a predicted ΔG of -10.5 kcal/mol, while SgrS pairing with site II has a ΔG of -1.1

kcal/mol (Fig. 5B, structured). The ΔG for interactions between SgrS and the isolated

binding sites, are -18 kcal/mol for site I and -7.4 kcal/mol for site II (Fig. 5B, isolated).

These predictions suggest that SgrS interaction with site II is less favorable, particularly

in the context of the longer structured *asd* mRNA.

We investigated the structure of *asdI-II* with selective 2'-hydroxyl acylation

analyzed by primer extension (SHAPE), where flexible nts are modified by N-

methylisotoic anhydride (NMIA), while nts constrained in helices are not reactive. In the

absence of SgrS, the sequence encompassing the *asd* ribosome binding site (+44 to

+50) is located within a structured loop (+36 to +69) on top of a short stem (+31 to +35

pairing with +70 to +74) (Fig. 5C, Fig. S3). The nts in site I (+31 to +49, Fig. 5A) are

located on the 5' side of the stem-loop structure. Most of the nts in this structure are

reactive, which is indicative of a flexible conformation that is accessible for ribosome

binding or base pairing with the seed sequence of SgrS (Fig. 5C). The seed interaction

of SgrS likely promotes opening of the structure. Downstream of the site I stem-loop

structure is a highly structured second stem (+83 to +155) that contains site II in the

apical region (+110 to +129) (Fig. 5C, Fig. S3). Site II is sequestered in a helix and

would not be accessible to base pair with SgrS (Fig. 5C). In light of binding cooperativity

observed in Fig. 4E, we hypothesize that SgrS pairing with site I induces rearrangement

of *asd* mRNA secondary structure to facilitate interaction with site II.

We next used SHAPE to probe changes to the *asdI-II* structure in the presence

of SgrS. The reactivity of site I nt +31 to +49 decreased as the concentration of SgrS

increased (Fig. 5D), with the exception of nt +41 which is not predicted to base pair with

SgrS (Fig. 5A). The SHAPE reactivity plateaued between 5 and 10-fold excess SgrS

(Fig. S3E,F), which suggests that binding to site I was saturated. This is consistent with

a strong base-paring interaction between SgrS and site I. In contrast, the reactivity of

the site II nts +110 to +129 decreased more slowly and to a lesser extent (Fig. 5D),

consistent with a weaker and cooperative interaction. Fewer site II nts showed changes

in SHAPE reactivity upon addition of SgrS; this is likely due to the highly structured

nature of site II in the absence of SgrS.

The reactivity of nts outside of the SgrS binding sites also changed in the

presence of SgrS (Fig. 5E). When a mutant SgrS that is not predicted to bind to *asdI-II*

was used, minimal changes in SHAPE reactivity were observed, which suggests that

the changes in the presence of wild-type SgrS are due to the interactions between SgrS

and *asdI-II* mRNA and not due to the presence of additional RNA in the system (Fig.

5E). This indicates that SgrS binding changes the overall structure of the *asd* RNA. A

secondary structure predicted using the SHAPE data suggests that these changes are

limited to opening the SgrS binding sites and extending the site II stem (Fig. 5C). It is

worth noting an important caveat to these analyses. The structure prediction algorithms

were not designed to account for intermolecular interactions, so this analysis may not

be able to capture the in vivo relevant structure of *asd* mRNA in complex with SgrS.

Nonetheless, SHAPE data are consistent with other analyses in demonstrating binding

of SgrS to *asd* mRNA, prominently at site I and to a lesser extent at site II.

Optimal repression by SgrS involves both pairing sites within *asd* mRNA. To

further investigate the role of the two SgrS pairing sites on *asd*, we performed stochastic

optical reconstruction microscopy (STORM) coupled with single-molecule *in situ*

hybridization (smFISH) to monitor SgrS regulation of *asd-lacZ* variants *asdI, asdII,* and

asdI-II (Fig. 6A) at single molecule resolution. In these experiments bacteria were grown

in the presence of L-arabinose to induce expression of chromosomal *asd-lacZ* variants.

Glucose-phosphate stress was induced for 10 min by the addition of 1% α-methyl D-

glucopyranoside (αMG). 3D super-resolution images show *asd-lacZ* mRNAs (Fig. 6B-D,

green) and SgrS (Fig. 6B-D, red), as projected on 2D planes, with cells outlined.

Numbers of *asd-lacZ* mRNAs and SgrS sRNAs were counted and represented as "copy

number per cell" in histograms, with average copy number per cell indicated above the

histogram (Fig. 6B-D). SgrS induction reduced the copy number of *asdI-lacZ* mRNA by

3-fold (Fig. 6B, green) when SgrS was induced to high levels after 10 min treatment with

αMG (Fig. 6B, red and S4A, B). On the contrary, the copy number of *asdII-lacZ* mRNAs

(Fig. 6C, green) was not strongly affected in the presence of high SgrS levels after αMG

treatment (Fig. 6C, red and Fig. S4C, D). Copy numbers of *asdI-II-lacZ* mRNA (Fig. 6D,

green) were reduced by ~8-fold after 10 min of αMG induction (Fig. 6D, red, Fig.

S6E,F). These data demonstrate that both binding sites on *asd* mRNA are important for

efficient SgrS-dependent regulation of mRNA stability.

We next examined the roles of the two SgrS binding sites in the efficiency of

translational regulation. SgrS regulation of an *asdI-II* translational fusion was compared

to regulation of an *asdI* fusion (Fig. 7A). By plotting regulated activity as a function of

basal activity as described above, we determined that SgrS repression of *asdI-II* was

more efficient than repression of *asdI* across the entire range of SgrS expression levels

(Fig. 7B), a result in line with our previous study (17). Comparison to other targets

indicated that *asdI-II* is regulated more efficiently than *manX*, *asdI* and *purR*, at all

concentrations of SgrS (Fig. 7C).

We then compared SgrS regulation of *asdI* and *asdI-II* in the *rne701* mutant

strain deficient in degradosome assembly (28). We determined basal activity (Fig. S5A)

and regulated activity (Fig. S5B-F) of *asdI* and *asdI-II* translational *gfp* fusions at

different levels of SgrS induction. Reminiscent of our data in the wild-type strain (Fig.

7A), SgrS regulated *asdI-II* more efficiently compared to *asdI* in the *rne701* mutant (Fig.

7D. Moreover, when compared to SgrS regulation of other targets, *asdI-II* was

repressed most efficiently (Fig. 7E) in the *rne701* mutant. Taken together the data

indicate that the second binding site on *asd* mRNA enhances the stringency of SgrS-

mediated regulation. Moreover, addition of the second binding site on *asd* changes its

regulatory priority relative to other targets in the SgrS regulon.

Transcriptional regulation of *asd* by SgrS. We observed that the *asdI-II* transcript is

more abundant than *asdI* (Fig. 6B, D). Since the constructs used in that experiment

were expressed from a heterologous promoter, we postulate that increased levels of

asdI-II mRNA compared to *asdI* mRNA must be due to increased mRNA stability or

transcription elongation of the *asdI-II* construct compared to *asdI.* We constructed *asdI*

and *asdI-II* transcriptional fusions to *lacZ* expressed from an inducible promoter (Fig.

8A) to test whether SgrS can regulate *asd* transcripts independent of translational

regulation. Consistent with observations from smFISH, the *asdI-II-lacZ* transcriptional

fusion had substantially higher activity compared to *asdI-lacZ* (Fig. 8B). While SgrS

regulated both reporter fusions, *asdI-II* repression was more efficient (3.3-fold

repression) than *asdI* (2.1-fold repression) (Fig. 8B). SgrS still regulated both fusions in

the *rne701* mutant strain (Fig. 8B). Importantly, SgrS-dependent degradation of other

SgrS targets *ptsG* mRNA (29), and *manXYZ* mRNA (15, 25) was abolished in

degradosome mutants. Together with our previous work, these observations suggest

that SgrS regulates *asd* by two independent mechanisms: translational repression by

pairing at site I (directly occluding the ribosome binding site) and reducing mRNA levels

by promoting mRNA turnover and/or inhibiting transcription elongation.

DISCUSSION

In this study, we set out to define the hierarchy of regulation by a model bacterial

sRNA. SgrS is a good model for this study because it has a modestly-sized regulon,

and the mechanisms of regulation of several targets have been characterized in detail

(16, 17, 25, 30). Our results demonstrate a clear pattern of prioritized regulation of

mRNA targets (Fig. 2B-F, Fig. 3A-E). Two targets in particular, *ptsG* and *yigL,* were

"high-priority" targets that were efficiently regulated even at low levels of SgrS

production. Other targets, *manX, purR,* and *asd,* were less impacted by SgrS and were

only regulated when SgrS was produced at higher levels.

We investigated features of sRNA-mRNA interactions that could impact the

overall efficiency of SgrS-mediated regulation. In vitro SgrS-mRNA interactions as

measured by EMSA defined K_Ds that were not well-correlated with in vivo regulatory

efficiency (Fig. 4A-F, Fig. 3A-E). Two targets stood out in the comparison of in vivo

regulation and in vitro SgrS-mRNA interactions. The *yigL* mRNA interaction with SgrS

was barely detectable in vitro (Fig. 4A), but in vivo, *yigL* translation was maximally

activated at low SgrS levels (Fig. 2F). Conversely, the translation of *asdI* was modestly

regulated by SgrS in vivo (Fig. 2E), but the in vitro SgrS-*asdI* interaction was

comparable to that of SgrS-*ptsG,* the strongest in vivo regulatory effect. These apparent

contradictions between in vitro interactions and in vivo regulatory efficiency led us to

further explore SgrS regulation of *asd.*

Previous work demonstrated that SgrS has two binding sites on *asd* mRNA: site I

overlaps that *asd* ribosome binding site and site II is ~60 nt downstream in the *asd*

coding sequence ((17) and Fig. 5A). EMSAs demonstrated SgrS pairing at site I alone,

but pairing at site II alone was undetectable. Binding of SgrS to an *asd* mRNA

containing both sites I and II was cooperative (Fig. 4E,F). Structural analyses of *asd*

mRNA in the absence and presence of SgrS demonstrated that SgrS indeed pairs

preferentially at site I over site II and induces substantial structural rearrangement in the

mRNA (Fig. 5C-E, Fig. S3). Quantification of SgrS-dependent degradation of *asd* mRNA

showed that site I is important, but sites I and II together promote the most efficient

regulation (Fig. 6B-D, Fig. S4). Similar to binding and regulation of mRNA degradation,

SgrS interactions at both sites I and II on *asd* mRNA improve the efficiency of

translational regulation (Fig. 7B,C, Fig. S2). These results suggest that the number and

position of sRNA binding sites on mRNA targets control regulation in vivo in ways that

could not be predicted based on in vitro characterization of sRNA-mRNA binding.

In many cases, sRNA-mediated regulation of translation is thought to indirectly

affect mRNA stability by making untranslated mRNA more susceptible to degradation by

RNase E. There are also examples of sRNA regulation, including SgrS regulation of

yigL (16), where modulation of mRNA stability is translation-independent. Truncation of

RNase E (encoded by *rne*), removing the C-terminal scaffold for degradosome

assembly, often prevents sRNA-mediated degradation of mRNA targets (15, 31, 32). If

translational regulation is the primary function of an sRNA on a given mRNA target, the

regulation should be preserved in *rne* mutant backgrounds. For SgrS targets, the

regulatory hierarchy is mostly preserved in an *rne701* degradosome mutant background

(Fig. 7, compare C and E), suggesting that for most targets, regulation of RNA stability

is not the primary mechanism of control by SgrS. Interestingly, the high-priority target

ptsG was a notable exception. In the wild-type background, *ptsG* is the most efficiently-

regulated target (Fig. 7C), whereas in the *rne701* host, it is weakly regulated. This

defect could be overcome by increasing SgrS expression levels (Fig. S6B). This result

suggests that RNase E-dependent degradation of *ptsG* mRNA is more important for its

efficient regulation by SgrS compared to other targets, where efficient regulation does

not depend on subsequent target degradation. This is consistent with the fact that *ptsG*

mRNA levels decrease at least 10-fold whereas other targets exhibit a modest 2-fold

decrease in mRNA levels upon SgrS expression (17). Our recent study quantifying

SgrS-dependent mRNA degradation at single molecule resolution indicated that *ptsG*

mRNA exhibits faster degradation kinetics than *manXYZ* mRNA (31), which could

enhance the efficiency of regulation in a wild-type but not *rne701* mutant background

where translational regulation and mRNA degradation are uncoupled.

One of our ultimate goals is to define at a molecular level the mechanisms by

which sRNAs select and prioritize regulation of their targets. The current study

implicates features of sRNA-mRNA interactions such as number and strength of sRNA

binding sites on each mRNA target and accessory factors such as RNase E in dictating

regulatory hierarchy. Another factor that is likely to play an important role in setting

regulatory priority is the RNA chaperone Hfq. EMSAs demonstrated Hfq binding to

ptsG, *manX*, *purR*, *yigL*, *asdI*, *asdII* and *asdI-II* mRNAs (Fig. S7A) with similar K_D values

for all targets (Fig. S7B). Previous work has shown that sRNAs compete for binding to

Hfq, and this competition affects their regulatory ability (33, 34). Different sRNAs can

bind to distinct sites on Hfq and this impacts their regulation of particular targets (34,

35). Additional work will be required to determine what role Hfq plays in establishing the

hierarchy of regulatory effects in sRNA regulons.

Most sRNA-mRNA interactions are conceived of as single binding site

interactions, but we have already identified two SgrS targets that deviate from this

model and have shown that additional binding sites can play important roles in

regulation and change regulation efficiency (17, 25). We have not yet discovered the

specific mechanism of regulation of *asd* mRNA by SgrS, but have shown definitively

that both binding sites are required for strong regulation. SgrS-dependent control of

both transcriptional and translational *asd* reporter fusions is not impacted in RNase E

degradosome deficient strains (Fig. 7B,D, Fig. 8B), suggesting that the regulation is not

strictly dependent on translation or mRNA turnover. Future work will test the hypothesis

that SgrS acts on *asd* mRNA at the level of transcription elongation, perhaps by an

attenuation mechanism.

In *Vibrio*, quorum sensing-regulated Qrr sRNAs regulate multiple targets by

distinct mechanisms and differences in those mechanisms influence the dynamics and

strength of regulation (23). Strong and rapid regulation is achieved by sRNAs acting

catalytically where the sRNA pairs with and promotes mRNA degradation but is then

recycled for use on another mRNA target. A sequestration mechanism, where formation

of the sRNA-mRNA complex is the terminal outcome of regulation, results in slower and

weaker sRNA-dependent regulation of the target mRNA. For Qrr sRNAs, these

regulatory mechanisms seem to depend on which region of the sRNA is pairing with a

given target and whether the sRNA-mRNA interaction is strong or weak (23). While

some of the same features of SgrS-mRNA interactions may be relevant in determining

regulatory efficiency, we note that the SgrS seed sequence responsible for pairing with

all mRNA targets characterized thus far is encompassed by a short (~20 nt) mostly

single-stranded region of SgrS (12, 15-17). Moreover, we did not see a good correlation

between strong versus weak binding in vitro and in vivo regulatory efficiency. It may be

true that the "rules" governing regulatory efficiency and specific outcomes are different

for each individual sRNA. Work on more model sRNAs will be needed to illuminate

broad general principles.

Beyond defining interesting molecular features of sRNA-mRNA interactions,

defining regulatory hierarchy for sRNA regulons is important for understanding bacterial

physiology. The vast majority of sRNA regulons remain undefined, and thus sRNA

functions unknown. For novel sRNAs, distinguishing high-priority from weaker targets

may provide crucial clues to the predominant role of the sRNA in cell physiology. For

SgrS, the regulatory hierarchy we have defined here is perfectly consistent with growth

studies demonstrating the primary importance of SgrS regulation of sugar transport and

efflux under glucose-phosphate stress conditions (36). The hierarchy of regulation by

sRNAs likely evolved to promote rapid and efficient responses to environmental signals

that would provide cells with a competitive growth advantage in their specific niche. It

will be critical to develop tools to more rapidly characterize sRNA regulatory hierarchy to

better understand functions of the hundreds of uncharacterized sRNAs in diverse

bacteria.

MATERIALS AND METHODS

Strain and plasmid construction. List of strains and plasmids used in this study are

listed in Table S1. All strains used in this study are derivatives of *E. coli* K-12 strain

MG1655. Oligonucleotide primers and 5' biotinylated probes used in this study are listed

in Table S2 and were acquired from Integrated DNA Technologies. Chromosomal

alleles were moved between strains by P1 *vir* transduction (37) and inserted using λ

Red recombination (38, 39).

Translational reporter fusion alleles P_{BAD}-*asdI-II-lacZ* (MBP151F/MBP193R

primers), P_{BAD}-*asdI-lacZ* (MBP151F/MBP151R primers) and P_{BAD}-*asdII-lacZ*

(MBP193F/MBP193R primers) were constructed by PCR amplifying desired fragments

using primers containing homologies to P_{BAD} and *lacZ.* Similarly, transcriptional fusions

P_{BAD}-*asdI-II-lacZ* (MBP151F/MBP206R3 primers) and P_{BAD}-*asdI-lacZ*

(MBP151F/MBP206R1 primers) were generated by PCR amplification using forward

primer with homology to P_{BAD} and reverse primers containing *lacZ* RBS and *lacZ*

homology. PCR products were then recombined into PM1205 strain using λ Red

homologous recombination.

Plasmid harboring SgrS under the control of P*tet0-1* promoter was constructed by

PCR amplifying *sgrS* from *E. coli* MG1655 chromosomal DNA using oligos containing

NdeI and BamHI restriction sites. Resulting PCR product and vector pZA31R (24) were

digested by NdeI and BamHI (New England Biolabs) restriction endonucleases.

Digestion products were ligated using DNA Ligase (New England Biolabs) to produce

pZAMB1 plasmid containing P$_{tet0-1}$-*sgrS* allele.

Plasmid pZEMB8 containing P$_{lac0-1}$-*ptsG-gfp* was constructed by PCR amplifying

ptsG from MG1655 chromosomal DNA using oligos containing KpnI and EcoRI

restriction sites. Resulting PCR products and vector pZE12S (24) were digested by KpnI

and EcoRI restriction endonucleases. Digestion products were ligated using DNA

Ligase to produce pZEMB2. Superfolder *gfp* (from now on just *gfp*) was amplified from

pXG10-SF (40) using oligos containing KpnI and XbaI restriction sites. pZEMB2 and the

resulting PCR product were digested with KpnI and XbaI, and ligated with DNA Ligase

to produce pZEMB8. Plasmids with translational reporter fusions P$_{lac0-1}$-*manX-gfp*

(pZEMB10), P$_{lac0-1}$-*yigL-gfp* (pZEMB15)*,* P$_{lac0-1}$-*purR-gfp* (pZEMB25), P$_{lac0-1}$-*asdI-gfp*

(pZEMB26) and P$_{lac0-1}$-*asdI-II-gfp* (pZEMB27) were constructed by restriction cloning

into pZEMB8 using KpnI and EcoRI restriction endonucleases.

Media and reagents. Bacteria were cultured in Luria-Bertani (LB) broth medium or LB

agar plates (37) at 37°C, unless stated otherwise. Bacteria were grown in MOPS

(morpholine-propanesulfonic acid) rich defined medium (Teknova) with 0.2% fructose

for reporter fluorescence assays. Where necessary, media was supplemented with

antibiotics at following concentrations: 100 µg/ml ampicillin (Amp), 25 µg/ml

chloramphenicol (Cm), 25 µg/ml kanamycin (Kan) and 50 µg/ml spectinomycin (Spec).

Isopropyl β-D-1-thiogalactopyranoside (IPTG) was used at concentrations of 0.1-1.5

mM for induction of P$_{lac0-1}$ promoter, anhydrotetracycline (aTc) 0-50 ng/ml for induction

of P$_{tet0-1}$ promoter and 0.000002%-0.2% L-arabinose for induction of P$_{BAD}$ promoter,

unless otherwise noted. To induce glucose-phosphate stress, 0.5% α-methylglucoside

(αMG) was added to the growth medium.

Reporter fluorescence assay. Bacterial strains were cultured overnight in MOPS rich

medium supplemented with 0.2% fructose, Amp, Cm and subcultured 1:100 to fresh

medium with appropriate inducers (IPTG, aTc) in 48 well plates. Relative fluorescence

units (RFU) and optical density (OD$_{600}$) were measured over time. "GFP expression"

was calculated by plotting RFU over OD$_{600}$ and determining the slopes of linear

regression plots for each IPTG concentration in exponentially growing cells in the

presence of aTc to induce SgrS expression. "Promoter activity" was calculated by

plotting RFU over OD$_{600}$ and determining the slopes of linear regression plots for each

IPTG concentration in exponentially growing cells in the absence of aTc.

***In vitro* transcription and radiolabeling**. Template DNA for *in vitro* transcription was

generated by PCR using gene-specific oligonucleotides containing the T7 promoter

sequence. Following oligonucleotides were used to generate specific template DNA:

MBP84F/MBP213R–*ptsG* (+1 to +240), O-JH218/MBP214R–*manX* (+1 to +240),

MBP56F/MBP215R–*asdl-II* (+1 to +240), MBP56F/MBP222R–*asdl* (+1 to +110),

MBP226F/MBP226R–*asdll* (+71 to +310), MBP65F/MBP174R–*purR* (+1 to +230),

MBP216F/MBP216R–*yigL* (-191 to +50 relative to ATG translation start of *yigL*)

MBP234F/MBP234R–*gfp* (+1 to +240) and O-JH219/O-JH119 were used to generate

full-length *sgrS* template DNA. *In vitro* transcription of DNA templates was performed

according to specifications of MEGAscript T7 Kit (Ambion). *In vitro* transcribed RNA was

5'-end labeled with radioisotope ^{32}P using the KinaseMax Kit (Ambion). Samples were

cleaned by passing through Illustra ProbeQuant G-50 Micro Columns (GE Healthcare).

Than samples were cleaned once more with phenol-chloroform: isoamyl alcohol

(Ambion) and labeled RNA precipitated with Ethanol:3M NaAc (30:1).

RNA-RNA gel electrophoretic mobility shift assay*. Different concentrations of

unlabeled mRNA were mixed with 0.02 pmol of 5'-end labeled SgrS. Samples were

denatured at 95°C for 1 min, placed on ice for 5 min, and incubated at 37°C for 30 min

in 1x binding buffer (20mM Tris-HCL (pH 8.0), 1mM DTT, 1mM MgCl$_2$, 20 mM KCl,

10mM Na$_2$HPO$_4$ (pH 8.0)) (41). Non-denaturing loading buffer was added and samples

resolved for 6 h at 40 V on native 5.6% PAGE.

Protein-RNA gel electrophoretic mobility shift assay. 0.02 pmol of 5'-end labeled

mRNA was denatured at 95°C for 1 min., placed on ice for 5 min. Different

concentrations of purified Hfq protein (His-tagged) were added. Samples were

incubated at 37°C for 30 min in 1x binding buffer (20mM Tris-HCL (pH 8.0), 1mM DTT,

1mM MgCl$_2$, 20 mM KCl, 10mM Na$_2$HPO$_4$ (pH 8.0)). Non-denaturing loading buffer was

added and samples resolved for 1h 30 min at 20 mA on native 4.0% PAGE (41).

SHAPE. The *asdI-II* RNA (0.15 µM) and SgrS RNA (0.075 µM, 0.15 µM, 0.30 µM, 0.75

µM, 1.5 µM, or 3.0 µM) were folded separately as in (42) using a modified SHAPE

buffer (100 mM HEPES [pH 8.0], 2 mM MgCl$_2$, 40 mM NaCl). For each SgrS

concentration, the SgrS RNA or the equivalent volume of 0.5X TE was added to the

asdI-II RNA and the samples were incubated at 37°C for 30 min. The RNAs were

modified with *N*-methylisatoic anhydride (NMIA, 6.5 mM; Sigma-Aldrich) and collected

by ethanol precipitation as in (42). Parallel primer extension inhibition and sequencing

reactions were performed using fluorescently labeled primers complementary to the 3'

end of the *asdI-II* RNA (5′-AGATCAAAGGCATCCTGAAG, 22.5 nM; Applied

Biosystems, ThermoFisher Scientific) as in (43) with minor modifications. Prior to primer

binding the RNAs were denatured and snap cooled and the reactions were carried out

for 20 min at 52°C, followed by 5 min at 65°C. The cDNAs were analyzed on a 3730

DNA Analyzer (Applied Biosystems, Inc.). The data were processed and SHAPE

reactivity (difference between the frequency of primer extension products at each

nucleotide in +NMIA vs. -NMIA samples) was derived using the QuShape software (44).

Data for each nucleotide were averaged with statistical outliers removed and normalized

using the 2-8% rule (45). Relative reactivity was calculated by subtracting normalized

SHAPE reactivity in the absence of the SgrS RNA from reactivity in the presence of the

WT or MT SgrS RNA.

Single-molecule fluorescence *in situ* hybridization (smFISH). The *asdI-lacZ*

(MB170), *asdII-lacZ* (MB183) and *asdI-II-lacZ* (MB171) strains were grown overnight at

37 ºC, 250 rpm in LB Broth Miller (EMD) with 25 µg/ml kanamycin (Kan) and 50 µg/ml

spectinomycin (Spec). The next day, the overnight culture was diluted 100-fold into

MOPS EZ rich defined medium (Teknova) with 0.2% (w/w) sodium succinate, 0.02%

glycerol and 0.01% L-arabinose, for *asdI-lacZ* and *asdII-lacZ* strains, and was allowed

to grow at 37 ºC till the OD_{600} reached 0.15-0.25. The concentration of L-arabinose used

for *asdI-II-lacZ* was 0.002%. α-methyl D-glucopyranoside (αMG) (Sigma-Aldrich) was

added to the culture to a desired concentration to introduce sugar phosphate stress and

induce SgrS sRNA expression. After 10 minutes of induction, the culture was taken out

and fixation was performed by mixing with formaldehyde (Fisher Scientific) at a final

concentration of 4%.

Δ*sgrS* and Δ*lacZ* strains were grown in LB Broth Miller (EMD) at 37 ºC, 250 rpm

overnight. Then the cultures were diluted 100-fold into MOPS EZ rich defined medium

(Teknova) with 0.2% glucose and allowed to grow at 37 ºC till the OD_{600} reached 0.2.

The cells were then fixed by mixing with formaldehyde (Fisher Scientific) at a final

concentration of 4%. TK310 cells were grown overnight, similar to the knockout strains.

The overnight culture was then diluted 100-fold into MOPS EZ rich defined medium

(Teknova) with 0.2% glucose and 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG,

Sigma-Aldrich) and allowed to grow at 37 ºC for 30 minutes. The cells were then fixed in

the same procedure as mentioned before.

The fixation and permeabilization of the cells were done using the methods

published previously (46). After fixing with 4% formaldehyde, the cells were incubated at

room temperature for 30 minutes. The cells were then centrifuged at 600 g for 7 minutes

and the pellets were washed with 1X PBS 3 times. The cells were then permeabilized

with 70% ethanol for 1 hour at room temperature and stored at 4 ºC before fluorescence

in situ hybridization.

The smFISH probes were designed using Stellaris Probe Designer and the

orders were placed from Biosearch Technologies (https://www.biosearchtech.com/).

The labeling of the probes was performed using equal volumes of each probe. The final

volume of sodium bicarbonate was adjusted to 0.1 M by adding 1/9 reaction volume of 1

M sodium bicarbonate (pH = 8.5). The probe solution was mixed with 0.05-0.25 mg of

Alexa Fluor 647 or Alexa Fluor 568 succinimidyl ester (Life Technologies) dissolved in 5

µL DMSO. The dye was kept about 20-25 fold in molar excess relative to the probes.

After incubation with gentle vortexing in the dark at 37 ºC overnight, the reaction was

quenched by adding 1/9 reaction volume of 3 M sodium acetate (pH = 5). Unconjugated

dyes were removed by ethanol precipitation first and then by P-6 Micro Bio-Spin

Column (Bio-Rad).

A previously published protocol (46) was used for the hybridization procedure. 60

µl of permeabilized cells were washed with FISH wash solution (10% formamide in 2X

SSC (Saline Sodium Citrate) buffer) and resuspended in 15 µl hybridization buffer (10%

dextran sulfate, 1 mg/ml *E. Coli* tRNA, 0.2 mg/ml BSA, 2 mM vanadyl ribonucleoside

complexes, 10% formamide in 2X SSC) with probes. The number of probes used for

sRNA SgrS was 9 and they were labeled with Alexa Fluor 647. The number of probes

for mRNA *lacZ* was 24 and they were labeled with Alexa Fluor 568. The concentration

of the labeled probes for SgrS and *lacZ* mRNA were 50 nM and 15 nM each. The

reactions were incubated in the dark at 30 ºC overnight. The cells were then

resuspended in 20X volume FISH wash solution and centrifuged. They were then

resuspended in FISH wash solution, incubated for 30 minutes at 30 ºC and centrifuged

and this was repeated 3 times. The cells were pelleted after the final washing step and

resuspended in 20 µl 4X SSC and stored at 4 ºC for imaging. The labeled cells were

immobilized in poly-L-lysine (Sigma-Aldrich) treated 1.0 borosilicate chambered

coverglass (Thermo Scientific™ Nunc™ Lab-Tek™). They were then imaged with

imaging buffer (50 mM Tris-HCl (pH = 8.0), 10% glucose, 1% β-mercaptoethanol

(Sigma-Aldrich), 0.5 mg/ml glucose oxidase (Sigma-Aldrich) and 0.2% catalase

(Calbiochem) in 2X SSC).

Single-molecule localization-based super-resolution imaging. An Olympus IX-71

inverted microscope with a 100X NA 1.4 SaPo oil immersion objective was used for the

3D super-resolution imaging. The lasers used for two-color imaging were Sapphire 568-

100 CW CDRH, Coherent (568nm) and DL640-100-AL-O, Crystalaser (647nm) and

DL405-025, Crystalaser (405nm) was used for the reactivation of Alexa 647 and Alexa

568 fluorophores. The laser excitation was controlled using mechanical shutters

(LS6T2, Uniblitz). A dichroic mirror (Di01-R405/488/561/635, Semrock) was used to

reflect the laser lines to the objective. The objective collected the emission signals and

then they made their way through an emission filter (FF01-594/730-25, Semrock for

Alexa 647 or HQ585/70M 63061, Chroma for Alexa 568) and excitation laser was

cleaned up using notch filters (ZET647NF, Chroma, NF01-568/647-25x5.0 and NF01-

568U-25, Semrock). They were then imaged on a 512x512 Andor EMCCD camera

(DV887ECS-BV, Andor Tech). Astigmatism was introduced by placing a cylindrical lens

with a focal length of 2 m (SCX-50.8-1000.0-UV-SLMF-520-820, CVI Melles Griot) in

the emission path between two relay lenses with focal lengths of 100 mm and 150 mm

each and this helped us to do 3D imaging. In this setup, each pixel corresponded to 100

nm. We used the CRISP (Continuous Reflective Interface Sample Placement) system

(ASI) to keep the z-drift of the setup to a minimum. The image acquisition was

controlled using the storm-control software written in Python by Zhuang's group and

available at GitHub.

599   The imaging of the sample began with a DIC image of the sample area.

600 Subsequently two-color super-resolution imaging was performed. 647nm excitation was

601 used first and after image acquisition was completed for Alexa Fluor 647, 568nm

602 excitation was used to image Alexa Fluor 568. 405nm laser power was increased slowly

603 to compensate for fluorophore bleaching and also to maintain moderate signal density.

604 We stopped imaging when most of the fluorophores had photobleached and the highest

605 reactivation laser power was reached.

606   The raw data acquired using the acquisition software was analyzed using the

607 same method as described in previously published work (31), which was a modification

608 of the algorithm published by Zhuang's group (47, 48). The clustering analysis on the

609 localization data was performed using MATLAB codes in the same method as described

610 previously (31). Background signal was estimated using $\Delta sgrS$ and $\Delta lacZ$ strains and

611 they were prepared, imaged and analyzed as described before. TK310 cells were

612 prepared, imaged and analyzed in the same way as a low copy *lacZ* mRNA sample for

613 copy number calculation. The copy number calculation was also performed using

614 MATLAB codes as described previously (31).

615 **ACKNOWLEDGEMENTS**

616   We would like to extend a special thank you to Erel Levine for providing

617 plasmids. We are grateful to Jennifer Rice, Rich Yemm, Divya Balasubramanian,

618 Chelsea Lloyd, Alisa King, Jessica Kelliher and other current and past members of the

619 Vanderpool lab for strains, plasmids and valuable advice. We appreciate and thank

620 Prof. James Slauch and members of his lab for fruitful discussions.

621 **FUNDING**

622   National Institutes of Health R01 GM092830 (M.B. and C.K.V.), R01 GM112659 (M.B.,

623   M.S.A., T.H., J.Z., and A.P.), R35 GM122569 (T.H., J.Z., and A.P.), R01 GM047823

624   (T.H.), T32 GM086252 (J.F.); National Science Foundation PHY 1430124 (T.H., J.Z.,

625   and A.P); University of Illinois Department of Microbiology James R. Beck Fellowship

626   (M. B.).

REFERENCES

1. Gisela Storz RH. 2011. Bacterial Stress Responses, Second Edition doi:10.1128/9781555816841. ASM Press, Washington DC.
2. Fang FC, Frawley ER, Tapscott T, Vazquez-Torres A. 2016. Bacterial Stress Responses during Host Infection. Cell Host Microbe 20:133-43.
3. Georg J, Hess W. 2011. cis-antisense RNA, another level of gene regulation in bacteria. Microbiology and molecular biology reviews : MMBR 75:286-300.
4. Storz G, Vogel J, Wassarman KM. 2011. Regulation by small RNAs in bacteria: expanding frontiers. Mol Cell 43:880-91.
5. Carroll RK, Weiss A, Broach WH, Wiemels RE, Mogen AB, Rice KC, Shaw LN. 2016. Genome-wide Annotation, Identification, and Global Transcriptomic Analysis of Regulatory or Small RNA Gene Expression in Staphylococcus aureus. MBio 7:e01990-15.
6. Koo JT, Lathem WW. 2012. Global discovery of small noncoding RNAs in pathogenic Yersinia species. Adv Exp Med Biol 954:305-14.
7. Zhang A, Wassarman KM, Rosenow C, Tjaden BC, Storz G, Gottesman S. 2003. Global analysis of small RNA and mRNA targets of Hfq. Mol Microbiol 50:1111-24.
8. Salvail H, Masse E. 2012. Regulating iron storage and metabolism with RNA: an overview of posttranscriptional controls of intracellular iron homeostasis. Wiley Interdiscip Rev RNA 3:26-36.
9. Sharma CM, Papenfort K, Pernitzsch SR, Mollenkopf HJ, Hinton JC, Vogel J. 2011. Pervasive post-transcriptional control of genes involved in amino acid metabolism by the Hfq-dependent GcvB small RNA. Mol Microbiol 81:1144-65.
10. Bobrovskyy M, Vanderpool CK. 2013. Regulation of bacterial metabolism by small RNAs using diverse mechanisms. Annu Rev Genet 47:209-32.
11. Bobrovskyy M, Vanderpool CK, Richards GR. 2015. Small RNAs Regulate Primary and Secondary Metabolism in Gram-negative Bacteria. Microbiol Spectr 3.
12. Vanderpool CK, Gottesman S. 2004. Involvement of a novel transcriptional activator and small RNA in post-transcriptional regulation of the glucose phosphoenolpyruvate phosphotransferase system. Mol Microbiol 54:1076-89.
13. Vanderpool CK, Gottesman S. 2007. The novel transcription factor SgrR coordinates the response to glucose-phosphate stress. J Bacteriol 189:2238-48.
14. Rice JB, Balasubramanian D, Vanderpool CK. 2012. Small RNA binding-site multiplicity involved in translational regulation of a polycistronic mRNA. Proceedings of the National Academy of Sciences of the United States of America 109:E2691-8.
15. Rice JB, Vanderpool CK. 2011. The small RNA SgrS controls sugar-phosphate accumulation by regulating multiple PTS genes. Nucleic Acids Res 39:3806-19.
16. Papenfort K, Sun Y, Miyakoshi M, Vanderpool CK, Vogel J. 2013. Small RNA-mediated activation of sugar phosphatase mRNA regulates glucose homeostasis. Cell 153:426-37.
17. Bobrovskyy M, Vanderpool CK. 2016. Diverse mechanisms of post-transcriptional repression by the small RNA regulator of glucose-phosphate stress. Mol Microbiol 99:254-73.
18. Chevance FF, Hughes KT. 2008. Coordinating assembly of a bacterial macromolecular machine. Nat Rev Microbiol 6:455-65.
19. Syed KA, Beyhan S, Correa N, Queen J, Liu J, Peng F, Satchell KJ, Yildiz F, Klose KE. 2009. The Vibrio cholerae flagellar regulatory hierarchy controls expression of virulence factors. J Bacteriol 191:6555-70.
20. Tonner PD, Pittman AM, Gulli JG, Sharma K, Schmid AK. 2015. A regulatory hierarchy controls the dynamic transcriptional response to extreme oxidative stress in archaea. PLoS Genet 11:e1004912.

678   21.   Yu H, Gerstein M. 2006. Genomic analysis of the hierarchical structure of regulatory
679   networks. Proc Natl Acad Sci U S A 103:14724-31.
680   22.   Gao R, Stock AM. 2015. Temporal hierarchy of gene expression mediated by
681   transcription factor binding affinity and activation dynamics. MBio 6:e00686-15.
682   23.   Feng L, Rutherford ST, Papenfort K, Bagert JD, van Kessel JC, Tirrell DA, Wingreen NS,
683   Bassler BL. 2015. A qrr noncoding RNA deploys four different regulatory mechanisms to
684   optimize quorum-sensing dynamics. Cell 160:228-40.
685   24.   Levine E, Zhang Z, Kuhlman T, Hwa T. 2007. Quantitative characteristics of gene
686   regulation by small RNA. PLoS Biol 5:e229.
687   25.   Rice JB, Balasubramanian D, Vanderpool CK. 2012. Small RNA binding-site multiplicity
688   involved in translational regulation of a polycistronic mRNA. Proc Natl Acad Sci U S A
689   109:E2691-8.
690   26.   Busch A, Richter AS, Backofen R. 2008. IntaRNA: efficient prediction of bacterial sRNA
691   targets incorporating target site accessibility and seed regions. Bioinformatics 24:2849-
692   56.
693   27.   Wright PR, Georg J, Mann M, Sorescu DA, Richter AS, Lott S, Kleinkauf R, Hess WR,
694   Backofen R. 2014. CopraRNA and IntaRNA: predicting small RNA targets, networks and
695   interaction domains. Nucleic Acids Res 42:W119-23.
696   28.   Morita T, Kawamoto H, Mizota T, Inada T, Aiba H. 2004. Enolase in the RNA
697   degradosome plays a crucial role in the rapid decay of glucose transporter mRNA in the
698   response to phosphosugar stress in Escherichia coli. Mol Microbiol 54:1063-75.
699   29.   Morita T, Maki K, Aiba H. 2005. RNase E-based ribonucleoprotein complexes:
700   mechanical basis of mRNA destabilization mediated by bacterial noncoding RNAs.
701   Genes Dev 19:2176-86.
702   30.   Kawamoto H, Koide Y, Morita T, Aiba H. 2006. Base-pairing requirement for RNA
703   silencing by a bacterial small RNA and acceleration of duplex formation by Hfq. Mol
704   Microbiol 61:1013-22.
705   31.   Fei J, Singh D, Zhang Q, Park S, Balasubramanian D, Golding I, Vanderpool CK, Ha T.
706   2015. RNA biochemistry. Determination of in vivo target search kinetics of regulatory
707   noncoding RNA. Science 347:1371-4.
708   32.   Morita T, Mochizuki Y, Aiba H. 2006. Translational repression is sufficient for gene
709   silencing by bacterial small noncoding RNAs in the absence of mRNA destruction. Proc
710   Natl Acad Sci U S A 103:4858-63.
711   33.   Moon K, Gottesman S. 2011. Competition among Hfq-binding small RNAs in Escherichia
712   coli. Mol Microbiol 82:1545-62.
713   34.   Santiago-Frangos A, Kavita K, Schu DJ, Gottesman S, Woodson SA. 2016. C-terminal
714   domain of the RNA chaperone Hfq drives sRNA competition and release of target RNA.
715   Proc Natl Acad Sci U S A 113:E6089-E6096.
716   35.   Schu DJ, Zhang A, Gottesman S, Storz G. 2015. Alternative Hfq-sRNA interaction
717   modes dictate alternative mRNA recognition. EMBO J 34:2557-73.
718   36.   Sun Y, Vanderpool CK. 2011. Regulation and function of Escherichia coli sugar efflux
719   transporter A (SetA) during glucose-phosphate stress. J Bacteriol 193:143-53.
720   37.   Miller JH. 1972. Experiments in Bacterial Genetics. Cold Spring Harbor Laboratory
721   Press, Cold Spring Harbor, NY.
722   38.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
723   Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640-5.
724   39.   Yu D, Ellis HM, Lee EC, Jenkins NA, Copeland NG, Court DL. 2000. An efficient
725   recombination system for chromosome engineering in Escherichia coli. Proc Natl Acad
726   Sci U S A 97:5978-83.

727   40.   Corcoran CP, Podkaminski D, Papenfort K, Urban JH, Hinton JC, Vogel J. 2012.
728   Superfolder GFP reporters validate diverse new mRNA targets of the classic porin
729   regulator, MicF RNA. Mol Microbiol 84:428-45.
730   41.   Morita T, Maki K, Aiba H. 2012. Detection of sRNA-mRNA interactions by electrophoretic
731   mobility shift assay. Methods Mol Biol 905:235-44.
732   42.   Wilkinson KA, Merino EJ, Weeks KM. 2006. Selective 2'-hydroxyl acylation analyzed by
733   primer extension (SHAPE): quantitative RNA structure analysis at single nucleotide
734   resolution. Nat Protoc 1:1610-6.
735   43.   Mortimer SA, Weeks KM. 2009. Time-resolved RNA SHAPE chemistry: quantitative
736   RNA structure analysis in one-second snapshots and at single-nucleotide resolution. Nat
737   Protoc 4:1413-21.
738   44.   Karabiber F, McGinnis JL, Favorov OV, Weeks KM. 2013. QuShape: rapid, accurate,
739   and best-practices quantification of nucleic acid probing information, resolved by
740   capillary electrophoresis. RNA 19:63-73.
741   45.   McGinnis JL, Duncan CD, Weeks KM. 2009. High-throughput SHAPE and hydroxyl
742   radical analysis of RNA structure and ribonucleoprotein assembly. Methods Enzymol
743   468:67-89.
744   46.   So LH, Ghosh A, Zong C, Sepulveda LA, Segev R, Golding I. 2011. General properties
745   of transcriptional time series in Escherichia coli. Nat Genet 43:554-60.
746   47.   Huang B, Wang W, Bates M, Zhuang X. 2008. Three-dimensional super-resolution
747   imaging by stochastic optical reconstruction microscopy. Science 319:810-3.
748   48.   Rust MJ, Bates M, Zhuang X. 2006. Sub-diffraction-limit imaging by stochastic optical
749   reconstruction microscopy (STORM). Nat Methods 3:793-5.
750   49.   Lorenz R, Luntzer D, Hofacker IL, Stadler PF, Wolfinger MT. 2016. SHAPE directed
751   RNA folding. Bioinformatics 32:145-7.
752   50.   Ishikawa H, Otaka H, Maki K, Morita T, Aiba H. 2012. The functional Hfq-binding module
753   of bacterial sRNAs consists of a double or single hairpin preceded by a U-rich sequence
754   and followed by a 3' poly(U) tail. RNA 18:1062-74.
755

FIGURE LEGENDS

Figure 1. Model for SgrS target prioritization during glucose-phosphate stress.

Glucose or the analogs αMG and 2-deoxyglucose are phosphorylated during transport

through the phosphotransferase system proteins EIICB^{Glc} (PtsG) or EIICD^{Man} (ManYZ).

If sugar-phosphates are not metabolized, the glucose-phosphate stress response is

triggered, and the transcription factor SgrR becomes active and promotes *sgrS*

transcription. The RNA chaperone Hfq promotes SgrS-mediated translational repression

of *ptsG* and *manXYZ* mRNAs, reducing synthesis of sugar transporters. SgrS stabilizes

yigL mRNA, promoting sugar phosphatase (YigL) synthesis. SgrS-mediated repression

of *asd, purR, folE* and *adiY* likely reroutes metabolism to restore homeostasis during

stress recovery. The hypothetical sequence of regulatory events following stress

induction is represented from left to right as SgrS levels increase over time. When SgrS

concentrations are low, only the highest priority targets are regulated. When stress

persists and concentrations of SgrS increase, lower priority targets are regulated.

Figure 2. Efficiency of target regulation by SgrS. A) Representation of genetic

constructs in two compatible plasmids used to study target regulation by SgrS. One

plasmid contains full-length SgrS under the control of the aTc-inducible P_{tet} promoter. A

second plasmid contains a P_{lac} promoter and the relevant region encoding each SgrS

target (including the SgrS binding site) translationally fused to a superfolder *gfp* (*sfgfp*)

reporter gene. B-F) Regulated activity was plotted as a function of basal activity (see

text for description) for (B) *ptsG*, (C) *manX*, (D) *purR,* (E) *asdI,* and (F) *yigL* fused to

sfgfp reporter gene. Without SgrS-mediated regulation we obtained a line with a slope

=1. The plots with slopes <1, indicate repression of (B) *ptsG*, (C) *manX*, (D) *purR* and

(E) *asdI* by SgrS. The plot with a slope of >1 are indicative of activation of (F) *yigL*.

Figure 3. Regulatory hierarchy established by SgrS. Regulated activity was plotted

as a function of basal activity for *ptsG*, *manX*, *purR, asdI,* and *yigL* fusions. Lack of

SgrS regulation is indicated by a line with a slope =1. The plots with slopes <1, indicate

repression (*ptsG*, *manX*, *purR* and *asdI*) by SgrS. The plot with slope >1 indicates

activation (*yigL*). Target fusion activity was monitored at different levels of SgrS

induction by aTc: (A) 10 ng/ml, (B) 20 ng/ml, (C) 30 ng/ml, (D) 40 ng/ml, (E) 50 ng/ml.

Figure 4. SgrS binding with target mRNAs *in vitro*. A) SgrS was labeled with [32]P and

incubated with unlabeled target transcripts at final concentrations of 0µM - 16 µM .

Electrophoretic mobility shift assays (EMSAs) were performed after incubating full-

length SgrS (+1 to +227) with its target transcripts (A) *ptsG* (+1 to +240), *manX* (+1 to

+240), *purR* (+1 to +230), *yigL* (-191 to +50 relative to ATG translation start of *yigL*),

asdI (+1 to +110), and *asdII* (+71 to +310). B-D) Targets transcripts (B) *ptsG* (+1 to

+240), (C) *manX* (+1 to +240), (D) *asdI* (+1 to +110) were labeled with ^{32}P and

incubated with unlabeled SgrS. EMSAs were performed to resolve complex formation.

Band densities were measured for biological replicates (n, top left) and plotted to

determine dissociation constant (K_D, bottom right) values for (B) *ptsG*, (C) *manX*, and

(D) *asdI*. E) EMSA of radiolabeled SgrS in the presence of increasing concentrations of

asdI-II transcript. Shift in mobility corresponding to one or two SgrS bound to *asdI-II* is

denoted as Site I-SgrS* and Sites I-II-SgrS* respectively. F) Quantification of SgrS

binding with radiolabeled *asdI-II* (+1 to +240), as described above.

Figure 5. Secondary structure of 5' end of *asd*. A) Diagram showing base-pairing

interactions of SgrS with binding sites I and II of *asd* mRNA. B) Energy of interaction

predicted by IntaRNA (26). "Structured" indicates pairing between full length SgrS (+1 to

+227) and *asdI-II* (+1 to +180). Plotted is the energy of interactions at either site I (*asdI*)

or site II (*asdII*). "Isolated" indicates interactions between isolated binding sites: SgrS

(+158 to +176) with *asdI* (+31 to +49) and SgrS (+158 to +178) pairing with *asdII* (+110

to +129). C) The structure of the *asdI-II* RNA alone or in complex with SgrS was probed

with NMIA and the modified RNA was analyzed by primer extension inhibition. SHAPE

reactivity (difference between the frequency of primer extension products at each

nucleotide in +NMIA vs. -NMIA samples) was then used as a parameter in the Vienna

RNAprobing WebServer (49) to predict the secondary structure of the *asdI-II* RNA.

Colors indicate SHAPE reactivity as following: red, highly reactive (≥ 0.8); gold, reactive

(0.4-0.79); green, moderately reactive (0.2-0.39); blue, minimally reactive (0.1-0.19);

grey, unreactive (< 0.01). Distinct structures were observed in the absence of SgrS and

in the presence of saturating concentrations of SgrS. The SHAPE reactivity of *asdI-II*

RNA alone (left) or in the presence of 5-fold excess SgrS (right) is mapped to the

predicted secondary structures. (D) SHAPE reactivity as a function of SgrS

concentration for each binding site (top, site I; bottom, site II). Only nucleotides with a

significant (≥ 0.1) change in reactivity are shown. Error bars denote SEM, n = 9. (E)

Relative SHAPE reactivity (difference in the SHAPE reactivity in the presence of SgrS

vs. the absence of SgrS) of the *asdI-II* RNA in the presence of wild-type (top) or mutant

(bottom) SgrS. Error bars denote SEM, n = 9 (WT), 6 (MT). The *asdI-II* RNA

nucleotides are numbered below the X-axis and the SgrS binding sites are indicated.

Figure 6. STORM imaging of SgrS regulation of *asd* variants. A) Illustration of *asdI-*

II, *asdI* and *asdII* translationally fused to *lacZ* reporter with SgrS binding sites I and II

marked. B-D) 2D projection of 3D super-resolution images of SgrS and *lacZ* mRNA for

the different *asd-lacZ* variants, labeled by smFISH, before and after 10 min induction

with 1% αMG. (B), (C) and (D) correspond to *asdI-lacZ*, *asdII-lacZ*, and *asdI-II-*

lacZ shown in (A). Probability distributions of RNA copy numbers in individual cells for

30-250 cells are plotted next to the representative images.

Figure 7. SgrS binding cooperativity allows for improved repression of *asd*

translation. A) Illustration of *asdI* and *asdI-II* constructs with SgrS binding sites marked.

Graphs show comparison of SgrS regulation of *asdI* and *asdI-II* variants in (B) wild-type

and (D) *rne701* mutant by plotting regulated activity over basal activity at various SgrS

expression levels (20-100 ng/ml aTc). Regulatory hierarchy of SgrS targets in (C) wild-

type and (E) *rne701* mutant strains. Regulation of target genes at one SgrS expression

level (40 ng/ml aTc) are compared by plotting regulated activity over basal activity of the

gfp reporter.

Figure 8. SgrS regulation of transcriptional *asd-lacZ* fusions. β-galactosidase

activity of (A) *asdI-lacZ* (+1 to +64) and *asdI-II-lacZ* (+1 to +277) was (B) assayed in

response to SgrS expression from a plasmid (and vector control) in WT and *rne701*

background strains.

Figure S1. Regulation of *ptsG* fusion by SgrS. Example plots of *ptsG-gfp*

translational fusion activity (RFU) over growth (OD_{600}) at various IPTG inducer

concentrations at (A) basal (uninduced) or (B-F) increasing SgrS expression levels.

Slopes of the linear regression plots for each IPTG concentration were calculated to

obtain (A) "basal expression" and (B-F) "regulated activity" values.

Figure S2. Inducer concentration-dependent activity of target fusions. (A) Basal

activity (0 ng/ml aTc) or (B-F) regulated activity (10-50 ng/ml aTc) of *ptsG*, *manX*, *purR*,

asdI, *asdI-II* and *yigL* fusions at varying IPTG concentrations (0-1.5 mM IPTG).

Figure S3: SHAPE analysis of *asdI-II* RNA. (A-G) SHAPE reactivity of the *asdI-II* RNA

alone and in complex with increasing concentrations of SgrS (0.5, 1, 2, 5, 10, or 20X).

The structure of the *asdI-II* RNA in the absence and presence of SgrS was probed with

NMIA and the modified RNA was analyzed by primer extension inhibition. SHAPE

reactivity is the difference between the frequency of primer extension products at each

nucleotide in +NMIA vs. -NMIA samples. Colors indicate SHAPE reactivity as following:

red, highly reactive (≥ 0.8); gold, reactive (0.4-0.79); green, moderately reactive (0.2-

0.39); blue, minimally reactive (0.1-0.19); grey, unreactive (< 0.01). (H) Relative SHAPE

reactivity of the *asdI-II* RNA in the presence of wild-type vs mutant SgrS. Relative

reactivity is the difference in the SHAPE reactivity in the presence of wild-type and

mutant SgrS. Error bars denote SEM, n = 9. The *asdI-II* RNA nucleotides are numbered

below the X-axis and the SgrS binding sites are indicated. (I-J) SHAPE reactivity as a

function of mutant SgrS concentration for each binding site (I, site I; J, site II). The same

nucleotides as in Fig. 5D are shown. Error bars denote SEM, n = 6.

Figure S4. Quantification of SgrS and *asd* mRNA variants using STORM. Copy

number of *lacZ* mRNA vs. SgrS in 30-250 individual cells for the different *asd-*

lacZ variants, (A-B) *asdI-lacZ*, (C-D) *asdII-lacZ*, and (E-F) *asdI-II-lacZ*, before (A,C,E)

and after (B,D,F) 10 min 1% αMG induction.

Figure S5. Inducer concentration-dependent activity of target-*gfp* fusions. (A)

Basal activity (0 ng/ml aTc) or (B-F) regulated activity (20-100 ng/ml aTc) of *ptsG*,

manX, *purR*, *asdI*, *asdI-II* and *yigL* fusions at varying IPTG concentrations (0-1.5 mM

IPTG).

Figure S6. Regulatory hierarchy established by SgrS in the RNase E mutant

strain. A) Regulated activity was plotted as a function of basal activity (see text for

description) for *ptsG*, *manX*, *purR, yigL*, *asdI* and *asdI-II* fused to *sfgfp* reporter gene in

the *rne701* mutant strain. Target fusion activity was monitored at different levels of SgrS

induction by aTc: 20 ng/ml, 40 ng/ml, 60 ng/ml, 80 ng/ml, 100 ng/ml. B) Target

regulation at different SgrS expression levels in the *rne701* mutant strain was compared

to determine regulatory hierarchy. Without SgrS-mediated regulation we obtained a line

with a slope =1. The plots with slopes <1, indicate repression of *ptsG*, *manX*, *purR, asdI*

and *asdI-II* by SgrS. The plot with the slope of >1 is indicative of activation of *yigL*. B)

Regulated activity was plotted as a function of basal activity for *ptsG*, *manX*, *purR, asdI,*

and *yigL* fusions.

Figure S7. Hfq binding with target mRNAs *in vitro.* A) Target transcripts *ptsG* (+1 to

+240), *manX* (+1 to +240), *purR* (+1 to +230), *yigL* (-191 to +50 relative to ATG

translation start of *yigL*) *, asdI* (+1 to +110)*, asdII* (+71 to +310) and *asdI-II* (+1 to +240)

were labeled with ^{32}P and incubated with appropriate concentrations of Hfq protein.

EMSAs were performed to resolve complex formation. Complexes of increasing size are

marked I-IV, as was previously noted for Hfq (50). B) Band densities were measured for

biological replicates (n, top left) and plotted to determine dissociation constant (K_D,

bottom right) values.

Table S1. Strains and plasmids used in this study.

Table S2. Oligonucleotides used in this study.

Figure 1. Model for SgrS target prioritization during glucose-phosphate stress.
Glucose or the analogs αMG and 2-deoxyglucose are phosphorylated during transport
through the phosphotransferase system proteins EIICB^{Glc} (PtsG) or EIICD^{Man} (ManYZ).
If sugar-phosphates are not metabolized, the glucose-phosphate stress response is
triggered, and the transcription factor SgrR becomes active and promotes *sgrS*
transcription. The RNA chaperone Hfq promotes SgrS-mediated translational repression
of *ptsG* and *manXYZ* mRNAs, reducing synthesis of sugar transporters. SgrS stabilizes
yigL mRNA, promoting sugar phosphatase (YigL) synthesis. SgrS-mediated repression
of *asd, purR, folE* and *adiY* likely reroutes metabolism to restore homeostasis during
stress recovery. The hypothetical sequence of regulatory events following stress
induction is represented from left to right as SgrS levels increase over time. When SgrS
concentrations are low, only the highest priority targets are regulated. When stress
persists and concentrations of SgrS increase, lower priority targets are regulated.

Figure 2. Efficiency of target regulation by SgrS. A) Representation of genetic constructs in two compatible plasmids used to study target regulation by SgrS. One plasmid contains full-length SgrS under the control of the aTc-inducible P_tet promoter. A second plasmid contains a P_lac promoter and the relevant region encoding each SgrS target (including the SgrS binding site) translationally fused to a superfolder *gfp* (*sfgfp*) reporter gene. B-F) Regulated activity was plotted as a function of basal activity (see text for description) for (B) *ptsG*, (C) *manX*, (D) *purR,* (E) *asdI,* and (F) *yigL* fused to *sfgfp* reporter gene. Without SgrS-mediated regulation we obtained a line with a slope =1. The plots with slopes <1, indicate repression of (B) *ptsG*, (C) *manX*, (D) *purR* and (E) *asdI* by SgrS. The plot with a slope of >1 are indicative of activation of (F) *yigL*.

Figure 3. Regulatory hierarchy established by SgrS. Regulated activity was plotted as a function of basal activity for *ptsG*, *manX*, *purR, asdI,* and *yigL* fusions. Lack of SgrS regulation is indicated by a line with a slope =1. The plots with slopes <1, indicate repression (*ptsG*, *manX*, *purR* and *asdI*) by SgrS. The plot with slope >1 indicates activation (*yigL*)*.* Target fusion activity was monitored at different levels of SgrS induction by aTc: (A) 10 ng/ml, (B) 20 ng/ml, (C) 30 ng/ml, (D) 40 ng/ml, (E) 50 ng/ml.

A

Target [µM]   0 ————————— 16   0 ————————— 16

ptsG-SgrS*   SgrS*-*yigL*

SgrS*   SgrS*

manX-SgrS*   SgrS*-*asdI*

SgrS*   SgrS*

purR-SgrS*   SgrS*-*asdII*
                                                    (expected)

SgrS*   SgrS*

B   *ptsG*   C   *manX*   D   *asdI*

n=3   n=3   n=2

Fraction Bound

1.0   1.0   1.0
0.8   0.8   0.8
0.6   0.6   0.6
0.4   0.4   0.4
0.2   Kd=0.11±0.01   0.2   Kd=19.7±2.78   0.2   Kd=0.15±0.04
0.0   0.0   0.0
 0.01 0.1   1   10   0.1   1   10 100   0.01 0.1   1   10
                                        SgrS [µM]

E   F   *asdI-II*

Target [µM]   0 ——————— 16   n=3

sites I-II-SgrS*   1.0
site I-SgrS*   0.8
                                           0.6
                                           0.4
                                           0.2   Kd=0.07±0.01
SgrS*   0.0
                                            0.01 0.1   1   10
                                                  [SgrS] µM

Figure 4 legend on the next page

Figure 4. SgrS binding with target mRNAs *in vitro*. A) SgrS was labeled with ^{32}P and incubated with unlabeled target transcripts at final concentrations of 0µM - 16 µM . Electrophoretic mobility shift assays (EMSAs) were performed after incubating full-length SgrS (+1 to +227) with its target transcripts (A) *ptsG* (+1 to +240), *manX* (+1 to +240), *purR* (+1 to +230), *yigL* (-191 to +50 relative to ATG translation start of *yigL*), *asdI* (+1 to +110), and *asdII* (+71 to +310). B-D) Targets transcripts (B) *ptsG* (+1 to +240), (C) *manX* (+1 to +240), (D) *asdI* (+1 to +110) were labeled with ^{32}P and incubated with unlabeled SgrS. EMSAs were performed to resolve complex formation. Band densities were measured for biological replicates (n, top left) and plotted to determine dissociation constant (K_D, bottom right) values for (B) *ptsG*, (C) *manX*, and (D) *asdI*. E) EMSA of radiolabeled SgrS in the presence of increasing concentrations of *asdI-II* transcript. Shift in mobility corresponding to one or two SgrS bound to *asdI-II* is denoted as Site I-SgrS* and Sites I-II-SgrS* respectively. F) Quantification of SgrS binding with radiolabeled *asdI-II* (+1 to +240), as described above.

Figure 5 legend on the next page

Figure 5. Secondary structure of 5' end of *asd*. A) Diagram showing base-pairing interactions of SgrS with binding sites I and II of *asd* mRNA. B) Energy of interaction predicted by IntaRNA (26). "Structured" indicates pairing between full length SgrS (+1 to +227) and *asdI-II* (+1 to +180). Plotted is the energy of interactions at either site I (*asdI*) or site II (*asdII*). "Isolated" indicates interactions between isolated binding sites: SgrS (+158 to +176) with *asdI* (+31 to +49) and SgrS (+158 to +178) pairing with *asdII* (+110 to +129). C) The structure of the *asdI-II* RNA alone or in complex with SgrS was probed with NMIA and the modified RNA was analyzed by primer extension inhibition. SHAPE reactivity (difference between the frequency of primer extension products at each nucleotide in +NMIA vs. -NMIA samples) was then used as a parameter in the Vienna RNAprobing WebServer (55) to predict the secondary structure of the *asdI-II* RNA. Colors indicate SHAPE reactivity as following: red, highly reactive ($\geq$ 0.8); gold, reactive (0.4-0.79); green, moderately reactive (0.2-0.39); blue, minimally reactive (0.1-0.19); grey, unreactive (< 0.01). Distinct structures were observed in the absence of SgrS and in the presence of saturating concentrations of SgrS. The SHAPE reactivity of *asdI-II* RNA alone (left) or in the presence of 5-fold excess SgrS (right) is mapped to the predicted secondary structures. (D) SHAPE reactivity as a function of SgrS concentration for each binding site (top, site I; bottom, site II). Only nucleotides with a significant ($\geq$ 0.1) change in reactivity are shown. Error bars denote SEM, n = 9. (E) Relative SHAPE reactivity (difference in the SHAPE reactivity in the presence of SgrS vs. the absence of SgrS) of the *asdI-II* RNA in the presence of wild-type (top) or mutant (bottom) SgrS. Error bars denote SEM, n = 9 (WT), 6 (MT). The *asdI-II* RNA nucleotides are numbered below the X-axis and the SgrS binding sites are indicated.

Figure 6 legend on the next page

Figure 6. STORM imaging of SgrS regulation of *asd* variants. A) Illustration of *asdI-II*, *asdI* and *asdII* translationally fused to *lacZ* reporter with SgrS binding sites I and II marked. B-D) 2D projection of 3D super-resolution images of SgrS and *lacZ* mRNA for the different *asd-lacZ* variants, labeled by smFISH, before and after 10 min induction with 1% αMG. (B), (C) and (D) correspond to *asdI-lacZ*, *asdII-lacZ*, and *asdI-II-lacZ* shown in (A). Probability distributions of RNA copy numbers in individual cells for 30-250 cells are plotted next to the representative images.

Figure 7. SgrS binding cooperativity allows for improved repression of *asd* translation. A) Illustration of *asdI* and *asdI-II* constructs with SgrS binding sites marked. Graphs show comparison of SgrS regulation of *asdI* and *asdI-II* variants in (B) wild-type and (D) *rne701* mutant by plotting regulated activity over basal activity at various SgrS expression levels (20-100 ng/ml aTc). Regulatory hierarchy of SgrS targets in (C) wild-type and (E) *rne701* mutant strains. Regulation of target genes at one SgrS expression level (40 ng/ml aTc) are compared by plotting regulated activity over basal activity of the *gfp* reporter.

Figure 8. SgrS regulation of transcriptional *asd-lacZ* fusions. β-galactosidase activity of (A) *asdI-lacZ* (+1 to +64) and *asdI-II-lacZ* (+1 to +277) was (B) assayed in response to SgrS expression from a plasmid (and vector control) in WT and *rne701* background strains.