Determinants of target prioritization and regulatory hierarchy for the bacterial small RNA SgrS

Maksym Bobrovskyy[1,2] Jane K. Frandsen[3,4], Jichuan Zhang[5], Anustup Poddar[5], Muhammad S. Azam[1], Tina M. Henkin[3], Taekjip Ha[5,6] and Carin K. Vanderpool[1]*

[1]Department of Microbiology, University of Illinois at Urbana-Champaign, 601 S. Goodwin Ave., Urbana, Illinois 61801
[2]Present address: Department of Microbiology, The University of Chicago, 920 E. 58[th] St., Chicago, Illinois 60637
[3]Department of Microbiology and Center for RNA Biology, The Ohio State University, Columbus, Ohio 43210
[4]Biochemistry Program, The Ohio State University
[5]Department of Biophysics and Biophysical Chemistry, Johns Hopkins University, Baltimore, Maryland, USA 21205
[6]Howard Hughes Medical Institute, Baltimore, Maryland, USA 21205

*Corresponding author
Carin K. Vanderpool, Ph.D.
Department of Microbiology
University of Illinois at Urbana-Champaign
C213 CLSL, MC-110
601 S. Goodwin Ave.
Urbana, IL 61801
(t) 217-333-7033
(f) 217-244-6697
cvanderp@life.illinois.edu

Running title: Target prioritization by small RNAs

Abstract word count: 241

Text word count: 6,690

ABSTRACT

The mechanisms by which small RNA (sRNA) regulators select and prioritize target mRNAs remain poorly understood, but serve to promote efficient responses to environmental cues and stresses. We sought to uncover mechanisms that establish regulatory hierarchy for a model sRNA, SgrS, found in enteric bacteria and produced under conditions of metabolic stress when sugar transport and metabolism are unbalanced. SgrS post-transcriptionally controls a nine-gene regulon to restore growth and homeostasis under stress conditions. An *in vivo* reporter system was used to quantify SgrS-dependent regulation of target genes and established that SgrS exhibits a clear preference for certain targets, and regulates those targets efficiently even at low SgrS levels. Higher SgrS concentrations are required to regulate other targets. The position of targets in the regulatory hierarchy is not well-correlated with the predicted thermodynamic stability of SgrS-mRNA interactions or the SgrS-mRNA binding affinity as measured *in vitro*. Detailed analyses of SgrS interaction with *asd* mRNA demonstrate that SgrS binds cooperatively to two sites and remodels *asd* mRNA secondary structure. SgrS binding at both sites increases the efficiency of *asd* mRNA regulation compared to mutants that have only a single SgrS binding site. Our results suggest that sRNA selection of target mRNAs and regulatory hierarchy are influenced by several molecular features. The sRNA-mRNA interaction, including the number and position of sRNA binding sites on the mRNA and cofactors like the RNA chaperone Hfq, seem to tune the efficiency of regulation of specific mRNA targets.

IMPORTANCE

To survive, bacteria must respond rapidly to stress and simultaneously maintain

metabolic homeostasis. The small RNA (sRNA) SgrS mediates the response to stress

that arises from imbalanced sugar transport and metabolism. To coordinate the stress

response, SgrS regulates genes involved in sugar uptake and metabolism. Intrinsic

properties of sRNAs such as SgrS allow them to regulate extensive networks of genes.

To date, sRNA regulation of targets has largely been studied in the context of "one

sRNA-one target", and little is known about coordination of multi-gene regulons and

sRNA regulatory network structure. Here, we explore the molecular basis for regulatory

hierarchy in sRNA regulons. Our results reveal a complex interplay of factors, including

the number and location of sRNA binding sites on mRNA targets and the participation of

an RNA chaperone, that dictate prioritized regulation of targets to promote an efficient

response to stress.

INTRODUCTION

Bacteria live in diverse niches, often encountering rapidly changing and stressful environments. Bacterial stress responses can mitigate the negative effects of stress on cell structure and function. Stress responses are usually coordinated by molecules, either RNAs or proteins, that alter expression of a regulon comprised of multiple genes. Coordinated control of the regulon prepares the cell to survive or adapt to the stress (1). Proteins control expression of target regulons by binding to DNA sequences and modulating the frequency of transcription initiation, whereas RNAs often modulate gene expression post-transcriptionally. A prevalent type of RNA regulator in bacteria is referred to simply as small RNA (sRNA). The sRNAs are often produced in response to a particular stress, and regulate target mRNAs through base pairing interactions that modify mRNA translation or stability (2, 3). Hundreds of sRNAs have been identified in diverse bacteria (4-6). While the majority of sRNAs have not been characterized, many studies suggest that sRNA regulatory networks are as extensive and complex as those controlled by proteins (7, 8).

A large body of work has illuminated base pairing-dependent molecular mechanisms of post-transcriptional regulation by sRNAs (9, 10). The sRNA SgrS (sugar-phosphate stress sRNA) has been an important model for discovery of both negative and positive mechanisms of target mRNA regulation. SgrS is induced in response to metabolic stress associated with disruption of glycolytic flux and intracellular accumulation of sugar phosphates (also referred to as glucose-phosphate stress) (11, 12). SgrS regulates at least 9 genes and promotes recovery from glucose-phosphate stress. SgrS-dependent repression of mRNAs encoding sugar transporters

(*ptsG, manXYZ*) reduces uptake of sugars to prevent further sugar-phosphate

accumulation (Fig. 1) (11, 13, 14). Activation of a sugar phosphatase (*yigL*) mRNA

promotes dephosphorylation and efflux of accumulated sugars (15), and repression of

other mRNAs is hypothesized to reroute metabolism to promote recovery from stress

(Fig. 1) (16). Each target of SgrS is regulated by a distinct molecular mechanism. How

different mechanisms of regulation yield effects of variable magnitude with respect to

mRNA stability and translation is an open question.

Temporally-ordered and hierarchical patterns of gene regulation carried out by

protein transcription factors have been characterized in many systems (17-20). These

regulatory patterns allow cells to respond efficiently to environmental signals by

prioritizing induction or repression of products needed to respond to those signals.

Protein regulators establish a hierarchy of regulation based on their affinities for binding

sites in the operator regions of different target genes. As the concentration of active

regulator increases, genes are sequentially regulated based on binding site affinity (21).

There is growing evidence that sRNAs also regulate their target genes hierarchically

(22, 23). However, the mechanisms involved in establishing and maintaining prioritized

regulation of sRNA targets are not known.

We hypothesize that the temporal progression of target regulation by SgrS is

optimized to promote efficient recovery from glucose-phosphate stress (Fig. 1). To test

this hypothesis, we first defined the efficiency of SgrS regulation of each target and

found that SgrS indeed prioritizes regulation of some targets over others. We examined

the factors that determine regulatory efficiency, including the arrangement and strength

of SgrS target binding sites and the roles of other factors like RNase E and Hfq.

Detailed characterization of a specific SgrS-mRNA target interaction revealed

cooperative binding of SgrS to two binding sites and a requirement for both binding sites

for maximal SgrS-dependent regulation. Collectively, our results uphold the hypothesis

that sRNAs regulate expression of genes in their target regulons hierarchically, and that

this is influenced by features of each sRNA-mRNA pair and different molecular

mechanisms of regulation that precisely determine the regulatory priority for each target.

RESULTS

SgrS differentially regulates targets at the level of translation. Previous studies

suggested the possibility of a regulatory hierarchy for the small RNA SgrS, which

regulates a diverse set of mRNA targets (10, 11, 13-15). To study this, we used a two-

plasmid system to control expression of SgrS and target translational fusions (Fig. 2A).

All target transcript fragments fused to *gfp* contain experimentally-confirmed SgrS

binding sites. To quantify translational regulation by SgrS and facilitate comparisons of

regulatory efficiency among targets, we analyzed the data as described previously (23).

Activity of reporter fusions was measured by monitoring GFP fluorescence over time. By

plotting the GFP fluorescence (RFU) as a function of growth (OD_{600}) for target-*gfp*

fusions in the absence of SgrS, we defined "basal activity" at different inducer

concentrations (example in Fig. S1A). This method of quantifying fusion activity

accounts for the fact that fluorescence levels are not directly proportional to inducer

concentrations ((23) and Fig. S1A). While the absolute values for basal activity differ

among different fusions, all fusions responded to induction in a dose-dependent manner

(Fig. S2A). Similar plots (RFU/OD_{600}) were generated for each fusion in the presence of

SgrS (examples Fig. S1B-F). We define "regulated activity" as the slope of the curve

(RFU/OD$_{600}$) under conditions where both the fusion and SgrS are induced (example in

Fig. S1B). As levels of SgrS increase, clear patterns of repression or induction are seen

for all target fusions (Figs S1B-F and S2B-F).

To define the efficiency of regulation of each target, we plotted regulated activity

as a function of basal activity for *ptsG*, *manX, asdI, purR,* and *yigL*. When there is no

SgrS-mediated regulation, a line with a slope of 1 is seen for all targets (Fig. 2B-F).

Slopes less than 1 indicate that the fusion is repressed by SgrS. This is true for *ptsG*,

manX, *asdI* and *purR* reporter fusions (Figure 2B-E). Slopes greater than 1 are

indicative of activation by SgrS, as seen for *yigL* (Fig. 2F). The magnitude of regulation

was responsive to SgrS levels over the range of SgrS inducer (aTc) concentrations (Fig.

2B-E). In contrast, for for *yigL,* the magnitude of activation did not increase beyond a

maximal level obtained at 20 ng/mL of inducer (Fig. 2F). While the basis for this

difference is unclear, it is likely to reflect the inherently different molecular mechanisms

of regulation, *i.e.,* mRNA stabilization for *yigL* and translational repression for other

targets.

We then compared regulatory efficiency of targets at different levels of SgrS

induction. At the two lowest levels of SgrS induction (10ng/mL and 20 ng/mL aTc), only

ptsG and *yigL* showed substantial repression and activation, respectively (Fig. 3A, B). In

contrast, *manX*, *asdI* and *purR* fusions yielded curves whose slopes remained at ~1,

indicating no regulation at lower levels of SgrS. Our interpretation is that *ptsG* and *yigL*

are high-priority targets of SgrS, since they are regulated preferentially when SgrS

levels are low. With increasing SgrS levels (20-50 ng/ml aTc), regulation of "weaker"

targets *manX, asdI* and *purR* became apparent (Fig. 3C, D, E). As SgrS levels

increased, *ptsG* repression became more efficient up to a maximal repression at 40

ng/mL of aTc, and it remained the most strongly repressed target at all levels of SgrS.

Collectively, these data suggest that SgrS targets are preferentially regulated in the

following order: 1/2) *ptsG* and *yigL*, 3) *manX,* 4) *asdI*, and 5) *purR* (Fig. 3A-E). We

hypothesize that the position of each target within the regulatory hierarchy is determined

by characteristics of SgrS-target mRNA interactions and the mechanism of SgrS-

dependent regulation.

Differences in binding affinity *in vitro* are not correlated with regulation efficiency.

One of the initial steps in sRNA-mediated regulation is formation of base-pairing

interactions with the target mRNA. Binding of the sRNA with its target mRNA is

dependent on sequence complementarity and RNA secondary structure. We examined

the characteristics of SgrS-target mRNA binding *in vitro* to determine whether the

strength of binding is correlated with the target hierarchy.

Electrophoretic mobility shift assays (EMSAs) were performed to measure

binding of SgrS to its target mRNAs *ptsG*, *manX*, *purR*, *yigL* and *asd*. SgrS bound to

ptsG with a K_D of 0.11 ± 0.01 µM (Fig. 4A, B), which was lower than K_Ds for SgrS

binding to most of the other targets (Fig. 4A-E). SgrS-*manX* mRNA binding had a K_D of

19.7 ± 2.78 µM (Fig. 4A, C), which is weaker than the interaction with *ptsG* (Fig. 4B), but

stronger compared to *purR* and *yigL* (Fig. 4A). Three different fragments of *asd* mRNA

were tested, because previous work demonstrated that SgrS pairs at two distinct sites

on *asd* mRNA (16). The first site, *asdI*, is adjacent to the ribosome binding site and is

sufficient for modest SgrS-dependent translational repression. The second site, *asdII,* is

in the coding sequence of *asd,* 60-nt downstream of the start codon. When both sites

are present (RNA *asdI-II),* stronger SgrS-dependent translational repression was

observed (16). Surprisingly, while *asdI* RNA (containing only the upstream SgrS binding

site) regulation was less efficient compared to that of *manX* (Fig. 3A-E), *in vitro* it bound

SgrS more strongly with a K_D of 0.15 ± 0.04 µM (Fig. 4A, D), which is comparable to

SgrS-*ptsG* binding (Fig. 4A, B). SgrS interaction with *asdII* was very weak (Fig. 4A). We

could not determine K_D values for SgrS interaction with *asdII*, *purR* and *yigL*, due to

limitations in obtaining high enough concentrations of RNA, but it is apparent that SgrS

binding to these targets is much weaker compared to *ptsG*, *manX* and *asdI* (Fig. 4A).

Results of EMSAs with SgrS and *asdI-II* (containing both SgrS binding sites)

revealed apparent binding cooperativity. SgrS binding to *asdI-II* has a K_D of 0.07 ± 0.01

µM (Fig. 4E, F), slightly lower than that of SgrS-*ptsG* mRNA binding. Moreover, we

observed two shifted species that correspond to binding of either one or two SgrS

sRNAs to a single *asdI-II* RNA (Fig. 4E).

Structural analyses of SgrS-*asd* mRNA interactions. Our data thus far indicate that

SgrS regulates mRNA targets in a hierarchical fashion (Figs. 2, 3). However, SgrS-

mRNA binding affinities do not explain the target hierarchy, as SgrS-*ptsG* mRNA and

SgrS-*asd* mRNA interactions have very similar K_Ds, but *ptsG* is much more efficiently

regulated than *asd* at all concentrations of SgrS (Fig. 3). To further understand the

features that influence the efficiency of target regulation, we performed more detailed

analyses of SgrS-*asd* mRNA interactions.

Previous work demonstrated that SgrS binding site I encompasses nt +31 to +49

and site II encompasses nt +110 to +127 ((16), Fig. 5A). We used IntaRNA (24, 25) to

predict the free energy (ΔG) for SgrS interactions with *asd* mRNA segments containing

both sites, or each site individually (Fig. 5B). IntaRNA accounts for the energy of

opening double-stranded RNA secondary structure, to make sequences accessible for

pairing. We first analyzed SgrS interactions with *asdI-II* mRNA (nt +1 to +240), which we

denote as "structured." Interaction of SgrS with *asd* site I has a predicted ΔG of -10.5

kcal/mol, while SgrS pairing with site II has a predicted ΔG of -1.1 kcal/mol (Fig. 5B,

structured). The ΔG for interactions between SgrS and the isolated binding sites are -18

kcal/mol for site I and -7.4 kcal/mol for site II (Fig. 5B, isolated). These predictions

suggest that SgrS interaction with site II is less favorable, particularly in the context of

the longer structured *asd* mRNA.

We investigated the structure of *asdI-II* with selective 2'-hydroxyl acylation

analyzed by primer extension (SHAPE), in which flexible nts are modified by N-

methylisotoic anhydride (NMIA), while nts constrained in helices are not reactive. In the

absence of SgrS, the sequence encompassing the *asd* ribosome binding site (+44 to

+50) is located within a structured loop (+36 to +69) on top of a short stem (+31 to +35

pairing with +70 to +74) (Fig. 5C, Fig. S3). The nts in site I (+31 to +49, Fig. 5A) are

located on the 5' side of the stem-loop structure. Most of the nts in this structure are

reactive, which is indicative of a flexible conformation that is accessible for ribosome

binding or base pairing with the seed sequence of SgrS (Fig. 5C). The seed interaction

of SgrS is likely to promote opening of the structure. Downstream of the site I stem-loop

structure is a highly structured second stem (+83 to +155) that contains site II in the

apical region (+110 to +129) (Fig. 5C, Fig. S3). Site II is sequestered in a helix and

would not be accessible to base pair with SgrS (Fig. 5C). In light of binding cooperativity

observed in Fig. 4E, we hypothesize that SgrS pairing with site I induces rearrangement

of *asd* mRNA secondary structure to facilitate interaction with site II.

We next used SHAPE to probe changes in the *asdI-II* structure in the presence of

SgrS. The reactivity of site I nt +31 to +49 decreased as the concentration of SgrS

increased (Fig. 5D), with the exception of nt +41 which is not predicted to base pair with

SgrS (Fig. 5A). The SHAPE reactivity plateaued between 5- and 10-fold excess SgrS

(Fig. S3E,F), which suggests that binding to site I was saturated. This is consistent with

a strong base-paring interaction between SgrS and site I. In contrast, the reactivity of

the site II nts +110 to +129 decreased more slowly and to a lesser extent (Fig. 5D),

consistent with a weaker and cooperative interaction. Fewer site II nts showed changes

in SHAPE reactivity upon addition of SgrS; this is likely to be due to the highly

structured nature of site II in the absence of SgrS.

The reactivity of nts outside of the SgrS binding sites also changed in the

presence of SgrS (Fig. 5E). In contrast, when a mutant SgrS that is not predicted to bind

to *asdI-II* was used, minimal changes in SHAPE reactivity were observed, which

suggests that the changes in the presence of wild-type SgrS are due to the specific

interactions between SgrS and *asdI-II* mRNA and not to the presence of additional RNA

in the system (Fig. 5E). These results indicate that SgrS binding changes the overall

structure of the *asd* RNA. A secondary structure predicted using the SHAPE data

suggests that these changes are limited to opening the SgrS binding sites and

extending the site II stem (Fig. 5C). We note an important caveat to these analyses. The

structure prediction algorithms were not designed to account for intermolecular

interactions, so this analysis may not be able to capture the *in vivo* relevant structure of

asd mRNA in complex with SgrS. Nonetheless, SHAPE data are consistent with other

analyses in demonstrating binding of SgrS to *asd* mRNA, prominently at site I and to a

lesser extent at site II.

Optimal repression by SgrS involves both pairing sites within *asd* mRNA. To

further investigate the role of the two SgrS pairing sites on *asd*, we performed stochastic

optical reconstruction microscopy (STORM) coupled with single-molecule *in situ*

hybridization (smFISH) to monitor SgrS regulation of *asd-lacZ* variants *asdI, asdII,* and

asdI-II (Fig. 6A) at single molecule resolution. In these experiments, bacteria were

grown in the presence of L-arabinose to induce expression of chromosomal *asd-lacZ*

variants, and glucose-phosphate stress was induced for 10 min by the addition of 1% α-

methyl D-glucopyranoside (αMG). 3D super-resolution images show *asd-lacZ* mRNAs

(Fig. 6B-D, green) and SgrS (Fig. 6B-D, red), as projected on 2D planes, with cells

outlined. Numbers of *asd-lacZ* mRNAs and SgrS sRNAs were counted and represented

as "copy number per cell" in histograms, with average copy number per cell indicated

above the histogram (Fig. 6B-D). SgrS induction reduced the copy number of *asdI-lacZ*

RNA by 3-fold (Fig. 6B, green) when SgrS was induced to high levels after 10 min

treatment with αMG (Fig. 6B, red and S4A, B). In contrast, the copy number of *asdII-*

lacZ RNAs (Fig. 6C, green) was not strongly affected in the presence of high SgrS

levels after αMG treatment (Fig. 6C, red and Fig. S4C, D). Copy numbers of *asdI-II-lacZ*

RNA (Fig. 6D, green) were reduced by ~8-fold after 10 min of αMG induction (Fig. 6D,

red, Fig. S6E,F). These data demonstrate that both binding sites on *asd* mRNA are

important for efficient SgrS-dependent regulation of mRNA stability.

We next examined the roles of the two SgrS binding sites in the efficiency of

translational regulation. SgrS regulation of an *asdI-II* translational fusion was compared

to regulation of an *asdI* fusion (Fig. 7A). By plotting regulated activity as a function of

basal activity as described above, we determined that SgrS repression of *asdI-II* was

more efficient than repression of *asdI* across a range of SgrS expression levels (Fig.

7B), a result in line with our previous study (16). Comparison to other targets indicated

that *asdI-II* is regulated more efficiently than *manX*, *asdI* and *purR*, at all concentrations

of SgrS (Fig. 7C).

We then compared SgrS regulation of *asdI* and *asdI-II* in the *rne701* mutant

strain deficient in degradosome assembly (26). We determined basal activity (Fig. S5A)

and regulated activity (Fig. S5B-F) of *asdI* and *asdI-II* translational *gfp* fusions at

different levels of SgrS induction. Similar to our data for the wild-type strain (Fig. 7A),

SgrS regulated *asdI-II* more efficiently compared to *asdI* in the *rne701* mutant (Fig. 7D).

Moreover, when compared to SgrS regulation of other targets, *asdI-II* was repressed

most efficiently (Fig. 7E) in the *rne701* mutant. Taken together the data indicate that the

second binding site on *asd* mRNA enhances the stringency of SgrS-mediated

regulation. Moreover, addition of the second binding site on *asd* changes its regulatory

priority relative to other targets in the SgrS regulon.

Transcriptional regulation of *asd* by SgrS. We observed that the *asdI-II* RNA is more

abundant than *asdI* (Fig. 6B, D). Since the constructs used in that experiment were

expressed from a heterologous promoter, we postulated that increased levels of *asdI-II*

compared to *asdI* mRNA must be due to increased mRNA stability or transcription

elongation of the *asdI-II* construct compared to *asdI*. We therefore constructed *asdI* and

asdI-II transcriptional fusions to *lacZ* expressed from an inducible promoter (Fig. 8A) to

test whether SgrS can regulate *asd* transcripts independent of translational regulation.

Consistent with observations from smFISH, the *asdI-II-lacZ* transcriptional fusion had

substantially higher activity compared to *asdI-lacZ* (Fig. 8B). Repression of *asdI-II* by

SgrS was more efficient (3.3-fold repression) than was observed for *asdI* (2.1-fold

repression) (Fig. 8B). SgrS still regulated both fusions in the *rne701* mutant strain (Fig.

8B). Importantly, SgrS-dependent degradation of other SgrS targets (*ptsG* mRNA (27)

and *manXYZ* mRNA (13, 14)) was abolished in degradosome mutants. Together with

our previous work, these observations suggest that SgrS regulates *asd* by two

independent mechanisms, translational repression by pairing at site I (directly occluding

the ribosome binding site) and reducing mRNA levels by promoting mRNA turnover

and/or inhibiting transcription elongation.

DISCUSSION

In this study, we set out to define the hierarchy of regulation by a model bacterial

sRNA. SgrS is a good model for this study because it has a modestly-sized regulon,

and the mechanisms of regulation of several targets have been characterized in detail

(13, 15, 16, 28). Our results demonstrate a clear pattern of prioritized regulation of

mRNA targets (Fig. 2B-F, Fig. 3A-E). Two targets in particular, *ptsG* and *yigL,* were

"high-priority" targets that were efficiently regulated even at low levels of SgrS. Other

targets, *manX, purR,* and *asd,* were less impacted by SgrS and were regulated only

when SgrS was produced at higher levels.

We investigated features of sRNA-mRNA interactions that could impact the

overall efficiency of SgrS-mediated regulation. SgrS-mRNA interactions *in vitro*, as

measured by EMSA defined K_Ds, were not well-correlated with *in vivo* regulatory

efficiency (Fig. 4A-F, Fig. 3A-E). Two targets stood out in the comparison of *in vivo*

regulation and *in vitro* SgrS-mRNA interactions. The *yigL* mRNA interaction with SgrS

was barely detectable *in vitro* (Fig. 4A), but *in vivo*, *yigL* translation was maximally

activated at low SgrS levels (Fig. 2F). Conversely, the translation of *asdI* was modestly

regulated by SgrS *in vivo* (Fig. 2E), but the *in vitro* SgrS-*asdI* interaction was

comparable to that of SgrS-*ptsG*, which had the strongest *in vivo* regulatory effect.

These apparent contradictions between *in vitro* interactions and *in vivo* regulatory

efficiency led us to further explore SgrS regulation of *asd.*

Previous work demonstrated that SgrS has two binding sites on *asd* mRNA: site I

overlaps that *asd* ribosome binding site and site II is ~60 nt downstream in the *asd*

coding sequence ((16) and Fig. 5A). EMSAs demonstrated SgrS pairing at site I alone,

but pairing at site II alone was undetectable. Binding of SgrS to an *asd* mRNA

containing both sites I and II was cooperative (Fig. 4E,F). Structural analyses of *asd*

mRNA in the absence and presence of SgrS demonstrated that SgrS indeed pairs

preferentially at site I over site II and induces substantial structural rearrangement in the

mRNA (Fig. 5C-E, Fig. S3). Quantification of SgrS-dependent degradation of *asd* mRNA

showed that site I is important, but sites I and II together promote the most efficient

regulation (Fig. 6B-D, Fig. S4). Similar to binding and regulation of mRNA degradation,

SgrS interactions at both sites I and II on *asd* mRNA improve the efficiency of

translational regulation (Fig. 7B,C, Fig. S2). These results suggest that the number and

position of sRNA binding sites on mRNA targets control regulation *in vivo* in ways that

could not be predicted based on *in vitro* characterization of sRNA-mRNA binding.

In many cases, sRNA-mediated regulation of translation is thought to indirectly

affect mRNA stability by making untranslated mRNA more susceptible to degradation by

RNase E. There are also examples of sRNA regulation, including SgrS regulation of

yigL (15), where modulation of mRNA stability is translation-independent. Truncation of

RNase E (encoded by *rne*), removing the C-terminal scaffold for degradosome

assembly, often prevents sRNA-mediated degradation of mRNA targets (14, 29, 30). If

translational regulation is the primary function of an sRNA on a given mRNA target, the

regulation should be preserved in *rne* mutant backgrounds. For SgrS targets, the

regulatory hierarchy is mostly preserved in an *rne701* degradosome mutant background

(Fig. 7, compare C and E), suggesting that for most targets, regulation of RNA stability

is not the primary mechanism of control by SgrS. Interestingly, the high-priority target

ptsG was a notable exception. In the wild-type background, *ptsG* is the most efficiently-

regulated target (Fig. 7C), whereas in the *rne701* host, it is weakly regulated. This

defect could be overcome by increasing SgrS expression levels (Fig. S6B). This result

suggests that RNase E-dependent degradation of *ptsG* mRNA is more important for its

efficient regulation by SgrS compared to other targets, where efficient regulation does

not depend on subsequent target degradation. This is consistent with the fact that *ptsG*

mRNA levels decrease at least 10-fold whereas other targets exhibit a modest 2-fold

decrease in mRNA levels upon SgrS expression (16). Our recent study quantifying

SgrS-dependent mRNA degradation at single molecule resolution indicated that *ptsG*

mRNA exhibits faster degradation kinetics than *manXYZ* mRNA (29), which could

enhance the efficiency of regulation in a wild-type but not *rne701* mutant background

where translational regulation and mRNA degradation are uncoupled.

One of our ultimate goals is to define at a molecular level the mechanisms by

which sRNAs select and prioritize regulation of their targets. The current study

implicates features of sRNA-mRNA interactions such as number and strength of sRNA

binding sites on each mRNA target and accessory factors such as RNase E in dictating

regulatory hierarchy. Another factor that is likely to play an important role in setting

regulatory priority is the RNA chaperone Hfq. EMSAs demonstrated Hfq binding to

ptsG, *manX*, *purR*, *yigL*, *asdI*, *asdII* and *asdI-II* mRNAs (Fig. S7A) with similar K_D values

for all targets (Fig. S7B). Previous work has shown that sRNAs compete for binding to

Hfq, and this competition affects their regulatory ability (31, 32). Different sRNAs can

bind to distinct sites on Hfq and this impacts their regulation of particular targets (32,

33). Additional work will be required to determine what role Hfq plays in establishing the

hierarchy of regulatory effects in sRNA regulons.

Most sRNA-mRNA interactions are conceived of as single binding site

interactions, but we have already identified two SgrS targets that deviate from this

model and have shown that additional binding sites can play important roles in

regulation and change regulation efficiency (13, 16). We have not yet discovered the

specific mechanism of regulation of *asd* mRNA by SgrS, but have shown definitively

that both binding sites are required for strong regulation. SgrS-dependent control of

both transcriptional and translational *asd* reporter fusions is not impacted in RNase E

degradosome deficient strains (Fig. 7B,D, Fig. 8B), suggesting that the regulation is not

strictly dependent on translation or mRNA turnover. Future work will test the hypothesis

that SgrS acts on *asd* mRNA at the level of transcription elongation, perhaps by an

attenuation mechanism.

In *Vibrio*, quorum sensing-regulated Qrr sRNAs regulate multiple targets by

distinct mechanisms and differences in those mechanisms influence the dynamics and

strength of regulation (22). Strong and rapid regulation is achieved by sRNAs acting

catalytically, whereby the sRNA pairs with and promotes mRNA degradation but is then

recycled for use on another mRNA target. A sequestration mechanism, where formation

of the sRNA-mRNA complex is the terminal outcome of regulation, results in slower and

weaker sRNA-dependent regulation of the target mRNA. For Qrr sRNAs, these

regulatory mechanisms seem to depend on which region of the sRNA is pairing with a

given target and whether the sRNA-mRNA interaction is strong or weak (22). While

some of the same features of SgrS-mRNA interactions may be relevant in determining

regulatory efficiency, we note that the SgrS seed sequence responsible for pairing with

all mRNA targets characterized thus far is encompassed by a short (~20 nt) mostly

single-stranded region of SgrS (11, 14-16). Moreover, we did not see a good correlation

between strong versus weak binding *in vitro* and *in vivo* regulatory efficiency. It is

possible that the "rules" governing regulatory efficiency and specific outcomes are

different for each individual sRNA. Detailed analysis of more model sRNAs will be

needed to illuminate broad general principles.

Beyond defining interesting molecular features of sRNA-mRNA interactions,

defining regulatory hierarchy for sRNA regulons is important for understanding bacterial

physiology. The vast majority of sRNA regulons remain undefined, and thus many

sRNA functions are unknown. For novel sRNAs, distinguishing high-priority from weaker

targets may provide crucial clues to the predominant role of the sRNA in cell physiology.

For SgrS, the regulatory hierarchy we have defined here is perfectly consistent with

growth studies that demonstrate the primary importance of SgrS regulation of sugar

transport and efflux under glucose-phosphate stress conditions (34). The hierarchy of

regulation by sRNAs is likely to have evolved to promote rapid and efficient responses

to environmental signals that would provide cells with a competitive growth advantage in

their specific niche. It will be crucial to develop tools to more rapidly characterize sRNA

regulatory hierarchy to better understand the functions of the hundreds of

uncharacterized sRNAs in diverse bacteria.

MATERIALS AND METHODS

Strain and plasmid construction. Strains and plasmids used in this study are listed in

Table S1. All strains used in this study are derivatives of *E. coli* K-12 strain MG1655.

Oligonucleotide primers and 5' biotinylated probes used in this study are listed in Table

S2 and were purchased from Integrated DNA Technologies. Chromosomal alleles were

moved between strains by P1 *vir* transduction (35) and inserted using λ Red

recombination (36, 37).

Translational reporter fusion alleles P_{BAD}-*asdI-II-lacZ* (MBP151F/MBP193R

primers), P_{BAD}-*asdI-lacZ* (MBP151F/MBP151R primers) and P_{BAD}-*asdII-lacZ*

(MBP193F/MBP193R primers) were constructed by PCR amplification of desired

fragments using primers containing homologies to P_{BAD} and *lacZ.* Similarly,

transcriptional fusions P_{BAD}-*asdI-II-lacZ* (MBP151F/MBP206R3 primers) and P_{BAD}-*asdI-*

lacZ (MBP151F/MBP206R1 primers) were generated by PCR amplification using a

forward primer with homology to P_{BAD} and reverse primers containing *lacZ* RBS and

lacZ homology. PCR products were then recombined into PM1205 strain using λ Red

homologous recombination.

A plasmid harboring SgrS under the control of P_{tet0-1} promoter was constructed

by PCR amplification of *sgrS* from *E. coli* MG1655 chromosomal DNA using primers

containing NdeI and BamHI restriction sites. The resulting PCR product and vector

pZA31R (23) were digested with NdeI and BamHI (New England Biolabs) restriction

endonucleases. Digestion products were ligated using DNA Ligase (New England

Biolabs) to produce plasmid pZAMB1 containing P_{tet0-1}-*sgrS* allele.

Plasmid pZEMB8 containing P_{lac0-1}-*ptsG-gfp* was constructed by PCR

amplification of *ptsG* from MG1655 chromosomal DNA using primers containing KpnI

and EcoRI restriction sites. The resulting PCR products and vector pZE12S (23) were

digested with KpnI and EcoRI restriction endonucleases. Digestion products were

ligated using DNA Ligase to produce pZEMB2. Superfolder *gfp* (referred to as *gfp*) was

amplified from pXG10-SF (38) using primrs containing KpnI and XbaI restriction sites.

pZEMB2 and the resulting PCR product were digested with KpnI and XbaI, and ligated

with DNA ligase to produce pZEMB8. Plasmids with translational reporter fusions P_{lac0-1}-

manX-gfp (pZEMB10), P_{lac0-1}-*yigL-gfp* (pZEMB15)*,* P_{lac0-1}-*purR-gfp* (pZEMB25), P_{lac0-1}-

asdI-gfp (pZEMB26) and P_{lac0-1}-*asdI-II-gfp* (pZEMB27) were constructed by restriction

cloning into pZEMB8 using KpnI and EcoRI restriction endonucleases.

Media and reagents. Bacteria were cultured in Luria-Bertani (LB) broth medium or LB

agar plates (35) at 37°C, unless stated otherwise. Bacteria were grown in MOPS

(morpholine-propanesulfonic acid) rich defined medium (Teknova) with 0.2% fructose

for reporter fluorescence assays. When necessary, media were supplemented with

antibiotics at the following concentrations: 100 µg/ml ampicillin (Amp), 25 µg/ml

chloramphenicol (Cm), 25 µg/ml kanamycin (Kan) and 50 µg/ml spectinomycin (Spec).

Isopropyl β-D-1-thiogalactopyranoside (IPTG) was used at concentrations of 0.1-1.5

mM for induction of P_{lac0-1} promoter, anhydrotetracycline (aTc) was used at 0-50 ng/ml

for induction of P_{tet0-1} promoter and L-arabinose was used at 0.000002%-0.2% for

induction of P_{BAD} promoter, unless otherwise noted. To induce glucose-phosphate

stress, 0.5% α-methylglucoside (αMG) was added to the growth medium.

Reporter fluorescence assay. Bacterial strains were cultured overnight in MOPS rich

medium supplemented with 0.2% fructose, Amp, and Cm, and diluted 1:100 into fresh

medium with appropriate inducers (IPTG, aTc) in 48 well plates. Relative fluorescence

units (RFU) and optical density (OD_{600}) were measured over time. "GFP expression"

was calculated by plotting RFU over OD_{600} and determining the slopes of linear

regression plots for each IPTG concentration in exponentially growing cells in the

presence of aTc to induce SgrS expression. "Promoter activity" was calculated by

plotting RFU over OD_{600} and determining the slopes of linear regression plots for each

IPTG concentration in exponentially growing cells in the absence of aTc.

***In vitro* transcription and radiolabeling**. Template DNA for *in vitro* transcription was

generated by PCR using gene-specific oligonucleotides containing the T7 RNA

polymerase promoter sequence. The following oligonucleotides were used to generate

specific template DNA: MBP84F/MBP213R–*ptsG* (+1 to +240), O-JH218/MBP214R–

manX (+1 to +240), MBP56F/MBP215R–*asdI-II* (+1 to +240), MBP56F/MBP222R–*asdI*

(+1 to +110), MBP226F/MBP226R–*asdII* (+71 to +310), MBP65F/MBP174R–*purR* (+1

to +230), MBP216F/MBP216R–*yigL* (-191 to +50 relative to ATG translation start of

yigL) MBP234F/MBP234R–*gfp* (+1 to +240) and O-JH219/O-JH119 were used to

generate full-length *sgrS* template DNA. *In vitro* transcription of DNA templates was

performed according to specifications of the MEGAscript T7 Kit (Ambion). *In vitro*

transcribed RNA was 5'-end labeled with radioisotope ^{32}P using the KinaseMax Kit

(Ambion), according to the manufecturer's instructions. Samples were purified by

passing through Illustra ProbeQuant G-50 Micro Columns (GE Healthcare) followed by

extraction with phenol-chloroform:isoamyl alcohol (Ambion), and labeled RNA was

precipitated with ethanol:3M NaAc (30:1).

RNA-RNA gel electrophoretic mobility shift assay. Different concentrations of

unlabeled mRNA were mixed with 0.02 pmol of 5'-end labeled SgrS. Samples were

denatured at 95°C for 1 min, placed on ice for 5 min, and incubated at 37°C for 30 min

in 1x binding buffer (20 mM Tris-HCL [pH 8.0], 1mM DTT, 1 mM MgCl$_2$, 20 mM KCl, 10

mM Na$_2$HPO$_4$ [pH 8.0]) (39). Non-denaturing loading buffer was added and samples

were resolved for 6 h at 40 V on native 5.6% PAGE.

Protein-RNA gel electrophoretic mobility shift assay. 0.02 pmol of 5'-end labeled

mRNA was denatured at 95°C for 1 min, and placed on ice for 5 min. Different

concentrations of purified Hfq protein (His-tagged) were added. Samples were

incubated at 37°C for 30 min in 1x binding buffer. Non-denaturing loading buffer was

added and samples were resolved for 90 min at 20 mA on native 4.0% PAGE (39).

SHAPE. The *asdI-II* RNA (0.15 µM) and SgrS RNA (0.075 µM, 0.15 µM, 0.30 µM, 0.75

µM, 1.5 µM, or 3.0 µM) were folded separately as in (40) using a modified SHAPE

buffer (100 mM HEPES [pH 8.0], 2 mM MgCl$_2$, 40 mM NaCl). For each SgrS

concentration, the SgrS RNA or the equivalent volume of 0.5X TE was added to the

asdI-II RNA and the samples were incubated at 37°C for 30 min. The RNAs were

modified with *N*-methylisatoic anhydride (NMIA, 6.5 mM; Sigma-Aldrich) and collected

by ethanol precipitation as in (40). Parallel primer extension inhibition and sequencing

reactions were performed using fluorescently labeled primers complementary to the 3'

end of the *asdI-II* RNA (5′-AGATCAAAGGCATCCTGAAG, 22.5 nM; Applied

Biosystems, ThermoFisher Scientific) as in (41) with minor modifications. Prior to primer

binding, the RNAs were denatured and snap cooled and the reactions were carried out

for 20 min at 52°C, followed by 5 min at 65°C. The cDNAs were analyzed on a 3730

DNA Analyzer (Applied Biosystems, Inc.). The data were processed and SHAPE

reactivity (difference between the frequency of primer extension products at each

nucleotide in +NMIA vs. -NMIA samples) was derived using the QuShape software (42).

Data for each nucleotide were averaged with statistical outliers removed and normalized

using the 2-8% rule (43). Relative reactivity was calculated by subtracting normalized

SHAPE reactivity in the absence of the SgrS RNA from reactivity in the presence of the

WT or MT SgrS RNA.

Single-molecule fluorescence *in situ* hybridization (smFISH). The *asdI-lacZ*

(MB170), *asdII-lacZ* (MB183) and *asdI-II-lacZ* (MB171) strains were grown overnight at

37ºC with shaking at 250 rpm in LB Broth Miller (EMD) containing Kan and Spec. The

next day, the overnight cultures were diluted 100-fold into MOPS EZ rich defined

medium (Teknova) with 0.2% (w/w) sodium succinate, 0.02% glycerol and L-arabinose

(0.01% for the *asdI-lacZ* and *asdII-lacZ* strains, 0.002% for *asdI-II-lacZ*) and were

allowed to grow at 37ºC till the OD_{600} reached 0.15-0.25. αMG was added to the culture

to a desired concentration to introduce sugar phosphate stress and induce SgrS sRNA

expression. After 10 min of induction, the cells were fixed by mixing with formaldehyde

(Fisher Scientific) at a final concentration of 4%.

$\Delta sgrS$ and $\Delta lacZ$ strains were grown in LB Broth Miller (EMD) at 37ºC with

shaking at 250 rpm overnight. The cultures were diluted 100-fold into MOPS EZ rich

defined medium (Teknova) with 0.2% glucose and allowed to grow at 37ºC until the

OD_{600} reached 0.2. The cells were then fixed by mixing with formaldehyde at a final

concentration of 4%. TK310 cells were grown overnight, similar to the knockout strains.

The overnight culture was then diluted 100-fold into MOPS EZ rich defined medium

(Teknova) with 0.2% glucose and 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG,

Sigma-Aldrich) and allowed to grow at 37ºC for 30 minutes. The cells were then fixed as

described above.

The fixation and permeabilization of the cells were done using the methods

published previously (44). After fixing with 4% formaldehyde, the cells were incubated at

room temperature for 30 min. The cells were then centrifuged at 600 x g for 7 min and

the pellets were washed with three times with 1X PBS 3. The cells were then

permeabilized with 70% ethanol for 1 h at room temperature and stored at 4ºC before

fluorescence *in situ* hybridization.

The smFISH probes were designed using Stellaris Probe Designer and

purchased from Biosearch Technologies (https://www.biosearchtech.com/). The labeling

of the probes was performed using equal volumes of each probe. The final volume of

sodium bicarbonate was adjusted to 0.1 M by adding 1/9 reaction volume of 1 M sodium

bicarbonate (pH = 8.5). The probe solution was mixed with 0.05-0.25 mg of Alexa Fluor

647 or Alexa Fluor 568 succinimidyl ester (Life Technologies) dissolved in 5 µL DMSO.

The dye was kept at 20-25 fold in molar excess relative to the probes. After incubation

with gentle vortexing in the dark at 37ºC overnight, the reaction was quenched by

adding 1/9 reaction volume of 3 M sodium acetate (pH 5). Unconjugated dye was

removed by ethanol precipitation followed by P-6 Micro Bio-Spin Column (Bio-Rad).

A previously published protocol (44) was used for the hybridization procedure. 60

µl of permeabilized cells were washed with FISH wash solution (10% formamide in 2X

SSC [Saline Sodium Citrate] buffer) and resuspended in 15 µl hybridization buffer (10%

dextran sulfate, 1 mg/ml *E. coli* tRNA, 0.2 mg/ml BSA, 2 mM vanadyl ribonucleoside

complexes, 10% formamide in 2X SSC) with probes. Nine probes labeled with Alexa

Fluor 647 were used for sRNA SgrS and 24 probes labeled with Alexa Fluor 568 were

used for mRNA *lacZ*. The concentration of the labeled probes for SgrS and *lacZ* mRNA

were 50 nM and 15 nM, respectively. The reactions were incubated in the dark at 3 ºC

overnight. The cells were then resuspended in 20X volume FISH wash solution and

centrifuged. They were then resuspended in FISH wash solution, incubated for 30 min

at 30ºC and centrifuged and this was repeated 3 times. The cells were pelleted after the

final washing step and resuspended in 20 µl 4X SSC and stored at 4ºC for imaging. The

labeled cells were immobilized on a poly-L-lysine (Sigma-Aldrich) treated 1.0

borosilicate chambered coverglass (Thermo Scientific™ Nunc™ Lab-Tek™). They were

then imaged in imaging buffer (50 mM Tris-HCl [pH = 8.0], 10% glucose, 1% β-

mercaptoethanol [Sigma-Aldrich], 0.5 mg/ml glucose oxidase [Sigma-Aldrich] and 0.2%

catalase [Calbiochem] in 2X SSC).

Single-molecule localization-based super-resolution imaging. An Olympus IX-71

inverted microscope with a 100X NA 1.4 SaPo oil immersion objective was used for the

3D super-resolution imaging. The lasers used for two-color imaging were Sapphire 568-

100 CW CDRH, Coherent (568nm) and DL640-100-AL-O, Crystalaser (647nm) and

DL405-025. Crystalaser (405nm) was used for the reactivation of Alexa 647 and Alexa

568 fluorophores. The laser excitation was controlled using mechanical shutters

(LS6T2, Uniblitz). A dichroic mirror (Di01-R405/488/561/635, Semrock) was used to

reflect the laser lines to the objective. The objective collected the emission signals and

then they made their way through an emission filter (FF01-594/730-25, Semrock for

Alexa 647 or HQ585/70M 63061, Chroma for Alexa 568) and excitation laser was

cleaned up using notch filters (ZET647NF, Chroma, NF01-568/647-25x5.0 and NF01-

568U-25, Semrock). Samples were then imaged on a 512x512 Andor EMCCD camera

(DV887ECS-BV, Andor Tech). Astigmatism was introduced by placing a cylindrical lens

with a focal length of 2 m (SCX-50.8-1000.0-UV-SLMF-520-820, CVI Melles Griot) in

the emission path between two relay lenses with focal lengths of 100 mm and 150 mm

each, which facilitated 3D imaging. In this setup, each pixel corresponded to 100 nm.

We used the CRISP (Continuous Reflective Interface Sample Placement) system (ASI)

to keep the z-drift of the setup to a minimum. The image acquisition was controlled

using the storm-control software written in Python by Zhuang and coworkers (45, 46)

and available at GitHub.

The imaging of the sample began with a DIC image of the sample area.

Subsequently two-color super-resolution imaging was performed. 647nm excitation was

used first and after image acquisition was completed for Alexa Fluor 647, 568nm

excitation was used to image Alexa Fluor 568. 405 nm laser power was increased

slowly to compensate for fluorophore bleaching and also to maintain moderate signal

density. Imaging was stopped when most of the fluorophores had photobleached and

the highest reactivation laser power was reached.

The raw data acquired using the acquisition software were analyzed using the

method described in previously published work (29), which was a modification of the

algorithm published by Zhuang and coworkers (45, 46). The clustering analysis on the

localization data was performed using MATLAB codes as described previously (29).

Background signal was estimated using Δ*sgrS* and Δ*lacZ* strains and they were

prepared, imaged and analyzed as described above. TK310 cells were prepared,

imaged and analyzed in the same way as a low copy *lacZ* mRNA sample for copy

number calculation. The copy number calculation was also performed using MATLAB

codes as described previously (29).

ACKNOWLEDGEMENTS

609   We would like to extend a special thank you to Erel Levine for providing

610   plasmids. We are grateful to Jennifer Rice, Rich Yemm, Divya Balasubramanian,

611   Chelsea Lloyd, Alisa King, Jessica Kelliher and other current and past members of the

612   Vanderpool lab for strains, plasmids and valuable advice. We appreciate and thank

613   Prof. James Slauch and members of his lab for fruitful discussions.

FUNDING

615   National Institutes of Health R01 GM092830 (M.B. and C.K.V.), R01 GM112659 (M.B.,

616   M.S.A., T.H., J.Z., and A.P.), R35 GM122569 (T.H., J.Z., and A.P.), R01 GM047823

617   (T.M.H.), T32 GM086252 (J.K.F.); National Science Foundation PHY 1430124 (T.H.,

618   J.Z., and A.P); University of Illinois Department of Microbiology James R. Beck

619   Fellowship (M.B.).

REFERENCES

1. Fang FC, Frawley ER, Tapscott T, Vazquez-Torres A. 2016. Bacterial stress responses during host infection. Cell Host Microbe 20:133-43.
2. Georg J, Hess W. 2011. *Cis*-antisense RNA, another level of gene regulation in bacteria. Microbiology and molecular biology reviews : MMBR 75:286-300.
3. Storz G, Vogel J, Wassarman KM. 2011. Regulation by small RNAs in bacteria: expanding frontiers. Mol Cell 43:880-91.
4. Carroll RK, Weiss A, Broach WH, Wiemels RE, Mogen AB, Rice KC, Shaw LN. 2016. Genome-wide annotation, identification, and global transcriptomic analysis of regulatory or Small RNA Gene Expression in Staphylococcus aureus. MBio 7:e01990-15.
5. Koo JT, Lathem WW. 2012. Global discovery of small noncoding RNAs in pathogenic *Yersinia* species. Adv Exp Med Biol 954:305-14.
6. Zhang A, Wassarman KM, Rosenow C, Tjaden BC, Storz G, Gottesman S. 2003. Global analysis of small RNA and mRNA targets of Hfq. Mol Microbiol 50:1111-24.
7. Salvail H, Masse E. 2012. Regulating iron storage and metabolism with RNA: an overview of posttranscriptional controls of intracellular iron homeostasis. Wiley Interdiscip Rev RNA 3:26-36.
8. Sharma CM, Papenfort K, Pernitzsch SR, Mollenkopf HJ, Hinton JC, Vogel J. 2011. Pervasive post-transcriptional control of genes involved in amino acid metabolism by the Hfq-dependent GcvB small RNA. Mol Microbiol 81:1144-65.
9. Bobrovskyy M, Vanderpool CK. 2013. Regulation of bacterial metabolism by small RNAs using diverse mechanisms. Annu Rev Genet 47:209-32.
10. Bobrovskyy M, Vanderpool CK, Richards GR. 2015. Small RNAs regulate primary and secondary metabolism in gram-negative bacteria. Microbiol Spectr 3.
11. Vanderpool CK, Gottesman S. 2004. Involvement of a novel transcriptional activator and small RNA in post-transcriptional regulation of the glucose phosphoenolpyruvate phosphotransferase system. Mol Microbiol 54:1076-89.
12. Vanderpool CK, Gottesman S. 2007. The novel transcription factor SgrR coordinates the response to glucose-phosphate stress. J Bacteriol 189:2238-48.
13. Rice JB, Balasubramanian D, Vanderpool CK. 2012. Small RNA binding-site multiplicity involved in translational regulation of a polycistronic mRNA. Proc Natl Acad Sci U S A 109:E2691-8.
14. Rice JB, Vanderpool CK. 2011. The small RNA SgrS controls sugar-phosphate accumulation by regulating multiple PTS genes. Nucleic Acids Res 39:3806-19.
15. Papenfort K, Sun Y, Miyakoshi M, Vanderpool CK, Vogel J. 2013. Small RNA-mediated activation of sugar phosphatase mRNA regulates glucose homeostasis. Cell 153:426-37.
16. Bobrovskyy M, Vanderpool CK. 2016. Diverse mechanisms of post-transcriptional repression by the small RNA regulator of glucose-phosphate stress. Mol Microbiol 99:254-73.
17. Chevance FF, Hughes KT. 2008. Coordinating assembly of a bacterial macromolecular machine. Nat Rev Microbiol 6:455-65.
18. Syed KA, Beyhan S, Correa N, Queen J, Liu J, Peng F, Satchell KJ, Yildiz F, Klose KE. 2009. The *Vibrio cholerae* flagellar regulatory hierarchy controls expression of virulence factors. J Bacteriol 191:6555-70.
19. Tonner PD, Pittman AM, Gulli JG, Sharma K, Schmid AK. 2015. A regulatory hierarchy controls the dynamic transcriptional response to extreme oxidative stress in archaea. PLoS Genet 11:e1004912.
20. Yu H, Gerstein M. 2006. Genomic analysis of the hierarchical structure of regulatory networks. Proc Natl Acad Sci U S A 103:14724-31.

21.   Gao R, Stock AM. 2015. Temporal hierarchy of gene expression mediated by
transcription factor binding affinity and activation dynamics. MBio 6:e00686-15.
22.   Feng L, Rutherford ST, Papenfort K, Bagert JD, van Kessel JC, Tirrell DA, Wingreen NS,
Bassler BL. 2015. A qrr noncoding RNA deploys four different regulatory mechanisms to
optimize quorum-sensing dynamics. Cell 160:228-40.
23.   Levine E, Zhang Z, Kuhlman T, Hwa T. 2007. Quantitative characteristics of gene
regulation by small RNA. PLoS Biol 5:e229.
24.   Busch A, Richter AS, Backofen R. 2008. IntaRNA: efficient prediction of bacterial sRNA
targets incorporating target site accessibility and seed regions. Bioinformatics 24:2849-
56.
25.   Wright PR, Georg J, Mann M, Sorescu DA, Richter AS, Lott S, Kleinkauf R, Hess WR,
Backofen R. 2014. CopraRNA and IntaRNA: predicting small RNA targets, networks and
interaction domains. Nucleic Acids Res 42:W119-23.
26.   Morita T, Kawamoto H, Mizota T, Inada T, Aiba H. 2004. Enolase in the RNA
degradosome plays a crucial role in the rapid decay of glucose transporter mRNA in the
response to phosphosugar stress in *Escherichia coli*. Mol Microbiol 54:1063-75.
27.   Morita T, Maki K, Aiba H. 2005. RNase E-based ribonucleoprotein complexes:
mechanical basis of mRNA destabilization mediated by bacterial noncoding RNAs.
Genes Dev 19:2176-86.
28.   Kawamoto H, Koide Y, Morita T, Aiba H. 2006. Base-pairing requirement for RNA
silencing by a bacterial small RNA and acceleration of duplex formation by Hfq. Mol
Microbiol 61:1013-22.
29.   Fei J, Singh D, Zhang Q, Park S, Balasubramanian D, Golding I, Vanderpool CK, Ha T.
2015. RNA biochemistry. Determination of *in vivo* target search kinetics of regulatory
noncoding RNA. Science 347:1371-4.
30.   Morita T, Mochizuki Y, Aiba H. 2006. Translational repression is sufficient for gene
silencing by bacterial small noncoding RNAs in the absence of mRNA destruction. Proc
Natl Acad Sci U S A 103:4858-63.
31.   Moon K, Gottesman S. 2011. Competition among Hfq-binding small RNAs in *Escherichia*
coli. Mol Microbiol 82:1545-62.
32.   Santiago-Frangos A, Kavita K, Schu DJ, Gottesman S, Woodson SA. 2016. C-terminal
domain of the RNA chaperone Hfq drives sRNA competition and release of target RNA.
Proc Natl Acad Sci U S A 113:E6089-E6096.
33.   Schu DJ, Zhang A, Gottesman S, Storz G. 2015. Alternative Hfq-sRNA interaction
modes dictate alternative mRNA recognition. EMBO J 34:2557-73.
34.   Sun Y, Vanderpool CK. 2011. Regulation and function of *Escherichia coli* sugar efflux
transporter A (SetA) during glucose-phosphate stress. J Bacteriol 193:143-53.
35.   Miller JH. 1972. Experiments in bacterial genetics. Cold Spring Harbor Laboratory Press,
Cold Spring Harbor, NY.
36.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640-5.
37.   Yu D, Ellis HM, Lee EC, Jenkins NA, Copeland NG, Court DL. 2000. An efficient
recombination system for chromosome engineering in *Escherichia coli*. Proc Natl Acad
Sci U S A 97:5978-83.
38.   Corcoran CP, Podkaminski D, Papenfort K, Urban JH, Hinton JC, Vogel J. 2012.
Superfolder GFP reporters validate diverse new mRNA targets of the classic porin
regulator, MicF RNA. Mol Microbiol 84:428-45.
39.   Morita T, Maki K, Aiba H. 2012. Detection of sRNA-mRNA interactions by electrophoretic
mobility shift assay. Methods Mol Biol 905:235-44.

40.   Wilkinson KA, Merino EJ, Weeks KM. 2006. Selective 2'-hydroxyl acylation analyzed by
primer extension (SHAPE): quantitative RNA structure analysis at single nucleotide
resolution. Nat Protoc 1:1610-6.
41.   Mortimer SA, Weeks KM. 2009. Time-resolved RNA SHAPE chemistry: quantitative
RNA structure analysis in one-second snapshots and at single-nucleotide resolution. Nat
Protoc 4:1413-21.
42.   Karabiber F, McGinnis JL, Favorov OV, Weeks KM. 2013. QuShape: rapid, accurate,
and best-practices quantification of nucleic acid probing information, resolved by
capillary electrophoresis. RNA 19:63-73.
43.   McGinnis JL, Duncan CD, Weeks KM. 2009. High-throughput SHAPE and hydroxyl
radical analysis of RNA structure and ribonucleoprotein assembly. Methods Enzymol
468:67-89.
44.   So LH, Ghosh A, Zong C, Sepulveda LA, Segev R, Golding I. 2011. General properties
of transcriptional time series in *Escherichia coli*. Nat Genet 43:554-60.
45.   Rust MJ, Bates M, Zhuang X. 2006. Sub-diffraction-limit imaging by stochastic optical
reconstruction microscopy (STORM). Nat Methods 3:793-5.
46.   Huang B, Wang W, Bates M, Zhuang X. 2008. Three-dimensional super-resolution
imaging by stochastic optical reconstruction microscopy. Science 319:810-3.
47.   Lorenz R, Luntzer D, Hofacker IL, Stadler PF, Wolfinger MT. 2016. SHAPE directed
RNA folding. Bioinformatics 32:145-7.
48.   Ishikawa H, Otaka H, Maki K, Morita T, Aiba H. 2012. The functional Hfq-binding module
of bacterial sRNAs consists of a double or single hairpin preceded by a U-rich sequence
and followed by a 3' poly(U) tail. RNA 18:1062-74.
742

FIGURE LEGENDS

Figure 1. Model for SgrS target prioritization during glucose-phosphate stress.

Glucose or the analogs αMG and 2-deoxyglucose are phosphorylated during transport

through the phosphotransferase system proteins EIICB^{Glc} (PtsG) or EIICD^{Man} (ManYZ).

If sugar-phosphates are not metabolized, the glucose-phosphate stress response is

triggered, and the transcription factor SgrR becomes active and promotes *sgrS*

transcription. The RNA chaperone Hfq promotes SgrS-mediated translational repression

of *ptsG* and *manXYZ* mRNAs, reducing synthesis of sugar transporters. SgrS stabilizes

yigL mRNA, promoting sugar phosphatase (YigL) synthesis. SgrS-mediated repression

of *asd, purR, folE* and *adiY* likely reroutes metabolism to restore homeostasis during

stress recovery. The hypothetical sequence of regulatory events following stress

induction is represented from left to right as SgrS levels increase over time. When SgrS

concentrations are low, only the highest priority targets are regulated. When stress

persists and concentrations of SgrS increase, lower priority targets are regulated.

Figure 2. Efficiency of target regulation by SgrS. A) Representation of genetic

constructs in two compatible plasmids used to study target regulation by SgrS. One

plasmid contains full-length SgrS under the control of the aTc-inducible P_{Tet} promoter.

A second plasmid contains a P_{Lac} promoter and the relevant region encoding each SgrS

target (including the SgrS binding site) translationally fused to a superfolder *gfp* reporter

gene. B-F) Regulated activity was plotted as a function of basal activity (see text for

description) for (B) *ptsG*, (C) *manX*, (D) *purR,* (E) *asdI,* and (F) *yigL* fused to *sfgfp*

reporter gene. Without SgrS-mediated regulation, we obtained a line with a slope =1.

The plots with slopes <1 indicate repression of (B) *ptsG*, (C) *manX*, (D) *purR* and (E)

asdI by SgrS. The plot with a slope of >1 is indicative of activation of (F) *yigL*.

Figure 3. Regulatory hierarchy established by SgrS. Regulated activity was plotted

as a function of basal activity for *ptsG*, *manX*, *purR, asdI,* and *yigL* fusions. Lack of

SgrS regulation is indicated by a line with a slope =1. The plots with slopes <1 indicate

repression (*ptsG*, *manX*, *purR* and *asdI*) by SgrS. The plot with slope >1 indicates

activation (*yigL*)*.* Target fusion activity was monitored at different levels of SgrS

induction by aTc: (A) 10 ng/ml, (B) 20 ng/ml, (C) 30 ng/ml, (D) 40 ng/ml, (E) 50 ng/ml.

Figure 4. SgrS binding to target mRNAs *in vitro*. A) SgrS was labeled with [32]P and

incubated with unlabeled target transcripts at final concentrations of 0 µM - 16 µM .

Electrophoretic mobility shift assays (EMSAs) were performed after incubating full-

length SgrS (+1 to +227) with its target transcripts (A) *ptsG* (+1 to +240), *manX* (+1 to

+240), *purR* (+1 to +230), *yigL* (-191 to +50 relative to ATG translation start of *yigL*)*,*

asdI (+1 to +110)*, and *asdII* (+71 to +310). B-D) Target transcripts (B) *ptsG* (+1 to

+240), (C) *manX* (+1 to +240), (D) *asdI* (+1 to +110) were labeled with ^{32}P and

incubated with unlabeled SgrS. EMSAs were performed to resolve complex formation.

Band densities were measured for replicate experiments (n, top left) and plotted to

determine dissociation constant (K_D, bottom right) values for (B) *ptsG*, (C) *manX*, and

(D) *asdI*. E) EMSA of radiolabeled SgrS in the presence of increasing concentrations of

asdI-II transcript. Shift in mobility corresponding to one or two SgrS bound to *asdI-II* is

denoted as Site I-SgrS* and Sites I-II-SgrS* respectively. F) Quantification of SgrS

binding with radiolabeled *asdI-II* (+1 to +240), as described above.

Figure 5. Secondary structure of 5' end of *asd*. A) Diagram showing base-pairing

interactions of SgrS with binding sites I and II of *asd* mRNA. B) Energy of interaction

predicted by IntaRNA (26). "Structured" indicates pairing between full length SgrS (+1 to

+227) and *asdI-II* (+1 to +180)*. Plotted is the energy of interactions at either site I (*asdI*)

or site II (*asdII*). "Isolated" indicates interactions between isolated binding sites: SgrS

(+158 to +176) with *asdI* (+31 to +49) and SgrS (+158 to +178) pairing with *asdII* (+110

to +129). C) The structure of the *asdI-II* RNA alone or in complex with SgrS was probed

with NMIA and the modified RNA was analyzed by primer extension inhibition. SHAPE

reactivity (difference between the frequency of primer extension products at each

nucleotide in +NMIA vs. -NMIA samples) was then used as a parameter in the Vienna

RNAprobing WebServer (47) to predict the secondary structure of the *asdI-II* RNA.

Colors indicate SHAPE reactivity as follows: red, highly reactive (≥ 0.8); gold, reactive

(0.4-0.79); green, moderately reactive (0.2-0.39); blue, minimally reactive (0.1-0.19);

grey, unreactive (< 0.01). Distinct structures were observed in the absence of SgrS and

in the presence of saturating concentrations of SgrS. The SHAPE reactivity of *asdI-II*

RNA alone (left) or in the presence of 5-fold excess SgrS (right) is mapped to the

predicted secondary structures. (D) SHAPE reactivity as a function of SgrS

concentration for each binding site (top, site I; bottom, site II). Only nucleotides with a

significant (≥ 0.1) change in reactivity are shown. Error bars denote SEM, n = 9. (E)

Relative SHAPE reactivity (difference in the SHAPE reactivity in the presence of SgrS

vs. the absence of SgrS) of the *asdI-II* RNA in the presence of wild-type (top) or mutant

(bottom) SgrS. Error bars denote SEM, n = 9 (WT), 6 (MT). The *asdI-II* RNA

nucleotides are numbered below the X-axis and the SgrS binding sites are indicated.

Figure 6. STORM imaging of SgrS regulation of *asd* variants. A) Illustration of *asdI-*

II, *asdI* and *asdII* translationally fused to *lacZ* reporter with SgrS binding sites I and II

marked. B-D) 2D projection of 3D super-resolution images of SgrS and *lacZ* mRNA for

the different *asd-lacZ* variants, labeled by smFISH, before and after 10 min induction

with 1% αMG. (B), (C) and (D) correspond to *asdI-lacZ*, *asdII-lacZ,* and *asdI-II-*

lacZ shown in (A). Probability distributions of RNA copy numbers in individual cells for

30-250 cells are plotted next to the representative images.

Figure 7. SgrS binding cooperativity allows for improved repression of *asd*

translation. A) Illustration of *asdI* and *asdI-II* constructs with SgrS binding sites marked.

Graphs show comparison of SgrS regulation of *asdI* and *asdI-II* variants in (B) wild-type

and (D) *rne701* mutant by plotting regulated activity over basal activity at various SgrS

expression levels (20-100 ng/ml aTc). Regulatory hierarchy of SgrS targets in (C) wild-

type and (E) *rne701* mutant strains. Regulation of target genes at one SgrS expression

level (40 ng/ml aTc) are compared by plotting regulated activity over basal activity of the

gfp reporter.

Figure 8. SgrS regulation of transcriptional *asd-lacZ* fusions. β-galactosidase

activity of (A) *asdI-lacZ* (+1 to +64) and *asdI-II-lacZ* (+1 to +277) was (B) assayed in

response to SgrS expression from a plasmid (and vector control) in WT and *rne701*

background strains.

Figure 1. Model for SgrS target prioritization during glucose-phosphate stress.
Glucose or the analogs αMG and 2-deoxyglucose are phosphorylated during transport through the phosphotransferase system proteins EIICB^{Glc} (PtsG) or EIICD^{Man} (ManYZ). If sugar-phosphates are not metabolized, the glucose-phosphate stress response is triggered, and the transcription factor SgrR becomes active and promotes *sgrS* transcription. The RNA chaperone Hfq promotes SgrS-mediated translational repression of *ptsG* and *manXYZ* mRNAs, reducing synthesis of sugar transporters. SgrS stabilizes *yigL* mRNA, promoting sugar phosphatase (YigL) synthesis. SgrS-mediated repression of *asd, purR, folE* and *adiY* likely reroutes metabolism to restore homeostasis during stress recovery. The hypothetical sequence of regulatory events following stress induction is represented from left to right as SgrS levels increase over time. When SgrS concentrations are low, only the highest priority targets are regulated. When stress persists and concentrations of SgrS increase, lower priority targets are regulated.

Figure 2. Efficiency of target regulation by SgrS. A) Representation of genetic constructs in two compatible plasmids used to study target regulation by SgrS. One plasmid contains full-length SgrS under the control of the aTc-inducible P_{Tet} promoter. A second plasmid contains a P_{Lac} promoter and the relevant region encoding each SgrS target (including the SgrS binding site) translationally fused to a superfolder *gfp* reporter gene. B-F) Regulated activity was plotted as a function of basal activity (see text for description) for (B) *ptsG*, (C) *manX*, (D) *purR,* (E) *asdI,* and (F) *yigL* fused to *sfgfp* reporter gene. Without SgrS-mediated regulation, we obtained a line with a slope =1. The plots with slopes <1 indicate repression of (B) *ptsG*, (C) *manX*, (D) *purR* and (E) *asdI* by SgrS. The plot with a slope of >1 is indicative of activation of (F) *yigL*.

Figure 3. Regulatory hierarchy established by SgrS. Regulated activity was plotted as a function of basal activity for *ptsG*, *manX*, *purR, asdI,* and *yigL* fusions. Lack of SgrS regulation is indicated by a line with a slope =1. The plots with slopes <1, indicate repression (*ptsG*, *manX*, *purR* and *asdI*) by SgrS. The plot with slope >1 indicates activation (*yigL*)*.* Target fusion activity was monitored at different levels of SgrS induction by aTc: (A) 10 ng/ml, (B) 20 ng/ml, (C) 30 ng/ml, (D) 40 ng/ml, (E) 50 ng/ml.

A

Target [µM] 0 ———————— 16   0 ———————— 16

ptsG*-SgrS

SgrS*

manX*-SgrS

SgrS*

purR*-SgrS

SgrS*

SgrS*-***yigL***

SgrS*

SgrS*-***asdI***

SgrS*

SgrS*-***asdII***
(expected)

SgrS*

B   C   D

ptsG   **manX**   **asdI**

n=3   n=3   n=2

Fraction Bound

Kd=0.11±0.01   Kd=19.7±2.78   Kd=0.15±0.04

SgrS [µM]

E   F

Target [µM] 0 ———————— 16   **asdI-II**

sites I-II*-SgrS   n=3

site I*-SgrS

Fraction bound

SgrS*

Kd=0.07±0.01

[SgrS] µM

Figure 4 legend on the next page

Figure 4. SgrS binding to target mRNAs *in vitro*. A) SgrS was labeled with ^{32}P and incubated with unlabeled target transcripts at final concentrations of 0 µM - 16 µM . Electrophoretic mobility shift assays (EMSAs) were performed after incubating full-length SgrS (+1 to +227) with its target transcripts (A) *ptsG* (+1 to +240), *manX* (+1 to +240), *purR* (+1 to +230), *yigL* (-191 to +50 relative to ATG translation start of *yigL*), *asdI* (+1 to +110), and *asdII* (+71 to +310). B-D) Target transcripts (B) *ptsG* (+1 to +240), (C) *manX* (+1 to +240), (D) *asdI* (+1 to +110) were labeled with ^{32}P and incubated with unlabeled SgrS. EMSAs were performed to resolve complex formation. Band densities were measured for replicate experiments (n, top left) and plotted to determine dissociation constant (K_D, bottom right) values for (B) *ptsG*, (C) *manX*, and (D) *asdI*. E) EMSA of radiolabeled SgrS in the presence of increasing concentrations of *asdI-II* transcript. Shift in mobility corresponding to one or two SgrS bound to *asdI-II* is denoted as Site I-SgrS* and Sites I-II-SgrS* respectively. F) Quantification of SgrS binding with radiolabeled *asdI-II* (+1 to +240), as described above.

Figure 5 legend on the next page

Figure 5. Secondary structure of 5' end of *asd*. A) Diagram showing base-pairing interactions of SgrS with binding sites I and II of *asd* mRNA. B) Energy of interaction predicted by IntaRNA (26). "Structured" indicates pairing between full length SgrS (+1 to +227) and *asdI-II* (+1 to +180). Plotted is the energy of interactions at either site I (*asdI*) or site II (*asdII*). "Isolated" indicates interactions between isolated binding sites: SgrS (+158 to +176) with *asdI* (+31 to +49) and SgrS (+158 to +178) pairing with *asdII* (+110 to +129). C) The structure of the *asdI-II* RNA alone or in complex with SgrS was probed with NMIA and the modified RNA was analyzed by primer extension inhibition. SHAPE reactivity (difference between the frequency of primer extension products at each nucleotide in +NMIA vs. -NMIA samples) was then used as a parameter in the Vienna RNAprobing WebServer (55) to predict the secondary structure of the *asdI-II* RNA. Colors indicate SHAPE reactivity as following: red, highly reactive (≥ 0.8); gold, reactive (0.4-0.79); green, moderately reactive (0.2-0.39); blue, minimally reactive (0.1-0.19); grey, unreactive (< 0.01). Distinct structures were observed in the absence of SgrS and in the presence of saturating concentrations of SgrS. The SHAPE reactivity of *asdI-II* RNA alone (left) or in the presence of 5-fold excess SgrS (right) is mapped to the predicted secondary structures. (D) SHAPE reactivity as a function of SgrS concentration for each binding site (top, site I; bottom, site II). Only nucleotides with a significant (≥ 0.1) change in reactivity are shown. Error bars denote SEM, n = 9. (E) Relative SHAPE reactivity (difference in the SHAPE reactivity in the presence of SgrS vs. the absence of SgrS) of the *asdI-II* RNA in the presence of wild-type (top) or mutant (bottom) SgrS. Error bars denote SEM, n = 9 (WT), 6 (MT). The *asdI-II* RNA nucleotides are numbered below the X-axis and the SgrS binding sites are indicated.

Figure 6. STORM imaging of SgrS regulation of *asd* variants. A) Illustration of *asdI-II*, *asdI* and *asdII* translationally fused to *lacZ* reporter with SgrS binding sites I and II marked. B-D) 2D projection of 3D super-resolution images of SgrS and *lacZ* mRNA for the different *asd-lacZ* variants, labeled by smFISH, before and after 10 min induction with 1% αMG. (B), (C) and (D) correspond to *asdI-lacZ*, *asdII-lacZ,* and *asdI-II-lacZ* shown in (A). Probability distributions of RNA copy numbers in individual cells for 30-250 cells are plotted next to the representative images.

Figure 7. SgrS binding cooperativity allows for improved repression of *asd* translation. A) Illustration of *asdI* and *asdI-II* constructs with SgrS binding sites marked. Graphs show comparison of SgrS regulation of *asdI* and *asdI-II* variants in (B) wild-type and (D) *rne701* mutant by plotting regulated activity over basal activity at various SgrS expression levels (20-100 ng/ml aTc). Regulatory hierarchy of SgrS targets in (C) wild-type and (E) *rne701* mutant strains. Regulation of target genes at one SgrS expression level (40 ng/ml aTc) are compared by plotting regulated activity over basal activity of the *gfp* reporter.

Figure 8. SgrS regulation of transcriptional *asd-lacZ* fusions. β-galactosidase activity of (A) *asdI-lacZ* (+1 to +64) and *asdI-II-lacZ* (+1 to +277) was (B) assayed in response to SgrS expression from a plasmid (and vector control) in WT and *rne701* background strains.