SELECTION FOR POLLEN COMPETITIVE ABILITY IN MIXED MATING SYSTEMS

ABSTRACT

Co-expression of genes in plant sporophytes and gametophytes allows correlated gametic and

sporophytic selection. Theory predicts that, under outcrossing, an allele conferring an advantage

to pollen competitive ability should rapidly fix within a population unless antagonistic pleiotropy

with the sporophyte stage is strong. However, under strong selfing, pollen competitiveness is

immaterial because superior and inferior competitors are deposited on opposite stigmas,

producing assortative competition. Because many plant species have mixed-mating systems,

selfing should be critical in the spread and maintenance of pollen-expressed genes affecting

competitiveness. We present two one-locus, two-allele population genetic models for the

evolution of a locus controlling pleiotropic antagonism between pollen competitiveness and

diploid fitness. Analytical solutions provide minimum and maximum selfing rates allowing

invasion of alleles with greater fitness in diploid and haploid stages respectively. Further,

polymorphism is only maintained when diploid selection is recessive. Numerically, we show

fixation of the allele conferring greater pollen competitiveness may be prevented, even with

weak counter-selection in the sporophyte stage, when selfing is sufficiently high. Finally, selfing

expands and limits the range of haploid-diploid selection coefficients that allow for

polymorphism, depending on dominance and mode of self-pollination.

Key Words: Assortative competition, gametic selection, sporophytic selection, population

genetics, mixed mating, genetic correlation

 INTRODUCTION

More than 7000 genes are expressed in male gametophytes—pollen grains, pollen tubes and the

sperm they transmit—as they compete in the pistil for access to ovules (Page and Grossniklaus

2002; Honys and Twell 2003; Rutley and Twell 2015). As many as 60% of all known plant

genes are expressed in both the gametophyte and sporophyte (Borges et al. 2009; Arunkumar et

al. 2013). For instance, a gene regulating cell polarity and elongation, *PIP5K4*, is expressed in

both pollen tubes and root hairs (Sousa et al. 2008). Over 100 cases of co-expression in pollen

tubes and root hairs have been identified (Hadif et al. 2012). This high degree of co-expression

presents ample opportunity for selection acting on pollen to affect sporophyte evolution, and *vice*

versa. Haldane (1932) was among the first to recognize this potential, writing "Clearly a higher

plant species is at the mercy of its pollen grains. A gene which greatly accelerates pollen tube

growth will spread through a species even if it causes moderately disadvantageous changes in the

adult plant." Put simply, genes conferring a competitive advantage in the gametophyte fix unless

there is strong antagonistic pleiotropy of these genes with sporophyte fitness.

Selection should be strong on alleles with concordant impacts on the gametophyte and

sporophyte stages. Deleterious recessive mutations, which are masked in the heterozygous

sporophyte, are fully exposed to selection in the haploid gametophyte. This facilitates purging,

thereby diminishing inbreeding depression (Charlesworth and Charlesworth 1992). Some have

suggested that alleles promoting pollen competition will increase offspring quality; large

stigmatic pollen loads lead to intense competition in the style, favouring the genetically superior

gametophytes, which then contribute their superior genes to the seed (Mulcahy and Mulcahy

1987; Winsor et al. 1987; Winsor et al. 2000; but see Baskin and Baskin 2015; Pélabon et al.

2016). Walsh and Charlesworth (1992) argue that strong selection on alleles with concordant

effects on the gametophyte and sporophyte will soon deplete genetic variation in pollen

competitive ability. But, as Haldane suggested, polymorphism may be maintained if an

advantage in pollen competitive ability is counterbalanced by fitness costs to the sporophyte.

Immler, Arnqvist and Otto (2012) explored the maintenance of genetic variation under

discordant selection between haploid and diploid phases, as well as under sex-specific selection.

Considering an autosomal gene, they demonstrated polymorphism is most easily maintained

under negative ploidy-by-sex interactions, under which sex-specific selection acts in opposite

directions in haploids and diploids, i.e., is ploidally antagonistic. More generally, they conclude

that polymorphism can be maintained, even in the absence of sex differences in selection, when

selection is opposing between sporophytic and gametophytic stages.

High rates of self-pollination diminish the advantage of pollen competitive ability (Mazer

et al. 2010), which may in turn restrict the conditions for stable polymorphism. This is due to

assortative competition under self-pollination. Imagine a one locus system, with allele C_g

conferring a competitive advantage to pollen (i.e., in the gametophytic phase) and the alternate

allele C_s advantages the sporophytic phase. When a $C_g C_g$ homozygote self-pollinates, the C_g

allele gets passed to the offspring by necessity and not due to its competitive superiority. On the

opposite homozygote, the C_s allele is likewise successful by necessity, despite its competitive

inferiority. The competitive advantage of C_g- over C_s-bearing pollen is realized when

heterozygotes self-pollinate, but because selfing reduces heterozygote frequency, over time there

are fewer and fewer opportunities for C_g to exercise its superiority. Thus, with obligate selfing,

the competitively inferior C_s allele gains refuge from competition, and since it confers a fitness

benefit in the sporophyte, this allele quickly goes to fixation. Indeed, a comparison between two

populations of *Clarkia tembloriensis* (Onagraceae) revealed slower pollen tube growth rates in

the highly selfing population than in the highly outcrossing one (Smith-Huerta 1996), consistent

with predictions. Additional examples are reviewed by Mazer et al. (2010).

While a little over half of plant species are self-incompatible, and thus obligate

outcrossers (Igic and Kohn 2006), about 40% of species have mixed mating systems—

individuals typically receive mixtures of self and outcross pollen (Schemske and Lande 1985;

Goodwillie et al. 2005). Of the 155 studies of mixed mating systems in plants reviewed by

Barrett and Eckert (2012), two-thirds report selfing rates in excess of 20%. How do intermediate

selfing rates influence the fate of a new mutation increasing pollen competitiveness at the

expense of sporophyte?

An important paper by Jordan and Connallon (2014) considered the impact of self-

fertilization on sexual conflict in hermaphrodites. They found that selfing narrows the conditions

for invasion of new alleles conferring male-benefit in the haploid phase if they impose a cost on

female function during the diploid phase. Their model considered the particular case in which

gamete quality is determined by the genotype of the adult producing it. In this situation a C_g-

bearing gamete from a homozygous adult differs from a C_g-bearing gamete produced by a

heterozygote, weakening the correspondence of gamete phenotype to haplotype. Gene expression

in animal sperm is limited (Parker and Begon 1993), and so it is reasonable to assume genetic

paternal effects on sperm quality for these species. In plants, adult diploid condition can affect

pollen size and quality (Young and Stanton 1990; Mazer and Gorchov 1996; Galloway 2001;

Smith-Huerta et al. 2007). Pollen size, which could well be mediated by genetically influenced

resource allocation by the sporophyte, readily responds to artificial selection (Sarkissian and

Harder 2001; McCallum and Chang 2016). Here we explore a different and potentially important

situation, specifically, when pollen competitive ability is affected by genes expressed directly in

the male gametophyte. Here one may expect stronger selection responses because of the stronger

correspondence of haplotype to phenotype.

The paper employs a one-locus model in which allele C_g increases pollen completive

ability at the expense of sporophyte fitness. Analytical solutions show there are maximum and

minimum selfing rates allowing invasion of C_g and C_s alleles, respectively, and that stable

polymorphism is possible only when C_g is recessive. Numerical analyses show that as selfing

rate increases, C_g's competitive advantage is negated by increasingly smaller diploid cost.

99   METHODS

THE MODELS

The departure from Hardy-Weinberg equilibrium caused by selfing necessitates that selection be

modelled in terms of genotype frequencies rather than allele frequencies (Crow and Kimura

1970). We refer to the frequencies of the C_gC_g, C_gC_s and C_sC_s genotypes as X_0, X_1 and X_2,

respectively, where the subscript denotes the number of C_s alleles. Allele frequencies are p and q,

for C_g and C_s, respectively.

Assume a pleiotropic effect by which the C_g allele increases success in pollen

competition but decreases the number of pollen grains and ovules that the diploid plant

contributes to the gamete pool (either by reduced survival or lower flower production).

Thus, selection favours the C_s allele through both sex functions in the diploid stage, disfavours it

in haploid pollen, and has no effect on ovules. Let the fitness of sporophyte C_sC_s be 1, and that of

C_gC_g to be 1-s, while the heterozygote C_gC_s is 1-hs, where h is the degree of dominance for

fitness. Similarly, define the absolute competitive ability of pollen bearing the C_g allele as 1 and

that of pollen bearing the C_s allele as 1-t. Finally, selfing rate, designated by r, varies between 0

and 1. All individual plants are expected to have many flowers, and to have the same selfing rate.

Using these parameterisations, we modelled two forms of selfing: fixed selfing and mass-action

selfing, defined below.

Fixed selfing

Under fixed selfing, a stigma receives either all self-pollen or all outcross pollen. However, the

proportion of flowers on a given plant that receive self-pollen can vary from zero to 1. We can

produce recursion equations for genotype frequencies by considering diploid selection and

selfing separately. After selection among diploids, genotype frequencies are:

$$x_0 = \frac{(1-s)X_0}{\overline{W}} \tag{1a}$$

$$x_1 = \frac{(1-hs)X_1}{\overline{W}} \tag{1b}$$

$$x_2 = \frac{X_2}{\overline{W}} \tag{1c}$$

122   where $\overline{W}$ is the mean diploid fitness:

$$\overline{W} = (1-s)X_0 + (1-hs)X_1 + X_2. \tag{2}$$

123   Next, genotype frequencies following mating are

125    $$X_0' = r\left[x_0 + \frac{1}{2}x_1\left(\frac{1}{2-t}\right)\right] + (1-r)\left[\left(x_0 + \frac{1}{2}x_1\right)\alpha\right] \tag{3a}$$

       $$X_1' = r\left[\frac{1}{2}x_1\right] + (1-r)\left[x_0(1-\alpha) + \frac{1}{2}x_1 + x_2\alpha\right] \tag{3b}$$

128    $$X_2' = r\left[x_2 + \frac{1}{2}x_1\left(\frac{1-t}{2-t}\right)\right] + (1-r)\left[\left(x_2 + \frac{1}{2}x_1\right)(1-\alpha)\right] \tag{3c}$$

130   where α reflects the relative competitiveness of C_g-bearing pollen as a function of the absolute

131   competitiveness conferred by the C_g allele and the frequency of the C_g allele in the pollen pool as

132   a result of genotype frequency following selection:

$$\alpha = \frac{x_0 + \frac{1}{2}x_1}{x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)}. \tag{4}$$

It follows that the relative competitiveness of C_s-bearing pollen is (1- α).

To understand Equations (3a-c), consider separately the offspring produced via selfing (*r*)

and those produced via outcrossing (1-*r*). Under selfing, competition among pollen tubes

carrying opposite alleles only occurs within heterozygote stigmas and styles. Assuming

heterozygotes produce C_g and C_s pollen in equal numbers, ½ of their offspring will likewise be

heterozygous, with the remaining offspring divided between C_gC_g and C_sC_s homozygotes in

proportions of $\left(\frac{1}{2-t}\right)$ and $\left(\frac{1-t}{2-t}\right)$ respectively. We assume that each selfing flower produces enough

pollen to fertilise its available ovules, such that only female diploid fitness terms are included.

Conversely, offspring produced through outcrossing will be a function of female diploid fitness

as well as male diploid fitness and the pollen competitive ability conferred by each allele, as

specified by α. For sake of simplicity, we make no provision for inbreeding depression;

increasing failure rate selfed offspring has the same effect as reducing the basic selfing rate (see

Jordan and Connallon 2014).

Mass-action selfing

Under mass-action selfing (Holsinger 1991), all individual stigmas can receive both self- and

outcross-pollen, resulting in direct competition for ovules. In such a case, the amount of self-

pollen received by a stigma is a function of the following: the proportion of the focal flower's

pollen that is dedicated to selfing (i.e., that is deposited on its own stigma), *r*; the quantity of

pollen produced by the individual relative to that available through outcrossing; and, the relative

competitive ability of the self-pollen.

Assuming diploid selection occurs according to Equations (1a-c), we arrive at the

following genotype frequencies after mating:

$$X_0' = x_0 \left[\frac{rW_0 + (1-r)(x_0 + \frac{1}{2}x_1)}{rW_0 + (1-r)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right)} \right]$$

$$+ \frac{1}{2}x_1 \left[\frac{r\frac{1}{2}W_1 + (1-r)(x_0 + \frac{1}{2}x_1)}{r\frac{1}{2}W_1(2-t) + (1-r)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right)} \right] \tag{5a}$$

$$X_1' = \frac{1}{2}x_1 + x_0 \left[\frac{(1-r)(x_2 + \frac{1}{2}x_1)(1-t)}{rW_0 + (1-r)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right)} \right]$$

$$+ x_2 \left[\frac{(1-r)(x_0 + \frac{1}{2}x_1)}{r\frac{1}{2}W_1(1-t) + (1-r)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right)} \right] \tag{5b}$$

$$X_2' = x_2 \left[\frac{rW_2(1-t) + (1-r)(x_0 + \frac{1}{2}x_1)(1-t)}{rW_2(1-t) + (1-r)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right)} \right]$$

$$+ \frac{1}{2}x_1 \left[\frac{r\frac{1}{2}W_1(1-t) + (1-r)(x_2 + \frac{1}{2}x_1)(1-t)}{r\frac{1}{2}W_1(2-t) + (1-r)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right)} \right] \tag{5c}$$

According to Eq. 5a, the two terms represent C_gC_g offspring seed produced by C_gC_g and C_gC_s

plants, respectively, with post-selection frequencies of x_0 and x_1. All ovules on the homozygote

carry C_g, whereas on heterozygotes only half do. The terms inside the square brackets are the

probability that an ovule receives C_g-bearing sperm, which depends on the amount of successful

C_g pollen (numerator) divided by the total amount of pollen arriving at the stigma (denominator).

SOFTWARE AND COMPUTATION

Conditions for stable polymorphisms were explored analytically through invasion analyses using

Wolfram Mathematica 11.0.1.0. We derived leading eigenvalues for the Jacobian Matrix using

the genotype recursions (Eqs. 3a-c, 5a-c). We then determined the conditions that allowed

invasion at a boundary by solving for parameters that would produce a leading eigenvalue of an

absolute value greater than 1. Invasion analyses were performed for two boundaries: the X_0

boundary, at which the C_s allele appears in a population fixed for the C_g allele (i.e., $p = 1$), and

the X_2 boundary, at which the C_g allele appears in a population fixed for the C_s allele (i.e., $p = 0$).

All numerical results were produced using Visual Basic in Microsoft Excel 2011 Version

1610, from which we generated and recorded genotype and allele frequencies over thousands of

generations for combinations of selfing, diploid fitness and pollen competitiveness. Long-term

allele frequencies were taken from the 20,000th generation and verified by a change in allele

frequency, i.e., Δp, of less than 10^{-6}. Hereafter we will refer to these long-term allele frequencies

as equilibrium frequencies, as we assume that, with such significantly small changes in allele

frequencies, drift will dominate and these long-term frequencies are good approximations of true

equilibrium frequencies, where $\Delta p = 0$.

176   RESULTS

ANALYTICAL RESULTS

We explored the impact of selfing rate on the potential for both C_g and C_s to invade when rare.

Invasion analyses at both the X_0 and X_2 boundaries using Equations (3a-c) as well as Equations

(5a-c) in their full forms produces extremely complicated leading eigenvalues, limiting their

usefulness for predicting population dynamics and drawing clear biological conclusions. To

derive more meaningful interpretations, we assumed weak selection, with the diploid phase

selection coefficient being considerably smaller than the haploid (i.e., $0 < s \ll t \ll 1$).

Fixed selfing

Under fixed selfing, boundary selfing rates permitting invasion of the C_g and C_s alleles,

respectively, are as follows:

$$\text{For } C_g: \ r < 1 - \frac{s}{(1-2h)s+t} \tag{6a}$$

$$\text{For } C_s: \ r > 1 + \frac{s}{(1-2h)s-t} \tag{6b}$$

These analytical solutions are visualised in Figure (1a-c). When mating is random (i.e., $r = 0$),

and $s \ll t$, the C_g allele can always invade, regardless of the degree of dominance h. This is

because, without selfing, the C_s allele has no refuge from competition. Conversely, under

complete selfing ($r = 1$), each individual sporophyte is a separate mating pool. Heterozygotes are

quickly lost by segregation. With a loss of heterozygotes follows decreasing opportunity for

competition between C_g- and C_s-bearing pollen. Consequently, the allele conferring higher

relative sporophyte fitness in the homozygous state can increase when rare.

The analytical solutions indicate that polymorphism can be maintained in mixed mating

systems ($0 < r < 1$), but only if the deleterious effect of the C_g allele in the diploid phase is

recessive ($h < 0.5$). When completely recessive ($h = 0$), there is a range of selfing rates in which

both alleles can invade when rare, implying protected polymorphism. The C_g allele invades

because, when rare, it occurs exclusively in heterozygotes. As such, its negative effect on diploid

fitness is entirely masked from selection, and when selfing occurs at moderate levels, there are

enough opportunities to outcompete C_s-bearing pollen through greater fertilisation success.

Within that same range of selfing rates, the C_s allele can also increase when rare because, 1)

heterozygotes, where C_s occurs, have higher relative fitness than the C_gC_g homozygote, and 2)

selfing reduces exposure of C_s pollen to competition with the superior C_g pollen.

Additive fitness effects ($h = 0.5$) prohibit polymorphism (Figure 1b). Under this

condition, the critical r values for invasion reduce to $1 - \frac{s}{t}$ (Eqn. 6a-b) for both alleles. As h

increases, the relative fitness of heterozygotes decreases. It then follows that the initial advantage

enjoyed by both alleles when rare also diminishes. To invade, C_g requires low r values to win

enough contests to offset the lower fitness of heterozygotes. Conversely, C_s will require higher r

values to avoid competitive losses and produce enough high-fitness X_2 homozygotes to offset

lower fitness of heterozygotes.

When the disadvantage of the C_g allele in the diploid sporophyte is entirely dominant ($h =$

1), there is a range of selfing rates that prohibit either allele from invading when rare (Figure 1c).

For a given value of s, the maximum selfing rate allowing C_g invasion is lowered because its

adverse effect on sporophyte fitness is fully expressed by heterozygotes. On the other hand, the

critical r for C_s invasion increases when C_g is dominant. When rare, the positive diploid fitness

effects of the C_s allele are concealed in heterozygotes. Thus, only very high selfing rates, can

provide sufficient refuge from competition to allow C_s to spread.

Mass-action selfing

Considering mass-action selfing, the maximum and minimum r values allowing for invasion of

the C_g and C_s alleles can be solved analytically. However, the expressions are are complex

difficult to interpret biologically; these are present in Appendix A (online supplemental

material). As above, these solutions are visualised in Figure (1d-f).

The analytical solutions for mass-action selfing lead to the qualitatively the same

outcomes as fixed selfing, i.e., at any given value of dominance (h), the selfing rate (r)

prohibiting invasion by C_g decreases as cost increases (Figures 1a-c versus 1d-f).

The modest quantitative differences in outcome between the two selfing modes stem

from the different impact that diploid cost for competitive ability gets paid through male

function. Under mass-action selfing each stigma can receive a mixture of C_g- and C_s- bearing

pollen (except when $r = 1$, i.e., all individuals are entirely selfing). A given selfing rate between

0 and 1 thus affords less refuge for C_s from competition with C_g under the mass-action model,

compared to fixed selfing. However, this greater opportunity for C_g pollen to outcompete C_s is

partly offset through male diploid fitness differences. Under fixed selfing, male diploid fitness

(pollen quantity) is moot, since all flowers of each genotype are assumed to produce enough

pollen to fertilise all their ovules. Under mass-action selfing, in contrast, the quantity of pollen

produced by an individual affects its realized fertilisation success through self-pollen. This

limitation is similar to the relative fertilisation success of the C_g allele under fixed selfing, as

parameterised by α (Equation 4). The different impacts on male function shift the invasion

boundary upward unde mass action selfing. For example, when $h = 0$ (Figures 1a, 1d) and $s =$

0.01, the maximum selfing rate allowing the C_g allele to invade increases from $r = 0.5$ to $r = 0.6$.

Likewise, the minimum selfing rate allowing C_s to invade increases from $r = 0.75$ to $r = 0.8$.

NUMERICAL RESULTS FOR INVASION, FIXATION, AND PROTECTED POLYMORPHISM

The analytical results rest on the assumption that, when an allele is rare, its increase can

be modelled by considering only the frequencies of heterozygotes and the common homozygote,

while ignoring the frequency of the *very* rare homozygote. This assumption is strained at high

selfing rates because self-pollination by heterozygotes quickly increases the frequency of the rare

homozygote to a level that is no longer negligible, especially if selection during the diploid stage

is strong. To arrive at a tractable solution, we assumed that $s \ll t < 1$. Here we present numerical

results for the recursion equations (Equations 3a-c, 5a-c) to probe the potential for polymorphism

under broader conditions.

We start by running Equations (3a-c) with $t = 0.03$ and $s = 0.01$; these are the values that

are illustrated along the top edge of the panes in Figure 1. Figure 2 plots the equilibrium

frequency of C_g as a function of selfing rate, r. These numerical results agree with the analytical

solution when C_g is recessive ($h = 0$). Figure 2a shows that both alleles are maintained within the

range of selfing rates ($r = 0.48$ to 0.74) predicted in Figure (1a). The numerical results depart

slightly from the analytical solution when the two alleles have additive effects ($h = 0.5$). Rather

than an abrupt transition from fixation of C_g to fixation of C_s at a critical selfing rate (Figure 1b),

there is a narrow window permitting polymorphism ($r = 063 - 0.69$; Figure 2b). When the

negative impact of C_g on sporophyte fitness is dominant ($h = 1$), analytical and numerical results

agree that high selfing rates prevent invasion by the C_g allele when rare, while low rates prevent

C_s invasion. Over an intermediate range of selfing rates, neither allele can increase in frequency

when rare (Figure 2c).

The numerical results confirm that fixed and mass-action selfing modes have

qualitatively the same impact on invasion, fixation and polymorphism (Figure 1, 2). Again, for a

given value of h, the minimum and maximum selfing rates allowing for invasion of the C_g and C_s

alleles respectively increases under mass-action selfing (Figure 2d-f), as the C_g allele is allotted

more opportunities to outcompete the C_s allele. Additionally, the parameter space allowing both

polymorphism (Figure 2d) and unstable equilibrium (Figure 2f) contracts under mass action

selfing relative to fixed (Figures 2a, c).

Returning to Haldane's conjecture that plants are at the mercy of their pollen grains, how

strong must sporophyte selection against a pollen-competition allele be to prevent its fixation?

The fate of the C_g allele—fixation, elimination or stable polymorphism—depends on the strength

of selection during the two life stages, which we can consider as a ratio of diploid cost to haploid

benefit, i.e., s/t. At any given s/t ratio, increasing the rate of self-pollination will restrict

opportunities for C_g pollen to exercise their competitive superiority, muting the selective

advantage. Against this, at any given ratio, Mendelian dominance alters how often the cost of

competitive ability gets paid. If C_g is recessive ($h = 0$) heterozygotes pay no diploid cost. If

dominant ($h = 1$), it masks the C_s benefit and heterozygotes pay full cost; however, since selfing

decreases heterozygote frequency, the dominance issue becomes moot. These conditions produce

the general outcome that s/t ratios allowing C_g to invade, and to be maintained in stable

polymorphism, tend to decrease as both r and s increase (Figure 3).

281   DISCUSSION

Classical theory predicts that an allele with an advantage to pollen competitiveness will fix in a

population unless a trade-off with the sporophyte stage sufficiently counter-balances this haploid

advantage. Our model predicts that even moderate rates of selfing, can slow or prevent the

spread of a pollen-expressed allele that confers competitive superiority (Table 1). Self-

pollination leads to assortative competition among pollen grains, restricting opportunity for such

an allele to exercise its superiority. When homozygotes self, only one haplotype is deposited on

the stigma, so by necessity that haplotype cannot be out-competed by another. Alternate

haplotypes compete only on heterozygotes. In addition, selfing depresses heterozygote frequency

through the generations such that the proportion of the pollen pool engaged in competitive

contests falls in proportion to the selfing rate.

The dominance relationships between the competitive and non-competitive alleles in the

diploid stage also influences invasion success. When the fitness costs to the sporophyte of

competitive allele is recessive, it is shielded from negative selection during the diploid phase.

Full dominance fully exposes it to negative selection. Thus the conditions for invasion narrow

with increase in the dominance coefficient (Table 1, Figures 1 and 2).

Increased dominance also narrows the condition for stable polymorphism (Table 1).

When costs are recessive, heterozygotes gain success as fathers because 50% of their pollen is

competitively superior, and they pay no cost. This can lead to overall heterozygote superiority.

Under random mating, this over-dominance in total fitness can maintain stable polymorphism

(Figures 1-3). But selfing diminished the impact of over-dominance (Figure 3) by eroding

heterozygote frequency: heterozygosity lost to segregation is not replenished by matings between

opposite homozygotes, and so the competitive allele is eventually carried mostly by the

homozygote, who pay full cost. Dominance of the allele precludes stable polymorphism.

Although heterozygotes get half the paternal benefit, they but pay full cost (Table 1, Figure 3),

making over-dominance for total fitness unachievable.

These results are qualitatively similar to the case explored by Jordan and Connallon

(2014), where pollen competitive ability is under the control of the paternal genotype. They

considered sexually antagonistic selection in a one-locus model, whereby in the diploid, one

allele increases the number or quality of sperm produced at the expense of ova, while its

alternate promotes ova over sperm. They also found that selfing and dominance restricts the

potential for stable polymorphism and ultimately diminish selection through male function.

Our results extend these findings to the potentially important case where pollen

competitive ability is determined by gene expression in the gametophyte itself, and not by gene

products and transcripts carried over during meiosis. With gametophyte gene expression pollen

phenotype (competitive ability) is more strongly correlated with haplotype. Specifically,

heterozygotes always produce a 50:50 mix of phenotypically and genetically 'strong' and 'weak'

pollen competitors. Under paternal control, in contrast, heterozygotes can produce

phenotypically all strong ($h = 0$), all weak ($h = 1$), or all intermediate ($h = 0.5$) competitors, even

though they always produce a 50:50 genetic mix. In the early stages of invasion, when pollen

carrying the competitive allele is produced only by heterozygotes, it will always confer

superiority over the alternate allele when expressed by the gametophyte. This clear genetic signal

thus expands conditions for a costly allele to invade and spread, and a somewhat stronger selfing

rate is required to extinguish the advantage.

The models indicate that under outcrossing, where opportunities to express selective superiority

are maximized, selection on the gametophyte can overcome greater sporophyte costs than under

selfing. Genomic evidence on loci expressed exclusively in pollen, where diploid costs are

necessarily zero, bear out relaxed selection under selfing. In the self-incompatible *Capsella*

grandiflora, Arukumar et al. (2011) found evidence for stronger purifying selection at sites

expressed solely in the pollen tubes than those expressed exclusively in seedling. Looking at the

highly selfing *Arabidopsis thaliana*, on the other hand, Harrison et al. (2015) detected weaker

purifying selection in pollen-exclusive versus sporophyte-exclusive genes, as evidenced by a

higher frequency of premature stop codons in the former. Greater scrutiny of loci expressed

during both stages are needed to clarify the general importance of negative pleiotropic

constraints on the evolution of gametophyte and sporophyte traits.

REFERENCES

Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013. Pollen-specific, but not

sperm-specific, genes show stronger purifying selection and higher rates of positive

selection than sporophytic genes in *Capsella grandiflora*. Mol. Biol. Evol. 30:2475-2486.

Barrett, S. C. and C. G. Eckert. 2012. 14 Variation and Evolution of Mating Systems in Seed

Plants. Biological approaches and evolutionary trends in plants :229.

Barrett, S. C. H. 1998. The evolution of mating strategies in flowering plants. Trends Plant Sci.

3:335-341.

Baskin, J. M. and C. C. Baskin. 2015. Pollen (microgametophyte) competition: an assessment of

its significance in the evolution of flowering plant diversity, with particular reference to

seed germination. Seed Sci. Res. 25:1-11.

Borg, M., L. Brownfield, and D. Twell. 2009. Male gametophyte development: a molecular

perspective. J. Exp. Bot. 60:1465-1478.

Campbell, S. A. 2015. Ecological mechanisms for the coevolution of mating systems and

defence. New Phytol. 205:1047-1053.

Charlesworth, D. and B. Charlesworth. 1992. The effects of selection in the gametophyte stage

on mutational load. Evolution 46:703-720.

Crow, J. F. and M. Kimura. 1970. Introduction to Population Genetics Theory. Harper & Row,

New York.

Damgaard, C. and V. Loeschcke. 1994. Genetic-variation for selfing rate the dependence of

selfing rate on mating history in *Brassica napus* (rape seed). Heredity 72:570-573.

Galloway, L. F. 2001. The effect of maternal and paternal environments on seed characters in the

herbaceous plant *Campanula americana* (Campanulaceae). Am. J. Bot. 88:832-840.

Goodwillie, C., S. Kalisz, and C. G. Eckert. 2005. The evolutionary enigma of mixed mating

systems in plants: occurrence, theoretical explanations, and empirical evidence. Annu. Rev.

Ecol. Evol. Syst. 36:47-79.

Hafidh, S., K. Breznenová, P. Růžička, J. Feciková, V. Čapková, and D. Honys. 2012.

Comprehensive analysis of tobacco pollen transcriptome unveils common pathways in

polar cell expansion and underlying heterochronic shift during spermatogenesis. BMC

Plant Biol. 12(1): 24.

Haldane, J. B. S. 1932. The Causes of Evolution. Reprinted 1990, Princeton University Press.

Harrison, M. C., E.B. Mallon, D. Twell, and R.L. Hammond. 2015. Deleterious mutation

accumulation in *Arabidopsis thaliana* pollen genes: a role for a recent relaxation of

selection. bioRxiv, 016626.

Holsinger, K. E. 1991. Mass-action models of plant mating systems: the evolutionary stability of

mixed mating systems. Am. Nat. 138:606-622.

Igic, B. and J. R. Kohn. 2006. Bias in the studies of outcrossing rate distributions. Evolution 60

(5): 1098–1103.

Immler, S., G. Arnqvist, and S. P. Otto. 2012. Ploidally antagonistic selection maintains stable

genetic polymorphism. Evolution 66:55-65.

Jarne, P. and D. Charlesworth. 1993. The evolution of the selfing rate in functionally

hermaphrodite plants and animals. Annu. Rev. Ecol. Evol. Syst. 24:441-466.

Jordan, C. Y. and T. Connallon. 2014. Sexually antagonistic polymorphism in simultaneous

hermaphrodites. Evolution 68:3555-3569.

Mazer, S. J., A. A. Hove, B. S. Miller, and M. Barbet-Massin. 2010. The joint evolution of

mating system and pollen performance: Predictions regarding male gametophytic evolution

in selfers vs. outcrossers. Perspect. Plant Ecol. Evol. Syst. 12:31-41.

McCallum, B. and S. M. Chang. 2016. Pollen competition in style: Effects of pollen size on

siring success in the hermaphroditic common morning glory, *Ipomoea purpurea*. Am. J.

Bot. 103(3):460-470.

Mulcahy, D. L. and G. B. Mulcahy. 1987. The effects of pollen competition. Am. Sci. 75:44-50.

Parker, G. and M. Begon. 1993. Sperm competition games: sperm size and number under

gametic control. Proc. R. Soc. London, Ser. B. 253:255-262.

Pélabon, C., L. Hennet, G. H. Bolstad, E. Albertsen, Ø. H. Opedal, R. K. Ekrem, and W. S.

Armbruster. 2016. Does stronger pollen competition improve offspring fitness when pollen

load does not vary? Am. J. Bot. 103:522-531.

Sarkissian, T. S. and L.D. Harder. 2001. Direct and indirect responses to selection on pollen size

in *Brassica rapa* L. J. Evol. Biol. 14(3):456-468

Schemske, D. W. and R. Lande. 1985. The evolution of self-fertilization and inbreeding

depression in plants. II. Empirical observations. Evolution 39:41-52.

Smith-Huerta, N. L. 1996. Pollen germination and tube growth in selfing and outcrossing

populations of *Clarkia tembloriensis* (Onagraceae). Int. J. Plant Sci. 157:228-233.

Sousa, E., B. Kost, and R. Malhó. (2008). *Arabidopsis* phosphatidylinositol-4-monophosphate 5-

kinase 4 regulates pollen tube growth and polarity by modulating membrane recycling.

Plant Cell, 20(11), 3050-3064.

Walsh, N. and D. Charlesworth. 1992. Evolutionary interpretations of differences in pollen tube

growth rates. Q. Rev. Biol. 67:19-37.

Winsor, J., L. Davis, and A. Stephenson. 1987. The relationship between pollen load and fruit

maturation and the effect of pollen load on offspring vigor in *Cucurbita pepo*. Am. Nat.

129:643-656.

Winsor, J. A., S. Peretz, and A. G. Stephenson. 2000. Pollen competition in a natural population

of *Cucurbita foetidissima* (Cucurbitaceae). Am. J. Bot. 87:527-532.

Young, H. J. and M. L. Stanton. 1990. Influence of environmental quality on pollen competitive

ability in wild radish. Science 248:1631-1633.

Tables

413   TABLE 1.   Summary of model outcomes for mutation to an allele that increases pollen (haploid

414   gametophyte) competitive ability, but reduces survivorship or fertility in the (diploid)

415   sporophyte.

	MATING SYSTEM		
EXPRESSION OF C_g **IN DIPLOID SPOROPHYTE**	**Obligate Outcrossing ($r = 0$)** *Ample opportunities* to exercise pollen competitive superiority	**Mixed ($0 < r <1$)** *Fewer opportunities* to exercise pollen competitive superiority	**Obligate Selfing ($r = 1$)** *No opportunities* to exercise pollen competitive superiority
Recessive Fitness cost paid only by homozygote	C_g goes to fixation if sporophyte cost is less than ½ the pollen competitive benefit, and stable polymorphism unless costs exceed ~10×benefit—*conditions for stable polymorphism broad*	C_g goes to fixation only if sporophyte cost is *substantially* less than ½ the competitive benefit; upper limit for cost to permit stable polymorphism is reduced even more—*conditions for stable polymorphism narrowed*	C_g unable to invade.
Dominant Fitness cost paid *both* by homozygote and heterozygote	C_g goes to fixation if sporophyte cost is less than ½ the pollen competitive benefit, but is eliminated if costs are higher— *no stable polymorphism.*	C_g goes to fixation only if sporophyte cost is *substantially* less than ½ the benefit, but *eliminated* if costs are any higher—*no stable polymorphism.*	C_g unable to invade.

Figure legends

Figure 1

Conditions producing fixation, polymorphism and unstable equilibrium (i.e., fixation is

dependent on initial frequencies) of two alleles C_g and C_s at an autosomal gene under

antagonistic pleiotropy between pollen competitive ability and diploid fitness. All results are for t

= 0.03. (a-c) Analytical results for the fixed-selfing model. Potential for polymorphism

diminishes as the dominance of the C_g allele increases, eventually being replaced by unstable

equilibrium when $h > 0.5$. As selfing rate increases, polymorphism requires greater diploid

selection (*s)* against the C_g allele to offset its greater pollen competitive ability (*t*). (d-f)

Analytical results for the mass-action model. Conditions allowing polymorphism versus unstable

equilibrium are similar to those of the fixed-selfing model, though markedly narrower, as

individual stigmas never receive entirely self-pollen except when $r = 1$, thus increasing

opportunity for competition between C_g- and C_s-bearing pollen.

Figure 2

Numerical results of equilibrium frequency of the C_g allele when $s = 0.01$ and $t = 0.03$ across

selfing rates $0 < r < 1$. Solid lines indicate equilibrium frequencies starting from the X_0 boundary,

i.e., when the C_s allele invades a population fixed for C_g, while dashed lines indicted the opposite

boundary. (a-c) Under the fixed-selfing model, the window of selfing rates contracts as

dominance increases. (d-f) Under the mass-action model, C_s pollen is never entirely protected

from competition with C_g pollen as individual stigmas receive a mixture of both self and outcross

pollen for intermediate selfing rates. Thus, maximum and minimum selfing rates allowing

fixation or maintenance of C_g and C_s alleles respectively are right-shifted compared to under

fixed selfing, reflecting the greater opportunity for the C_g allele to exercise its advantage in

pollen competition with the C_s allele.

Figure *3*

Numerical results for equilibrium frequency of the *Cg* allele across a range of *s/t*, i.e., ratios of

diploid and haploid fitness values. Series specify selfing rates *r* ranging between 0 and 1. Under

both (a-c) fixed and (d-f) mass-action selfing models, increases in dominance of the *Cg* allele

sharply decrease the maintenance of polymorphism, particularly at lower selfing rates. Noticably,

for a given combination of selfing rate *r* and dominance *h,* the *Cg* allele persists at higher ratios

of *s* and *t* under mass-action selfing relative to fixed-selfing , reflecting the greater opportunity

for *Cg*-bearing pollen to compete *Cs*-bearing pollen under mass-action selfing.

Appendix A

Conditions for stable polymorphisms were explored analytically through invasion analyses. We

derived leading eigenvalues for the Jacobian Matrix using the genotype recursions (Eqs. 3a-c,

5a-c) in the main text. We then determined the conditions that allowed invasion at a boundary by

solving for parameters that would produce a leading eigenvalue of an absolute value greater than

1. Invasion analyses were performed for two boundaries: the X_0 boundary, at which the C_s allele

appears in a population fixed for the C_g allele (i.e., $p = 1$), and the X_2 boundary, at which the C_g

allele appears in a population fixed for the C_s allele (i.e., $p = 0$).

The analytical solutions specifying the selfing rates required for invasion C_g and C_s

alleles are provided below:

For C_g: $r <$ (A1)

$$-\frac{-3s+4hs+t}{3s-6hs}+(2^{1/3}\,((9-30h+28h^2)\,s^2+(-9+14h)\,st+t^2))/(3\,(-1+2h)\,s\,((54-270h+396h^2-160h^3)\,s^3-3\,(-9+12h+4h^2)\,s^2\,t+3\,(-9+14h)\,s\,t^2+2\,t^3+3\,\sqrt{3}$$

$$\sqrt{-(s-2hs)^2\,(576h^4\,s^4-32h^3\,s^3\,(27s-37t)+12hst\,(135s^2-100st+17t^2)+4h^2\,s^2\,(81s^2-558st+205t^2)+t\,(-432s^3+441s^2\,t-134st^2+9t^3))}\,)^{1/3})+\frac{1}{3\times2^{1/3}\,(-1+2h)\,s}((54-270h+396h^2-160h^3)$$

$$s^3-3\,(-9+12h+4h^2)\,s^2\,t+3\,(-9+14h)\,s\,t^2+2\,t^3+3\,\sqrt{3}\,\sqrt{-(s-2hs)^2\,(576h^4\,s^4-32h^3\,s^3\,(27s-37t)+12hst\,(135s^2-100st+17t^2)+4h^2\,s^2\,(81s^2-558st+205t^2)+t\,(-432s^3+441s^2\,t-134st^2+9t^3))}\,)^{1/3}$$

For C_s: $r >$ (A2)

$$\frac{1}{6}\,(\frac{6\,(s-4hs+t)}{3s-6hs}+(2\,2^{1/3}\,((7-26h+28h^2)\,s^2+(5-14h)\,st+t^2))/((-1+2h)\,s\,((-20+42h+84h^2-160h^3)\,s^3+3\,(7-20h+4h^2)\,s^2\,t+3\,(-5+14h)\,s\,t^2-2\,t^3+3\,\sqrt{3}$$

$$\sqrt{-(s-2hs)^2\,(36\,(1-5h+4h^2)^2\,s^4-4\,(-35+177h-330h^2+296h^3)\,s^3\,t+(61-440h+820h^2)\,s^2\,t^2+2\,(35-102h)\,st^3+9t^4)}\,)^{1/3})+\frac{1}{(-1+2h)\,s}2^{2/3}\,((-20+42h+84h^2-160h^3)\,s^3+3$$

$$(7-20h+4h^2)\,s^2\,t+3\,(-5+14h)\,s\,t^2-2\,t^3+3\,\sqrt{3}\,\sqrt{-(s-2hs)^2\,(36\,(1-5h+4h^2)^2\,s^4-4\,(-35+177h-330h^2+296h^3)\,s^3\,t+(61-440h+820h^2)\,s^2\,t^2+2\,(35-102h)\,st^3+9t^4)}\,)^{1/3})$$

Figures

466   Figure 1

468   Figure 2

470   Figure 3