SELECTION FOR POLLEN COMPETITIVE ABILITY IN MIXED MATING SYSTEMS

ABSTRACT

Co-expression of genes in plant sporophytes and gametophytes allows correlated gametic and sporophytic selection. Theory predicts that, under outcrossing, an allele conferring greater pollen competitive ability should fix within a population unless antagonistic pleiotropy with the sporophyte stage is strong. However, under strong selfing, pollen competitiveness is immaterial as superior and inferior competitors are deposited on opposite stigmas, producing assortative competition. Because many plant species have mixed-mating systems, selfing should be critical in the spread and maintenance of pollen-expressed genes affecting competitiveness. We present two one-locus, two-allele population genetic models for the evolution of a locus controlling pleiotropic antagonism between pollen competitiveness and diploid fitness. Analytical solutions provide minimum and maximum selfing rates allowing invasion of alleles with greater diploid and haploid fitness respectively. Further, polymorphism is only maintained when diploid selection is recessive. Fixation of the allele conferring greater pollen competitiveness may be prevented, even with weak sporophytic counter-selection, with sufficiently high selfing. Finally, selfing expands and limits the range of haploid-diploid selection coefficients allowing polymorphism, depending on dominance and selfing mode.

Key Words: Assortative competition, gametic selection, sporophytic selection, population genetics, mixed mating, genetic correlation

 INTRODUCTION

Over 7000 genes are expressed in male gametophytes—pollen grains and tubes and the sperm

they transmit—as they compete in the pistil for ovules (Page and Grossniklaus 2002; Honys and

Twell 2003; Rutley and Twell 2015). Upwards of 60% of known plant genes are expressed in

both the gametophyte and sporophyte (Borges et al. 2009; Arunkumar et al. 2013). For example,

over 100 cases of co-expression in pollen tubes and root hairs have been identified in tobacco

(Hadif et al. 2012). This high degree of co-expression presents opportunity for pollen selection to

affect sporophyte evolution, and *vice versa*. Haldane (1932) recognized this potential, writing

"Clearly a higher plant species is at the mercy of its pollen grains. A gene which greatly

accelerates pollen tube growth will spread through a species even if it causes moderately

disadvantageous changes in the adult plant." Simply, genes conferring an advantage in the

gametophyte should fix unless there is strong antagonistic pleiotropy with sporophyte fitness.

Selection should be strong on alleles with concordant impacts on both gametophyte and

sporophyte stages. Deleterious recessive mutations, which are masked in heterozygous

sporophytes, are fully exposed to selection in haploid gametophytes. This facilitates purging,

diminishing inbreeding depression (Charlesworth and Charlesworth 1992). Some have suggested

that alleles promoting pollen competition will increase offspring quality: large stigmatic pollen

loads lead to intense competition in the style, favouring genetically superior gametophytes,

which contribute their genes to the seed (Mulcahy and Mulcahy 1987; Winsor et al. 1987;

Winsor et al. 2000; but see Baskin and Baskin 2015; Pélabon et al. 2016). Walsh and

Charlesworth (1992) found that when fitness effects on the gametophyte and sporophyte are

concordant, selection will quickly eliminate deleterious mutations and drive favourable ones to

fixation. But, as Haldane suggested, polymorphism may be maintained if increased pollen

competitive ability is counterbalanced by costs to the sporophyte.

Ploidally antagonistic selection has been considered previously (Ewing 1977). Immler,

Arnqvist and Otto (2012) explored the maintenance of genetic variation under discordant

selection between haploid and diploid phases, as well as under sex-specific selection.

Considering an autosomal gene, they demonstrated polymorphism is most easily maintained

under negative ploidy-by-sex interactions, under which sex-specific selection acts in opposite

directions in haploids and diploids. They conclude that polymorphism can be maintained, even in

the absence of sex differences in selection, when selection is opposing between sporophytic and

gametophytic stages.

High rates of self-pollination diminish the advantage of pollen competitiveness due to

assortative competition (Mazer et al. 2010). Imagine a one locus system, with allele C_g

conferring a competitive advantage to pollen (i.e., in the gametophytic phase) and the alternate

allele C_s advantages in the sporophytic phase. When a C_gC_g homozygote self-pollinates, the C_g

allele gets passed to the offspring by necessity. On the opposite homozygote, the C_s allele is

likewise successful by necessity, despite its competitive inferiority. The competitive advantage

of C_g- over C_s-bearing pollen is realized when heterozygotes self-pollinate, but because selfing

reduces heterozygote frequency, over time there are increasingly fewer opportunities for C_g to

exercise its superiority. Thus, with obligate selfing, the C_s allele gains refuge from competition

and quickly goes to fixation due to its higher sporophytic fitness. Consistent with these

predictions, Mazer et al. (2018) demonstrated faster pollen tube growth rates in the

predominantly outcrossing *Clarkia unguiculate* (Onagraceae) compared to the facultatively

selfing *C. exilis*. A similar result was found for selfing and outcrossing populations of *Clarkia*

tembloriensis (Smith-Huerta 1996).

70   About 40% of plant species have mixed-mating systems: individuals receive mixtures of

self and outcross pollen (Schemske and Lande 1985; Goodwillie et al. 2005). Of the 155 studies

of mixed-mating systems in plants reviewed by Barrett and Eckert (2012), two-thirds report

selfing rates in excess of 20%. In Whitehead et al. (2018), it was found that 63% of 105

surveyed species had at least one population with a mixed-mating system. How do intermediate

selfing rates influence the fate of a new mutation increasing pollen competitiveness at the

expense of the sporophyte?

77   An important paper by Jordan and Connallon (2014) considered the impact of self-

fertilization on sexual conflict in hermaphrodites. They found that selfing narrows invasion

conditions of new alleles conferring male-benefit in the haploid phase if they impose a cost on

diploid female function. Their model considered the case in which gamete quality is determined

by parental genotype. In this situation, a C_g-bearing gamete from a homozygous adult differs

from a C_g-bearing gamete produced by a heterozygote, weakening the gamete haplotype-

phenotype correspondence. Gene expression in animal sperm is limited (Parker and Begon

1993), and so it is reasonable to assume genetic paternal effects on sperm quality for these

species. In plants, adult diploid condition can affect pollen size and quality (Young and Stanton

1990; Mazer and Gorchov 1996; Galloway 2001; Smith-Huerta et al. 2007). Pollen size, which

could well be mediated by genetically influenced resource allocation in the sporophyte, readily

responds to artificial selection (Sarkissian and Harder 2001; McCallum and Chang 2016).

89   We explore a different situation, in which pollen competitiveness is affected by genes

expressed directly in the male gametophyte. Here one may expect stronger selection responses

due to stronger haplotype-phenotype correspondence. In our one-locus model, allele C_g increases

pollen competitive ability at the expense of sporophyte fitness. Analytical solutions show there

are maximum and minimum selfing rates allowing invasion of C_g and C_s alleles, respectively,

and that stable polymorphism is possible only when C_g is recessive. Further analyses show that

as selfing rate increases, C_g's competitive advantage is negated by increasingly smaller diploid

cost.

METHODS

The models

Departures from Hardy-Weinberg equilibrium via selfing necessitate that selection be modelled

in terms of genotype frequencies (Crow and Kimura 1970). We refer to the frequencies of the

C_gC_g, C_gC_s and C_sC_s genotypes as X_0, X_1 and X_2, respectively, where subscripts denote the

number of C_s alleles. Allele frequencies are p and q for C_g and C_s, respectively.

Assume a pleiotropic effect by which the C_g allele increases pollen competitive ability

but decreases the number of pollen grains and ovules that the diploid plant contributes to the

gamete pool, either by reduced survival or lower flower production. The C_s allele is favoured

through both sex functions in the diploid stage, disfavoured in haploid pollen and has no effect

on ovules. Fitness of sporophyte C_sC_s is $W_2 = 1$, and that of C_gC_g is $W_0 = 1\text{-}s$, while the

heterozygote C_gC_s is $W_1 = 1\text{-}hs$, where h is the degree of dominance for fitness. As selection may

act either through survival or fertility, we assume a common dominance coefficient for male and

female function. (For a discussion of the effects of sex-specific dominance, see Appendix A).

The absolute competitive ability of pollen bearing the C_g allele is 1 and that of pollen bearing the

C_s allele is 1-t. All individual plants are expected to have many flowers. Population size is

effectively infinite.

The level of self-pollination in the model depends on r, the proportion of pollen allocated

to self-mating, which reflects the potential rate of self-fertilization. Absent selection, r equals the

proportion of zygotes that are produced by self-pollination, *i.e.*, the realized selfing rate. As

shown below, potential and realized selfing rates can differ when selection acts on the

gametophyte. For sake of brevity, r is hereafter referred to as selfing rate. We assume all plants

have the same r value. We model two selfing forms: fixed selfing and mass-action selfing,

defined below. The models we present assume haploid genetic control over pollen competitive

ability. As we present our results, we will for comparison also present results using models by

Jordan and Connallon (2014; see Appendix B), which assume diploid control.

FIXED SELFING

The fixed selfing model echoes the biology of species such as *Impatiens* and *Viola*, where

cleistogamous flowers are obligate selfers, and chastogamous flowers outcross (Schemske 1978,

Winn and Moriuchi 2009). In this model, proportion r of flowers on each plant is dedicated to

selfing while 1-r of flowers is allocated to outcrossing. Thus, a stigma receives either only self-

pollen or only outcross-pollen. Selection for pollen competitiveness occurs on all outcrossing

flowers, but among selfing flowers, occurs only on heterozygotes. Our recursion equations for

genotype frequencies consider selection during the diploid and haploid stages in separate steps.

After diploid selection, diploid genotype frequencies are:

$$x_0 = \frac{(1-s)X_0}{\overline{W}} \tag{1a}$$

$$x_1 = \frac{(1-hs)X_1}{\overline{W}} \tag{1b}$$

$$x_2 = \frac{X_2}{\overline{W}} \tag{1c}$$

132   where $\overline{W}$ is the mean diploid fitness:

$$\overline{W} = (1 - s)X_0 + (1 - hs)X_1 + X_2. \tag{2}$$

After mating and haploid selection, offspring genotype frequencies are

$$X_0' = r\left[x_0 + \tfrac{1}{2}x_1\left(\tfrac{1}{2-t}\right)\right] + (1-r)\left[\left(x_0 + \tfrac{1}{2}x_1\right)\alpha\right] \tag{3a}$$

$$X_1' = r\left[\tfrac{1}{2}x_1\right] + (1-r)\left[x_0(1-\alpha) + \tfrac{1}{2}x_1 + x_2\alpha\right] \tag{3b}$$

$$X_2' = r\left[x_2 + \tfrac{1}{2}x_1\left(\tfrac{1-t}{2-t}\right)\right] + (1-r)\left[\left(x_2 + \tfrac{1}{2}x_1\right)(1-\alpha)\right] \tag{3c}$$

where α reflects the relative competitiveness of C_g-bearing pollen as a function of the absolute

competitiveness conferred by the C_g allele and the frequency of the C_g allele in the pollen pool as

a result of genotype frequency following selection:

$$\alpha = \frac{x_0 + \tfrac{1}{2}x_1}{x_0 + \tfrac{1}{2}x_1(2-t) + x_2(1-t)}. \tag{4}$$

It follows that the relative competitiveness of C_s-bearing pollen is (1- α).

To understand Equations (3a-c), consider separately the offspring produced on selfing (r)

and outcrossing (1-r) flowers. Under selfing, competition among pollen tubes carrying opposite

alleles only occurs within heterozygote pistils. Assuming heterozygotes produce C_g and C_s pollen

in equal numbers, ½ of their offspring will likewise be heterozygous, with the remaining

offspring divided between C_gC_g and C_sC_s homozygotes in proportions of $\left(\dfrac{1}{2-t}\right)$ and $\left(\dfrac{1-t}{2-t}\right)$

(Appendix C). We assume each selfing flower produces enough pollen to fertilise its ovules,

such that only female diploid fitness terms, and not α (*i.e.*, male diploid fitness terms), are

included. Simply, the number of offspring produced via selfing is limited by seed production, not

pollen production. Offspring produced through outcrossing will be a function of both female and

male diploid fitness, as well as allelic differences in pollen competitive ability, specified by α.

For simplicity, we make no provision for inbreeding depression; increasing failure rate of selfed

offspring has the same effect as reducing the basic selfing rate (see Jordan and Connallon 2014).

Mass-action selfing

Mass-action selfing (Holsinger 1991) echoes more general modes of mixed-mating. Here r still

denotes the proportion of pollen individuals allocate to selfing, but in this case all stigmas

receive a mix of self- and outcross-pollen, resulting in inter-allelic competition for all ovules.

Under this scenario, the amount of self-pollen received by a stigma is a function of two factors,

one being r, and the other being the quantity of pollen produced by the individual relative to that

it receives through outcrossing. Thus, the true selfing rate is dependent on genotype frequency.

We make no provisions for pollen discounting here but consider this elaboration in Appendix D.

163   Assuming diploid selection occurs according to Equations (1a-c), we arrive at the

164   following genotype frequencies after mating and haploid selection:

$$X_0' = x_0 \left[\frac{rW_0 + (1-r)\left(x_0 + \frac{1}{2}x_1\right)}{P_0} \right] + \frac{1}{2}x_1 \left[\frac{r\frac{1}{2}W_1 + (1-r)\left(x_0 + \frac{1}{2}x_1\right)}{P_1} \right] \tag{5a}$$

$$X_1' = \frac{1}{2}x_1 + x_0 \left[\frac{(1-r)\left(x_2 + \frac{1}{2}x_1\right)(1-t)}{P_0} \right] + x_2 \left[\frac{(1-r)\left(x_0 + \frac{1}{2}x_1\right)}{P_2} \right] \tag{5b}$$

$$X_2' = x_2 \left[\frac{rW_2(1-t) + (1-r)\left(x_2 + \frac{1}{2}x_1\right)(1-t)}{P_2} \right]$$

$$+ \frac{1}{2}x_1 \left[\frac{r\frac{1}{2}W_1(1-t) + (1-r)\left(x_2 + \frac{1}{2}x_1\right)(1-t)}{P_1} \right] \tag{5c}$$

165   where P_0, P_1 and P_2 give the total pollen received by X_0, X_1 and X_2 individuals respectively and

166   are defined as:

$$P_0 = rW_0 + (1-r)\left(x_0 + \tfrac{1}{2}x_1(2-t) + x_2(1-t)\right) \tag{6a}$$

$$P_1 = r\frac{1}{2}W_1(2-t) + (1-r)\left(x_0 + \tfrac{1}{2}x_1(2-t) + x_2(1-t)\right) \tag{6b}$$

$$P_2 = rW_2(1-t) + (1-r)\left(x_0 + \tfrac{1}{2}x_1(2-t) + x_2(1-t)\right). \tag{6c}$$

In Eq. 5a, the two terms represent C_gC_g offspring seed produced by C_gC_g and C_gC_s plants with

post-diploid selection frequencies of x_0 and x_1. All ovules on the homozygote carry C_g, whereas

on heterozygotes only half do. The terms inside the square brackets are the probability that an

ovule receives C_g-bearing sperm, which depends on the amount of successful C_g pollen

(numerator) divided by the total amount of pollen arriving at the stigma (denominator). Eq. 5c

follows similar logic for C_sC_s offspring. Turning to Eq. 5b, the three terms represent C_gC_s

offspring produced by C_gC_s, C_gC_g and C_sC_s plants, respectively. Half of all seed produced by

C_gC_s individuals are also heterozygotes. The proportion of heterozygotes produced by C_gC_g

individuals are a function of the relative abundance and competitiveness of C_s pollen (numerator)

compared to the total pollen arriving on C_gC_g stigmas. This proportion is given by the term

inside the first square bracket. C_gC_s offspring from C_sC_s individuals follow identical logic but for

C_g pollen.

Software and computation

All analytical analyses were derived using Wolfram Mathematica 11.0.1.0. To find conditions

for stable polymorphism, we derived leading eigenvalues for Jacobian Matrices using genotype

recursions for gametophytic control (Eqs. 3a-c, 5a-c) and sporophytic control (Eqs. B9-B11,

Appendix B) of pollen competitiveness and determined the conditions that allowed invasion at a

boundary by solving for parameters producing a leading eigenvalue of absolute value greater

than 1. Invasion analyses were performed for two boundaries: the X_0 boundary, at which the C_s

allele appears in a population fixed for the C_g allele (i.e., $p = 1$), and the X_2 boundary, at which

the C_g allele appears in a population fixed for the C_s allele (i.e., $p = 0$). Code for analytical

analyses is provided in the online supplemental materials.

To find expressions for equilibrium allele frequencies, we generated power series

expansions (assuming weak selection for simplification) for expressions giving the single-

generation change in genotype frequencies. We then solved for the equilibrium by finding the p

value yielding zero change. Predicted equilibrium frequencies were verified numerically.

Numerical methods and results are presented in Appendix E.

194   **RESULTS**

Analytical invasion conditions

We explored the impact of selfing rate on invasion potential for both C_g and C_s when rare. For

fixed selfing, analytical conditions for invasion at the X_0 and X_2 boundaries are derived using

Equations (3a-c). Likewise, analytical conditions for mass-action selfing use Equations (5a-c).

For comparison with diploid control models (Jordan and Connallon 2014) and, in the case of

mass-action selfing, to produce tractable solutions, we assume weak selection (i.e., $0 < s, t \ll 1$),

though we do find good agreement between invasion conditions predicted under fixed selfing

both with and without weak selection (Appendix E).

Fixed selfing

204   Under fixed selfing, boundary selfing rates permitting invasion of the C_g and C_s alleles,

205   respectively, are as follows:

$$r < \frac{t+2hs}{s+t-2hs} \qquad \text{(7a)}$$

$$r > \frac{t+2s(1-h)}{t+s(1+2h)} \qquad \text{(7b)}$$

These analytical solutions are visualised in Figure (1a-c). Under random mating (i.e., $r = 0$), and

when either fully dominant or recessive ($h = 0$ or 1), the C_g allele invades and goes to fixation if

$s/t < 0.5$ (*i.e.*, when the haploid competitive benefit is more than twice the diploid fitness cost).

When diploid fitness effects are additive ($h = 0.5$), fixation occurs even as s/t approaches 1.0. C_g

wins under random mating as the C_s allele has no refuge from competition. Under complete

selfing ($r = 1$), C_g never invades. Pollen competition occurs only on heterozygotes, and since

selfing reduces heterozygote frequencies, opportunities for C_g-bearing pollen to outcompete C_s

effectively vanish. As C_s confers higher sporophyte fitness in the homozygous state, it increases

when rare.

215   The analytical solutions indicate that polymorphism can be maintained in mixed-mating

systems ($0 < r < 1$) with fixed selfing only if the deleterious effect of the C_g allele in the diploid

phase is recessive ($h < 0.5$) (Figures 1a-c, 2). When completely recessive ($h = 0$), there is a range

of selfing rates in which both alleles can invade, implying protected polymorphism. The C_g allele

invades because, when rare, it occurs exclusively in heterozygotes, where its negative effect on

diploid fitness is masked from selection and moderate selfing allow opportunities to outcompete

C_s-bearing pollen for ovules on heterozygotes. Within this range of selfing rates, the C_s allele

also increases when rare because, 1) heterozygotes have higher relative fitness than C_gC_g

homozygotes and 2) selfing reduces exposure of C_s pollen to competition with superior C_g

pollen.

225   Additive fitness effects ($h = 0.5$) largely prohibit polymorphism (Figures 1b, 2). When h

$= 0.5$, the critical r values for invasion (Eq. 7a-b) reduce to approximately $1 - \frac{s}{t}$ for both alleles.

As *h* increases, the relative fitness of heterozygotes decreases. It then follows that the initial

advantage enjoyed by both alleles when rare also diminishes. To invade, C_g requires low *r* values

to win enough contests to offset lower heterozygote fitness. C_s will require higher *r* values to

avoid competitive losses and produce enough high-fitness X_2 homozygotes to offset lower fitness

of heterozygotes.

When the diploid fitness disadvantage of the C_g allele is dominant ($h = 1$), there is a range

of selfing rates that prohibit either allele from invading (Figures 1c, 2). For a given value of *s*,

the maximum selfing rate allowing C_g invasion is lowered because its adverse effect on

sporophyte fitness is fully expressed by heterozygotes. Oppositely, the critical *r* for C_s invasion

increases. When rare, the positive diploid fitness effects of the C_s allele are concealed in

heterozygotes. Only very high selfing rates can produce C_s homozygotes quickly enough to

provide sufficient refuge from competition, allowing C_s to spread.

Mass-action selfing

Considering mass-action selfing, the maximum and minimum *r* values allowing invasion of the

C_g and C_s alleles can be solved analytically. However, even under weak selection the expressions

are complex and difficult to interpret biologically; these are presented in Appendix F and are

visualised in Figure (1d-f).

While the expressions for critical *r* values are complex, similarly derived expressions for

critical diploid selection *s* values can be used to understand how selfing rate affects invasion

success. Expressions for maximum and minimum *s* values permitting invasion of C_g and C_s

alleles respectively are given below:

$$s < \frac{(-2+r+r^2)t}{(2h(r-1)-r)(r-2)(1+r)} \tag{8a}$$

$$s > \frac{(-2+r+r^2)t}{(2+2h(r-1)-r)(r-2)(1+r)}.$$ (8b)

Taking the partial derivative of Eqs. 8a and 8b with respect to *r* lends insight into how these

critical *s* values change in response to selfing. (For a discussion of these partial derivative

expressions, see Appendix G). These show an increase in selfing rate always decreases the

maximum and minimum *s* values permitting invasion of C_g and C_s alleles respectively (Figure

1d-f). It then follows that increased selfing contracts the parameter space allowing invasion of an

allele conferring a benefit to pollen competitiveness at the expense of diploid fitness.

Comparing invasion conditions for fixed and mass-action selfing

Invasion conditions for the two selfing modes were qualitatively similar, with modest

quantitative differences. We compared sensitivity of the minimum and maximum *s* values

permitting invasion of C_s and C_g alleles to increases in *r* for both systems. (Partial derivative

expressions for fixed selfing are also provided in Appendix G). For a given set of *h, s* and *t*

values, *r* values allowing C_g and C_s invasion are slightly lower under mass-action selfing relative

to those of fixed-selfing (Figure 2), narrowing the permissible parameter space. Thus, under

mass-action selfing, the C_g allele can persist at higher selfing rates. Additionally, the range of *r*

values permitting polymorphism (when $h < 0.5$) or unstable equilibrium (when $h > 0.5$) contracts

under mass-action selfing, which is consistent with narrowed parameter space regions for

polymorphism and unstable equilibrium under mass-action selfing observed in Figure 2.

Under fixed selfing, male diploid fitness (pollen quantity) is moot since all flowers of

each genotype are assumed to produce enough pollen to fertilise their ovules. Under mass-action

selfing, the quantity of pollen produced by an individual affects its realized fertilisation success

through self-pollen, as each stigma can receive a mixture of C_g- and C_s- bearing pollen,

restricting the competition refuge for C_s. While each C_g-bearing grain has a greater chance of

success, this advantage is partially countered by lower pollen production in C_g homozygotes. The

different impacts on male function shift the invasion boundary upward under mass-action

selfing. For example, when $h = 0$, $s = 0.01$ and $t = 0.04$ (visualised as $s/t = 0.25$ on Figure 1), the

maximum selfing rate allowing the C_g allele to invade increases from roughly $r = 0.8$ to $r = 0.84$.

Likewise, the minimum selfing rate allowing C_s to invade increases from $r = 0.67$ to $r = 0.75$.

Fixation and protected polymorphism

Having established invasion conditions for the C_g and C_s alleles, this section explores their

equilibrium frequencies. We first present analytical solutions to the recursion equations (Eqns.

3a-c, 5a-c) assuming weak selection ($0 < s$, $t << 1$). We then highlight the difference in outcomes

when pollen competitiveness is under gametophyte genetic control, as Haldane speculated,

versus sporophyte control (seen in Jordan and Connallon 2014). Under fixed selfing, the

equilibrium value ($\hat{p}$) of the C_g allele is given as:

$$\hat{p} = \text{Min}\left[1, \text{Max}\left[0, \frac{1}{2} + \frac{t(1-r) - s}{(2\,(-1 + 2\,h)(-1 + r)s)}\right]\right]. \tag{10}$$

When C_g is recessive ($h < 0.5$) and mating is mixed ($0 < r < 1$), the denominator of the far-right

term is always positive. Whether $\hat{p}$ falls above or below 0.5 then depends on the sign of the

numerator. If $r < 1 - s/t$, then $\hat{p}$ will fall above 0.5 but fall below if the inverse is true. Simply,

the higher the selfing rate (r) the higher the diploid cost (s) needed to prevent C_g fixation, which

is implied by a $\hat{p}$ value < 0.5. When C_g is dominant ($h > 0.5$), the above expression gives the

repelling equilibrium frequency (above $\hat{p}$ the C_g allele goes to fixation and below is eliminated).

When $h > 0.5$ and mating is mixed, the denominator of the far-right term is always negative.

Thus, the repelling $\hat{p}$ value falls above 0.5 when the numerator is negative and below when

positive. Looking at the effect of selfing rate on $\hat{p}$, the threshold $\hat{p}$ allowing C_g invasion

decreases as r decreases, as the numerator will become increasingly positive. When $h = 0.5$, the

denominator of the far-right terms goes to zero, putting the whole term to infinity. Threshold

selfing rates permitting or preventing C_g invasion are thus given by the sign of infinity, which is

determined by if r is greater than or less than $1 - s/t$ respectively.

Under mass-action selfing, the equilibrium frequency of the C_g allele is:

$$\hat{p} = \mathrm{Min}\left[1, \mathrm{Max}\left[0, \frac{(2\,h\,(-1+r)-r)(-2+r)(1+r)s + (-1+r)(2+r)t}{2\,(-1+2\,h)\,(-2+r)\,(-1+r)\,(1+r)\,s}\right]\right]. \tag{11}$$

When $h < 0.5$, Eqs. 10 and 11 provide the equilibrium frequency of the C_g allele. When $h > 0.5$,

these expressions provide the 'repelling' allele frequency. Figure 3 plots the equilibrium

frequency of C_g as a function of selfing rate, r, under relatively weak selection. (For a

comparison of analytical results to numerically derived frequencies under strong selection, see

Appendix E).

When the negative diploid effect of C_g is dominant ($h = 1$), high selfing prevents C_g

invasion when rare, while low selfing prevents C_s invasion. Under intermediate selfing rates,

neither allele can increase when rare (Figure 3c, f). Following the appearance of the C_g allele in a

population by mutation, its initial frequency will almost inevitably fall below the repelling

frequencies predicted by Eqs. 10 or 11 and thus it will be eliminated.

Comparison of Eqs. 10 and 11 reveals that when $0 < h < 0.5$, the equilibrium frequency

$\hat{p}$ is higher under mass-action selfing relative to fixed selfing; however, when $h > 0.5$, the inverse

is true (for derivation, see online supplemental materials). As suggested by invasion conditions,

under mass-action selfing, minimum and maximum selfing rates allowing invasion of C_g and C_s

alleles respectively increase for given h, s and t values (Figure 3d-f). Additionally, parameter

space allowing both polymorphism (Figures 3d) and unstable equilibrium (Figures 3f) contracts

under mass-action selfing relative to fixed (Figures 3a, 3c).

As is visualised in Figures 1 and 3, invasion conditions and equilibrium allele frequencies

under both selfing modes assuming haploid control of pollen competitiveness are qualitatively

similar to those derived using models provided in Jordan and Connallon (2014), where

competitiveness is under paternal genotype control (for derivation of analytical results, see

online supplemental materials; for models and discussion of analytical solutions, see Appendix

B). They also found selfing and dominance restrict potential for stable polymorphism and

ultimately diminish selection through male function. Our results show, however, that

gametophytic control substantially expands the conditions favouring an allele promoting pollen

competitiveness. With gametophyte gene expression, pollen phenotype is more strongly

correlated with haplotype. Heterozygotes always produce a 50:50 mix of phenotypically and

genetically 'strong' and 'weak' pollen competitors. Under sporophytic control, heterozygotes can

produce phenotypically all strong ($h = 0$), all weak ($h = 1$), or all intermediate ($h = 0.5$)

competitors, despite always producing a 50:50 genetic mix. Further, under gametophytic control,

variance in competitive ability between pollen carrying opposite alleles is constant. Under

paternal control, variance in competitive ability changes with genotype frequency: increasing

heterozygote frequency decreases the difference in mean competitiveness of the two pollen

haplotypes. Consequentially, C_g invasion occupies a markedly smaller parameter space when

under diploid genetic control (Figure 1). By extension, we see that the C_g allele reaches a lower

equilibrium frequency, *i.e.*, invades with more difficulty, for a given selfing rate when it is

expressed in the sporophyte rather than the gametophyte (Figure 3).

Returning to Haldane's conjecture that plants are at the mercy of their pollen grains, how

strong must sporophyte selection against a pollen-competition allele be to prevent its fixation?

The fate of the C_g allele depends on selection strength during the two life stages, which can be

considered as a ratio of diploid and haploid selection coefficients, s/t. At a given s/t ratio,

increasing the selfing rate will restrict opportunities for C_g pollen to exercise its superiority,

muting the selective advantage. Against this, at any given ratio, Mendelian dominance alters how

often the diploid cost is paid. If C_g is recessive ($h = 0$), heterozygotes pay no diploid cost. If

dominant ($h = 1$), it masks the C_s benefit and heterozygotes pay full cost; however, since selfing

decreases heterozygote frequency, the dominance issue becomes moot. These conditions produce

the general outcome that s/t ratios allowing C_g to invade, and to be maintained in polymorphism,

tend to decrease as both r and s increase (Figure 4).

DISCUSSION

Classical theory, which assumes random mating, predicts an allele conferring an advantage in

pollen competitiveness fixes unless a trade-off with sporophyte fitness sufficiently counters this

haploid advantage. Our models predict that even moderate selfing can prevent the spread of a

pollen-expressed allele that confers competitive superiority (Table 1). Increased selfing increases

assortative competition among pollen, restricting opportunity for a competitive allele to exercise

its superiority. When homozygotes self completely, they deposit only one pollen haplotype per

stigma, so by necessity that haplotype cannot be out-competed by another. Alternate haplotypes

compete only on heterozygotes. Further, selfing depresses heterozygote frequency such that the

proportion of the pollen pool engaged in competitive contests falls in proportion to the selfing

rate.

The dominance relationships between the competitive and non-competitive alleles in the

diploid stage also influence invasion. When the sporophyte fitness cost of the C_g allele is

recessive, it is shielded from negative selection during the diploid phase. Complete dominance

fully exposes it to negative selection. Thus, the conditions for invasion narrow with increases in

the dominance coefficient (Table 1, Figures 1-4).

Increased dominance also narrows the condition for stable polymorphism (Table 1,

Figures 1-4). When costs are recessive, heterozygotes gain success as fathers because 50% of

their pollen is competitively superior and they pay no cost. This can lead to overall heterozygote

superiority. Under random mating, this over-dominance in total fitness can maintain stable

polymorphism; however, selfing diminishes the impact of over-dominance by eroding

heterozygote frequency: heterozygosity lost to segregation is not replenished by matings between

opposite homozygotes, and so the competitive allele is eventually carried mostly by the

homozygote, who pay full cost. If the competitive allele is dominant, stable polymorphism is

precluded. In this case, the heterozygotes still get half the paternal benefit but pay full cost

(Table 1, Figures 1-4), making over-dominance for total fitness unachievable.

By comparing invasion conditions and equilibrium allele frequencies under haploid and

diploid control (as seen in Jordan and Connallon 2014) of pollen ability (Figures 1, 3), we

demonstrate that the form of genetic control overseeing competitive ability has important

implications for the fate of an allele conferring an advantage to pollen ability. The stronger

haplotype-phenotype correlation in our model is important during the early stages of invasion,

when pollen carrying the competitive allele is produced only by heterozygotes: it assures that the

pollen grains carrying the competitive allele are indeed competitive. This clear genetic signal

expands conditions for a costly allele to invade and spread; stronger selfing is required to counter

the advantage.

Our fixed and mass-action selfing models are caricatures representing two extremes in

pollination systems, but real systems can be intermediate. In *Impatiens palida*, cleistogamous

flowers always self whereas there is a small incidence of geitenogamous selfing in chastogamous

flowers (Schemske 1978). Gradations in selfing prior to bud opening or delayed self-pollination

when pollinator visitation is low (Loyd 1979) can generate variation in the proportion of self-

pollen involved with competitive contests with outcross. However, we have shown strong

qualitative agreement between the outcomes of the two extreme models, both leading to the

general prediction that selection favouring increased pollen competitive ability is weakened by

selfing.

Genomic analysis of exclusively pollen-expressed loci expressed, where diploid costs are

necessarily zero, bears out the prediction of relaxed selection under self-pollination. In the self-

incompatible *Capsella grandiflora*, evidence points to stronger purifying selection at sites

expressed solely in the pollen tubes than at sites expressed exclusively in the seedling (Arukumar

et al., 2011). In the highly selfing *Arabidopsis thaliana*, by contrast, purifying selection on

pollen-exclusive genes is weaker than on sporophyte-exclusive genes, as evidenced by a higher

frequency of premature stop codons in the former (Harrison et al. 2015). Greater scrutiny of loci

expressed during both stages are needed to clarify the general importance of negative pleiotropic

constraints on the evolution of gametophyte and sporophyte traits.

398   **References**

Arunkumar, R., E. B. Josephs, R. J. Williamson, and S. I. Wright. 2013. Pollen-specific, but not

sperm-specific, genes show stonger purifying selection and higher rates of positive

selection than sporophytic genes in *Capsella grandiflora*. Mol. Biol. Evol. 30:2475-2486.

Barrett, S. C. and C. G. Eckert. 2012. 14 Variation and Evolution of Mating Systems in Seed

Plants. Biological approaches and evolutionary trends in plants :229.

Barrett, S. C. H. 1998. The evolution of mating strategies in flowering plants. Trends Plant Sci.

3:335-341.

Baskin, J. M. and C. C. Baskin. 2015. Pollen (microgametophyte) competition: an assessment of

its significance in the evolution of flowering plant diversity, with particular reference to

seed germination. Seed Sci. Res. 25:1-11.

Borg, M., L. Brownfield, and D. Twell. 2009. Male gametophyte development: a molecular

perspective. J. Exp. Bot. 60:1465-1478.

Campbell, S. A. 2015. Ecological mechanisms for the coevolution of mating systems and

defence. New Phytol. 205:1047-1053.

Charlesworth, D. and B. Charlesworth. 1992. The effects of selection in the gametophyte stage

on mutational load. Evolution 46:703-720.

Crow, J. F. and M. Kimura. 1970. Introduction to Population Genetics Theory. Harper & Row,

New York.

Ewing, E. 1977. Selection at the haploid and diploid phases: clinical variation. Genetics. 87:195-

208.

Damgaard, C. and V. Loeschcke. 1994. Genetic-variation for selfing rate the dependence of

selfing rate on mating history in *Brassica napus* (rape seed). Heredity 72:570-573.

Galloway, L. F. 2001. The effect of maternal and paternal environments on seed characters in the

herbaceous plant *Campanula americana* (Campanulaceae). Am. J. Bot. 88:832-840.

Goodwillie, C., S. Kalisz, and C. G. Eckert. 2005. The evolutionary enigma of mixed mating

systems in plants: occurrence, theoretical explanations, and empirical evidence. Annu. Rev.

Ecol. Evol. Syst. 36:47-79.

Hafidh, S., K. Breznenová, P. Růžička, J. Feciková, V. Čapková, and D. Honys. 2012.

Comprehensive analysis of tobacco pollen transcriptome unveils common pathways in

polar cell expansion and underlying heterochronic shift during spermatogenesis. BMC

Plant Biol. 12(1): 24.

Haldane, J. B. S. 1932. The Causes of Evolution. Reprinted 1990, Princeton University Press.

Harrison, M. C., E.B. Mallon, D. Twell, and R.L. Hammond. 2015. Deleterious mutation

accumulation in *Arabidopsis thaliana* pollen genes: a role for a recent relaxation of

selection. bioRxiv, 016626.

Holsinger, K. E. 1991. Mass-action models of plant mating systems: the evolutionary stability of

mixed mating systems. Am. Nat. 138:606-622.

Igic, B. and J. R. Kohn. 2006. Bias in the studies of outcrossing rate distributions. Evolution 60

(5): 1098–1103.

Immler, S., G. Arnqvist, and S. P. Otto. 2012. f. Evolution 66:55-65.

Jarne, P. and D. Charlesworth. 1993. The evolution of the selfing rate in functionally

hermaphrodite plants and animals. Annu. Rev. Ecol. Evol. Syst. 24:441-466.

Jordan, C. Y. and T. Connallon. 2014. Sexually antagonistic polymorphism in simultaneous

hermaphrodites. Evolution 68:3555-3569.

Lloyd, D. G. 1979. Some reproductive factors affecting the selection of self-fertilization in

plants. The American Naturalist 113(1):67-79.

Mazer, S. J., A. A. Hove, B. S. Miller, and M. Barbet-Massin. 2010. The joint evolution of

mating system and pollen performance: Predictions regarding male gametophytic evolution

in selfers vs. outcrossers. Perspect. Plant Ecol. Evol. Syst. 12:31-41.

Mazer, S. J., B. T. Hendrickson, J. P. Chellew, L. J. Kim, J. W. Liu, J. Shu, and M. V. Sharma.

2018. Divergence in pollen performance between *Clarkia* sister species with contrasting

mating systems supports predictions of sexual selection. Evolution 72(3):453-472.

McCallum, B. and S. M. Chang. 2016. Pollen competition in style: Effects of pollen size on

siring success in the hermaphroditic common morning glory, *Ipomoea purpurea*. Am. J.

Bot. 103(3):460-470.

Mulcahy, D. L. and G. B. Mulcahy. 1987. The effects of pollen competition. Am. Sci. 75:44-50.

Parker, G. and M. Begon. 1993. Sperm competition games: sperm size and number under

gametic control. Proc. R. Soc. London, Ser. B. 253:255-262.

Pélabon, C., L. Hennet, G. H. Bolstad, E. Albertsen, Ø. H. Opedal, R. K. Ekrem, and W. S.

Armbruster. 2016. Does stronger pollen competition improve offspring fitness when pollen

load does not vary? Am. J. Bot. 103:522-531.

Sarkissian, T. S. and L.D. Harder. 2001. Direct and indirect responses to selection on pollen size

in *Brassica rapa* L. J. Evol. Biol. 14(3):456-468.

Schemske, D. W. 1978. Evolution of reproductive characteristics in Impatiens (Balsaminaceae):

the significance of cleistogamy and chasmogamy. Ecology 59(3):596-613.

Schemske, D. W. and R. Lande. 1985. The evolution of self-fertilization and inbreeding

depression in plants. II. Empirical observations. Evolution 39:41-52.

Smith-Huerta, N. L. 1996. Pollen germination and tube growth in selfing and outcrossing

populations of *Clarkia tembloriensis* (Onagraceae). Int. J. Plant Sci. 157:228-233.

Sousa, E., B. Kost, and R. Malhó. (2008). *Arabidopsis* phosphatidylinositol-4-monophosphate 5-

kinase 4 regulates pollen tube growth and polarity by modulating membrane recycling.

Plant Cell, 20(11), 3050-3064.

Walsh, N. and D. Charlesworth. 1992. Evolutionary interpretations of differences in pollen tube

growth rates. Q. Rev. Biol. 67:19-37.

Winn, A. A., & Moriuchi, K. S. 2009. The maintenance of mixed mating by cleistogamy in the

perennial violet *Viola septemloba* (Violaceae). American Journal of Botany 96(11):2074-

2079.

Winsor, J., L. Davis, and A. Stephenson. 1987. The relationship between pollen load and fruit

maturation and the effect of pollen load on offspring vigor in *Cucurbita pepo*. Am. Nat.

129:643-656.

Winsor, J. A., S. Peretz, and A. G. Stephenson. 2000. Pollen competition in a natural population

of *Cucurbita foetidissima* (Cucurbitaceae). Am. J. Bot. 87:527-532.

Young, H. J. and M. L. Stanton. 1990. Influence of environmental quality on pollen competitive

ability in wild radish. Science 248:1631-1633.

483   **Tables**

484 TABLE 1.   Summary of model outcomes for mutation to an allele that increases pollen (haploid

485 gametophyte) competitive ability but reduces survivorship or fertility in the (diploid) sporophyte.

EXPRESSION OF C_g IN DIPLOID SPOROPHYTE	MATING SYSTEM		
	Obligate Outcrossing ($r = 0$) *Ample opportunities* to exercise pollen competitive superiority	Mixed ($0 < r < 1$) *Fewer opportunities* to exercise pollen competitive superiority	Obligate selfing ($r = 1$) *No opportunities* to exercise pollen competitive superiority
Recessive Fitness cost paid only by homozygote	C_g goes to fixation if sporophyte cost is less than ½ the pollen competitive benefit, otherwise reaches stable polymorphism– *conditions for stable polymorphism broad.*	C_g goes to fixation at increasingly lower sporophyte costs as selfing rate increases; upper limit for cost to permit polymorphism is increasingly reduced at higher selfing rates– *conditions for stable polymorphism narrowed.*	C_g unable to invade.
Dominant Fitness cost paid by *both* homozygote and heterozygote	C_g goes to fixation if sporophyte cost is less than ½ the pollen competitive benefit, but eliminated if costs are higher – *no stable polymorphism.*	C_g goes to fixation at increasingly lower sporophyte costs as selfing rate increases, but *eliminated* if costs are any higher – *no stable polymorphism.*	C_g unable to invade.

487   **Figure captions**

Figure 1

Conditions producing fixation, polymorphism and unstable equilibrium (i.e., fixation is

dependent on initial frequencies) of two alleles C_g and C_s at an autosomal gene under

antagonistic pleiotropy between pollen competitive ability and diploid fitness. Solid lines

delineate conditions under gametophytic genetic control and dashed lines under sporophytic

genetic control. (a-c) Analytical results for the fixed-selfing model assuming weak selection.

Potential for polymorphism diminishes as the dominance of the C_g allele increases, eventually

being replaced by unstable equilibrium when $h > 0.5$. As selfing rate (*i.e.*, proportion pollen

dedicated to selfing) increases, polymorphism requires greater diploid selection s against the C_g

allele to offset its greater pollen competitive ability t. (d-f) Analytical results assuming weak

selection for the mass-action model, where r is the proportion pollen dedicated to selfing and

stigmas receive a mixture of self- and outcross pollen. Conditions allowing invasion of the C_g

allele are notably greater than under fixed selfing, as individual stigmas never receive entirely

self-pollen except when $r = 1$, thus increasing opportunity for competition between C_g- and C_s-

bearing pollen. For both fixed and mass-action selfing, relative to haploid control, diploid control

expands conditions under which the C_s allele can be maintained while contracting those under

which the C_g allele can be maintained.

Figure 2

Conditions producing fixation, polymorphism and unstable equilibrium of C_g and C_s alleles

under variable diploid selection s and dominance h for both fixed and mass-action selfing.

Analytical solutions producing these conditions do not assume weak selection. Solid lines

indicate boundaries under fixed-selfing; dashed lines indicate those for mass-action selfing.

Potential for polymorphism diminishes as the dominance of the C_g allele increases, eventually

being replaced by unstable equilibrium when $h > 0.5$. As the diploid selection coefficient s

increases relative to t, maintenance of the C_g allele requires either lower dominance h or less

allocation of reproductive structures to selfing, given by r. Compared to fixed-selfing, mass-

action selfing conditions permitting polymorphism versus unstable equilibrium are qualitatively

similar but quantitatively more limited. The placement of mass-action lines to the right of those

for fixed selfing are consistent with individual stigmas never receiving solely self-pollen except

when $r = 1$, which increases the opportunity for C_g to exercise its competitive superiority.

Figure 3

Analytical results of equilibrium frequency of the C_g allele across selfing rates $0 < r < 1$ under

gametophytic (haploid) and sporophytic (diploid) genetic control. Analytical solutions were

derived assuming weak selection. Solid lines indicate equilibrium allele frequencies, while

dashed lines indicated the analytical repelling frequency (i.e., the starting p frequency below

which p will be lost and above which it will fix). (a-c) Under the fixed-selfing model, the

window of selfing rates contracts as dominance increases. (d-f) Under the mass-action model, C_s

pollen is never entirely protected from competition with C_g pollen as individual stigmas receive a

mixture of both self and outcross pollen for intermediate selfing rates. Thus, maximum and

minimum selfing rates allowing fixation or maintenance of C_g and C_s alleles respectively are

right-shifted compared to under fixed selfing, reflecting the greater opportunity for the C_g allele

to exercise its advantage in pollen competition with the C_s allele. Notably, for both selfing

modes, the range of selfing rates at which the C_g allele can invade contracts under diploid genetic

control relative to haploid control.

Figure 4

Analytical results for equilibrium frequency of the C_g allele across a range of s/t, i.e., ratios of

diploid and haplod fitness values. Series specify selfing rates r ranging between 0 and 1. Under

both (a-c) fixed and (d-f) mass-action selfing models, increases in dominance of the C_g allele

sharply decrease the maintenance of polymorphism, particularly at lower selfing rates. Noticably,

for a given combination of selfing rate r and dominance h, the C_g allele persists at higher ratios

of s and t under mass-action selfing relative to fixed-selfing , reflecting the greater opportunity

for C_g-bearing pollen to compete C_s-bearing pollen under mass-action selfing. For both selfing

models, regardless of dominance, the C_g allele cannot invade under complete selfing, i.e., $r = 1$.

541   **Appendices**

Appendix A

Implications of sex-specific dominance

Given we assume diploid selection acts either through lower flower production or reduced

survival, we also assume a common dominance coefficient h for female and male function (i.e.,

quantities of ovule and pollen produced). However, it is possible to consider the effects of

different dominance coefficients for male and female function, given as h_m and h_f. In such a case,

absolute fitness via female function for C_gC_s individuals is $W_{1f} = 1-h_f s$ and that via male function

is $W_{1m} = 1-h_m s$. Absolute fitness terms for homozygote individuals remain unchanged from the

main text. Using these sex-specific dominance coefficients, we can derive new expressions for

equilibrium frequency of the C_g allele under weak selection for both fixed (Eq. A1) and mass-

action (Eq. A2) selfing:

$$\hat{p} = \mathrm{Min}\left[1, \mathrm{Max}\left[0, \frac{-h_m\,(-1+r)^2\,s + h_f\,(-1+r^2)\,s + t - r\,(s+t)}{(2\,(-1+r)\,(-1+h_f+h_m+h_f\,r-h_m\,r)\,s)}\right]\right] \qquad (A1)$$

$$\hat{p} = \mathrm{Min}\left[1, \mathrm{Max}\left[0, \frac{((1+r)\,(2\,h_f\,(-1+r) - 2\,h_m\,(-1+r)^2 + (-2+r)\,r)\,s - (-2+r+r^2)\,t)}{(2\,(-2+2\,h_f - 2\,h_m\,(-1+r)+r)\,(-1+r^2)\,s)}\right]\right] \qquad (A2)$$

553   To understand the relative impacts of changing h_f and h_m on $\hat{p}$, we can take the partial derivative

554   of the above equations first with respect to h_f and then with respect to h_m. For fixed selfing, these

555   partial derivatives are as follows:

$$\frac{\partial \hat{p}}{\partial h_f} = -\frac{(1+r)(s+(-1+r)t)}{2(1-r)(-1+h_f+h_m+h_f r - h_m r)^2 s} \tag{A3}$$

$$\frac{\partial \hat{p}}{\partial h_m} = -\frac{s+(-1+r)t}{2(-1+h_f+h_m+h_f r - h_m r)^2 s} \tag{A4}$$

556   With simple arranging, it can be shown that

$$\frac{\partial \hat{p}}{\partial h_f} = \frac{(1+r)}{(1-r)}\frac{\partial \hat{p}}{\partial h_m} \tag{A5}$$

In a mixed mating system where $0 < r < 1$, the far-left term of the right-hand side of Eq. A5 is

always greater than 1 and increasing as selfing rate increases, meaning an increase in h_f always

yields a greater change in $\hat{p}$ than a comparable increase in h_m. This is expected, as under fixed

selfing, the cost to male diploid fitness is only paid through outcrossing, as we assume all plants

produce enough pollen to self-fertilise. Conversely, diploid fitness effects on female function are

recognised both in selfing and outcrossing. Thus, the effects of changing female and male

dominance terms are only equivalent under complete outcrossing ($r = 0$) and Eq. A5 reduces to

$\frac{\partial \hat{p}}{\partial h_f} = \frac{\partial \hat{p}}{\partial h_m}.$

We can also compare the effect of increasing h_f relative to a shared dominance coefficient

h. Returning to Eq. A3, it is evident that the denominator is always positive, while the sign of the

numerator changes from positive to negative when r switches from less than $\frac{-s+t}{t}$ to more than

$\frac{-s+t}{t}$. This particular value, $\frac{-s+t}{t}$, gives the selfing rate r at which $\hat{p}$ is 0.5. Looking specifically

at the effects of dominance, the magnitude of the slope of $\hat{p}$ in response to r can be controlled by

adjusting h_f in the denominator. Increasing the value of h_f increases the magnitude of the

denominator, thus yielding a more positive slope when $r < \frac{-s+t}{t}$ and a more negative slope when

$r > \frac{-s+t}{t}$. This effect is visualised in Figure A1. Compared to $h_f = h_m = 0$, increasing h_f to 0.25

allows p to remain fixed at higher selfing rates but also to be lost more quickly, thus contracting

the space for polymorphism. An identical effect, but for space allowing unstable equilibrium, is

observed when all dominance coefficients are above 0.5 (Fig. A2).

576   When h_f increases and the C_s allele is rare, the C_s allele confers a lesser advantage to

female diploid fitness as this advantage is increasingly masked by the C_g allele in heterozygotes

(in which it is predominantly found when rare). As an increase in h_f means the C_s allele enjoys

less of a diploid fitness benefit over the C_g allele when found in heterozygotes, the C_s allele

becomes increasingly dependent on the production of homozygotes to facilitate its invasion.

Thus, as h_f increases, C_g remains fixed up to higher values of r. A similar logic follows for

invasion of the C_g allele. When h_f increases, the C_g allele also suffers in the heterozygote. It thus

requires lower selfing rates so that it can exercise its pollen superiority competitive over the C_s

allele under outcrossing.

585   When h_m increases, qualitatively similar but quantitatively different effects occur.

Grouping h_f terms together in the denominators of Eqs. A3 and A4 produces $h_f(1+r)$, while

grouping h_m terms together produces $h_m(1-r)$. This means that an increase in h_m produces a

smaller change in the size of the denominator than an equivalent change in h_f. This is expected, as

diploid fitness effects through male function are only experienced under outcrossing, unlike

female fitness effects, which are experienced both under selfing and outcrossing. Otherwise, the

logic of how dominance affects invasion of both C_g and C_s alleles described above for h_f also

applies to h_m, but to a lesser magnitude. This difference is visualized in Figure A1, as the lines

depicting $h_f = 0.25$ and $h_m = 0.25$ both remain fixed for p up to higher selfing rates, but also drop

(and thus lose p entirely) at lower selfing rates, when compared to $h_f = h_m = 0$.

Turning to mass-action selfing, the partial derivatives of Eq. A2 with respect to h_f (Eq.

A6) and h_m (Eq. A7) are as follows:

$$\frac{\partial \hat{p}}{\partial h_f} = -\frac{(2 + r - r^2)\,s + (-2 + r + r^2)\,t}{\left(-2 + 2\,h_f - 2\,h_m(-1 + r) + r\right)^2 (-1 + r^2)\,s} \tag{A6}$$

$$\frac{\partial \hat{p}}{\partial h_m} = -\frac{2\,t + (1 + r)\left((-2 + r)\,s - r\,t\right)}{(1 + r)\left(-2 + 2\,h_f - 2\,h_m(-1 + r) + r\right)^2 s} \tag{A7}$$

597   Rearrangement of Eqs. A6 yields:

$$\frac{\partial \hat{p}}{\partial h_f} = \frac{1}{(1-r)}\frac{\partial \hat{p}}{\partial h_m} \tag{A8}$$

As seen under fixed selfing, an increase to h_f will yield greater change to $\hat{p}$ than an equivalent

change to h_m. Interestingly, the ratio of these changes is not the same between fixed and mass-

action selfing. Relative to a change in h_m, an equivalent increase in h_f under mass-action selfing

has a weaker effect on $\hat{p}$ than under fixed selfing, as evidenced by comparing the coefficients of

Eqs. A5 and A8, $\frac{(1+r)}{(1-r)}$ and $\frac{1}{(1-r)}$ respectively (assuming $r > 0$). This can be understood by

considering where diploid selection on male function acts in mass-action selfing. Unlike fixed

selfing, costs to pollen production are paid both through both self-pollen and outcross-pollen, as

both pollen types compete on the same stigma. Differences in pollen competitive ability offset

costs through diploid male fitness, thus allowing an increase in h_m to still have less impact on $\hat{p}$

than an increase in h_f. The lesser effect of altering h_m on $\hat{p}$ can be seen in Figure A1. Compared

to fixed selfing, increasing h_m produces less deviation from baseline series (*i.e.*, those with

equivalent h_m and h_f values).

Appendix B

Models for sporophytic control of pollen competitive ability (Jordan and Connallon 2014)

The relative frequencies of each genotype among the available ovules are given as:

$$x_{0f} = \frac{(1-s)X_0}{\overline{W_f}} \tag{B1}$$

$$x_{1f} = \frac{(1-hs)X_1}{\overline{W_f}} \tag{B2}$$

$$x_{2f} = \frac{X_2}{\overline{W_f}} \tag{B3}$$

615   Where *s* is the selection coefficient acting on female function, *h* is the dominance coefficient and

616   $\overline{W_f}$ is defined as

$$\overline{W_f} = (1-s)X_0 + (1-hs)X_1 + X_2 \tag{B4}$$

617   The relative frequencies of each genotype among the pollen pool are given as:

$$x_{0m} = \frac{X_0}{\overline{W_m}} \tag{B5}$$

$$x_{1m} = \frac{(1-ht)X_1}{\overline{W_m}} \tag{B6}$$

$$x_{2m} = \frac{(1-t)X_2}{\overline{W_m}} \tag{B7}$$

618   Where t is the selection coefficient acting on male function, h is the dominance coefficient and

619    $\overline{W_m}$ is defined as

$$\overline{W_m} = X_0 + (1-ht)X_1 + (1-t)X_2 \tag{B8}$$

620   Using these fitness expressions, the genotype recursion equations are:

$$X_0' = \rho\left[x_{0f} + \frac{1}{4}x_{1f}\right] + (1-\rho)\left[\left(x_{0f} + \frac{1}{2}x_{1f}\right)q_m\right] \tag{B9}$$

$$X_1' = \rho\left[\frac{1}{2}x_{1f}\right] + (1-\rho)\left[x_{0f}(1-q_m) + \frac{1}{2}x_{1f} + x_{2f}q_m\right] \tag{B10}$$

$$X_2' = \rho\left[x_{2f} + \frac{1}{4}x_{1f}\right] + (1-\rho)\left[\left(x_{2f} + \frac{1}{2}x_{1f}\right)(1-q_m)\right] \tag{B11}$$

621   where q_m is the frequency of the C_g allele in the pollen pool:

$$q_m = x_{0m} + \frac{1}{2}x_{1m} \tag{B12}$$

622   In the case of fixed selfing, selfing rate ρ is equivalent to r (i.e., proportion pollen allocated to

623   selfing). Under mass-action selfing, ρ is frequency-dependent and genotype-specific:

$$\rho = \frac{rW_{im}}{rW_{im} + (1-r)\overline{W_m}} \tag{B13}$$

624   where $W_{0m} = 1, W_{1m} = (1-ht)$ and $W_{2m} = 1 - t$ and r is the proportion of pollen dedicated

625   to selfing.

626   Equilibrium allele frequencies for fixed and mass-action selfing assuming sporophytic

627   control were derived using the same methods provided in the main text and are as follows:

$$\hat{p} = \text{Min}[1, \text{Max}\left[0, \frac{-r(1+r)s + (r-2)(r-1)t + 2h(r-1)(s+rs+t-rt)}{2(2h-1)(r-1)(s+rs+t-rt)}\right]] \quad \text{(B14)}$$

$$\hat{p} = \text{Min}[1, \text{Max}\left[0, \frac{rs - (r-2)(r-1)t - 2h(r-1)(s+t-rt)}{2(2h-1)(r-1)(s+t-rt)}\right]] \quad \text{(B15)}$$

628   We can understand the conditions under which the C_g allele will perform better under

629   sporophytic control relative to haploid and *vice versa* by taking the difference between Eqs. 10

630   (main text) and B16 as well as Eqs. 11 (main text) and B15. Under fixed selfing:

$$\Delta\hat{p} = \frac{(1-r)(s-t)t}{2(2h-1)s(s(1+r)+t(1-r))} \quad \text{(B16)}$$

In Eq. B16, when $h < 0.5$, the denominator is always negative. The sign of the numerator is

dependent on the size of s relative to t. When diploid selection s is larger than haploid selection t,

the numerator is positive and thus $\Delta\hat{p}$ is negative, revealing the C_g allele is more successful under

sporophytic than gametophytic genetic control. Inversely, when $s < t$ (as presented in the main

text), $\Delta\hat{p}$ is positive and C_g reaches a higher frequency under gametophytic control.

Under mass-action selfing:

$$\Delta\hat{p} = \frac{t((r^2+r-2)t - (2r^2-r-2)s)}{2(2h-1)(r-2)(1+r)s(s+t-rt)} \quad \text{(B17)}$$

When $h < 0.5$ and $0 \leq r \leq 1$, the denominator of B17 is negative. Thus, the sign of $\Delta\hat{p}$ is

dependent on the sign of the numerator. Given $0 \leq r \leq 1$, (r^2+r-2) is always negative and

$(2r^2-r-2)$ is always positive, meaning the numerator is always negative and $\Delta\hat{p}$ is always

positive. In other words, when $h < 0.5$, the C_g allele will always do better under gametophytic

control than under sporophytic control. Conversely, when $h > 0.5$, the denominator is positive

while the numerator remains unchanged. In such a case, $\Delta\hat{p}$ is always negative and the C_g allele

reaches a higher equilibrium frequency under sporophytic control. These analytical results are

visualised in Figure 3, in which, when $h = 0$, the equilibrium allele frequency $\hat{p}$ is higher for

gametophytic (i.e., haploid) control relative to sporophytic and, when $h = 1$, the repelling allele

frequency below which C_g cannot invade is higher under sporophytic control (meaning fixation

requires a higher initial allele frequency). This result can be understood easily by considering the

haplotype-phenotype correspondence under both gametophytic and sporophytic genetic control.

Under gametophytic control, this correspondence is complete (i.e., phenotype reflects haplotype

and the Cg allele always confers a benefit to pollen). Under sporophytic control, this

correspondence is a function of the paternal genotype (homozygote or heterozygote) and the

dominance coefficient h, and thus, the Cg allele will not always confer its maximum advantage

to pollen ability, weakening its invasion ability.

Likewise, invasion conditions for fixed and mass-action selfing assuming sporophytic

control were derived using the methods found in the main text. Under fixed selfing, maximum

and minimum selfing rates permitting invasion of C_g and C_s alleles respectively are given as:

$$r < \frac{s + (3 - 4h)t - \sqrt{(1 - 4h)^2 s^2 + 2(7 - 12h)st + t^2}}{2(2h - 1)(s - t)} \tag{B18}$$

$$r > -\frac{-s + (1 - 4h)t + \sqrt{(3 - 4h)^2 s^2 + 2(12h - 5)st + t^2}}{2(2h - 1)(s - t)} \tag{B19}$$

657   The corresponding expressions for mass-action selfing are:

$$r < \frac{(2h - 1 + 2h)s + (4h - 3)t + \sqrt{(1 - 2h)^2 s^2 + 6(1 - 2h)st + t^2}}{2(2h - 1)t} \tag{B20}$$

$$r < \frac{-s + 2hs + (4h - 1)t - \sqrt{(1 - 2h)^2 s^2 + 6(1 - 2h)st + t^2}}{2(2h - 1)t} \tag{B21}$$

659 **Appendix C**

660 *Production of heterozygote offspring via selfing with variable pollen competitiveness*

661 With a simple Punnet square, it can be demonstrated that even with differential competitive

662 ability between pollen haplotypes, selfing of heterozygotes produces ½ heterozygote offspring:

	C_g **pollen** $\frac{1}{2-t}$ *fertilisation success given 50% allele frequency*	C_s **pollen** $\frac{1-t}{2-t}$ *fertilisation success given 50% allele frequency*
C_g **ovule** *0.5 allele frequency*	$\frac{1}{2(2-t)} C_g C_g$	$\frac{1-t}{2(2-t)} C_g C_s$
C_s **ovule** *0.5 allele frequency*	$\frac{1}{2(2-t)} C_s C_g$	$\frac{1-t}{2(2-t)} C_s C_s$

TABLE C 1

Together, the frequency of heterozygote offspring produced via heterozygote selfing is $\frac{1}{2(2-t)} +$

$\frac{1-t}{2(2-t)}$, which reduces to ½.

Appendix D

Analytical results for mass-action selfing with pollen discounting

Pollen discounting, i.e., loss of pollen for outcrossing during transport, can be incorporated by

inclusion of an additional term, *d,* which gives the proportion of pollen lost and is identical to the

pollen discounting term included in Jordan and Connallon (2014). Equations 5a-c are thus

modified to be:

$$X_0' = x_0 \left[\frac{rW_0 + (1-r)(1-d)\left(x_0 + \frac{1}{2}x_1\right)}{P_0} \right]$$
$$+ \frac{1}{2}x_1 \left[\frac{r\frac{1}{2}W_1 + (1-r)(1-d)\left(x_0 + \frac{1}{2}x_1\right)}{P_1} \right] \tag{D1}$$

$$X_1' = \frac{1}{2}x_1 + x_0 \left[\frac{(1-r)(1-d)\left(x_2 + \frac{1}{2}x_1\right)(1-t)}{P_0} \right] + x_2 \left[\frac{(1-r)(1-d)\left(x_0 + \frac{1}{2}x_1\right)}{P_2} \right] \tag{D2}$$

$$X_2' = x_2 \left[\frac{rW_2(1-t) + (1-r)(1-d)\left(x_2 + \frac{1}{2}x_1\right)(1-t)}{P_2} \right]$$

$$+ \frac{1}{2}x_1 \left[\frac{r\frac{1}{2}W_1(1-t) + (1-r)(1-d)\left(x_2 + \frac{1}{2}x_1\right)(1-t)}{P_1} \right] \tag{D3}$$

672   Similarly, Eqs. 6a-c are also modified:

$$P_0 = rW_0 + (1-r)(1-d)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right) \tag{D4}$$

$$P_1 = r\frac{1}{2}W_1(2-t) + (1-r)(1-d)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right) \tag{D5}$$

$$P_2 = rW_2(1-t) + (1-r)(1-d)\left(x_0 + \frac{1}{2}x_1(2-t) + x_2(1-t)\right) \tag{D6}$$

673   Using Eqs. D1-6, and assuming weak selection, we can derive the equilibrium frequency of the

674   C_g allele. For the full expression, see the corresponding online supplemental materials. To

675   understand the effects of increasing d on $\hat{p}$, we take the partial derivative of the $\hat{p}$ expression

676   with respect to d:

$$\frac{\partial \hat{p}}{\partial d} = -\frac{r(-2 + d^2(-2+r)(-1+r)^2 + r - 3r^2 - 2d(-1+r)(2 + (-1+r)r))t}{2(-1+2h)(-2 - 2d(-1+r) + r)\wedge 2(1 + d(-1+r) + r)\wedge 2 s} \tag{D7}$$

677   To get a better handle on the above expression, we can isolate and rearrange the terms in which d

678   appears to:

$$d(r-1)((r-2)*(2(r+1) - d*(r-1)) + 8) \tag{D8}$$

679   The term $d(r-1)$ is always negative (when $0 < r < 1$), so how d affects the equilibrium allele

680   frequency is dependent on the sign of the following term:

$$((r-2)*(2(r+1)-d*(r-1))+8) \tag{D9}$$

681   With further rearrangement, it can be shown that:

$$\frac{\partial \hat{p}}{\partial d} = \begin{cases} < 0, & d < 2 + \dfrac{8}{r-2} - \dfrac{4}{r-1} \\[2mm] > 0, & d > 2 + \dfrac{8}{r-2} - \dfrac{4}{r-1} \end{cases} \tag{D10}$$

As both d and r are constrained to between 0 and 1, $\frac{\partial \hat{p}}{\partial d}$ is always positive and thus an increase in

d always lowers the equilibrium frequency of the C_g allele. This is not surprising as inclusion of

a pollen discounting term $d > 0$ should decrease the relative proportion of outcross- versus self-

pollen that a stigma receives, therefore decreasing the realised competitive advantage of C_g

pollen, under the mass-action selfing model.

In other words, under mass-action selfing, intermediate r values indicate a stigma

receives both self and outcross pollen. Successful invasion and fixation of either the C_g or C_s

allele depends on the ratio of self to outcross pollen produced, with higher values of r favouring

the C_s allele. Pollen discounting essentially alters this ratio, allowing the C_s allele to invade and

fix at lower values of r, which thus implies limited invasion of the C_g allele.

Figure D1 displays this effect of pollen discounting on the success of the C_g allele using t

$= 0.1$, $s = 0.05$ and d values of 0, 0.2 and 0.5. Indeed, as d is increased, the equilibrium frequency

of the C_g allele becomes increasingly left-shifted, implying a decreased range of selfing rates at

which it can successfully invade.

Appendix E

Numerical validation and consideration of strong selection

Equilibrium and repelling allele frequencies presented in the main text are entirely analytical,

i.e., derived using Equations 10 and 11. Confirmation of these analytically derived frequencies

was done numerically. All numerical results were produced using RStudio Version 1.1.353, in

which we generated and recorded genotype and allele frequencies over generations for different

combinations of selfing, diploid fitness and pollen competitiveness. For invasion from X_0 and X_2

boundaries, initial frequency of the invading allele was always set to 0.0001, and initial genotype

frequencies were at Hardy-Weinberg equilibrium. Equilibrium allele frequencies presented in

Figures E1 were evaluated at selfing rates r ranging from 0 to 1 in increments of 0.01. Long-term

allele frequencies were taken from the 50,000th generation and verified by a change in allele

frequency, i.e., Δp, of less than 10^{-6}. We consider these long-term allele frequencies to be

equilibrium frequencies, as we assume that, with such significantly small changes in allele

frequencies, drift will dominate, and these long-term frequencies are good approximations of true

equilibrium frequencies, where $\Delta p = 0$. As both models are deterministic, numerical results are

the outcome of a single run.

714    As stated in the main text, analytical solutions assume weak (s, $t \ll 1$). However, we

715 found that analytical results under relatively strong selection still provided a qualitatively good

716 fit to their corresponding numerical results under both haploid and diploid control of pollen

717 competitive ability (grey series, Figure E1), suggesting interpretations presented in the main text

718 are broadly applicable across a wide range of selection strengths. In Figure E1, weak selection

719 indicates $s = 0.01$ and $t = 0.03$ strong selection indicates $s = 0.05$ and $t = 0.1$.

721                     FIGURE E 1

722    Importantly, when $h = 1.0$, drops in equilibrium allele frequencies from 1 to 0 for all

723 combinations of selection strength and selfing mode are immediate, i.e., there are no

intermediate equilibrium values (i.e., $0 < \hat{p} < 1$). Any slopes are the result of the increment

chosen for the x axis ($0 \leq r \leq 1$ by 0.01).

In addition to a comparison with numerical results under strong selection, we can also

compare invasion conditions under fixed selfing assuming weak selection to conditions derived

without assuming weak selection. For example, Eqs. 7a and 7b in the main text give maximum

and minimum selfing rates for C_g and C_s invasion respectively under weak selection. Without

this assumption, these maximum and minimum r values are:

$$r < \frac{hs(-2+t)+t}{-(1+h(-2+s)s+(-1+s)(1-hs)t}$$ (E1)

$$r > \frac{(-1+s)(s(2+h(-2+t))-t)}{hs^2(-1+t)+t-s(1+h(-2+t)+t)}$$ (E2)

731   Comparison with expressions 7a and 7b in the main text can be easily considered visually

732   (Figure E2), in which series were generated using $t = 0.1$.

734   FIGURE E 2

Perhaps most notable in Figure E2 is the small space permitting polymorphism when $h = 0.5$ at

low values of r. This result is consistent with Figure E1, in which under strong selection there are

true numerically-derived intermediate equilibrium allele frequencies (i.e., $0 < \hat{p} < 1$) when $h =$

0.5, unlike under weak selection. Overall, we find strong qualitative agreement between invasion

conditions predicted both with and without weak selection, suggesting our results likely hold for

a broader range of selection parameters than those presented in the main text.

Appendix F

Critical selfing rates under mass-action selfing

Using the methods described in the main text, the analytical solutions specifying the selfing rates

required for invasion C_g and C_s alleles under mass-action selfing are as follows:

For C_g: $r <$

$$-\frac{s - 4\,h\,s + t}{3\,(-s + 2\,h\,s)} - \frac{\left(2^{1/3}\left(-(s - 4\,h\,s + t)^2 + 3\,(-s + 2\,h\,s)\,(2\,s - 2\,h\,s + t)\right)\right)}{3\,(-s + 2\,h\,s)\left(-20\,s^3 + 42\,h\,s^3 + 84\,h^2\,s^3 - 160\,h^3\,s^3 + 21\,s^2\,t - 60\,h\,s^2\,t + 12\,h^2\,s^2\,t - 15\,s\,t^2 + 42\,h\,s\,t^2 - 2\,t^3 + \sqrt{\left(\left(-20\,s^3 + 42\,h\,s^3 + 84\,h^2\,s^3 - 160\,h^3\,s^3 + 21\,s^2\,t - 60\,h\,s^2\,t + 12\,h^2\,s^2\,t - 15\,s\,t^2 + 42\,h\,s\,t^2 - 2\,t^3\right)^2 + 4\left(-(s - 4\,h\,s + t)^2 + 3\,(-s + 2\,h\,s)\,(2\,s - 2\,h\,s + t)\right)^3\right)}\right)^{1/3}} +$$
$$\frac{1}{3 \times 2^{1/3}\,(-s + 2\,h\,s)}\left(-20\,s^3 + 42\,h\,s^3 + 84\,h^2\,s^3 - 160\,h^3\,s^3 + 21\,s^2\,t - 60\,h\,s^2\,t + 12\,h^2\,s^2\,t - 15\,s\,t^2 + 42\,h\,s\,t^2 - 2\,t^3 + \sqrt{\left(\left(-20\,s^3 + 42\,h\,s^3 + 84\,h^2\,s^3 - 160\,h^3\,s^3 + 21\,s^2\,t - 60\,h\,s^2\,t + 12\,h^2\,s^2\,t - 15\,s\,t^2 + 42\,h\,s\,t^2 - 2\,t^3\right)^2 + 4\left(-(s - 4\,h\,s + t)^2 + 3\,(-s + 2\,h\,s)\,(2\,s - 2\,h\,s + t)\right)^3\right)}\right)^{1/3}$$
(F1)

746 For C_s: $r >$

$$-\frac{-3\,s + 4\,h\,s + t}{3\,s - 6\,h\,s} + \left(2^{1/3}\left(\left(9 - 30\,h + 28\,h^2\right)s^2 + (-9 + 14\,h)\,s\,t + t^2\right)\right)\Big/$$
$$\left(3\,(-1 + 2\,h)\,s\left(\left(54 - 270\,h + 396\,h^2 - 160\,h^3\right)s^3 - 3\left(-9 + 12\,h + 4\,h^2\right)s^2\,t + 3\,(-9 + 14\,h)\,s\,t^2 + 2\,t^3 + 3\sqrt{3}\sqrt{-(s - 2\,h\,s)^2\left(576\,h^4\,s^4 - 32\,h^3\,s^3\,(27\,s - 37\,t) + 12\,h\,s\,t\left(135\,s^2 - 100\,s\,t + 17\,t^2\right) + 4\,h^2\,s^2\left(81\,s^2 - 558\,s\,t + 205\,t^2\right) + t\left(-432\,s^3 + 441\,s^2\,t - 134\,s\,t^2 + 9\,t^3\right)\right)}\right)^{1/3}\right) +$$
$$\frac{1}{3 \times 2^{1/3}\,(-1 + 2\,h)\,s}\left(\left(54 - 270\,h + 396\,h^2 - 160\,h^3\right)s^3 - 3\left(-9 + 12\,h + 4\,h^2\right)s^2\,t + 3\,(-9 + 14\,h)\,s\,t^2 + 2\,t^3 + 3\sqrt{3}\sqrt{-(s - 2\,h\,s)^2\left(576\,h^4\,s^4 - 32\,h^3\,s^3\,(27\,s - 37\,t) + 12\,h\,s\,t\left(135\,s^2 - 100\,s\,t + 17\,t^2\right) + 4\,h^2\,s^2\left(81\,s^2 - 558\,s\,t + 205\,t^2\right) + t\left(-432\,s^3 + 441\,s^2\,t - 134\,s\,t^2 + 9\,t^3\right)\right)}\right)^{1/3}$$
(F2)

747 **Appendix G**

748 *Critical diploid selection coefficients under mass-action and fixed selfing*

749 Taking the partial derivative of in-text Eqs. 8a and 8b with respect to r produces Eqs. G1 and G2

750 respectively, which in turn provide insights into the effect of increasing selfing rate on the

751 conditions permitting invasion:

$$\frac{\partial s}{\partial r} = \frac{\left(-4 + r\left(4 - 2h(r-1)^2(4+r) - r(-5 + r(2+r))\right)\right)t}{(-2 - 2h(r-1) + r)^2(2 + r - r^2)^2}$$
(G1)

$$\frac{\partial s}{\partial r} = \frac{\left(-4 + r\left(4 - 2h(r-1)^2(4+r) + r(-9 + r(2+r))\right)\right)t}{(-2 - 2h(r-1) + r)^2(2 + r - r^2)^2} \tag{G2}$$

Assuming h, s, t and r all fall between 0 and 1, the above expressions are always negative,

implying the critical value of s always decreases as r increases (Figure 1). When $h < 0.5$, the

maximum value of s permitting C_g invasion decreases more slowly as r increases than the

minimum s value permitting C_s invasion. This results in a range of r values in which s is below

the maximum value for C_g invasion and above the minimum value for C_s value, producing the

potential for polymorphism. Conversely, when $h > 0.5$, the minimum s for C_s invasion decreases

more quickly in response to r than the maximum value for C_g invasion. This produces a range of

r values in which neither allele can invade (i.e., unstable equilibrium). Focusing on the effect of

selfing on the C_g allele, increasing r contracts the parameter space allowing invasion of an allele

conferring a benefit to pollen competitiveness.

To compare effects of fixed versus mass-action selfing on invasion, we can compare how

minimum and maximum s values permitting invasion of C_s and C_g alleles respond to increases in

r in these two selfing systems. Below are expressions for how the maximum and minimum s

values change in response to r under fixed selfing:

$$\frac{\partial s}{\partial r} = -\frac{t}{(-2\,h\,(-1+r)+r)^2} \tag{G3}$$

$$\frac{\partial s}{\partial r} = -\frac{t}{(-2-2\,h\,(-1+r)+r)^2} \tag{G4}$$

We can easily visualise differences between Eqs. G1,2 and G3,4, and thus how critical s values

respond to r under mass-action versus fixed selfing, by considering the case in which $h = 0.25$

and $t = 0.03$ (Figure G1).

FIGURE G 1

For both maximum and minimum s values, it is clear that the critical s value decreases more

rapidly for fixed selfing than mass-action as r increases from 0, as evidenced by the initially

lower (*i.e.*, more negative) values for the fixed $\frac{\partial s}{\partial r}$ series than for mass-action series. This

difference implies that for a given s value, the critical r value allowing C_g invasion will be lower

than that of mass-action selfing. Indeed, we find this to be true.

FIGURE G 2

For a given *s* value (in this case 0.01, as given by the green series in Figure G2), both maximum

and minimum *s* values permitting invasion are right-shifted along the *r* axis. This indicates the C_g

allele can be maintained at higher selfing rates under mass-action selfing than fixed selfing.

Returning to Figure G1, the difference in $\frac{\partial s}{\partial r}$ between fixed and mass-action selfing for a given *r*

value is greater when considering the maximum *s* permitting C_g invasion than the minimum *s*

permitting C_s invasion. While the minimum *s* for C_s invasion is right-shifted under mass-action

selfing compared to fixed selfing, the magnitude of this shift is less than that of the

corresponding maximum *s* values. Put together, this results in a contracted range of *r* values

permitting polymorphism, which can be seen in Figure G2 as the space in which the minimum *s*

series falls above the s = 0.01 series and the maximum *s* series falls below ($0.69 < r < 0.77$ under

mass-action selfing; $0.56 < r < 0.73$ under fixed selfing).

A similar analysis when $h > 0.5$ yields qualitatively the same results but with unstable

equilibria, not protected polymorphism.

803   **Figures**

FIGURE 2

FIGURE 3

810   FIGURE 4

FIGURE 5