Within host selection for faster replicating bacterial symbionts

Ewa Chrostek[1,2] and Luis Teixeira[1,3]*

1 Instituto Gulbenkian de Ciência, Oeiras, Portugal

2 Current Address: Max Planck Institute for Infection Biology, Berlin, Germany

3 Faculdade de Medicina da Universidade de Lisboa, Lisboa, Portugal

 * Corresponding author:

E-mail: lteixeira@igc.gulbenkian.pt

Abstract

Wolbachia is a widespread, intracellular symbiont of arthropods, able to induce reproductive distortions and antiviral protection in insects. *Wolbachia* can also be pathogenic, as is the case with *w*MelPop, a virulent variant of the endosymbiont of *Drosophila melanogaster*. An extensive genomic amplification of the 20kb region encompassing eight *Wolbachia* genes, called Octomom, is responsible for *w*MelPop virulence. The Octomom copy number in *w*MelPop can be highly variable between individual *D. melanogaster* flies, even when comparing siblings arising from a single female. Moreover, Octomom copy number can change rapidly between generations. These data suggest an intra-host variability in Octomom copy number between *Wolbachia* cells. Since *w*MelPop *Wolbachia* with different Octomom copy numbers grow at different rates, we hypothesized that selection could act on this intra-host variability. Here we tested if total Octomom copy number changes during the lifespan

of individual *Drosophila* hosts, revealing selection for different *Wolbachia*

populations. We performed a time course analysis of Octomom amplification in flies

whose mothers were controlled for Octomom copy number. We show that despite the

Octomom copy number being relatively stable it increases slightly throughout *D.*

melanogaster adult life. This indicates that there is selection acting on the intra-host

variation in the Octomom copy number over the lifespan of individual hosts. This

within host selection for faster replicating bacterial symbionts may be in conflict with

between host selection against highly pathogenic *Wolbachia*.

Introduction

Gene copy number variation is one of the mechanisms allowing rapid evolution

across the tree of life [1–3]. In bacteria, growth inhibition by nutrient limitation or

antibiotic presence may be overcome by increasing copy number of genes

functionally related to these challenges [4]. Moreover, amplified genomic regions

allow accumulation of mutations without the risk of loss of the original function. This

can lead to the generation of more beneficial variants and subsequent loss of extra

copies or repurposing of the new copies for a new function [4]. Thus, genomic

amplifications generate extensive and reversible genetic variation, which can either

increase the fitness of an individual directly or be a substrate on which adaptive

evolution can act.

We have previously found that a genomic amplification affects the biology of the

intracellular, maternally transmitted bacterium *Wolbachia* [5]. *Wolbachia* is a

widespread endosymbiont of insects, causing an array of phenotypes, including

reproductive manipulations [6] and antiviral protection [7,8]. Moreover, some

Wolbachia strains can strongly reduce the host lifespan. This was first described for

*w*MelPop, a laboratory *Wolbachia* variant, in *Drosophila melanogaster* [9]. The

Octomom genomic region, which contains eight *Wolbachia* genes, is amplified in

*w*MelPop, while it is present as a single copy in closely related non-pathogenic

Wolbachia variants [10,11]. The number of copies of this region varies greatly

between individual *w*MelPop-infected flies from the same population, ranging from

two to ten copies [5]. We have previously established *D. melanogaster* lines carrying

defined and different Octomom copy numbers and observed that the higher the

Octomom copy number, the higher *Wolbachia* levels and the shorter the lifespan of its

D. melanogaster host [5]. Moreover, a *w*MelPop that reverted to carrying only one

Octomom copy proliferates at the same rate as the control *w*MelCS_b variant and is

not pathogenic [5]. Thus, we identified Octomom copy number as a pathogenicity

determinant of *w*MelPop [5].

The high variation in Octomom copy numbers between individual flies can also

be observed in the progeny of single *w*MelPop-carrying females [5]. This variation

between siblings could be explained by *Wolbachia* variation within a female and

differential symbiont assortment to the progeny. The fact that the variability

decreased under selection argues for initial high variation within single flies, which is

pruned over a few generations of selection for either the highest or the lowest

Octomom copy number. However, even under constant selection some variation was

either maintained or continuously generated, since reversing the direction of the

selection or relaxing it could rapidly change the Octomom copy numbers in these

lines [5].

We hypothesized that variation in Octomom copy number between *Wolbachia*

cells within an individual host could lead to a differential growth of these cells. If

Wolbachia cells with higher Octomom copy number proliferate more, their frequency

in the pool of *Wolbachia* within a host will increase over time, and the average

Octomom copy number of the within-host population increases over the host lifespan.

We tested this hypothesis through a time course analysis of Octomom copy number in

individual *w*MelPop flies originating from mothers with controlled Octomom copy

number.

Results

We examined the stability of Octomom copy number over the adult life (from

eclosion to the onset of high mortality) in single *w*MelPop-carrying *D. melanogaster*

(Fig 1A). The flies were the offspring of parents carrying *w*MelPop with low

Octomom copy number (median of 4.5 Octomom copies, range from 4 to 5.5) or high

Octomom copy number (median of 9.5, range from 8.5 to 11.5). The low Octomom

cohort had, on average, 5.45 (standard deviation, SD = 0.82) Octomom copies per

genome and the high Octomom cohort had an average of 10.29 (SD = 1.90) Octomom

copies per genome. The fitted generalized additive model (GAM) clearly shows an

increase in Octomom copy number in the first few days (six to eight days), which

subsequently levels off at the later timepoints (Fig 1A and B). The trend is highly

significant (the smooth term for time, $p < 0.001$). This shows that Octomom copy

number in the *w*MelPop population increases during most of the adult host lifespan.

Figure 1. Octomom copy number and *Wolbachia* levels during adult *D. melanogaster* life. (A)

Each dot represents *WD0513* genomic levels in a single fly, as we have previously shown that this gene

can be used to estimate Octomom region copy number [5]. The values were obtained using the Pfaffl

method with *wsp* as a reference gene and calibrated using the median of three samples of control

*w*MelCS_b flies (CTR). The lines represent the fit of the generalized additive model (GAM) and

shaded area - the 95% confidence interval. (B) Fitted GAM smooth of Octomom copy number in

response to the age of adults. The 95% confidence interval is indicated by the dashed lines. Note the Y-

axis is standardized so that average is zero. (C) Each dot represents *wsp* levels in a single fly. Lines are

medians of the replicates. *wsp* levels were obtained using Pfaffl method with *Rpl32* as a reference gene

and calibrated using the median of the low copy number samples at time zero. (A) and (C) represent

data from single females derived from the 3rd generation of selection for Octomom copy number in the

DrosDel isogenic *w*[1118] genetic background [5]. Females were raised and kept at 25ºC (survival of their

siblings is shown in Fig. S5A of [5]). Supporting data can be found in S1 Data.

In parallel, we tested *Wolbachia* titer in the same flies. *Wolbachia* levels increase with the age of flies (log-linear mixed-effect model (lme), $p < 0.001$) and flies carrying *w*MelPop with higher Octomom copy numbers have 1.85 times higher *Wolbachia* levels compared to flies carrying *w*MelPop with lower Octomom copy numbers (lme , $p < 0.001$), confirming previous results (Fig 1C). However, we do not see a significant interaction between cohort and growth rate (lme, $p = 0.102$). We also tested if Octomom copy number could be an explanatory variable for *Wolbachia* levels within each cohort. This variable was not significant (lme, $p = 0.48$).

Discussion

By analyzing cohorts of individuals from mothers carrying *Wolbachia* with controlled Octomom copy number we observed that Octomom copy number is relatively stable over the life of the host. However, we detected a small, but clear and statistically significant, increase in Octomom copy number over time in the first six to eight days of the adult *D. melanogaster* life. This could be explained by selection acting on the heterogeneity of *w*MelPop copy number between *Wolbachia* cells within a single host. Since bacteria with higher copy numbers grow faster over time, they contribute more to the total pool of *w*MelPop in an older fly and therefore increase total Octomom copy number. In our dataset, the Octomom copy number stops increasing at the last days of the host life. This is surprising and could be explained by the initial heterogeneity in the flies being reduced in the course of selection. If *Wolbachia* cells with the maximal Octomom copy number within each fly reach a very high frequency or are fixed, there is no genetic variation for selection to act on,

and the Octomom copy number does not continue to increase. Alternatively, the

differential fitness between *Wolbachia* harboring different Octomom copy numbers

may decrease with the age of the fly, weakening the strength of the selection and

preventing continuous Octomom copy number growth.

Importantly, our data indicate that the selective pressures acting on *w*MelPop

within and between hosts are different. Within a host, there may be a selection for

Wolbachia that grow fast. On the other hand, competition between flies could select

for *Wolbachia* that grow slower and have a lower cost for the host [5]. These

opposing selective pressures may play a role in shaping the evolution of *Wolbachia* in

natural populations.

Although the results presented here are compatible with our prediction of within

host heterogeneity and selection, other forces may contribute to the changes in

Octomom copy number during host lifespan. Amplification and deletion of Octomom

copies may also occur throughout *w*MelPop host's adult life. The rates of these two

opposing mutations will determine an overall tendency of Octomom copy number to

increase or decrease over time. Therefore, the dynamics of Octomom copy number

changes may be the combination of selection and mutation. A mutation bias for

deletion of Octomom copies could temper the effect of selection for higher Octomom

copy numbers. On the other hand, a bias for amplification could explain the observed

increase in Octomom copy number, even in the absence of selection. Nonetheless, this

would also result in an increased proportion of more pathogenic *w*MelPop with host

age. The influence of mutation and selection on Octomom copy number change

during host lifespan may vary with intrinsic and extrinsic factors. For instance,

mutation bias towards amplification or deletion could change with Octomom copy

number itself. On the other hand, since temperature has a strong effect on *w*MelPop

pathogenicity [9,12], it may also affect the differential growth of *Wolbachia* with

different Octomom copy numbers and, therefore, within host selection on this trait. A

mechanistic understanding of how Octomom copy number changes and influences

Wolbachia phenotype and which factors modulate it will help disentangling the

relative contribution of selection and mutation to Octomom copy number changes

during *D. melanogaster* life.

Wolbachia with higher Octomom copy number proliferate more and are more

pathogenic to their hosts [5]. The within host selection for bacteria that proliferate

more and have a higher potential of being deleterious to the host may be a common

phenomenon. For instance, *Staphylococcus aureus* variants that cause blood or deep

tissue infection are, in the majority, the result of within-host selection from non-

pathogenic nose colonizing variants [13]. The adaptations conferring high virulence

identified in this study do not favor *S. aureus* dissemination and onward transmission

(discussed in [13]). Thus, the selective pressure acting on the bacteria within a single

host may be in conflict with the selective pressure acting on the entire bacterial

population. This implies that although the more pathogenic bacterial variants arise

throughout the life of the hosts, they are constantly purged from the overall bacterial

population by selection either on the fitness of the host (vertically transmitted

symbionts) or on the bacterial transmission capacity (horizontally transmitted

symbionts).

A recent report suggested that *w*MelPop Octomom copy numbers change

drastically during *D. melanogaster* lifespan, increasing more than two-fold in the first

ten days of adult life and then decreasing more than four-fold over the next thirty days

[14]. These results differ from the ones we present here and may be explained by the

lack of experimental control for the Octomom copy number in these flies and

determination of the Octomom copy numbers after the onset of mortality. Using data

from Chrostek and Teixeira 2015 [5] we constructed a model showing that in a mixed

population of flies the differential growth of *Wolbachia* with different Octomom copy

numbers, combined with differential death of flies carrying *Wolbachia* with different

Octomom copy numbers, leads to initial increase in Octomom copy number, followed

by a decrease due to death of the flies carrying *Wolbachia* with higher Octomom copy

number at the host population level [15].

Here we confirmed that *w*MelPop with higher Octomom copy number has higher

Wolbachia titers, supporting our conclusion that this amplification controls *w*MelPop

levels. However, the difference in growth rate between these lines was not statistically

significant. This may be due to the high variability in the data and the differential

growth between these lines being potentially small. We have previously shown

different growth rates between *w*MelPop carrying one and two copies of Octomom

and between these and *w*MelPop carrying 12 or 15 copies [5]. However, the growth

rate of *w*MelPop carrying 12 and 15 copies was not significantly different [5]. This

indicates that the relationship between growth rate and Octomom copy number is not

linear and that differences in *w*MelPop carrying higher copy numbers may have a

smaller impact on growth. Therefore, the difference in growth between *w*MelPop

carrying five and ten copies may indeed be small. The difference in *Wolbachia* titers

despite the lack of measurable difference in the growth rate might be the result of the

cumulative effect of small growth rate differences throughout the fly development

from egg to adult, the result of a differential growth rate at different development

stages, or even the accumulation of small differences in growth for more than one

generation.

The labile nature of the Octomom amplification and the resulting phenotypes of

this amplification make *w*MelPop an interesting case study to understand genome

dynamics and selective forces acting on endosymbionts. This system may be further

used in the future to reveal general principles in host-bacteria symbiosis.

Material and methods

Fly strains

213   *D. melanogaster* DrosDel isogenic background (*iso*) flies with *w*MelCS_b and

214 *w*MelPop were described before [5,10]. Selection on *D. melanogaster* lines carrying

215 *w*MelPop with different Octomom copy number was described in [5].

Experimental setup for time-course analysis of *WD0513* and *Wolbachia* levels

Female progeny of females from the 3rd generation of selection for Octomom

copy number in the *D. melanogaster* DrosDel isogenic background (*iso*) [5] was

collected at eclosion (ten females per tube), allowed to mate with brothers for 24 h

(five males per tube), separated from males, and 20 females were sacrificed every

second day for *WD0513* and *Wolbachia* density quantification. Females were

maintained at 25°C on a standard cornmeal diet without live yeast and were passed to

fresh vials every 3 days. We sampled only until the onset of high mortality in the

different lines in order to avoid sampling bias for surviving, low Octomom copy

number bearing flies.

DNA extractions

DNA was extracted from single flies (*w*MelPop) or pools of ten flies (*w*MelCS_b controls). Each fly or pool of flies was squashed in 250 µl of 0.1 M Tris HCl, 0.1 M EDTA, and 1% SDS (pH 9.0) and incubated 30 min at 70°C. Next, 35 µl of 8 M CH_3CO_2K was added, and samples were mixed by shaking and incubated on ice for 30 min. Subsequently, samples were centrifuged for 15 min at 13,000 rpm at 4°C, and the supernatant was diluted 100× for qPCR.

Real-time quantitative PCR

The real-time qPCR reactions were performed using CFX384 Real-Time PCR Detection System (Bio-Rad) as described before [5,10]. Each reaction contained 6 µl of iQ SYBR Green Supermix (Bio-Rad), 0.5 µl of each primer (3.6 mM), and 5 µl of diluted DNA. We performed two technical replicates for each sample for each set of primers. Primer sequences were described before [5]. For all three genes assayed: *Wolbachia WD0513* and *wsp,* and *Drosophila Rpl32* the following thermal cycling protocol was applied: 2 min at 50°C, 10 min at 95°C, and 40 cycles of 30 s at 95°C, 1 min at 59°C, and 30 s at 72°C. Melting curves were examined to confirm the specificity of amplified products. Ct values were obtained using Bio-Rad CFX Manager software with default threshold settings. Ct values were subjected to a quality check - samples with standard deviation between technical replicates exceeding 0.5 for one of the genes were discarded. The experiment spanned six qPCR plates and three samples of ten wMelCS_b flies (extracted and aliquoted beforehand and assayed on every qPCR plate) were used to normalize between plates. Relative amounts of genes were calculated by the Pfaffl method [16]. To apply the method, the

efficiency of each primer set was predetermined in a separate experiment. For relative

Octomom copy number quantification, *WD0513* was the target gene and the single-

copy *wsp* gene was used as a reference. The medians of three samples of pools of ten

wMelCS_b flies were used as control values for the Pfaffl method. *w*MelCS_b has

one copy of the Octomom region in the genome, determined by the coverage analysis

of sequencing data [10]. This sample, with known Octomom copy number, is required

to estimate Octomom copy number of the remaining samples [17]. For *Wolbachia*

quantification, *wsp* was the target gene and *Drosophila Rpl32* gene was used as a

reference. The levels of *wsp* are relative to the median of the samples of the low

Octomom cohort at time zero.

264 **Statistical analysis**

The statistical analysis was performed in R [18]. The script of the analysis is

provided in S1 Text. Graphs were generated using the package ggplot2 [19].

Since the temporal trend over time of the number of Octomom copies was not

linear we analyse it by fitting a Generalized Additive Model (GAM, package mgcv in

R [20]). We included time and line as independent variables and PCR plate as a

random effect. The smooth terms for the interaction between time and lines were non-

significant ($p > 0.114$) and were removed form the final model.

Analysis of *wsp* levels over time was performed with log-linear mixed-effect

model fits (package lme4 in R [21]). The effect of interaction between factors was

determined by an ANOVA comparing models fit to the data with and without the

interaction.

Financial Disclosure

LT lab is funded by the Fundação para a Ciência e Tecnologia (www.fct.pt) grant PTDC/BEX-GMG/3128/2014. EC is supported by EMBO Long Term Fellowship EMBO ALTF 1497-2015 (http://www.embo.org), co-funded by Marie Curie Actions by the European Commission (LTFCOFUND2013, GA-2013-609409) (ec.europa.eu/research/mariecurieactions). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgments

We thank Tiago Marques for advice on the statistical analysis.

References

1. Ohno S. Evolution by Gene Duplication. Berlin, Heidelberg: Springer Berlin Heidelberg; 1970. doi:10.1007/978-3-642-86659-3

2. Bergthorsson U, Andersson DI, Roth JR. Ohno's dilemma: evolution of new genes under continuous selection. Proc Natl Acad Sci U S A. 2007;104: 17004–9. doi:10.1073/pnas.0707158104

3. Romero D, Palacios R. Gene Amplification and Genomic plasticity in Prokaryotes. Annu Rev Genet. 1997;31: 91–111. doi:10.1146/annurev.genet.31.1.91

4. Andersson DI, Hughes D. Gene amplification and adaptive evolution in bacteria. Annu Rev Genet. 2009;43: 167–95. doi:10.1146/annurev-genet-102108-134805

5.   Chrostek E, Teixeira L. Mutualism breakdown by amplification of Wolbachia

genes. PLoS Biol. 2015;13: e1002065. doi:10.1371/journal.pbio.1002065

6.   Werren JH, Baldo L, Clark ME. Wolbachia: master manipulators of

invertebrate biology. Nat Rev Microbiol. 2008;6: 741–751.

doi:10.1038/nrmicro1969

7.   Hedges LM, Brownlie JC, O'Neill SL, Johnson KN. Wolbachia and virus

protection in insects. Science. 2008;322: 702. doi:10.1126/science.1162418

8.   Teixeira L, Ferreira A, Ashburner M. The bacterial symbiont Wolbachia

induces resistance to RNA viral infections in Drosophila melanogaster. PLoS

Biol. 2008;6: e2. doi:10.1371/journal.pbio.1000002

9.   Min KT, Benzer S. Wolbachia, normally a symbiont of Drosophila, can be

virulent, causing degeneration and early death. Proc Natl Acad Sci U S A.

1997;94: 10792–6.

10.   Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM,

et al. Wolbachia variants induce differential protection to viruses in Drosophila

melanogaster: a phenotypic and phylogenomic analysis. PLoS Genet. 2013;9:

e1003896. doi:10.1371/journal.pgen.1003896

11.   Woolfit M, Iturbe-Ormaetxe I, Brownlie JC, Walker T, Riegler M, Seleznev A,

et al. Genomic Evolution of the Pathogenic Wolbachia Strain, wMelPop.

Genome Biol Evol. 2013;5: 2189–204. doi:10.1093/gbe/evt169

12.   Reynolds KT, Thomson LJ, Hoffmann AA. The effects of host age, host

nuclear background and temperature on phenotypic effects of the virulent

Wolbachia strain popcorn in Drosophila melanogaster. Genetics. 2003;164:

1027–34.

13-   Young BC, Wu Ch, Gordon NC, Cole K, Price JR, Liu E, et al. Severe

infections emerge from commensal bacteria by adaptive evolution. Elife.

2017;6: e30637. doi: 10.7554/eLife.30637

14.   Rohrscheib CE, Frentiu FD, Horn E, Ritchie FK, Swinderen B Van, Ii MWW,

et al. Intensity of Mutualism Breakdown Is Determined by Temperature Not

Amplification of Wolbachia Genes. PLoS Pathog. 2016; 9: e1005888.

doi:10.1371/journal.ppat.1005888

15.   Chrostek E, Teixeira L. Comment on Rohrscheib et al . 2016 " Intensity of

mutualism breakdown is determined by temperature not amplification of

Wolbachia genes ." PLoS Pathog. 2017; 13: e1006540.

doi:10.1371/journal.ppat.1006540

16.   Pfaffl MW. A new mathematical model for relative quantification in real-time

RT-PCR. Nucleic Acids Res. 2001;29: e45.

17.   D 'haene B, Vandesompele J, Hellemans J. Accurate and objective copy

number profiling using real-time quantitative PCR. Methods. 2010;50: 262–

270. doi:10.1016/j.ymeth.2009.12.007

18.   Team R. A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. 2012.

19.   Wickham H. ggplot2 Elegant Graphics for Data Analysis. Media. 2009.

doi:10.1007/978-0-387-98141-3

20.   Wood SN. Generalized additive models : an introduction with R. Texts Stat

Sci. 2006; xvii, 392 . doi:10.1111/j.1541-0420.2007.00905_3.x

21.   Bates D, Machler, M, Bolker BM, Walker, SC. Fitting Linear Mixed-Effects

Models using lme4. Journal of Statistical Software. 2015;67.

doi:10.18637/jss.v067.i01

Supporting information

352   S1 Data – Relative levels of *WD0513* and *wsp* in single females carrying

353   wMelPop.

355   S1 Text – R script with the statistical analysis of the data.