A perturbation model of the gut microbiome's response to antibiotics

Liam P. Shaw[1,2,3], Chris P. Barnes[4,5], A. Sarah Walker[6], Nigel Klein[3], Francois Balloux[1]

1: UCL Genetics Institute, UCL, London

2: CoMPLEX, UCL, London

3: UCL Institute of Child Health, UCL, London

4: Cell and Developmental Biology, UCL, London

5: Department of Genetics, Evolution and Environment (GEE), UCL, London

6: MRC Clinical Trials Unit at UCL, UCL, London

Abstract

Even short courses of antibiotics are known to reduce gut microbiome diversity. However, there has been limited mathematical modelling of the associated dynamical time-response. Here, we take inspiration from a 'stability landscape' schematic and develop an impulse-response model of antibiotic perturbation. We fit this model to previously published data where individuals took a ten-day course of antibiotics (clindamycin or ciprofloxacin) and were sampled up to a year afterwards. By fitting an extended model allowing for a transition to an alternative stable state, we find support for a long-term transition to an alternative community state one year after taking antibiotics. This implies that a single treatment of antibiotics not only reduces the diversity of the gut flora for up to a year but also alters its composition, possibly indefinitely. Our results provide quantitative support for a conceptual picture of the gut microbiome and demonstrate that simple models can provide biological insight.

Introduction

The human gut microbiome is a complex ecosystem and, as such, can be thought of in ecological terms. The relative stability of the gut microbiome in the absence of large perturbations has been suggested to indicate the presence of restoring forces within a dynamical system (Relman 2012). While stability appears to be the norm, disturbances to this ecosystem are also important when considering the impact of the gut microbiome on human health. One example of a major perturbation is a course of antibiotics, which typically leads to a marked reduction in species diversity before subsequent recovery (Modi *et al.* 2014). Even a brief course of antibiotics can result in long-term effects on microbial community composition, with species diversity remaining lower than its baseline value up to a year afterwards (Zaura *et al.* 2015). However, the nature of the reconstitution of the gut microbiome remains an active area of research.

Artificial perturbation experiments are widely used to explore the underlying dynamics of macro-ecological systems (Wootton 2010). In the context of the gut microbiome, the response after antibiotics has been extensively investigated (Sullivan *et al.* 2001; Dethlefsen *et al.* 2008; Dethlefsen & Relman 2011). However, despite interest in the application of ecological theory to the gut microbiome (Pepper & Rosenfeld 2012) there has been limited quantitative or mechanistic modelling of this response. While this may be because responses can appear individualized (Dethlefsen & Relman 2011), this does not preclude the possibility of generalized models that are applicable at the population level. Additionally, recent work suggests that alterations due to specific antibiotics are predictable and reproducible (Raymond *et al.* 2015).

Applying mathematical models to other ecological systems subject to perturbation has a long tradition of giving useful insight into their behaviour (Skellam 1951; May 1973; Scheffer *et al.* 2001). Crucially, it allows the comparison of different models based on different hypotheses about the subsequent behaviour of the system. Additionally, developing a consistent mathematical framework for quantifying the long-term effects of antibiotic use would facilitate comparisons between different antibiotics and different regimens, with the potential to inform approaches to antibiotic stewardship (Doron & Davidson 2011). Some previous work has attempted to model species interactions in the context of antibiotics using Lotka-Volterra models (Stein *et al.* 2013), but such models require dense temporal sampling and restriction to a small number of species to make meaningful inference, limiting their applicability to broader ecological questions. Furthermore, it has recently been shown that pairwise microbial interactions in different scenarios cannot be captured by a single equation, suggesting that pairwise modelling will often fail to predict microbial dynamics (Momeni *et al.* 2017).

In one popular schematic picture taken from classical ecology, the state of the gut microbiome is represented by a ball sitting in a stability landscape (Holling 1973; Lemon *et al.* 2012; Relman 2012; Lloyd-Price *et al.* 2016). Perturbations can be thought of either as forces acting on the ball to displace it from its equilibrium position (Lloyd-Price *et al.* 2016), or alterations of the stability landscape (Costello *et al.* 2012). While this image is usually provided only as a conceptual model to aid thinking about the complexity of the ecosystem, we used it to derive a mathematical model to investigate whether it could provide mechanistic insight.

The model we outline here, based on simple ecological concepts, allows quantitative hypotheses about the effect of antibiotics on the gut microbiome to be tested. We model the effect of a brief course of antibiotics on the microbial community's phylogenetic diversity as the impulse response of an overdamped harmonic oscillator (Figure 1; see Materials and Methods), and compare parameters for two widely-used antibiotics by fitting to empirical data previously published by Zaura et al. (2015). We find that a variant of the model with an extra parameter accounting for the possibility of an altered equilibrium value of diversity is better supported, providing evidence from a sparse dataset that antibiotics can produce transitions to alternative stable states.

Figure 1. An impulse response model of antibiotic perturbation to the gut microbiome. We represent the gut microbiome as a unit mass on a stability landscape, where height corresponds to phylogenetic diversity. (A) The healthy human microbiome can be conceptualized as resting in the equilibrium of a stability landscape of all possible states of the microbiome. Perturbations can displace it from this equilibrium value into alternative states (adapted from Lloyd-Price et al. (2016)). (B) Choosing to parameterize this stability landscape using diversity, we assume that there are just two states: the healthy baseline state and an alternative stable state. (C) Perturbation to the microbiome (e.g. by antibiotics) is then modelled as an impulse, which assumes the duration of the perturbation is short relative to the overall timescale of the experiment. We consider the form of the diversity time-response under two scenarios: a return to the baseline diversity; and a transition to a different value of a diversity (i.e. an alternative stable state).

Results

An impulse response model for the effect of antibiotics

Our mechanistic model (Figure 1) assumes that a short course of antibiotics can be modelled as an impulse on the gut microbiome. With some additional simplifying assumptions about the form of the stability landscape (see Materials and Methods), we derive an analytical form for this overdamped impulse response in terms of the phylogenetic diversity of the gut microbiome (eq. 6).

We fit the model to published data from Zaura et al. (2015) where 30 individuals received a ten-day course of either a placebo, ciprofloxacin, or clindamycin (Table 1). Clindamycin is a lincosamide with a broad spectrum of activity against Gram-positive aerobes and anaerobes Gram-negative anaerobes (Guay 2007). Ciprofloxacin is a quinolone which targets bacterial DNA topoisomerase and DNA gyrase, making it active against a range of Gram-positive and Gram-negative bacteria (Mustaev *et al.* 2014). Faecal samples were taken at baseline (i.e. before treatment), then subsequently at ten days, one month, two months, four months, and one year after treatment.

Group	n	% males	% Caucasian	Average age, years (SD)	Average weight, kg (SD)	Average height, cm (SD)
Placebo	10	50	100	26 (4)	74 (9)	179 (10)
Ciprofloxacin	10*	50	80	26 (3)	69 (13)	176 (10)
Clindamycin	9**	56	100	24 (5)	67 (11)	175 (9)

Table 1. Demographic data of study participants by treatment group. Adapted from Zaura *et al.* (2015). * On reanalysis after downloading data from SRA Run Selector, we found that participant KI17 was missing 2/6 faecal samples, so they were excluded from analysis i.e. leaving $n = 9$ for in our reanalysis of ciprofloxacin as well as clindamycin. However, these summary statistics apply before the exclusion of KI17. ** One female participant who was initially recruited dropped out of the study after enrolment.

The model appeared to adequately describe the initial response to antibiotics (Figure 2), where diversity decreases (i.e. displacement from equilibrium increases) before returning gradually towards equilibrium. Despite large variability between samples from the same treatment group, reassuringly the placebo group clearly did not warrant an impulse response model whereas data from individuals receiving ciprofloxacin and clindamycin was qualitatively in agreement with the model.

However, the residuals suggested that diversity after a year was not well-captured by the model. In their analysis, Zaura et al. (2015) noted significantly ($p < 0.05$) reduced Shannon diversity when comparing samples a year after receiving 10 days' ciprofloxacin to baseline, but this could have in principle merely been due to slow reconstitution and return to original equilibrium under the dynamics we have described.

Figure 2. An impulse response model captures the dynamics of the effect of antibiotics on the gut microbiome. Bayesian fits with Stan for participants taking either a placebo (n=10), ciprofloxacin (n=9), or clindamycin (n=9). The mean phylogenetic diversity from 100 bootstraps for each sample (black points) and median and 95% credible interval from the posterior distribution (bold and dashed coloured lines, respectively). The grey line indicates the equilibrium diversity value, defined on a per-individual basis relative to the mean baseline diversity. The biased positive skew of residuals after a year suggests the possibility of a transition to an alternative stable state with persistently reduced diversity.

Fitting the impulse model to the data and taking into account the whole temporal response suggests that the lack of return to the initial equilibrium state is not due to slow reconstitution of the initial microbiome species community. Instead, the distribution of residuals indicates that, while the initial response fits a standard impulse response model well, the longer-term dynamics of the system did not − as might be expected under a scenario involving a long-term transition to an alternative community state (Figure 1). We therefore developed a variant of the model (eq. 7) to take into account potential shifts to alternative stable states.

Support for an antibiotic-induced state transition

To test the hypothesis that the course of antibiotics could have moved individuals' gut microbiomes into alternative states, we fit an extended version of our model that allowed a potential non-zero asymptotic value (model 2; eq. 7), representing a new long-term value of diversity. We assumed a normally distributed prior for the asymptote parameter A centred at zero (i.e. return to original equilibrium) with a variance given by the variance of the displacement of placebo samples from baseline after a year.

Qualitatively, this slightly more complex model gave a similar fit (Figure 3) but with a positive displacement
from equilibrium, corresponding to an alternative equilibrium state with lower diversity. We compared
models with the Bayes factor BF, where $BF > 1$ indicates support for one model over another. There was no
support for model 2 over model 1 for the placebo ($BF = 0.96$) but support for ciprofloxacin ($BF = 3.36$)
and clindamycin ($BF = 3.99$). The posterior estimates for the asymptote parameter for ciprofloxacin and
clindamycin were substantially positively skewed (Figure 4), providing evidence of a transition to a state
with lower phylogenetic diversity than the baseline.

Figure 3: A model with a possible state transition improves the fit to empirical data. Bayesian fits with Stan for participants taking either a placebo (n=10), ciprofloxacin (n=9), or clindamycin (n=9). The mean phylogenetic diversity from 100 bootstraps for each sample (black points) and median and 95% credible interval from the posterior distribution (bold and dashed coloured lines, respectively). The grey line indicates the equilibrium diversity value, defined on a per-individual basis relative to the mean baseline diversity. The biased positive skew of residuals after a year suggests the possibility of a transition to an alternative stable state with persistently reduced diversity. The non-zero-centred asymptote indicates support for a state transition.

Comparison of parameters between antibiotics

Comparing the posterior distribution of parameters for model 2 fits between treatment groups (Figure 4), the
strength of the perturbation parameter D was not substantially different between antibiotics. The asymptotic
equilibrium parameter A was positively skewed for both antibiotics (median (95% CI): $A_{\text{clinda}} = 0.66$ (-
0.13-1.41); $A_{\text{cipro}} = 0.58$ (-0.14-1.27),   strongly suggesting persistent detrimental effects on microbiome
diversity and a transition to an alternative stable state.

The parameters b and k were both greater in clindamycin compared to ciprofloxacin. The damping ratio
$\zeta = b/(2\sqrt{k})$ summarises how perturbations decay over time, and is an inherent property of the system
independent of the perturbation itself. Therefore, if our modelling framework and ecological assumptions
were valid we would expect to find a consistent damping ratio across both the clindamycin and ciprofloxacin
groups. This is indeed what we observed, with median (95% CI) damping ratios of ζ_{clinda}=1.07 (1.00-1.65)
and ζ_{cipro}=1.07 (1.00-1.66), substantially different from both the prior and the posterior distribution in the
placebo group of $\zeta_{\text{placebo}} = 1.21(1.00-3.00)$, supporting the view of the gut microbiome as a damped
harmonic oscillator.

Figure 4: Posterior parameter estimates for model with a possible transition to an alternative stable state. The posterior distributions from Bayesian fits of model 2 (eq. 7) to empirical data for ciprofloxacin (green) and clindamycin (red). Each posterior distribution represents 400,000 iterations in total.

129
A complex, individualized antibiotic response does not prevent modelling

While it is not our intention to repeat a comprehensive description of the precise nature of the response for
the different antibiotics, we note some interesting qualitative observations from our reanalysis that highlight
the complexity of the antibiotic response in order to make the point that, while modelling these interactions
is far beyond the scope of our model, our approach is unaffected by this underlying complexity. We discuss
here observations at the level of taxonomic family (Supplementary Figure 1).

Despite their different mechanisms of action, both clindamycin and ciprofloxacin caused a dramatic decrease
in the Gram-negative anaerobes *Rikenellaceae*, which was most marked a month after the end of the course.
However, for ciprofloxacin this decrease had already started immediately after treatment, whereas for
clindamycin the abundance after treatment was unchanged in most participants. The different temporal
nature of this response perhaps reflects the bacteriocidal nature of ciprofloxacin (Mustaev *et al.* 2014)
compared to the bacteriostatic effect of clindamycin, although concentrations *in vivo* can produce
bacteriocidal effects (Spížek & Řezanka 2004).

There were clear differences in response between antibiotics. For example, clindamycin caused a decrease in
the anaerobic Gram-positives *Ruminococcaceae* after a month*,* whereas ciprofloxacin had no effect.

Conversely ciprofloxacin caused lower levels of *Barnesiellaceae* which was largely unaffected by
clindamycin.

Some families appeared unaffected by antibiotics: the *Bacteroidaceae* were largely unaffected in most
individuals. Furthermore, while overall diversity decreased, this can still be consistent with increases in the
relative abundance of certain taxa. For example, ciprofloxacin led to increases in *Erysipelotrichaceae,* which
were dramatic in some individuals. Interestingly, for these individuals these increases coincided with marked
decreases in *Bacteroidaceae*, suggesting the relevance of inter-family microbial interactions (Supplementary
Figure 1). The individualized nature of the ciprofloxacin response was also noticeable in *Lachnospiraceae* –
which was largely unaffected by clindamycin – as its abundance dropped below detectable levels in some
individuals after a month but remained unchanged in other individuals.

Comparing relative abundances at the family level, there were few differences between community states of
different treatment groups after a year. Equal phylogenetic diversity can be produced by different community
composition, and this suggests against consistent trends in the long-term dysbiosis associated with each
antibiotic. However, we did find that *Peptostreptococcaceae*, a member of the order *Clostridiales*, was
significantly more abundant in the clindamycin group when compared to both the ciprofloxacin group and
the placebo group separately ($p < 0.05$, Wilcoxon rank sum test). In a clinical setting, clindamycin is well-
established to lead to an increased risk of a life-threatening infection caused by another member of
Clostridiales: Clostridium difficile (Thomas *et al.* 2003). The long-term reduction in diversity may well
similarly increase the risk of colonization and overgrowth of pathogenic species.

Discussion

Starting from a common qualitative conceptual picture of the gut microbiome as resting within a stability
landscape, we have developed a simple mathematical model of its response to perturbation. With a few
simplifying ecological assumptions, most notably that the phylogenetic diversity of the gut microbiome
relative to its baseline value in some way parameterises this stability landscape, we have demonstrated that
the response of the gut microbiome to a short course of antibiotics can be modelled as an impulse acting on a
damped harmonic oscillator. Crucially, the simplifications involved appear to be justified at some
fundamental level, as this model proves to successfully capture dynamics of empirical data. From this, we
suggest that the restoring forces that contribute to the gut microbiome's resilience to perturbation are
proportional to displacement from equilibrium and that the system is overdamped.

Our approach uses a simple conceptual model to give mechanistic insight. Zaura et al. (2015) made the
observation from their dataset that the lowest diversity was observed after a month rather than immediately
after treatment stopped. This cannot be due to a persistence of the antibiotic effect, as clindamycin and
ciprofloxacin only have short half-lives of the order of hours (Leigh 1981; Bergan *et al.* 1987). Our model
gives us a mechanistic framework for thinking about this temporal delay: the full effects of the transient
impulse take time to be realized due to the overdamped nature of the system, and we found a consistent
damping ratio for both antibiotics analyzed.

We have also demonstrated how this modelling framework could be used to compare different hypotheses
about the long-term effect of antibiotic perturbation on the gut microbiome by fitting different models and
using Bayesian model selection. Our modelling work provides an additional line of evidence that while
short-term restoration obeys a simple impulse response model, the underlying long-term community state
can be fundamentally altered by a brief course of antibiotics, as suggested previously by others (Dethlefsen
& Relman 2011), raising concerns about the long-term impact of antibiotic use on the gut microbiome.
Despite the noisiness of the dataset and reliance on uninformative priors, we still found evidence that a
model with a state transition was better supported, which was not observed in individuals taking a placebo.
The transition to a new state with reduced diversity may increase the risk of colonization and overgrowth of
pathogenic species. Even if only marginal, when considered at a population level this may mean that

antibiotics have substantial negative health consequences that could support reductions in the length of
antibiotic courses, in addition to concerns about antibiotic resistance (Llewelyn *et al.* 2017). Modelling the
long-term impact on the microbiome of different doses and courses could help to influence the use of
antibiotics in routine clinical care.

While the evidence for a long-term state transition is weak at present, we can at the very least conclude that
the restoration of diversity after a year does not seem to obey the same underlying dynamics that govern the
initial response, even if we remain agnostic about the most appropriate model refinement. This disparity
between the short- and long-term time-evolution of the system is relevant to the distinction between different
definitions of resilience. Implicit in some definitions of ecological resilience is the assumption that the
fundamental shape of the stability landscape remains unaltered (Gunderson 2000), which we also adopt here,
but it is possible that this assumption is invalid and should also be explicitly modelled.

Our sample size is small so the precise posterior estimates for parameters that we obtain should not be over-
interpreted, but comparing antibiotics using these estimates represents another practical application of such
simple models. However, these posterior estimates for the model parameters were fairly wide, which is to be
expected with a sparse and small dataset. Hierarchical mixed effects models may offer an improved fit,
particularly if they take into account other covariates; however, here we lacked metadata on the participants
from the original study (Table 1).

A single metric clearly fails to capture all the complexity of the microbial community and its interactions.
Nevertheless, the observation that treating phylogenetic diversity as the 'height' in the stability landscape
leads to a reasonable fit of a simple model is interesting, as it supports observations of functional redundancy
in the gut microbiome (Turnbaugh *et al.* 2007). An interesting extension of this work would be to
systematically fit the model to a variety of diversity metrics and assess the model fit to see which metric, or
combination of metrics, is most appropriately interpreted as the state variable parameterizing the stability
landscape. A possible complementary approach could consider the diversity of the gut resistome (van Schaik
2015).

We would not expect the behavior with longer or repeated courses of antibiotics to be well-described by an
impulse response model, but it would be possible to use the mathematical framework given here to obtain an
analytic form for the possible system response by convolving any given perturbation function with the
impulse response. It remains to be seen whether this simple model would break down in such circumstances.

The detailed nature of the gut microbiome's response to clindamycin and ciprofloxacin was individualized in
our dataset, as others have also observed with shotgun sequencing of samples from healthy participants given
a second-generation cephalosporin (Raymond *et al.* 2016). We believe it would be a mistake to react to this
complexity by assuming that no simplified model can capture general details of the ecosystem. At this stage
of our understanding, creating a comprehensive inter-species model of the hundreds of members of the gut
microbiome appears intractable. We recommend that microbiome research instead starts with ecologically-
informed simple models and believe there is a place for both 'bottom-up' models using pairwise interactions
for systems of reduced complexity like bioreactors, and 'top-down' models using general ecological
principles, as we have attempted to demonstrate here.

We have shown that comparing different hypotheses about the response of the gut microbiome to antibiotics
is possible by using a simple model derived from minimal assumptions about the nature of its equilibrium
diversity. Future mathematical models of the gut microbiome, in conjunction with carefully designed
longitudinal studies, will offer many more opportunities to rigorously test ecological hypotheses.

Materials and methods

Ecological assumptions

We represent the state of the gut microbiome as a unit mass resting in a stability landscape (Figure 1A). Choosing to mathematically model the state of the gut microbiome in this way also requires choosing a mathematical representation with reference to an equilibrium value. While earlier studies sought to identify a core set of 'healthy' microbes, the disturbance of which would indicate displacement from equilibrium, it has become apparent that this is not a practical definition due to high inter-individual variability in taxonomic composition (Lloyd-Price *et al.* 2016). More recent concepts of a healthy 'functional core' appear more promising, but characterization is challenging, particularly as many gut microbiome studies use 16S rRNA marker gene sequencing rather than whole-genome shotgun sequencing.

Therefore, we choose to use a metric that offers a proxy for the general functional potential of the gut microbiome: phylogenetic diversity (Lloyd-Price *et al.* 2016). Higher diversity has previously been associated with health (Turnbaugh *et al.* 2007) and temporal stability (Flores *et al.* 2014). For these reasons, we assume the equilibrium position to have higher diversity than the points immediately surrounding it, forming a potential well (Figure 1B). However, there may be alternative stable states that represent possible 'dysbiotic' states (Figure 1B), which are of interest when considering the effect of perturbations (Figure 1C).

The model

We treat the local stability landscape as a harmonic potential, with a 'restoring' force proportional to the displacement x from the equilibrium position ($-kx$). We also assume the presence of a 'frictional' force acting against the direction of motion ($-b\dot{x}$). This system is equivalent to a damped harmonic oscillator (Riley *et al.* 1997) with the following equation of motion:

(1) $\quad \dfrac{d^2x}{dt^2} + b\dfrac{dx}{dt} + kx = 0$

Additional forces acting on the system now appear on the right-hand side of this equation as perturbations. Consider a course of antibiotics of duration τ. If we are interested in the behaviour of the system at timescales $T \gg \tau$, we can assume for simplicity that this perturbation is of infinitesimal duration and model it as an impulse of magnitude D acting at time $t = 0$:

(2) $\quad \dfrac{d^2x}{dt^2} + b\dfrac{dx}{dt} + kx = D\delta(t)$

To solve this second order differential equation, we assume that $b^2 > 4k$ (the 'overdamped' case) based on the lack of any oscillatory behaviour previously observed in the gut microbiome, to the best of our knowledge. Then, subject to the initial conditions $x(0^+) = 0$ and $\dot{x}(0^+) = D$ we obtain the following equation describing the system's trajectory:

(3) $\quad x(t) = \dfrac{D}{2\cdot\sqrt{\left(\frac{b}{2}\right)^2 - k}}\left(e^{-\left(\frac{b}{2} - \sqrt{\left(\frac{b}{2}\right)^2 - k}\right)t} - e^{-\left(\frac{b}{2} + \sqrt{\left(\frac{b}{2}\right)^2 - k}\right)t}\right)$

Fitting the model therefore requires fitting three parameters: b (the damping on the system), k (the strength of the restoring force), and D (how strong the perturbation is). For the purposes of fitting the model, we choose to reparameterise the model using the following definitions:

(4) $\quad b = e^{\phi_1} + e^{\phi_2}$

(5) $\quad k = e^{\phi_1 + \phi_2}$

Resulting in the following model (Model 1, Figure 1C):

271   **(6)** $x_1(t) = \frac{De^{\phi_1}e^{\phi_2}}{e^{\phi_2}-e^{\phi_1}} \cdot \left(e^{-e^{\phi_1}t} - e^{-e^{\phi_2}t}\right)$

272   Antibiotics may lead not just to displacement from equilibrium, but also state transitions to new equilibria
273   (Modi *et al.* 2014). To investigate this possibility, we also consider a model where the value of equilibrium
274   diversity asymptotically tends to a new value *A* (Model 2, Figure 1C).

275   **(7)** $x_2(t) = \frac{De^{\phi_1}e^{\phi_2}}{e^{\phi_2}-e^{\phi_1}} \cdot \left(e^{-e^{\phi_1}t} - e^{-e^{\phi_2}t}\right) + A \cdot \left(1 - e^{-e^{\phi_1}t}\right)$

Empirical dataset

To validate our model and test whether antibiotic perturbation caused a state transition we fitted both models
to an empirical dataset and compared the results. Zaura *et al.* (2015) conducted a study on the long-term
effect of antibiotics on the gut microbiome which provides an ideal test dataset. As part of this study, 30
Swedish individuals (15 males and 15 females, average age 26 years, range 18-45 years) were randomly
assigned to either ciprofloxacin, clindamycin, or a placebo. The antibiotics (150 mg clindamycin four times a
day, 500 mg ciprofloxacin twice a day) and placebo were administered for $\tau = 10$ days and longitudinal
faecal samples collected until $T = 1$ year afterwards (i.e. $\frac{\tau}{T} \sim 0.027 \ll 1$) at baseline, after treatment, one
month, two months, four months, and one year. Samples underwent 16S rRNA gene amplicon sequencing,
targeting the V5-V7 region (SRA: SRP057504). We reanalysed this data, doing de novo clustering into
operational taxonomic units (OTUs) at 97% similarity with VSEARCH v1.1.1 (Rognes *et al.* 2016) with
chimeras removed against the 16S gold database (http://drive5.com/uchime/gold.fa). Taxonomy was
assigned with RDP (Wang *et al.* 2007).

Phylogenetic diversity

There are many possible diversity metrics that could be used to compute the displacement from equilibrium.
Because of our assumption that phylogenetic diversity approximates functional potential, which is itself a
proxy for ecosystem 'health' (see 'Ecological assumptions'), we chose to use Faith's phylogenetic diversity
(Faith 1992) calculated with the `pd()` function in the 'picante' R package v1.6-2 (Kembel *et al.* 2010).
Calculating Faith's phylogenetic diversity requires a phylogeny, which we produced with RaxML v8.1.15
(Stamatakis 2014) after aligning 16S rRNA V5-V7 OTU sequences with Clustal Omega v1.2.1 (Sievers *et
al.* 2011). To obtain values for fitting the model, we used mean bootstrapped values ($n = 100$, sampling
depth $r = 2000$) of phylogenetic diversity d_i relative to the baseline phylogenetic diversity d_0 for each
individual, representing the displacement from equilibrium in our model:

(8) $\bar{d}_i = d_i - d_0$

Model fitting

We used a Bayesian framework to fit models 1 and 2 (eq. 6 and 7) using Stan (Carpenter *et al.* 2017) and
RStan (Stan Development Team 2017) to the three separate groups: placebo, ciprofloxacin, and clindamycin.
In brief, our approach used 4 chains with a burn-in period of 10,000 iterations and 100,000 subsequent
iterations, verifying that all chains converged ($\hat{R} = 1$) and the effective sample size for each parameter was
sufficiently large ($n_{eff} > 10,000$).

We used uninformative priors for the three parameters in the original model 1 without a state transition (eq.
6). For ciprofloxacin and clindamycin we used the same uniformly distributed prior for *D*, and uniform
priors for ϕ_1, ϕ_2. For model 2 with a state transition (eq. 7) we used the same priors, with a normal prior
centred at zero for the new equilibrium value *A* with a standard deviation given by the standard deviation of
the displacement of placebo samples from baseline after a year, with bounds between -2 and 2. The priors
are as follows:

(9.1) $D \sim \text{uniform}(0, 15)$

(9.2) $\phi_1 \sim \text{uniform}(-1.99, 1.99)$

(9.3) $\phi_2 \sim \text{uniform}(-2, 2)$

(9.4) $A \sim \text{normal}(0, 1.263)$

For the placebo group, we expected no perturbation response so used a uniform prior for D centred at zero:

(10) $D \sim \text{uniform}(-5, 5)$

We compared models 1 and 2 for each treatment group using the Bayes factor (Aitkin 1991; Kass & Raftery
1995) after extracting the model fits using bridge sampling with the bridgesampling R package v0.2-2
(Gronau *et al.* 2017). A prior sensitivity analysis showed that choice of priors did not affect the conclusion
that model 2 outperformed model 1 for the two antibiotics, although the strength of the Bayes factor varied.

Full code for fitting the models to empirical data is available as a zipped archive (Supplemental Code 1).

Acknowledgements

Author contributions

325   LPS conceived the model, performed analyses, and wrote the paper. All authors contributed to discussion of
326   the model and gave comments on the paper.

Data accessibility statement

328   Datasets and code necessary to reproduce the results and figures are available as Supporting Information. All
329   sequence data reported in this paper has been previously deposited in the NCBI Sequence Read Archive as
330   part of another publication (SRA accession SRP057504).

Funding

332   LPS is supported by the Engineering and Physical Sciences Research Council [EP/F500351/1] and the
333   Reuben Centre for Paediatric Virology and Metagenomics. CPB is supported by the Wellcome Trust
334   [097319/Z/11/Z].

References

1.Aitkin, M. (1991). Posterior Bayes Factors. *J. R. Stat. Soc. Ser. B*

2.Bergan, T., Thorsteinsson, S.B., Solberg, R., Bjornskau, L., Kolstad, I.M. & Johnsen, S. (1987). Pharmacokinetics of ciprofloxacin: intravenous and increasing oral doses. *Am. J. Med.*, 82, 97–102

3.Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., *et al.* (2017). *Stan* : A Probabilistic Programming Language. *J. Stat. Softw.*, 76, 1–32

4.Costello, E.K., Stagaman, K., Dethlefsen, L., Bohannan, B.J.M. & Relman, D.A. (2012). The application of ecological theory toward an understanding of the human microbiome. *Science*, 336, 1255–62

5.Dethlefsen, L., Huse, S., Sogin, M.L. & Relman, D.A. (2008). The Pervasive Effects of an Antibiotic on the Human Gut Microbiota, as Revealed by Deep 16S rRNA Sequencing. *PLoS Biol.*, 6, e280

6.Dethlefsen, L. & Relman, D.A. (2011). Incomplete recovery and individualized responses of the human distal gut microbiota to repeated antibiotic perturbation. *Proc. Natl. Acad. Sci. U. S. A.*, 4554–61

7.Doron, S. & Davidson, L.E. (2011). Antimicrobial stewardship. *Mayo Clin. Proc.*, 86, 1113–23

8.Faith, D.P. (1992). Conservation evaluation and phylogenetic diversity. *Biol. Conserv.*, 61, 1–10

9.Flores, G.E., Caporaso, J.G., Henley, J.B., Rideout, J.R., Domogala, D., Chase, J., *et al.* (2014). Temporal variability is a personalized feature of the human microbiome. *Genome Biol.*, 15, 531

10.Gronau, Q.F., Singmann, H. & Wagenmakers, E.-J. (2017). bridgesampling: An R Package for Estimating Normalizing Constants

11.Guay, D. (2007). Update on clindamycin in the management of bacterial, fungal and protozoal infections. *Expert Opin. Pharmacother.*, 8, 2401–2444

12.Gunderson, L.H. (2000). Ecological Resilience—In Theory and Application. *Annu. Rev. Ecol. Syst.*, 31, 425–439

13.Holling, C.S. (1973). Resilience and Stability of Ecological Systems. *Annu. Rev. Ecol. Syst.*, 4, 1–23

14.Kass, R.E. & Raftery, A.E. (1995). Bayes Factors. *J. Am. Stat. Assoc.*, 90, 773–795

15.Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D., *et al.* (2010). Picante: R tools for integrating phylogenies and ecology. *Bioinformatics*, 26, 1463–1464

16.Leigh, D.A. (1981). Antibacterial activity and pharmacokinetics of clindamycin. *J. Antimicrob. Chemother.*, 7, 3–9

17.Lemon, K.P., Armitage, G.C., Relman, D.A. & Fischbach, M.A. (2012). Microbiota-Targeted Therapies: An Ecological Perspective. *Sci. Transl. Med.*, 4, 137rv5-137rv5

383 18. Llewelyn, M.J., Fitzpatrick, J.M., Darwin, E., SarahTonkin-Crine, Gorton, C., Paul, J., *et al.* (2017). The
384 antibiotic course has had its day. *BMJ*, 358, j3418

385
386 19. Lloyd-Price, J., Abu-Ali, G. & Huttenhower, C. (2016). The healthy human microbiome. *Genome Med.*,
387 8, 51

388
389 20. May, R.M. (Robert M. (1973). *Stability and complexity in model ecosystems*. Princeton University Press

390
391 21. Modi, S.R., Collins, J.J., Relman, D.A., D'Costa, V., Bhullar, K., Costello, E., *et al.* (2014). Antibiotics
392 and the gut microbiota. *J. Clin. Invest.*, 124, 4212–4218

393
394 22. Momeni, B., Xie, L. & Shou, W. (2017). Lotka-Volterra pairwise modeling fails to capture diverse
395 pairwise microbial interactions. *Elife*, 6, e25051

396
397 23. Mustaev, A., Malik, M., Zhao, X., Kurepina, N., Luan, G., Oppegard, L.M., *et al.* (2014).
398 Fluoroquinolone-Gyrase-DNA Complexes. *J. Biol. Chem.*, 289, 12300–12312

399
400 24. Pepper, J.W. & Rosenfeld, S. (2012). The emerging medical ecology of the human gut microbiome.
401 *Trends Ecol. Evol.*, 27, 381–384

402
403 25. Raymond, F., Ouameur, A.A., Déraspe, M., Iqbal, N., Gingras, H., Dridi, B., *et al.* (2015). The initial
404 state of the human gut microbiome determines its reshaping by antibiotics. *ISME J.*

405
406 26. Raymond, F., Ouameur, A.A., Déraspe, M., Iqbal, N., Gingras, H., Dridi, B., *et al.* (2016). The initial
407 state of the human gut microbiome determines its reshaping by antibiotics. *ISME J.*, 10, 707–720

408
409 27. Relman, D.A. (2012). The human microbiome: ecosystem resilience and health. *Nutr. Rev.*, 70 Suppl 1,
410 S2-9

411
412 28. Riley, K.F. (Kenneth F., Hobson, M.P. (Michael P. & Bence, S.J. (Stephen J. (1997). *Mathematical
413 methods for physics and engineering : a comprehensive guide*. Cambridge University Press

414
415 29. Rognes, T., Flouri, T., Nichols, B., Quince, C. & Mahé, F. (2016). VSEARCH: a versatile open source
416 tool for metagenomics. *PeerJ*, 4, e2584

417
418 30. van Schaik, W. (2015). The human gut resistome. *Philos. Trans. R. Soc. B Biol. Sci.*, 370, 20140087–
419 20140087

420
421 31. Scheffer, M., Carpenter, S., Foley, J.A., Folke, C. & Walker, B. (2001). Catastrophic shifts in ecosystems.
422 *Nature*, 413, 591–596

423
424 32. Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., *et al.* (2011). Fast, scalable
425 generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol. Syst. Biol.*, 7,
426 539

427
428 33. Skellam, J.G. (1951). Random dispersal in theoretical populations. *Biometrika*, 38, 196–218

429
34. Spížek, J. & Řezanka, T. (2004). Lincomycin, clindamycin and their applications. *Appl. Microbiol.
Biotechnol.*, 64, 455–464

35. Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics*, 30, 1312–1313

36. Stan Development Team. (2017). RStan: the R interface to Stan. R package version 2.16.2

37. Stein, R.R., Bucci, V., Toussaint, N.C., Buffie, C.G., Rätsch, G., Pamer, E.G., *et al.* (2013). Ecological
modeling from time-series inference: insight into dynamics and stability of intestinal microbiota. *PLoS*
Comput. Biol., 9, e1003388

38. Sullivan, Å., Edlund, C. & Nord, C.E. (2001). Effect of antimicrobial agents on the ecological balance of
human microflora. *Lancet Infect. Dis.*, 1, 101–114

39. Thomas, C., Stevenson, M. & Riley, T. V. (2003). Antibiotics and hospital-acquired Clostridium difficile-
associated diarrhoea: a systematic review. *J. Antimicrob. Chemother.*, 51, 1339–1350

40. Turnbaugh, P.J., Ley, R.E., Hamady, M., Fraser-Liggett, C.M., Knight, R. & Gordon, J.I. (2007). The
Human Microbiome Project. *Nature*, 449, 804–810

41. Wang, Q., Garrity, G.M., Tiedje, J.M. & Cole, J.R. (2007). Naive Bayesian classifier for rapid assignment
of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.*, 73, 5261–7

42. Wootton, J.T. (2010). Experimental species removal alters ecological dynamics in a natural ecosystem.
Ecology, 91, 42–8

43. Zaura, E., Brandt, B.W., Teixeira de Mattos, M.J., Buijs, M.J., Caspers, M.P.M., Rashid, M.-U., *et al.*
(2015). Same Exposure but Two Radically Different Responses to Antibiotics: Resilience of the Salivary
Microbiome versus Long-Term Microbial Shifts in Feces. *MBio*, 6, e01693-15-e01693-15

Figure 1. An impulse response model of antibiotic perturbation to the gut microbiome. We represent the gut microbiome as a unit mass on a stability landscape, where height corresponds to phylogenetic diversity. (A) The healthy human microbiome can be conceptualized as resting in the equilibrium of a stability landscape of all possible states of the microbiome. Perturbations can displace it from this equilibrium value into alternative states (adapted from Lloyd-Price et al. (2016)). (B) Choosing to parameterize this stability landscape using diversity, we assume that there are just two states: the healthy baseline state and an alternative stable state. (C) Perturbation to the microbiome (e.g. by antibiotics) is then modelled as an impulse, which assumes the duration of the perturbation is short relative to the overall timescale of the experiment. We consider the form of the diversity time-response under two scenarios: a return to the baseline diversity; and a transition to a different value of a diversity (i.e. an alternative stable state).

Figure 2. An impulse response model captures the dynamics of the effect of antibiotics on the gut microbiome. Bayesian fits with Stan for participants taking either a placebo (n=10), ciprofloxacin (n=9), or clindamycin (n=9). The mean phylogenetic diversity from 100 bootstraps for each sample (black points) and median and 95% credible interval from the posterior distribution (bold and dashed coloured lines, respectively). The grey line indicates the equilibrium diversity value, defined on a per-individual basis relative to the mean baseline diversity. The biased positive skew of residuals after a year suggests the possibility of a transition to an alternative stable state with persistently reduced diversity.

Figure 3: A model with a possible state transition improves the fit to empirical data. Bayesian fits with Stan for participants taking either a placebo (n=10), ciprofloxacin (n=9), or clindamycin (n=9). The mean phylogenetic diversity from 100 bootstraps for each sample (black points) and median and 95% credible interval from the posterior distribution (bold and dashed coloured lines, respectively). The grey line indicates the equilibrium diversity value, defined on a per-individual basis relative to the mean baseline diversity. The biased positive skew of residuals after a year suggests the possibility of a transition to an alternative stable state with persistently reduced diversity. The non-zero-centred asymptote indicates support for a state transition.

Figure 4: Posterior parameter estimates for model with a possible transition to an alternative stable state. The posterior distributions from Bayesian fits of model 2 (eq. 7) to empirical data for ciprofloxacin (green) and clindamycin (red). Each posterior distribution represents 400,000 iterations in total.

Supplementary Figure 1: Differences in individual response over time for the top twelve most abundant taxonomic families for each treatment group. Relative abundances (log-scale) of the top twelve most abundant bacterial families plotted at each sampled timepoint. Observations are linked by coloured lines for each individual. Despite some consistency in changes between antibiotics across individuals, there is inter-individual variability and evidence of possible interactions between bacterial families.