1 **Experimental assay of a fitness landscape on a macroevolutionary scale**
2

3 Victoria O. Pokusaeva[1+], Dinara R. Usmanova[2+], Ekaterina V. Putintseva[3,4+], Lorena Espinar[3,4],
4 Karen S. Sarkisyan[3,4], Alexander S. Mishin[5], Natalya S. Bogatyreva[3,4], Dmitry N. Ivankov[3,4],
5 Guillaume J. Filion[3,4], Lucas B. Carey[4*], Fyodor A. Kondrashov[1,3,4,6*]
6

[1]Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria.
[2]Department of Systems Biology, Columbia University, New York, 10032, NY, USA.
[3]Bioinformatics and Genomics Programme, Centre for Genomic Regulation (CRG), 88 Dr.
Aiguader, 08003 Barcelona, Spain.
[4]Universitat Pompeu Fabra (UPF), 08003 Barcelona, Spain.
[5]Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Miklukho-Maklaya 16/10, 117997
Moscow, Russia.
[6]ICREA, 23 Pg. LluísCompanys, 08010 Barcelona, Spain.
[+]Equal contribution
[*]Authors for correspondence

Characterizing the fitness landscape, a representation of fitness for a large set of genotypes,
is key to understanding how genetic information is interpreted to create functional
organisms. Here we determined the evolutionarily-relevant segment of the fitness landscape
of His3, a gene coding for an enzyme in the histidine synthesis pathway, focusing on
combinations of amino acid states found at orthologous sites of extant species. The His3
fitness landscape is dominated by synergistic epistasis. For 80% of substitutions, the effect
on fitness depended on the genetic background. Furthermore, at 67% of sites, substitutions
are under sign epistasis, having both strongly positive and negative effect on different genetic
background. 46% of sites were under reciprocal sign epistasis. Sign epistasis affected few
genotypes but involved interaction of multiple sites which makes multidimensionality of the
His3 fitness landscape essential, with the number of dimensions of the landscape
approximately one quarter of the sequence length.

Predicting function and fitness of organisms from their genotypes is the ultimate goal of
many fields in biology, from medical genetics to systems biology to the study of evolution[1-5].
Among the conceptual frameworks for understanding the genotype to phenotype connection is the
fitness landscape, which assigns a fitness (phenotype) to every possible genotype (sequence) of a
gene or genome under consideration[4,6]. The recognition of the importance of the fitness landscape
stimulated the development of a variety of theoretical approaches to its description, including its
general shape and epistatic interactions between alleles, a key property which determines the
complexity of the fitness landscape[7-10]. Before the advent of next-generation sequencing,
experimental assays of the fitness landscape were few and could not address the issue at the
sequence level (e.g. [11]). Recently, large-scale experimental assays described the shape of the
fitness landscape a few mutations away from a local fitness peak[12-21]. Also, some assays involving
a smaller number of genotypes considered combinations of mutations with established
functional[22-27] or evolutionary[28-33] significance.

Empirical evidence of the nature of large-scale fitness landscapes mostly comes from the
study of genotypes incorporating random mutations[13-21] (also see [4] for review), a majority of
which were deleterious[14,18-21,34]. Thus, our present knowledge of fitness landscapes is primarily
driven by the study of deleterious mutations and their interactions, although local adaptive
trajectories have also been considered[2,4,27,35-37]. Deleterious mutations were found to engage in
synergistic epistasis, whereby the joint effect of multiple mutations was stronger than the sum of
their individual effects[11,14,15,17-21,27]. Furthermore, sign epistasis was mostly rare[5,17-21,27,38],
although some of these conclusions differ from study to study (i.e. see [4,38]).

Unfortunately, there are fundamental limitations to assaying the fitness landscape on a
large or macroevolutionary level by random mutation libraries. The number of genotypes
underlying the fitness landscape is the combinatorial set of all amino acids across the length of the
protein[4,6]. For example, for the 220 amino acid protein coded by His3 gene in *Saccharomyces*
cerevisiae, the fitness landscape is a 220 dimensional genotype space with 20^{220} different possible
sequences. Such immense spaces are intractable computationally to say nothing of an experimental
assay. Fortunately, it may not be necessary to survey all genotypes to study the evolutionary-
relevant section of the fitness landscape. Because a vast majority of mutations in protein sequence
are deleterious[34], a randomly sampled sequence is non-functional, especially a protein
sequence[39,40], although this limitation may be less severe for DNA[41].

Here we propose an evolutionary approach for assaying fitness landscapes on a
macroevolutionary scale in a high-throughput manner avoiding random sampling of mostly non-
functional sequences. We reasoned that the functionally and evolutionarily relevant section of the
fitness landscape can be represented by the combination of extant amino acid states, those found
in extant species. This approach, applied previously on a limited scale[28-33] mitigates the problem
of chartering a prohibitively large fitness landscape while highlighting the relationships between
evolutionarily-relevant genotypes (**Figure 1a**). Crucially, substitutions that have been fixed in
evolution are fundamentally different from random mutations, the former representing the driving
force of molecular evolution being either neutral or beneficial in at least some genetic contexts
while the latter, in contrast, are mostly deleterious and are primarily relevant on a
microevolutionary scale. Therefore, current empirical data do not shed much light on the impact
of interactions between substitutions that fixed in the course of evolution by natural selection.
Combinations of extant amino acid states allow one to assay a much wider functionally relevant
area of the sequence space than approaches based on random mutagenesis of a single sequence
(**Figure 1b,c**).

Estimating fitness of evolutionary-relevant genotypes
78
We studied His3, a gene coding for imidazoleglycerol-phosphate dehydratase (IGPD,
His3p), an enzyme essential for histidine synthesis. In a multiple alignment of His3 orthologues
from 21 yeast species we identified 686 extant amino acid states (**Supplementary Data File 1**),
which were evenly distributed across the His3p structure (**Figure 1d**). These 686 substitutions,
which occurred over the course of ~400 million years of yeast evolution[42] (**Figure 1b,c**),
correspond to ~10^{83} sequences, even a tiny fraction of which would be too many to survey. Thus,
we sectioned His3 into 12 independent segments such that the full combinatorial set of
substitutions that have occurred in His3 during yeast evolution comprised 10,000-100,000
genotypes per segment (see **Methods** and **Extended Data Figure 1a**). The 12 segments were of
similar length, constrained by the molecular methods employed for library construction (see
Methods), and covering a diverse range of secondary structure and functional elements. For each
of the 12 segments of His3 we performed an independent experiment surveying its fitness
landscape. For each segment we used degenerate oligonucleotides to construct genotypes
consisting of combinations of amino acids present in extant His3 sequences, and determined the
fitness conferred by these genotypes by expressing them in a Δ*his3* strain of *S. cerevisiae* and
measuring the rate of growth (**Extended Data Figure 1b**). This way we assayed the fitness
landscape of the genotype space that was traversed over the course of the last ~400 million years
of evolution[42].
97
98   Across 12 experiments, we measured fitness for a total of 4,170,503 genotypes (956,648
99 unique amino acid sequences) with high accuracy. Of these, 452,709 consist solely of
100 combinations of extant amino acid states from His3 orthologues, while the remaining genotypes
101 incorporate other amino acid substitutions (**Supplementary Table 1** and **Methods**). For each
102 segment we measured fitness for 60% - 99.8% of all possible genotypes from the combinatorial

set of selected extant amino acid states found in 21 yeast species and a smaller fraction of combinations found across all domains of life (**Supplementary Table 1**), characterizing the evolutionary relevant fitness landscape (**Figure 1b**). For example, for segment 3 consisting of 17 amino acid sites, for 11 sites more than one extant amino acid state was selected from the yeast species (L145=2, L147=2, Q148=3, K151=2, V152=2, D154=3, L164=3, E165=4, A168=2, E169=4, A170=4), with the full yeast combinatorial set consisting of 2*2*3*2*2*3*3*4*2*4*4 = 55,296 genotypes out of which we determined the fitness for 48,198, or 87% of the possible yeast extant states combinations in our library.

A substantial proportion of combinations of extant amino acid states led to genotypes with low fitness (**Figure 2, Figure 3a,b, Extended Data Figure 2**). This observation could be explained by i) some extant amino acids having a universally deleterious effect, ii) some amino acid states exerting a negative effect on fitness because of intergenic interactions with other *S. cerevisiae* genes, or iii) by epistatic interactions between the extant amino acid states within His3[45]. We exclude the possibility that some extant amino acid states had a universally deleterious effect because no extant amino acid states were present only in unfit genetic backgrounds (**Figure 3c**). We exclude the possibility that some extant amino acid states disrupt intergenic interactions because the complete His3 coding sequences from extant species fully complemented a His3 deletion in *S. cerevisiae* (**Extended Data Figure 3**). Thus, the observed genotypes with low fitness can only be explained by epistatic interactions among extant amino acid states in the same or different segment of the His3 gene. Remarkably, more than 80% of substitutions between extant amino acid states had substantially different effects on fitness in different backgrounds (**Figure 3d**). In contrast, only 20% of amino acid substitutions that occurred in His3 evolution are truly neutral, in the sense that do not exert strong influence on fitness in any genetic background. More than half of the universally neutral substitutions were observed in the disordered region of the protein (48/78). Thus, the His3 fitness landscape across the 12 segments was strongly influenced by epistasis on a macroevolutionary scale, whereby the impact of extant amino acid state on fitness being dependent on the background in which it occurs[43-45]. An epistatic fitness landscape is rugged in the sense that evolving genotypes must avoid fitness valleys that emerge through deleterious combinations of amino acid states that may also be found in fit genotypes[29,30,46-48]. Characterizing the ruggedness and the mechanisms that determine the underlying epistasis becomes the primary challenge in understanding the fitness landscape of His3.

Unidimensional epistasis of the His3 fitness landscape

The ruggedness of the fitness landscape can be characterized by the complexity of underlying epistatic interactions. If the epistatic interactions are high-order[49-51] then the fitness landscape can only be projected in a complex multidimensional space, i.e. fitness can only be described by a multidimensional function of the genotype[52]. Alternatively, if epistasis is unidimensional the fitness landscape can be described as a function of an intermediate variable, called the fitness potential[53], determined by additive contribution of amino acid states.

To quantitatively determine how well fitness differences between genotypes can be explained by unidimensional epistasis we used a deep learning approach. We used a sequential neural network architecture composed of three layers. Each neuron in the architecture performs a linear transformation of its input and then applies a nonlinear (sigmoid) function. Hence, by utilizing one neuron in the first layer, we obtain a linear combination of the contributions of each amino acid state to fitness potential, which is then non-linearly mapped to fitness by the three layers of the neural network architecture. (see **Methods**; **Extended Data Figure 4**). Ten segments were described by a sigmoid function in which organismal fitness remains constant with decreasing fitness potential and then is rapidly reduced to lethal after a certain threshold (**Figure 4a**). The ability of the cliff-like threshold fitness function[54] to predict fitness from genotype varied

155   between the His3 segments from near perfect (R2=0.99) in segment 7, to relatively poor (R2=0.66)
156   in segment 5 (**Extended Data Figure 5**).
157
158
159   **Multidimensional epistasis of the His3 fitness landscape**
160
Many genotypes were predicted poorly by any function of a unidimensional fitness
potential (**Extended Data Figure 5a**), and many substitutions can both increase or decrease fitness
depending on the background (**Figure 3d**). Theoretically, the same deep learning approach could
be used to detect non-additive multidimensional interactions by including pairwise (n-wise) terms
as features of the samples in the binary feature matrix (see **Methods** and **Extended Data Figure
4**). However, the number of pairwise features is the square of the number of individual features
and even ~100,000 unique genotype fitness measurements were too few to provide a reasonable
sampling of pairwise interactions. Thus, the neural network could not be used to estimate the
prevalence of multidimensional epistasis with the available number of data points. To determine
the contribution of multidimensional forms of epistasis to the ruggedness of the His3 fitness
landscape we identified instances of sign epistasis: substitutions between extant amino acid states
that were strongly beneficial or strongly deleterious (change in fitness of > 0.4 in absolute value)
depending on the background in which they occurred[55]. Some of these instances may be due to
miscalled fitness of few genotypes, therefore, we confirmed pairs of extant amino acid states to be
under sign epistasis only when sign epistasis was a significantly predictable function of the genetic
background (see **Methods**).
177
An example of sign epistasis is the substitution C141S in the second segment that had an
opposite effect on fitness depending on amino acid at site 143 (I, V or T). The substitution I143T
in turn exhibits sign epistasis depending on the amino acid at site 163 (F, I, V or L) (**Figure 4b,d**).
These epistatic interactions can be represented by a graph in which nodes represent a pair of extant
amino acid states at a specific site and nodes are connected by edges if strong sign epistasis has
been detected between them (C141S - I143T - I163F) (**Figure 4d**). We found that 86 out of 128
(67%) sites in our library exhibit sign epistasis and 46% (59/128) exhibit reciprocal sign epistasis.
Most sites interacted by sign epistasis with multiple other sites (**Figure 4e,f**) demonstrating that,
although sign epistasis effects few genotypes, it leads to a fitness landscape that requires multiple
genotype dimensions for a proper characterization.
188
The dimensionality of the fitness landscape can then be determined by the dimensionality
of the graph of all pairs of sites under sign epistasis. The Euclidean dimensionality of the fitness
landscape was calculated as the fractal dimensionality of the graph of interacting pairs[56], which
ranged from two to seven (**Extended Data Figure 6**). Mathematically, the dimensionality of the
fitness landscape corresponds to the minimal number of dimensions in which the epistatic
interactions shown in (**Extended Data Figure 6**) can be drawn with the lines having equal lengths
(representing a Hamming distance of 1) and without intersections. Thus, strong sign epistasis at
affected sites leads to a high dimensionality of the landscape, at least seven in segment 5, i.e. it is
necessary to consider the simultaneous interaction of seven sites to accurately predict the fitness
of genotypes consisting of extant amino acid states at these sites[50]. Consequently, high
dimensionality decreases the ability of an additive unidimensional model to predict fitness of some
genotypes from sequence (**Figure 4f, inset**).
201
202   **Inference of inter-segmental epistatic interactions in the His3 gene sequence**
203
Epistasis may be caused by interaction among positions within a segment (intra-segmental
epistasis) or by interaction of the segment with the rest of the *S. cerevisiae* His3 sequence (inter-
segmental epistasis). The contribution of inter- versus intra-segmental interactions can be

decoupled. Given two fit genotypes (eg: ABC & abc in one His3 segment), any unfit intermediate
states (eg: aBc in the same His3 segment) must be due to intra-segmental epistasis because the rest
of the protein remains constant. For each segment we took as a measurement of intra-segmental
epistasis all pairs of fit genotypes and calculated the proportion of unfit intermediate genotypes as
a function of the Hamming distance between the two fit genotypes. We then compared this
proportion with the total proportion of all unfit genotypes as a function of Hamming distance from
S. cerevisiae, a measurement that includes both inter- and intra-segmental epistasis. We found
three times more inter-segmental than intra-segmental epistasis (**Extended Data Figure 7**), likely
because a single segment provides a much smaller target space for interactions than the entire His3
protein. The proportion of sites under epistatic interactions increased exponentially with Hamming
distance (**Extended Data Figure 7**), analogous to Orr's snowball, the accumulation of genetic
incompatibilities in the course of speciation[43,57].
219
220 **Evolutionary trajectories on the His3 fitness landscape**
221
The presence of deleterious intermediate genotypes between highly fit ones leads to
inaccessibility of some evolutionary trajectories between extant or ancestral sequences[6,7,31-33,58]
(**Figure 5a**). The sigmoid fitness function provides a mechanism for the restriction of available
evolutionary trajectories in fitness space. Combinations of substitutions, all of which are neutral
in some genetic backgrounds, can take a genotype beyond the fitness threshold through their
additive effect on fitness potential, making some genotypes inaccessible for evolution (**Figure 5a**).
Between any two fit genotypes, the fraction of intermediate genotypes that are unfit depends on
the fitness potential of the two parental genotypes (**Figure 5b**). Evolution between two fit
genotypes with high fitness potential can proceed unhindered because all intermediate genotypes
also have high fitness potential and, consequently, high fitness. Conversely, when both fit
genotypes are located close to the threshold, many of the intermediate genotypes between them
have low fitness and many evolutionary paths between them are inaccessible (**Figure 5c**). Thus,
the cliff-like threshold fitness function is the major determinant of the observation that not all paths
between two fit genotypes are accessible to evolution (**Figure 5b**). We find that unfit intermediate
genotypes are in genetic proximity with each other and are on a limited number of paths; the
fraction of inaccessible paths is smaller than if the same number of unfit genotypes were distributed
randomly in genotype space (**Figure 5d,e,f**).
239
The effect of synergistic epistasis dominates the His3 fitness landscape, affecting over 80%
of amino acid substitutions from our library that occurred in His3 evolution. This synergistic
epistasis may reflect the free energy of the protein[21,59,60], as evidenced by a correlation between
the impact of substitutions on the free energy of His3p and fitness potential (**Extended Data
Figure 8**). Similarly, instances of sign epistasis may also be explained by changes in protein
stability; for example, in the 141T background C141S increased fitness and also had a positive
effect on stability (**Figure 4c**). Consistent with protein stability contributing to the observed sign
epistasis we find that sites that exhibited reciprocal sign epistasis are close together in the His3p
structure (**Extended Data Figure 8)**. However, an additive contribution to free energy can lead
only to a unidimensional fitness function[59], indicating that other non-additive mechanisms, such
as catalytic activity or inter-subunit interactions, or a non-additive model of free energy, must be
responsible for the multidimensionality of the His3 fitness landscape.
252
253 **Conclusions**
254
255   The concept of the fitness landscape introduced by Sewall Wright (Figures 1 and 2 in [6])
256 is an indispensable tool for understanding multiple biological phenomena[1-5]. Experimental high-
257 throughput assays of random mutations have begun to unravel some local properties of fitness
258 landscapes[4]. Here, we described a fitness landscape on a macroevolutionary scale by focusing on

amino acid states that have been put through the sieve of natural selection. We found that only 20% of substitutions that fixed in the evolution of His3 are universally neutral. For the remaining 80%, substitutions from His3 evolution had a profound influence on each other's effect on fitness, providing the first experimental confirmation that epistasis is a staple feature of molecular evolution[45]. Substitutions that occur in evolution have properties vastly different from those of random mutations, which are mostly deleterious[34]. Therefore, the way in which combinations of extant amino acid states affect fitness may also be different from that of combinations of random mutations. Unexpectedly, we found that the interaction of extant amino acid states was dominated by synergistic epistasis in a manner similar to that previously found for random mutations[11,14,15,17-21,27]. However, the accumulation of random mutations leads to low fitness much faster than the accumulation of extant amino acid states (compare Figure 2 from [27] and Figure 3b from [21] to **Figure 2a**).

A key difference of the fitness landscapes assayed by random mutations and through combinations of extant amino acid states is dimensionality. We found that up to 67% of sites with an extant amino acid state participate in multidimensional forms of epistatic interaction forming the basis of a multidimensional structure of the fitness landscape for most His3 segments. However, the presentation of the multidimensional sign epistasis was highly background-specific, affecting a relatively small number of genotypes. For example, the 5-dimensional fitness landscape of His3 segment 2 means that its genotype space cannot be described by fewer than five dimensions (such as the 5-dimensional sequences space in **Figure 1a**). Extrapolating from 20 amino acid long segments to a 220 amino acid long protein, the fitness landscape of the entire protein is expected to be a ~50 dimensional space. By contrast, large-scale fitness landscapes incorporating multiple mutations away from the wildtype sequence in a constant test environment have not displayed pervasive sign epistasis[19-21,27]. Multidimensional epistasis appears to be a more prevalent factor among substitutions that have been subject to positive selection[9,23-27,29-33,35-38], which may have been a factor in the fixation of some of the extant amino acid states in His3.

The evolutionary-relevant section of the His3 fitness landscape is best described as a fitness ridge, with the crest of the ridge defined by its high fitness potential. Evolution can proceed unhindered along the crest **(Figure 5c)**, however, pathway availability declines rapidly when evolution proceeds close to the edge of the fitness ridge. Occasionally, due to the influence of multidimensional sign epistasis a genotype will confer a fitness value not predicted by the fitness potential. The influence of sign epistasis on the His3 fitness ridge is that it is peppered with multidimensional holes and peaks along the landscape; however, their appearance on the landscape is, at present, unpredictable.

Acknowledgements

We thank Ben Lehner for feedback on the design of our study and Julia Domingo for technical assistance with preliminary analysis of our data, Jochen Hecht and the CRG Genomics Unit, and Svetlana Belogorodtdeva and Roman Belogorodtsev for technical assistance. The work was supported by HHMI International Early Career Scientist Program (55007424), the MINECO (BFU2012-31329 and BFU2015-68723-P), Spanish Ministry of Economy and Competitiveness Centro de ExcelenciaSevero Ochoa 2013-2017 grant (SEV-2012-0208), Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement de la Generalitat's AGAUR program (2014 SGR 0974), the CERCA Programme of the Generalitat de Catalunya, and the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013, ERC grant agreement 335980_EinME). L.B.C was supported by MINECO (BFU2015-68351-P) and AGAUR (2014 SGR 0974).

Methods

Study design
        The His3 gene was selected for three principle reasons, it is short, conditionally essential and was not known to be involved in protein-protein interactions. Studying 20^{220} variants of His3 is impossible, thus, we have chosen an approach to survey the fitness landscape in a manner that would elucidate the area most relevant to His3 evolution while managing the technical limitations of our experimental design. We considered amino acid states found in extant species, focusing on yeast species, which translated into a full combinatorial set of $\sim 10^{83}$ unique genotypes. Technically it is feasible to measure fitness of up to 100,000 unique genotypes in a single growth experiment. Therefore, we split the His3 gene into 12 independent segments such that the full combinatorial set of extant amino acid states from 21 yeast species in each segment was 10,000 – 100,000 genotypes. We then considered the combinatorial library for each segment in an independent growth experiment, which allowed us to study a tractable section of the sequence space while considering trajectories across a vast part of the space connecting extant species (**Figure 1a**). We constructed these combinations in 12 plasmid libraries and transformed them into a haploid His3 knockout *S*. *cerevisiae* strain. Growth rate (fitness) of yeast carrying different mutations in His3 was measured using serial batch culture in the absence of histidine.
        We split the His3 gene sequence into segments in a manner agnostic to the structure of the His3 protein (**Extended Data Figure 1a**). For technical reasons, a segment consisted of two variable regions with a constant region between them (**Extended Data Figure 1b**). All growth experiments were performed independently for each segment, with the exception of one experiment on a limited group of genotypes from each segment which was done for the normalization of fitness values across different segments (**Extended Data Figure 3**).
        As a control, we measured the rate of growth of *S*. *cerevisiae* whose entire His3 gene sequence came from another distant species. We found that the replacement of an entire gene sequence of His3 leads to wild-type rates of growth of *S*. *cerevisiae* even when the His3 sequence comes from very distant yeasts, as far as *S*. *pombe* (**Extended Data Figure 3**). Therefore, His3 appears to be an independent unit of the fitness landscape and is a good model for the study of fitness landscapes of an isolated gene.

Measuring fitness
Plasmid construction
        The His3 open reading frame of *S*. *cerevisiae* was PCR amplified with its regulatory region from 622 base pairs (bp) upstream of the open reading frame (ORF) to 237 bp downstream of the ORF, using primers 126 and 127 (see **Supplementary Table 1**) from the wild-type prototroph strain FY4. The PCR product was cloned into vector pRS416 using Gibson assembly (NEB, E2611S). The His3 orthologues from other species were amplified from genomic DNA using designed primers (**Supplementary Table 1**) and were cloned into the vector pRS416_his3, replacing the ORF of *S*. *cerevisiae* by Gibson assembly (NEB, E2611S). Since the His3 orthologue from *A*. *nidulans* contains an intron, the whole open reading frame was initially cloned into the vector, and the intron was later removed by PCR-amplifying the whole plasmid without this sequence, followed by recircularization.

Genomic DNA extraction
        Genomic DNA from fungi (*Saccharomyces cerevisiae, Saccharomyces bayanus, Candida glabrata, Saccharomyces castellii, Kluyveromyces lactis, Eremotheciumgossypii, Debaryomyces*

363 *hansenii, Lodderomycese longosporus, Aspergillus nidulans, Schizosaccharomyces pombe,*
364 *Candida guilliermondii, Saccharomyces kluyveri, Kluyveromyces waltii*) was extracted using
365 MasterPure™ Yeast DNA Purification Kit according to the manufacturer's instructions
366 (Epicentre, MPY80200).

368 *Mutant library construction*
369    Twelve independent mutant libraries, each for different regions of His3 (**Supplementary**
370 **Table 1**), were generated based on the results of multiple alignment of His3 orthologues. The
371 alignment was built using the ClustalW alignment feature of the MEGA 6.0 software package[61]
372 and user-corrected.
373    Mutant libraries were constructed by fusion-PCR, leaving two variable regions separated
by a constant region. For each library, two contiguous fragments of His3 were amplified
independently, using 1 μg of *S. cerevisiae* (strain FY4) genomic DNA in separate Phusion
polymerase reaction mixes (Thermo Fisher Scientific, F530S) in GC buffer. For each PCR, one of
the primers was a degenerate oligonucleotide with a constant part at the 5' end required for the
fusion-PCR; the other primer was either 126 or 127. The cycling conditions for the PCR were
98°C for 30 s; 98°C for 20 s, 60 °C for 30s and 72°C for 1 min (25 cycles); and 72°C for 5 min.
The products were column-purified (QIAGEN, QIAquick PCR purification kit, 28104), eluted in
50 μl and mixed in equimolar proportion. The fusion-PCR was carried out by diluting 10 μl of the
mix to 25 μL of standard Phusion polymerase reaction mix in GC buffer. The cycling conditions
of the fusion-PCR were 98°C for 30 s; 98°C for 30 s, 60°C for 2 min and 72°C for 1 min (25
cycles); and 72°C for 5 min. The product of fusion was purified from agarose gel (Qiagen,
MinElute Gel Extraction Kit, 28604) and eluted in 10 μl of water. 10 μl of the product was used
as a template for additional 5 cycles of PCR reaction in Phusion polymerase reaction mix (Thermo
Fisher Scientific, F530S) in GC buffer, using primers 126 and 127. The cycling conditions were
as follows: 98°C for 30 s; 98°C for 20 s, 60°C for 30 s and 72°C for 1 min (5 cycles); and 72°C
for 5 min. The product was column-purified (QIAGEN, QIAquick PCR purification kit, 28104),
and used as an insert for Gibson assembly.
391    To create a library of His3 mutants, pRS416 plasmid was amplified using primers 128 and
129. The insert was cloned into the vector using Gibson assembly (NEB, E2611S). Ligated
products (200 - 300 ng/μL) were desalted by drop dialysis using 13 mm diameter, Type-VS
Millipore membrane (Merck Millipore, VSWP01300). 20 μL ElectroMAX DH10B competent
cells (Invitrogen, 18290015) were electroporated with 3 μL ligated products. 0.01% of the
electroporated bacteria were plated on ampicillin-containing medium in order to estimate the
complexity of the library; the remaining culture was grown overnight in 100 ml of liquid medium,
and the plasmid was extracted the next day. For each library, the maximum number of protein
sequences that can be generated was computed. Libraries were generated until to total complexity
reached at least 3 times this value.

Yeast transformation and yeast library generation
403    For each segment, yeast strain LBCY47 (*his3*:KanMX*leu2*Δ0 *met15*Δ0 *ura3*Δ0, derived
404 from BY4741) was transformed with 50 μg of pRS416_His3 mutant library using lithium acetate
405 transformation and plated onto glucose synthetic complete dropout plates lacking uracil. After 40
406 hours' growth at 30°C, approximately 0.5 million yeast colonies were scraped off the plates, mixed
407 together and washed 2 times with 100 ml of PBS.

409 *Bulk competition*
410    4×10^9 cells were inoculated into 500 ml of glucose synthetic complete dropout medium
lacking uracil with 200 mg/L of G418, and grown at 30°C at 220 RPM for 6-8 h in order to
eliminate clones with low fitness irrespective of histidine biosynthesis. Cells were later pelleted
and washed with 50 ml of PBS. Approximately 10^{10} cells were inoculated into 1 L of synthetic
complete dropout medium lacking histidine, and grown at 30°C at 220 RPM for 168 h with 12 h

between bottlenecks: ~10^{10} cells were transferred into fresh medium ~10^{8} cells from the culture
were kept as sample for the given time point. Bulk competition for each library of mutants was
done in two replicates to account for biological variability.
418
NGS library preparation
The relative abundance of yeast mutants was measured in 3 samples: 1) the initial
population before selection was applied (t0), 2) the population after 12 h of growth in the selective
medium (t1), and 3) the final population after 168 h of growth in the selective medium (t14). In
order to extract plasmid DNA, $5x10^{9}$ cells from each sample were incubated in 300 μL of
zymolyase buffer (1 M sorbitol, 0.1 M sodium acetate, 60mM EDTA (pH 7.0), 2 mg/ml
zymolyase, 1% 2-Mercaptoethanol) at 37°C for 3 h. The plasmid DNA was purified from the
obtained spheroplasts using QIAprep Spin Miniprep Kit (QIAGEN, 27104) according to the
manufacturer's protocol. The obtained DNA was used as a template in a 25 μL of Q5 DNA
polymerase reaction mix (NEB, M0491S), using staggered primers for demultiplexing in the
following cycling conditions: 98°C for 30s; 98°C for 10s, 60°C for 30s and 72°C for 30s (18
cycles); and 72°C for 2 min. PCR products were purified using Agencourt AM Pure XP beads
(Beckman Coulter, A63880), and eluted in 40 μL of TE buffer (pH 8.0). DNA extraction and PCR-
amplification were repeated twice for every sample to account for the technical variability.
NGS libraries were prepared from 100 ng of the purified DNA amplicons using Ovation Rapid
DR System (Nugen, 0319-32) according to manufacturer's instructions. Each library was
visualized on a Bioanalyzer (Agilent Technologies) and quantified by qPCR with a Kapa Library
Quantification Kit (Kapa Biosystems, KK4835). Twelve samples were pooled together
(accounting for two biological replicates, two technical replicates and three time points) at the final
concentration of 4 nM, and sequenced in the same lane. Samples were sequenced as 125-bp paired-
end reads on a HiSeq2500 sequencer (Illumina) with v4 sequencing chemistry.
440
Yeast growth assay
Mutant strains were grown overnight in complete dropout medium lacking uracil. The
cultures were diluted to 0.05 OD 600 nm, and grown for 5 h in the same medium. 2 μL of each
culture were transferred into 96-well plates in 125 μL of complete dropout medium lacking
histidine. Growth of the strains was monitored by measuring OD 600 nm every 60 min using
Infinite M200 microplate reader (Tecan). Growth rate was defined as the slope of the logOD (time)
line.
448
Growth rates of isolated strains
We isolated 197 strains from all segment libraries of extant amino acid combinations (9-
26 strain per segment) and used Sanger sequencing to determine the sequence. For each strain we
performed 6 repeats of growth assay and calculated the average growth rate. Fitness values from
competition and growth rates are highly correlated (r=0.82 p=10^-48). Correlation was significant
and greater than 0.6 for all segments except S9, where all selected genotypes appeared to be neutral
(**Extended Data Figure 3**).
456
Initial data filtering
The individual sequences of the variants were recovered from pair-end reads with the
following steps: The constant region between the two variable regions was identified by inexact
matching allowing up to 20% errors using the Seeq library version 1.1.2
(https://github.com/ezorita/seeq). The reads are not oriented because the Illumina sequencing
adapters were added by ligation, so the constant regions were searched on both reads. Forward and
reverse reads were swapped when a match was found on the reverse read. This ensured that all of
the sequences are in the same orientation. For multiplexing purposes, the sample identity was
encoded in the left and right primers used to PCR-amplify the variants. To demultiplex the reads,
we used inexact matching with the candidate primers, allowing up to 20% errors. This approach

was faster and less error-prone than using FLASH[62]. To merge the reads, the sequence of the
reverse reads was reverse complemented and the constant region was searched by inexact matching
allowing up to 20% errors. The position of the constant part in each read indicated how they must
be stitched together. In the region of overlap, the consensus sequence was determined by picking
the nucleotide with highest quality as indicated in the quality line of the fastq files. If 'N' persisted
in the final sequence, the reads were discarded. The PCR primers were trimmed so that all the
sequences of the same competition would start and end at the same location.
Reads that did not have the constant region, that could not be oriented or that could not be
demultiplexed were discarded. The remaining errors in the reads were corrected by sequence
clustering. We used starcode version 1.0 ref. [63] with default parameters and allowing up to two
errors. The corrected reads were translated using the genetic code. Variants encoding the same
proteins were not merged; they were kept separate for downstream analyses. A running Docker
virtual machine with commented scripts to replay the whole the process is available for download
at https://hub.docker.com/r/gui11aume/epi/.
481

DNA sequence variant frequency calculation and data filtering
The total number of reads for 12 segments, 3 time points and 4 replicas are shown in
Supplementary Table 1. Genotypes frequencies are defined as the number of reads for a given
genotype divided by the total number of reads in that replicate. Mean frequency was calculated
over 4 replicas to be used in further analysis. However, to eliminate influence of outliers the
median was taken instead of mean if absolute difference between mean and median was greater
than the median value. Only genotypes present in both technical replicas of both biological
replicas with at least ten reads (summed across all time points) in each of them were kept.
490

Noise estimation
The major factors causing noise in genotype frequency measurements are sampling errors,
PCR amplification errors and genetic drift during the competition. For all of these factors, the
amount of error depends on the genotype frequency. Therefore, we estimated measurement errors
as the function of genotype frequency.
For a given segment, time point and a pair of biological or technical replicas for each
genotype we calculated the mean frequency and the squared difference of frequencies from these
two replicas. We sorted genotypes by mean frequency and grouped them such that each bin
contains 5000 genotypes. We calculated the average frequency and the average squared difference
in each bin. Additionally, squared error for frequency 0 was set equal to $\frac{1}{2} \cdot \left(\left(\frac{0.5}{N_i} \right)^2 + \left(\frac{0.5}{N_j} \right)^2 \right)$,
where N_i and N_j are total read numbers in replicas i and j. Finally, by linear interpolation we
obtained dependencies of squared differences as a function of frequency, $s_{ij}^2(f)$, where i and j are
different replicas.
Using squared differences from pairwise comparison of replicas we can estimate variance
of mean frequency over four replicas. Let numerate replicas 1, 2, 3, 4 where 1, 2 are technical
replicas of the first biological repeat and 3, 4 are the technical replicas of the second biological
repeat. Errors coming from the competition (e.g.: genetic drift) are shared for replicas 1, 2 and for
replicas 3, 4. Let's call them Δf_{b_1} and Δf_{b_2} and their variances $\sigma_{b_1}^2$ and $\sigma_{b_2}^2$ respectively. Technical
errors of sampling from the population and from PCR are unique for each replica. Let's call them
$\Delta f_{t_i}, i = 1..4$ and their variances $\sigma_{t_i}^2, i = 1..4$ respectively. All variances are function of frequency
and writing $\sigma_{X_i}^2$ we assume $\sigma_{X_i}^2(f)$.
In the introduced notations the mean frequency over 4 replicas is:

$$f = \frac{1}{4} \cdot (f_1 + f_2 + f_3 + f_4) =$$

$$\frac{1}{4} \cdot \left(\left(f^* + \Delta f_{b_1} + \Delta f_{t_1} \right) + \left(f^* + \Delta f_{b_1} + \Delta f_{t_2} \right) + \left(f^* + \Delta f_{b_2} + \Delta f_{t_3} \right) + \left(f^* + \Delta f_{b_2} + \Delta f_{t_4} \right) \right) =$$

515
$$f^* + \frac{1}{2} \cdot \left(\Delta f_{b_1} + \Delta f_{b_2}\right) + \frac{1}{4} \cdot \left(\Delta f_{t_1} + \Delta f_{t_2} + \Delta f_{t_3} + \Delta f_{t_4}\right),$$

516 where f^* is the true frequency. Applying basic properties of variance the variance of mean
517 frequency:

519
$$\sigma^2 = \frac{1}{4} \cdot \left(\sigma_{b_1}^2 + \sigma_{b_2}^2\right) + \frac{1}{16} \cdot \left(\sigma_{t_1}^2 + \sigma_{t_2}^2 + \sigma_{t_3}^2 + \sigma_{t_4}^2\right)$$

521 To estimate $\sigma_{b_1}^2, \sigma_{b_2}^2, \sigma_{t_1}^2, \sigma_{t_2}^2, \sigma_{t_3}^2, \sigma_{t_4}^2$ we used squared differences from pairwise
522 comparison of replicas calculated above $s_{12}^2, s_{13}^2, s_{14}^2, s_{23}^2, s_{24}^2, s_{34}^2$:

523
$$\mathbb{E}[s_{12}^2] = \mathbb{E}[(\Delta f_{t_1} - \Delta f_{t_2})^2] = \sigma_{t_1}^2 + \sigma_{t_2}^2$$

524
$$\mathbb{E}[s_{13}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_1} - \Delta f_{b_2} - \Delta f_{t_3})^2] = \sigma_{b_1}^2 + \sigma_{t_1}^2 + \sigma_{b_2}^2 + \sigma_{t_3}^2$$

525
$$\mathbb{E}[s_{14}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_1} - \Delta f_{b_2} - \Delta f_{t_4})^2] = \sigma_{b_1}^2 + \sigma_{t_1}^2 + \sigma_{b_2}^2 + \sigma_{t_4}^2$$

526
$$\mathbb{E}[s_{23}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_2} - \Delta f_{b_2} - \Delta f_{t_3})^2] = \sigma_{b_1}^2 + \sigma_{t_2}^2 + \sigma_{b_2}^2 + \sigma_{t_3}^2$$

527
$$\mathbb{E}[s_{24}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_2} - \Delta f_{b_2} - \Delta f_{t_4})^2] = \sigma_{b_1}^2 + \sigma_{t_2}^2 + \sigma_{b_2}^2 + \sigma_{t_4}^2$$

528
$$\mathbb{E}[s_{34}^2] = \mathbb{E}[(\Delta f_{t_3} - \Delta f_{t_4})^2] = \sigma_{t_3}^2 + \sigma_{t_4}^2$$

529 Therefore, the variance of mean frequency f can be found as:

530
$$\sigma^2 = \frac{1}{16} \cdot \left((s_{13}^2 + s_{14}^2 + s_{23}^2 + s_{24}^2) - (s_{12}^2 + s_{34}^2)\right)$$

531 Recalling that variance and squared differences are a function of frequency:

532
$$\sigma^2(f) = \frac{1}{16} \cdot \left((s_{13}^2(f) + s_{14}^2(f) + s_{23}^2(f) + s_{24}^2(f)) - (s_{12}^2(f) + s_{34}^2(f))\right)$$

For each segment and time point we calculated the numerical function $\sigma^2(f)$. Then for each
genotype having mean frequency f_x we estimated its variance as $\sigma^2(f_x)$

Merging amino acid genotypes
We merged nucleotide genotypes that corresponded to the same amino acid sequence and
summed their frequencies and variances. We filtered out all genotypes x which had any of
following patterns:
$f_x^{t_0} = 0, f_x^{t_1} = 0, f_x^{t_2} > 0$ or $f_x^{t_0} = 0, f_x^{t_1} > 0, f_x^{t_2} = 0$ or $f_x^{t_0} > 0, f_x^{t_1} = 0, f_x^{t_2} > 0$. Fraction of
such genotypes were <0.5% for all segments except S9, for which it was 4.5%
For further analysis, this amino acid dataset was used except when specified.

Fitness estimation
Number of cells in a pool with particular genotype x after time interval t increases
exponentially
547
$$n_x^t = n_x^0 \cdot Exp[s_x \cdot t],$$

548 where s_x is absolute fitness. Frequency of genotype x as well depends exponentially on absolute
549 fitness with an additional multiplicative factor:
550
$$f_x^t = \frac{n_x^t}{N^t} = \frac{n_x^0 \cdot Exp[s_x \cdot t]}{N^t} = \frac{f_x^0 \cdot Exp[s_x \cdot t]}{N^t/N^0},$$

551 where N^t and N^0 are total cell numbers in a pool at time points 0 and t. Factor $\frac{1}{N^t/N^0}$ reflects the
552 total growth of population, it changes with time but is the same for all genotypes. Therefore, we
553 can rewrite genotype frequency at time t as:
554
$$f_x^t = f_x^0 \cdot Exp[(s_x - \overline{s^{0t}}) \cdot t],$$

555 where $\overline{s^{0t}} = \frac{1}{t} \cdot Log\left(\frac{N^t}{N^0}\right)$

556   In the measured dataset for each genotype x we have 3 measurements of frequency $f_x^{t_0}$, $f_x^{t_1}$,

557    $f_x^{t_2}$ and their errors $\sigma^2\left(f_x^{t_0}\right), \sigma^2\left(f_x^{t_1}\right), \sigma^2\left(f_x^{t_2}\right)$. To estimate genotype fitness we minimize relative

558   squared errors of exponential fit as function of fitness s_x and initial frequency f_x^0:

559
$$(s_x, f_x^0) = \underset{s_x, f_x^0}{\operatorname{argmin}} \left(\frac{\left(f_x^{t_0} - f_x^0\right)^2}{\sigma^2\left(f_x^{t_0}\right)} + \frac{\left(f_x^{t_1} - f_x^0 \cdot Exp[(s_x - \overline{s^{01}}) \cdot t_1]\right)^2}{\sigma^2\left(f_x^1\right)} + \frac{\left(f_x^{t_2} - f_x^0 \cdot Exp[(s_x - \overline{s^{02}}) \cdot t_2]\right)^2}{\sigma^2\left(f_x^2\right)} \right) \quad (1)$$

This formula contains four parameters common for all genotypes from one segment:

$\overline{s^{01}}, \overline{s^{02}}, t_1, t_2$. Further we will perform additional shifting and scaling of fitness values (see next

section), therefore, without loss of generality we could set $\overline{s^{01}} = 0$ and $t_1 = 1$. Ideally, t_2/t_1 should

equal 14, however, we noticed that this ratio does not hold for many segments and fitted $k = t_2/t_1$

from data instead of using value 14.

To find specific $\overline{s^{02}}$ and k for each segment we selected genotypes with high frequencies

at t_0 $(t_0 > 25 \cdot 10^{-6})$ which corresponds to ~500-1000 reads per technical replicate. Each

segment contains 10^3-10^4 genotypes that meet this criterion. We minimized eq. (1) for selected

genotypes trying all possible combinations of $(\overline{s^{02}}, k)$ from a grid where $\overline{s^{02}} \epsilon [0,1.2]$ with step 0.01

and $k \epsilon [1,14]$ with step 0.1 and choose $(\overline{s^{02}}, k)$ which gives minimal (*).

Finally, given $(\overline{s^{02}}, k)$ for each segment we found s_x for each genotype. Errors for fitness

values, s_x^{std}, were estimated as standard error of best-fit parameter.

For genotypes with frequencies pattern $f_x^{t_0} > 0, f_x^{t_1} = 0, f_x^{t_2} = 0$ fit of eq. (1) cannot be

obtained. Therefore we defined upper boundary for their fitness value as $s_x^{boundary} =$

$Log\left(\frac{1}{\max(N_1^{t_1}, N_2^{t_1}, N_3^{t_1}, N_4^{t_1})}\right)$, where $N_i^{t_1}, i = 1..4$ are total read numbers at time point t_1 in i replica.

Fitness rescaling

We scaled fitness such that lethal genotypes have fitness 0 and neutral genotypes have

fitness 1. We assumed that genotypes with a stop codon or frame shift are lethal. Thus, for each

segment we linearly rescaled the fitness distribution so that 95% of genotypes with nonsense

mutations have a fitness of 0 and so that the local maximum of the fitness distribution of genotypes

with extant amino acids is 1. The scaling around the local maximum led to the shift of fitness

values of less than +/- 0.025 in each of the 12 segments compared to the measured wildtype strains

and did not affect our results (for scale, we called a substitution non-neutral if its effect on fitness

was > 0.4). All fitness values which became smaller than 0 were set to 0.

Control by comparison of synonymous sequences

We used nucleotide synonymous sequences as an internal control. The error rate for a

measurement of fitness of an amino acid sequence depends on the number of synonymous

sequences, *n*, that were used to estimate it. Therefore, we estimated the false discovery rates

separately for categories with *n=1,..10* variants. For each amino acid genotype with more than *n*

synonymous variants we merged random combination of *n* of its nucleotide genotypes and

estimated fitness. We then calculated the difference between this fitness and the fitness of the

corresponding amino acid sequence. We classified case as "false unfit" if difference was <-0.4 and

as "false fit" if difference was >0.4. The fraction of such cases gives us false discovery rates for

genotypes having *n* synonymous variants. To get total false discovery rates for each segment we

averaged "false unfit" and "false fit" rates for different *n* with weights equal to the fractions of

genotypes in amino acid dataset which have *n* synonymous variant (**Supplementary Table 1**).

Predicting fitness using deep learning

To predict the unidimensional fitness function based on additive contribution of extant amino

acid states we used deep learning, a powerful machine learning technique, capable of constructing

virtually any function, even with a simple neural network architecture. To convert amino acid
sequences into a binary feature matrix we used one-hot encoding in which each feature (column
in the matrix) is the presence or absence of a particular amino acid state.

To optimize the accuracy/overfitting ratio, we tested over a hundred of different neural
network architectures and parameters. As a starting point, selected several complicated
architectures, which described our data but were prone to overfitting due to a large number of
parameters. We then gradually reduced the number of layers and neurons to reduce the overfitting,
while controlling for accuracy.

Our final architecture consists of three layers and four neurons in total (**Extended Data**
Figure 4). Each of the neurons performs a linear transformation of the input and subsequently
applies a non-linear activation function (a sigmoid) to the result. The output of the first layer,
therefore, is a sigmoid of a linear transformation of the feature vector ($\mathbf{w}_1^T$x). The second layer
decompresses the hidden nonlinear representation into 2 sigmoids, the combination of which is
further linearly transformed with the only neuron of the third layer and wrapped into another
sigmoid function:

$$F(x) = \sigma(w_{3,1}\sigma(w_{2,1}\sigma(\mathbf{w}_1^T x + b_1) + b_{2,1}) + w_{3,2}\sigma(w_{2,2}\sigma(\mathbf{w}_1^T x + b_1) + b_{2,2}) + b_3)$$

where $\sigma(t) = \frac{1}{(1+exp(-t))}$, x − is the feature vector, w_1 – the vector of weights, corresponding to the
first layer, $w_{n,k}$- the weight corresponding to the n-th layer and the k-th neuron, b_n - bias
corresponding to the n-th layer. This architecture is capable of fitting not only to sigmoid shapes,
but other more complex functions.

In the first hidden layer, a single neuron was used to obtain the predicted fitness potential.
Thus, each amino acid state was assigned a single unique weight. Multiplication of the feature and
weight vectors gave the predicted fitness potential, which was transformed by a nonlinear phase
shift function constructed by the four neurons of the neural network. The architecture simplicity
avoids overfitting, which was further prevented by using 95:5 cross-validation. To ensure
reproducibility, each of the models was constructed five independent times using random train-
test splits.

Each model was trained for 500 epochs using mean squared error as the loss function. An
unpublished adaptive learning rate method proposed by Geoff Hinton, RMSProp, was used as the
optimizer. This algorithm is a version of a mini-batch stochastic gradient descent, utilizing the
gradient magnitude of the recent gradients in order to normalize the current ones. All the weights
were initialized using Xavier normal initializer[64].

Paths between pairs of fit genotypes
For analysis in Figure 5d, we first choose two fit "parental" genotypes, one randomly chosen
genotype (eg: ABE) and the other parental genotype that is either *S. cerevisiae* wildtype genotype
(inter-segmental) or another random fit genotype in the data (intra-segmental) (eg: abe). The two
genotypes in this example are Hamming Distance 3 apart (HD=3). We next compute all (HD2-2)
intermediate genotypes (eg: AbC, aBc, *et cetera*) and retain the subset that were experimentally
measured. We represent the two parental genotypes and all measured intermediate genotypes as
an undirected graph in which each genotype is a vertex. All genotypes one substitution apart are
connected by an unweighted edge. The shortest possible path for a given pair of genotypes is of
length HD. We find all shortest paths between the two parental genotypes using a breadth-first
search. We next remove all vertices (genotypes) that are unfit, and recompute the number of
shortest between the two parental genotypes. For example, in Fig5a, there are six paths of length
three if you take into account all genotypes, but only three paths of length three if you take into
account only fit genotypes.

Clustering of unfit genotypes in sequence space

For the analysis in Figure 5e,f, we first represent the two parental genotypes and all measured intermediate genotypes as an undirected graph in which each genotype is a vertex. All genotypes one substitution apart are connected by an unweighted edge. We can then compute the degree (# of genotypes of distance one) for each vertex (genotype). We do so using all measured genotypes or using only unfit genotypes or using the same number but randomly chosen genotypes. For the randomly chosen genotypes, the value is the average over 1000 runs.

Quantifying sign epistasis

For each substitution (eg: C -> S at position 141), we considered only those that exhibit a large fitness effect (abs. difference > 0.4) comprising a set of substitutions with large effects. For each substitution we divided the genetic backgrounds into two categories: those in which the substitution caused a > 0.4 increase in fitness, and those backgrounds in which the substitution caused > 0.4 decrease in fitness. A single substitution can cause a large increase in fitness in some backgrounds and a large decrease in others due to two possible reasons: sign epistasis or experimental error. To differentiate the two cases, we identified secondary substitutions that significantly alter the ratio of large increases to large decreases in fitness (Fisher's exact test, Bonferroni corrected p-value < 0.05). We only consider a site to be under sign epistasis if there is a second site that alters the frequency of sign epistasis in a statistically significant manner, i.e. more frequently than expected by chance alone.

Structural analysis

Structure prediction

An initial model was obtained with the I-TASSER server[60]. The list of top 10 PDB structural templates picked up by the I-TASSER included high-quality crystal structures of imidazoleglycerol-phosphate dehydratases from *Arabidopsis thaliana* and *Cryptococcus neoformans*. Coordinates of the top-scoring model (C-score=0.21, estimated TM-score = 0.74±0.11, estimated RMSD = 5.1±3.3Å) and the predicted normalized B-factor[66] were used for further analysis. The value of the model quality metric (TM-score >0.5) indicates a model of correct topology. The proteins structurally close to the final model (RMSD 0.6 - 1.7Å are PDB IDs 4MU0, 4GQU, 1RHY, 5DNL and 2AE8 from *Arabidopsis thaliana*, *Mycobacterium tuberculosis*, *Cryptococcus neoformans*, *Pyrococcusfuriosus* and *Staphylococcus aureus*.

We measured the distribution of distances (in angstroms) between pairs of residues that exhibit strong sign epistasis (**Supplementary Table 1**, ReallyPositivePair == TRUE), and compared it with the distribution of pairwise distances among residues for which we have sufficient data to be certain that a given pair does not exhibit sign epistasis (**Supplementary Table 1**, ReallyNegativePair == TRUE).

ΔΔG prediction

Cartesian_ddg application[67] from Rosetta version 2017.08.59291 was used for ΔΔG predictions. Top-scoring I-TASSER model was pre-minimized using the Relax[68] application in dual-space[69] with the flags: -relax:dualspace true; -ex1; -ex2; -use_input_sc; -flip_HNQ; -no_optH false; -relax:min_typelbfgs_armijo_nonmonotone; -nonideal. The best scoring model from 1000 structures was selected. The effect of up to 4 mutations (54,500 genotypes in total) was assessed in cartesian space with the Talaris_2014 score function, and the -fa_max_dis 9.0 flag. ΔΔG was estimated as a difference of mean score for 3 independent runs for every mutant and the wild-type score.

Figures

Figure 1. Combinatorial approach to the study of fitness landscapes. A fitness landscape is
the representation of fitness for all possible genotypes composed of a specific set of loci. **a,**
Following Figure 1 from Sewall Wright ref. [6] consider the genotype space consisting of 5 loci,
each with two allele states (lower and uppercase letters). The entire genotype space is 5-
dimensional consisting of 2^5 genotypes. Given two genotypes found in extant species (abCde and
ABCdE in this example) surveying combinations of extant alleles substantially reduces the
dimensionality, concomitantly reducing the number of genotypes to assay. The surveyed area (blue
cube) considers all combinations of allele substitutions that have occurred in the course of
evolution between the two sequences (red line), avoiding the sampling of combinations with less
relevance to the evolutionary trajectory (black genotypes). **b,** Given the entire multidimensional
genotype space (black circle) our approach considers an N-dimensional subspace consisting of the
combinatorial set of amino acid states form extant species (blue line represents the yeast phylogeny
and the surrounding blue space represents combinations of extant amino acids). By contrast,
random mutagenesis studies consider only a local segment of the genotype space surrounding a
specific genotype (green circle). **c**, A multiple alignment of orthologous sequences of His3 for
segment 2 for which we incorporated almost all extant amino acid states from 21 yeast species
(blue bars) and 10-100% extant states from a set of 396 orthologues (grey bars). **d**, The predicted
structure of His3p with amino acid residues that were substituted in our library.

725
Figure 2. Visual representations of the fitness landscape. a, The fitness landscape for all
assayed genotypes in segment 7. Nodes represent unique amino acid sequences with edges
connecting those separated by a single amino acid substitution. Colour saturation represents the
minimum fitness of the two connected nodes. **b,** On the yeast phylogeny (black lines), fitness of
ancestral and extant nodes and genotypes one substitution away from these nodes are shown.

Figure 3. Fitness distributions. a, The distribution of fitness for genotypes composed of combination of extant amino acid states (red) and non-extant amino acid states (blue) at the same positions. **b**, The fraction of unfit genotypes per segment among genotypes consisting entirely from extant amino acid states (red) and those incorporating non-extant amino acid states (blue). **c**, The number of genotypes with high fitness that incorporates specific amino acid states. For each amino acid state, the number of genetic backgrounds that contain that amino acid state and are fit (fitness > 0) are shown. **d**, The percent of backgrounds in which a specific substitution is neutral (white), beneficial (blue) or deleterious (grey). The region marked in green shows substitutions that never have large effects (> 0.4) on fitness. Beneficial and deleterious effects are shown only if the frequency for a given substitution was higher than the false discovery rate.

Figure 4. Epistasis and the His3 fitness landscape. **a**, Fitness as a function of predicted fitness potential (blue curve). **b,** Substitution C->S at site 141 in segment 2 more frequently has a positive effect on fitness in the background of T at site 143, a negative effect in the background of 143I and is equally likely to be strongly deleterious or strongly beneficial in the background of 143V. **c,** Predicted change in free energy upon a C141S substitution in all genetic backgrounds with either an I or T at 143 and that are fewer than four mutations away from *S. cerevisiae*. **d,** A network depiction of sign epistasis between amino acid substitutions. Colour coded sites with reciprocal sign epistasis (black lines) and unidirectional interactions (grey arrows) are shown. **e,** Distribution of the number of substitutions at each site under sign (yellow) and reciprocal sign (orange) epistasis. Sites with 0 interactions do not exhibit sign epistasis. **f,** The fraction of genotypes in which the substitution under sign epistasis has the less frequent effect on fitness. The inset shows that the ability of an additive model to accurately predict the fitness impact of a substitution decreases with increasing dimensionality of sign-epistasis.

Figure 5. Analysis of evolutionary pathway accessibility. a, A threshold fitness potential function can lead to some paths being inaccessible between two genotypes of high fitness (abe, ABE) if the joint contribution of several alleles to the fitness potential (abE, aBE) leads to the fitness potential below the threshold. **b,** The fraction of unfit intermediate genotypes between two fit genotypes as a function of their average fitness potential. **c,** The grey area represents all genotypes in segment 7. When two fit genotypes (red dots) have high fitness potential, many paths between them will be accessible because many intermediate genotypes will also have high fitness potential and fitness (blue dots). **d,** The fraction of accessible shortest paths between two fit genotypes with unfit genotypes from data (blue) or the same number of randomly selected genotypes (orange), shown as a function of Hamming distance between two fit genotypes. Error bars are standard deviation. **e,** On a graph with genotypes connected by edges if they are one substitution apart we calculate the degree of connectivity (number of edges for each node) for all genotypes (grey), only unfit (fitness = 0) genotypes (red) and a graph with the same number of nodes as the graph with unfit genotypes but with nodes chosen at random (blue). **f,** For all segments except 9 the graph of only unfit genotypes is more interconnected than random graphs with the same number of genotypes.

Extended Data Figures

Extended Data Figure 1. Experimental design. a, The sequence of the His3 protein from *S. cerevisiae* was separated into 12 independent segments of similar lengths, such that the full combinatorial set of extant amino acid substitutions was less than 100,000 possible genotypes. These segments represented different combinations of structural elements of the His3 protein structure. **b**. For each of the 12 segments from His3, we selected extant amino acid states using a multiple alignment of His3 orthologues from 396 species, preferentially incorporating states from

21 yeast species, the variability is shown in section3 as an example. Mutant degenerate codon
libraries were constructed by fusion PCR of two synthesized variable halves of each segment.
These high-complexity plasmid libraries were transformed into haploid His3 knockout *S*.
cerevisiae strain. The growth rate of yeast carrying different extant amino acid state combinations
in His3 gene was measured using serial batch culture in the absence of histidine with 12 hours
between ~100-fold dilutions. To estimate the fitness of yeast mutants their relative abundance was
measured at three points: in the initial population before selection (t0), in the population after 12
hours of growth in the selective medium (t1), and in the final population after 168 hours of growth
in the selective medium (t14). To assess the fitness of individual mutants the segments from three
populations were amplified and sequenced. The relative abundance of each sequence was used as
a proxy for abundance of the associated yeast mutant, which in turn determines its fitness.
804

805
806
Extended Data Figure 2. Segment-specific fitness distributions for extant and non-extant
amino acid states. a, The fitness distribution for each segment for genotypes consisting only of
extant amino acid states (red) or that contain one or more non-extant amino acid states (blue)only
at positions with a substitution in the extant library. **b,** The fitness distribution for each segment
for genotypes consisting only of extant amino acid states (red) and genotypes with mutations at
other positions in that segment (green).
813

814
Extended Data Figure 3. Manual measurements of growth rates of isolated strains.a,His3
orthologues from different species complement a *Δhis3* deletion. Growth rates of fourteen *Δhis3*
strains of *S. cerevisiae*, each of which contains a plasmid with the whole His3 orthologous gene
from another yeast species. **b,** Comparison of fitness values from the pooled competition assay
with growth rates of isolated strains as measured in 96 well plates in a plate reader. **c,** Pearson
correlation coefficients between fitness values from competition and growth rates of isolated
strains for each segment. Error bars for growth rates show s.e.m. from 6 replicas. Error bars for
fitness from competition show standard error of best-fit parameter from fitness estimation, **
signifies p-value < 0.005, * signifies p-value <0.05.

824
825
826 **Extended Data Figure 4. Schematic representation of the deep learning approach.** Each
827 genotype was encoded as a binary vector (x). During training, each of the substitutions was
828 assigned a weight (w_i), comprising a vector of weights (w). The multiplication of these two vectors
829 is the fitness potential of the genotype. After going through three layers, each with a sigmoid
830 activation function, the predicted genotype fitness was obtained.
831

832
833 **Extended Data Figure 5. The fitness potential of the His3 fitness landscape. a**, Fitness potential
834 predicted by the neural network as a function of the measured fitness for all 12 segments. **b**, The
835 correlation between the fitness predicted by the fitness potential and the measured fitness. **c,**
836 Training and test R^2 for each segment for 20-fold cross-validation.
837
838

839
Extended Data Figure 6. Sign epistasis dimensionality graphs for all twelve segments. Each node represents a substitution, with multiple substitutions at the same site having the same colour. Substitutions under reciprocal sign epistasis are indicated by black lines while grey arrows indicate unidirectional sign epistasis.

845
846
Extended Data Figure 7. Decoupling inter- and intra-segmental epistasis. a, The fraction of
unfit genotypes between *S. cerevisiae* and any other genotype consisting of extant amino acid
states with high (blue) or any (red) fitness, and genotypes in the latter but not the former category
(black) as a function of the Hamming distance between the two boundary genotypes. Points
indicate median, the bars and lines indicate 50% of the genotypes and genotypes 2.7 sigmas from
the mean, respectively. **b,** The neural network model assigns higher weights to amino acid states
that that first occur in His3 orthologues farther from *S cerevisiae,* indicating the presence of
intrasegmental interactions.
855

856
Extended Data Figure 8. Protein stability and the fitness potential. a, A comparison of
correlation coefficients between predicted and measured values across segments. **b,c**, correlations
between the estimated impact of substitutions on free energy (ΔΔG), fitness potential and fitness.
ΔΔG correlates better with fitness potential than with fitness. **d**, Pairs of sites that exhibit sign
(connected by a light edge in Extended Data Figure 6) and those that exhibit reciprocal sign
epistasis (connected by a dark edge in Extended Data Figure 6) are closer together in the His3p
structure that randomly chosen non-connected pairs of positions that exhibit sign epistasis.
864

References

1. Dean, A. M. & Thornton, J. W. Mechanistic approaches to the study of evolution: the functional
synthesis. *Nature Rev. Genet.* **8**, 675–688 (2007).

2. Kogenaru, M., de Vos, M. G. J., & Tans, S. J. Revealing evolutionary pathways by fitness
landscape reconstruction. *Crit. Rev. Biochem. Mol. Biol.* **44**, 169–174 (2009).

3.Mackay, T. F. C. Epistasis and quantitative traits: using model organisms to study gene-gene
interactions. *Nature Rev. Genet.* **15**, 22–33 (2014).

4. de Visser, J. A. G. M. & Krug, J. Empirical fitness landscapes and the predictability of evolution.
Nature Rev. Genet. **15**, 480–490 (2014).

5. de Visser, J. A. G. M., Cooper, T. F., & Elena, S. F. The causes of epistasis. *Proc. Biol. Sci.*
278, 3617–3624 (2011).

6. Wright, S. The roles of mutation, inbreeding, crossbreeding and selection in evolution. *Proc.*
Sixth Int. Congr. Genet. **1**, 356–366 (1932).

7. Franke, J., Klözer, A., de Visser. J. A. & Krug, J. Evolutionary accessibility of mutational
pathways. *PLoS Comput. Biol.* **7**, e1002134 (2011).

8. Szendro, I. G., Schenk, M. F., Franke, J., Krug, J. & de Visser, J. A. G. M. Quantitative analyses
of empirical fitness landscapes. *J. Stat. Mech.* P01005 (2013).

9. Schenk, M. F., Szendro, I. G., Salverda, M. L. M., Krug, J., de Visser, J. A. G. M. Patterns of
epistasis between beneficial mutations in an antibiotic resistance gene. *Mol. Biol. Evol.* **30**, 1779–
1787 (2013).

10. Ferretti. L. et al. Measuring epistasis in fitness landscapes: The correlation of fitness effects of
mutations. *J. Theor. Biol.* **396**, 132–143 (2016).

11. de Visser, J. A., Hoekstra, R. F., & van den Ende, H. An experimental test for synergistic
epistasis and its application in Chlamydomonas. *Genetics* **145**, 815–819 (1997).

12. Fowler, D. M. et al. High-resolution mapping of protein sequence-function relationships.
Nature Methods **7**, 741–746 (2010).

13. Jacquier, H. et al. Capturing the mutational landscape of the beta-lactamase TEM-1. *Proc. Natl*
Acad. Sci. USA **110**, 13067–13072 (2013).

14. Melamed, D., Young, D. L., Gamble, C. E., Miller, C. R. & Fields, S. Deep mutational
scanning of an RRM domain of the Saccharomyces cerevisiae poly(A)-binding protein. *RNA* **19**,
1537–1551 (2013).

15. Roscoe, B. P., Thayer, K. M., Zeldovich, K. B., Fushman, D. & Bolon, D. N. Analyses of the
effects of all ubiquitin point mutants on yeast growth rate. *J. Mol. Biol.* **425**, 1363–1377 (2013).

16. Firnberg, E., Labonte, J. W., Gray, J. J. & Ostermeier, M. A comprehensive, high-resolution
map of a gene's fitness landscape. *Mol. Biol. Evol.* **31**, 1581–1592 (2014).

17. Olson, C. A., Wu, N. C. & Sun, R. A comprehensive biophysical description of pairwise
epistasis throughout an entire protein domain. *Curr. Biol.* **24**, 2643–2651 (2014).

18. Bank, C., Hietpas, R. T., Jensen, J. D. & Bolon, D. N. A systematic survey of an intragenic
epistatic landscape. *Mol. Biol. Evol.* **32**, 229–238 (2015).

19. Puchta, O., Cseke, B., Czaja, H., Tollervey, D., Sanguinetti, G., and Kudla, G. Network of
epistatic interactions within a yeast snoRNA. *Science.* **352**, 840–844 (2016).

20. Li, C. Qian, W. Maclean, C. J. & Zhang, J. The fitness landscape of a tRNA gene. *Science* **352**,
837–840 (2016).

21. Sarkisyan, K.S., et al. Local fitness landscape of the green fluorescent protein. *Nature* **533**,
397–401 (2016.)

22. Hietpas, R. T., Bank, C., Jensen, J. D. & Bolon, D. N. A. Shifting fitness landscapes in response
to altered environments. *Evolution* **67**, 3512–3522 (2013).

23. Parera, M. & Martinez, M. A. Strong epistatic interactions within a single protein. *Mol. Biol.*
Evol. **31**, 1546–1553 (2014).

24. Podgornaia, A. I. & Laub, M. T. Pervasive degeneracy and epistasis in a protein-protein
interface. *Science* **347**, 673–677 (2015).

25. Anderson, D. W. McKeown, A. N. & Thornton, J. W. Intermolecular epistasis shaped the
function and evolution of an ancient transcription factor and its DNA binding sites. *Elife* **4**, e07864
(2015).

26. Meini, M. R., Tomatis, P. E., Weinreich, D. M. & Vila, A. J. Quantitative description of a
protein fitness landscape based on molecular features. *Mol. Biol. Evol.* **32**, 1774–1787 (2015).

27. Bank, C., Matuszewski, S., Hietpas, R. T. & Jensen, J. D. On the (un)predictability of a large
intragenic fitness landscape. *Proc. Natl. Acad. Sci. USA.* 113, 14085–14090 (2016).

28. Poelwijk, F.J., Kiviet, D.J., Weinreich, D.M., & Tans, S.J. Empirical fitness landscapes reveal
accessible evolutionary paths. *Nature* **445**, 383–386(2007).

29. Gong, L. I. Suchard, M. A.& Bloom, J.D. Stability-mediated epistasis constrains the evolution
of an influenza protein. *Elife* **2**, e00631(2013).

30. Gong, L. I.& Bloom, J.D. Epistatically interacting substitutions are enriched during adaptive
protein evolution. *PLoS Genet.* **10**, e1004328(2014).

31. Starr, T. N. & Thornton, J. W. Epistasis in protein evolution. *Protein Sci.* **25**, 1204–1218
(2016).

32. Kumar, A. Natarajan, C. Moriyama, H. Witt, C.C. Weber, R.E. Fago, A.& Storz, J.F. Stability-
mediated epistasis restricts accessible mutational pathways in the functional evolution of avian
hemoglobin. *Mol. Biol. Evol.* **34**, 1240–1251(2017).

967
968   33. Tufts, D. M., *et al*. Epistasis constrains mutational pathways of hemoglobin adaptation in high-
969   altitude pikas. *Mol*. *Biol*. *Evol*. **32**, 287–298 (2015).
970
971   34. Eyre-Walker A. & Keightley P. D. The distribution of fitness effects of new mutations. *Nat*.
972   *Rev*. *Genet*. 8, 610–618 (2007).
973
974   35. Draghi, J. A. & Plotkin, J. B. Selection biases the prevalence and type of epistasis along
975   adaptive trajectories. Evolution 67, 3120–3131 (2013).
976
977   36. Lunzer, M., Miller, S. P., Felsheim, R. & Dean, A. M. The biochemical architecture of an
978   ancient adaptive landscape. Science 310, 499–501 (2005).
979
980   37. Palmer, A. C., Toprak, E., Baym, M., Kim, S., Veres, A., Bershtein, S. & Kishony, R. Delayed
981   commitment to evolutionary fate in antibiotic resistance fitness landscapes. *Nat*. *Commun*. **6**, 7385
982   (2015).
983
984   38. Lalić, J. & Elena, S. F. Magnitude and sign epistasis among deleterious mutations in a positive-
985   sense plant RNA virus. *Heredity* **109**, 71–77 (2012).
986
987   39. Keefe, A. D. & Szostak, J. W. Functional proteins from a random-sequence library. *Nature*
988   **410**, 715–718 (2001).
989
990   40. du Plessis, L. Leventhal, G.E. & Bonhoeffer, S. How good are statistical models at
991   approximating complex fitness landscapes? *Mol*. *Biol*. *Evol*. **33**, 2454–2468(2016).
992
993   41. Neme, R., Amador, C., Yildirim, B., McConnell, E. & Tautz, D. Random sequences are an
994   abundant source of bioactive RNAs or peptides. *Nat*. *Ecol*. *Evol*. **1**, 0217 (2017).
995
996   42. Lum, P. Y. Edwards, S.& Wright, R. Molecular, functional and evolutionary characterization
997   of the gene encoding HMG-CoA reductase in the fission yeast, *Schizosaccharomycespombe*. *Yeast*
998   **12**, 1107–1124(1996).
999
1000   43. Kondrashov, A.S. Sunyaev, S. & Kondrashov, F. A. Dobzhansky-Muller incompatibilities in
1001   protein evolution. *Proc*. *Natl*. *Acad*. *Sci*. *USA* **99**, 14878–14883 (2002).
1002
1003   44. Kvitek, D. J. & Sherlock, G. Reciprocal sign epistasis between frequently experimentally
1004   evolved adaptive mutations causes a rugged fitness landscape. *PLoS Genet*.**7**, e1002056 (2011).
1005
1006   45. Breen, M. S., Kemena, C., Vlasov, P. K., Notredame, C. & Kondrashov, F. A. Epistasis as the
1007   primary factor in molecular evolution. *Nature* **490**, 535–538 (2012).
1008
1009   46. Gavrilets, S. Evolution and speciation on holey adaptive landscapes. *Trends Ecol*. *Evol*. **12**,
1010   307–312(1997).
1011
1012   47. Salverda, M. L. M., Dellus, E., Gorter, F. A., Debets, A.J. M., van der Oost, J. Hoekstra, R. F.
1013   Tawfik, D. S. & de Visser, J.A.G. M. Initial mutations direct alternative pathways of protein
1014   evolution. *PLoS Genet*. **7**, e1001321(2011).
1015
1016   48. Weinreich, D. M., Delaney, N. F., de Pristo, M. A. & Hartl, D.L. Darwinian evolution can
1017   follow only very few mutational paths to fitter proteins. *Science* **312**, 111–114(2006).
1018

1019 49. Weinreich, D. M. Lan, Y. Wylie, C. S.&Heckendorn, R. B. Should evolutionary geneticists
1020 worry about higher-order epistasis? *Curr. Opin. Genet. Dev.* **23**, 700–707(2013).
1021
1022 50. Poelwijk, F.J. Krishna, V. & Ranganathan, R. The Context-Dependence of Mutations: A
1023 Linkage of Formalisms. *PLoS Comput. Biol.* **12**, e1004771(2016).
1024
1025 51. Sailer, Z. R. & Harms, M. J. High-order epistasis shapes evolutionary trajectories.
1026 *PLoSComput. Biol.* **13**, e1005541 (2017).
1027
1028 52. Kondrashov, F. A. & Kondrashov, A. S. Multidimensional epistasis and the disadvantage of
1029 sex. *Proc. Natl. Acad. Sci. USA* **98**, 12089–12092 (2001).
1030
1031 53. Crow, J. F. Twenty-five years ago in Genetics: identical triplets. *Genetics* **130**, 395–398(1992).
1032
1033 54. Crow, J. F. & Kimura, M. Efficiency of truncation selection. *Proc. Natl. Acad. Sci. USA* **76**,
1034 396–399(1979).
1035
1036 55. Weinreich, D. M. Watson, R. A. & Chao, L. Perspective: Sign epistasis and genetic constraint
1037 on evolutionary trajectories. *Evolution* **59**, 1165–1174(2005).
1038
1039 56. Song, C. Gallos, L. K. Havlin, S. & Makse, H. A. How to calculate the fractal dimension of a
1040 complex network: the box covering algorithm. *J. Stat. Mech.* **2007**, P03006(2007).
1041
1042 57. Orr, H. A. The population genetics of speciation: the evolution of hybrid incompatibilities.
1043 *Genetics* **139**, 1805–1813 (1995).
1044
58. Bridgham, J. T., Ortlund, E. A. & Thornton, J. W. An epistatic ratchet constrains the
direction of glucocorticoid receptor evolution. *Nature* **461**, 515–519(2009).
59. De Pristo, M. A. Weinreich, D. M. & Hartl, D. L. Missense meanderings in sequence space:
a biophysical view of protein evolution. *Nature Rev. Genet.* **6**, 678–687 (2005).
60. Bershtein, S., Segal, M., Bekerman, R., Tokuriki, N. & Tawfik, D. S. Robustness epistasis
link shapes the fitness landscape of a randomly drifting protein. *Nature* **444**, 929–932 (2006).
61. Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. MEGA6: Molecular
Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol.* **30**, 2725–2729 (2013).
62. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve
genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).
63. Zorita, E., Cuscó, P. & Filion, G. J. Starcode: sequence clustering based on all-pairs search.
Bioinformatics **31**, 1913–1919(2015).
64. Glorot, X. & Bengio, Y. Understanding the difficulty of training deep feedforward neural
networks. in Proceedings of the 13th International Conference on Artificial Intelligence and
Statistics (2010).
1060
1061 65. Yang, J. & Zhang, Y. I-TASSER server: new development for protein structure and function
1062 predictions. *Nucleic Acids Res.* **43**, W174–W181(2015).
1063 66. Yang, J., Wang, Y. & Zhang, Y. ResQ: An Approach to Unified Estimation of B-Factor and
1064 Residue-Specific Error in Protein Structure Prediction. *J. Mol. Biol.* **428,** 693–701 (2016).
1065

67. Park, H., Bradley, P., Greisen, P. Jr., Liu, Y., Mulligan, V. K., Kim, D.E., Baker, D. &
DiMaio, F. Simultaneous optimization of biomolecular energy functions on features from small
molecules and macromolecules. *J. Chem. Theory Comput.* **12**, 6201–6212(2016).

68. Nivón, L. G., Moretti, R. & Baker, D. A Pareto-optimal refinement method for protein
design scaffolds. *PLoS One* **8**, e59004(2013).

69. Conway, P., Tyka, M. D., DiMaio, F., Konerding, D. E. & Baker, D. Relaxation of backbone
bond geometry improves protein energy landscape modeling. *Protein Sci.* **23**, 47–55(2014).

1073