1 **Experimental assay of a fitness landscape on a macroevolutionary scale**
2

3 Victoria O. Pokusaeva[1+], Dinara R. Usmanova[2+], Ekaterina V. Putintseva[1+], Lorena Espinar[3,4],
4 Karen S. Sarkisyan[1,5], Alexander S. Mishin[5], Natalya S. Bogatyreva[3,4,6], Dmitry N. Ivankov[1,6],
5 Arseniy V. Akopyan[1], Sergey Ya. Avvakumov[1], Inna S. Povolotskaya[4], Guillaume J. Filion[3,4],
6 Lucas B. Carey[4*], Fyodor A. Kondrashov[1*]
7

[1]*Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria.*
[2]*Department of Systems Biology, Columbia University, New York, 10032, NY, USA.*
[3]*Bioinformatics and Genomics Programme, Centre for Genomic Regulation (CRG), 88 Dr.*
Aiguader, 08003 Barcelona, Spain.
[4]*Universitat Pompeu Fabra (UPF), 08003 Barcelona, Spain.*
[5]*Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Miklukho-Maklaya 16/10, 117997*
Moscow, Russia.
[6]*Laboratory of Protein Physics, Institute of Protein Research of the Russian Academy of*
Sciences, Pushchino, Moscow region, Russia
[+]*Equal contribution*
[*]*Authors for correspondence*

Characterizing the fitness landscape, a representation of fitness for a large set of genotypes,
is key to understanding how genetic information is interpreted to create functional
organisms. Here we determined the evolutionarily-relevant segment of the fitness
landscape of His3, a gene coding for an enzyme in the histidine synthesis pathway, focusing
on combinations of amino acid states found at orthologous sites of extant species. Just 15%
of amino acids found in yeast His3 orthologues were always neutral while the impact on
fitness of the remaining 85% depended on the genetic background. Furthermore, at 67%
of sites, substitutions are under sign epistasis, having both strongly positive and negative
effect in different genetic backgrounds. 46% of sites were under reciprocal sign epistasis.
Sign epistasis affected few genotypes but involved interaction of multiple sites, shaping a
rugged fitness landscape in which many of the shortest paths between highly fit genotypes
are inaccessible.

Predicting function and fitness of organisms from their genotypes is the ultimate goal of
many fields in biology, from medical genetics to systems biology to the study of evolution[1-5].
Among the conceptual frameworks for understanding the genotype to phenotype connection is
the fitness landscape, which assigns a fitness (phenotype) to every possible genotype (sequence)
of a gene or genome under consideration[4,6]. The recognition of the importance of the fitness
landscape stimulated the development of a variety of theoretical approaches to its description,
including its general shape and epistatic interactions between alleles, a key property which
determines the complexity of the fitness landscape (see [ref. 4] and references within). Before
the advent of next-generation sequencing, experimental assays of the fitness landscape were few
and could not address the issue at the sequence level. Recently, large-scale experimental assays
described the shape of the fitness landscape a few mutations away from a local fitness peak (see
[ref. 7-10] and references within). Also, some assays involving a smaller number of genotypes
considered combinations of mutations with established functional[11-16] or evolutionary[17-22]
significance.

Empirical evidence of the nature of large-scale fitness landscapes mostly comes from the
study of genotypes incorporating random mutations[4,7-10], the majority of which are deleterious[7-
10,23]. Thus, our present knowledge of fitness landscapes is primarily driven by the study of
deleterious mutations and their interactions, although local adaptive trajectories have also been
considered[2,4,16,24-26]. Deleterious mutations were found to engage in synergistic epistasis,

whereby the joint effect of multiple mutations was stronger than the sum of their individual
effects[4,7-10,16]. Furthermore, sign epistasis among random mutations was mostly rare[5,7-10,16,27],
although some of these conclusions differ from study to study (*e.g.* see [4,27]).

Unfortunately, there are fundamental limitations to assaying the fitness landscape on a
large or macroevolutionary level with random mutation libraries. The number of genotypes
underlying the fitness landscape is the combinatorial set of all amino acids across the length of
the protein[4,6]. For example, for the 220 amino acid protein coded by the His3 gene in
Saccharomyces cerevisiae, the fitness landscape is a 220 dimensional genotype space with 20^{220}
different possible sequences. Such immense spaces are both computationally and experimentally
intractable. Fortunately, it may not be necessary to survey all genotypes to study the
evolutionary-relevant section of the fitness landscape. Because the vast majority of mutations in
protein sequences are deleterious[23], a randomly sampled protein sequence is non-functional[28,29].

Here we propose an evolutionary approach for assaying fitness landscapes on a
macroevolutionary scale in a high-throughput manner that avoids the random sampling of mostly
non-functional sequences. The functionally and evolutionarily relevant section of the fitness
landscape can be represented by the combination of extant amino acid states, those found in
extant species. This approach, applied previously on a limited scale[17-22] mitigates the problem of
exploring a prohibitively large fitness landscape while highlighting the relationships between
evolutionarily-relevant genotypes **(Fig. 1a)**. Crucially, substitutions that have been fixed in
evolution are fundamentally different from random mutations, the former are either neutral or
beneficial in at least some genetic contexts and represent the driving force of molecular
evolution, while the latter are mostly deleterious and are primarily relevant on a
microevolutionary scale. Therefore, current empirical data do not shed much light on the impact
of interactions between substitutions that fixed in the course of evolution by natural selection.
Combinations of extant amino acid states allow one to assay a much wider functionally relevant
area of the sequence space than approaches based on random mutagenesis of a single sequence
(Fig. 1b,c).

Estimating fitness of evolutionary-relevant genotypes

We studied His3, a gene coding for imidazoleglycerol-phosphate dehydratase (IGPD,
His3p), an enzyme essential for histidine synthesis. In a multiple alignment of His3 orthologues
from 21 yeast species we identified 686 extant amino acid states (**Supplementary Information
1**), which were evenly distributed across the His3p structure (**Fig. 1d**). These 686 substitutions,
which occurred over the course of ~400 million years of evolution[30] (**Fig. 1b,c**) correspond to
~10^{83} sequences, even a tiny fraction of which would be too many to survey. Thus, we sectioned
His3 into 12 independent segments such that the full combinatorial set of substitutions that have
occurred in His3 during yeast evolution comprised 10,000-100,000 genotypes per segment (see
Methods and **Supplementary Fig. 1a**). The 12 segments were of similar length, constrained by
the molecular methods employed for library construction (see **Methods**), and covered a diverse
range of secondary structures and functional elements (**Supplementary Fig. 1c**). For each of the
12 segments of His3 we performed an independent experiment surveying its fitness landscape.
For each segment we used degenerate oligonucleotides to construct genotypes consisting of
combinations of amino acids present in extant His3 sequences, and determined the fitness
conferred by these genotypes by expressing them in a Δ*his3* strain of *S. cerevisiae* and
measuring the rate of growth **(Supplementary Fig. 1b)**. This way we assayed the fitness
landscape of the genotype space that was traversed over the course of the last ~400 million years
of evolution[30].

Across 11 experiments we measured fitness for a total of 4,018,105 genotypes (875,151
unique amino acid sequences) with high accuracy. Of these, 422,717 consist solely of

combinations of extant amino acid states from His3 orthologues, while the remaining genotypes incorporate other amino acid substitutions (**Supplementary Table 1** and **Methods**). For one segment, 9, the accuracy of our experiment was low, and it was not used in cumulative analyses. For each segment we measured fitness for 60% - 99.8% of all possible genotypes from the combinatorial set of selected extant amino acid states found in 21 yeast species and a smaller fraction of combinations found across all domains of life (**Supplementary Table 1**), characterizing the evolutionary relevant fitness landscape (**Fig. 1b**). For segment 3 for instance, 11 out of 17 amino acid sites had more than one extant amino acid state: L145=2, L147=2, Q148=3, K151=2, V152=2, D154=3, L164=3, E165=4, A168=2, E169=4, A170=4, with the full yeast combinatorial set consisting of 2*2*3*2*2*3*3*4*2*4*4 = 55,296 genotypes out of which we determined the fitness for 48,198, or 87% of the possible yeast extant states combinations in our library.

A substantial proportion of combinations of extant amino acid states led to genotypes with low fitness (**Fig. 2, Fig. 3a,b, Supplementary Fig. 2**), an observation that takes into account the false discovery rate in our data (**Supplementary Table 1**). This observation could be explained by i) some extant amino acids having a universally deleterious effect, ii) some amino acid states exerting a negative effect on fitness because of intergenic interactions with other *S. cerevisiae* genes, or iii) by epistatic interactions between the extant amino acid states within His3[31]. We exclude the possibility that some extant amino acid states had a universally deleterious effect because no extant amino acid states were present only in unfit genetic backgrounds, genotypes conferring a fitness of zero (**Fig. 3c**). We exclude the possibility that some extant amino acid states disrupt intergenic interactions because the complete His3 coding sequences from extant species fully complemented a His3 deletion in *S. cerevisiae* (**Supplementary Fig. 3c**). Thus, the observed genotypes with low fitness can only be explained by epistatic interactions among extant amino acid states within the His3 gene in the same or different segments. Remarkably, 85% (330/389) of substitutions between extant amino acid states had substantially different effects on fitness in different backgrounds (**Fig. 3d**). By contrast, only 15% of amino acid substitutions that occurred in His3 evolution are truly neutral, in the sense that they do not exert strong influence on fitness in any genetic background. Three quarters of the universally neutral substitutions were observed in the disordered region of the protein (44/59). Thus, the His3 fitness landscape across the 11 segments with high accuracy was strongly influenced by epistasis on a macroevolutionary scale, i.e. the impact of an extant amino acid state on fitness often depends on the background in which it occurs[31-34]. An epistatic fitness landscape is rugged in the sense that evolving genotypes must avoid fitness valleys that emerge through deleterious combinations of amino acid states that may also be found in fit genotypes[18,19,34-36]. Characterizing the ruggedness and the mechanisms that determine the underlying epistasis becomes the primary challenge in understanding the fitness landscape of His3.

Unidimensional epistasis of the His3 fitness landscape

The ruggedness of the fitness landscape can be characterized by different measures of complexity of the underlying epistatic interactions. In the simplest case, epistasis may be unidimensional, in the sense that the fitness landscape can be described as a function of an intermediate variable, the fitness potential[37-39]. The fitness landscape is a function from the space of genotypes to fitness. In analogy with a scalar field, we can characterize the ruggedness of this function with standard measures of complexity if genotypes are arranged in a linear space. The simplest case is that of a linear predictor called the fitness potential: $p = c_1 x_1 + c_1 x_2 + \ldots + c_n x_n$, where c_i is a coefficient and x_i is a binary variable that signifies the presence (1) or absence (0) of a given amino acid at a given position. By definition, e^p describes a non-epistatic fitness landscape because the effect of every substitution is multiplicative and it depends only on the

associated c. Any other function of p leads to epistasis. If the $f(p)$ function is "simple", meaning
that is has a small number of local extremas, such as a bimodal function, the epistasis is called
unidimensional[39]. The limitation of simplicity of $f(p)$ is necessary because any function $f_0(x_1, \ldots,$
$x_n)$ can be represented by a function $f'(p)$ and choosing appropriate coefficients $c_1, \ldots, c_n$ in p.
Thus, a simple $f(p)$ leads to unidimensional epistasis because the entire genotype space can be
reduced to a single dimension[39].
160
To quantitatively determine how well fitness differences between genotypes can be
explained by unidimensional epistasis we used a deep learning approach to estimate the
coefficients c for each allele x in the fitness potential and determine the best unidimensional
function of p that best approximated the fitness landscape. We used a dense neural network
architecture composed of three layers. Each neuron in the architecture performed a linear
transformation of its input and then applied a nonlinear (sigmoid) function. Hence, by using one
neuron in the first layer we obtained a linear combination of the contributions of each amino acid
state to fitness potential, which was then non-linearly mapped to fitness by the three layers of the
neural network architecture (see **Methods**; **Supplementary Fig.4**). Ten segments were described
by a threshold function in which organismal fitness remains constant with decreasing fitness
potential and then is rapidly reduced to lethal after a certain threshold (**Fig. 4a**). The ability of
the cliff-like threshold fitness function[40] to predict fitness from genotype varied between the
His3 segments from near perfect (r^2=0.97) in segment 7, to relatively poor (r^2=0.44) in segment 5
(**Supplementary Fig. 5**). Thus, while the fitness landscape of His3 is approximately
unidimensional for some segments, it has a higher degree of complexity for others.
176
177 **Ruggedness and multidimensional epistasis of the His3 fitness landscape**
178
Ruggedness is a general property of fitness landscapes that quantifies the accessible paths
of high fitness that connect fit genotypes[41-43]. A path between highly fit genotypes is inaccessible
when one of the intermediate genotypes has low fitness[6,20-22,41,44] (*e.g.* for genotypes AB and ab,
the intermediate are aB and Ab). Such instances also manifest in sign epistasis on the fitness
landscape, that the same substitution may be beneficial or deleterious when occurring in a
different genetic background[44,45]. To quantify the ruggedness of the His3 fitness landscape we
identified instances of sign epistasis: substitutions between extant amino acid states that were
strongly beneficial or strongly deleterious (change in fitness of > 0.4 in absolute value)
depending on the background in which they occurred[44]. Some of these instances may be due to
miscalled fitness of very few genotypes. Therefore, we considered a pair of extant amino acid
states to be under sign epistasis only when sign epistasis was observed in a statistically
significant number of different genetic backgrounds (see **Methods**).
191
An example of sign epistasis is the substitution C141S in the second segment that had an
opposite effect on fitness depending on amino acid at site 143 (I, V or T). The substitution I143T
in turn exhibits sign epistasis depending on the amino acid at site 163 (F, I, V or L) (**Fig. 5a**).
These epistatic interactions can be represented by a graph in which nodes represent a pair of
extant amino acid states at a specific site and nodes are connected by edges if strong sign
epistasis has been detected between them (C141S - I143T - I163F) (**Fig. 4b**). We found that 86
out of 128 (67%) sites in our library exhibit sign epistasis and 46% (59/128) exhibit reciprocal
sign epistasis. Most sites showed a sign epistatic interaction with multiple other sites (**Fig. 5c,d**)
demonstrating that, although sign epistasis affects few genotypes, it leads to a fitness landscape
that requires the interaction of multiple sites for proper characterization.
202
The complexity of interaction of sites can be estimated by using the graph of sign
epistasis where vertices represent a substitution and edges connect vertices with sign epistasis
between them. If only few substitutions display sign epistasis then such a graph would signify

that the fitness landscape is relatively smooth, alternatively, a highly-interconnected graph of
such interactions signifies a more rugged landscape[4,5,41-43]. To measure the relative fitness
complexity, we used the maximum clique size of a graph, which approximates the maximum
number of simultaneously interacting substitutions. In our data, this measure ranged from two to
seven depending on the segment (**Supplementary Fig. 6**). The ruggedness of the fitness
landscape of His3 is high for most segments, such as segment 5, where it is necessary to consider
the simultaneous interaction of at least seven sites to accurately predict the fitness of genotypes
consisting of extant amino acid states at these sites[38,40].

Sign epistasis can appear when fitness is described by a unidimensional function of the
fitness potential, for example, when the fitness landscape is a unimodal function with an
optimum in an intermediate range of the fitness potential[41,45]. However, sign epistasis may also
be a sign of multidimensional epistasis, when a unidimensional function of the fitness potential
cannot fully describe genotype fitness[39]. Many genotypes were predicted poorly by a
unidimensional function of the fitness potential (**Supplementary Fig. 5a**). Two lines of evidence
suggest that such genotypes reveal the presence of multidimensional epistasis. First, genotypes
with a higher number of substitutions influenced by sign epistasis were less well-predicted by a
unidimensional fitness function (**Fig. 4c and Supplementary Fig. 7b**). Second, we explain a
larger fraction of genotypes by using a more complex neural network architecture
accommodating multiple fitness potentials instead of one. We found that increasing the amount
of neurons in the first layer of the neural network architecture, which is equivalent to increasing
the number of independent fitness potentials, gradually improves the prediction power of the
obtained models for most of the segments (**Fig. 4d**). Thus, adding dimensions to the function of
fitness potential increases the prediction power of the model. For example, for a two-
dimensional case fitness was described by $f_1(p_1, p_2)$ with $p_1 = a_1x_1 + a_1x_2 \ldots a_nx_n$ and $p_2 = b_1x_1 +$
$b_1x_2 \ldots b_nx_n$. For several His3 segments, a fitness function with multiple underlying fitness
potentials described the fitness landscape more accurately than a simple unidimensional function
of a single fitness potential (**Supplementary Fig. 7a**). For instance, for these segments, fitness
function of two fitness potentials described the shape with a higher degree of accuracy than a
function of a single fitness potential (**Fig. 4d,e**). By contrast, epistasis in segment 7 is entirely
unidimensional (**Fig. 4d,e and Supplementary Information 2**); we do not see any improvement
in the model's predictive power when adding extra dimensions.

Evolutionary trajectories on the His3 fitness landscape

On a smooth fitness landscape, evolution can proceed along any of the evolutionary paths
connecting two fit genotypes, as none of the intermediate genotypes confer low fitness (see Box
2 in [46]). Alternatively, the fitness landscape is rugged when it contains non-connected fitness
peaks, such that there are no viable paths between some pairs of genotypes that confer high
fitness[4,5]. In other words, the presence of deleterious intermediate genotypes between highly fit
ones leads to inaccessibility of some evolutionary trajectories between extant or ancestral
sequences[6,20-22,44]. The simplest explanation for the substantial ruggedness of the landscape
observed in many of the His3 segments lies in the unidimensional threshold fitness function
(**Fig. 6a**). On a threshold function a combinations of substitutions, all of which are neutral in
some genetic backgrounds, can take a genotype beyond the fitness threshold through their
additive effect on fitness potential, making some genotypes inaccessible for evolution (**Fig. 6a**).
Between any two fit genotypes, the fraction of intermediate genotypes that are unfit depends on
the fitness potential of the two parental genotypes (**Fig. 6b**). Evolution between two fit
genotypes with high fitness potential can proceed unhindered because all intermediate genotypes
also have high fitness potential and, consequently, high fitness. Conversely, when both fit
genotypes are located close to the threshold, many of the intermediate genotypes between them
have low fitness and many evolutionary paths between them are inaccessible (**Fig. 6c**). Thus, the

cliff-like threshold fitness function is the major determinant of the observation that not all paths
between two fit genotypes are accessible to evolution (**Fig. 6b**). We find that unfit intermediate
genotypes are in genetic proximity with each other and are on a limited number of paths; the
fraction of inaccessible paths is smaller than if the same number of unfit genotypes were
distributed randomly in genotype space (**Fig. 6d,e**).
263
The effect of synergistic epistasis dominates the His3 fitness landscape, affecting over
85% of amino acid substitutions from our library that occurred in His3 evolution. This
synergistic epistasis may reflect the free energy of the protein[10,47,48], as evidenced by a
correlation between the fitness potential and the impact of substitutions on the free energy of
His3p (**Supplementary Fig. 8**). Similarly, instances of sign epistasis may also be explained by
changes in protein stability; for example, in the 143T background C141S increased fitness and
also had a positive effect on stability (**Fig. 5b**). Consistent with protein stability contributing to
the observed sign epistasis we find that sites that exhibited reciprocal sign epistasis are close
together in the His3p structure (**Supplementary Fig. 8**). However, an additive contribution to
free energy can lead only to a unidimensional fitness function[47], indicating that other non-
additive mechanisms, such as catalytic activity or inter-subunit interactions, or a non-additive
model of free energy, must be responsible for the multidimensionality of the His3 fitness
landscape.
277
278 **Inference of inter-segmental epistatic interactions in the His3 gene sequence**
279
Epistasis may be caused by interaction among positions within a segment (intra-
segmental epistasis) or by interaction of the segment with the rest of the *S. cerevisiae* His3
sequence (inter-segmental epistasis). The contribution of inter- versus intra-segmental
interactions can be decoupled. Given two fit genotypes (*e.g.* ABC & abc in one His3 segment),
any unfit intermediate states (*e.g.* aBc in the same His3 segment) must be due to intra-segmental
epistasis because the rest of the protein remains constant. For each segment we took as a
measurement of intra-segmental epistasis all pairs of fit genotypes and calculated the proportion
of unfit intermediate genotypes as a function of the Hamming distance between the two fit
genotypes. We then compared this proportion with the total proportion of all unfit genotypes as a
function of Hamming distance from *S. cerevisiae*, a measurement that includes both inter- and
intra-segmental epistasis. We found three times more inter-segmental than intra-segmental
epistasis (**Supplementary Fig. 9**), likely because a single segment provides a much smaller
target space for interactions than the entire His3 protein. The proportion of sites under epistatic
interactions increased exponentially with Hamming distance (**Supplementary Fig. 9**), analogous
to Orr's snowball, the accumulation of genetic incompatibilities in the course of speciation[31,49].
295
296 **Conclusions**
297
The concept of the fitness landscape introduced by Sewall Wright (Figures 1 and 2 in [6])
is an indispensable tool for understanding multiple biological phenomena[1-5]. Experimental high-
throughput assays of random mutations have begun to unravel some local properties of fitness
landscapes[4]. Here, we described a fitness landscape on a macroevolutionary scale by focusing on
amino acid states that have been put through the sieve of natural selection. We found that only
15% of substitutions that fixed in the evolution of His3 are universally neutral. For the remaining
85%, substitutions from His3 evolution had a profound influence on each other's effect on
fitness, providing an experimental confirmation that epistasis is one of the defining features of
molecular evolution[33]. Substitutions that occur in evolution have properties vastly different from
those of random mutations, which are mostly deleterious[23]. Therefore, the way in which
combinations of extant amino acid states affect fitness may also be different from that of
combinations of random mutations. Unexpectedly, we found that the interaction of extant amino

acid states was dominated by synergistic epistasis in a manner similar to that previously found
for random mutations[7-10,16]. However, the accumulation of random mutations leads to low fitness
much faster than the accumulation of extant amino acid states (compare Figure 2 from [16] and
Figure 3b from [10] to **Fig. 2a**).
314

The experimental data showing that 85% of amino acid states found in extant species
confer low fitness in a different genetic background lends strong support to the notion that
epistasis is a key factor in protein evolution[31,33]. We showed that the fitness landscape of several
segments of the His3 gene cannot be reduced to a single unidimensional forms of epistasis, with
a function of multiple fitness potentials providing a more accurate description of the fitness
landscape. By contrast, large-scale fitness landscapes incorporating multiple random mutations
away from the wildtype sequence in a constant test environment have not displayed evidence of
multidimensional epistasis[8-10,16]; however, it appears to be a more prevalent factor among
substitutions that have been subject to positive selection[12-16,18-22,34-36]. We also found that up to
67% of sites with an extant amino acid state were influenced by sign epistasis, resulting in a
rugged fitness landscape and a limited number of fitness ridges connecting extant sequences for
most His3 segments. Overall, the evolutionary-relevant section of the His3 fitness landscape is
best described as a fitness ridge, with the crest of the ridge defined by a fitness potential. In some
cases, the crest is multidimensional requiring several independent underlying fitness potentials.
Evolution can proceed unhindered along the crest **(Fig. 6a,c)**, however, pathway availability
declines rapidly when evolution proceeds close to the edge of the fitness ridge.
331

Acknowledgements
We thank Ben Lehner for a thorough study of the concept behind our experiments and Julia
Domingo for technical assistance with preliminary analysis of our data. Jochen Hecht of the
CRG Genomics Unit and Svetlana Belogorodtdeva and Roman Belogorodtsev provided
technical assistance. The work was supported by HHMI International Early Career Scientist
Program (55007424), the MINECO (BFU2012-31329, BFU2012-37168, BFU2015-68351-P and
BFU2015-68723-P), Spanish Ministry of Economy and Competitiveness Centro de Excelencia
Severo Ochoa 2013-2017 grant (SEV-2012-0208), the Unidad de Excelencia María de Maeztu
funded by the MINECO (MDM-2014-0370), Secretaria d'Universitats i Recerca del Departament
d'Economia i Coneixement de la Generalitat AGAUR program (2014 SGR 0974), the CERCA
Programme of the Generalitat de Catalunya, Russian Foundation for Basic Research grant (18-
04-01173), the European Union's Horizon 2020 research and innovation programme under the
Marie Skłodowska-Curie programme (665385) and the European Research Council under the
European Union's Seventh Framework Programme (FP7/2007-2013, ERC grant agreement
335980_EinME and Synergy Grant 609989).

Author contribution. ISP, LBC and FAK conceived the general approach of the study. FAK,
LBC, VOP, LE and GJF participated in detailed experimental design. VOP, DRU and EVP
spearheaded the experiment, data analysis and main data interpretation, respectively. LE
performed a large fraction of the experimental work. AVA and SYaA participated in working on
dimensionality. KSS, ASM, NSB, DNI and GJF participated in data analysis. LBC and FAK
wrote the draft.

Methods
356

Data access
The raw and processed data have been submitted to the NCBI Gene Expression Omnibus
(GEO; http://www.ncbi.nlm.nih.gov/geo/) under accession number GSE99990. A virtual machine
containing a running version of the data processing pipeline is available as a Docker image

https://hub.docker.com/r/gui11aume/epi/. The scripts to reproduce the figures are on Github at
https://github.com/Lcarey/HIS3InterspeciesEpistasis.
363

Study design

The His3 gene was selected for three principal reasons, it is short, conditionally essential
and was not known to be involved in protein-protein interactions. Studying 20^{220} variants of His3
is impossible, thus, we have chosen an approach to survey the fitness landscape in a manner that
would elucidate the area most relevant to His3 evolution while managing the technical
limitations of our experimental design. We considered amino acid states found in extant species,
focusing on yeast species, which translated into a full combinatorial set of ~10^{83} unique
genotypes. Technically it is feasible to measure fitness of up to 100,000 unique genotypes in a
single growth experiment. Therefore, we split the His3 gene into 12 independent segments such
that the full combinatorial set of extant amino acid states from 21 yeast species in each segment
was 10,000 – 100,000 genotypes. We then considered the combinatorial library for each segment
in an independent growth experiment, which allowed us to study a tractable section of the
sequence space while considering trajectories across a vast part of the space connecting extant
species (**Fig. 1a**). We constructed these combinations in 12 plasmid libraries and transformed
them into a haploid His3 knockout *S. cerevisiae* strain. Growth rate (fitness) of yeast carrying
different mutations in His3 was measured using serial batch culture in the absence of histidine.
We split the His3 gene sequence into segments in a manner agnostic to the structure of
the His3 protein (**Supplementary Fig. 1a**). For technical reasons, a segment consisted of two
variable regions with a constant region between them (**Supplementary Fig. 1b**). All growth
experiments were performed independently for each segment, with the exception of one
experiment on a limited group of genotypes from each segment which was done for the
normalization of fitness values across different segments (**Supplementary Fig. 3**).
As a control, we measured the rate of growth of *S. cerevisiae* whose entire His3 gene
sequence came from another distant species. We found that the replacement of an entire gene
sequence of His3 leads to wild-type rates of growth of *S. cerevisiae* even when the His3
sequence comes from very distant yeasts, as far as *S. pombe* (**Supplementary Fig. 3**). Therefore,
His3 appears to be an independent unit of the fitness landscape and is a good model for the study
of fitness landscapes of an isolated gene.
392

Measuring fitness

Plasmid construction
The His3 open reading frame of *S. cerevisiae* was PCR amplified with its regulatory
region from 622 base pairs (bp) upstream of the open reading frame (ORF) to 237 bp
downstream of the ORF, using primers 126 and 127 (see **Supplementary Table 1**) from the
wild-type prototroph strain FY4. The PCR product was cloned into vector pRS416 using Gibson
assembly (NEB, E2611S). The His3 orthologues from other species were amplified from
genomic DNA using designed primers (**Supplementary Table 1**) and were cloned into the
vector pRS416_his3, replacing the ORF of *S. cerevisiae* by Gibson assembly (NEB, E2611S).
Since the His3 orthologue from *A. nidulans* contains an intron, the whole open reading frame
was initially cloned into the vector, and the intron was later removed by PCR-amplifying the
whole plasmid without this sequence, followed by recircularization.
405

Genomic DNA extraction
Genomic DNA from fungi (*Saccharomyces cerevisiae, Saccharomyces bayanus, Candida*
glabrata, Saccharomyces castellii, Kluyveromyces lactis, Eremotheciumgossypii, Debaryomyces
hansenii, Lodderomycese longosporus, Aspergillus nidulans, Schizosaccharomyces pombe,
Candida guilliermondii, Saccharomyces kluyveri, Kluyveromyces waltii) was extracted using
MasterPure™ Yeast DNA Purification Kit according to the manufacturer's instructions
(Epicentre, MPY80200).

413
Mutant library construction
Twelve independent mutant libraries, each for different regions of His3 **(Supplementary**
Table 1), were generated based on the results of multiple alignment of His3 orthologues. The
alignment was built using the ClustalW alignment feature of the MEGA 6.0 software package[50]
and user-corrected.
Mutant libraries were constructed by fusion-PCR, leaving two variable regions separated
by a constant region. For each library, two contiguous fragments of His3 were amplified
independently, using 1 μg of *S. cerevisiae* (strain FY4) genomic DNA in separate Phusion
polymerase reaction mixes (Thermo Fisher Scientific, F530S) in GC buffer. For each PCR, one
of the primers was a degenerate oligonucleotide with a constant part at the 5' end required for
the fusion-PCR; the other primer was either 126 or 127. The degenerate primer approach led to
the integration of non-extant amino acid sequences due to the redundancy of the genetic code.
Consider the amino acid Phe in *S. cerevisiae* coded by the codon TTT. When incorporating an
extant orthologous state Trp (TGG) two independent T -> G nucleotide mutations will be
incorporated creating the codons TTG (Leu) and TGT (Cys). If these two amino acids were not
found in other species then they would be non-extant. The cycling conditions for the PCR were
98°C for 30 s; 98°C for 20 s, 60 °C for 30s and 72°C for 1 min (25 cycles); and 72°C for 5 min.
The products were column-purified (QIAGEN, QIAquick PCR purification kit, 28104), eluted in
50 μl and mixed in equimolar proportion. The fusion-PCR was carried out by diluting 10 μl of
the mix to 25 μL of standard Phusion polymerase reaction mix in GC buffer. The cycling
conditions of the fusion-PCR were 98°C for 30 s; 98°C for 30 s, 60°C for 2 min and 72°C for 1
min (25 cycles); and 72°C for 5 min. The product of fusion was purified from agarose gel
(Qiagen, MinElute Gel Extraction Kit, 28604) and eluted in 10 μl of water. 10 μl of the product
was used as a template for additional 5 cycles of PCR reaction in Phusion polymerase reaction
mix (Thermo Fisher Scientific, F530S) in GC buffer, using primers 126 and 127. The cycling
conditions were as follows: 98°C for 30 s; 98°C for 20 s, 60°C for 30 s and 72°C for 1 min (5
cycles); and 72°C for 5 min. The product was column-purified (QIAGEN, QIAquick PCR
purification kit, 28104), and used as an insert for Gibson assembly.
To create a library of His3 mutants, pRS416 plasmid was amplified using primers 128
and 129. The insert was cloned into the vector using Gibson assembly (NEB, E2611S). Ligated
products (200 - 300 ng/μL) were desalted by drop dialysis using 13 mm diameter, Type-VS
Millipore membrane (Merck Millipore, VSWP01300). 20 μL ElectroMAX DH10B competent
cells (Invitrogen, 18290015) were electroporated with 3 μL ligated products. 0.01% of the
electroporated bacteria were plated on ampicillin-containing medium in order to estimate the
complexity of the library; the remaining culture was grown overnight in 100 ml of liquid
medium, and the plasmid was extracted the next day. For each library, the maximum number of
protein sequences that can be generated was computed. Libraries were generated until to total
complexity reached at least 3 times this value.
452
Yeast transformation and yeast library generation
For each segment, yeast strain LBCY47 (*his3*:KanMX*leu2*Δ0 *met15*Δ0 *ura3*Δ0, derived
from BY4741) was transformed with 50 μg of pRS416_His3 mutant library using lithium acetate
transformation and plated onto glucose synthetic complete dropout plates lacking uracil. After 40
hours' growth at 30°C, approximately 0.5 million yeast colonies were scraped off the plates,
mixed together and washed 2 times with 100 ml of PBS.
459
Bulk competition
4×10^9 cells were inoculated into 500 ml of glucose synthetic complete dropout medium
lacking uracil with 200 mg/L of G418, and grown at 30°C at 220 RPM for 6-8 h in order to
eliminate clones with low fitness irrespective of histidine biosynthesis. Cells were later pelleted
and washed with 50 ml of PBS. Approximately 10^{10} cells were inoculated into 1 L of synthetic

complete dropout medium lacking histidine, and grown at 30°C at 220 RPM for 168 h with 12 h
between bottlenecks: ~10^{10} cells were transferred into fresh medium ~10^8 cells from the culture
were kept as sample for the given time point. Bulk competition for each library of mutants was
done in two replicates to account for biological variability.
469
470   *NGS library preparation*
471   The relative abundance of yeast mutants was measured in 3 samples: 1) the initial
population before selection was applied (t0), 2) the population after 12 h of growth in the
selective medium (t1), and 3) the final population after 168 h of growth in the selective medium
(t14). In order to extract plasmid DNA, $5x10^9$ cells from each sample were incubated in 300 μL
of zymolyase buffer (1 M sorbitol, 0.1 M sodium acetate, 60mM EDTA (pH 7.0), 2 mg/ml
zymolyase, 1% 2-Mercaptoethanol) at 37°C for 3 h. The plasmid DNA was purified from the
obtained spheroplasts using QIAprep Spin Miniprep Kit (QIAGEN, 27104) according to the
manufacturer's protocol. The obtained DNA was used as a template in a 25 μL of Q5 DNA
polymerase reaction mix (NEB, M0491S), using staggered primers for demultiplexing in the
following cycling conditions: 98°C for 30s; 98°C for 10s, 60°C for 30s and 72°C for 30s (18
cycles); and 72°C for 2 min. PCR products were purified using Agencourt AM Pure XP beads
(Beckman Coulter, A63880), and eluted in 40 μL of TE buffer (pH 8.0). DNA extraction and
PCR-amplification were repeated twice for every sample to account for the technical variability.
484
NGS libraries were prepared from 100 ng of the purified DNA amplicons using Ovation Rapid
DR System (Nugen, 0319-32) according to manufacturer's instructions. Each library was
visualized on a Bioanalyzer (Agilent Technologies) and quantified by qPCR with a Kapa Library
Quantification Kit (Kapa Biosystems, KK4835). Twelve samples were pooled together
(accounting for two biological replicates, two technical replicates and three time points) at the
final concentration of 4 nM, and sequenced in the same lane. Samples were sequenced as 125-bp
paired-end reads on a HiSeq2500 sequencer (Illumina) with v4 sequencing chemistry.
492
493   *Yeast growth assay*
494   Mutant strains were grown overnight in complete dropout medium lacking uracil. The
495   cultures were diluted to 0.05 OD 600 nm, and grown for 5 h in the same medium. 6 μL of each
496   culture were transferred into 96-well plates in 125 μL of complete dropout medium lacking
497   histidine. Growth of the strains was monitored by measuring OD 600 nm every 10 min using
498   Tecan Infinite M1000 PRO microplate reader equipped with an integrated Stacker module.
499
500   The growth rate of individual curves was measured as the inverse of the time to grow
from OD = 0.135 = exp(-2) to OD = 0.368 = exp(-1). If the curve did not reach 0.368, the growth
was set to 0. Curves that crossed 0.135 or 0.368 were excluded. The growth rate of a clone was
measured as the median of 6 independent growth experiments. We excluded from the analysis
clones with discordance between growth in solid and liquid medium, clones that could not be
sequenced or that showed evidence of contamination by sequencing, and clones such that the
Kullback-Leibler divergence of their read counts compared to all synonymous clones was greater
than 0.22. The later criterion ensured that the selected clones were not outliers compared to other
variants encoding the same protein.
509
510   **Growth rates of isolated strains**
511   We isolated 197 strains from all segment libraries of extant amino acid combinations (9-
512   26 strain per segment) and used Sanger sequencing to determine the sequence. For each strain
513   we performed 6 repeats of growth assay and calculated the average growth rate. Fitness values
514   from competition and growth rates are highly correlated (r=0.82 p=10^{-48}). Correlation was
515   significant and greater than 0.6 for all segments except S9, where all selected genotypes
516   appeared to be neutral (**Supplementary Fig. 3**).

517
Initial data filtering

The individual sequences of the variants were recovered from pair-end reads with the
following steps: the constant region between the two variable regions was identified by inexact
matching allowing up to 20% errors using the Seeq library version 1.1.2
(https://github.com/ezorita/seeq). The reads are not oriented because the Illumina sequencing
adapters were added by ligation, so the constant regions were searched on both reads. Forward
and reverse reads were swapped when a match was found on the reverse read. This ensured that
all of the sequences are in the same orientation. For multiplexing purposes, the sample identity
was encoded in the left and right primers used to PCR-amplify the variants. To demultiplex the
reads, we used inexact matching with the candidate primers, allowing up to 20% errors. This
approach was faster and less error-prone than using FLASH[51]. To merge the reads, the sequence
of the reverse reads was reverse complemented and the constant region was searched by inexact
matching allowing up to 20% errors. The position of the constant part in each read indicated how
they must be stitched together. In the region of overlap, the consensus sequence was determined
by picking the nucleotide with highest quality as indicated in the quality line of the fastq files. If
'N' persisted in the final sequence, the reads were discarded. The PCR primers were trimmed so
that all the sequences of the same competition would start and end at the same location.

Reads that did not have the constant region, that could not be oriented or that could not
be demultiplexed were discarded. The remaining errors in the reads were corrected by sequence
clustering. We used starcode version 1.0 [ref. 52] with default parameters and allowing up to two
errors. The corrected reads were translated using the genetic code. Variants encoding the same
proteins were not merged; they were kept separate for downstream analyses. A running Docker
virtual machine with commented scripts to replay the whole the process is available for
download at https://hub.docker.com/r/gui11aume/epi/.

542
DNA sequence variant frequency calculation and data filtering

The total number of reads for 12 segments, 3 time points and 4 replicas are shown in
Supplementary Table 1. Genotypes frequencies are defined as the number of reads for a given
genotype divided by the total number of reads in that replicate. Mean frequency was calculated
over 4 replicas to be used in further analysis. However, to eliminate influence of outliers the
median was taken instead of mean if absolute difference between mean and median was greater
than the median value. Only genotypes present in both technical replicas of both biological
replicas with at least ten reads (summed across all time points) in each of them were kept.

551
Noise estimation

The major factors causing noise in genotype frequency measurements are sampling
errors, PCR amplification errors and genetic drift during the competition. For all of these factors,
the amount of error depends on the genotype frequency. Therefore, we estimated measurement
errors as the function of genotype frequency.

For a given segment, time point and a pair of biological or technical replicas for each
genotype we calculated the mean frequency and the squared difference of frequencies from these
two replicas. We sorted genotypes by mean frequency and grouped them such that each bin
contains 5000 genotypes. We calculated the average frequency and the average squared
difference in each bin. Additionally, squared error for frequency 0 was set equal to $\frac{1}{2}\cdot$

$\left(\left(\frac{0.5}{N_i}\right)^2 + \left(\frac{0.5}{N_j}\right)^2\right)$, where N_i and N_j are total read numbers in replicas i and j. Finally, by linear

interpolation we obtained dependencies of squared differences as a function of frequency, $s_{ij}^2(f)$,
where i and j are different replicas.

Using squared differences from pairwise comparison of replicas we can estimate variance
of mean frequency over four replicas. Let numerate replicas 1, 2, 3, 4 where 1, 2 are technical

replicas of the first biological repeat and 3, 4 are the technical replicas of the second biological
repeat. Errors coming from the competition (e.g.: genetic drift) are shared for replicas 1, 2 and
for replicas 3, 4. Let's call them Δf_{b_1} and Δf_{b_2} and their variances $\sigma_{b_1}^2$ and $\sigma_{b_2}^2$ respectively.
Technical errors of sampling from the population and from PCR are unique for each replica.
Let's call them $\Delta f_{t_i}, i = 1..4$ and their variances $\sigma_{t_i}^2, i = 1..4$ respectively. All variances are
function of frequency and writing $\sigma_{X_i}^2$ we assume $\sigma_{X_i}^2(f)$.
573   In the introduced notations the mean frequency over 4 replicas is:

$$f = \frac{1}{4} \cdot (f_1 + f_2 + f_3 + f_4) =$$

$$\frac{1}{4} \cdot \left(\left(f^* + \Delta f_{b_1} + \Delta f_{t_1}\right) + \left(f^* + \Delta f_{b_1} + \Delta f_{t_2}\right) + \left(f^* + \Delta f_{b_2} + \Delta f_{t_3}\right) + \left(f^* + \Delta f_{b_2} + \Delta f_{t_4}\right) \right) =$$

574    $$f^* + \frac{1}{2} \cdot \left(\Delta f_{b_1} + \Delta f_{b_2}\right) + \frac{1}{4} \cdot \left(\Delta f_{t_1} + \Delta f_{t_2} + \Delta f_{t_3} + \Delta f_{t_4}\right),$$

575   where f^* is the true frequency. Applying basic properties of variance the variance of mean
576   frequency:
577

$$\sigma^2 = \frac{1}{4} \cdot \left(\sigma_{b_1}^2 + \sigma_{b_2}^2\right) + \frac{1}{16} \cdot \left(\sigma_{t_1}^2 + \sigma_{t_2}^2 + \sigma_{t_3}^2 + \sigma_{t_4}^2\right)$$

578
579   To estimate $\sigma_{b_1}^2, \sigma_{b_2}^2, \sigma_{t_1}^2, \sigma_{t_2}^2, \sigma_{t_3}^2, \sigma_{t_4}^2$ we used squared differences from pairwise comparison
580   of replicas calculated above $s_{12}^2, s_{13}^2, s_{14}^2, s_{23}^2, s_{24}^2, s_{34}^2$:

$$\mathbb{E}[s_{12}^2] = \mathbb{E}[(\Delta f_{t_1} - \Delta f_{t_2})^2] = \sigma_{t_1}^2 + \sigma_{t_2}^2$$
$$\mathbb{E}[s_{13}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_1} - \Delta f_{b_2} - \Delta f_{t_3})^2] = \sigma_{b_1}^2 + \sigma_{t_1}^2 + \sigma_{b_2}^2 + \sigma_{t_3}^2$$
$$\mathbb{E}[s_{14}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_1} - \Delta f_{b_2} - \Delta f_{t_4})^2] = \sigma_{b_1}^2 + \sigma_{t_1}^2 + \sigma_{b_2}^2 + \sigma_{t_4}^2$$
$$\mathbb{E}[s_{23}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_2} - \Delta f_{b_2} - \Delta f_{t_3})^2] = \sigma_{b_1}^2 + \sigma_{t_2}^2 + \sigma_{b_2}^2 + \sigma_{t_3}^2$$
$$\mathbb{E}[s_{24}^2] = \mathbb{E}[(\Delta f_{b_1} + \Delta f_{t_2} - \Delta f_{b_2} - \Delta f_{t_4})^2] = \sigma_{b_1}^2 + \sigma_{t_2}^2 + \sigma_{b_2}^2 + \sigma_{t_4}^2$$
$$\mathbb{E}[s_{34}^2] = \mathbb{E}[(\Delta f_{t_3} - \Delta f_{t_4})^2] = \sigma_{t_3}^2 + \sigma_{t_4}^2$$

581   Therefore, the variance of mean frequency f can be found as:

$$\sigma^2 = \frac{1}{16} \cdot \left((s_{13}^2 + s_{14}^2 + s_{23}^2 + s_{24}^2) - (s_{12}^2 + s_{34}^2) \right)$$

582   Recalling that variance and squared differences are a function of frequency:

$$\sigma^2(f) = \frac{1}{16} \cdot \left((s_{13}^2(f) + s_{14}^2(f) + s_{23}^2(f) + s_{24}^2(f)) - (s_{12}^2(f) + s_{34}^2(f)) \right)$$

583   For each segment and time point we calculated the numerical function $\sigma^2(f)$. Then for each
584   genotype having mean frequency f_x we estimated its variance as $\sigma^2(f_x)$
585
586   ***Merging amino acid genotypes***
587   We merged nucleotide genotypes that corresponded to the same amino acid sequence and
588   summed their frequencies and variances. We filtered out all genotypes x which had any of
589   following patterns:
590    $f_x^{t_0} = 0, f_x^{t_1} = 0, f_x^{t_2} > 0$ or $f_x^{t_0} = 0, f_x^{t_1} > 0, f_x^{t_2} = 0$ or $f_x^{t_0} > 0, f_x^{t_1} = 0, f_x^{t_2} > 0$. Fraction of such
591   genotypes were <0.5% for all segments except S9, for which it was 4.5%
592   For further analysis, this amino acid dataset was used except when specified.
593
594   ***Fitness estimation***
595   Number of cells in a pool with particular genotype x after time interval t increases
596   exponentially

597    $$n_x^t = n_x^0 \cdot Exp[s_x \cdot t],$$

598   where s_x is absolute fitness. Frequency of genotype x as well depends exponentially on absolute
599   fitness with an additional multiplicative factor:

$$600 \qquad f_x^t = \frac{n_x^t}{N^t} = \frac{n_x^0 \cdot Exp[s_x \cdot t]}{N^t} = \frac{f_x^0 \cdot Exp[s_x \cdot t]}{N^t/N^0},$$

601 where N^t and N^0 are total cell numbers in a pool at time points 0 and t. Factor $\frac{1}{N^t/N^0}$ reflects the
602 total growth of population, it changes with time but is the same for all genotypes. Therefore, we
603 can rewrite genotype frequency at time t as:

$$604 \qquad f_x^t = f_x^0 \cdot Exp[(s_x - s^{0t}) \cdot t],$$

605 where $s^{0t} = \frac{1}{t} \cdot Log\left(\frac{N^t}{N^0}\right)$

606   In the measured dataset for each genotype x we have 3 measurements of frequency $f_x^{t_0}$,
607 $f_x^{t_1}$, $f_x^{t_2}$ and their errors $\sigma^2(f_x^{t_0}), \sigma^2(f_x^{t_1}), \sigma^2(f_x^{t_2})$. To estimate genotype fitness we minimized
608 relative squared errors of exponential fit as function of fitness s_x and initial frequency f_x^0:

$$609 \qquad (s_x, f_x^0) = \text{argmin}_{s_x, f_x^0}\left(\frac{\left(f_x^{t_0} - f_x^0\right)^2}{\sigma^2\left(f_x^{t_0}\right)} + \frac{\left(f_x^{t_1} - f_x^0 \cdot Exp[(s_x - s^{01}) \cdot t_1]\right)^2}{\sigma^2\left(f_x^1\right)} + \frac{\left(f_x^{t_2} - f_x^0 \cdot Exp[(s_x - s^{02}) \cdot t_2]\right)^2}{\sigma^2\left(f_x^2\right)}\right) \qquad (1)$$

This formula contains four parameters common for all genotypes from one segment:
s^{01}, s^{02}, t_1, t_2. Further we will perform additional shifting and scaling of fitness values (see next
section), therefore, without loss of generality we could set $s^{01} = 0$ and $t_1 = 1$. Ideally, t_2/t_1 should
equal 14, however, we noticed that this ratio does not hold for many segments and fitted
$k = t_2/t_1$ from data instead of using value 14.

To find specific s^{02} and k for each segment we selected genotypes with high frequencies
at t_0 ($t_0 > 25 \cdot 10^{-6}$) which corresponds to ~500-1000 reads per technical replicate. Each
segment contains 10^3-10^4 genotypes that meet this criterion. We minimized eq. (1) for selected
genotypes trying all possible combinations of (s^{02}, k) from a grid where $s^{02}\epsilon[0,1.2]$ with step
0.01 and $k\epsilon[1,14]$ with step 0.1 and choose (s^{02}, k) which gives minimal (*).

Finally, given (s^{02}, k) for each segment we found s_x for each genotype. Errors for fitness
values, s_x^{std}, were estimated as standard error of best-fit parameter.

For genotypes with frequencies pattern $f_x^{t_0} > 0, f_x^{t_1} = 0, f_x^{t_2} = 0$ fit of eq. (1) cannot be
obtained. Therefore we defined upper boundary for their fitness value as $s_x^{boundary} =$
$Log\left(\frac{1}{\max(N_1^{t_1}, N_2^{t_1}, N_3^{t_1}, N_4^{t_1})}\right)$, where $N_i^{t_1}, i = 1..4$ are total read numbers at time point t_l in i replica.

626
Fitness rescaling
We scaled fitness such that lethal genotypes have fitness 0 and neutral genotypes have
fitness 1. We assumed that genotypes with a stop codon or frame shift are lethal. Thus, for each
segment we linearly rescaled the fitness distribution so that 95% of genotypes with nonsense
mutations have a fitness of 0 and so that the local maximum of the fitness distribution of
genotypes with extant amino acids is 1. The scaling around the local maximum led to the shift of
fitness values of less than +/- 0.025 in each of the 12 segments compared to the measured
wildtype strains and did not affect our results (for scale, we called a substitution non-neutral if its
effect on fitness was > 0.4). All fitness values which became smaller than 0 were set to 0.

636
Quality control and comparison of synonymous sequences
We used nucleotide synonymous sequences as an internal control. The error rate for a
measurement of fitness of an amino acid sequence depends on the number of synonymous
sequences, *n*, that were used to estimate it. Therefore, we estimated the false discovery rates
separately for categories with *n=1,..10* variants. For each amino acid genotype with more than *n*

synonymous variants we merged random combination of n of its nucleotide genotypes and
estimated fitness. We then calculated the difference between this fitness and the fitness of the
corresponding amino acid sequence. We classified case as "false unfit" if difference was <-0.4
and as "false fit" if difference was >0.4. The fraction of such cases gives us false discovery rates
for genotypes having n synonymous variants. To get total false discovery rates for each segment
we averaged "false unfit" and "false fit" rates for different n with weights equal to the fractions
of genotypes in amino acid dataset which have n synonymous variant (**Supplementary Table 1**).
The high correlation between biological replicas (**Supplementary Table 6**) confirms high
accuracy of our high-throughput experiments, with the exception of segment 9.
651
Predicting fitness using deep learning
To predict the unidimensional fitness function based on additive contribution of extant
amino acid states we used deep learning, a powerful machine learning technique, capable of
constructing virtually any function, even with a simple neural network architecture. To convert
amino acid sequences into a binary feature matrix we used one-hot encoding strategy, in which
each feature (column in the matrix) indicates the presence or absence of a particular amino acid
state.
To optimise the accuracy/overfitting ratio, we tested over a hundred of different neural
network architectures and parameters. As a starting point, we selected a number of complicated
architectures, which describe our data, but are prone to overfitting due to a large number of
parameters. We then gradually reduced the number of layers and neurons to reduce the
overfitting, while controlling for accuracy.
Our final architecture consists of three layers and 22 neurons in total (**Supplementary Fig.**
4). Each of the neurons performs a linear transformation of the input and subsequently applies a
non-linear activation function (a sigmoid) to the result. The output of the first layer, therefore, is
a sigmoid of a linear transformation of the feature vector ($c_1^T x$). The second layer decompresses
the hidden nonlinear representation into 20 sigmoids, the combination of which is further
linearly transformed with the only neuron of the third layer and wrapped into another sigmoid
function:
671

$$F(x) = \sigma(\sum_{i=1}^{20} c_{3,i}\sigma(c_{2,i}\sigma(c_1^T x + b_1) + b_{2,i}))$$

where $\sigma(t) = \frac{1}{(1+exp(-t))}$, x – is the feature vector, c_1 – the vector of coefficients, corresponding to
the first layer, $c_{n,k}$ - coefficient corresponding to the n-th layer and the k-th neuron, b_n - bias
corresponding to the n-th layer. Crucially, this relatively simple architecture is capable of fitting
virtually any function[53], thus, in contrast to conventional logistic regression, in our approach we
select the correct model from a vast variety of functions.
The key idea of our approach is that the number of neurons in the first layer of the neural
network determines the number of linear combinations of mutations (or fitness potentials) used
in order to predict mutant fitness. In other words, each neuron in the first layer assigns a single
unique weight to every amino acid state in the dataset (**Supplementary Fig. 4**). Multiplication of
such weight vectors and binary genotype vectors result in fitness potential. Thus, the number of
neurons in the first layer of the architecture basically determines the dimensionality of epistasis
we assume. The obtained fitness potentials are then transformed by a nonlinear phase shift
function constructed by the 22 neurons of the neural network.
The architecture simplicity avoids overfitting, which was further prevented by using early
stopping and keeping 10% of data as a test set. The loss function that is being optimised in our
experiments is not convex, which leads to a high probability of getting stuck in different local
minima. To ensure reproducibility, each of our models was constructed ten independent times
using random train-test splits.

Each model was trained for under 100 epochs using mean squared error as the loss
function. An unpublished adaptive learning rate method proposed by Geoff Hinton, RMSProp,
was used as the optimiser. This algorithm is a version of a mini-batch stochastic gradient
descent, utilising the gradient magnitude of the recent gradients in order to normalise the current
ones. All the weights were initialised using Xavier normal initialiser[54].

Paths between pairs of fit genotypes

For analysis in **Fig. 6d**, we first choose two fit "parental" genotypes, one randomly
chosen genotype (eg: ABE) and the other parental genotype that is either *S. cerevisiae* wildtype
genotype (inter-segmental) or another random fit genotype in the data (intra-segmental) (eg:
abe). The two genotypes in this example are Hamming Distance 3 apart (HD=3). We next
compute all (HD2-2) intermediate genotypes (eg: AbC, aBc, *et cetera*) and retain the subset that
were experimentally measured. We represent the two parental genotypes and all measured
intermediate genotypes as an undirected graph in which each genotype is a vertex. All genotypes
one substitution apart are connected by an unweighted edge. The shortest possible path for a
given pair of genotypes is of length HD. We find all shortest paths between the two parental
genotypes using a breadth-first search. We next remove all vertices (genotypes) that are unfit,
and recompute the number of shortest between the two parental genotypes. For example, in **Fig.**
6a, there are six paths of length three if you take into account all genotypes, but only three paths
of length three if you take into account only fit genotypes.

Clustering of unfit genotypes in sequence space

For the analysis in **Fig. 6e**, we first represent the two parental genotypes and all measured
intermediate genotypes as an undirected graph in which each genotype is a vertex. All genotypes
one substitution apart are connected by an unweighted edge. We can then compute the degree
(number of genotypes of distance one) for each vertex (genotype). We do so randomly drawing
from all measured genotypes and using only unfit genotypes or using the same number but
randomly chosen genotypes. For the randomly chosen genotypes, the value is the average over
1000 runs.

Quantifying sign epistasis

For each substitution (eg: C -> S at position 141), we considered only those that exhibit a
large fitness effect (abs. difference > 0.4) comprising a set of substitutions with large effects. For
each substitution we divided the genetic backgrounds into two categories: those in which the
substitution caused a > 0.4 increase in fitness, and those backgrounds in which the substitution
caused > 0.4 decrease in fitness. A single substitution can cause a large increase in fitness in
some backgrounds and a large decrease in others due to two possible reasons: sign epistasis or
experimental error. To differentiate the two cases, we identified secondary substitutions that
significantly alter the ratio of large increases to large decreases in fitness (Fisher's exact test,
Bonferroni corrected p-value < 0.05). We only consider a site to be under sign epistasis if there
is a second site that alters the frequency of sign epistasis in a statistically significant manner, i.e.
more frequently than expected by chance alone.

Structural analysis

Structure prediction

An initial model was obtained with the I-TASSER server[55]. The list of top 10 PDB
structural templates picked up by the I-TASSER included high-quality crystal structures of
imidazoleglycerol-phosphate dehydratases from *Arabidopsis thaliana* and *Cryptococcus*
neoformans. Coordinates of the top-scoring model (C-score=0.21, estimated TM-score =
0.74±0.11, estimated RMSD = 5.1±3.3Å) and the predicted normalized B-factor[56] were used for
further analysis. The value of the model quality metric (TM-score >0.5) indicates a model of

correct topology. The proteins structurally close to the final model (RMSD 0.6 - 1.7Å are PDB
IDs 4MU0, 4GQU, 1RHY, 5DNL and 2AE8 from *Arabidopsis thaliana*, *Mycobacterium*
tuberculosis, *Cryptococcus neoformans*, *Pyrococcus furiosus* and *Staphylococcus aureus*.
749   We measured the distribution of distances (in angstroms) between pairs of residues that
750   exhibit strong sign epistasis (**Supplementary Table 1**, ReallyPositivePair == TRUE), and
751   compared it with the distribution of pairwise distances among residues for which we have
752   sufficient data to be certain that a given pair does not exhibit sign epistasis (**Supplementary**
753   **Table 1**, ReallyNegativePair == TRUE).
754
755   *ΔΔG prediction*
756   Cartesian_ddg application[57] from Rosetta version 2017.08.59291 was used for ΔΔG
predictions. Top-scoring I-TASSER model was pre-minimized using the Relax[58] application in
dual-space[59] with the flags: -relax:dualspace true; -ex1; -ex2; -use_input_sc; -flip_HNQ; -
no_optH false; -relax:min_typelbfgs_armijo_nonmonotone; -nonideal. The best scoring model
from 1000 structures was selected. The effect of up to 4 mutations (54,500 genotypes in total)
was assessed in Cartesian space with the Talaris_2014 score function, and the -fa_max_dis 9.0
flag. ΔΔG was estimated as a difference of mean score for 3 independent runs for every mutant
and the wild-type score.
764
765

766
767 **Figures**
768
Figure 1. Combinatorial approach to the study of fitness landscapes. A fitness landscape is
the representation of fitness for all possible genotypes composed of a specific set of loci. **a,**
Following Figure 1 from Sewall Wright ref. [6] consider the genotype space consisting of 5 loci,
each with two allele states (lower and uppercase letters). The entire genotype space is 5-
dimensional consisting of 2^5 genotypes. Given two genotypes found in extant species (abCde and
ABCdE in this example), surveying combinations of extant alleles substantially reduces the
dimensionality of the genotype space, concomitantly reducing the number of genotypes to assay.
The surveyed area (blue cube) considers all combinations of allele substitutions that have
occurred in the course of evolution between the two sequences (red line), avoiding the sampling
of combinations with less relevance to the evolutionary trajectory (black lines). **b,** Given the
entire multidimensional genotype space (black circle) our approach considers an
multidimensional subspace consisting of the combinatorial set of amino acid states from extant
species. The blue line represents the yeast phylogeny and the surrounding blue space represents a
multidimensional set of combinations of extant amino acids of the sequence under consideration,
one His3 gene segment in our study. By contrast, random mutagenesis studies consider only a
local segment of the genotype space surrounding a specific genotype (green circle). **c**, A multiple
alignment of orthologous sequences of His3 for segment 2 for which we incorporated almost all
extant amino acid states from 21 yeast species (blue bars) and 10-100% extant states from a set
of 396 orthologues (grey bars). **d**, The predicted structure of His3p with amino acid residues that
were substituted in our library.
789
Figure 2. Visual representations of the fitness landscape. a, The fitness landscape for all
assayed genotypes in segment 7. Nodes represent unique amino acid sequences with edges
connecting those separated by a single amino acid substitution. Colour saturation represents the
minimum fitness of the two connected nodes. **b,** For segment 7, fitness of ancestral and extant
nodes and genotypes one substitution away from the nodes in the background of *S. cerevisiae*
gene on the yeast phylogeny (black lines), are shown in colour ranging from grey (lowest fitness)
to blue (highest fitness).
797

798
799
Figure 3. Fitness distributions. a, The distribution of fitness for genotypes composed of
combination of extant amino acid states (green) and non-extant amino acid states (purple) at the
same positions. **b**, The fraction of unfit genotypes per segment among genotypes consisting
entirely from extant amino acid states (green) and those incorporating non-extant amino acid
states (purple). **c**, The number of genotypes with high fitness that incorporates specific amino
acid states. For each amino acid state, the number of genetic backgrounds that contain that amino
acid state and are fit (fitness > 0) are shown. **d**, The percent of backgrounds in which a specific
substitution is neutral (white), beneficial (blue) or deleterious (grey). The region marked in green
shows substitutions that never have large effects (> 0.4) on fitness. Beneficial and deleterious
effects are shown only if the frequency for a given substitution was higher than the false
discovery rate (**Supplementary Table 1**). Data from segment 9 were excluded for this figure.
811
812

813
814
Figure 4. Epistasis and the His3 fitness landscape for segments 2, 5 and 7. **a**, Fitness as a
function of a single fitness potential (black curve, the fitness of individual genotypes is orange).
b, A network depiction of sign epistasis between amino acid substitutions. Colour coded sites
with reciprocal sign epistasis (black lines) and unidirectional interactions (grey arrows) are
shown. **c**, Genotypes containing substitutions with a higher number of sign epistatic interactions
are less likely to be fit by the threshold function of the fitness potential. **d**, Increasing the number
of neurons in the first layers of the neural network, which is equivalent to increasing the number
of underlying fitness potentials, leads to more accurate models for segments with detected sign
epistasis. Each dot corresponds to an independent optimization of model parameters. **e**, Fitness
as a function of two fitness potentials (black dots, measured fitness is depicted in orange).
825
Figure 5. Sign epistasis. a, Substitution C->S at site 141 in segment 2 more frequently has a
positive effect on fitness in the background of T at site 143, a negative effect in the background
of 143I and is equally likely to be strongly deleterious or strongly beneficial in the background
of 143V. **b**, Predicted change in free energy following a C141S substitution in all genetic
backgrounds with an I or T at 143 and that are closer than four mutations away from *S.*
cerevisiae. **c**, Distribution of the number of substitutions at each site under sign (yellow) and
reciprocal sign (orange) epistasis. Sites with 0 interactions do not exhibit sign epistasis. **d**, The
fraction of genotypes in which the substitution under sign epistasis has the less frequent effect on
fitness.
835

836
837
Figure 6. Analysis of evolutionary pathway accessibility. a, A threshold fitness potential
function can lead to some paths being inaccessible between two genotypes of high fitness (abe,
ABE) if the joint contribution of several alleles to the fitness potential (abE, aBE) leads to the
fitness potential below the threshold. **b,** The fraction of unfit intermediate genotypes between
two fit genotypes as a function of their average fitness potential. **c**, The grey area represents all
genotypes in segment 7. When two fit genotypes (red dots) have high fitness potential, many
paths between them will be accessible because many intermediate genotypes will also have high
fitness potential and fitness (blue dots). **d**, The fraction of accessible shortest paths between two
fit genotypes with unfit genotypes from data (orange) or the same number of randomly selected
genotypes (grey), shown as a function of Hamming distance between two fit genotypes. Error
bars are standard deviation. **e**, On a graph with genotypes connected by edges if they are one
substitution apart we calculate the degree of connectivity (number of edges for each node) for all
genotypes (blue), only unfit (fitness = 0) genotypes (orange) and a graph with the same number
of nodes as the graph with unfit genotypes but with nodes chosen at random (grey).
852
853

854
855 **Supplementary Figures**
856
Supplementary Figure 1. Experimental design. a, The sequence of the His3 protein from S.
cerevisiae was separated into 12 independent segments of similar lengths, such that the full
combinatorial set of extant amino acid substitutions was less than 100,000 possible genotypes.
These segments represented different combinations of structural elements of the His3 protein
structure. **b,** For each of the 12 segments from His3, we selected extant amino acid states using a
multiple alignment of His3 orthologues from 396 species, preferentially incorporating states
from 21 yeast species, the variability is shown in segment 3 as an example. Mutant degenerate
codon libraries were constructed by fusion PCR of two synthesized variable halves of each
segment. These high-complexity plasmid libraries were transformed into haploid His3 knockout
S. cerevisiae strain. The growth rate of yeast carrying different extant amino acid state
combinations in His3 gene was measured using serial batch culture in the absence of histidine
with 12 hours between ~100-fold dilutions. To estimate the fitness of yeast mutants their relative
abundance was measured at three points: in the initial population before selection (t0), in the
population after 12 hours of growth in the selective medium (t1), and in the final population after
168 hours of growth in the selective medium (t14). To assess the fitness of individual mutants
the segments from three populations were amplified and sequenced. The relative abundance of
each sequence was used as a proxy for abundance of the associated yeast mutant, which in turn
determines its fitness. **c**, Secondary structure of His3 mapped to the segments in our
experiments.
876
Supplementary Figure 2. Segment-specific fitness distributions for extant and non-extant
amino acid states. a, The fitness distribution for each segment for genotypes consisting only of
extant amino acid states (green) or that contain one or more non-extant amino acid states
(purple) only at positions with a substitution in the extant library. **b,** The fitness distribution for
each segment for genotypes consisting only of extant amino acid states (green) and genotypes
with mutations at other positions in that segment (red).
883
Supplementary Figure 3. Growth rate measurement of isolated strains. a, Comparison of
fitness values from the pooled competition assay with growth rates of isolated strains as
measured in a microplate reader. Error bars for growth rates show s.e.m. of 6 replicates. **b**,
Pearson correlation coefficients between fitness values from competition and growth rates of
isolated strains for each segment. ** signifies p-value < 0.005 (correlation test). **c**, His3p
orthologues from different species complement a Δ his3 deletion in *S. cerevisiae*. Growth rates
of transformants containing whole HIS3 orthologous genes from other yeast species. Error bars
for growth rates show s.e.m. of ≥ 7 replicates.
892
893 **Supplementary Figure 4. Schematic representation of the deep learning approach.** Each
894 genotype was encoded as a binary vector (x). During training, each of the substitutions was
895 assigned a coefficient (c_i), comprising a vector of coefficients (c). The multiplication of these
896 two vectors is the fitness potential of the genotype. After going through three layers, each with a
897 sigmoid activation function, the predicted fitness is obtained.
898
899 **Supplementary Figure 5. The fitness potential of the His3 fitness landscape. a**, Fitness
900 potential predicted by the neural network as a function of the measured fitness for all 12
901 segments. **b**, The correlation between the fitness predicted by the fitness potential and the
902 measured fitness. **c,** Training and test R^2 for each segment for 20-fold cross-validation.
903
904

905
906 **Supplementary Figure 6. Sign epistasis dimensionality graphs for all twelve segments.** Each
907 node represents a substitution, with multiple substitutions at the same site having the same
908 colour. Substitutions under reciprocal sign epistasis are indicated by black lines while grey
909 arrows indicate unidirectional sign epistasis.
910

911
Supplementary Figure 7. Multidimensional description of epistasis in His3 segments. a,
Increasing the number of neurons in the first layer of the neural network, which is equivalent to
increasing the number of underlying fitness potentials, leads to more accurate models for
segments with detected sign epistasis. Each dot corresponds to an independent optimization of
model parameters. **b,** Number of sign epistatic interactions of certain substitutions against
average model prediction power for mutants including these substitutions.
918
Supplementary Figure 8. Protein stability and the fitness potential. a, A comparison of
correlation coefficients between predicted and measured values across segments. **b,c,**
correlations between the estimated impact of substitutions on free energy ($\Delta\Delta G$), fitness
potential and fitness. $\Delta\Delta G$ correlates better with fitness potential than with fitness. **d**, Pairs of
sites that exhibit sign (connected by a light edge in Supplementary Figure 6) and those that
exhibit reciprocal sign epistasis (connected by a dark edge in Supplementary Figure 6) are closer
together in the His3p structure that randomly chosen non-connected pairs of positions that
exhibit sign epistasis.
927
Supplementary Figure 9. Decoupling inter- and intra-segmental epistasis. a, The fraction of
unfit genotypes between *S. cerevisiae* and any other genotype consisting of extant amino acid
states with high (blue) or any (red) fitness, and genotypes in the latter but not the former
category (black) as a function of the Hamming distance between the two boundary genotypes.
Points indicate median, the bars and lines indicate 50% of the genotypes and genotypes 2.7
sigmas from the mean, respectively. **b,** The neural network model assigns higher weights to
amino acid states that that first occur in His3 orthologues farther from *S cerevisiae,* indicating
the presence of intrasegmental interactions.
936
937 **Supplementary Information 1. Multiple alignment of His3 orthologues.**
938
939 **Supplementary Information 2. Multidimensional description of epistasis in His3**
940 **segments.** Fitness as a function of two fitness potentials (black dots, measured fitness is depicted
941 in orange).
942
943

References

1. Dean, A. M. & Thornton, J. W. Mechanistic approaches to the study of evolution: the functional synthesis. *Nature Rev. Genet.* **8**, 675–688 (2007).

2. Kogenaru, M., de Vos, M. G. J., & Tans, S. J. Revealing evolutionary pathways by fitness landscape reconstruction. *Crit. Rev. Biochem. Mol. Biol.* **44**, 169–174 (2009).

3. Mackay, T. F. C. Epistasis and quantitative traits: using model organisms to study gene-gene interactions. *Nature Rev. Genet.* **15**, 22–33 (2014).

4. de Visser, J. A. G. M. & Krug, J. Empirical fitness landscapes and the predictability of evolution. *Nature Rev. Genet.* **15**, 480–490 (2014).

5. de Visser, J. A. G. M., Cooper, T. F., & Elena, S. F. The causes of epistasis. *Proc. Biol. Sci.* **278**, 3617–3624 (2011).

6. Wright, S. The roles of mutation, inbreeding, crossbreeding and selection in evolution. *Proc. Sixth Int. Congr. Genet.* **1**, 356–366 (1932).

7. Olson, C. A., Wu, N. C. & Sun, R. A comprehensive biophysical description of pairwise epistasis throughout an entire protein domain. *Curr. Biol.* **24**, 2643–2651 (2014).

8. Puchta, O., Cseke, B., Czaja, H., Tollervey, D., Sanguinetti, G., and Kudla, G. Network of epistatic interactions within a yeast snoRNA. *Science*. **352**, 840–844 (2016).

9. Li, C. Qian, W. Maclean, C. J. & Zhang, J. The fitness landscape of a tRNA gene. *Science* **352**, 837–840 (2016).

10. Sarkisyan, K.S., et al. Local fitness landscape of the green fluorescent protein. *Nature* **533**, 397–401 (2016.)

11. Hietpas, R. T., Bank, C., Jensen, J. D. & Bolon, D. N. A. Shifting fitness landscapes in response to altered environments. *Evolution* **67**, 3512–3522 (2013).

12. Parera, M. & Martinez, M. A. Strong epistatic interactions within a single protein. *Mol. Biol. Evol.* **31**, 1546–1553 (2014).

13. Podgornaia, A. I. & Laub, M. T. Pervasive degeneracy and epistasis in a protein-protein interface. *Science* **347**, 673–677 (2015).

14. Anderson, D. W. McKeown, A. N. & Thornton, J. W. Intermolecular epistasis shaped the function and evolution of an ancient transcription factor and its DNA binding sites. *Elife* **4**, e07864 (2015).

15. Meini, M. R., Tomatis, P. E., Weinreich, D. M. & Vila, A. J. Quantitative description of a protein fitness landscape based on molecular features. *Mol. Biol. Evol*. **32**, 1774–1787 (2015).

994 16. Bank, C., Matuszewski, S., Hietpas, R. T. & Jensen, J. D. On the (un)predictability of a large
995 intragenic fitness landscape. *Proc. Natl. Acad. Sci. USA*. 113, 14085–14090 (2016).
996
997 17. Poelwijk, F.J., Kiviet, D.J., Weinreich, D.M., & Tans, S.J. Empirical fitness landscapes
998 reveal accessible evolutionary paths. *Nature* **445**, 383–386(2007).
999
1000 18. Gong, L. I. Suchard, M. A.& Bloom, J.D. Stability-mediated epistasis constrains the
1001 evolution of an influenza protein. *Elife* **2**, e00631(2013).
1002
1003 19. Gong, L. I.& Bloom, J.D. Epistatically interacting substitutions are enriched during adaptive
1004 protein evolution. *PLoS Genet.* **10**, e1004328(2014).
1005
1006 20. Starr, T. N. & Thornton, J. W. Epistasis in protein evolution. *Protein Sci.* **25**, 1204–1218
1007 (2016).
1008
1009 21. Kumar, A. Natarajan, C. Moriyama, H. Witt, C.C. Weber, R.E. Fago, A.& Storz, J.F.
1010 Stability-mediated epistasis restricts accessible mutational pathways in the functional evolution
1011 of avian hemoglobin. *Mol. Biol. Evol.* **34**, 1240–1251(2017).
1012
1013 22. Tufts, D. M., *et al.* Epistasis constrains mutational pathways of hemoglobin adaptation in
1014 high-altitude pikas. *Mol. Biol. Evol.* **32**, 287–298 (2015).
1015
1016 23. Eyre-Walker A. & Keightley P. D. The distribution of fitness effects of new mutations. *Nat.*
1017 *Rev. Genet.* 8, 610–618 (2007).
1018
1019 24. Draghi, J. A. & Plotkin, J. B. Selection biases the prevalence and type of epistasis along
1020 adaptive trajectories. *Evolution* **67**, 3120–3131 (2013).
1021
1022 25. Lunzer, M., Miller, S. P., Felsheim, R. & Dean, A. M. The biochemical architecture of an
1023 ancient adaptive landscape. *Science* **310**, 499–501 (2005).
1024
1025 26. Palmer, A. C., Toprak, E., Baym, M., Kim, S., Veres, A., Bershtein, S. & Kishony, R.
1026 Delayed commitment to evolutionary fate in antibiotic resistance fitness landscapes. *Nat.*
1027 *Commun.* **6**, 7385 (2015).
1028
1029 27. Lalić, J. & Elena, S. F. Magnitude and sign epistasis among deleterious mutations in a
1030 positive-sense plant RNA virus. *Heredity* **109**, 71–77 (2012).
1031
1032 28. Keefe, A. D. & Szostak, J. W. Functional proteins from a random-sequence library. *Nature*
1033 **410**, 715–718 (2001).
1034
1035 29. du Plessis, L. Leventhal, G.E. & Bonhoeffer, S. How good are statistical models at
1036 approximating complex fitness landscapes? *Mol. Biol. Evol.* **33**, 2454–2468(2016).
1037
1038 30. Lum, P. Y. Edwards, S.& Wright, R. Molecular, functional and evolutionary characterization
1039 of the gene encoding HMG-CoA reductase in the fission yeast, *Schizosaccharomyces pombe.*
1040 *Yeast* **12**, 1107–1124 (1996).
1041
1042 31. Kondrashov, A.S. Sunyaev, S. & Kondrashov, F. A. Dobzhansky-Muller incompatibilities in
1043 protein evolution. *Proc. Natl. Acad. Sci. USA* **99**, 14878–14883 (2002).
1044

1045 32. Kvitek, D. J. & Sherlock, G. Reciprocal sign epistasis between frequently experimentally
1046 evolved adaptive mutations causes a rugged fitness landscape. *PLoS Genet.* **7**, e1002056 (2011).
1047
1048 33. Breen, M. S., Kemena, C., Vlasov, P. K., Notredame, C. & Kondrashov, F. A. Epistasis as
1049 the primary factor in molecular evolution. *Nature* **490**, 535–538 (2012).
1050
1051 34. Gavrilets, S. Evolution and speciation on holey adaptive landscapes. *Trends Ecol. Evol.* **12**,
1052 307–312(1997).
1053
1054 35. Salverda, M. L. M., Dellus, E., Gorter, F. A., Debets, A.J. M., van der Oost, J. Hoekstra, R.
1055 F. Tawfik, D. S. & de Visser, J.A.G. M. Initial mutations direct alternative pathways of protein
1056 evolution. *PLoS Genet.* **7**, e1001321(2011).
1057
1058 36. Weinreich, D. M., Delaney, N. F., de Pristo, M. A. & Hartl, D.L. Darwinian evolution can
1059 follow only very few mutational paths to fitter proteins. *Science* **312**, 111–114(2006).
1060
1061 37. Milkman, R. Selection differentials and selection coefficients. *Genetics* **88**, 391–403 (1978).
1062
1063 38. Kimura, M. & Crow J. F. Effect of overall phenotypic selection on genetic change at
1064 individual loci. *Proc. Natl. Acad. Sci. USA*. **75**, 6168–6171 (1978).
1065
1066 39. Kondrashov, F. A. & Kondrashov, A. S. Multidimensional epistasis and the disadvantage of
1067 sex. *Proc. Natl. Acad. Sci. USA*. **98**, 12089–12092 (2001).
1068
1069 40. Crow, J. F. & Kimura, M. Efficiency of truncation selection. *Proc. Natl. Acad. Sci. USA* **76**, 396–
1070 399 (1979).
1071
1072 41. Weinreich, D. M. Lan, Y. Wylie, C. S.& Heckendorn, R. B. Should evolutionary geneticists
1073 worry about higher-order epistasis? *Curr. Opin. Genet. Dev.* **23**, 700–707 (2013).
1074
1075 42. Poelwijk, F.J. Krishna, V. & Ranganathan, R. The Context-Dependence of Mutations: A
1076 Linkage of Formalisms. *PLoS Comput. Biol.* **12**, e1004771 (2016).
1077
1078 43. Sailer, Z. R. & Harms, M. J. High-order epistasis shapes evolutionary trajectories. *PLoS*
1079 *Comput. Biol.* **13**, e1005541 (2017).
1080
1081 44. Weinreich, D. M. Watson, R. A. & Chao, L. Perspective: Sign epistasis and genetic
1082 constraint on evolutionary trajectories. *Evolution* **59**, 1165–1174 (2005).
1083
1083 45. Rokyta, D. R. et al. Epistasis between beneficial mutations and the phenotype-to-fitness Map
1084 for a ssDNA virus. *PLoS Genet*. **7**, e1002075 (2011).
1085
1086 46. Kondrashov, D. A. & Kondrashov, F. A. Topological features of rugged fitness landscapes in
1087 sequence space. *Trends Genet*. **31**, 24–33 (2015).
1088
1089 47. De Pristo, M. A. Weinreich, D. M. & Hartl, D. L. Missense meanderings in sequence space:
1090 a biophysical view of protein evolution. *Nature Rev. Genet.* **6**, 678–687 (2005).
1091
1091 48. Bershtein, S., Segal, M., Bekerman, R., Tokuriki, N. & Tawfik, D. S. Robustness epistasis
1092 link shapes the fitness landscape of a randomly drifting protein. *Nature* **444**, 929–932 (2006).
1093
49. Orr, H. A. The population genetics of speciation: the evolution of hybrid incompatibilities.
Genetics **139**, 1805–1813 (1995).

50. Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. MEGA6: Molecular
Evolutionary Genetics Analysis version 6.0. *Mol Biol Evol*. **30**, 2725–2729 (2013).

51. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve
genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

52. Zorita, E., Cuscó, P. & Filion, G. J. Starcode: sequence clustering based on all-pairs search.
Bioinformatics **31**, 1913–1919 (2015).

53. LeCun, Y., Bengio, Y. & Hinton, G. Deep learning. *Nature*. **521**, 436–444 (2015).

54. Glorot, X. & Bengio, Y. Understanding the difficulty of training deep feedforward neural
networks. In Proceedings of the 13th International Conference on Artificial Intelligence and
Statistics (2010).
1105
1106   55. Yang, J. & Zhang, Y. I-TASSER server: new development for protein structure and function
1107   predictions. *Nucleic Acids Res*. **43**, W174–W181(2015).

1108   56. Yang, J., Wang, Y. & Zhang, Y. ResQ: An Approach to Unified Estimation of B-Factor and
1109   Residue-Specific Error in Protein Structure Prediction. *J. Mol. Biol.* **428,** 693–701 (2016).
1110
57. Park, H., Bradley, P., Greisen, P. Jr., Liu, Y., Mulligan, V. K., Kim, D.E., Baker, D. &
DiMaio, F. Simultaneous optimization of biomolecular energy functions on features from small
molecules and macromolecules. *J. Chem. Theory Comput.* **12**, 6201–6212 (2016).

58. Nivón, L. G., Moretti, R. & Baker, D. A Pareto-optimal refinement method for protein
design scaffolds. *PLoS One* **8**, e59004 (2013).

59. Conway, P., Tyka, M. D., DiMaio, F., Konerding, D. E. & Baker, D. Relaxation of backbone
bond geometry improves protein energy landscape modeling. *Protein Sci.* **23**, 47–55 (2014).

A

B

His3

Random mutagenesis

C

His3 extant sequence variant library

% of variation measured from
- all orthologues
- yeast species

Fraction of sampled extant amino acid states

D

- 1 amino acid states assayed
- 2 amino acid states assayed
- 3 amino acid states assayed
- 4 amino acid states assayed
- 5 amino acid states assayed
- 6 amino acid states assayed
- 9 amino acid states assayed

A

B r = 0.77

% of unfit intermediates

parental genotypes
mean fitness potential

C

98%
2%
0%

Fitness

parent genotypes
intermediate genotypes

84%
14%
2%

Fitness

40%
26%
33%

Fitness

Fitness potential

D

% of shortest paths that are accessible

remove :
unfit genotypes
random genotypes

Hamming distance
between fit genotypes

E

all genotypes
only unfit
random

Mean node degree
(# connections of distance=1)

Segment

F

log2(unfit / random)

Segment

A

His3 gene divided into 12 interleaved segments

His3 gene of *S. cerevisiae*

Preparation of 12 libraries

Full combinatorial set of selected extant amino acids

36 864 in Segment 12
18 432 in Segment 11
32 768 in Segment 10
49 152 in Segment 9
65 536 in Segment 8
4 320 in Segment 7
55 296 in Segment 6
62 208 in Segment 1
55 296 in Segment 2
55 296 in Segment 3
73 728 in Segment 4
55 296 in Segment 5

B

1. Selection of one of the 12 His3 segments

His3 gene of *S. cerevisiae*

Segment 3

2. Selection of variability based on an alignment of sequences from extant species

L G L R E K Y G D L S C E M I P H F L E S F A Q A

Library complexity: 2 (L145) x 2 (L147) x 3 (Q148) x 2 (K151) x 2 (V152) x 3 (D154) x 3 (L164) x 4 (E165) x 2 (A168) x 4 (E169) x 4 (A170) = 55 296

Library construction technique

His3 gene
Fusion-PCR
His3 gene mutant

3. Library generation

(55 296 extant amino acid combinations constructed for segment 3)

4. Competition experiments in yeast

t0   t1   t14

12 h   156 h

5. Sequencing

A   A   C
B   A   C
C   C   C
D   C   C
E   C   C
F   C   C

(fitness measured for 48 198 genotypes from segment 3)

C

– Disordered region
– Alpha helix
– Beta-sheet
– Minimal distance to Mn less than 5 Å
– Minimal distance to substrate less than 5 Å

S5
S4
S5
S4
S3
S2
S3
S2
S1
S6
S1
S6
S7
S8
S7
S8
S9
S10
S9
S10
S11
S12
S11
S12

Positions according to Uniprot sequence

A

B C

A

B