Non-coding Transcriptome Maps Across Twenty Tissues of the Korean Black Chicken, Yeonsan Ogye

Hyosun Hong[1,6], Han-Ha Chai[3,4,6], Kyoungwoo Nam[1], Dajeong Lim[3], Kyung-Tai Lee[3], Yoon Jung Do[3], Chang-Yeon Cho[5] and Jin-Wu Nam[1,2,*]

[1]Department of Life Science, College of Natural Sciences, Hanyang University, Seoul 133791, Republic of Korea

[2]Research Institute for Convergence of Basic Sciences, Hanyang University, Seoul 133791, Republic of Korea

[3]Department of Animal Biotechnology & Environment of National Institute of Animal Science, RDA, Wanju 55365, Republic of Korea

[4]College of Pharmacy, Chonnam National University, Kwangju 61186, Republic of Korea

[5]Animal Genetic Resource Research Center of National Institute of Animal Science, RDA, Namwon 55717, Republic of Korea

[6]These authors contributed equally to this paper.

*Corresponding author

Tel: +82-2-2220-2428, Fax: +82-2-2298-0319, Email: jwnam@hanyang.ac.kr

Abstract

The Yeonsan Ogye (Ogye) is a rare chicken breed that populates the Korean peninsula. The

entire body of this bird, including its feathers and skin, has a unique black coloring. Although

some protein-coding genes related to this unique feature have been examined, non-coding

elements have not been globally investigated. In this study, high-throughput RNA

sequencing (RNA-seq) and reduced representation bisulfite sequencing (RRBS) were

performed to construct whole non-coding transcriptome maps across twenty different Ogye

tissues. The resulting maps included 6900 long non-coding RNA (lncRNA) genes comprising

1290 known and 5610 novel lncRNA genes. Compared to lncRNAs previously annotated in

the *Galllus gallus red junglefowl*, a considerable number were either fragments of protein-

coding genes or not expressed in Ogye tissues. Newly annotated Ogye lncRNA genes

showed tissue-specific expression and simple gene structures containing 2 or 3 exons.

Systematic analyses of sequencing data and other genomic data demonstrated that about

39% of the tissue-specific lncRNAs displayed evidence of function. In particular, heat shock

transcription factor 2 (HSF2)-associated lncRNAs were discovered to be functionally linked

to protein-coding genes that are specifically expressed in black skin tissues, tended to be

more syntenically conserved in mammals, and were differentially expressed in black tissues

relative to white tissues. Our findings and resulting maps provide not only a comprehensive

catalogue of lncRNAs but also a set of functional lncRNAs that will facilitate understanding

how the non-coding genome regulates unique phenotypes. Furthermore, our results should

be of use for future genomic breeding of chickens.

Keywords

lncRNA, Chicken, Black tissues, Tissue-specific genes, Non-coding transcriptome

Background

The Yeonsan Ogye (Ogye) chicken is one of the rarest breeds of *Gallus gallus domesticus*. Domesticated in the Korean peninsula, it probably originated from the Indonesian Ayam Cemani black chicken, which populates tropical, high-temperature areas [1]. Ogye shares common features—such as black plumage, skin, shank, and fascia—with Ayam Cemani [1], although it has a smaller comb and shorter legs. Silkie fowl (Silkie), one of the most popular black-bone chickens, also has black skin but has white or varied color plumage [2]. Several genes involved in Silkie skin hyperpigmentation have been reported in previous studies [2-4]. Recently, transcriptomes from Chinese native black chickens were compared with those from white chickens to globally identify hyperpigmentation-related genes [5]. However, studies of the molecular mechanisms and pathways related to black chicken hyperpigmentation have been restricted to coding genes.

A major part of the non-coding transcriptome corresponds to long non-coding RNAs (lncRNAs), which originate from intergenic, intervening, or antisense-overlapping regions of protein-coding genes [6-8]. lncRNAs are defined as transcripts longer than 200nt and are mostly untranslated because they lack an open reading frame; however, they interact with RNA binding proteins and have diverse intrinsic RNA functions [9-11]. They tend to be localized to subcellular areas, particularly the nucleus, and often interact with heterochromatin remodelers and DNA methylation regulators to regulate gene expression at the epigenetic level. For instance, DNMT1-associated colon cancer repressed lncRNA-1 (DACOR1) is localized to genomic sites, known to be differentially methylated, and regulates methylation at least 50 CpG sites by recruiting DNMT1 in colon cancers [12].

lncRNAs are also known to regulate gene expression at other levels: transcriptional, post-transcriptional, translational, and post-translational [9, 10, 13-15]. They regulate distant genes by modulating the recruitment of transcription factors (TFs) to target genes. Only a

few lncRNAs, however, have been experimentally validated as functional; most candidates

remain unvalidated. In particular, some lncRNAs have been shown to regulate the

expression of neighboring genes in a *cis*-acting manner [16-20]. Enhancer-associated

lncRNAs (eRNAs) are a well-known group in this class that regulate the expression of

downstream genes. Knockdown of eRNAs reduces target gene expression, suggesting their

function as *cis*-acting elements [21-23]. eRNA regulatory roles are known to be achieved via

several mechanisms: trapping transcription factors, directing chromatin roofing, and inducing

DNA methylation [9, 24-28]. On the other hand, lncRNAs that associate with post-

transcriptional regulators control target splicing and stability. For instance, antisense lncRNA

from the *FGFR2* locus promotes cell-type specific alternative splicing of FGFR2 by

interacting with polycomb complex [29].

Despite their regulatory roles, only a few lncRNAs are highly conserved across

vertebrates [30]. lncRNAs generally exhibit either poor conservation at the nucleotide level or

conservation in a short region only, particularly compared to protein-coding genes [31-33].

Although sequence conservation is often likely to indicate related function, sometimes it is

difficult to detect conservation across multiple genome sequences because of technical

challenges. lncRNAs, however, appear to be syntenically conserved with protein-coding

genes, which suggests that lncRNAs could have evolutionarily conserved roles in similar

genomic contexts [34-36]. A zebrafish lncRNA, linc-*oip5*, which has a short region of

sequence conservation with mammalian orthologs in the last exon, also exhibits preserved

genomic architecture in its size and arrangement of exons; furthermore, linc-*oip5* loss of

function disrupts zebrafish embryonic development, which can be rescued by the

mammalian orthologs [37]. Thus, examining the genomic context and/or short regions of

conservation in a lncRNA may be necessary for understanding lncRNA function.

lncRNA expression signatures also provide hints about lncRNA functional roles at

the cellular level. Global lncRNA profiling demonstrated that lncRNAs generally exhibit lower

expression than protein-coding genes [31, 38, 39] but tend to be uniquely or specifically

expressed in distinct tissues, developmental stages, conditions, or disease states [30-32, 38,

40-42]. For instance, one lncRNA, SAMMSON, is specifically expressed in melanoma cells

during melanogenesis and is known to regulate the process at the epigenetic level [43]. In

addition, large-scale analyses of lncRNA and protein-coding gene co-expression led to the

finding that a considerable number of paired genes are actually co-regulated by common

TFs [44, 45]. Often common TF binding motifs have been discovered in the promoters of the

co-expressed lncRNA and protein-coding genes, suggesting that the co-regulated genes

could share functional roles [46, 47]. Thus, to predict lncRNA biological functions, co-

expression networks of lncRNAs and protein-coding genes from large scale transcriptomic

data have been constructed and used for the inference of function [48-50].

Although genome and transcriptome maps of livestock animals, such as rainbow

trout, cow, goat, and chicken [51-55], have been recently constructed, only a few non-coding

transcriptome studies have been done in those genomes. To date, 9,681 lncRNAs have

been annotated in the red jungle fowl *Gallus gallus* genome, but these studies have been

limited to a few tissues and many lncRNAs seem to be missing. Thus, a comprehensive non-

coding transcriptome map of Ogye will help us understand phenotypic similarities and

differences between Ogye and *Gallus gallus.*

Data description

Total RNA samples and Bisulphite-treated DNA samples were collected from twenty

different tissues (Breast, Liver, Bone marrow, Fascia, Cerebrum, Gizzard, Immature egg,

Comb, Spleen, Mature egg, Cerebellum, Gall bladder, Kidney, Heart, Uterus, Pancreas,

Lung, Skin, Eye, and Shank) from 8-month-old Ogye (Fig. 1a). About 1.5 billion RNA-seq

reads (843 million single-end reads and 638 million paired-end reads) and 123 million

reduced representation bisulfite sequencing (RRBS) reads were analyzed (Table 1-2). The

RNA-seq and RRBS data from this study have been submitted to NCBI Gene Expression

Omnibus under SuperSeries accession number GSE104358.

Analyses

A comprehensive Ogye lncRNA catalogue

To construct an Ogye transcriptome map, pooled single- and paired-end RNA-seq reads of

each tissue were mapped to the Ogye draft genome (Genome;

https://www.ncbi.nlm.nih.gov/genome/; PRJNA412424) using STAR (ver 2.4.2) [56], and

subjected to transcriptome assembly using Cufflinks (ver 2.1.1) [57], leading to the

construction of transcriptome maps for twenty tissues. The resulting maps were combined

using Cuffmerge (ver 1.0.0) and, in total, 206,084 transcripts from 103,405 loci were

reconstructed in the Ogye genome (see the "Annotation of protein-coding and lncRNA genes"

section in the Methods for more details). In the unified Ogye transcriptome map, in addition

to 15,766 protein-coding genes, 1290 known (3266 transcripts) and 5610 novel (6263

transcripts) lncRNA genes were confidently annotated using our lncRNA annotation pipeline,

adopted from our previous study [58] (Additional file 1: Figure S1a). Compared to previously

annotated chicken lncRNAs from *Gallus gallus*, only 34% were redetected in the Ogye

lncRNA catalogue. Because approximately 92% previously annotated chicken protein-coding

genes were redetected in our Ogye protein-coding gene annotations (data not shown here),

the remainder that are missing may not result from genome mis-assembly. In fact, the

remainder were mainly either fragments of protein-coding genes in which exon junctions

were missed during transcriptome assembly or not expressed in all twenty Ogye tissues

(Additional file 1: Figure S1b). Only 276 previously annotated lncRNAs were missed in our

Ogye lncRNA annotations. Consistent with other species [39, 51, 52, 55], the median gene

length and the median exon number of Ogye lncRNAs were less than those of protein-

coding genes (Additional file 1: Figure S1c-d).

To profile the expression of protein-coding and lncRNA genes across tissues,

fragments per kilobase of exons per million mapped reads (FPKM) were measured for

transcripts using RSEM (v1.2.25) [59]. 6,565 lncRNA genes were expressed with FPKM $\geq$ 1

in at least one tissue, whereas 13,765 protein-coding genes were. As previously reported [32,

60, 61], Ogye lncRNAs generally displayed a tissue-specific expression pattern and some

lncRNAs were solely expressed in a single tissue, although a few displayed ubiquitous

expression across tissues (Additional file 1: Figure S1e). Tissue-specific genes with a four-

fold higher maximum expression value than the mean value over twenty tissues were

depicted on the genome using a Circos plot (Fig. 1c, green track). About 75% of lncRNA

genes (5191 loci) were tissue-specific, a significantly higher proportion than that of protein-

coding genes (45%; Fig. 1d; $P < 2.2e^{-16}$; Wilcoxon rank sum test). The fractions of lncRNAs

that were tissue-specific ranged from 2.4 % (Fascia) to 12.5% (Kidney), much higher

percentages than those of protein-coding genes, which ranged from 0.4% (Fascia) to 4.2%

(Kidney) (Fig. 1e). Hierarchical clustering of commonly expressed lncRNA genes among

tissues using the PHYLIP package (ver 3.6) [62] (see the "Hierarchical clustering of

expressed lncRNAs across tissues" section in the Methods for more details) defined

functionally and histologically-related tissue clusters well. In particular, 2,317 lncRNAs were

specifically expressed in the comb, skin, and shank, which are black tissues in Ogye (Fig.

1e). Only 780 lncRNAs were ubiquitously expressed across all tissues (Fig. 1e).

Tissue-specific DNA methylation landscape of the Ogye genome

To correlate tissue-specific lncRNA expression with its epigenetic status in the respective

tissue, reduced representation bisulfite sequencing (RRBS) reads were mapped to the Ogye

draft genome (Fig. 1b; Table2). The DNA methylation signals (C to T changes in CpGs)

across chromosomes were, then, calculated using Bismark in each sample (version 0.7.0)

[63] (see the "DNA methylation profiling" section in the Methods for more details). A

significant correlation (nominal $P \leq 0.05$) between the expression levels and the methylation

signals in the region 2kb upstream of genes across twenty tissues was demonstrated along

with a variation of the signals (Fig. 1c). The variability was measured as the relative

standard deviation. Of lncRNAs and protein-coding genes with tissue-specific differentially

methylated CpG sites (tDMC) that include ≥ five reads with C to T changes in the promoter

region in ≥ 10 tissues, 6.4% of the lncRNAs and 9.3% of the protein-coding genes displayed

a significant negative correlation (nominal $P \leq 0.05$) between their promoter methylation

levels and   their expression levels, percentages that were significantly higher than those of

random-pair controls (Additional file 1: Figure S1f; $P = 1.30 \times 10^{-6}$ for lncRNAs; $P = 7.93 \times$

10^{-36} for protein-coding genes; Fisher's exact test). However, only about 3% of genes

showed a positive correlation between their expression and methylation signals, which is

comparable or less than the control (Additional file 1: Figure S1g; $P = 0.87$ for lncRNAs; $P =$

0.013 for protein-coding genes). Collectively, these results show that CpG methylation in the

promoters represses the expression of target genes.

Co-expression analyses of lncRNAs specify tissue-specific functional clusters

As lncRNAs tend to be specifically expressed in a tissue or in related tissues, they could be

better factors for defining genomic characteristics of tissues than protein-coding genes. To

prove this idea, principle component analyses (PCA) were performed with tissue-specific

lncRNAs and protein-coding genes (Additional file 1: Figure S2). As expected, the 1st, 2nd,

and 3rd PCs of lncRNAs enabled us to predict the majority of variances, and better discerned

distantly-related tissues and functionally and histologically-related tissues (i.e., black tissues

and brain tissues) (Additional file 1: Figure S2a) than those of protein-coding genes

(Additional file 1: Figure S2b).

To identify functional clusters of lncRNAs, pairwise correlation coefficients between

tissue-specific lncRNAs were calculated and the co-expression patterns across 20 tissues

were clustered, defining 16 co-expression clusters (Fig. 2). As expected, each co-expression

cluster was defined as a functional group, highly expressed in a certain tissue (kidney, eye,

pancreas, uterus, mature egg, immature egg, breast, heart, liver, lung, gall bladder, gizzard,

bone marrow, or spleen) or related tissues (brain and black tissues) (Additional file 4: Table

S3). In particular, the largest co-expression cluster, the brain-specific group, included 930 co-

expressed lncRNAs, highly expressed in cerebrum and cerebellum. The second largest

cluster, the black tissue-specific group, included 479 co-expressed lncRNAs, highly

expressed in fascia, comb, skin, and shank (Fig. 2). Clusters of related tissues also display

distinct sub-modules corresponding to each tissue. For instance, lncRNA clusters specific to

black tissues displayed sub-clusters including sub-cluster 1 specific to shank and sub-cluster

2 specific to comb, although the sub-clusters shared skin-specific expression (Additional file

1: Figure S 2c).

The functional role of each co-expressed lncRNA cluster can be indirectly inferred

by a set of co-expressed mRNAs [48-50]. Thus, mRNAs that are exclusively co-expressed

with each lncRNA cluster were identified with the following criteria: a mean Pearson's

correlation ($\bar{r}$) ≥ 0.5 with members within a cluster and the differences between the

corresponding $\bar{r}$ and the mean correlation ($\bar{r}_i$) with all other groups ≥ 0.3, and were

subsequently subjected to gene ontology (GO) analyses using DAVID [64] (Fig. 2; Additional

file 5:Table S4). In particular, 1617 mRNAs exclusively correlated to the brain-specific

lncRNA group (930 lncRNAs) were identified and had been associated with brain-function

specific terms, such as neuroactive ligand-receptor interaction (q = 2.18 X 10^{-12}; False

discovery rate, FDR correction). In contrast, 748 mRNAs exclusively correlated to spleen-

specific lncRNAs were identified and had been associated with immune-related terms, such

as leukocyte activation (q = 2.37X 10^{-12}). Likewise, 10 out of 16 co-expression clusters of

lncRNAs had functional evidence, with significantly enriched GO terms and KEGG pathways

(Fig. 2).

lncRNAs as epigenetic activators

The coherent expression of two different RNA classes could be in part the outcome of either

active regulation by lncRNAs in *cis* and *trans*, or co-regulation by common regulators, such

as TFs or epigenetic regulators, in *cis* and *trans* (Additional file 1: Figure S3). Regulation of

gene expression by lncRNAs often involves engagement with chromatin remodelers, such

as polycomb repressive complexes (PRCs) that mediate the suppression of target mRNA

expression [65, 66] or demethylases that open the chromatin structure to enhance the

expression of target mRNAs [67, 68] (Additional file 1: Figure S3a). Remote co-expression of

lncRNAs and mRNAs can be also regulated by common TFs [44, 45] (Additional file 1:

Figure S3b). Co-expressed genes tend to have common TF binding motifs in their promoters.

However, *cis*-regulation of mRNA expression by lncRNAs is known to be associated with

common epigenetic factors (Additional file 1: Figure S3c) or enhancers (Additional file 1:

Figure S3d).

To find lncRNAs that act as epigenetic activators that reduce methylation levels,

lncRNAs with expression levels that are significantly negatively correlated with the

methylation level in the promoters of co-expressed protein-coding genes (nominal $P \leq 0.01$)

were examined in each co-expression cluster. In this case, the lncRNAs are thought to

reduce the methylation level in the promoters of the co-expressed protein-coding genes. Of

the lncRNAs in clusters, the expression of 15.0%~72.9% displayed significantly negative

correlation with methylation levels in the promoters of co-expressed protein-coding genes,

which were compared to those of random protein-coding gene cohorts (Fig. 3a). Clusters

specific to brain, kidney, mature egg, breast, heart and spleen included significantly more

lncRNAs with a significant correlation than did the random controls ($P = 0.026 \sim 7.71 \times 10^{-13}$)

but this was not true for the black tissue cluster. To identify DNA methylation activators with

more confidence, we also examined whether the expression and methylation of the co-

expressed coding genes were correlated (nominal $P \leq 0.01$). 820 lncRNAs in the clusters

were identified as confident DNA methylation activator candidates (Fig. 3b). Genes encoding

lncRNAs that act as DNA methylation regulators of protein-coding genes were mostly 100kb

apart, and only five were within 100kb from target genes, suggesting that lncRNAs that

function as epigenetic activators mostly play their roles in *trans*-form rather than *cis*-form.

Transcriptional regulation by common TFs

To identify co-expressed pairs of lncRNAs and mRNAs regulated by common TFs, TF

binding sites (TFBSs) enriched in the promoters of the co-expressed genes were examined.

For this analysis, sequences 2kb upstream of the co-expressed genes were extracted and

enriched sequence motifs were identified using the multiple expectation-maximization for

motif elicitation (MEME) suite [69] (see the "Prediction of TFBSs" section in the Methods for

more details). The resulting motifs were subjected to analysis by the TOMTOM program [70]

to annotate TFBSs based on TRANSFAC database v3.2 [71]. As a result, 14 common TFs

that have significantly abundant binding sites in the promoters of lncRNA and protein-coding

genes were detected (Additional file 1: Figure S4; corresponding to model 2). To discern TFs

available in chicken genomes, PANTHER [72, 73] was used to examine whether there are

chicken orthologs of the TFs and whether the orthologs are expressed in the corresponding

tissues (FPKM $\geq$ 1). Finally, five TFs, including HSF2 and SP1, were identified as candidates

(Fig. 4a). HSF2 and SP1 binding sites were more recurrently detected across tissues than

others and were significantly enriched in the promoters of 478 lncRNAs and 634 protein-

coding genes. Although the binding motifs were slightly degenerated from the annotated

motifs, the HSF2 motifs were similar in the promoters of lncRNA genes and protein-coding

genes (Figs. 4b).

To examine further whether the respective TFs actually affect the expression of

lncRNAs and protein-coding genes, the correlation between the expression of each TF and

co-expressed genes in each cluster was examined. Interestingly, HSF2 expression had a

strong positive correlation with expression of genes in black tissues but not in other tissues

(Fig. 4c). The expression pattern for each of the five lncRNAs and protein-coding genes that

were highly correlated with that of HSF2 was specific for skin, shank, and comb compared to

other tissues (Fig. 4d). Thus, HSF2 is a promising candidate for regulating the black tissue-

specific expression of lncRNAs and protein coding genes. Taken together, our data indicate

that of a total of 3466 lncRNA in ten clusters, 615 (17.74%) appear to be co-regulated with

co-expressed protein-coding genes by common TFs, such as HSF2.

Coherent expression of neighboring lncRNA and protein-coding genes

Previous studies showed that lncRNAs and their neighboring protein-coding genes are

highly correlated in their expression across tissues and developmental stages [35, 38]. To

examine how the co-expressed lncRNAs and mRNAs in our study are co-localized in

chromosomes, lncRNAs from each group were first classified based on the closest distances

(≤10kb, ≤100kb, >100kb, and other chromosomes) from the significantly co-expressed

protein-coding genes (nominal $P ≤ 0.01$; Pearson's correlation) (Fig. 5a). Genes encoding

co-expressed pairs of lncRNAs and mRNAs are significantly proximally co-localized within

10kb (Fig. 5a left; $P ≤ 0.05$, Fisher's exact test), compared to random controls (Fig. 5a right)

but not those of lncRNAs and mRNAs in the range of 10~100kb or in the 100kb outside.

Overall, 2 ~ 15 % of the co-expressed pairs in the clusters tended to be proximally co-

regulated within 10kb.

To examine how neighboring lncRNAs and protein-coding genes are tissue-

specifically co-regulated, the pairs within 10kb were classified into three categories on the

basis of their relative orientations (head-to-tail, tail-to-tail, or head-to-head). The correlation

coefficients of the pairs in each category were compared to those of lncRNA and random

protein-coding gene controls from tissue-specific gene sets (Fig. 5b) or from ubiquitously

expressed gene sets (Additional file 1: Figure S5a). Both neighboring lncRNA and protein-

coding gene pairs displayed significantly greater correlation than did random controls,

regardless of the category, in both sets (Fig. 5b; Additional file 1: Figure S5a). The

correlations were also compared to those of neighboring protein-coding gene pairs. Whereas

the correlations of the ubiquitously expressed, neighboring lncRNAs and protein-coding

genes were significantly lower than those of ubiquitously expressed neighboring protein-

coding gene pairs in the head-to-tail and head-to-head categories (Additional file 1: Figure

S5a), the correlation coefficients of the tissue-specific pairs were slightly yet insignificantly

higher than those of neighboring protein-coding gene pairs (Fig. 5b).

To dissect factors that affect the co-regulation of tissue-specific neighboring lncRNA

and protein-coding gene pairs, the pairs with a high correlation ($P \leq 0.05$) between the

methylation levels of their promoters (methylation-related group – model 3) and those with

no correlation (methylation-unrelated group) were divided. Tissue-specific neighboring

lncRNA and protein-coding gene pairs showed no more expression correlation than did

neighboring protein-coding genes in the methylation-related group (Fig. 5c; $P = 0.71$,

Wilcoxon rank sum test), whereas they did show a significantly higher correlation in the

methylation-unrelated group (Fig. 5d; $P \leq 0.001$ for head-to-tail, $P \leq 0.05$ for head-to-head,

Wilcoxon rank sum test), which suggests that neighboring lncRNAs and protein-coding

genes in the methylation-unrelated group have a regulatory interaction between them.

Enhancer-associated RNA-mediated gene regulation

Previous studies showed that lncRNAs associated with enhancers could regulate their

neighboring protein-coding genes [74]. Genomic association between lncRNAs and

enhancers, detected in embryonic developmental stages in the chicken [75], revealed that

lncRNAs in the methylation-unrelated group are more significantly associated with

enhancers than those in the other group (Fig. 5e; P = 2.72 X 10^{-6}; Fisher's exact test). As a

result, 136 head-to-tail lncRNAs, 67 tail-to-tail lncRNAs and 124 head-to-head lncRNAs were

considered as enhancer-associated lncRNA candidates (eRNAs; Supplementary Table S5).

The eRNAs (corresponding to model 4) had a greater correlation with neighboring protein-

coding genes only in the head-to-tail group (Fig. 5f), whereas non-eRNAs displayed a

greater correlation in the head-to-head orientation, which could allow sharing of common

promoters (Fig. 5g). A few eRNAs were discovered to have strong bi-directional

transcriptional activity (Additional file 1: Figure S5b; see the "Transcriptional activity of

eRNAs" section in the Methods for more details), as previously reported [76, 77]

Next, to identify TFs binding to genomic regions that transcribe eRNAs, TF binding

sites detected from all the genomic regions associated with enhancers were profiled and

were compared to those of TFs detected from the enhancers specific to a certain tissue (Fig.

5h). Oct1 and HSF2 binding sites were significantly localized in eRNAs specific to black

tissues (P < 3.09 X 10^{-5} for Oct1; P < 3.11 X 10^{-7} for HSF2; binomial test). Besides the TFs

specific to black tissues, GR, YY1, RAP1 and GATA1, and HSF3 binding sites were localized

in eRNAs specific to heart, eye, spleen and bone marrow, respectively (Fig. 5h). Interestingly,

HSF2 was a common TF candidate for co-regulating lncRNAs and protein-coding genes at a

distance (Fig. 5d).

Conserved black skin-specific lncRNAs

As mentioned earlier, unlike other chicken breeds, both the plumage and skin of the Ogye

are black. To identify lncRNAs potentially functionally related to this trait, lncRNAs

specifically co-expressed in black tissues (Fig. 2) were further investigated by comparing to

those in non-black skin of other chicken breeds. Of 479 lncRNAs specific to black tissues, 47

were significantly two-fold up- (29) or down-regulated (18) in Ogye black skin, compared to

those in brown leghorn skin (Fig. 6a; Additional file 7:Table S6; FDR < 0.05).

To find functionally conserved lncRNAs, the 47 differentially expressed lncRNAs

were examined for synteny and sequence conservation in human and mouse genomes.

Synteny conservation considers whether orthologs of a certain lncRNA's neighboring genes

are positionally conserved in these mammalian genomes (Fig. 6b). As a result of this

analysis, about 10% of lncRNAs were found to be syntenically conserved in both the human

and mouse genomes and about 25% were syntenically conserved in at least one genome

(Fig. 6c; Additional file 8:Table S7), percentages that are comparable to those of the protein-

coding genes (Fig. 6d). However, sequence similarity analyses by the BLAST showed that

only 6% of the syntenically conserved lncRNAs had conserved sequences relative to

sequences in either the human or mouse genomes (Fig. 6c; Additional file 8:Table S7), which

is much lower than that of protein-coding genes (56%). Taken together, our data showed that

16 lncRNAs were syntenically or sequentially conserved and differentially expressed in black

tissue (Fig. 6e).

Of the 16 lncRNAs that have evidence of black tissue-specific function, four,

including eRNAs, were associated with HSF2 binding motifs, whereas of the 104 that have

synteny and sequence conservation but are not differentially expressed in black tissues, only

one was associated with HSF2. The presence of HSF2 binding motifs appears to be

significantly related to black tissue-specific expression (Fig 6f; $P \leq 0.0008$, Fisher's exact

test). For instance, linc-THEM184c is significantly up-regulated in black tissue (Fig. 7b), its

locus is syntenically conserved with neighboring genes, TMEM184C and EDNRA, in both

human and mouse genomes, and its promoter includes a HSF2 binding motif (Fig. 7a). In

addition, the protein-coding genes that are co-expressed with this lncRNA are enriched for

GO terms that are functionally relevant for black skin: keratinocyte differentiation,

angiogenesis, and ECM-receptor-interaction (Fig. 7c). Among the co-expressed genes, 31

have HSF2 binding sites in their promoters (Fig. 7a). As another example, black-tissue

specific linc-FAM204A is syntenically conserved with the RAB11FIP2 and FAM204A genes in

the human and mouse genomes (Additional file 1: Figure S6a). This lncRNA was highly

expressed in black tissues including the skin, shank, and comb but had no expression in

other tissues except for the eye (Additional file 1: Figure S6b). The co-expressed protein-

coding genes are enriched for functionally relevant GO terms melanogenesis, ECM-receptor

interaction, and Wnt signaling (Additional file 1: Figure S6c). Interestingly, the human and

Ogye lncRNA orthologs share a conserved sequence of 389 nt, which includes multiple

miRNA 7-mer target sites (Additional file 1: Figure S6a).

Discussion

In this study, 6900 multiple-exon lncRNAs were identified from twenty tissues of Ogye; about

18% had been previously annotated in *Galllus gallus red junglefowl*. The remainder of the

previously annotated lncRNAs were mostly not expressed in Ogye or were false annotations,

suggesting that the current chicken lncRNA annotations should be reconstructed more

carefully. Our Ogye lncRNAs resembled previously annotated lncRNAs in mammals in their

genomic characteristics, including transcript length, exon number, and tissue-specific

expression pattern, providing evidence for the accuracy of the new annotations. Hence, the

Ogye lncRNA catalogue may help us to improve lncRNA annotations in the chicken

reference genome.

The majority of lncRNAs showed a tissue-specific expression pattern, defining

functionally coherent co-expression clusters. The tissue-specific expression and the

coherent expression of lncRNA genes with other protein-coding genes could be attributed to

common epigenetic and transcriptional regulation. In fact, of the lncRNAs in clusters, 39.3%

had evidence associating them with at least one model (Additional file 1: Figure S7a); most

commonly, these involved lncRNAs that act as epigenetic activators of protein-coding gene

expression and common TFs that bind to the lncRNA and protein-coding gene promoters

(Additional file 1: Figure S7b). Interestingly, 126 lncRNAs had evidence supporting both the

epigenetic activator and TF models (Additional file 1: Figure S7c). 79 lncRNAs had functional

evidence supporting their identity as eRNAs. Although lncRNAs are known to be mostly

involved in epigenetic repression of genes, our study intentionally focused on lncRNAs as

epigenetic activators by correlating the level of lncRNAs and the methylation in target gene

promoters. Furthermore, because only a subset of CpG sites are sometimes related to the

chromatin state and transcriptional activity of target genes, averaging CpG methylation

signals in the promoter might underestimate the fraction of epigenetically activating lncRNAs

in our study.

lncRNA and protein-coding genes co-expressed in black tissues had HSF2 binding

sites in their promoters and were specifically correlated with the level of HSF2 across tissues,

supporting that the genes are co-regulated by HSF2. Moreover, enhancers that included

HSF2 binding sites were associated with eRNAs specific to black tissue, indicating that

HSF2 is the most likely regulator of black tissue-specific expression. Because the ancestor

of Ogye appears to have originated in the rainforest, it makes sense that heat shock-related

factors could be involved in melanogenesis and hyper-pigmentation processes, which would

help avoid the absorption of too much heat. One of the black skin-specific lncRNAs, lnc-

THMEM184c, is most abundantly expressed in comb, and HSF2 appears to co-regulate lnc-

THMEM184c and its co-expressed protein-coding genes, which are related to keratinocyte

differentiation and ECM-receptor interaction (Fig. 7).

In addition, several previous studies that also focused on animal coat color showed that the

color can be determined by the amount and type of melanin produced and released by

melanocytes present in the skin [78, 79]. Melanin is produced by melanosomes, large

organelles in melanocytes, in a process called melanogenesis. Wnt signaling has a

regulatory role in the melanogenesis pathway and is also required for the developmental

process that leads to melanocyte differentiation from neural crest cells [80, 81]. One of the

candidate lncRNAs related to the process is linc-FAM204A, whose co-expressed protein-

coding genes are associated with GO terms melanogenesis, ECM-receptor interaction, and

Wnt signaling pathway (Additional file 1: Figure S6c). linc-FAM204A, which contains a short-

conserved motif, is broadly preserved in mammalian genomes, including the human, rhesus

macaque, mouse, dog, and elephant genomes. Among these orthologs, the human ortholog

is known as CASC2, and is suppressed in lung, colorectal, renal and other cancers by miR-

21-5p targeting via the conserved 7-mer site (Additional file 1: Figure S6a).

Taken together, these results indicate that coding and non-coding RNAs functionally

relevant to black and other tissues could help explain unique genomic and functional

characteristics of a Korean domestic chicken breed, Yeonsan Ogye. Additionally, these

findings could provide unprecedented insight for future studies with industrial and agricultural

applications, as well as for scientific analysis of chicken genomes.

Methods

Acquisition and care of Yeonsan Ogye

Yeonsan Ogye chickens (object number: 02127), obtained from the Animal Genetic

Resource Research Center of the National Institute of Animal Science (Namwon, Korea),

were used in the study. The care and experimental use of Ogye was reviewed and approved

by the Institutional Animal Care and Use Committee of the National Institute of Animal

Science (IACUC No.: 2014-080). Ogye management, treatment, and sample collection and

further analysis of all raw data were performed at the National Institute of Animal Science.

Preparation of RNA-seq libraries

Total RNAs were extracted from twenty Ogye tissues using 80% EtOH and TRIzol. The RNA

concentration was checked by Quant-IT RiboGreen (Invitrogen, Carlsbad, USA). To assess

the integrity of the total RNA, samples were run on a TapeStation RNA screentape (Agilent,

Waldbronn, Germany). Only high quality RNA samples (RIN ≥ 7.0) were used for RNA-seq

library construction. Each library was independently prepared with 300ng of total RNA using

an Illumina TruSeq Stranded Total RNA Sample Prep Kit (Illumina, San Diego, CA, USA).

The rRNA in the total RNA was depleted using a Ribo-Zero kit. After rRNA depletion, the

remaining RNA was purified, fragmented and primed for cDNA synthesis. The cleaved RNA

fragments were copied into the first cDNA strand using reverse transcriptase and random

hexamers. This step was followed by second strand cDNA synthesis using DNA Polymerase

I, RNase H and dUTP. The resulting cDNA fragments then underwent an end repair process,

the addition of a single 'A' base, after which adapters were ligated. The products were

purified and enriched with PCR to create the final cDNA library. The libraries were quantified

using qPCR according to the qPCR Quantification Protocol Guide (KAPA Library

Quantificatoin kits for Illumina Sequecing platforms) and qualified using the TapeStation

D1000 ScreenTape assay (Agilent Technologies, Waldbronn, Germany).

Preparation of RRBS libraries

Preparation of reduced representation bisulfite sequencing (RRBS) libraries was done

following Illumina's RRBS protocol. 5ug of genomic DNA that had been digested with the

restriction enzyme MspI and purified with a QIAquick PCR purification kit (QIAGEN, Hilden,

Germany) was used for library preparation, which was done using a TruSeq Nano DNA

Library Prep Kit (Illumina, San Diego, USA). Eluted DNA fragments were end-repaired,

extended on the 3′ end with an 'A', and ligated with Truseq adapters. After ligation had been

assessed, the products, which ranged from 175 to 225bp in length (insert DNA of 55–105 bp

plus adaptors of 120 bp), were excised from a 2%(w/v) Low Range Ultra Agarose gel (Biorad,

Hercules, USA) and purified using the QIAquick gel extraction protocol. The purified DNA

underwent bisulfite conversion using an EpiTect Bisulfite Kit (Qiagen, 59104). The bisulfite-

converted DNA libraries were amplified by PCR (four cycles) using PfuTurbo Cx DNA

polymerase (Agilent, 600410). The final product was then quantified using qPCR and

qualified using the Agilent Technologies 2200 TapeStation assay (Agilent, Waldbronn,

Germany). The final product was sequenced using the HiSeq™ 2500 platform (Illumina, San

Diego, USA).

Annotations of protein-coding and lncRNA genes

To annotate protein-coding genes in the Ogye genome, *Gallus gallus* (red junglefowl)

protein-coding   genes   downloaded   from   Ensembl   biomart   (release   81;

http://www.ensembl.org/biomart) were mapped onto the Ogye draft genome v1.0 using

GMAP (v2015-07-23)[82]. Genes that had greater than 90% coverage and identity were

selected as Ogye protein-coding genes. As a result, 14,264 protein-coding genes were

subjected to further analysis.

For lncRNA gene annotations, RNA-seq data from twenty different tissues (Breast, Liver,

Bone marrow, Fascia, Cerebrum, Gizzard, Immature egg, Comb, Spleen, Mature egg,

Cerebellum, Gall bladder, Kidney, Heart, Uterus, Pancreas, Lung, Skin, Eye, and Shank)

were produced in both single end and paired-end types. Sequences were preprocessed to

filter nucleotides with low quality scores using FASTQC (v 0.10.1) [83] and were mapped to

the Ogye draft genome using STAR (v2.4.2)[56] with the options '--runMode alignReads --

alignIntronMin 67 --alignIntronMax 36873 –outReadsUnmapped Fastx –outFilterMismatch_

Nmax 999 --outFilterMismatchNoverLmax 0.02 -alignMatesGapMax 1000000 –outSAMtyp_

e BAM SortedByCoordinate --outFilterMultimapNmax 5 --outWigType wiggle –outWigStran_

d Stranded --outWigNorm RPM'. Initial transcriptome assemblies from twenty tissues were

performed with Cufflinks (ver 2.1.1)[57] with the parameter '--library-type fr-firststrand' and

the resulting assemblies were combined using Cuffmerge (ver 1.0.0)[57] with the default

option. In total, 206,084 transcripts from 103,405 loci were annotated in the Ogye genome.

To distinguish lncRNAs from other biotypes of RNAs, such as mRNAs, tRNAs, rRNAs,

snoRNAs, miRNAs, and other small non-coding RNAs, these sequences were downloaded

from ENSEMBL biomart and aligned with the resulting transcript sequences. Any transcripts

overlapping at least 1nt with known RNAs were excluded. Of the remainder, those of less

than 200nt in length and within 200 bp of protein-coding genes on the same strand were

further excluded to avoid the inclusion of fragmented RNAs. 54,760 lncRNA candidate loci

(60257 transcripts) were retained and compared with a chicken lncRNA annotation of

NONCODE (v2016) [84]. Of the candidates, 2094 loci (5215 transcripts) overlapped with

previously annotated chicken lncRNAs. 52,666 non-overlapping loci (55,042 transcripts)

were further examined to determine whether they had coding potential using coding potential

calculator (CPC) scores [85]. Those with a score greater than -1 were filtered out, and the

remainder (14,108 novel lncRNA candidate loci without coding potential) were subjected to

the next step. Because many candidates still appeared to be fragmented, those with a single

exon but with neighboring candidates within 36,873bp, which is the intron length of the 99[th]

percentile, were re-examined using both exon-junction reads consistently presented over

twenty tissues and the maximum entropy score [86], as done in our previous study [58]. If

there were at least two junction reads spanning two neighboring transcripts or if the entropy

score was greater than 4.66 in the interspace, two candidates were reconnected, and those

with a single exon were discarded. In the final version, 9529 transcripts from 6900 lncRNA

loci (5610 novel and 1290 known) were annotated as lncRNAs.

DNA methylation profiling

RRBS reads with a low quality score (Phred quality score < 20) were discarded using

FastQC (v0.10.1). The remaining reads were aligned to the Ogye draft genome (v1.0) using

Bismark [63]. The methylation level of each cytosine in a CpG region was calculated using

Bismark methylation extractor. Tissue-specific, differentially methylated CpG sites (tDMC),

covered with at least five reads in a promoter, were considered for downstream analysis. A

tissue specific site is defined as one in which its mean methylation across tissues is at least

four time greater than the signal in a certain tissue. A promoter region is defined as the

region 2 kb upstream of the 5' end of genes.

Expression profiling

The expression values (FPKM) of lncRNA and protein-coding genes were estimated using

RSEM (v1.2.25) in each tissue. The values across tissues were normalized using the

quantile normalization method. In all downstream analyses, lncRNA or protein-coding genes

with FPKM ≥ 1 in at least one tissue were used. lncRNAs for which the maximum expression

value across twenty tissues was at least four-fold higher than the mean value were

considered to exhibit tissue-specific expression. In total, 5,191 (75%) lncRNAs were

considered to be tissue-specific across twenty different tissues.

Hierarchical clustering of expressed lncRNAs across tissues

To perform hierarchical clustering of commonly expressed lncRNA genes among tissues, the

list of expressed lncRNAs in each tissue was used as a input vector for phylogenetic

clustering. The clustering was done using the PHYLIP package. lncRNAs with FPKM ≥ 1 in a

certain tissue were considered to be expressed in a certain tissue. As two tissues share

more common genes, they become more closely clustered.

Clustering of co-expressed lncRNAs

Hierarchical clustering was performed to search for expression clusters of lncRNAs across

twenty tissues using Pearson's correlation coefficient metrics. Clusters in which more than

80% of their members are most highly expressed in the same or related tissues (brain and

black tissues) were regarded as tissue-specific. Sub-clusters in the brain and black tissue

clusters were further defined with the same criterion mentioned above.

Defining coding genes co-expressed with lncRNAs in a cluster

Protein-coding genes with a high mean correlation with lncRNAs in a cluster (Pearson's

correlation $\geq$ 0.5), but for which the mean correlation to the cluster is at least 0.3 greater

than those of other clusters, were assigned to the co-expressed set of the cluster. Each set

of mRNAs was used to perform gene ontology (GO) term and pathway enrichment analyses

using DAVID [64]. Terms were only selected when the false discovery rate (FDR) q value

was ≤ 0.05.

Correlation of the methylation level of neighboring lncRNA and protein-coding genes.

The methylation levels at CpG sites in the promoters of neighboring lncRNA and protein-

coding genes were correlated with each other over twenty tissues (using Pearson's

correlation coefficients). Only tissues in which a certain position had sufficient read coverage

(at least five) were considered for measuring the correlation. If the nominal P value was ≤

0.05, then the pair of lncRNA and protein-coding genes was considered as having a

significantly correlated interaction.

Correlating the expression level of lncRNAs with the methylation level of protein-

coding genes

To identify lncRNAs as potential epigenetic activators, the expression of lncRNAs and the

methylation at CpG sites in the promoters of protein-coding genes were correlated over

twenty tissues using a non-parametric correlation method (Spearman's correlation). Only

pairs of lncRNA and protein-coding genes exhibiting a nominal P value ≤ 0.01 were

considered as having a significantly correlated interaction. Of the resulting pairs, if the

protein-coding mRNAs had a significant correlation (nominal P value ≤ 0.01) between their

expression level and the methylation level in their promoter, its paired lncRNA was regarded

as an epigenetic activator.

Prediction of TFBSs

To identify enriched TFBSs in the promoters of the co-expressed lncRNAs in each tissue-

specific cluster and in the promoters of the co-expressed protein-coding genes within the

cluster, the promoter sequences were examined using the MEME suite (V4.9.0). Motifs that

exhibit an E-value ≤ 1 X 10^{-5} were selected as enriched motifs, associated with the

corresponding tissue. The resulting motifs were searched for in the Transfac database [71]

using TomTom [70]. As a result, 14 TFBSs significantly enriched in a certain tissue or in a set

of similar tissues were detected (P ≤ 0.01), of which 6 had associated TF orthologs (SP1,

HEN1, HSF2, HB, AP-2, Oct1) encoded in the genome. However, HEN1 was not expressed

in a corresponding tissue (FPKM ≤ 1). In addition, to confirm TFs related to enhancers,

enhancer sequences were compared with the resulting TFBSs.

Identification of enhancer regions

To annotate enhancer regions in the Ogye draft genome, annotation files including all

enhancers in the *Gallus gallus* (*red junglefowl*) genome were downloaded from the NCBI

gene expression omnibus (GEO, GSE75480). Enhancer sequences extracted using our in-

house script were aligned to the Ogye draft genome using BLAST (-p blastn). Regions

that significantly matched the original enhancers (E-value ≤ 1 X 10^{-5}) and with high

coverage of more than 80% were annotated as Ogye enhancers.

Transcriptional activity of eRNAs

To examine bi-directional transcriptional activity of eRNAs, total mapped reads in the range

spanning 1kb upstream to 1 kb downstream of the eRNA transcription start site (TSS) were

re-examined on both forward and reverse strands.

Correlation of expression between neighboring lncRNA and protein-coding genes

Pairs consisting of a lncRNA and its closest neighboring protein-coding gene within 10kb

were classified into three groups based on their genomic orientations: head-to-head (can be

divergently overlapped), head-to-tail (including only independent lncRNAs with evidence of a

TSS and cleavage and polyadenylation site; otherwise, these lncRNAs must be at least 1kb

apart from each other), and tail-to-tail (can be convergently overlapped). The correlation of

the expression of these pairs was calculated over twenty tissues using Pearson's correlation

method. The average correlation coefficient values and their standard errors were calculated

in the respective groups. As a random control, the expression of 1000 random pairs of

lncRNA and protein-coding genes were correlated using the same method. As another

control, number-matched pairs of neighboring protein-coding genes were also correlated

with each other.

Synteny and sequence conservation

To examine the conservation of synteny of a lncRNA, its closest downstream and upstream

neighboring protein-coding genes in the Ogye genome were matched to their orthologous

genes in the mouse and human genomes. If a lncRNA is located between the two

orthologous genes, regardless of direction, that lncRNA was regarded as syntenically

conserved. GENCODE lncRNA annotations (v25 for human and vM11 for mouse) were

analyzed for this study. To check for sequence conservation, Ogye lncRNA sequences were

aligned to lncRNA sequences from other species, intronic sequences, and their flanking

sequences (up to 4 Mb) using BLAST. For a significant match, an E-value 1 X 10^{-6} was

used as a cutoff.

Analysis of lncRNA differential expression

To identify lncRNAs that are differentially expressed between Ogye and Brown leghorn skin

tissues, Brown leghorn skin RNA-seq data were downloaded from the NCBI SRA

(ERR1298635, ERR1298636, ERR1298637, ERR1298638, ERR1298639, ERR1298640,

and ERR1298641). Reads were mapped to the *Gallus gallus* Galgal4 reference genome

using Bowtie (V1.0.0), and the average mismatch rates were estimated across read

positions. If the mismatch rate was greater than 0.1 at a certain position, sequences on high

mismatch side of the position were trimmed using seqtk (https://github.com/lh3/seqtk), and

then sickle was used with the default option for read quality control. Preprocessed reads

from RNA-seq data were mapped onto the chicken Galgal4 reference genome using STAR.

The read counts of lncRNAs were performed using HTSeq (v0.6.0) and the differential

expression analysis was performed using DESeq [87]. Genes with a greater than two-fold

difference in expression and a FDR q value ≤ 0.05 were considered to be differentially

expressed.

Tables

Table 1. Sequencing and mapping summary of RNA-seq reads

Samples	Paired-end sequencing reads			Single-end sequencing reads		
	Sequencing reads	Mapping rate	SRA accession	Sequencing reads	Mapping rate	SRA accession

Tissue	Reads	%	Accession	Reads	%	Accession
Breast	34,893,064	92.05%	SRX3223583	43,294,022	90.70%	SRX3223603
Liver	33,476,266	85.75%	SRX3223584	48,032,813	85.81%	SRX3223604
Bone marrow	30,975,506	85.00%	SRX3223585	40,286,974	87.99%	SRX3223605
Fascia	33,316,764	84.61%	SRX3223586	42,425,452	87.93%	SRX3223606
Cerebrum	30,887,821	89.95%	SRX3223587	46,455,658	92.32%	SRX3223607
Gizzard	31,537,118	84.00%	SRX3223588	38,689,871	85.82%	SRX3223608
Immature egg	32,009,437	87.73%	SRX3223589	32,048,703	87.80%	SRX3223609
Comb	31,936,332	85.34%	SRX3223590	37,985,049	87.76%	SRX3223610
Spleen	28,946,777	89.70%	SRX3223591	38,704,448	89.33%	SRX3223611
Mature egg	30,873,699	91.98%	SRX3223592	40,650,664	92.17%	SRX3223612
Cerebellum	30,798,145	93.53%	SRX3223593	39,940,946	93.34%	SRX3223613
Gall bladder	35,862,229	84.83%	SRX3223594	35,423,339	87.06%	SRX3223614
Kidney	29,953,007	87.25%	SRX3223595	39,894,009	89.99%	SRX3223615
Heart	30,986,431	94.14%	SRX3223596	45,951,338	91.49%	SRX3223616
Uterus	33,444,002	91.89%	SRX3223597	46,650,355	90.63%	SRX3223617
Pancreas	30,595,568	82.52%	SRX3223598	47,361,192	84.35%	SRX3223618
Lung	31,533,498	87.63%	SRX3223599	45,552,982	92.34%	SRX3223619
Skin	34,442,464	82.36%	SRX3223600	41,934,970	84.00%	SRX3223620
Eye	33,006,509	89.21%	SRX3223601	44,044,630	91.82%	SRX3223621
Shank	28,643,334	94.07%	SRX3223602	47,716,995	79.86%	SRX3223622

Table 2. **Sequencing and mapping summary of RRBS reads**

Samples	Sequencing reads	Mapping rate	SRA accession
Breast	6,042,106	68.90%	SRX3223667
Liver	6,744,208	74.20%	SRX3223668
Bone marrow	5,736,011	72.00%	SRX3223669
Fascia	5,720,194	68.90%	SRX3223670
Cerebrum	6,078,989	70.00%	SRX3223671
Gizzard	5,731,878	69.40%	SRX3223672
Immature egg	6,741,258	67.70%	SRX3223673
Comb	5,948,687	72.90%	SRX3223674
Spleen	6,307,517	77.60%	SRX3223675
Mature egg	6,246,607	69.20%	SRX3223676
Cerebellum	6,291,610	68.20%	SRX3223677
Gall bladder	5,738,180	70.10%	SRX3223678
Kidney	5,470,502	68.60%	SRX3223679
Heart	5,462,739	69.40%	SRX3223680
Uterus	6,046,764	67.90%	SRX3223681
Pancreas	7,100,215	70.30%	SRX3223682
Lung	5,640,120	67.60%	SRX3223683
Skin	7,226,309	72.40%	SRX3223684
Eye	6,956,141	71.90%	SRX3223685
Shank	5,924,463	74.20%	SRX3223686

Availability of supporting data and materials

Raw RNA-seq and RRBS data from twenty different Ogye tissues have been submitted to

the NCBI Gene Expression Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/; GSE104351)

under SuperSeries accession number GSE104358. All lncRNA catalogs and expression

tables from this study have also been submitted to NCBI GEO (GSE104351) under the same

SuperSeries accession number, GSE104358.

Declarations

Consent for publication

Not applicable

Competing interests

Not applicable

Funding

This work was supported by the Program for Agriculture Science & Technology Development

(Project No. PJ01045301; PJ01045303) of the Rural Development Administration.

Author's contributions

JWN and HHC conceived and supervised the project. CYC provided materials. DJL, KTL,

and YJD performed RNA-seq experiments. HSH performed the analyses. JWN and HSH

wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank all members of the BIG lab for helpful comments and discussions.

Figure legends

Figure 1. Comprehensive coding and non-coding transcriptome maps of Yeonsan Ogye. **a.**

Yeonsan Ogye. **b.** A schematic flow for the analyses of coding and non-coding

transcriptomes and DNA methylation from twenty different tissues. **c.** A Circos plot

illustrating the expression variability (green bars) of lncRNA and protein-coding genes, the

methylation variability (red bars) at tissue-specific, differentially methylated CpG sites in

the promoters, and the correlation coefficients between expression and methylation levels

across chromosomes (heatmaps). **d.** Shown are the distributions of the maximum versus

mean expression values of lncRNA (red line) and protein-coding genes (black line) across

tissues (top), and the distributions of the maximum versus mean methylation levels of

each cytosine in the promoter of lncRNAs (red line) and protein-coding genes (black line)

(bottom). The vertical dotted lines indicate the median value of the respective distribution

(black for protein-coding genes and red for lncRNAs). **e.** Numbers of commonly or uniquely

expressed lncRNAs across tissues are shown in the phylogenetic tree of tissues. The

numbers at the leaf nodes indicate lncRNAs expressed in the indicated tissue (FPKM ≥ 1)

and the numbers at the internal nodes indicate those commonly expressed in the indicated

tissues. Of the expressed genes in a certain tissue, the fraction of the tissue-specific genes

(red for lncRNA and black for protein-coding genes) and the fraction of genes with a

differentially methylated region (DMR) in the promoters are indicated as bar graphs. Of the

genes with a DMR, tissue-specific genes (dark) and others (light) were distinguished and

the enrichment of the tissue-specific genes was tested using Fisher's exact test (* P ≤ 1

X e^{-5}, ** P ≤ 1 X e^{-10}, *** P ≤ 1 X e^{-20}). The scale bar represents 10.0, which is the unit of

120 differentially expressed genes across tissues.

Figure 2. Co-expression clusters of lncRNAs and functional annotations. Co-expression

clustering of lncRNAs across twenty tissues defines sixteen clusters and two sub-clusters

specific to a tissue or a set of similar tissues. The boxes outlined in a color indicate

clusters that have significant GO biological processes (orange bars) or KEGG pathway

terms (cyan bars) associated with the protein-coding genes co-expressed with lncRNAs in

the respective cluster. The significant enrichment of terms was tested using the

hypergeometric test and adjusted by FDR, indicated with a logarithmic scale on the X-axis

in the box. Clusters outlined in black are those that had neither a significant association

with any GO term nor any co-expressed protein-coding genes. Sub-clusters in the

clusters are indicated where appropriate. The number in each cluster indicates the

number of lncRNAs in the cluster and the number in the boxes with functional terms

indicates the number of co-expressed protein-coding genes.

Figure 3. lncRNAs as epigenetic activators. **a.** The proportions of lncRNAs with expression

levels that are correlated with the methylation level in the promoter of co-expressed protein-

coding genes (dark green) in each cluster are shown in bar graphs. The numbers were

compared to the mean methylation level of randomly selected protein-coding genes. To test

the significance of the enrichment of lncRNAs as epigenetic activator candidates, 1000

number-matched random cohorts were compared to the original numbers (* $P \leq 0.05$, ** $P \leq$

0.01, *** $P \leq 0.001$). **b.** lncRNAs as epigenetic activators whose expression levels are

negatively correlated with the methylation level in the promoters of protein-coding genes,

which in turn are negatively correlated with the level of protein-coding gene expression, as

shown in heatmaps. The key indicates the z-score range of the expression values. White

indicates N.A.

Figure 4. Co-transcriptional regulation of lncRNA and protein-coding genes by common

TFs. **a.** TFs (Sp1, Ap-2, Oct1, HSF2, and HB) with binding motifs that are significantly co-

enriched in the promoters of lncRNAs in a tissue-specific cluster and their co-expressed

protein-coding genes are shown in the heatmap. The TFs are expressed in the indicated

tissues. The significance of the motif enrichment was tested using MEME and E values

are presented with color codes (blue: more significance, yellow: less significance) in the

key. PCG indicates protein-coding gene. **b.** The HSF2 binding motif. A known motif is

shown in the top panel, a motif in lncRNA promoters is shown in the middle panel, and a

motif in protein-coding gene promoters is shown in the bottom panel. **c.** The expression

correlation between co-regulated genes (red boxes for lncRNAs and green boxes for protein-

coding genes) and HSF2 across tissues. Red lines indicate the significance level of the

correlation coefficient ($P \leq 0.05$). **e.** Expression pattern of HSF2 and its target genes that

have the top 5 correlations with HSF2.

Figure 5. Co-regulation of neighboring lncRNA and protein-coding genes. **a.** Shown are

the numbers of lncRNAs, classified by the distance from the closest protein-coding gene

(red for the $\leq$ 10kb group, orange for the $\leq$ 100kb group, and green for the > 100kb or on

another chromosome group) (left). *, **, and *** indicate $P \leq 0.05$, ≤ 0.01, and ≤ 0.001,

respectively. **b.** The average correlation coefficients of tissue-specific lncRNA and protein-

coding gene pairs in close neighborhoods ($\leq$ 10kb) are shown based on their relative

orientations (head-to-tail, tail-to-tail, and head-to-head) (red bars). The average

correlation coefficients of random pairs are also shown (blue bars) and those of tissue-

specific protein-coding gene pairs in close neighborhoods ($\leq$ 10kb) are shown with green

bars. *, **, and *** indicate $P \leq 0.05$, ≤ 0.01, and ≤ 0.001, respectively. Error bars indicate

the standard error. The number in the bars indicates the number of analyzed pairs. **c.** The

average correlation coefficients of neighboring lncRNA and protein-coding genes with

similar methylation levels in their promoters (methylation-related) are shown in bar graphs.

Otherwise, as in **b**. **d.** The average correlation coefficients of tissue-specific lncRNA and

protein-coding genes (methylation-unrelated), except for those of **c**. Otherwise, as in **b**. **e.**

The proportion of eRNAs (red) in the methylation-related group (**c**) and -unrelated group (**d**).

** indicates $P \leq 0.01$. **f.** The average correlation coefficients of tissue-specific eRNAs.

Otherwise, as in **b**. **g.** The average correlation coefficients of tissue-specific lncRNAs not

associated with enhancers. Otherwise, as in **b**. **h.** TF binding motifs significantly associated

with the eRNAs. The total count of the indicated TF binding sites in eRNAs is indicated in the

heatmap (left) and the significance of the association over the total background is indicated

with color-coded *P* values across tissues. The significance of a specific TF binding motif was

tested using a binomial test in each tissue.

Figure 6. Black tissue-specific lncRNAs with sequence and synteny conservation. **a.** The

expression patterns of differentially expressed lncRNAs in Ogye skin, compared to brown

leghorn skin samples. Expression levels are indicated with a color-coded Z-score (red for

low and blue for high expression) as shown in the key. **b.** A cartoon showing a lncRNA that is

syntenically conserved with up- and down-stream protein-coding genes in the human and/or

mouse genome. **c.** The fraction of lncRNAs with syntenic conservation in the human (blue),

mouse (green) or both (red) genomes is shown in the pie chart. Of the syntenically

conserved lncRNAs, the fraction of lncRNAs with sequence conservation (purple) in the

human or mouse genome is indicated in the secondary pie charts. **d.** The fraction of protein-

coding genes with synteny conservation is indicated in the pie chart. Otherwise, as in **c**. **e.**

The numbers of differentially expressed lncRNAs in black skin with evidence of sequence

and synteny conservation are indicated in a Venn diagram. **f.** Evidence for differential

expression (DE) + synteny + sequence (red), DE + synteny conservation (purple), or DE +

sequence conservation (blue) for 16 black-skin specific lncRNAs is shown in a heatmap. 104

non-specific lncRNAs with evidence of sequence + synteny conservation are indicated in

gray. The co-regulation models associated with a certain lncRNA are indicated to the left with

color codes (orange for HSF2 binding and green for eRNAs). * indicates the eRNA

associated with HSF2. The expression level is indicated with a color-coded z-score, as

shown in the key.

Figure 7. An example of black skin-specific lncRNAs with synteny conservation, which is

transcriptionally regulated by HSF2. **a.** Ogye lncRNA (lnc-TMEM184C) with synteny

conservation in human and mouse genomes (top). The lncRNA has an HSF2 binding motif in

its promoter; this motif is also present in the promoters of protein-coding genes with

correlated expression (below). Gray bar plots indicate the expression correlation between

the lncRNA and the protein-coding genes. **b.** The lnc-TMEM184C expression pattern across

20 tissues. **c.** GO terms that are significantly associated with the protein-coding genes that

are co-expressed with lnc-TMEM184C.

Additional files

Additional file 1: Supplementary Figures S1-S7.

Additional file 2: Table S1. Expression levels of all lncRNAs across twenty tissues.

Additional file 3: Table S2. Tissue-specific lncRNAs and ubiquitously expressed lncRNAs.

Additional file 4: Table S3. lncRNAs corresponding to each cluster (16 clusters).

Additional file 5: Table S4. Enriched biological process GO terms and KEGG pathways for

protein-coding genes co-expressed with each cluster (FDR ≤ 0.05).**Additional file 6:** Table

S5. lncRNAs associated with enhancers and neighboring protein-coding genes in terms of

their genomic orientations: head-to-tail, tail-to-tail, head-to-head.

Additional file 7: Table S6. lncRNAs that are differentially expressed between Ogye and

Brown leghorn skin.

Additional file 8: Table S7. Synteny-conserved lncRNAs and sequence-conserved lncRNAs.

References

References

1.   Dharmayanthi A.B., Terai Y., Sulandari S., Zein M.S.A., Akiyama T.,Satta Y., The origin and
evolution of fibromelanosis in domesticated chickens: Genomic comparison of Indonesian
Cemani and Chinese Silkie breeds. PLOS ONE, 2017: 12(4); e0173147.

2.   Dorshorst B., Molin A.-M., Rubin C.-J., Johansson A.M., Strömstedt L., Pham M.-H., et al., A
Complex Genomic Rearrangement Involving the Endothelin 3 Locus Causes Dermal
Hyperpigmentation in the Chicken. PLOS Genetics, 2011: 7(12); e1002412.

3.   Shinomiya A., Kayashima Y., Kinoshita K., Mizutani M., Namikawa T., Matsuda Y., et al., Gene
duplication of endothelin 3 is closely correlated with the hyperpigmentation of the internal
organs (Fibromelanosis) in silky chickens. Genetics, 2012: 190(2); 627-638.

4.   Li Y., Zhu X., Yang L., Li J., Lian Z., Li N., et al., Expression and network analysis of genes
related to melanocyte development in the Silky Fowl and White Leghorn embryos. Molecular
Biology Reports, 2011: 38(2); 1433-1441.

5.   Zhang J., Liu F., Cao J.,Liu X., Skin Transcriptome Profiles Associated with Skin Color in
Chickens. PLOS ONE, 2015: 10(6); e0127301.

6.   Ponting C.P., Oliver P.L.,Reik W., Evolution and Functions of Long Noncoding RNAs. Cell,
2009: 136(4); 629-641.

7.   Simon M.D., Pinter S.F., Fang R., Sarma K., Rutenberg-Schoenberg M., Bowman S.K., et al.,
High-resolution Xist binding maps reveal two-step spreading during X-chromosome
inactivation. Nature, 2013: 504(7480); 465-469.

8.   Morris K.V.,Mattick J.S., The rise of regulatory RNA. Nat Rev Genet, 2014: 15(6); 423-437.

9.   Wang X., Arai S., Song X., Reichart D., Du K., Pascual G., et al., Induced ncRNAs
allosterically modify RNA-binding proteins in cis to inhibit transcription. Nature, 2008:
454(7200); 126-30.

10.   Hellwig S.,Bass B.L., A starvation-induced noncoding RNA modulates expression of Dicer-
regulated genes. Proc Natl Acad Sci U S A, 2008: 105(35); 12897-902.

11.   Ferre F., Colantoni A.,Helmer-Citterich M., Revealing protein-lncRNA interaction. Brief
Bioinform, 2016: 17(1); 106-16.

12.   Merry C.R., Forrest M.E., Sabers J.N., Beard L., Gao X.H., Hatzoglou M., et al., DNMT1-
associated long non-coding RNAs regulate global gene expression and DNA methylation in
colon cancer. Hum Mol Genet, 2015: 24(21); 6240-53.

13.   Hirota K., Miyoshi T., Kugou K., Hoffman C.S., Shibata T.,Ohta K., Stepwise chromatin
remodelling by a cascade of transcription initiation of non-coding RNAs. Nature, 2008:
456(7218); 130-4.

14.   Zalfa F., Giorgi M., Primerano B., Moro A., Di Penta A., Reis S., et al., The fragile X syndrome
protein FMRP associates with BC1 RNA and regulates the translation of specific mRNAs at
synapses. Cell, 2003: 112(3); 317-27.

15.   Liu X., Li D., Zhang W., Guo M.,Zhan Q., Long non-coding RNA gadd7 interacts with TDP-43
and regulates Cdk6 mRNA decay. Embo j, 2012: 31(23); 4415-27.

16.   Ulitsky I.,Bartel D.P., lincRNAs: genomics, evolution, and mechanisms. Cell, 2013: 154(1); 26-
46.

17.   Garding A., Bhattacharya N., Claus R., Ruppel M., Tschuch C., Filarsky K., et al., Epigenetic
upregulation of lncRNAs at 13q14. 3 in leukemia is linked to the In Cis downregulation of a
gene cluster that targets NF-kB. PLoS Genet, 2013: 9(4); e1003373.

18.   Zhang B., Arun G., Mao Y.S., Lazar Z., Hung G., Bhattacharjee G., et al., The lncRNA Malat1
is dispensable for mouse development but its transcription plays a cis-regulatory role in the
adult. Cell reports, 2012: 2(1); 111-123.

19.   Jariwala N.,Sarkar D., Emerging role of lncRNA in cancer: a potential avenue in molecular
medicine. Ann Transl Med, 2016: 4(15); 286.

20.   Sahu A., Singhal U.,Chinnaiyan A.M., Long noncoding RNAs in cancer: from function to
translation. Trends in cancer, 2015: 1(2); 93-109.

21.   Kim T.-K., Hemberg M., Gray J.M., Costa A.M., Bear D.M., Wu J., et al., Widespread
transcription at neuronal activity-regulated enhancers. Nature, 2010: 465(7295); 182-187.

22.   Marques A.C., Hughes J., Graham B., Kowalczyk M.S., Higgs D.R.,Ponting C.P., Chromatin

signatures at transcriptional start sites separate two equally populated yet distinct classes of intergenic long noncoding RNAs. Genome Biology, 2013: 14(11); R131.

23. De Santa F., Barozzi I., Mietton F., Ghisletti S., Polletti S., Tusi B.K., et al., A large fraction of extragenic RNA pol II transcription sites overlap enhancers. PLoS Biol, 2010: 8(5); e1000384.

24. Orom U.A., Derrien T., Beringer M., Gumireddy K., Gardini A., Bussotti G., et al., Long noncoding RNAs with enhancer-like function in human cells. Cell, 2010: 143(1); 46-58.

25. Lai F., Orom U.A., Cesaroni M., Beringer M., Taatjes D.J., Blobel G.A., et al., Activating RNAs associate with Mediator to enhance chromatin architecture and transcription. Nature, 2013: 494(7438); 497-501.

26. Feng J., Bi C., Clark B.S., Mady R., Shah P.,Kohtz J.D., The Evf-2 noncoding RNA is transcribed from the Dlx-5/6 ultraconserved region and functions as a Dlx-2 transcriptional coactivator. Genes Dev, 2006: 20(11); 1470-84.

27. Bertani S., Sauer S., Bolotin E.,Sauer F., The noncoding RNA Mistral activates Hoxa6 and Hoxa7 expression and stem cell differentiation by recruiting MLL1 to chromatin. Mol Cell, 2011: 43(6); 1040-6.

28. Dimitrova N., Zamudio J.R., Jong R.M., Soukup D., Resnick R., Sarma K., et al., LincRNA-p21 activates p21 in cis to promote Polycomb target gene expression and to enforce the G1/S checkpoint. Mol Cell, 2014: 54(5); 777-90.

29. Gonzalez I., Munita R., Agirre E., Dittmer T.A., Gysling K., Misteli T., et al., A lncRNA regulates alternative splicing via establishment of a splicing-specific chromatin signature. Nat Struct Mol Biol, 2015: 22(5); 370-6.

30. Mercer T.R., Dinger M.E., Sunkin S.M., Mehler M.F.,Mattick J.S., Specific expression of long noncoding RNAs in the mouse brain. Proc Natl Acad Sci U S A, 2008: 105(2); 716-21.

31. Derrien T., Johnson R., Bussotti G., Tanzer A., Djebali S., Tilgner H., et al., The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. Genome Res, 2012: 22(9); 1775-89.

32. Cabili M.N., Trapnell C., Goff L., Koziol M., Tazon-Vega B., Regev A., et al., Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. Genes Dev, 2011: 25(18); 1915-27.

33. Quinn J.J.,Chang H.Y., Unique features of long non-coding RNA biogenesis and function. Nat Rev Genet, 2016: 17(1); 47-62.

34. Djebali S., Davis C.A., Merkel A., Dobin A., Lassmann T., Mortazavi A., et al., Landscape of transcription in human cells. Nature, 2012: 489(7414); 101-8.

35. Ulitsky I., Shkumatava A., Jan C.H., Sive H.,Bartel D.P., Conserved function of lincRNAs in vertebrate embryonic development despite rapid sequence evolution. Cell, 2011: 147(7); 1537-50.

36. Li L.,Chang H.Y., Physiological roles of long noncoding RNAs: insight from knockout mice. Trends in Cell Biology, 2014: 24(10); 594-602.

37. Ulitsky I., Shkumatava A., Jan Calvin H., Sive H.,Bartel David P., Conserved Function of lincRNAs in Vertebrate Embryonic Development despite Rapid Sequence Evolution. Cell, 2011: 147(7); 1537-1550.

38. Nam J.W.,Bartel D.P., Long noncoding RNAs in C. elegans. Genome Res, 2012: 22(12); 2529-40.

39. Pauli A., Valen E., Lin M.F., Garber M., Vastenhouw N.L., Levin J.Z., et al., Systematic identification of long noncoding RNAs expressed during zebrafish embryogenesis. Genome Res, 2012: 22(3); 577-91.

40. Iyer M.K., Niknafs Y.S., Malik R., Singhal U., Sahu A., Hosono Y., et al., The landscape of long noncoding RNAs in the human transcriptome. Nat Genet, 2015: 47(3); 199-208.

41. Dinger M.E., Amaral P.P., Mercer T.R., Pang K.C., Bruce S.J., Gardiner B.B., et al., Long noncoding RNAs in mouse embryonic stem cell pluripotency and differentiation. Genome Res, 2008: 18(9); 1433-45.

42. Faghihi M.A., Modarresi F., Khalil A.M., Wood D.E., Sahagan B.G., Morgan T.E., et al., Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of beta-secretase. Nat Med, 2008: 14(7); 723-30.

43. Leucci E., Vendramin R., Spinazzi M., Laurette P., Fiers M., Wouters J., et al., Melanoma addiction to the long non-coding RNA SAMMSON. Nature, 2016: 531(7595); 518-522.

44. Ghosh S., Sati S., Sengupta S.,Scaria V., Distinct patterns of epigenetic marks and
transcription factor binding sites across promoters of sense-intronic long noncoding RNAs.
Journal of genetics, 2015: 94(1); 17-25.
45. Guttman M., Amit I., Garber M., French C., Lin M.F., Feldser D., et al., Chromatin signature
reveals over a thousand highly conserved large non-coding RNAs in mammals. Nature, 2009:
458(7235); 223-227.
46. Pang K.C., Dinger M.E., Mercer T.R., Malquori L., Grimmond S.M., Chen W., et al., Genome-
wide identification of long noncoding RNAs in CD8+ T cells. J Immunol, 2009: 182(12); 7738-
48.
47. Liao Q., Liu C., Yuan X., Kang S., Miao R., Xiao H., et al., Large-scale prediction of long non-
coding RNA functions in a coding-non-coding gene co-expression network. Nucleic Acids Res,
2011: 39(9); 3864-78.
48. Lv Y., Liang Z., Ge M., Qi W., Zhang T., Lin F., et al., Genome-wide identification and
functional prediction of nitrogen-responsive intergenic and intronic long non-coding RNAs in
maize (Zea mays L.). BMC Genomics, 2016: 17(1); 350.
49. Wang Y., Xue S., Liu X., Liu H., Hu T., Qiu X., et al., Analyses of long non-coding RNA and
mRNA profiling using RNA sequencing during the pre-implantation phases in pig endometrium.
Scientific reports, 2016: 6.
50. Cogill S.B.,Wang L., Co-expression Network Analysis of Human lncRNAs and Cancer Genes.
Cancer Inform, 2014: 13(Suppl 5); 49-59.
51. Weikard R., Hadlich F.,Kuehn C., Identification of novel transcripts and noncoding RNAs in
bovine skin by deep next generation sequencing. BMC Genomics, 2013: 14(1); 789.
52. Billerey C., Boussaha M., Esquerré D., Rebours E., Djari A., Meersseman C., et al.,
Identification of large intergenic non-coding RNAs in bovine muscle using next-generation
transcriptomic sequencing. BMC Genomics, 2014: 15(1); 499.
53. Li T., Wang S., Wu R., Zhou X., Zhu D.,Zhang Y., Identification of long non-protein coding
RNAs in chicken skeletal muscle using next generation sequencing. Genomics, 2012: 99(5);
292-298.
54. Ren H., Wang G., Chen L., Jiang J., Liu L., Li N., et al., Genome-wide analysis of long non-
coding RNAs at early stage of skin pigmentation in goats (Capra hircus). BMC Genomics,
2016: 17(1); 67.
55. Al-Tobasei R., Paneru B.,Salem M., Genome-Wide Discovery of Long Non-Coding RNAs in
Rainbow Trout. PLOS ONE, 2016: 11(2); e0148940.
56. Dobin A., Davis C.A., Schlesinger F., Drenkow J., Zaleski C., Jha S., et al., STAR: ultrafast
universal RNA-seq aligner. Bioinformatics, 2013: 29(1); 15-21.
57. Trapnell C., Williams B.A., Pertea G., Mortazavi A., Kwan G., van Baren M.J., et al.,
Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and
isoform switching during cell differentiation. Nat Biotechnol, 2010: 28(5); 511-5.
58. You B.H., Yoon S.H.,Nam J.W., High-confidence coding and noncoding transcriptome maps.
Genome Res, 2017: 27(6); 1050-1062.
59. Li B.,Dewey C.N., RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC bioinformatics, 2011: 12(1); 323.
60. Greco S., Gorospe M.,Martelli F., Noncoding RNA in age-related cardiovascular diseases. J
Mol Cell Cardiol, 2015: 83; 142-55.
61. Cech T.R.,Steitz J.A., The noncoding RNA revolution-trashing old rules to forge new ones.
Cell, 2014: 157(1); 77-94.
62. Felsenstein J., PHYLIP - Phylogeny Inference Package (Version 3.2). Cladistics, 1989: 5;
164-166.
63. Krueger F.,Andrews S.R., Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics, 2011: 27(11); 1571-1572.
64. Huang da W., Sherman B.T.,Lempicki R.A., Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. Nat Protoc, 2009: 4(1); 44-57.
65. Wu L., Murat P., Matak-Vinkovic D., Murrell A.,Balasubramanian S., Binding interactions
between long noncoding RNA HOTAIR and PRC2 proteins. Biochemistry, 2013: 52(52); 9519-
9527.
66. Zhao J., Sun B.K., Erwin J.A., Song J.J.,Lee J.T., Polycomb proteins targeted by a short

repeat RNA to the mouse X chromosome. Science, 2008: 322(5902); 750-6.

67. Chalei V., Sansom S.N., Kong L., Lee S., Montiel J.F., Vance K.W., et al., The long non-
coding RNA Dali is an epigenetic regulator of neural differentiation. eLife, 2014: 3; e04530.

68. Di Ruscio A., Ebralidze A.K., Benoukraf T., Amabile G., Goff L.A., Terragni J., et al., DNMT1-
interacting RNAs block gene-specific DNA methylation. Nature, 2013: 503(7476); 371-376.

69. Machanick P.,Bailey T.L., MEME-ChIP: motif analysis of large DNA datasets. Bioinformatics,
2011: 27(12); 1696-1697.

70. Gupta S., Stamatoyannopoulos J.A., Bailey T.L.,Noble W.S., Quantifying similarity between
motifs. Genome Biology, 2007: 8(2); R24.

71. Wingender E., Kel A.E., Kel O.V., Karas H., Heinemeyer T., Dietze P., et al., TRANSFAC,
TRRD and COMPEL: towards a federated database system on transcriptional regulation.
Nucleic Acids Res, 1997: 25(1); 265-8.

72. Mi H., Dong Q., Muruganujan A., Gaudet P., Lewis S.,Thomas P.D., PANTHER version 7:
improved phylogenetic trees, orthologs and collaboration with the Gene Ontology Consortium.
Nucleic Acids Research, 2010: 38(suppl_1); D204-D210.

73. Thomas P.D., Campbell M.J., Kejariwal A., Mi H., Karlak B., Daverman R., et al., PANTHER: a
library of protein families and subfamilies indexed by function. Genome research, 2003: 13(9);
2129-2141.

74. Li W., Notani D.,Rosenfeld M.G., Enhancers as non-coding RNA transcription units: recent
insights and future perspectives. Nat Rev Genet, 2016: 17(4); 207-223.

75. Seki R., Li C., Fang Q., Hayashi S., Egawa S., Hu J., et al., Functional roles of Aves class-
specific cis-regulatory elements on macroevolution of bird-specific features. Nat Commun,
2017: 8; 14229.

76. Kim T.-K., Hemberg M., Gray J.M., Costa A.M., Bear D.M., Wu J., et al., Widespread
transcription at neuronal activity-regulated enhancers. Nature, 2010: 465(7295); 182-187.

77. Wang D., Garcia-Bassets I., Benner C., Li W., Su X., Zhou Y., et al., Reprogramming
transcription by distinct classes of enhancers functionally defined by eRNA. Nature, 2011:
474(7351); 390-394.

78. Ito S., Wakamatsu K.,Ozeki H., Chemical analysis of melanins and its application to the study
of the regulation of melanogenesis. Pigment Cell Res, 2000: 13 Suppl 8; 103-9.

79. Ito S.,Wakamatsu K., Chemistry of mixed melanogenesis--pivotal roles of dopaquinone.
Photochem Photobiol, 2008: 84(3); 582-92.

80. Dunn K.J., Williams B.O., Li Y.,Pavan W.J., Neural crest-directed gene transfer demonstrates
Wnt1 role in melanocyte expansion and differentiation during mouse development.
Proceedings of the National Academy of Sciences, 2000: 97(18); 10050-10055.

81. Guo H., Xing Y., Liu Y., Luo Y., Deng F., Yang T., et al., Wnt/β-catenin signaling pathway
activates melanocyte stem cells in vitro and in vivo. Journal of Dermatological Science, 2016:
83(1); 45-51.

82. Wu T.D.,Watanabe C.K., GMAP: a genomic mapping and alignment program for mRNA and
EST sequences. Bioinformatics, 2005: 21(9); 1859-1875.

83. Andrews S., FastQC A Quality Control tool for High Throughput Sequence Data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

84. Zhao Y., Li H., Fang S., Kang Y., wu W., Hao Y., et al., NONCODE 2016: an informative and
valuable data source of long non-coding RNAs. Nucleic Acids Research, 2016: 44(D1); D203-
D208.

85. Kong L., Zhang Y., Ye Z.-Q., Liu X.-Q., Zhao S.-Q., Wei L., et al., CPC: assess the protein-
coding potential of transcripts using sequence features and support vector machine. Nucleic
Acids Research, 2007: 35(suppl_2); W345-W349.

86. Yeo G.,Burge C.B., Maximum entropy modeling of short sequence motifs with applications to
RNA splicing signals. J Comput Biol, 2004: 11(2-3); 377-94.

87. Anders S.,Huber W., Differential expression analysis for sequence count data. Genome Biol,
2010: 11(10); R106.

1025

Figure 1

Figure 2

Cerebrum & Cerebellum (1617)

- Neuroactive ligand-receptorinteraction
- Ion transport
- Calcium signaling pathway
- Cation transport
- Neuron development

$-\log_{10}(\text{FDR})$

Mature egg (415)

- Extracellular matrix organization
- ECM-receptor interaction
- Extracellular structure organization
- Cell adhesion

$-\log_{10}(\text{FDR})$

Kidney (591)

- Oxidation reduction
- Transmembrane transport
- Sodium ion transport

$-\log_{10}(\text{FDR})$

Immature egg (606)

- Cell cycle phase
- M phase
- Cell cycle process
- Cell cycle
- Spliceosome

$-\log_{10}(\text{FDR})$

Eye (367)

- Sensory perception of light stimulus
- Visual perception
- Sensory perception

$-\log_{10}(\text{FDR})$

Black tissues (420)

- Appendage development
- Limb development
- Limb morphogenesis

$-\log_{10}(\text{FDR})$

Breast (973)

- Protein catabolic process
- Modification-macromolecule catabolic process
- Modification-protein catabolic process
- Proteasome

$-\log_{10}(\text{FDR})$

Bone marrow (902)

- M phase
- chromosome organization
- Cell cycle
- DNA replication
- organelle fission

$-\log_{10}(\text{FDR})$

Heart (491)

- Oxidative phosphorylation
- Generation of precursor metabolites
- -and energy
- Citrate cycle
- Cellular respiration
- Cofactor metabolic process

$-\log_{10}(\text{FDR})$

Spleen (748)

- Leukocyte activation
- Cell activation
- Lymphocyte activation
- Cytokine-cytokine receptor interaction
- Intestinal immune network
- -for IgA production

$-\log_{10}(\text{FDR})$

Pancreas

Uterus

Liver

Lung

Gall bladder

Black tissues

Gizzard

Figure 3

a.

Percentage (%)

Brain Kidney Eye Mature egg Immature egg Breast Heart Black tissues Bone marrow Spleen

*** ** ** * **

■ Co-expressed protein-coding genes
■ Random control

b.

Z−score
−4 −2 0 2 4

lncRNA expression

Methylation level of
protein-coding genes

Protein-coding gene
expression

Breast
Liver
Bone marrow
Fascia
Cerebrum
Gizzard
Immature egg
Crest
Spleen
Mature egg
Cerebellum
Gall bladder
Kidney
Heart
Uterus
Pancreas
Lungs
Skin
Eye
Shank

Figure 4

a.

pcg: protein-coding gene

b.

c.

d.

a.

Figure 6

Figure 7

a.

b.

c.