Effects of background and contour luminance on the hue and brightness of the Watercolor effect

Peggy Gerardin[2], Michel Dojat[3], Kenneth Knoblauch[2], Frédéric Devinck[1*],

[1] Université Rennes 2, LP3C, EA 1285, 35000 Rennes, France

[2] Univ Lyon, Université Claude Bernard Lyon 1, Inserm, Stem Cell and Brain Research Institute U1208, 69500 Bron, France. 3 Université Lyon 1, 69003 Lyon, France

[3] Univ. Grenoble Alpes, Inserm, CHU Grenoble Alpes, GIN, 38000 Grenoble, France

* Corresponding author. Tel.: +33-2-99-19-59; Fax: +33-2-99-14-19-54

e-mail address: frederic.devinck@univ-rennes2.fr

Present address: Département de Psychologie, Université Rennes 2, 35043 Rennes Cedex, France.

Abstract

Conjoint measurement was used to investigate the joint influences of the luminance of the background and the inner contour on hue- and brightness filling-in for a stimulus configuration generating a water-color effect (WCE), i.e., a wiggly bi-chromatic contour enclosing a region with the lower luminance component on the exterior. Two stimuli with the background and inner contour luminances covarying independently were successively presented, and in separate experiments, the observer judged which member of the pair's interior regions contained a stronger hue or was brighter. Braided-contour control stimuli that generated little or no perceptual filling-in were also used to assess whether observers were judging the interior regions and not the contours themselves. Three nested models of the contributions of the background and inner contour to the judgments were fit to the data by maximum likelihood and evaluated by likelihood ratio tests. Both stimulus components contributed to both the hue and brightness of the interior region with increasing luminance of the inner contour generating an assimilative filling-in for the hue judgments but a contrast effect for the brightness judgments. Control analyses showed negligible effects for the order of the luminance of the background or inner contour on the judgments. An additive contribution of both components was rejected in favor of a saturated model in which the responses depended on the levels of both stimulus components. For the hue judgments, increased background luminance led to

greater hue filling-in at higher luminances of the interior contour. For the

brightness judgments, the higher background luminance generated less brightness

filling-in at higher luminances of the interior contour. The results indicate

different effects of the inner contour and background on the induction of the

brightness and coloration percepts of the WCE, suggesting that they are mediated

by different mechanisms.

49
50

51   Keywords: filling-in, color assimilation, watercolor effect, color appearance,

52   scaling, conjoint measurement, MLCM.

1. Introduction.

Color appearance is not determined only by the local light signals from each object but is also influenced by global contextual features. The watercolor effect (WCE) is an interesting phenomenon for studying such processes (Pinna, 1987; Pinna et al., 2001). A pair of wiggly contours composed of a light chromatic contour (e.g., orange) surrounded by a darker chromatic contour (e.g., purple) bounding an achromatic surface area elicits a filling-in of the hue of the lighter contour over the entire enclosed area (Figure 1a). The WCE is distinguished from other assimilation illusions by its large spatial extent; the phenomenon has been observed over distances of up to 45 deg (Pinna et al., 2001). In addition to the assimilative color spreading, the subjectively colored area is perceived as figure while the surrounding area appears as ground (Pinna et al., 2003; Pinna & Tanca, 2008; Tanca & Pinna, 2008).

Figure 1 about here

Studies of the WCE have typically examined the effects of the inducer configuration producing the WCE. For example, the intensity of the filling-in percept appears greater with increases in luminance contrast between the inner and outer contours for an achromatic WCE (Cao et al., 2011) and for a WCE that has both luminance and chromatic components (Devinck et al., 2005; Devinck & Knoblauch, 2012). Devinck et al. (2005) noted that observers did

not need to modify significantly the luminance of the enclosed area in a

matching experiment. Other critical characteristics of the inducing contours that

modulate the strength of the WCE include the continuity and contiguity of the

contour pairs (Devinck & Spillmann, 2009; Devinck & Knoblauch, 2012).

Recent demonstrations of the sensitivity of the phenomenon to contour

adaptation provide additional support for a role of contour integration

mechanisms in the WCE (Coia & Crognale, 2017). The strength of the

phenomenon was found to be size-tuned with the strongest WCE observed for a

contour width of about 15 arcmin and was optimal for equal contour widths

(Devinck et al., 2014a). While the WCE has been reported for linear contours

(Pinna et al., 2001), its strength is nearly independent of the amplitude of

contour undulation but increases with contour frequency up to an asymptotic

level (Gerardin et al., 2014). Finally, Pinna et al. (2001) demonstrated that

several different color pairs can generate the coloration effect (see also Devinck

et al., 2005). Specifically, Devinck et al. (2006) demonstrated that the

coloration effect is stronger when the chromatic contrast is larger. Thus, the

coloration effect depends on a conjunction of chromatic and luminance contrasts

but also on spatial parameters of the inner and outer contours.

The WCE is perceptually salient but has proved difficult to quantify with

precision showing large variability within and across observers (Cao et al.,

2011; Devinck et al., 2005; von der Heydt & Pierson, 2006). More recently, the

WCE was quantified by using paired-comparison methods that have been

extended to estimate perceptual scales within a signal detection framework (Devinck & Knoblauch, 2012). Two such procedures are Maximum Likelihood Difference Scaling or MLDS (Maloney & Yang, 2003; Knoblauch & Maloney, 2008, 2012) and Maximum Likelihood Conjoint Measurement or MLCM (Ho et al., 2008; Knoblauch & Maloney, 2012). Difference scaling is useful for measuring perceptual strength along a single physical dimension, whereas conjoint measurement was conceived to assess the combined effects of several dimensions on appearance (Falmagne, 1985 ; Knoblauch & Maloney, 2012 ; Krantz et al., 1971; Luce & Tukey, 1964; Roberts, 1979). MLCM has been successfully applied to estimate perceptual scales associated with different sets of physical continua including surface material properties (Ho, Landy & Maloney, 2008; Qi, Chantler, Siebert & Dong, 2015; Hansmann-Roth & Mamassian, 2017), color appearance (Gerardin et al., 2014; Rogers, Knoblauch & Franklin, 2016) and time perception (Lisi & Gorea, 2016). The signal detection decision model allows specifying the perceptual scales in terms of the signal detection parameter d' (Gerardin et al., 2014; Knoblauch & Maloney, 2012).

The aim of the present study is to estimate perceptual scales for two dimensions, the luminance elevation of the inner contour and the luminance elevation of the background. While the luminance contrast between the inner and outer contours has been tested intensively in the WCE, experiments evaluating the influence of the background luminance are scarce. Indeed, the

WCE has generally been demonstrated for a background of higher luminance than both inner and outer contours. Although the surround (e.g., the background) is known to be an important influence of color appearance (Brenner & Cornelissen, 2002 ; Brown & MacLeod, 1997 ; Shevell, 1978 ; Walraven, 1976), it has not been systematically explored for the coloration effect in the WCE. In addition, most studies of the WCE focus solely on its coloration effect. Here, we also investigate the influences of the background and inner contour luminances on the perceived brightness of the interior region. In summary, we employed conjoint measurement to study how both the background and the inner contour luminances influence judgments of both the hue and brightness in the WCE.

2. General Methods

2.1. Observers

Four observers participated in these experiments. Three were naïve and the fourth was one of the authors. Observers ranged in age between 26 and 40 years. All had normal color vision as tested with the Farnsworth Panel D15, and had normal or corrected-to-normal visual acuity. Experiments were performed in accordance with the principles of the Declaration of Helsinki for the protection of human subjects.

2.2. Apparatus

Stimuli were presented on a NEC MultiSync FP2141sb color CRT monitor

driven by a Cambridge Research ViSaGe graphic board with a color resolution

of 14 bits per gun (Cambridge Research Systems, Rochester, United Kingdom).

The experimental software was written to generate all stimuli, control stimulus

presentation and collect responses in MATLAB 7.9 (MathWorks,

http://mathworks.com), using the CRS Toolbox extensions. The monitor was

calibrated using an OptiCal photometer with the calibration routines of

Cambridge Resarch Systems. Observer position was stabilized by a chinrest and

observer-to-screen distance was 80 cm. Experiments were performed in a dark

room. Both eyes were used for viewing.

2.3. Stimuli

The stimuli were constructed as Fourier descriptors (Zahn & Roskies, 1972).

Each stimulus was defined with respect to a circle of 3.2 deg diameter whose

radius, *r*, was modulated sinusoidally as a function of angle according to the

equation:

$$R(\theta) = r + A\sin(2\pi f\theta) \qquad (1)$$

where *R* is the stimulus radius at angle θ, *r* the average radius of the stimulus, *A*

the modulation and *f* the frequency in cycles per revolution (cpr). In the present

study, the frequency was fixed at *f* = 10 cpr and the amplitude of both contours

at *A* = 0.36 (Figure 1b, left).

All stimuli were composed of three colors: an orange inner contour ($x,y =$

0.44, 0.43) with the luminance varying from 30.02 cd/m^2 to 62.74 cd/m^2 and a

purple outer contour ($x,y = 0.31, 0.11$; $Y = 21.12$ cd/m^2), presented on a neutral

white background ($x,y = 0.29, 0.32$) with the luminance of the background (both

outside and inside of the contours) varying between 35.56 cd/m^2 and 65.56

cd/m^2. The contour pairs were each of width 16 arcmin, i.e., 8 arcmin for the

inner and outer contours, each.

Stimuli were specified in the DKL color space (MacLeod & Boynton,

1979; Krauskopf, Williams & Heeley, 1982; Derrington, Krauskopf & Lennie,

1984). DKL color space is a three-dimensional opponent-modulation space

based on the Smith and Pokorny (1975) cone fundamentals. The sum of L and

M cone excitations varies on one axis (luminance), while M cone excitation

subtracted from L cone excitation varies on the second axis (L - M); and the sum

of L and M cone excitations subtracted from S cone excitation varies on the

third axis (S - (L + M)). The DKL axes were scaled between -1 and 1, where +/-

1 corresponds to the maximum contrast for each axis on the monitor. The

stimuli were specified with the purple and orange contours at azimuth of 320

and 45 deg respectively. Luminance of the independent variables is specified as

elevation from the equiluminant plane. The luminance elevations of the orange

contour in DKL color space varied from -0.6 to 0 while the luminance elevation

of the background ranged from -0.5 to 0.

In the present study, five levels of luminance inner contour and five levels

of luminance background were used. All levels were crossed creating a 5 × 5

grid with a total of 25 stimuli. Figure 2 shows an example of the range of

stimuli used, with the inner contour luminance varying across rows and the

background luminance across columns.

191   **Figure 2 about here**

Control stimuli were also tested consisting of patterns that were identical

to the test stimuli except that the contours were interlaced (Figure 1b, right) and

generated little filling-in, as previously demonstrated (Devinck & Knoblauch,

2012; Devinck et al., 2014a; Gerardin et al., 2014). These control stimuli were

used to verify that observers responded to the filling-in appearance and not to

other stimulus features.

2.4. Procedure

On each trial, two different stimuli chosen randomly from the 5 × 5 grid were

presented in succession to the observer. Observers performed two tasks in

separate randomly ordered and counter-balanced sessions in which they

compared the interior regions of the two successively presented stimuli. In the

first task, observers were instructed to judge which central region evoked the

strongest orange hue. In the second task, observers were asked to judge which

central region appeared brighter. An equal number of test and control stimuli were interleaved in each session. With 5 levels along each of the dimensions varied, there are (25 * 24)/2 = 300 unordered pairs. Stimuli were randomly ordered for each presentation. On each trial, a randomly chosen pair of test or control stimuli was presented. A session consisted of the random presentation of all 600 test and control pairs. Each task was repeated five times, yielding 1500 test and 1500 control trials for each observer.

Prior to the experiment, observers were dark-adapted for 3 min. At the beginning of each trial, a fixation cross was presented in the center of the screen of duration 500 ms. At its extinction, the first pattern was presented during 500 ms followed by a fixation cross for 500 ms, and then the second pattern for 500 ms., followed by a blank screen. The observer's response initiated a 1 s pause before the start of the next trial. An initial practice block of 10 trials preceded the experiment. The experimental session started, when the observer felt at ease with the task, otherwise additional practice sessions were run. A free viewing procedure was used to ensure that observer's judgments were based on foveal views of the stimuli.

2.5. Model

The data were analyzed as a decision process within the framework of a signal detection model and fit by maximum likelihood (Ho et al., 2008; Knoblauch & Maloney, 2012). Three nested models of the decision process are fit to obtain

the best prediction of the set of observers' choices: an independence model, an

additive model and a saturated model. Each model yields estimates of

perceptual scale values or internal responses that have the property that equal

differences in response are perceptually equal. The independence model fits the

observer's judgments based on only one of the component dimensions. The

additive model fits the judgments based on the sum of component psychological

responses generated by the physical dimensions. The saturated model fits the

observer's judgments including an interaction term that depends on the specific

levels of the two components in addition to their simple additive combination.

The three models are then evaluated using a nested likelihood ratio test. This is

done separately for the experiments based on hue, and brightness judgments.

The formal description of the model is described next and follows similar

descriptions elsewhere (Gerardin et al., 2014; Ho et al., 2008; Knoblauch &

Maloney, 2012).

We represent the stimulus levels along the two dimensions by a variable

$\phi_{i,j}$, where i and j correspond to the luminance levels of the background and the

inner contour, respectively. In the decision models, each of the dimensions

contributes a response, ψ_i^1, ψ_j^2, to the intensity of the perceived filling-in

depending on the corresponding physical intensity levels, where the superscripts

correspond to the responses to the background and interior contour luminances,

respectively. In the additive model, when observers judge which central area is

the more saturated orange color or which appears to be brighter, we suppose that

the filling-in response is the sum of the component responses

$$\psi_{i,j} = \psi_i^1 + \psi_j^2. \qquad (2)$$

Observers compare the two presented central surfaces and the difference

between the filling-in strength of the stimulus $\phi_{i,j}$ and the stimulus $\phi_{k,l}$ is

computed as follows

$$\Delta(i,j,k,l) = (\psi_i^1 + \psi_j^2) - (\psi_k^1 + \psi_l^2) + \varepsilon \qquad (3)$$

where ε refers to additive noise in the decision process and is modeled as a

Gaussian random variable with $\mu = 0$ and variance $= \sigma^2$. In plots, we indicate

the stimulus level by the index and not by the physical units, allowing both

dimensions to be plotted together. With 5 levels along each dimension, there are

2 * 5 levels plus 1 variance = 11 parameters to estimate. To make the model

identifiable, however, the response at the lowest level along each dimension is

arbitrarily set to 0, $\psi_1^1 = \psi_1^2 = 0$, and the variance is fixed to 1 for each

estimated value, yielding only 8 parameters to estimate. The parameter values,

ψ_i^j are chosen to maximize the likelihood, $\mathcal{L}(\Psi; R)$, over the ensemble of

choices, R, made by the observer.

$$\mathcal{L}(\Psi; R) = \prod_i \Phi\left(\frac{\Delta_i}{2}\right)^{R_i} \left(1 - \Phi\left(\frac{\Delta_i}{2}\right)\right)^{1 - R_i}, \qquad (4)$$

where Φ is a cumulative distribution function for a Gaussian with mean 0 and

variance 1. The 2 in the denominator of the argument scales the variance for

each value of ψ to 1 so that the perceptual scale values are parameterized in

terms of d'. In practice, this is performed using a Generalized Linear Model

with the **MLCM** package in R (Knoblauch & Maloney, 2012, 2014).

If the observer's judgments depend on only one of the component

dimensions, we obtain the independence model, reducing the decision variable

to

$$\Delta(i,j,k,l) = \psi_i^1 - \psi_k^1 + \varepsilon \quad (5)$$

where the judgments depend on only dimension 1, here. In this reduced model,

the values of ψ_j^2 are fixed at 0 and there are only 4 free parameters to estimate.

Replacing the superscript 1 by 2 yields the independence model for the other

dimension.

Finally, the saturated observer model includes an interaction factor that

depend on the intensity levels of both dimensions; the decision variable is

defined as follows:

$$\Delta(i,j,k,l) = (\psi_i^1 + \psi_j^2 + \psi_{ij}^{12}) - (\psi_k^1 + \psi_l^2 + \psi_{kl}^{12}) + \varepsilon \quad (6)$$

Due to the interaction terms, the responses cannot be explained by a simple

additive combination of components as in the previous two models. With 5

levels along each dimension, only one cell in the 5x5 grid of responses is fixed

at 0 leading to 24 (the maximum) free parameters to estimate, which is the

origin of the term saturated.

We analysed the data using the MLCM package (Knoblauch & Maloney,

2012, 2014) in the open source software R (R Core Team, 2017) to estimate the

perceptual scale values and model the contribution of both dimensions. The

likelihood ratio tests were evaluated using a χ^2 statistic with degrees of freedom

the difference in number of parameters fit for each pair of models.

3. Results

Judgments based on color and on brightness from the observers are shown in

Figure 3 for test and control conditions in Conjoint Proportion Plots or CPP (Ho

et al., 2008 ; Knoblauch & Maloney, 2012). In the CPP, the raw data are

presented in a grid format in which each cell of the grid corresponds to one

stimulus pair comparison. Each CPP contains all stimulus combinations and

summarizes the proportion of times the stimulus S_{kl} was judged for one response

criterion, hue (a) or brightness (b), to show a greater filling-in than the stimulus

S_{ij}, coded according to the grey levels indicated by the color bar presented on the

right side of each graph. The levels of both dimensions are represented along

each axis where the 5×5 outer check indicates the stimulus levels along one

dimension and with each outer check subdivided into smaller 5×5 checks

indicating the stimulus levels for the second dimension. Figure 3c shows the

expected pattern of responses for an ideal observer who chooses only the higher

level along one of the two stimulus dimension. The CPP presented on the left

side indicates the results when the judgments depend on the first dimension

alone (here, the background luminance) and the CPP displayed on the right side

when the judgment depend on the second dimension (here, the inner contour luminance).

Results from the hue and brightness judgments for each observer are shown in Figure 3a and 3b respectively with the results displayed on the top row for the test condition and on the bottom row for the control condition in each figure. For the hue judgments, the CPP for the test stimuli resembles more closely the ideal CPP displayed for the second dimension, suggesting that the luminance of the inner contour contributed more strongly to the choices than the luminance of the background. For the brightness judgments, the CPP for the test stimuli is more similar to the ideal CPP displayed for the first dimension, indicating that the luminance of the background contributed more strongly to the choices in comparison with the luminance of the inner contour. Deviations from the ideal patterns, however, indicate contributions from both dimensions for both tasks.

In these experiments, observers were instructed to judge the appearance of the interior region of the stimulus. However, it is possible that observers attended to the experimental dimensions (e.g., the continuity of the color of the contour) instead of the appearance of the interior region. If this were the case, we should obtain the same pattern of responses between the test and the control conditions. However, the test CPP patterns are different, with the control CPP patterns showing little, if any, systematic structure.

 Figure 3 about here

Statistical comparisons of additive and independence models based on a

nested likelihood ratio test are shown in Table I for judgments based on hue and

in Table II for judgments based on brightness. In these tables, each row

indicates the test for an observer, and the last column corresponds to the

probability that the additive model fits no better than the independence model.

The degrees of freedom are obtained from the difference of the number of

coefficients estimates in each model (8 (additive) − 4 (independence) = 4).

Comparing the independence with the additive model indicates that the

independence model can be rejected for the test stimuli for both tasks and for all

observers ($p < 0.001$). The motivation for testing the nested decision models for

the control stimuli is less clear. Instead, we used a linear mixed-effects model to

test if the estimated perceptual scale values depended significantly on the

stimulus level with a random observer intercept (Bates et al., 2015). For both

the hue and the brightness judgments, no significant dependence was found

(hue: $\chi^2(30) = 38.5, p = 0.14$; brightness: $\chi^2(30) = 22.2, p = 0.85$). It could

be argued that 4 observers is not sufficient to estimate the variances accurately

in a mixed-effect model. We also performed the tests using a linear model with

Observer entering as a fixed-effect, with no significance obtained in either case.

The advantage of the mixed-effects model is that the results generalize to the

population rather than just the sample of 4 observers tested (Knoblauch &

Maloney, 2012; Moscatelli et al., 2013; Pinheiro & Bates, 2000).

Tables I and II about here

Figure 4 shows the estimated contributions of each dimension obtained

from fitting the additive model for each of the tasks. For judgments based on

hue, the average estimated scales for each pair of inner contour luminance and

background luminance elevation are shown in Figure 4 (a) for 4 observers. In

this figure, the column labels indicate the observer identification. The top row

shows the scale values estimated for the test stimulus and the bottom for the

control stimulus. Black circles indicate the inner contour contribution and white

the background contribution to the judgments. The values on the abscissa

indicate the five stimulus levels for each dimension coded from 1 to 5. These

values are indices to the 5 luminance elevations of the inner contour and the

background used in the experiment.

The results in Figure 4a indicate that the contribution of the luminance of

the interior contour dimension to the hue filling-in strength of the inner contour

as does the contribution of the background increases with luminance elevation

but with a smaller effect. The second row shows results obtained for the control

stimuli. Here, there appears no systematic influence of either dimension on the

strength of the coloration effect, because both dimension contributions are close

to zero at all stimulus levels. This result further supports that the observers

based their judgments on the perceived filled in color of the interior rather than

the luminance of the inner contour or the background.

Figure 4 about here

Figure 4 (b) shows the average estimated scales for each observer for each

pairing of inner contour and background luminance elevations when judgments

were based on the brightness of the interior region. The information in the

figure is organized in the same fashion as Figure 4a. The top row shows the

additive model fits to judgments of the brightness of the interior region. As the

inner contour increases in luminance, the estimated contribution for this

component decreases, indicating that the brightness of the interior region

decreases, i.e., generating a relative contrast rather than an assimilative effect.

The background dimension contributes positively with the luminance elevation

to judgments of brightness. These results contrast with the scales estimated for

the control stimuli that show no difference and little effect of both dimensions

on the strength of brightness filling-in.

Temporal forced choice experiments can be subject to order effects

(Yeshurun et al., 2008). To test for this, we compared the scales obtained for

trials in which the higher luminance was presented in the first interval with the

trials in which it was presented in the second interval. We performed the same

analysis with respect to the luminance of the inner contour. For the hue

judgments, the average estimated scales for each background luminance order

are shown in Figure 5 (a) for 4 observers, and judgments based on brightness are

displayed in Figure 5 (b). In these figures, the column labels indicate the

observer identification. The top row shows the estimated scale values for the

contribution of the inner contour luminance and the bottom for the background

dimension. White circles indicate the dimension contribution to the judgments

when the luminance elevation of the background is higher in the first stimulus

presentation and black circles the contribution to the judgments when the

luminance elevation of the background is higher in the second presentation. The

values on the abscissa indicate the five levels of each dimension coded from 1 to

5. For judgments based on hue, the results in Figure 5a indicate that the

contribution of the inner contour dimension is not different between both

background presentation orders. Moreover, the background dimension is

approximately the same between the presentation orders. Similar results were

obtained when judgments were based on brightness (Figure 5b). This shows

that any order effect based on the luminance background is very slight or absent

in our experiments.

For completeness, we show in Figure 6 the effect of the presentation order

of the luminance of the inner contour on the estimated scales for both judgments

organized in the same fashion as Figure 5. For both tasks, the scales for neither

dimension were influenced as a function of the presentation order.

425  Comparing the saturated and additive models with nested likelihood ratio

426 tests rejected the hypothesis that the saturated model provided a fit no better

427 then the additive model, thus, demonstrating that an interaction term is required

428 to describe the observers' judgments for both tasks (Tables III and IV). The

429 degrees of freedom indicate the difference of the number of coefficents

430 estimated in the 2 models (24 (saturated) – 8 (additive) = 16). The additive

431 model fit was rejected in all 8 tests. The estimated coefficients for the saturated

432 model are shown for each observer and both tasks in the panels of Figure 7. In

433 these displays, the estimated scale values are plotted as a function of the

434 stimulus index for the luminance of the inner contour with the index of the

435 background luminance specified as a parameter for each curve. The results for

436 the hue judgments are presented on the top row with the brightness judgments

437 on the bottom. The averages of the four observers for both judgment conditions

438 are summarized as three-dimensional surfaces in Figures 8a and b. If the

439 additive model was a good fit to the data, the curves for different background

440 levels would be parallel in Figure 7. However, for both the hue and brightness

441 responses, as the background luminance increases, the curves fan out. For the

442 hue judgments, this results in a larger range of hue responses to the range of

443 inner contour luminances tested for the higher than lower luminance

backgrounds. For the brightness judgments, in contrast, the range of response decreases at the highest luminance background. Thus, the background luminance produces different effects on both the type of filling-in (assimilation vs contrast) for hue and brightness judgments and on the dynamic range of the response, expanding it for hue judgments but compressing it for brightness.

Figures 7 and 8 about here

Tables III and Table IV about here

4. Discussion

In the present study, MLCM was used to quantify the contributions to the filling-in strength of the WCE of two stimulus dimensions: the background and inner contour luminances. We quantified how changes in these features affected perceived filling-in using two separate response criteria, linked to the strength of the hue (the conventional WCE) and the brightness of the interior region. Control experiments using a stimulus contour that generated statistically undetectable filling-in confirmed that observers judged the perceived attributes of the interior regions and not changes in the background and contour luminances, per se. We also found that the results were largely independent of the ordering of presentation in a trial of both the backgrounds and the inner contours.

As found previously, the strength of the coloration effect depends on the

luminance of the inner contour (Devinck & Knoblauch, 2012; Devinck et al.,

2014; Gerardin et al., 2014). Here, we show that the stimulus configuration

inducing the WCE generates both a hue and also a brightness filling-in of the

interior area and that these two phenomena are differentially affected by the

stimulus dimensions that we manipulated. The hue filling-in effect was

assimilative, and the hue became more saturated with increases in luminance of

the inner contour and the background. However, luminance of the interior

contour generated a contrast effect for brightness, in that the judged brightness

of the interior region decreased at higher contour luminances. As observers

compared the interior regions of the two successively displayed stimuli and

judged which central region appeared brighter, it is still possible that the

perceived filling-in was assimilative, i.e., the same contrast polarity as the inner

contour and that the effect of the contour luminance was simply to reduce the

lightness. This is difficult to assess since we cannot simply compare the interior

with the surround because both the surround and the interior vary in each

condition. Additionally, if we hold the surround constant, then we cannot rule

out its effect on the interior.

The hue effect

Both the hue and the brightness of the interior region were judged to be

greater with increases in the luminance elevation of the background. Pinna et al.

(2001) have previously reported that color spreading in the WCE occurs not

only with a white background but also with grey backgrounds and that even a faint spreading is perceived with dark backgrounds. Our results agree with these observations in that the background contribution to the hue judgments increased at higher luminances.

We also found that hue filling-in was more pronounced for bright than dim backgrounds. Previous studies have reported that color saturation diminishes when the brightness contrast between a colored object and its luminance background increases, a phenomenon named the gamut expansion effect (Brown & McLeod, 1997) subsequently confirmed and quantified by several investigators (Bimler et al., 2009; Faul et al., 2008; Xing et al., 2015). In contrast with these studies, our experiment shows that increasing the background luminance strengthens the assimilation hue, suggesting that we are observing a different phenomenon.

The brightness effect

The WCE has typically been described as a coloration effect. Brightness variations in the interior region of the stimulus have not typically been systematically quantified. One exception to this concerns studies that used an achromatic stimulus configuration (Cao et al., 2011; Coia & Crognale, 2017). In Coia and Crognale (2017), observers compared the filling-in region in the WCE to a reference stimulus with a physical luminance difference. Their matching results indicated that the test field shifted in the opposite direction

from the inner contour showing an assimilation effect. In Cao et al. (2011), the

luminances of the inner contour and the luminance of the background were fixed

while the outer contour varied between high and low luminance levels, thus

varying the contrast. Observers were asked to report which of two interior

surface stimuli appeared darker. Their results followed a U-shape with the

strength of the effect maximized for a range of medium luminance levels but not

for the extreme luminance levels. Consequently, the luminance contrast

between both contours affects the WCE but not linearly. An intriguing point is

the authors' description of their results: "Although there is an apparent

assimilation effect in the chromatic WCE, it is hard to tell whether it is actually

assimilation or some type of contrast effect happening here for the 'opposite

polarity' condition". As their method based the estimation of the WCE strength

on the probability of discriminating which stimulus appeared darker, we can

assume that if responses are inferior to 50%, then the surface appears lighter but

not equal (which would be the case for responses equal to 50%), indicating a

contrast rather than an assimilation effect. For our experiment based on

brightness judgment, when the inner contour increases in luminance, the

estimated contribution for this component decreases, indicating that the

brightness of the interior region decreases, also indicating a contrast

phenomenon. Thus, the two sets of results may be in agreement. Nevertheless,

the nature of the brightness effect could be influenced by the presence of the

chromatic component in our stimulus situation.

In Devinck et al. (2005), observers were allowed to adjust both the luminance and the chromaticity of a field in order to match the color of the WCE. The mean luminance match was near the luminance of the background, indicating that little or no luminance adjustment was required to make a perceptual match. The authors concluded that the WCE is predominantly a chromatic effect as originally suggested by Pinna et al. (2001). These results are not necessarily in conflict with the current study, in that we observed that the variation in brightness contrast with inner contour luminance is diminished at high backgrounds and might have been difficult to detect via matching. We would predict, then, that manipulating the luminance of the background would affect the luminance match, but this would require a more systematic study of the background than was performed by Devinck et al. (2005). Here, our data indicate that the perceptual effect is not limited only to a coloration phenomenon in that both the background and inner contour luminances influence observers' judgment of the brightness of the central surface. A simple hypothesis to account for the reduced brightness effect at high background luminances is to suppose that the background light added to the interior region cancels the contrast effect. This would not explain, however, the fact that higher background luminances led to a stronger hue percept.

Assimilation vs contrast

Whether one observes a contrast or assimilation effect may depend on the width of the contours (Fach & Sharpe 1986 ; Helson & Rohles, 1959; Helson, 1963). Other factors such as the luminance of the inducing stimuli can also influence our percept. Thus, de Weert and Spillmann (1995) indicated that assimilation or contrast occurred depending on whether the inducing contours of varied reflectance were darker or lighter than the gray background in using a pincushion pattern. A matching experiment for brightness judgements indicated that contrast occurred when the luminance level of the inducing contour was above the luminance level of the background and that assimilation occurred when the luminance of the inducing contour was below the luminance of the background. Given the different spatial dependencies of chromatic and luminance sensitive mechanisms, one might expect differences in the spatial domains over which chromatic and luminance components of a stimulus induce assimilation and contrast. Given the dependence of induction phenomena on stimulus configuration, however, (Fach & Sharpe 1986; de Weert & Spillmann 1995; Smith et al. 2001; Monnier & Shevell 2003, 2004), it is difficult to predict *a priori* whether the dimensions of the contours that we used should predict one or the other for the hue and brightness judgments.

In our study, observers compared the interior regions of the two successively displayed stimuli and judged which central region appeared brighter. Under this condition, our results indicated that the brightness of the

interior region decreases generating a relative contrast rather an assimilation

effect.   However, it still possible that different visual phenomena are perceived

in other circumstances. In making a brightness judgements of the central region

with respect to the outer region, we noted an assimilation phenomenon at the

lowest luminance level of the background and at the highest luminance level of

the inner contour. This condition corresponds to the lower left corner in Figure

2.

Unitary vs multiple mechanisms

The color of the central surface in the WCE is characterized by a spread

of color from the inner contour. Most previous studies of the WCE reported that

the coloration effect depends on both the chromatic and luminance contrasts of

the inner and outer contours. For example, most authors demonstrated that the

coloration effect increases with increasing luminance contrast between inner and

outer contours (Devinck & Knoblauch, 2012; Devinck et al., 2005).

Additionally, the coloration effect increases when the chromatic coordinates of

the inner and outer contours are approximately complementary in the color

diagram (Devinck et al., 2006). Thus, the main explanation assumes a filling-in

process in which a neuronal mechanism detects the contour and generalizes it

beyond the confines of the immediate stimulus. Most studies in the WCE have

reported an important role for several types of contour mechanisms generating a

long-range filling-in percept. Taken together, these data suggest that the filling-

in process involved in the WCE requires multiple levels of processing (Devinck

et al., 2014b; Pinna et al., 2001; Pinna & Grossberg, 2005; von der Heydt &

Pierson, 2006). The present results indicate that the mechanisms inducing the

brightness and coloration percept in the WCE are affected differently by the

luminance of the inner contour. These opposing responses due to the inner

contour suggest that multiple mechanisms contribute to the appearance of the

interior region. Different mechanisms could be activated or inhibited yielding to

color assimilation or brightness contrast effects, respectively. Future

experiments based on visual masking or contour adaptation could investigate

such phenomena.

Acknowledgements

609
610 This work was supported by a grant from the Agence Nationale de la Recherche

611 to FD (ANR-11-JSH-20021) and by LABEX CORTEX (ANR-11-LABX-0042).

References

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67(1), 1-48. doi:10.18637/jss.v067.i01.

Bimler, D.L., Paramei, G.V., & Izmailov, C.A. (2009) Hue and saturation shifts from spatially induced blackness. *Journal of the Optical Society of America A*, 26, 163–172.

Brown, R.O., & MacLeod, D.I.A. (1997). Color appearance depends on the variance of surround colors. *Current Biology*, 7, 844-849.

Brenner, E., & Cornelissen, F.W. (2002). The influence of chromatic and achromatic variability on chromatic induction and perceived colour. *Perception*, 31, 225-232.

Cao, B., Yazdanbakhsh, A., & Mingolla, E. (2011). The effect of contrast intensity and polarity in the achromatic watercolor effect. *Journal of Vision*, 11(3):18, 1–8.

Coia, A.J., & Crognale, M.A. (2017). Contour adaptation reduces the spreading of edge induced colors. *Vision Research*,

Derrington, A.M., & Krauskopf, J., & Lennie, P. (1984). Chromatic mechanisms in lateral geniculate nucleus of macaque. *The Journal of Physiology*, 357, 241-265.

Devinck, F., and Knoblauch, K. (2012). A Common Signal Detection Model for the Perception and Discrimination of the Watercolor Effect. *Journal of Vision*, 12, 1–14.

Devinck, F., and Spillmann, L. (2009). The watercolor effect: Spacing constraints. *Vision Research*, 49, 2911–2917.

Devinck, F., Gerardin, P., Dojat, M., and Knoblauch, K. (2014a). Spatial selectivity of the Watercolor Effect. *Journal of the Optical Society of America A*, 31, A1–A6.

Devinck, F., Gerardin, P., Dojat, M., and Knoblauch, K. (2014b). Quantifying the Watercolor effect: from stimulus properties to neural models. *Frontiers in Human Neurosciences*, 8:805.

638 Devinck, F., Delahunt, P.B., Hardy, J.L., Spillmann, L. and Werner, J.S. (2005). The

639    Watercolor effect: Quantitative evidence for luminance-dependent mechanisms of

640    long-range color assimilation. *Vision Research*, 45, 1413-1424.

641 Devinck, F., Hardy, J.L., Delahunt, P.B., Spillmann, L., & Werner, J.S. (2006). Illusory

642    spreading of Watercolor. *Journal of Vision*, 6, 625-633.

643 DeWeert, C.M.M., & Spillmann, L. (1995). Assimilation: asymmetry between brightness and

644    darkness? *Vision Research*, 35, 1413 - 1419.

645 Fach, C., & Sharpe, L. (1986). Assimilative hue shifts in color gratings depend on bar width.

646    *Perception & Psychophysics*, 40, 412 – 418.

647 Falmagne, J.-C. (1985). *Elements of psychophysical theory*. Oxford: Oxford University Press.

648 Faul, F., Ekroll, V., & Wendt, G. (2008) Color appearance: the limited role of chromatic

649    surround variance in the "gamut expansion effect." *Journal of Vision*, 8(3):30, 1-20.

650 Hansmann-Roth, S., & Mamassian, P. (2017). A glossy simultaneous contrast: Conjoint

651    measurement of gloss and lightness. *i-Perception*, 1-16.

652 Helson, H. (1960) Studies of anomalous contrast and assimilation. *Journal of the Optical*

653    *Society of America*, 53, 179-184.

654 Helson H. & Rohles F. (1959) A quantitative study of reversal of classical lightness-contrast ?

655    *The American Journal of Psychology*, 72, 530-538.

656 Ho, Y. X., Landy, M. S., & Maloney, L. T. (2008). Conjoint measurement of gloss and

657    surface texture. *Psychological Science*, 19, 196–204.

658 Kirschmann, A. (1890). Uber die quantitativen verhältnisse des simultanen helligkeits- und

659    farben-contrastes. *Philosophische Studien*, 6, 417-491.

660 Knoblauch, K., & Maloney, L. T. (2008). MLDS: Maximum likelihood difference scaling in

661    R. *Journal of Statistical Software*, 25, 1–26.

662 Knoblauch, K., & Maloney, L. T. (2012). *Modeling psychophysical data in R*. New York:

663    Springer.

664 Knoblauch, K., & Maloney, L. T. (2014). MLCM: Maximum Likelihood Conjoint

665  Measurement. R package version 0.4.1. https://CRAN.R-project.org/package=MLCM

666 Krantz, D. H., Luce, R. D., Suppes, P., & Tversky, A. (1971). Foundations of measurement

667  (vol. 1): Additive and polynomial representations. New York: Academic Press.

668 Krauskopf, J., Williams, D.R., & Heeley, D.W. (1982). Cardinal directions of color space.

669  *Vision Research*, 22, 1123-1131.

670 Lisi, M., & Gorea, A. (2016). Time constancy in human perception. *Journal of Vision*,

671  16(4):3, 1-12.

672 Luce, R. D., & Tukey, J. W. (1964). Simultaneous conjoint measurement: A new scale type of

673  fundamental measurement. *Journal of Mathematical Psychology*, 32, 466–473.

674 MacLeod, D.I., & Boynton, R.M. (1979). Chromaticity diagram showing cone excitation by

675  stimuli of equal luminance. *Journal of the Optical Society of America*, *69*, 1183-1186.

676 Maloney, L.T., & Yang, J.N. (2003). Maximum Likelihood Difference Scaling. *Journal of*

677  *Vision*, 3, 573–585.

678 Monnier, P., & Shevell, S.K. (2003). Large shifts in color appearance from patterned

679  chromatic backgrounds. *Nature Neuroscience*, 6, 801 - 802.

680 Monnier, P., & Shevell, S.K. (2004). Chromatic induction from S-cone patterns. *Vision*

681  *Research*, 44, 849 - 856.

682 Moscatelli, A., Mezzetti, M., & Lacquaniti, F. (2012). Modeling psychophysical data at the

683  population-level: The generalized linear mixed model. *Journal of Vision*, 12(11):26.

684 Pinheiro, J.C., & Bates, D.M. (2000). *Mixed-Effects Models in S and S-PLUS*. Springer, New

685  York.

686 Pinna, B. (1987). Un effeto di colorazione. In V. Majer & M. Santinello (Eds.), *Il laboratorio*

687  *e la citta`. XXI Congresso degli Psicologi Italiani* (p. 158). Edizioni SIPs, Milano:

688  Societa` Italiana di Psicologia.

Pinna, B., & Tanca, M. (2008). Perceptual organization reconsidered in the light of the

watercolor illusion: The problem of perception of holes and the object-hole effect.

Journal of Vision, 8, 1-15.

Pinna, B., Brelstaff, G. & Spillmann, L. (2001). Surface color from boundaries: a new

'watercolor' illusion. *Vision Research*, 41, 2669-2676.

Pinna, B., Werner, J.S. & Spillmann, L. (2003). The watercolor effect: a new principle of

grouping and figure–ground organization. *Vision Research*, 43, 43-52.

Qi, L., Chantler, M.J., Siebert, J.P., & Dong, J. (2015). The joint effect of mesoscale and

microscale roughness on perceived gloss. *Vision Research*, 115, Pt B:209-217.

R Core Team (2017). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Roberts, F. S. (1985). *Measurement theory*. Cambridge, UK: Cambridge University Press.

Rogers, M., Knoblauch, K., & Franklin, A. (2016). Maximum likelihood conjoint

measurement of lightness and chroma. *Journal of the Optical Society of America A*,

33, A184–A193.

Shevell, S. (1978). The dual role of chromatic background in color perception. *Vision

Research*, 18, 1649-1661.

Smith, V.C., Jin, P.Q., & Pokorny, J. (2001). The role of spatial frequency in color induction.

Vision Research, 41, 1007 - 1021.

Tanca, M., & Pinna, B. (2008). The phenomenal dissociation between coloration and object-

hole effects in the watercolor illusion. *Visual Neuroscience*, 25, 423-432.

von der Heydt, R., & Pierson, R. (2006). Dissociation of color and figure–ground effects in

the watercolor illusion. *Spatial Vision*, 19, 323–340.

Walraven, J. (1976). Discounting the background – the missing link in the explanation of

chromatic induction. *Vision Research*, 16, 289-295.

Wiebel, C.B., Aguilar, G., & Maertens, M. (2017). Maximum Likelihood Difference Scales

represent perceptual magnitudes and predict appearance matches. *Journal of Vision*,

17(4):1, 1-14.

Xing, D., Ouni, A., Chen, S., Sahmoud, H., Gordon, J., & Shapley, R. (2015). Brightness-

color interactions in human early visual cortex. *The Journal of Neuroscience*, 35,

2226-2232.

Yeshurun, Y., Carrasco, M., & Maloney, L.T. (2008). Bias and sensitivity in two-interval

forced choice procedures: Tests of the difference model. *Vision Research*, 48, 1837–

1851.

Zahn, C. T., & Roskies, R. Z. (1972). Fourier descriptors for plane close curves. *IEEE*

Transactions on Computers, C–21, 269–281.

 Figure Legends

733   Fig. 1: (a) Example of the Watercolor Effect. When a light orange contour is

734   surrounded by a dark purple contour, the enclosed area takes the tint of

735   the orange border. (b) Example of stimuli using Fourier descriptor as

736   test stimulus (presented on the left side) and using braided contour as

737   control stimulus (displayed on the right side).

739   Fig. 2: Examples of stimulus set used for a conjoint measurement experiment.

740   The figure indicates the set of stimuli used in both judgment tasks.

741   Each column corresponds to a different luminance elevation of the

742   background and each row to a different luminance elevation of the

743   inner orange contour.

745   Fig. 3: Conjoint proportion plots for judgments based on hue (a) and on

746   brightness (b). Each plot shows the proportion of stimulus S_{kl} judged to

747   have a greater filling-in than the stimulus represented in abscissa S_{ij} as

748   grey level according to the color bar on the right side. The luminance

749   elevation of the background is indicated by the large grids (i,k) and

750   each grid is subdivided into smaller 5×5 grid indicating the luminance

751   elevation of the inner contour (j,l). In each set of graphs, the top row

752   indicates the results for the test stimuli and the bottom for the control

stimuli for 4 observers. (c) Conjoint proportion plots for a simulated

observer. Expected responses for an observer who judges the stimuli

based on the contributions along only one of the dimensions.

Fig. 4: (a) Estimated scales for judgments based on hue. Additive model

average estimates for test (top row) and control stimuli (bottom row) as

a function of inner contour elevation (black circles) and luminance

background elevation (white circles) for four observers. Error bars

show 95% confidence intervals for estimates across the 5 runs. (b)

Results for judgments based on brightness. The solid lines indicate the

estimated contributions of each dimension under the additive model for

test (top row) and control patterns (bottom row) as a function of inner

contour elevation (black circles) and luminance background elevation

(white circles) for four observers. Error bars show 95% confidence

intervals for estimates across the 5 runs.

Fig. 5: (a) Results for judgments based on hue depending on the presentation

order for the luminance background of the test stimulus. The solid

lines indicate the estimated contributions under the additive model for

the inner contour dimension (top row) and the background dimension

(bottom row) as a function of the background order presentation. The

white circles are used to indicate that the luminance elevation of the

background is higher in the first interval than in the second interval and the black circles are used to represent a higher luminance elevation of the background in the second interval than in the first interval. Error bars are 95% confidence intervals based on a bootstrap procedure (Knoblauch & Maloney, 2012). (b) Results for judgments based on brightness depending of the luminance background presentation order for the test stimulus. Additive model average estimates for the inner contour dimension (top row) and the background dimension (bottom row) when the luminance elevation of the background is higher in the first interval than in the second interval (white circles) and when the luminance elevation of the background is higher in the second interval than in the first interval (black circles). Error bars are 95% confidence intervals based on a bootstrap procedure (Knoblauch & Maloney, 2012).

Fig. 6: Results for judgment based on hue (a) and on brightness (b) depending on the presentation order for the luminance contour of the test stimulus. The information in the figure is organized in the same fashion as Figure 5. The white circles are used to indicate that the luminance elevation of the inner contour is higher in the first interval than in the second interval and the black circles are used to represent a higher luminance

elevation of the inner contour in the second interval than in the first

interval. Error bars are 95% confidence intervals.

Fig. 7:   Results of the estimated contributions for each combination of the two

dimensions under the saturated model for four observers. The different

lines are used to code the index of the background dimension (for

indices 1 to 5). The top row represents the estimated contribution when

judgment is based on hue and the bottom row is the estimated

contribution when judgment is based on brightness.

Fig. 8   Three-dimensional surfaces for the average of the data over observers

from Figure 7, showing the contributions of the inner contour and

background to a) the hue and b) the brightness judgments under the

saturated model.

(a)

(b)

Figure 1

Background Luminance

Inner Contour Luminance

Figure 2

(a)

(b)

(c)

Figure 3

(a)

(b)

Figure 4

(a)

(b)

Figure 5

(a)

(b)

Figure 6

Figure 7

Figure 8

Table I: Comparison of independence and additive models for judgements based on hue

	Test stimuli (additive model × independent model)		
	Df	Deviance	p-value
Obs. 1	4	270.22	< 0.001
Obs. 2	4	568.41	< 0.001
Obs. 3	4	309.60	< 0.001
Obs. 4	4	406.38	< 0.001

Table II: Comparison of independence and additive models for judgement based on brightness

	Test stimuli (additive model × independent model)		
	Df	Deviance	p-value
Obs. 1	4	275.434	< 0.001
Obs. 2	4	912.11	< 0.001
Obs. 3	4	400.82	< 0.001
Obs. 4	4	430.64	< 0.001

Table III: Comparison of additive and saturated models for judgement based on hue

	Test stimuli (additive model × saturated model)		
	Df	Deviance	p-value
Obs. 1	16	105.44	< 0.001
Obs. 2	16	142.82	< 0.001
Obs. 3	16	104.81	< 0.001
Obs. 4	16	108.14	< 0.001

Table IV: Comparison of additive and saturated models for judgement based on brightness

	Test stimuli (additive model × saturated model)		
	Df	Deviance	p-value
Obs. 1	16	56.154	< 0.001
Obs. 2	16	74.01	< 0.001
Obs. 3	16	32.83	< 0.01
Obs. 4	16	122.78	< 0.001