**TITLE: Habitat complexity mediates benthic predator-prey interactions in Chesapeake**

**Bay**

**AUTHORS AND ADDRESSES**: Cassandra N. Glaspie*[†], Rochelle D. Seitz

Virginia Institute of Marine Science, Department of Biological Sciences, Gloucester Point,

Virginia 23062, U.S.A.

*Corresponding author contact information: glaspiec@oregonstate.edu; phone: 541-737-6105

†Present address: Oregon State University, Department of Fisheries and Wildlife, 104 Nash Hall,

Corvallis, OR 97731, USA

**ABSTRACT**

In Chesapeake Bay, the soft-shell clam *Mya arenaria* (thin-shelled, deep-burrowing)

exhibits population declines when predators are active and persists at low densities. In contrast,

the hard clam *Mercenaria mercenaria* (armored, shallow-burrowing) has a stable population and

age distribution. We examined the potential for habitat and predators to drive densities and

distributions of bivalves in a field caging experiment (*Mya* only) and laboratory mesocosm

experiments (*Mya* and *Mercenaria*). In the field, clams exposed to predators experienced 76.3%

greater mortality as compared to caged individuals, and blue crabs were likely responsible for

most of the mortality of juvenile *Mya*. In mesocosm experiments, *Mya* had lower survival in

sand and seagrass than in shell hash or oyster shell habitats. However, crabs often missed one or

more prey in seagrass, shell, and oyster shell habitats, suggesting that more complex habitat such

as seagrass may serve as refuge for low densities of *Mya*. Predator search times and encounter

rates declined when prey were at low densities, likely due to the added cost of inefficient

foraging causing crabs to give up; however, this effect was more pronounced for *Mya* than for

*Mercenaria*. *Mercenaria* had higher survival than *Mya* in mesocosm experiments, likely because

predators feeding on *Mercenaria* spent less time foraging than those feeding on *Mya*. *Mya* may

retain a low-density refuge from predation even with the loss of structurally complex habitats,

though a loss of habitat refuge may result in clam densities that are not sustainable. A better

understanding of density-dependent predator-prey interactions is necessary to prevent loss of

food-web integrity and to conserve marine resources.

**KEY WORDS**: bivalve, seagrass, functional response, density-dependent predation, optimal

foraging

**INTRODUCTION**

Predators exhibit top-down control on communities, influencing the abundance, size structure, and distribution of prey by restricting their survival or activity in time and space [1–3]. Predators also influence community function by preying upon dominant species [4–6]. To understand the structure and function of a community, it is important to consider the impact of the predators. Prey populations experience the effects of predation differently depending on how abundant the prey species is and, for actively foraging predators, how quickly the predator can find and consume prey [7]. The degree to which a predator can reduce prey abundance is a function of the probability of encountering a prey item, and the probability that the prey item will be eaten, given that it has been encountered. Both factors depend on the characteristics of the prey, the predator, and other environmental factors [6].

Bivalve mollusks exhibit a number of morphological and behavioral characteristics to defend against predators. Armor and aggregation decrease rates of predation, allowing predators and prey to coexist in the same space. For example, the hard-shell clam *Mercenaria mercenaria* (hereafter, *Mercenaria*) exhibits armor that protects it from predation by blue crabs *Callinectes sapidus*; clams larger than 40 mm cannot be crushed and therefore coexist with crabs [8]. Other bivalves must avoid predators to survive; the shell of a soft-shell clam *Mya arenaria* (hereafter, *Mya*) is thin and has a permanent gape, indicating that for this species, armor is not an important mode of protecting against attack by predators [9]. To avoid predation, *M. arenaria* achieves a non-coexistence refuge by burrowing 25-30 cm deep in the sediment, out of range of foraging predators, which rarely consume clams buried deeper than 10 cm [10].

Habitat also plays an important role in predator defense strategies of marine bivalves.

Predators in habitats that are not complex have a greater effect on prey than those in complex

habitats [11,12]. Vegetated or shell habitat provides a refuge from predation for many prey

[12,13], and increased sediment grain size allows infaunal species to avoid predators more

effectively than in fine sediments [10,14,15]. Complex habitats make foraging inefficient, and as

the cost to forage becomes too high, predators may opt to conserve energy or forage elsewhere

[16,17].

The functional response is a way to quantify predator foraging efficiency [7]. A

predator's functional response is the relationship between the number of prey consumed per

predator and prey density [18]. Predators that search for prey exhibit a density-dependent

functional response, because the encounter rate depends on prey density. In a type II density-

dependent response, handling rate and attack rate remain constant as prey density increases [7].

Prey consumed per predator increases with increasing prey density, but the rate of increase

declines to an upper asymptote. The asymptote is reached when the predator becomes satiated

and spends less time foraging, or when the predator is limited by the amount of time it takes to

consume prey [7]. A type III sigmoidal density-dependent response occurs when a predator

becomes more active as prey density rises, which means attack rate is a function of prey density

[7]. Type II and type III functional responses are very different biologically, since type III

functional responses create a refuge for prey at low densities, which may result in prey

persistence over time, even if a population is driven to low abundance [7,19,20].

The main parameters in a functional response model are encounter rate and handling time

[7], both of which change as a function of prey mortality, prey behavior, and habitat type. For the

purposes of this study, the encounter rate was defined as the number of encounters with prey

divided by the amount of time a predator spends foraging, or actively looking for prey; and the

handling time was defined as the amount of time a predator spends manipulating or eating a prey

item. For armored bivalves, the consumption rate of their predators is determined more by

handling time than encounter rate; in this case, a type II functional response is more likely [14].

For burrowing, thin-shelled bivalves, encounter rate is more important than handling time for

their predators [2], which means that a density-dependent sigmoidal (type III) response is likely

[14]. The biological mechanism behind a type III response is that low encounter rates often lead

to low activity levels or predators emigrating from the area [21]. The functional response of a

predator-prey interaction can also be habitat specific. Reduced sediment penetrability [14] or

increased vegetative cover [22] may lead to decreased encounter rate, and this may change the

functional response by creating or strengthening a low-density refuge from predation.

In the Chesapeake Bay, two commercially valuable clam species, the soft-shell clam *Mya*

and the hard clam *Mercenaria* have very different population dynamics. *Mya* exists in the Bay at

low abundance except immediately after spring recruitment, and juveniles are nearly completely

consumed by predators each year [23] (Fig 1). *Mercenaria* is fairly abundant throughout the

year, and all size classes persist in the Bay in all seasons [24] (Fig 1). The different dynamics of

these species may be due to predator-prey dynamics, since the two species exhibit different

predator-avoidance strategies. Specifically, the persistence of *Mya* at low abundance may be due

to a low-density refuge, especially in complex habitats that prevent efficient foraging by the

species' main predators, the blue crab *Callinectes sapidus* [25,26] and the cownose ray

*Rhinoptera bonasus* [27].

Fig 1. Size frequency histograms of *Mercenaria mercenaria* (left) and *Mya arenaria* (right) in

lower Chesapeake Bay. Samples were collected in spring (a-b), summer (c-d), and fall (e-f) for

two years starting in fall 2011. Sizes expressed are biomass (g AFDW m$^{-2}$) for *Mercenaria* [24]

and shell length (mm) for *Mya* [23].

This study aims to examine the nature of blue crab-bivalve predator-prey interactions for

these two infaunal bivalves, including the role of structural refuge (in the form of complex

habitat) on these interactions, using both field and laboratory experiments.  In field caging

experiments, we hypothesized the following: 1) blue crabs and cownose rays are both sources of

mortality for *Mya* (evidenced as a significant difference in *Mya* survival among all caging

treatments); and 2) the presence of seagrass increases clam survival rates as compared to sand

and mud (for all plots without a complete cage). In laboratory mesocosm experiments, we

hypothesized the following: 1) predators on *Mya* exhibit a type III functional response and

predators on *Mercenaria* exhibit a type II functional response (evidenced as a significant species-

density interaction); 2) complex (as compared to unstructured) habitats increase the extent of the

low-density refuge for species using density as a refuge, which manifests as increased

proportional survival in complex habitats as compared to sand, but only for *Mya* (evidenced as a

significant species-habitat interaction); 3) *Mercenaria*'s armor leads to increased handling time

compared to *Mya* (evidenced as a significant main effect of species on handling time); 4) low

densities, complex habitat, and deep-burrowing prey result in decreased blue crab search time,

due to the added cost of inefficient foraging (evidenced as a 3-way interaction between species,

density, and habitat), and 5) there is a decreased encounter rate at low densities of *Mya* compared

to high densities (evidenced as a significant species-density interaction).

## MATERIALS AND METHODS

### Field caging experiment

A caging study was conducted in patchy seagrass, sand, and mud near-shore habitats (1.5-2 m depth mean high water) in May 2014 near the mouth of the York River, VA (between 37.258323, -76.428047 and 37.275197, -76.370150). Ten replicate 0.25 $m^2$ plots were randomly assigned one of three caging treatments: full cage, stockade, or uncaged. Full cages were constructed of 13-mm galvanized wire mesh with PVC frames (0.6 m height, 0.5 m width, 0.5 m length) and were sunk into the sediment approximately 10 cm and secured with PVC legs that sunk into the ground an additional 30-40 cm. Stockades were constructed by placing 8 10-ft PVC poles around the plot at 25-cm intervals. Stockades kept cownose rays out of the plots, while still allowing for crab and fish predation. Uncaged plots were marked with two PVC poles on the diagonals. There were n = 10 replicates of each cage type in each habitat.

Juvenile soft-shell clams (*Mya*) were collected from the York River and held in flow-through tanks until experimentation. Clams were marked with permanent marker and transplanted towards the center of the plot at densities of 12 clams per plot (48 $m^{-2}$) [28]. A cage was placed over all transplanted clams to allow them to acclimate overnight and achieve a stable burrowing depth [21], and acclimation cages were removed from stockade and uncaged treatments. After 5 d, the contents of all plots were collected to a depth of 40 cm using a suction sampler [20]. Remaining bivalves were counted and shell fragments were noted as evidence of crab predation. Partial cages were not used to control for caging artifacts due to the short nature of this study and the tendency for partial cages to attract blue crabs. Only one density was used

in this study due to the presence of wild *Mya* in the area, and the consequent logistical

difficulties associated with creating reliable densities.

Proportional survival data were box-cox transformed ($\lambda = 0.51$) to achieve normality and

homogeneous variance (assessed using quantile-quantile and residual plots), and analyzed using

two-way ANOVA, with cage type (3 levels: full cage, stockade, and uncaged) and habitat (3

levels: sand, mud, and seagrass) as fixed factors, with $\alpha = 0.05$ for main effects and $\alpha = 0.20$ for

interaction terms [29]. Post-hoc pairwise comparisons were done using Tukey honest significant

difference (HSD) tests. From a pilot caging experiment in 2012, we used a simulation of

resampled data to determine that our sample size of $n = 10$ resulted in the following estimates of

statistical power: 1.00 for the main effect of cage type, 0.42 for the main effect of habitat, and

0.87 for the interaction effect.

**Laboratory mesocosm experiment**

*Mya* (thin-shelled deep infaunal) and *Mercenaria* (armored shallow infaunal) were

exposed to blue crab *C. sapidus* predation in mesocosm tanks of 0.87 m diameter and 0.59 m

height, which were partitioned with corrugated plastic to form a rectangular experimental arena

(40 cm x 70 cm). Sand was added to the tank to 25 cm depth, and an additional 25 cm of the tank

was filled with filtered water from the York River. Tank temperature was held constant at 26-27

°C with an aquarium heater and the water was aerated by air stone placed outside the

experimental arena. Trials were randomly assigned one of four substrate treatments: sand alone,

sand/shell hash, sand/oyster shell, or sand/seagrass. For trials receiving shell or oyster shell, a

constant volume of 0.5-L crushed shell hash (lightly crushed Baltic clam *Macoma balthica*,

ribbed mussel *Geukensia demissa*, and *Mercenaria* shell halves) or oyster shell halves was added

to the center of the mesocosm tank. Eelgrass (*Zostera marina*) and widgeongrass (*Ruppia*

*maritima*) shoots and rhizomes were collected from the York River and used to construct

seagrass mats for use in trials receiving seagrass. Seagrass mats were constructed with 0.5 liter of

natural seagrass blades tied onto plastic 1-cm Vexar mesh meant to simulate a rhizome mat.

Holes measuring approximately 25 cm$^2$ were cut approximately every 10 cm to allow crabs to

forage for clams buried under the simulated seagrass mat. The mesh and attached seagrass roots

were placed in the center of the tank and completely covered with sand.

Juvenile *Mya* were collected from the York River and held in flow-through tanks until

experimentation. Hard clams *Mercenaria* were obtained from Cherrystone Aqua-Farms in

Virginia. Only hard clams with shell lengths < 40 mm were used in the study, because blue crabs

are able to consume clams of this size [30]. Bivalves were placed in the sediment siphon up,

away from the edge of the tank to avoid edge effects, and allowed 24 h to achieve a stable burial

depth [21]. Each species was transplanted at two densities as determined from the literature, one

low and one medium density, which is sufficient to determine whether the functional response is

type II or III [21,31]. Low densities for both species were 4 clams per tank, and medium

densities were 11 clams per tank for *Mercenaria* and 16 clams per tank for *Mya* [16,31].

*Callinectes sapidus* were collected from the York River via baited crab pot. All crabs were

acclimated to the lab for 1 week or longer and fed fish or clam meat three times per week.

At the start of the experiment, one adult male blue crab with a carapace width ≥ 100 mm

was added to each tank receiving a predator treatment. Bivalves were exposed to blue crab

predation for 48 h, as is common for similar mesocosm studies [20].  Remaining bivalves were

excavated and counted upon termination of the experiment. There were six replicates of each

substrate/density combination, as well as an equal number of mesocosms set up without

predators, which served as controls (though only 0.6% of clams died in predator-free controls

and they are not analyzed or discussed further).

Proportional survival data were box-cox transformed ($\lambda = -0.14$) to achieve normality and

homogeneous variance (assessed using quantile-quantile and residual plots), and they were

analyzed using three-way ANOVA, with density (2 levels: low and medium), species (2 levels:

*Mya* and *Mercenaria*) and habitat (4 levels: sand, shell hash, oyster shell, and seagrass) as fixed

factors, with $\alpha = 0.05$ for main effects and $\alpha = 0.20$ for interaction terms [29]. Effect size and

standard error estimates from a previously conducted mesocosm experiment [21] were used to

calculate power to see a significant main effect of density, which was 0.95 for $n = 6$. Post-hoc

pairwise comparisons were done using Tukey HSD tests. Analysis of the number of mesocosm

trials with all of the clams eaten, a portion of the clams eaten, and none of the clams eaten was

completed using a chi-square test with Monte Carlo simulation of p values due to the presence of

zeroes in the contingency table.

It was not possible to use a different crab for each trial due to space requirements, nor

was it possible to use each crab the same number of times due to losses throughout the

experiment. Crabs were used between one and five times, and crabs were randomly assigned to

trials so there was no bias inherent in the re-use of crabs. An ANCOVA including individual

crab identity (51 levels), number of times a crab was used (continuous, 1-5), tank (4 levels), and

day of the experiment (continuous, standardized using z score transformation) as covariates

indicated that there was no difference in proportion of bivalves eaten based on crab identity ($F_{49,}$

$_{24} = 1.23$, $p = 0.30$), number of times the crabs were used ($F_{1, 24} = 1.56$, $p = 0.22$), tank ($F_{3, 24} =$

$0.48$, $p = 0.70$), or day of the experiment ($F_{1, 24} = 1.15$, $p = 0.29$). These results provided no

evidence that crabs exhibited learning behavior, and no evidence for tank effects or trends

through time; thus, each trial was treated as an independent replicate.

For half of the trials (n = 3 for each treatment) predator behavior was recorded using an

infrared-sensitive camera system. A red spotlight was used to improve night-time video quality

without disrupting crab behavior [32]. Videos were used to calculate search time, encounter rate,

and handling time. Search time (h) was defined as the total time spent exhibiting foraging

behavior, such as probing the sediment with legs or claws or lifting items to mouthparts.

Encounter rate $(hr^{-1})$ was defined as the number of encounters (picking up bivalve) divided by

the search time. Handling time (h) was defined as the total time spent manipulating or eating a

bivalve, divided by the number of encounters. Handling time, search time, and encounter rate

were fourth-root transformed to achieve homogeneity and compared for the two bivalve species

in different habitat treatments and at different densities using three-way ANOVAs with the same

factors as were used for analysis of proportional survival. Post-hoc pairwise comparisons were

done using Tukey HSD tests.

All analyses were completed using R statistical software [33], and data and R code files

are available in the Knowledge Network for Biocomplexity (KNB) repository [34].

**ETHICS STATEMENT**

Virginia Institute of Marine Science is statutorily-mandated as Virginia's scientific advisor on

marine- and coastal-related natural resources and exempt from having to obtain a scientific

collection permit for non-protected species in Virginia's waters.

**RESULTS**

**Field caging experiment**

Over the 5-day caging experiment, mean water temperature at the nearby YKTV2

weather buoy was 18.76 °C (± 1.63 SD). All replicates (n = 10) for the stockade and uncaged

plots lasted through the experiment and were subsequently sampled. At least one of the caged

plots was lost from each habitat, leaving n = 9 replicates in mud, n = 8 replicates in sand, and n =

8 replicates in seagrass.

As compared to full cages, there was a decrease in proportional survival of 75.6% in

stockades and 77.0% in uncaged plots (Fig 2), but the effect of one main effect depended on the

conditions of the other. Stockade and uncaged treatments had similar survival among habitats

(Table 1). Mud had significantly lower survival than sand or seagrass (Table 1). Seagrass and

sand had similar survival (Table 1). Due to a significant habitat x cage interaction, main effects

need to be interpreted with caution. The significant habitat x cage treatment interaction was

driven by the full cage treatment, which had different patterns of survival than the other caging

treatments (Supp. Table 1). Survival of clams in stockades placed in mud was lower than might

be expected with just main effects of substrate and cage type (Supp. Table 1).

Table 1. ANOVA summary table for field caging study proportional survival data.

Three types of caging treatments (full cage, stockade, and uncaged) were placed in three

substrate types (mud, sand, and seagrass); all were included in the ANOVA model as fixed

factors. Data were box-cox transformed ($\lambda$ = 0.51) prior to analysis. Significant p values (at $\alpha$ =

0.05 for main effects and $\alpha$ = 0.20 for interaction terms) are bolded.

| | Df | Sum Sq | Mean Sq | F value | Pr(>F) |
|---|---|---|---|---|---|
| *Substrate* | 2 | 2.65 | 1.32 | 10.35 | **0.0001** |
| *Cage* | 2 | 20.29 | 10.14 | 79.28 | **< 0.0001** |
| *Substrate x Cage* | 4 | 2.59 | 0.65 | 5.05 | **0.001** |
| *Residuals* | 75 | 9.60 | 0.13 | | |

Fig 2. Survival of transplanted juvenile *Mya arenaria* exposed to a natural suite of predators near

the mouth of the York River, VA. Shown are mean proportional survival (± 1 SE) after 5 d in the

field. Bivalves were placed in full cages (full), stockades, or uncaged plots. Plots were in

different habitats (denoted by different color bars). There were n = 10 replicates for the stockade

and uncaged plots, and n = 9, 7, and 8 replicates for cages in mud, sand, and seagrass,

respectively.

On average, 39.3% of missing clams were recovered as crushed shells within the plots.

Mean recovery of crushed shells varied little among caging types and habitats. The highest

occurred in stockade plots in sand, with 49.2% (± 28.7 SD) of missing clams recovered as

crushed shells, and lowest occurred in uncaged plots in mud, with 24.7% (± 26.5 SD) of missing

clams recovered as crushed shells.

**Laboratory mesocosm experiment**

In mesocosm experiments, mean proportional survival ranged from 0.27 (*Mya* in seagrass

at medium densities) to 1.00 (*Mercenaria* in seagrass at medium densities). Crabs ate at least one

*Mercenaria* in 18 out of 48 trials, and ate all offered *Mercenaria* in only one trial (low density in

shell). Predation of *Mya* was more common, with at least one *Mya* eaten in 27 out of 48 trials. In

the sand at low densities, crabs either ate all of the available *Mya* (occurred 3 times), or none of

them (occurred 3 times; Fig 3a). In the more-complex habitats (shell hash, oyster shell, and

seagrass), crabs offered low densities of clams usually ate none of them (occurred 13 out of 18

trials); only occasionally would a crab eat a portion (occurred 3 times) or all (occurred 2 times)

of the clams (Fig 3b, c, d). Frequency of occurrence of these events (all clams eaten, a portion of

the clams eaten, and no clams eaten) was significantly different between low and medium

bivalve densities for seagrass ($\chi^2 = 8.6$, p = 0.02), but not for sand ($\chi^2 = 6.0$, p = 0.13), shell ($\chi^2 =$

0.5, p = 1.00), or oyster ($\chi^2 = 3.3$, p = 0.24). There was no significant difference in the frequency

of occurrence of these events (all clams eaten, a portion of the clams eaten, and no clams eaten)

among habitat types ($\chi^2 = 5.3$, p = 0.57).

Fig 3. Foraging success for crabs feeding on clams in different habitats. Proportion of the

laboratory mesocosm trials in which all (black), some (partial, gray), or none (white) of the *Mya*

*arenaria* clams were eaten by crabs. Proportions are shown for low and medium densities of prey

in a) sand, b) shell, c) oyster, and d) seagrass.

*Mya* had significantly lower survival than *Mercenaria* (Fig 4; Table 2), but the effect of one

main effect depended on the conditions of the others. There was some evidence that bivalves had

lower proportional survival in trials with medium bivalve densities than in trials with low bivalve

densities (Table 2). There were no significant differences in survival by habitat type or bivalve

density (Table 2), but there were significant species x habitat interactions. *Mya* in medium

densities had lower survival than the other species x density combinations, driving a significant

species x density interaction (Supp. Table 2). In sand and seagrass, *Mya* had lower survival than

some other species x habitat combinations, driving a significant species x habitat interaction

(Supp. Table 3).

Table 2. ANOVA results for mesocosm study proportional survival of juvenile clams, as well as handling time (HT), search time (ST),

and encounter rate (ER) of blue crabs *Callinectes sapidus* feeding on juvenile clams. Two species (*Mya arenaria* and *Mercenaria*

*mercenaria*) were offered to blue crabs *Callinectes sapidus* at two densities (low and medium) in tanks with four different habitats

(sand, sand with shell hash, sand with oyster shell halves, and sand with live seagrass); all were included in the ANOVA model as

fixed factors. Data were box-cox transformed ($\lambda = -0.14$; survival only) or fourth-root transformed (HT, ST, and ER) prior to analysis.

Significant p values (at $\alpha = 0.05$ for main effects and $\alpha = 0.20$ for interaction terms) are bolded.

| | *Survival* | *HT* | *ST* | *ER* |
|---|---|---|---|---|
| *Species* | $\mathbf{F_{1,80} = 15.90, p = 0.0001}$ | $F_{1,32} = 2.87, p = 0.10$ | $F_{1,32} = 0.69, p = 0.41$ | $F_{1,32} = 0.07, p = 0.79$ |
| *Density* | $F_{1,80} = 3.68, p = 0.06$ | $\mathbf{F_{1,32} = 4.28, p = 0.05}$ | $\mathbf{F_{1,32} = 10.10, p = 0.003}$ | $\mathbf{F_{1,32} = 6.46, p = 0.02}$ |
| *Habitat* | $F_{3,80} = 1.86, p = 0.14$ | $F_{3,32} = 1.23, p = 0.32$ | $F_{3,32} = 0.31, p = 0.82$ | $F_{3,32} = 1.19, p = 0.33$ |
| *Species x Density* | $\mathbf{F_{1,80} = 7.17, p = 0.01}$ | $F_{1,32} = 0.03, p = 0.88$ | $\mathbf{F_{1,32} = 11.38, p = 0.002}$ | $F_{1,32} = 0.95, p = 0.34$ |
| *Species x Habitat* | $\mathbf{F_{3,80} = 2.19, p = 0.10}$ | $\mathbf{F_{3,32} = 2.01, p = 0.13}$ | $F_{3,32} = 1.13, p = 0.35$ | $F_{3,32} = 0.65, p = 0.59$ |
| *Density x Habitat* | $F_{3,80} = 0.65, p = 0.58$ | $F_{3,32} = 0.91, p = 0.45$ | $F_{3,32} = 1.47, p = 0.24$ | $F_{3,32} = 1.27, p = 0.30$ |
| *Species x Density x Habitat* | $F_{3,80} = 0.62, p = 0.61$ | $F_{3,32} = 0.25, p = 0.86$ | $\mathbf{F_{3,32} = 2.08, p = 0.12}$ | $F_{3,32} = 0.54, p = 0.66$ |

Fig 4. Density-dependent predation in different habitats. Mean juvenile *Mya arenaria* and

*Mercenaria mercenaria* proportional survival (± 1 SE) in mesocosms when exposed to blue crab

predation in a) sand, b) shell hash, c) oyster shell, and d) seagrass. Solid black lines are mean

proportional survival for *Mya* at two initial densities of 4 and 16 per tank, and dashed black lines

are mean proportional survival for *Mercenaria* at two initial densities of 4 and 11 per tank.

Handling time was significantly lower in low-density trials than in medium-density trials

(Fig 5a, b; Table 2), but the effect of one main effect depended on the conditions of the others.

The two treatments with the longest mean handling times were *Mercenaria* at medium density in

shell hash (1.31 h) and *Mercenaria* at medium density in sand (0.76 h). All other treatments had

mean handling times of 0.30 h or less. The overall mean handling times for *Mercenaria* and *Mya*

were 0.18 h and 0.03 h, respectively. In shell hash, *Mercenaria* had longer handling times than

the rest of the species x habitat combinations, driving a significant species x habitat interaction

(Supp. Table 4).

Fig 5. Behavior of blue crab *Callinectes sapidus* feeding on juvenile *Mya arenaria* and

*Mercenaria mercenaria*. Shown are means (± 1 SE) of a) handling time (HT) for crabs feeding

on *Mya*, b) HT for crabs feeding on *Mercenaria*, c) search time (ST) for crabs feeding on *Mya*,

d) ST for crabs feeding on *Mercenaria*, e) encounter rate (ER) for crabs feeding on *Mya*, and f)

ER for crabs feeding on *Mercenaria*. Lines of different colors and patterns represent different

habitat types (shell = shell hash; oyster = oyster shell), and means were calculated from n = 3

trials.

Search time was shorter in low-density trials than in medium-density trials (Fig 5c, d;

Table 2), but the effect of one main effect depended on the conditions of the others. The two

treatments with the longest mean search times were *Mya* at medium density in seagrass (5.67 h)

and *Mya* at medium density in oyster shell (5.56 h). The overall mean search times for

*Mercenaria* at low and medium densities were 1.22 h and 1.91 h, respectively. The overall mean

search times for *Mya* at low and medium densities were 0.89 h and 4.16 h, respectively. *Mya* at

medium densities had longer search times than the other species x density combinations, driving

a significant species x density interaction (Supp. Table 5). However, relatively long search times

for medium densities of *Mya* only occurred in certain habitats (sand, oyster shell, and seagrass),

resulting in a three-way interaction (Supp. Table 6).

Encounter rate was significantly lower in low-density trials than in medium-density trials

(Fig 5e, f; Table 2). The two treatments with the highest mean encounter rates were *Mya* at

medium density in sand (4.08 ind. $h^{-1}$) and *Mya* at medium density in seagrass (3.23 ind. $h^{-1}$).

The overall mean encounter rates for *Mercenaria* at low and medium densities were 0.79 ind. $h^{-1}$

and 1.80 ind. $h^{-1}$, respectively. The overall mean encounter rates for *Mya* at low and medium

densities were 0.81 ind. $h^{-1}$ and 2.85 ind. $h^{-1}$, respectively.

**DISCUSSION**

Blue crabs were the main predators of *Mya* in all habitats we examined, with no

significant difference between stockades and uncaged plots and high incidence of crushed shells,

which served as evidence of crab predation rather than fish predation. This was in line with our

hypothesis that crab predation would be important. Despite evidence in the literature that

schooling rays can result in mass mortality of bivalves [35], and evidence from gut content

analysis that cownose rays consume Mya [27], we did not observe evidence that cownose rays

increased predation in uncaged plots relative to stockade plots during the time frame of our field

experiment (May). These results were contrary to our hypothesis and indicate that over the time

and spatial scale of this study, rays were not a major source of mortality for *Mya*.

Predation-related mortality was high for juvenile *Mya* that were not protected by a cage.

Over a period of five days, exposure to predators decreased survival of juvenile *Mya* by 76.3% as

compared to caged individuals. Clam survival was habitat dependent, and both sand and seagrass

provided more refuge from predation than mud. *Mya arenaria* has previously been shown to

achieve a low-density refuge in sand [14,21]; however, our results went against our hypothesis

that the added complexity afforded by seagrass habitats provides an extended refuge for juvenile

*Mya*.

In the laboratory study, there was an effect of habitat on predator-related mortality only

for *Mya*, which had lower survival in sand and seagrass than in shell hash or oyster shell habitats.

However, in the case of a prey species that relies on achieving a low-density refuge for

persistence, proportional survival may not be the best measure of success. Shell hash, oyster

shell, and seagrass habitats had higher occurrence of trials with at least one clam remaining,

which may be biologically significant. Habitat that allows survival of one or a few clams may

maintain the low-density refuge for *Mya*.

Predators on *Mercenaria* (armored infaunal) and *Mya* (thin-shelled infaunal) had

significantly different functional responses. Predators on *Mya* had a type III sigmoidal functional

response, with a negative relationship between density and proportional survival, as has been

seen in previous studies [14]. Predators on *Mercenaria* had a type II hyperbolic functional

response, as has been seen previously [16], exhibiting either a positive relationship between

density and proportional mortality or no density dependence, depending on the habitat. This

difference is relevant to population dynamics and persistence of these two bivalve species

because a type II functional response is unstable and can lead to local extinction of prey if they

are driven to low densities, but a type III functional response may lead to prey persistence at low

density [7,36]. The type II functional response of predators feeding on *Mercenaria* means this

bivalve species must remain at relatively high densities to achieve population stability.

Conversely, the type III functional response of predators feeding on *Mya* allows the species to

persist, even at very low density,

The differences in functional response of predators feeding on *Mya* and *Mercenaria* were

likely due to differences in predator behavior. Predators had shorter search time and encounter

rate when prey were in low densities as compared to high densities, in agreement with our

hypotheses, as predators appeared to give up foraging. At low densities, encounter rate did not

differ between the two bivalve species, indicating blue crabs had less trouble finding deep-

burrowing clams than we hypothesized. There was no evidence that blue crabs spent less time

foraging in complex habitats or when exposed to deep-burrowing prey; on the contrary, blue

crabs spent more time searching for *Mya* at medium densities than they did searching for

*Mercenaria* at medium densities, indicating crabs may have a preference for *Mya* as prey. This

tendency of blue crabs to pass up on *Mercenaria* as prey may explain why handling times for

*Mercenaria* were not significantly greater than handling times for *Mya*; while some crabs spent

the extra time opening up the heavily armored clams (*Mercenaria*), many predators also gave up

without investing much time into the encounter.

Seagrass did not provide a refuge from predation for *Mya* in the field or in the laboratory

experiment. However, seagrass in both studies was patchy, mesocosms were small, and caging

sites were chosen so that the three habitat types (mud, sand, and seagrass) were in close

proximity. Fragmented seagrass may not be able to provide much protection from generalist

predators such as blue crabs, especially if they feed efficiently at patch edges [37]. Despite little

evidence for patchy seagrass as a refuge from predation from this study, *Mya* are more likely to

be found in seagrass than all other habitat types in the lower Chesapeake Bay [23]. This indicates

that dense, contiguous seagrass stands may still provide a refuge from predation for *Mya*. Future

research examining the effect of seagrass density or patch size on the survival of juvenile *Mya* is

warranted.

Declines in complex habitat will likely lead to declines in thin-shelled species such as

*Mya.* Oyster shell and shell hash provided juvenile *Mya* some protection from predation in

mesocosm trials; however, in the lower Chesapeake Bay, thick layers of shell are uncommon.

Loss of many bivalves in the Bay, including oysters [38,39] and large-bodied clams [23,40,41],

will make hard-bottom shell-hash habitat even more rare in the future. Seagrass is also

experiencing declines in the Chesapeake Bay [42], resulting in a decrease of many potential

sources of highly complex benthic habitat in the Bay and a subsequent decrease in refuge for

thin-shelled clams. *Mya* may retain a low-density refuge from predation even with the loss of

structurally complex habitats, though a loss of habitat-mediated refuge may eventually result in

clam densities that are not sustainable.

Loss of complex habitat in the Chesapeake Bay may have little impact on armored

species such as *Mercenaria*. We did not see an effect of habitat on *Mercenaria* survival in the

current study, yet in previous research, *Mercenaria* had higher survival in crushed oyster shell

habitats than in sand or mud [30]. This inconsistency is likely due to the use of larger clams in

the current study (~30 mm shell length) as compared to the previous study, which used clams 5-

10 mm shell length [30]. Ontogenetic shifts in functional response may drive spatial distributions

of hard-shelled bivalves in Chesapeake Bay, which are most dense in oyster shell habitats [43].

However, the effect of habitat on survival of recruits does not appear to impact population

dynamics of *Mercenaria*, which were present in multiple size classes throughout the year in

lower Chesapeake Bay. Future research should examine whether complex habitat reduces blue

crab encounter rates with small (< 10 mm) *Mercenaria* to determine the relationship between this

species and complex habitat over its entire ontogeny.

**Relevance for conservation**

Understanding the mechanism underlying bivalve refuges from predation is important in

a changing world. Loss of structured habitat such as seagrass, mangroves, coral reefs, and oysters

is occurring world-wide [44]. There is a current research need for models that can be used to

forecast the impacts of global change, such as habitat loss, on predator-prey interactions [45].

We demonstrated that understanding the effect of habitat loss on predator-prey interactions is

improved by understanding the mechanisms prey use to defend themselves against predators and

the effects of prey density.

Nonlinear predator-prey dynamics can result in catastrophic changes and regime shifts

[46,47]. An examination of the functional response is key in predicting the result of predator-

prey interactions over time, and determining if a population crash can be expected in a food web,

potentially leading to a regime shift. For instance, functional responses will be a major factor in

determining whether a species driven to low abundance is likely to become locally extinct, or if

it is likely to persist [19]. Documenting the functional response of bivalve species with a variety

of different physical characteristics can help ecosystem managers decide on which species to

focus conservation efforts, since species with a type II functional response are at higher risk of

local extinction [48,49], and populations exhibiting a type III functional response are generally

more stable over time [21,50,51].

A better understanding of density-dependent predator-prey interactions can be used to

inform a variety of ecosystem management decisions. For example, functional responses can be

used to determine a threshold density for reintroduction of endangered or depleted species [52],

stock enhancement, [12,13], and pest control [53,54]. Effective bivalve seeding efforts that take

into account predation may help restore marine bivalves, many of which have experienced severe

declines in the recent past [38,39,55,56]. A better understanding of density-dependent predator-

prey interactions will assist in the effort to maintain the integrity of marine trophic interactions

and the viability of marine resources.

**ACKNOWLEDGMENTS**

We gratefully acknowledge the assistance given by the students and staff of the

Community Ecology and Marine Conservation Biology labs at the Virginia Institute of Marine

Science. This paper is contribution number XXXX from the Virginia Institute of Marine Science,

College of William & Mary.

**REFERENCES**

1.  Garrity SD, Levings SC. A predator-prey interaction between two physically and biologically constrained tropical rocky shore gastropods: Direct, indirect and community effects. Ecol Monogr. 1981;51: 267–286. doi:10.2307/2937274

2.  Micheli F. Effects of predator foraging behavior on patterns of prey mortality in marine soft bottoms. Ecol Monogr. 1997;67: 203–224.

3.  Beal BF. Relative importance of predation and intraspecific competition in regulating growth and survival of juveniles of the soft-shell clam, *Mya arenaria* L., at several spatial scales. J Exp Mar Bio Ecol. 2006;336: 1–17. doi:10.1016/j.jembe.2006.04.006

4.  Randall JE. Overgrazing of algae by marine fishes. Ecology. 1961;42: 812.

5.  Dayton PK. Competition, disturbance, and community organization: The provision and subsequent utilization of space in a rocky intertidal community. Ecol Monogr. 1971;41: 351–389. doi:10.2307/1948498

6.  Lubchenco J, Gaines SD. A unified approach to marine plant-herbivore interactions. I. Populations and communities. Annu Rev Ecol Syst. 1981;12: 405–437.

7.  Hassell MP. The Dynamics of Arthropod Predator-Prey Systems. Princeton, New Jersey: Princeton University Press; 1978.

8.  Blundon JA, Kennedy VS. Mechanical and behavioral aspects of blue crab, *Callinectes sapidus* (Rathbun), predation on Chesapeake Bay bivalves. J Exp Mar Bio Ecol. 1982;65: 47–65.

9.  Vermeij GJ. Evolution and Escalation: An Ecological History of Life. Princeton, New Jersey: Princeton University Press; 1987.

10. Blundon JA, Kennedy VS. Refuges for infaunal bivalves from blue crab, *Callinectes*

*sapidus* (Rathbun), predation in Chesapeake Bay. J Exp Mar Bio Ecol. 1982;65: 67–81.

11.    Sih A, Crowley P, Mcpeek M, Petranka J, Strohmeier K. Predation, competition, and prey

communities: A review of field experiments. Annu Rev Ecol Syst. 1985;16: 269–311.

12.    Stoner AW. Habitat-mediated survival of newly settled red king crab in the presence of a

predatory fish: Role of habitat complexity and heterogeneity. J Exp Mar Bio Ecol.

Elsevier B.V.; 2009;382: 54–60. doi:10.1016/j.jembe.2009.10.003

13.    Long WC, Whitefleet-Smith L. Cannibalism in red king crab: Habitat, ontogeny, and the

predator functional response. J Exp Mar Bio Ecol. 2013;449: 142–148.

doi:10.1016/j.jembe.2013.09.004

14.    Seitz RD, Lipcius RN, Hines AH, Eggleston DB. Density-dependent predation, habitat

variation, and the persistence of marine bivalve prey. Ecology. 2001;82: 2435–2451.

15.    Quammen ML. Predation by shorebirds, fish, and crabs on invertebrates in intertidal

mudflats: An experimental test. Ecology. 1984;65: 529–537.

16.    Sponaugle S, Lawton P. Portunid crab predation on juvenile hard clams: Effects of

substrate type and prey density. Mar Ecol Prog Ser. 1990;67: 43–53.

doi:10.3354/meps067043

17.    Abrams PA. Functional responses of optimal foragers. Am Nat. 1982;120: 382–390.

18.    Solomon ME. The natural control of animal populations. J Anim Ecol. 1949;18: 1–35.

doi:10.2307/1578

19.    Hassell MP, May RM. Stability in insect host-parasite models. J Anim Ecol. 1973;42:

693–726. doi:10.2307/3133

20.    Eggleston DB, Lipcius RN, Hines AH. Density-dependent predation by blue crabs upon

infaunal clam species with contrasting distribution and abundance patterns. Mar Ecol Prog

Ser. 1992;85: 55–68. doi:10.3354/meps085055

21.   Lipcius RN, Hines AH. Variable functional responses of a marine predator in dissimilar

homogenous microhabitats. Ecology. 1986;67: 1361–1371.

22.   Lipcius RN, Eggleston DB, Miller DL, Luhrs TC. The habitat-survival function for

Carribean spiny lobster: An inverted size effect and non-linearity in mixed algal and

seagrass habitats. Mar Freshw Res. 1998;49: 807–816.

23.   Glaspie CN. Mya arenaria population and disease survey; 2017 [cited 2017 Nov 15].

Database: Knowledge Network for Biocomplexity. Available from

https://knb.ecoinformatics.org/#view/doi:10.5063/F1N29V34.

24.   Glaspie CN. Chesapeake Bay bivalve survey 2011-2013; 2017 [cited 2017 Nov 15].

Database: Knowledge Network for Biocomplexity. Available from

https://knb.ecoinformatics.org/#view/doi:10.5063/F11C1V1Q.

25.   Hines A, Haddon A, Wiechert L. Guild structure and foraging impact of blue crabs and

epibenthic fish in a sub-estuary of Chesapeake Bay. Mar Ecol Prog Ser. 1990;67: 105–

126. doi:10.3354/meps067105

26.   Lipcius RN, Eggleston DB, Heck KL, Seitz RD, van Monfrans J. Post-settlement

abundance, survival, and growth of postlarvae and young juvenile blue crabs in nursery

habitats. In: Kennedy VS, Cronin LE, editors. Biology and Management of the Blue Crab.

Baltimore, MD: University of Maryland Press; 2007. pp. 535–565.

27.   Fisher RA. Life history, trophic ecology, and prey handling by cownose ray, *Rhinoptera*

*bonasus*, from Chesapeake Bay. Gloucester Point, VA; 2010.

28.   Skilleter G. Refuges from predation and the persistence of estuarine clam populations.

Mar Ecol Prog Ser. 1994;109: 29–42. doi:10.3354/meps109029

29.  Underwood AJ. Experiments in Ecology: Their Logical Design and Interpretation Using

Analysis of Variance. Cambridge: Cambridge University Press; 1997.

30.  Arnold WS. The effects of prey size, predator size, and sediment composition on the rate

of predation of the blue crab, *Callinectes sapidus* Rathbun, on the hard clam, *Mercenaria*

*mercenaria* (Linné). J Exp Mar Bio Ecol. 1984;80: 207–219.

31.  Taylor DL, Eggleston DB. Effects of hypoxia on an estuarine predator-prey interaction:

Foraging behavior and mutual interference in the blue crab *Callinectes sapidus* and the

infaunal clam prey *Mya arenaria*. Mar Ecol Prog Ser. 2000;196: 221–237.

doi:10.3354/meps196221

32.  Cronin TW, Forward RB. The visual pigments of crabs 1. Spectral characteristics. J Comp

Physiol A. 1988;162: 463–478.

33.  R Core Team. R: A language and environment for statistical computing [Internet]. Vienna,

Austria: R Foundation for Statistical Computing; 2017. Available: http://www.r-

project.org/

34.  Glaspie CN. *Mya arenaria* and *Mercenaria mercenaria* predator-prey dynamics; 2017

[cited 2017 Nov 15]. Database: Knowledge Network for Biocomplexity. Available from

https://knb.ecoinformatics.org/#view/doi:10.5063/F1RV0KTP.

35.  Peterson CH, Fodrie FJ, Summerson HC, Powers SP. Site-specific and density-dependent

extinction of prey by schooling rays: generation of a population sink in top-quality habitat

for bay scallops. Oecologia. 2001;129: 349–356. doi:10.1007/s004420100742

36.  Murdoch WW, Oaten A. Predation and population stability. Adv Ecol Res. 1975;9: 1–131.

37.  Laurance WF, Yensen E. Predicting the impact of edge effects in fragmented habitats.

Biol Conserv. 1991;55: 77–92.

38.    Beck MW, Brumbaugh RD, Airoldi L, Carranza A, Coen LD, Crawford C, et al. Oyster

reefs at risk and recommendations for conservation, restoration, and management.

Bioscience. 2011;61: 107–116. doi:10.1525/bio.2011.61.2.5

39.    Rothschild BJ, Ault JS, Goulletquer P, Héral M. Decline of the Chesapeake Bay oyster

population: A century of habitat destruction and overfishing. Mar Ecol Prog Ser.

1994;111: 29–39. doi:10.3354/meps111029

40.    Dungan CF, Hamilton RM, Hudson KL, McCollough CB, Reece KS. Two epizootic

diseases in Chesapeake Bay commercial clams, *Mya arenaria* and *Tagelus plebeius*. Dis

Aquat Organ. 2002;50: 67–78.

41.    Homer ML, Dungan CF, Tarnowski ML. Assessment of Chesapeake Bay commercial

softshell clams *Mya arenaria* and *Tagelus plebeius*. Report to NOAA Chesapeake Bay

Fisheries Science Program, Award NA07NMF4570326; 2011.

42.    Orth RJ, Moore KA. Chesapeake Bay: An unprecedented decline in submerged aquatic

vegetation. Science (80- ). 1983;222: 51–53.

43.    Glaspie CN, Seitz RD. Role of habitat and predators in maintaining functional diversity of

estuarine bivalves. Mar Ecol Prog Ser. 2017;570: 113–125. doi:10.3354/meps12103

44.    Duarte CM, Dennison WC, Orth RJW, Carruthers TJB. The charisma of coastal

ecosystems: Addressing the imbalance. Estuaries and Coasts. 2008;31: 233–238.

doi:10.1007/s12237-008-9038-7

45.    Hunsicker ME, Ciannelli L, Bailey KM, Buckel JA, Wilson White J, Link JS, et al.

Functional responses and scaling in predator-prey interactions of marine fishes:

Contemporary issues and emerging concepts. Ecol Lett. 2011;14: 1288–1299.

doi:10.1111/j.1461-0248.2011.01696.x

46.    Sinclair ARE, Byrom AE. Understanding ecosystem dynamics for conservation of biota. J

602            Anim Ecol. 2006;75: 64–79. doi:10.1111/j.1365-2656.2006.01036.x

603    47.    Hughes TP, Bellwood DR, Folke C, Steneck RS, Wilson J. New paradigms for supporting

604            the resilience of marine ecosystems. Trends Ecol Evol. 2005;20: 380–386.

605            doi:10.1016/j.tree.2005.03.022

606    48.    Gascoigne JC, Lipcius RN. Allee effect driven by predation. J Appl Ecol. 2004;41: 801–

607            810. doi:10.1111/j.0021-8901.2004.00944.x

608    49.    Kramer AM, Drake JM. Experimental demonstration of population extinction due to a

609            predator-driven Allee effect. J Anim Ecol. 2010;79: 633–639. doi:10.1111/j.1365-

610            2656.2009.01657.x

611    50.    Bellmore JR, Baxter C V., Connolly PJ. Spatial complexity reduces interaction strengths

612            in the meta-food web of a river floodplain mosaic. Ecology. 2015;96: 274–283.

613            doi:10.1890/14-0733.1

614    51.    Uszko W, Diehl S, Pitsch N, Lengfellner K, Müller T. When is a type III functional

615            response stabilizing? Theory and practice of predicting plankton dynamics under

616            enrichment. Ecology. 2015;96: 3243–3256. doi:10.1890/15-0055.1

617    52.    Sinclair ARE, Pech RP, Dickman CR, Hik D, Mahon P, Newsome AE. Predicting effects

618            of predation on conservation of endangered prey. Conserv Biol. 1998;12: 564–575.

619    53.    Madadi H, Mohajeri Parizi E, Allahyari H, Enkegaard A. Assessment of the biological

620            control capability of *Hippodamia variegata* (Col.: Coccinellidae) using functional

621            response experiments. J Pest Sci (2004). 2011;84: 447–455. doi:10.1007/s10340-011-

622            0387-9

623    54.    Boukal DS, Sabelis MW, Berec L. How predator functional responses and Allee effects in

624   prey affect the paradox of enrichment and population collapses. Theor Popul Biol.

625   2007;72: 136–147. doi:10.1016/j.tpb.2006.12.003

626 55. Whetstone JM, Eversole AG. Effects of size and temperature on mud crab, *Panopeus*

627   *herbstii*, predation on hard clams, *Mercenaria mercenaria*. Estuaries. 1981;4: 153–156.

628 56. Beal BF, Kraus MG. Interactive effects of initial size, stocking density, and type of

629   predator deterrent netting on survival and growth of cultured juveniles of the soft-shell

630   clam, *Mya arenaria* L., in eastern Maine. Aquaculture. 2002;208: 81–111.

631   doi:10.1016/S0044-8486(01)00900-0

633 **SUPPORTING INFORMATION**

635 S1 Table. Summary of Tukey HSD results for the caging study interaction term between

636 substrate and cage type. For each pairwise comparison, 95% confidence intervals (CI) and

637 adjusted p values are presented. Data were box-cox transformed ($\lambda = 0.51$) prior to analysis and

638 are not back-transformed. Only interactions with significant p values at $\alpha = 0.20$ are shown.

640 S2 Table. Summary of Tukey HSD results for the mesocosm study proportional mortality

641 interaction term between species and density. For each pairwise comparison, 95% confidence

642 intervals (CI) and adjusted p values are presented. Data were box-cox transformed ($\lambda = -0.14$)

643 prior to analysis and are not back-transformed. Only interactions with significant p values at $\alpha =$

644 0.20 are shown.

S3 Table. Summary of Tukey HSD results for the mesocosm study bivalve proportional

mortality interaction term between species and habitat. For each pairwise comparison, 95%

confidence intervals (CI) and adjusted p values are presented. Data were box-cox transformed ($\lambda$

= -0.14) prior to analysis and are not back-transformed. Only interactions with significant p

values at $\alpha = 0.20$ are shown.

S4 Table. Summary of Tukey HSD results for the mesocosm study *Callinectes sapidus* handling

time interaction term between species and habitat. For each pairwise comparison, 95%

confidence intervals (CI) and adjusted p values are presented. Data were fourth-root transformed

prior to analysis and are not back-transformed. Only interactions with significant p values at $\alpha =$

0.20 are shown.

S5 Table. Summary of Tukey HSD results for the mesocosm study *Callinectes sapidus* search

time interaction term between species and density. For each pairwise comparison, 95%

confidence intervals (CI) and adjusted p values are presented. Data were fourth-root transformed

prior to analysis and are not back-transformed. Only interactions with significant p values at $\alpha =$

0.20 are shown.

S6 Table. Summary of Tukey HSD results for the mesocosm study *Callinectes sapidus* search

time interaction term between species, density, and habitat. For each pairwise comparison, 95%

confidence intervals (CI) and adjusted p values are presented. Data were fourth-root transformed

prior to analysis and are not back-transformed. Only interactions with significant p values at $\alpha =$

0.20 are shown.