1 **Empirical test of crab-clam predator-prey model predictions: storm-driven phase shift to a**
2 **low-density steady state**
3

4 **Authors: Cassandra N. Glaspie*†, Rochelle D. Seitz, Romuald N. Lipcius**
5
6 Affiliations: Virginia Institute of Marine Science, College of William & Mary, Gloucester Point,
7 Virginia 23062, U.S.A.
8
9 *Correspondence to: cglaspie1@lsu.edu
10
11 † Present address: Louisiana State University, Department of Oceanography and Coastal Sciences,
12 3195 Energy, Coast, and Environment Building, Baton Rouge, LA 70803, USA

14 **Keywords:** bivalve, *Mya arenaria*, *Callinectes sapidus*, blue crab, soft-shell clam, tropical storm
15 Agnes, alternate stable state, Lotka-Volterra, nonlinear dynamics, Chesapeake Bay

Abstract:

A dynamic systems approach can predict steady states in predator-prey interactions, but there are very few empirical tests of predictions from predator-prey models. Here, we examine the empirical evidence for the low-density steady state predicted by a Lotka-Volterra model of a crab-clam predator-prey system using data from long-term monitoring, a field survey, and a field experiment. We show that Tropical Storm Agnes in 1972 likely resulted in a phase shift to a low-density state for the soft-shell clam *Mya arenaria*, which was once a biomass dominant in Chesapeake Bay. This storm altered predator-prey dynamics between *M. arenaria* and the blue crab *Callinectes sapidus*, shifting from a system controlled from the bottom-up by prey resources, to a system controlled from the top-down by predation pressure on bivalves. Predator-prey models with these two species alone were capable of reproducing observations of clam densities and mortality rates, consistent with the idea that *C. sapidus* are a major driver of *M. arenaria* population dynamics. Over 40 y post-storm, *M. arenaria* densities hover near a low-density steady state predicted from the predator-prey model. Predation rates observed in the field were similar to modeled mortality rates. Predator-prey models can be used to predict dynamics of natural populations, including phase shifts and densities at steady states. The preponderance of multispecies interactions exhibiting nonlinear dynamics indicates that this may be a general phenomenon.

Significance statement:

Biotic interactions often contain nonlinearities that result in complex and unpredictable ecosystem responses, including phase shifts to alternative stable states. Predator-prey theory may be used to predict population phase shifts and prevent unwanted ecological surprises if these models are validated against empirical data, especially from field-based studies. We use a Lotka-Volterra predator-prey model to predict prey densities and predation rates in a natural environment. Agreement between model predictions and field data in this crab-clam predator-prey system indicates that relatively simple models can be used to predict phase shifts and identify alternative stable states. The approach described here can be adapted to a variety of predator-prey systems to predict the impact of phase shifts on ecological communities.

Introduction:

Predators play a key role in ecosystem stability and function by consuming dominant competitors (*1*, *2*). Predators can also destabilize ecosystems or collapse food webs if they become too abundant, or if their prey do not have natural defenses against predation. Generally, predators and their prey have evolved over time to coexist. Prey have anti-predator behaviors or morphological adaptations to avoid being eaten (*3*, *4*). Similarly, predators have adaptations or behaviors that help them to forage optimally and take advantage of prey when they are available (*5*, *6*).

One of the ways the balance between predator and prey adaptations manifests in nature is through density-dependent predation. Predators can exhibit a numerical response to prey densities by increasing reproduction rates due to an overabundance of prey (demographic response) or by gathering in areas with relatively high densities of prey (aggregative response) (*7*). An individual predator may also adjust its predation rate to prey density through a 'functional response' (changes in a predator's consumption rate in response to prey density). Density-dependent mechanisms tend to stabilize prey population dynamics (*8*, *9*) and can maintain population viability when a population is reduced to low levels (*10*).

Certain characteristics of a predator-prey system can help predict which functional response will
be observed. A linear relationship between consumption rate and prey density (type I functional
response) is expected for organisms that do not actively search for prey, such as filter feeders.
Most vertebrate and invertebrate predators exhibit a hyperbolic functional response that increases
to an asymptote due to limits associated with prey handling, ingestion, and metabolism (type II
functional response) (*11*). Predators that feed upon cryptic or otherwise hard-to-find prey exhibit
a sigmoidal functional response, where consumption rates increase slowly at low prey densities
(type III functional response) (*7*). Prey that avoid predators can achieve a low-density refuge; thus,
the functional response can explain the distribution of prey items, and it can be used to predict the
persistence of prey species at low densities (*12*).
There are many mathematical models that can be used to predict predator-prey dynamics (*13*).
These models contain nonlinear functions describing the density-dependent interactions between
predator and prey. Due to these nonlinearities, model behavior often includes shifts to alternative
stable states (*14*). These states may include extinction of one or both species, or coexistence steady
states where both predator and prey are able to coexist at densities predicted by the model. Multiple
stable coexistence states are possible in fairly simple predator-prey functions (*15*, *16*).
A dynamic systems approach can predict coexistence states in natural and managed systems.
Steady states analysis been used to develop optimum harvesting and conservation strategies for
forest and rangeland systems (*17*, *18*). The theory surrounding steady states and bifurcations is
well developed, but there are very few empirical tests. To date, empirical tests of coexistence
steady states in predator-prey systems have involved populations of small organisms such as
unicellular organisms, yeast, small crustaceans, or bacteria (*19–21*). Identifying or establishing
coexisting predator-prey systems for use in testing model predictions has proven difficult,
especially for macro-organisms. Of the few empirical tests of steady states that focus on macro-
organisms, none examines interactions between natural populations of predators and prey (*16*, *22*,
23). Here, we show that a severe storm resulted in a phase shift to a low-density state for the soft-
shell clam *Mya arenaria*, which was once a biomass dominant in Chesapeake Bay, in the face of
predation by the blue crab *Callinectes sapidus*. We examine the empirical evidence for the low-
density steady state predicted by a Lotka-Volterra model of this predator-prey system using data
from long-term monitoring, a field survey, and a field experiment.

History of predator-prey system:
Tropical Storm Agnes, which reached and remained in the Chesapeake Bay watershed 21-23
June 1972, has long been suspected of resulting in long-term changes for the Bay (*24*). Tropical
Storm Agnes was a "100-year storm" that caused sustained, extremely low salinities (Figure 1)
and increased sedimentation throughout Chesapeake Bay (*25*, *26*). This storm has been blamed for
the loss of seagrass in certain areas of Chesapeake Bay (*24*), high mortality rates and recruitment
failure in oysters *Crassostrea virginica* (*27*), and declines in abundance of the soft-shell clam *Mya*
arenaria, which suffered a mass mortality after the storm (*28*).

Fig. 1. Pre- and post- storm salinity profiles. Salinity profiles for post-Agnes (left) and average
summer (right) conditions. Post-Agnes salinity was measured over the period June 29 – July 3,
1972 (*25*). The summer salinity profile (right) is average surface salinity for 1985-2006 (*29*).

Mya arenaria was abundant enough to support a major commercial fishery throughout Chesapeake
Bay prior to 1972 (*30*). When the population declined abruptly after Tropical Storm Agnes, the

fishery never recovered in lower Chesapeake Bay (Virginia) (*31*). Attempts to revive a commercial
fishery in Virginia waters were never realized after the storm's passage. The commercial fishery
for soft-shell clams in the Maryland portion of the Bay is characterized by variable and low harvest
(*32*); the fishery declined by 89% after the storm and has been near collapse since (*33*).
The failure of *M. arenaria* to recover from storm-related declines has been attributed to
predation, habitat loss, disease, rising temperatures, and overfishing (*31*). The Virginia and
Maryland portions of Chesapeake Bay have different habitats, disease dynamics, climates, and
fishing pressure; therefore, these factors are unlikely to explain the inability of *M. arenaria* to
recover from low density in both regions (*31*, *32*). More recently, disease has been blamed for an
added minor decline in *M. arenaria* (*32*); however, there is no evidence that disease prevalence or
intensity are correlated with *M. arenaria* density (*31*).
Experimental evidence suggests that on a local scale, interactions between *M. arenaria* and their
major predator, the blue crab *C. sapidus* (*34*), are capable of keeping clams at low densities (*35*,
36). *Mya arenaria* burrow deeply in sediments, and when clams are at low densities, crabs are
unable to detect their presence (*35*). The result is a low-density refuge for *M. arenaria*, driven by
disproportionately low predation, which is characteristic of a sigmoidal functional response (*35*,
36). Given this evidence regarding a potential mechanism for the decline in *M. arenaria* and
maintenance of the population at low density, this study examines the empirical evidence for a
low-density steady state in this predator-prey system, and the impact of Tropical Storm Agnes on
basin-scale population dynamics of *M. arenaria*.
158
159 **Results:**
160   Tropical Storm Agnes in 1972 resulted in a phase shift for *M. arenaria*, which was maintained
161 at low abundance likely due to predation by the blue crab *C. sapidus*. An abrupt shift in clam
162
Fig. 2. Predator-prey time series. Time series for soft-shell clam *Mya arenaria* landings (red) and
adult female blue crab *Callinectes sapidus* abundance (blue). Blue crab data are log-transformed
average female abundance per tow (VIMS trawl survey). *Mya arenaria* data are fisheries landings
(1000 bushels) (*33*). Vertical dashed line represents Tropical Storm Agnes (1972), and the location
of the changepoint from time series analysis. Photo credits: C.N. Glaspie (clam), R.N. Lipcius
(crab).
169
abundance was identified in 1972, the year of Tropical Storm Agnes (Figure 2). Before the storm,
crab abundance was positively correlated with clam abundance at a lag of 1 y (r = 0.66, p = 0.01),
indicating that each year, clams were prey for juvenile crabs that recruited to the fishery at one
year of age (Figure 3a). After the storm, clam abundance was negatively correlated with crab
abundance with a lag of 1 y (r = -0.48, p = 0.04), indicating that each year, crabs were consuming
juvenile clams that would have recruited to the fishery a year later (Figure 3b). This is consistent
with a phase shift from a system controlled from the bottom-up by prey resources, to a system
controlled from the top-down by predation pressure on bivalves.
178
179
180 Fig. 3. Pre- and post- storm relationship between crab abundance (log-transformed average
181 female abundance per tow, VIMS trawl survey) and clam landings (1000 bushels, fisheries-
182 dependent data). (a) Before the storm, crab abundance was positively correlated with clam

183   abundance (1 y lag). (b) After the storm, clam abundance was negatively correlated with crab
184   abundance with (1 y lag).
185
Predator-prey modeling confirmed the presence of high-density (near carrying capacity at 173.99
clams m^{-2}) and low-density (at 1.41 clams m^{-2}) steady states separated by an unstable steady state
at 20.93 clams m^{-2} (Figure 4). We propose that *M. arenaria* existed in Chesapeake Bay at high
densities until perturbed past the unstable steady state in 1972 by Tropical Storm Agnes.
Thereafter, it was able to persist at low density due to the low-density refuge from blue crab
predation (*35*, *36*), rather than collapsing to local extinction. Unfortunately, *M. arenaria* is unlikely
to rebound to high abundance without a beneficial disturbance, such as a considerable recruitment
episode or substantial reduction in predation pressure, which propels it above the unstable steady
state (Figure 4) and concurrently allows it to overcome the exacerbated disease burden.
195
196   Fig. 4. Slope field diagrams for predator-prey models. Trajectories of *Mya arenaria* density
197   either approach a high-density stable steady state at carrying capacity (green and purple lines) or
198   a low-density stable steady state at 1.41 clams m^{-2}. Trajectories diverge from an unstable steady
199   state at 20.93 clams m^{-2}.
200
Predator-prey models with these two species alone were capable of reproducing observations of
clam densities and mortality rates, consistent with the idea that blue crabs are a major driver of *M.*
arenaria population dynamics (*34–36*). The low-density steady state predicted by the predator-
prey model is similar to observed densities of *M. arenaria* in Chesapeake Bay. *Mya arenaria*
persist in the lower Chesapeake Bay at average densities of 0.4 – 1.73 m^{-2} (95% CI), despite
episodes of high recruitment (*31*). In the field, juvenile *M. arenaria* exposed to predators suffered
76.3% higher mortality as compared to caged individuals (*37*). Predator exclusion treatments
confirmed that blue crabs were responsible for most of the mortality of juvenile *M. arenaria* (*37*),
and mortality rates observed in the field were comparable to mortality rates predicted by the model
for blue crab density 4.8 m^{-2}, which is a typical density for juvenile crabs in the summer months
in Chesapeake Bay (*38*).
212
Discussion:
The observations, theory, and mechanistic basis indicate that *M. arenaria* was subjected to a
storm-driven phase shift to low density, which has been maintained by blue crab predation in
Chesapeake Bay. Extreme weather events are costly, and they are likely to become even more
common with predicted increases in the intensity and frequency of extreme events due to
anthropogenic climate change (*39*). When examining the cost of extreme weather, ecological
impacts are rarely considered, even though the impacts of such events on the ecosystem may be
severe (*40*). Evidence for storm-driven phase shifts in coral reefs (*15*), kelp ecosystems (*41*), and
now soft-sediment communities (current study) suggests that management of ecosystems should
include an examination of nonlinear interactions and the potential for phase shifts.
To our knowledge, this is one of a handful of empirical tests of predator-prey theory as a
predictive tool in natural systems. More evidence is needed to fully evaluate the predictive
potential of steady-state analysis involving predator-prey models, and the value of a dynamics
systems approach to ecosystem modeling efforts. However, the approach described here can be
adapted to a variety of predator-prey systems with available estimates of life history parameters,

population density, and distribution. The approach may also be expanded to include more than two
species, or encompass additional complexity, such as metabolic functions.
The concepts encapsulated in the crab-clam predator-prey system, including population self-
limitation, consumer-resource oscillations, and the functional response are widespread in nature.
These concepts may be considered "laws" of population ecology (*42*). Each concept listed above
introduces nonlinear dynamics into population models, especially those with multiple interacting
species. Given the preponderance of multispecies interactions exhibiting nonlinear dynamics,
multiple steady states may be a general ecological phenomenon.
236
Methods:
Changepoint analysis of time series was conducted using R statistical software on *M. arenaria*
landings (*43*) and log-transformed average adult female *C. sapidus* abundance (VIMS trawl
survey) in the Chesapeake Bay from 1958-1992, with an AIC penalty and using the segment
neighbor algorithm (*44*). This time period was chosen for analysis because it begins when *M.*
arenaria landings data first became available and ends before the slow decline in landings in the
early 1990s due to fisheries collapse.
Predator-prey ordinary differential equation (ODE) models were modified with a type III
functional response:
246

247   $$N(t) = rN\left(1 - \frac{N}{K}\right) - f(N)P$$

249   where N is the density of prey, P is the density of predators, r is the intrinsic per capita growth rate,
250   K is the carrying capacity, and $f(N)$ takes the form of a sigmoidal functional response:
251

$$f(N) = \frac{N^2 bT}{1 + cN + bT_h N^2}$$

where T is the time available for foraging, T_h is handling time, and b and c are components of the
attack rate in a sigmoidal functional response (*11*).
Models were parameterized using data from the literature as follows: P = 0.06 m^{-2} (*45*), r = 1.75
y^{-1} (*46*), K = 200 m^{-2} (*47*), T = 1 y, T_h = 0.0015 y (*35*), b = 26.30 y^{-1} (*35*), and c = 0.14 (*35*).
Analytical solutions of steady states were calculated using Matlab statistical software. Stability of
each coexistence steady state was determined by examining the sign of eigenvalues.
To examine mortality rates, we solved the equation for number consumed:
261

262   $$N_E = N - f(N)\,P$$

264   where N_E = the number of clams eaten calculated for a period of 8 d (0.022 y) at an initial density
265   of N = 48 m^{-2} to match the field predation experiments (*37*). We then calculated mortality as:
266

267   $$M = \frac{N - N_E}{N} * 100\,\%$$

269   where M = percent mortality. Density of predators P was allowed to vary to achieve M = 76.3%
270   (*37*), and the resultant predator density that achieved observed mortality rates of juvenile *M.*

arenaria was compared to published *C. sapidus* densities for Chesapeake Bay. Data and R code files are available in the Knowledge Network for Biocomplexity (KNB) repository (*48*).

Acknowledgments: We gratefully acknowledge assistance by the students and staff of the Community Ecology and Marine Conservation Biology labs at the Virginia Institute of Marine Science. CNG performed the analysis and wrote the manuscript. CNG, RDS, and RNL contributed substantially to study conception and design, interpretation of the data, and manuscript revisions. This material is based upon work supported by the National Oceanic and Atmospheric Administration grant number NA11NMF4570218; the Environmental Protection Agency EPA STAR Fellowship under grant number FP91767501; and the National Science Foundation Administration GK-12 program under grant number DGE-0840804. This paper is contribution number XXXX from the Virginia Institute of Marine Science, College of William & Mary.

References:
1. J. Lubchenco, S. D. Gaines, A unified approach to marine plant-herbivore interactions. I. Populations and communities. *Annu. Rev. Ecol. Syst.* **12**, 405–437 (1981).
2. S. A. Boudreau, B. Worm, Ecological role of large benthic decapods in marine ecosystems: A review. *Mar. Ecol. Prog. Ser.* **469**, 195–213 (2012).
3. R. Bibby, P. Cleall-Harding, S. Rundle, S. Widdicombe, J. Spicer, Ocean acidification disrupts induced defences in the intertidal gastropod Littorina littorea. *Biol. Lett.* **3**, 699–701 (2007).
4. W. L. Whitlow, Changes in survivorship, behavior, and morphology in native soft-shell clams induced by invasive green crab predators. *Mar. Ecol.* **31**, 418–430 (2010).
5. P. M. Meire, A. Ervynck, Are oystercatchers (*Haemoptopus ostralegus*) selecting the most profitable mussels (*Mytilus edulis*)? *Anim. Behav.* **34**, 1427–1435 (1986).
6. R. R. Rindone, D. B. Eggleston, Predator-prey dynamics between recently established stone crabs (*Menippe* spp.) and oyster prey (*Crassostrea virginica*). *J. Exp. Mar. Bio. Ecol.* **407**, 216–225 (2011).
7. C. Holling, The components of predation as revealed by a study of small mammal predation of the European pine sawfly. *Can. Entomol.* **91**, 293–320 (1959).
8. T. Royama, *Analytical Population Dynamics* (Chapman & Hall, London, UK, 1992).
9. P. Turchin, *Complex Population Dynamics* (Princeton University Press, Princeton, New Jersey, 2003).
10. D. Cushing, *Marine Ecology and Fisheries* (Cambridge University Press, NY, 1975).
11. M. P. Hassell, J. H. Lawton, J. R. Beddington, Sigmoid functional responses by invertebrate predators and parasitoids. *J. Anim. Ecol.* **46**, 249–262 (1977).
12. D. B. Eggleston, R. N. Lipcius, A. H. Hines, Density-dependent predation by blue crabs upon infaunal clam species with contrasting distribution and abundance patterns. *Mar. Ecol. Prog. Ser.* **85**, 55–68 (1992).
13. C. J. Briggs, M. F. Hoopes, Stabilizing effects in spatial parasitoid-host and predator-prey models: A review. *Theor. Popul. Biol.* **65**, 299–315 (2004).
14. J. M. Drake, B. D. Griffen, Early warning signals of extinction in deteriorating environments. *Nature*. **467**, 456–459 (2010).
15. P. J. Mumby, A. Hastings, H. J. Edwards, Thresholds and the resilience of Caribbean coral reefs. *Nature*. **450**, 98–101 (2007).
16. A. M. Kramer, J. M. Drake, Experimental demonstration of population extinction due to a predator-driven Allee effect. *J. Anim. Ecol.* **79**, 633–639 (2010).

17. C. T. Bauch, R. Sigdel, J. Pharaon, M. Anand, Early warning signals of regime shifts in
coupled human–environment systems. *Proc. Natl. Acad. Sci.* **113**, 14560–14567 (2016).
18. N. Hritonenko, Y. Yatsenko, R. U. Goetz, A. Xabadia, Optimal harvesting in forestry:
Steady-state analysis and climate change impact. *J. Biol. Dyn.* **7**, 41–58 (2013).
19. J. M. Drake, B. D. Griffen, Early warning signals of extinction in deteriorating environments.
Nature. **467**, 456–459 (2010).
20. L. Luckinbill, Coexistence in laboratory populations of Paramecium aurelia and its predator
Didinium nasutum. *Ecology*. **54**, 1320–1327 (1973).
21. P. van den Ende, Predator-prey interactions in continuous culture. *Science (80-.)*. **181**, 562–
564 (1973).
22. J. Jiang *et al.*, Predicting tipping points in mutualistic networks through dimension reduction.
Proc. Natl. Acad. Sci., 201714958 (2018).
23. G. G. McNickle, W. D. Evans, Toleration games: Compensatory growth by plants in
response to enemy attack is an evolutionarily stable strategy. *AoB Plants*. **10**, 1–14 (2018).
24. R. J. Orth, K. A. Moore, Chesapeake Bay: An unprecedented decline in submerged aquatic
vegetation. *Science (80-.)*. **222**, 51–53 (1983).
25. J. R. Schubel, H. H. Carter, W. B. Cronin, in *The Effects of Tropical Storm Agnes on the*
Chesapeake Bay Estuarine System, E. P. Ruzecki, Ed. (The Johns Hopkins University Press,
Baltimore, MD, 1976), pp. 33–65.
26. J. R. Schubel, in *The Effects of Tropical Storm Agnes on the Chesapeake Bay Estuarine*
System, J. R. Schubel, Ed. (The Johns Hopkins University Press, Baltimore, MD, 1976), pp. 179–
200.
27. D. S. Haven, W. J. Hargis, J. G. Loesch, J. P. Whitcomb, in *The Effects of Tropical Storm*
Agnes on the Chesapeake Bay Estuarine System, A. M. Anderson, Ed. (The Johns Hopkins
University Press, Baltimore, MD, 1976), pp. 488–508.
28. R. L. Cory, M. J. Redding, in *The Effects of Tropical Storm Agnes on the Chesapeake Bay*
Estuarine System, A. M. Anderson, Ed. (The Johns Hopkins University Press, Baltimore, MD,
1976), pp. 478–487.
29. H. Weinberg, Map: Chesapeake Bay Mean Surface Salinity - Summer (1985-2006). *Chesap.*
Bay Progr. (2008), (available at
http://www.chesapeakebay.net/maps/map/https://www.chesapeakebay.net/what/maps/chesapeak
e_bay_mean_surface_salinity_summer_1985_2006).
30. D. S. Haven, "A study of the hard and soft clam resources of Virginia: Annual contract report
for the period 1 July 1969 through 30 June 1970" (Gloucester Point, VA, 1970).
31. C. N. Glaspie, R. D. Seitz, M. B. Ogburn, C. F. Dungan, A. H. Hines, Impacts of habitat,
predators, recruitment, and disease on soft-shell clams *Mya arenaria* and stout razor clams
Tagelus plebeius in Chesapeake Bay. *Mar. Ecol. Prog. Ser.* **603**, 117–133 (2018).
32. C. F. Dungan, R. M. Hamilton, K. L. Hudson, C. B. McCollough, K. S. Reece, Two epizootic
diseases in Chesapeake Bay commercial clams, *Mya arenaria* and *Tagelus plebeius*. *Dis. Aquat.*
Organ. **50**, 67–78 (2002).
33. NMFS, Annual Commercial Landing Statistics, (available at
https://www.st.nmfs.noaa.gov/commercial-fisheries/commercial-landings/annual-
landings/index).
34. C. J. Meise, L. L. Stehlik, Habitat use, temporal abundance variability, and diet of blue crabs
from a New Jersey estuarine system. *Estuaries*. **26**, 731–745 (2003).
35. R. N. Lipcius, A. H. Hines, Variable functional responses of a marine predator in dissimilar

homogenous microhabitats. *Ecology*. **67**, 1361–1371 (1986).

36. R. D. Seitz, R. N. Lipcius, A. H. Hines, D. B. Eggleston, Denity-dependent predation, habitat

variations, and the persistence of marine bivalve prey. *Ecology*. **82**, 2435–2451 (2001).

37. C. N. Glaspie, R. D. Seitz, Habitat complexity and benthic predator-prey interactions in

Chesapeake Bay. *PLoS One*. **13**, e0205162 (2018).

38. G. M. Ralph, R. N. Lipcius, Critical habitats and stock assessment: Age-specific bias in the

Chesapeake Bay blue crab population survey. *Trans. Am. Fish. Soc.* **143**, 889–898 (2014).

39. J. Settele *et al.*, in *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A:*

Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report

of the Intergovernmental Panel on Climate Change, C. B. Field et al., Eds. (Cambridge

University Press, New York, NY, 2014), pp. 271–359.

40. J. R. Thomson *et al.*, The influences of climatic variation and vegetation on stream biota:

Lessons from the Big Dry in southeastern Australia. *Glob. Chang. Biol.* **18**, 1582–1596 (2012).

41. J. E. Byrnes *et al.*, Climate-driven increases in storm frequency simplify kelp forest food

webs. *Glob. Chang. Biol.* **17**, 2513–2524 (2011).

42. P. Turchin, Does population ecology have general laws? *Oikos*. **94**, 17–26 (2001).

43. M. L. Homer, C. F. Dungan, M. L. Tarnowski, "Assessment of Chesapeake Bay commercial

softshell clams *Mya arenaria* and *Tagelus plebeius* with emphasis on abundance and disease

status" Report to the NOAA Chesapeake Bay Office Fisheries Science Program (2011).

44. I. E. Auger, C. E. Lawrence, Algorithms for the optimal identification of segment

neighborhoods. *Bull. Math. Biol.* **51**, 39–54 (1989).

45. MD DNR, 2017 Blue Crab Winter Dredge Survey, (available at

http://dnr.maryland.gov/fisheries/Pages/blue-crab/dredge.aspx).

46. D. J. Brousseau, Population dynamics of the soft-shell clam *Mya arenaria*. *Mar. Biol.* **50**,

63–71 (1978).

47. B. J. Abraham, P. L. Dillon, "Species profiles: Life histories and environmental requirements

of coastal fishes and invertebrates (Mid-Atlantic)- softshell clam" U.S. Fish and Wildlife Service

Biological Report 82(11.68) (1986).

48. C. N. Glaspie, Blue crab *Callinectes sapidus* and soft-shell clam *Mya arenaria* index of

abundance time series 1958-1992. *Knowl. Netw. Biocomplexity* doi:10.5063/F19S1PB9

(2019).

409   **Figures:**

410
411   Figure 1.

424
425   Figure 2.

426
427   Figure 3.
428

429
430   Figure 4.
431