Evidence against tetrapod-wide digit identities and for a limited frame shift in bird wings

Thomas A. Stewart[1,2]*, Cong Liang[3], Justin L. Cotney[4], James P. Noonan[5], Thomas J. Sanger[6], and Günter P. Wagner[1,3]*

[1]Department of Ecology and Evolutionary Biology, Yale University
[2]Minnesota Center for Philosophy of Science, University of Minnesota
[3]Systems Biology Institute, Yale University
[4]Department of Genetics and Genome Sciences, UConn Health
[5]Department of Genetics, Yale School of Medicine
[6]Department of Biology, Loyola University in Chicago
*To whom correspondence should be addressed

Summary: In crown group tetrapods, individual digits are homologized in relation to a pentadactyl ground plan. However, testing hypotheses of digit homology is challenging because it is unclear whether digits develop with distinct and conserved gene regulatory states. Here we show dramatic evolutionary dynamism in the gene expression profiles of digits, challenging the notion that five digit identities are conserved across amniotes. Transcriptomics of developing limbs shows diversity in the patterns of genetic differentiation of digits, although the anterior-most digit of the pentadactyl limb has a unique, conserved expression profile. Further, we identify a core set of transcription factors that are differentially expressed among the digits of amniote limbs; their spatial expression domains, however, vary between species. In light of these results, we reevaluate the frame shift hypothesis of avian wing evolution and conclude that only the identity of the anterior-most digit has shifted position, suggesting a 1,3,4 digit identity in the bird wing.

Limbs evolved from paired fins in the Late Devonian, and early tetrapods possessed more than five digits on the fore- and hindlimbs[1,2]. Later in the tetrapod stem, a pentadactyl pattern stabilized as the ground plan for the limb. Individual digits are homologized between species and between fore- and hindlimbs in reference to this pentadactyl ground plan[3]. However, it remains controversial whether such hypotheses of identity correspond to distinct developmental programs among the digits (developmental identities), or just the relative position of digits along the limb's anteroposterior axis (positional identities)[4-7]. Below we use the symbols D1, D2, etc. to indicate positional identities in the pentadactyl ground plan, rather than to indicate developmental

identities.

The anterior-most digit (D1) (*e.g.,* human thumb) appears to have a distinct

developmental identity as compared to the more posterior digits (D2-D5). D1 is marked by a

unique gene expression profile—low expression of *HoxD11* and *HoxD12* and high expression of

Zic3 relative to other digits[7-9]—and it appears able to develop independently of *Shh* signaling[9-11].

Additionally, analysis of morphological variation in primates identified a high degree of variational

independence of D1 relative to the more posterior digits[12]. Models of posterior digit identity have

been proposed according to the relative exposure of limb bud mesenchymal cells to *Shh,* which

emanates from the zone of polarizing activity prior to digit condensation[10,11]. However, broadly

conserved marker genes for individual posterior digits have not been identified in the interdigital

mesenchyme, the signaling center that patterns digits[13,14]. For instance, while the combinatorial

expression of *Tbx2* and *Tbx3* is necessary to generate the phenotypes of D3 and D4 in chicken

hindlimb[15], it is questionable whether these developmental identities are conserved in other

species, like mouse, with limited morphological differentiation of the posterior digits.

Debates of digit homology are especially challenging to resolve when limbs have fewer

than five digits. This problem has been most actively investigated in the tridactyl avian wing,

because of the appearance of conflict between paleontological and developmental data[16]. The

fossil record of theropod dinosaurs shows a clear pattern of reduction of the posterior two digits in

the lineage leading to birds, yet digits in the wing have been described as developing in the

middle three positions of a pentadactyl developmental groundplan[17-22]. To explain this

discrepancy, the frame shift hypothesis was proposed[16]. It posited that a homeotic shift occurred

in the avian stem such that the developmental programs that were once expressed in D1, D2,

and D3 are now executed in the digits that develop in positions D2, D3, and D4 respectively.

Comparative analyses of gene expression have found support for this hypothesis: *in situ*

hybridization and transcriptomics have revealed similarity between the anterior digit of the adult

avian wing, which develops in position D2, and D1 of other limbs[7,23].

Studies that aim to test digit homology assume that developmental identities (1) were

present in a common ancestor, (2) are conserved among the descendent lineages, and (3) are

reflected in gene expression profiles. Here we present comparative transcriptomic data from five

species that challenge these assumptions among amniotes by documenting a surprising diversity

of digital gene expression profiles. Analyses further reveal a core set of transcription factor genes

differentially expressed among digits and suggest a new model for the evolution of the bird wing.

Results

Disparity in digit expression profiles

75   To characterize the gene expression profiles of digits in pentadactyl amniote limbs, we

76 sequenced RNA of developing digits and their associated posterior interdigital mesenchyme from

77 the forelimbs of mouse, green anole (*Anolis*), and American alligator (Fig. 1 a). In each of these

78 species, hierarchical cluster analysis (HCA) and principal component analysis (PCA) of the

79 transcriptomes shows a weak signal of sample clustering by digit (Extended Data Fig. 1). The

80 strongest signals of digit-specific expression profiles are observed in D1 of mouse and D4 of the

81 alligator. Groupings of the other digit samples are not well supported. We hypothesized that this

82 result might imply that any signal of gene expression differentiation among digits is overwhelmed

83 by noise when all genes are considered, because most genes are likely irrelevant to the

84 developmental identity of digits. If such a signal exists, we predict that it will be reflected

85 preferentially in the expression of transcription factor and signaling genes. Therefore, we again

86 performed HCA and PCA on the samples of each species, this time using two gene lists: a

87 curated set of known limb patterning genes that are sensitive to *Shh* signaling (N=159)[24], and

88 transcription factor genes (N=2183)[25].

89   In mouse and alligator, HCA and PCA of known limb patterning genes results in

90 clustering of samples by digit (Fig. 1 b, c). In mouse, D1 is strongly differentiated from the other

91 digits. In alligator, an anterior cluster, comprised of digits D1, D2, and D3, is differentiated from a

92 posterior cluster, comprised of D4 and D5. By contrast, analysis of known limb patterning genes

93 in *Anolis* shows weak clustering of samples by digits (Fig. 1 d). This suggests a level of

94 homogeneity among *Anolis* digits that is not observed in either mouse or alligator. Analysis of all

95 transcription factors for these species yields comparable results to what is recovered for limb

96 patterning genes, but with generally lower adjusted uncertainty values in HCAs (Extended Data

97 Fig. 2).

98   To further test the hypothesis that there is limited gene expression differentiation among

99 *Anolis* digits as compared to the other pentadactyl limbs sampled, we took advantage of a result

100 from multiple testing theory[26]: If a differential expression analysis is conducted on two sample

101 types that are not genetically differentiated, then the resultant frequency distribution of *p* values

102 will be uniform within the [0, 1] interval. On the other hand, if there are truly differentially

103 expressed genes among the compared sample types, then the *p* value distribution is expected to

104 be biased towards *p*=0. We conducted differential expression analyses of adjacent digits of the

105 forelimbs of mouse, alligator, and *Anolis* using EdgeR[27,28] and inspected *p* value distributions

106 (Fig. 2). In *Anolis*, all comparisons of adjacent digits result in *p* value distributions that are close to

107 uniform, suggesting that there is very weak, if any, genetic differentiation of adjacent fingers. We

108 note that this result is independent of any *p* value significance threshold or false discovery

109 correction method. By contrast, most of adjacent pairwise digit comparisons for mouse and

110 alligator show a strongly biased *p* value distribution, the exception being D2 and D3 in mouse.

This is consistent with the idea that, in general, most digits in a limb are genetically differentiated,

while in *Anolis* genetic differentiation of digits is minimal or absent.

Given that these three limbs differ in their broad patterns of gene expression

differentiation of digits, we next asked whether individual genes show divergent or constrained

expression patterns across the forelimb in the different species. Specifically, we compared

adjacent digits, identified differentially expressed transcription factor genes, and then assessed

which differences are shared among mouse, alligator, and *Anolis*. Of the 1133 transcription factor

genes that are one-to-one orthologs in these three species, only four genes are differentially

expressed in a conserved pattern among corresponding adjacent digits (Fig. 3). There are three

genes that differentiate D1 from D2 (*Hoxd11, Hoxd12*, and *Sall1)*, and there is one that

differentiates D4 from D5 (*Tbx15*) in all three species. No transcription factors are differentially

expressed in all three species between the median digits (*i.e.*, differentiating D2 from D3, or D3

from D4).

If the homogeneity observed among *Anolis* forelimb digits is a derived condition, then this

could limit our ability to diagnose plesiomorphic developmental identities. Therefore, we also

considered the chicken hindlimb, which has digits D1-D4. We reanalyzed published

transcriptomic data for hindlimb digits[7], mapping reads to a new chicken genome (Galgal5.0)[29].

HCA and PCA of digits of the chicken hindlimb show a unique pattern of similarity as compared to

pentadactyl limbs: an anterior cluster, comprised of D1 and D2, is differentiated from the posterior

cluster, comprised of D3 and D4 (Extended Data Fig. 3). Similar to alligator, this pattern of

correspondence among the digits is stable across the developmental window sampled (st. 28–

31). As before, we tested for differential expression in adjacent digits and identified one-to-one

orthologous transcription factor genes that are differentially expressed at the same position

between mouse and alligator forelimb and chicken hindlimb (Fig. 4 a, Extended Data Fig. 4). Of

the 1049 transcription factor genes, ten differentiate D1 and D2 (n=10), none distinguish D2 and

D3, and one (*Tbx3*) differentiates D3 from D4 in all three species (Fig. 4 a).

Overall, data from these four species do not support the hypothesis that amniote digits

have conserved developmental identities. The exception appears to be D1, which likely had a

distinct developmental program in the most recent common ancestor of amniotes. We further

tested whether D1 has a conserved gene expression profile by sequencing RNA from developing

human fore- and hindlimb, which were partitioned into D1 and the posterior digital plate (D2-5). Of

the ten genes identified above as differentiating D1 and D2, six show conserved patterns of

expression change at this position: in all limbs sampled *Hand2, Hoxd11, Hoxd12, and Tfap2b* are

more highly expressed in D2 than D1, and *Alx1 and Pax9* are more lowly expressed in D2 than

D1 (Fig. 4 b)*.*

T-box family genes are predicted to regulate the identities of posterior digits[15]. Our data

provide some support for the hypothesis that this function is conserved across amniotes (Fig. 4

c). *Tbx2*, which was previously shown to regulate posterior digit identity in the chicken hindlimb[14],

shows divergent patterns of expression in the posterior digits of other species. *Tbx3* differentiates

D3 from D4 in mouse, alligator, and chicken hindlimb, and the likelihood that it was recovered by

chance alone is 7.2×10^{-6} (binomial test); however, it is not differentially expressed at this position

in *Anolis* forelimb. *Tbx15* differentiates D4 from D5 among pentadactyl limbs (Fig. 3), and the

likelihood that it was recovered by chance alone is 1.9×10^{-5} (binomial test).

154   Analyses aiming to identify genes that are conserved and differentially expressed at a

particular position within the limb (*e.g.*, between D1 and D2 in mouse, alligator, and *Anolis*) can

be affected by the threshold stringency of the false discovery rate (FDR). Binomial tests, as

presented above, are one means of accounting for this. We present a second strategy for

assessing whether genes identified as differentially expressed in one species behave similarly in

other species that does not depend on a particular FDR threshold being reached in all species.

Specifically, we consider the genes identified as differentially expressed in one species between

adjacent digits (*e.g.*, in mouse, 129 transcription factor genes are identified between D1 and D2).

Then we ask how expression fold change between the two digits in the original species compares

to expression fold change of the same genes and also a set of randomly selected genes of similar

expression levels in other species. To make these comparisons, we calculated Pearson's

correlation of the fold changes between the original genes *versus* each of the two gene sets

(orthologs and random genes) in other species. Results of this approach broadly mirror those

described, above.

168   Among the pentadactyl limbs sampled, genes differentially expressed between D1 and

D2 behave consistently between species and can be distinguished from random genes, and

comparisons of the more posterior digits do not clearly distinguish orthologs from random genes,

(Extended Fig. 5 a-d). If chicken hindlimb is considered instead of the *Anolis* forelimb, we again

obtain strong support for conserved behavior of genes at the position D1 and D2, weaker support

for conserved gene behavior between D2 and D3, and comparisons at the position D3 and D4 do

not clearly distinguish orthologs from random genes (Extended Fig. 5 e-g). Thus, testing for

genes that are differentially expressed at the same position can recover genes that behave

consistently across species (*i.e.*, *Tbx15* between D4 and D5 among pentadactyl limbs, and *Tbx3*

between D3 and D4 between mouse, alligator, and *Anolis*), while comparisons of all genes

differentially expressed for these species might not show evidence of broadly conserved profiles.

Conversely, while we might obtain modest evidence for shared behavior among differentially

expressed genes (*i.e.*, between digits D2 and D3 among mouse, alligator, and chicken), there

might be no individual genes recovered as differentially expressed among the taxa at that

position. However, both types of comparisons between digits D1 and D2 paint the consistent

picture that D1 exhibits a shared digit identity across these limbs.

A core set of digit patterning genes

Given our result that the gene expression profile of digits is evolutionarily dynamic, we

next tested whether a conserved set of genes might pattern amniote autopods, albeit in different

spatial patterns. Specifically, we reanalyzed transcriptomic data of mouse, alligator and *Anolis*

forelimbs and chicken hindlimb, conducting ANOVA to test for genes that were differentially

expressed between any two digits in the limb, not just adjacent digits. This analysis recovers

genes that are differentially expressed between some digits in the limb, but it does not indicate

between which digits a gene is differentially expressed. The number of differentially expressed

transcription factor genes differs greatly among species: 356 in mouse, 377 in alligator, 34 in

Anolis, and 144 in the chicken hindlimb (FDR <0.05, Fig. 5 a). This is consistent with previous

results (above) that showed the *Anolis* forelimb to be more homogeneous than other sampled

limbs. Therefore, we focused on transcription factor genes that are one-to-one orthologous

between mouse, alligator, and chicken and identified a set of 49 genes that are differentially

expressed in these three limbs (Fig. 5 b). We call these *conserved differentially expressed genes*

(CDEGs). The expected number of overlapping genes among these sets by chance alone is 7.57,

and the probability of observing an overlap of 49 genes or more by chance is $<10^{-6}$ (binomial

test). Thirteen of the CDEGs are included in the list of limb patterning genes sensitive to *Shh*

signaling[24]. To assess whether this gene set is biologically meaningful, we performed HCA and

PCA on the samples of each species using the 49 CDEGs. In *Anolis,* we considered the subset

(n=42) that are one-to-one orthologs across all four species. In combination, CDEGs can produce

unique expression profiles of each digit within a limb (Fig. 5 c) and show patterns similar to those

generated by analyses of known limb patterning genes (Fig. 1 b-d, Extended Fig. 3 c).

Analysis of amniote limbs showed that targeted gene lists generated either

experimentally (*i.e.*, known limb patterning genes[24]), by gene ontology (*i.e.*, all transcription

factors[25]), or statistically (*i.e.*, 49 CDEGs), can reveal distinct gene expression profiles among

digits of a limb, which are not observed in the full transcriptome. The spatial digit expression

profiles of these genes, however, is species specific. In light of these results, we reevaluated the

frame shift hypothesis of bird wing origin[16].

Reevaluating the frame shift hypothesis

The frame shift hypothesis predicts that the three digits of the adult avian forelimb, which

we refer to here as D2, D3, and D4 according to their developmental position[17-22], will express the

developmental programs observed in the digits D1, D2, and D3 of other limbs[16]. This hypothesis

was tested previously by analyzing the transcriptomes of chicken fore- and hindlimb digits[7]. That

study found correspondence between forelimb D2 and hindlimb D1, consistent with the frame

shift hypothesis. However, correspondence of more posterior digits was not detected[7].

We re-analyzed published transcriptomic data of digits from the chicken forelimb[7] and

compared them to digits of the chicken hindlimb. Surprisingly, when the 49 CDEGs are

considered, gene expression profiles of forelimb digits D2, D3, and D4 correspond to hindlimb

digits D1, D3, and D4, respectively (Fig. 6 a). Analyses of transcription factor genes and known

limb patterning genes show a consistent pattern (Extended Data Fig. 6). Similarity between the

posterior two digits of the chicken fore- and hindlimb (D3 and D4 in each limb) can also be

observed in the expression patterns of numerous individual genes that are known to be involved

in the patterning of digits (Fig. 6 b).

To assess whether spatial gene expression profiles can be conserved between the fore-

and hindlimbs of a species, even when they differ in digit number, we performed *in situ*

hybridization in alligator. We evaluated expression of *Tbx2*, *Tbx3*, and *Sall1*, three transcription

factor genes identified as differentially expressed between alligator forelimb D3 and D4. *In situ*

hybridization confirms their expression in the posterior interdigital mesenchyme (Fig. 6 c) and

shows conserved positional expression patterns for *Tbx3* and *Sall1* between the forelimb and

hindlimb. It is unclear whether the pattern also holds for *Tbx2*, where difference in expression

level detected from RNA sequencing appear to reflect the proximodistal extent of gene

expression.

We also tested the frame shift hypothesis by comparing the limbs of chicken to the

pentadactyl forelimbs of other species. For each pentadactyl species, PCA were run using the

CDEGs (49 in mouse and alligator, and 42 in *Anolis*), and chicken samples were projected into

the reference PCA plane as supplementary observations. CDEGs were used because they can

produce digit-specific expression profiles for mouse and alligator forelimb and chicken hindlimb,

and because these patterns are reflective of more inclusive gene lists, as described above. These

projections show that the digits D2, D3, and D4 of the bird wing consistently fall into regions of the

PCA plane corresponding to the digits D1, D3, and D4 of other limbs. Although it is difficult to

differentiate D3 and D4 expression in all species, it is clear that D3 of the chicken forelimb does

not correspond in its expression profile of these genes to the D2 of the other limbs sampled (Fig 6

d).

Some have proposed on the basis of *Shh* expression that the digits in the avian wing are

of positions D1, D2, and D3[30]. This model, like the frame-shift hypothesis, predicts wing digits

have developmental identities corresponding to the digits D1, D2, D3 of other limbs. The results

presented here as well, as the morphological evidence for five digit condensations[17-22], suggest

Shh expression is not a conserved marker of digit identity or position in limbs with highly reduced

digit number.

Discussion

Serial homologs are repeated body parts, generated by a common developmental

program. In the case of digits, chondrogenic condensations are generated by a reaction-diffusion

Turing-type mechanism[31,32]. Serial homologs can be developmentally identical (homomorph

parts) or they can assume distinct developmental identities through the differential expression of

regulatory genes (paramorph parts)[33]. The degree to which serial homologs are individuated can

be difficult to assess from morphology alone, because the same developmental program can lead

to different morphological outcomes depending on the developmental environment[34,35]. However,

detailed analyses of gene expression and regulation can identify developmentally individualized

body parts.

In this study, we performed a comparative analysis of whole genome expression data to

test the hypothesis that digits have conserved developmental identities. In interpreting our data,

we acknowledge that gene expression does not demonstrate gene function. Nevertheless, a lack

of differential gene expression between digits is evidence of a lack of developmental

individuation, and a high level of differential expression (particularly in transcription factor and

signaling genes) is evidence for distinct gene regulatory states.

The developmental stages studied here were selected on the basis of previous

experimental work. Dahn and Fallon[13] demonstrated in the chicken hindlimb that genes

expressed in the interdigital mesenchyme regulate digit-specific morphologies, including the

number of phalanges. Subsequent work in the chicken hindlimb showed that this signaling, in the

phalanx-forming region, is active between stages 27 and stage 30[14]. Here, we analyzed the

expression profiles chicken hindlimb digits of stages 28 and 31 and showed that expression

profiles of limb patterning genes and transcription factor genes are stable over this developmental

window (Fig. 4 b, c; Extended Data Fig. 3 b, c; Fig. 6 b). Thus, signals pertinent to digit patterning

continue to be expressed at late stages of limb development, even after phalanges have formed.

Analyses of alligator show a consistent pattern: between stages 18 and 19.5, the expression

patterns of limb patterning genes and transcription factor genes are stable as assessed by HCA,

PCA, and in profiles of genes of interest (Fig. 1 c, Extended Data Figure 2, Fig. 4 b, c, Fig. 6 c).

Although not all species were sampled at multiple time points, we argue on the basis of these

comparisons in chicken and alligator that it is unlikely our conclusions on the evolution and

development of digit identity are biased by temporal dynamism in gene expression within the

developmental window studied here.

Our analyses show that patterns of regulatory gene expression in digits are evolutionarily

dynamic (Fig. 7 a). The developmental identities of digits are evolving across amniotes and can

be lineage-specific. The exception is a conserved developmental identity that characterizes the

D1 of mouse, alligator and *Anolis* forelimbs, chicken hindlimb, and human fore- and hindlimbs

(Fig. 4 b). This digit identity is unlikely to be an edge-effect (*i.e.*, merely a corollary to which digit

occupies the most-anterior position in a limb). In the rabbit hindlimb, which has lost the digit D1,

this developmental identity is not observed in D2, despite that digit now occupying the anterior-

most position in the limb[36]. Additionally, in the hindlimb of Silkie chicken mutants, which have

additional anterior digit on their foot, developmental identity is preserved in the digit of the

morphology of the native D1, despite that digit no longer occupying the anterior-most position in

the limb[23].

In contrast to D1 we do not find support for conserved digit identities in the more posterior

digits. Among the pentadactyl limbs we studied, no genes consistently differentiate the median

digits (D2, D3, and D4) from one another. And when we consider the chicken hindlimb rather than

Anolis, because similarity among *Anolis* digits might be secondarily derived, we find no gene

differentiates D2 and D3, and only one gene (*Tbx3*) differentiates D3 and D4. There is limited

evidence for a conserved developmental identity for digit D5. A single gene (*Tbx15*) is

differentially expressed between D4 and D5 among mouse, alligator and *Anolis*, however more

genes are shared between just mouse and alligator (Fig. 3, Extended Data Fig 4 b).

Our analyses also identified a core set of regulatory genes, which we call CDEGs, that

are differentially expressed among digits, although species differ in which digits differentially

express the genes (Fig. 5). We propose that the CDEGs represent a "digit differentiation tool kit"

deployed for the individuation of different sets of digits in different lineages, depending on the

adaptive needs of the species. Between mouse and human, 28 the 49 CDEGs have

demonstrated roles in patterning distal limb skeleton (Extended Data Table 1). Of the CDEGs,

only 15 are differentially expressed across the *Anolis* forelimb. This homogeneity appears to be a

derived condition among the taxa sampled, as it is unlikely that the other 34 CDEGs reflect

homoplasy between mammals and archosaurs.

In *Anolis* most fingers, though they differ in number of phalanges, lack developmental

individuality and, thus, appear to be homomorphic. We consider a number of alternative, non-

biological explanations for the unique *Anolis* pattern; however, these do not adequately explain

homogeneity in the data. For example, it is possible that is the limbs were sampled at too-late a

stage, after signals pertinent to digit patterning were expressed. We regard this explanation as

unlikely because, as discussed above, in limbs sampled at multiple time points gene expression

profiles are stable over broad developmental window, through late stages of development.

Another possible alternative explanation is that variance among *Anolis* samples is greater as

compared to other data sets, and that this diminished our ability to detect differentially expressed

genes. We assessed this possibility in two ways. First, we repeated all differential expression

analyses considering only the two most highly correlated samples of each digit for mouse,

alligator and *Anolis*, which consistently had correlation values above 0.99 (Extended Data Fig. 7

a). Results of these two-sample comparisons are consistent with analyses of all three samples

(*e.g.*, compare Fig. 3 and Extended Data Fig. 7 b), indicating that the unique *Anolis* pattern is not

an artifact of sample quality.

Second, we evaluated the dispersion values of our samples. Dispersion is a measure of

variance among samples that is calculated by the software edgeR. This parameter affects the

sensitivity of differential expression analyses (*e.g.*, a set of samples with high dispersion will have

low sensitivity in tests of differential expression), and it can be impacted by specimen pedigree[37].

Anolis embryos were collected from non-siblings, whereas mouse and alligator samples were

collected from siblings. As expected, the mean dispersion value of *Anolis* samples is greater than

either mouse or alligator (Extended Data Fig. 8)*.* The *Anolis* mean dispersion value is consistent

with other data sets in which samples were collected across a population[37]. However, such

differences in dispersion cannot explain the unique *Anolis* pattern. Chicken hindlimb digits, which

were also collected from non-siblings and have dispersion values comparable to *Anolis*

(Extended Data Figure 8), show patterns of differential expression comparable to mouse and

alligator (Fig. 2, Extended Data Fig. 4 a, Fig. 5 a). Thus, neither timing, sample quality, nor

pedigree appears sufficient to explain the *Anolis* data. It appears that homogeneity among the

digits reflects biological reality, and digits in this lineage have undergone secondary

homogenization. Other lineages might have similarly experienced loss of digit identities (*e.g.*,

ichthyosaur forelimbs), and the secondary homogenization of paramorphic serial homologs has

been described in other anatomical systems (*e.g.*, the homodont dentition in cetaceans[38] and the

snake vertebral column[39]).

Finally, we reassessed the homology of fingers in the bird wing and obtain the novel

result that the three digits reflect a combination of translocated digit identities and conserved

identities. The anterior-most digit in the chicken wing, although it develops in position D2[17-22],

exhibits a gene expression profile seen in the D1 of the other limbs; this is consistent with

previous studies and the frame shift hypothesis[7,23]. The gene expression profiles of the posterior

wing digits (D3 and D4), however, do not show evidence of translocation. This is observed most

clearly by comparison to the hindlimb of the chicken, with the pattern recovered when three

different gene lists are considered (transcription factors, limb patterning genes, and CDEGs). As

discussed above, although we cannot diagnose conserved gene expression profiles for the digits

D3 and D4 across amniotes, we obtain indirect evidence for a correspondence of avian digits to

the digits D1, D3, and D4 of other amniote limbs (Fig. 6 d). The possibility of a 1-3-4 pattern of

digit identity in the bird wing has been proposed previously[40] on the basis of experimental

studies[41]. Still, this pattern of correspondence is surprising. It challenges the predominant

hypotheses of digit identity and suggests an alternative scenario for how limb development

evolved in the lineage leading to Aves (Fig. 7 b). Significantly, it indicates that diagnoses of digit

identity from the paleontological record and hypotheses of digit identity based upon gene

expression profiles have a more complex relationship than previously anticipated.

The frame shift hypothesis is an integrative model. It aimed to explain an apparent

incongruity between paleontological and neontological data sets by providing a developmental

account for evolutionary transformation rooted in a mechanistic basis of homology. Our results

show that any such integrative model will be more complicated than previously presumed. Moving

forward, we recommend systematic reappraisal of phalangeal and metacarpal characters along

the avian stem. It has been proposed that patterns of digit reduction in theropods might be more

complex than is generally assumed[40]. For example, study of the ceratosaur *Limusaurus* led to the

hypothesis that in basal tetanurans metacarpal characters correspond to identities 2-3-4, while

phalanges have identities 1-2-3[41], although specifically how this taxa informs the plesiomorphic

avian condition has been contested[42]. Additionally, we recommend continued, broad taxonomic

sampling in studies of limb development. Building expanded, comparative data sets will allow for

quantification of homoplasy between species and between the fore- and hindlimbs, which could

impact hypotheses of digit identity presented here. Finally, continued functional genetic studies

are required to understand how digit-specific phenotypes are regulated and to test the hypothesis

that CDEGs play privileged roles in establishing gene regulatory states in the interdigital

mesenchyme.

The question of how to diagnose the digits of the avian wing is among the oldest in

comparative morphology[3,43]. This study tests several assumptions that underlay many

contemporary studies of the homology and developmental identity of digits. Indeed, it is the first to

comparatively analyze the full gene expression profiles of digits of different species. Such data,

and a willingness to consider hypotheses that previously might have been regarded as heterodox,

is required for the testing and refinement of integrative theories on the nature of limbs.

Methods

Limbs of each species were sampled after digital condensations have formed and after inter-

digital webbing has begun to reduce. RNA was extracted from digits and their associated

posterior inter-digital webbing following the dissection strategy shown in Figure 1 a of Wang *et*

al.[7]. A summary of the taxonomic and tissue sampling strategy is presented in Extended Data

Fig. 9. Investigators were not blinded to the group allocation during the experiment or when

assessing outcomes.

Alligator mississipiensis. Fertilized eggs were collected from six nests of wild individuals at the

Rockefeller Wildlife Refuge in Grand Chenier, Louisiana (USA) in July 2015 by Dr. Ruth Elsey

and colleagues. Eggs were marked with pencil to indicate the side that was facing upwards in the

nest so that embryos would not be injured by rotation of during transfer. Eggs were transported to

Yale University in mesh wire boxes containing original nesting material, and they were incubated

in a temperature-controlled room at 32°C. Eggs were placed on a plastic rack, surrounded with

original nesting material. Racks were suspended four inches above the bottom of a 10 gallon

aquarium. The base of the aquarium was filled with three inches of water, which was heated to

90°F with a submerged aquarium heater. The top of the tank was covered with plexiglass

perforated with 1cm diameter holes to allow for airflow. Humidity within the tank was maintained

at 90%.

Embryos were collected at Ferguson[44] st.18 and 19.5. The left and right limbs of ten

individuals were dissected at each stage. For each stage, individuals sampled were of a single

nest and, therefore, at least half-siblings[45]. Embryos were extracted under sterile, RNAse-free

conditions. Individual digits and the associated posterior interdigital webbing were dissected with

fine scissors and forceps and placed immediately in room temperature RNAlater (Sigma-Aldrich).

Digits were pooled into a single vial (n=20 digits) and divided into four samples of five randomly

selected digits. RNA was extracted from each sample with TRIzol (Thermo Fisher Scientific)

following described methods[46]. For digit D3 stage 18, two of the extractions yielded too little RNA

for sequencing approach described below; therefore, there are only two replicates of this sample

type. RNA quality was assessed using an Agilent Technologies 2100 Bioanalyzer, and samples

with RIN scores above 8.5 were submitted for sequencing at the Yale Genome Sequencing

Center. Sample size (three replicates per sample type) was selected for downstream differential

expression analyses, according to References 27 and 28. To generate strand-specific

polyadenylated RNA libraries, samples were processed as follows: Approximately 500 ng of RNA

was purified with oligo-dT beads and the mRNA recovered was sheared by incubation at 94 C.

First strand synthesis was performed with random primers, and then second strand synthesis was

performed with dUTP to generate strand-specific libraries for sequencing. cDNA libraries were

end-paired, A-tailed adapters were ligated, and the second strand was digested with Uricil-DNA-

Glycosylase. qRT-PCR was performed using a commercially available kit (KAPA Biosystems) to

confirm library quality, and insert size distribution was determined with Agilent Bioanalyzer.

Samples were multiplexed on an Illumina Hiseq 2000. Each sample was sequenced to a depth of

approximately 50 million reads (single-stranded, 75 base pair length).

Reads were mapped to the American alligator genome assembly (allMis0.2) with genome

assembly described by Green *et* al.[47]. Sequenced reads were mapped to the genome using

Tophat2 v2.0.6 on Yale University's Ruddle computing cluster. In Tophat2, reads were first

mapped to the transcriptome, and the remaining reads were then mapped to the genome.

Mapped reads were assigned to genes with *HTSeq v0.5.3p*[48], which was implemented with

Python v2.7.2. In HTSeq, we required that reads be mapped to a specific strand, and to account

for reads that mapped to more than one feature, we ran with the setting "intersection-nonempty."

Mus musculus. Mice embryos (E13.5) were collected from a pregnant female of the strain

C57BL/6J (Jackson Laboratories) in accordance with Yale IACUC #2015-11-483. The female was

pregnant with nine embryos. Digits from the left and right forelimbs of each individual were

dissected as described for alligator and pooled. From these 18 digits, RNA was extracted for

three batches of five digits each. RNA extraction and sequencing methods are the same as

described above for alligator, with the exception of sequencing depth (30 million reads were

obtained for each mouse sample). Sequenced reads were mapped to the mouse genome

assembly GRCm38 with Ensembl annotation v85 and the same Bowtie2 and HTSeq settings as

described for alligator.

Anolis carolinensis. Animals were bred according to published protocols[49] and in accordance

with Loyola University's IACUC protocol #1992. Fertilized eggs were collected and transferred to

petri dishes containing vermiculate moistened by equal mass water. Embryos were shipped to

Yale University and incubated in a Digital Sportsman Incubator (No. 1502) at 26°C. Tissues were

extracted and dissected according to methods described for alligator. Stage 10[50] embryos were

sampled, and RNA was extracted using Qiagen RNeasy Micro Kit. RNA quality was assessed

using with a BioAnalyzer. Samples with RIN scores above 9.0 were submitted for sequencing at

the Yale Genome Sequencing Center. The RNAseq library was prepared with Clontech's Ultra

Low V4 kit (cat# 634890). Each sample was sequenced to a depth of approximately 30 million

reads (single-stranded, 75 base pair length). Sequenced reads were mapped to the *Anolis*

genome assembly (AnoCar2.0, GCA_000090745.1) with Ensembl annotation v85.

Gallus gallus. Published transcriptomes of the digits of the fore- and hindlimbs of chicken[7], were

mapped to the newest chicken genome version (GalGal5.0) with Ensembl annotation v86

following analytic methods described for alligator.

__Homo sapiens.__ Three individuals of Carnegie stage 18[51] were donated to Yale University's

Medical School. The fore- and hindlimbs were sampled, and the anterior-most digit and its

posterior interdigital webbing was dissected from the posterior digital plate. Dissections were

performed and RNA was extracted and sequenced as previously described[43]. Limbs at this stage

are similar to E12.5 of mouse[52]. Sequenced reads were mapped to the human genome assembly

GRCh37 with Ensembl annotation v82 using the same Bowtie2 and HTSeq settings as described

for alligator.

Hierarchical clustering analysis. To estimate relative mRNA abundance, we calculated

transcripts per million (TPM)[53] for the genes of a given gene list (*i.e.*, full trancriptome,

transcription factors, limb patterning genes, CDEGs). The TPM measurement standardizes for

sequencing depth and transcript length. If multiple transcripts are described for a gene, then the

median transcript length was used to calculate TPM; these lengths are available as a

supplementary data file. TPM measures were normalized by a square root transformation, and

hierarchical clustering was performed on the normalized TPM data with the R package

"pvclust"[54]. Clusters were generated from the correlation-based dissimilarity matrix using the

average-linked method. Adjusted uncertainty values were calculated from 1000 bootstrapping

analysis.

      If analyses involved comparisons between developmental stages or limbs, a bulk

correction was performed with a mean transformation (*i.e.*, mean-centering)[7]. In these instances,

Pearson's correlation coefficients range from [-1:1], rather than from [0:1]. Negative correlation

values arise because after bulk correction, a gene's expression values is negative for samples

with a sqrt(TPM) value less than the mean sqrt(TPM) value of that gene among all samples of the

bulk. Bulks were comprised of all samples from a particular stage or all samples of a particular

limb.

Principal component analyses. PCA were performed using the "prcomp" function in R for

various gene lists using square root TPMs as normalized measures of relative mRNA abundance.

As with HCA, if analyses included samples from multiple stages or from different types of limbs, a

bulk correction was performed with a mean transformation (*i.e.*, mean-centering). Bulks were

comprised of all samples from a particular stage or all samples of a particular limb. Loading

values for samples in PCAs and also bootstrap values, which were calculated using the with the

"bootPCA" function of the bootSVD package[55] with centerSamples=True and 1000 bootstrap

samples), are provided as a supplementary data file.

Differential expression testing of adjacent digits. EdgeR (Release 3.1)[27,28] was used to test

for differential expression of adjacent digits (*e.g.*, D1 vs. D2) of mouse, alligator, *Anolis* and

chicken. We used function glmFit and glmLRT in EdgeR, which implemented a generalized

regression model for differential expression test. Specifically, in alligator and chicken PCA and

PCA revealed stable grouping of samples by digit number across stages. Therefore, subsequent

analyses consider these data from two stages simultaneously.

In *Anolis*, pairwise testing of samples revealed nonconventional *p* value distribution, with

a decrease near zero and sometimes a bump near 0.5. Because correlation between replicates

was lower than what was observed in either mouse or alligator (Extended Data Fig. 7), and

because PCA of the full transcriptomes revealed two major clusters of data that did not

correspond to biological phenomena (Extended Data Fig. 1 c), we corrected for the artifact of the

non-biological clusters by including the first principle component in the regression model in

EdgeR. Analyses were also run without this PC1 correction. Results presented in the manuscript

are robust to both analytic approaches, although PC1 correction results in discovery of slightly

more differentially expressed genes for a given false discovery rate. (*e.g.*, compare Fig. 3 and

Extended Data Fig. 10).

Following analyses of differential expression, multiple hypothesis testing was accounted

for by adjusting *p* values following the Benjamini-Hochberg method[26]. We also considered a

second correction method, the *q*-value of Storey[56]. The major results presented in the study are

robust to both methods. Although the Storey method uniformly called more genes as significant at

the FDR threshold of 0.05, the same genes are recovered in the center of Venn diagrams (Fig. 3,

Fig 4a, Fig 5 b).

Comparing correlation of fold change between orthologs and random genes.

To assess whether genes differentially expressed at a given position in one species are behaving

similarly in other species, we compared the relative fold change of these genes to random genes

of similar expression level in the other species. For example, between D1 and D2 of alligator 46

genes are recovered as differentially expressed among the one-to-one orthologs of the three

pentadactyl species sampled (FDR threshold of 0.05) (Fig. 3). For these genes, we calculated the

fold change in TPM according to the equation in Fig. 4 b for all three species (*e.g.*, mouse,

alligator and *Anolis*). Next, for each of the 46 genes, we identified the gene most similar in its

TPM value at the position of the anterior digit (*e.g.*, for the comparisons of D1 vs D2, we matched

the TPM of D1) among the one-to-one orthologous transcription factor genes for the two other

species. Then, we calculated to Pearson's correlation for the vector comprised the gene fold

changes from the original species (alligator) and the orthologs of the other species (mouse and

Anolis). We also calculated Pearson's correlation between gene fold changes from the original

species (alligator) and the random list of random genes of similar expression level for each of the

other species. This was repeated for each the three species, and if a limb was sampled at

multiple time points, for each time point. Finally, to assess whether at a given position

orthologous genes could be distinguished in their behavior to random genes, we compared the

means of these correlations using two tests: t-test and a Mann-Whitney U test.

Differential expression testing between all digits. EdgeR was also used to test for genes that

differed between any combination of digits within a limb. This was done by specifying multiple

coefficients to function glmLRT. As with the pairwise tests, we considered both stages of alligator

and chicken simultaneously and included the first principle component in the regression model for

Anolis. Genes identified as CDEGs for each species are available as supplemental information.

Transcription factors. To identify transcription factor genes, we utilized a published atlas of

human and mouse transcription factors[25]. The published Entrez gene IDs were matched with

human Ensembl gene IDs in Ensembl assembly v85 using BioMart (N=2183). To recover the

species-specific lists of transcription factors, two approaches were taken. In mouse and *Anolis*,

orthologous genes were identified using BioMart's orthology predictions for Ensemble assembly

v85. In alligator and chicken, because these species were analyzed using different assembly

builds, orthology was determined by matching gene symbols to those of human from Ensembl

assembly v85. By this approach, we recovered transcription factors for each species as follows:

1838 in mouse, 1563 in *Anolis*, 1455 in Alligator, and 1217 in chicken. To identify human

orthologs of select genes identified by differential expression analyses, we first used gene

symbols and then confirmed that the ensemble IDs were consistent across Ensemble

assemblies. Genes identified as transcription factors for each species are available as

supplemental information.

To identify one-to-one orthologous transcription factor genes in mouse, alligator, and

Anolis, we used BioMart to generate a list of one-to-one orthologous genes between mouse and

Anolis in Ensembl assembly v85 that correspond to the published transcription factor Entrez

IDs[31]. This gene list was then matched to alligator and chicken by gene symbol to recover

transcription factor genes that are one-to-one orthologs in multiple species.

Limb patterning genes. A Ph.D. dissertation by Carkett[24] identified genes that are sensitive to

Shh signaling by experimental pertubations and *in silico* analyses. From these studies, a

summary list of genes that pattern the autopod was produced (pg. 172). This gene list includes

transcription factor and signaling genes. These gene symbols were matched with each species to

identify the subset of genes present in the genome assemblies that we considered for mouse

(n=151), alligator (n=142), *Anolis* (n=140), and chicken (n=136). Genes identified as limb

patterning genes for each species are available as supplemental information.

***In situ* hybridization.** RNA from a stage 18 alligator limb was extracted, as described above for

sequencing, and cDNA was generated with the High Capacity cDNA Reverse Transcription kit

(Applied Biosystems). Primers were designed with Primer3[57] to amplify fragments of *Tbx2*

(forward, GACCTTGGGCCTTCTCCTAC; reverse, GGGAGTTGTTTGGGGTTTT)*, Tbx3* (forward,

ACCAGGGGTGGATGAACATA; reverse, GCCCTAAAGCAGAGACATGC)*,* and *Sall1* (forward,

CTCACAGCTCAACAACCCAC; reverse, AAACCACCAGCCTCTACCTC). PCR products were

purified with the QIAquick Gel Extraction kit (Qiagen) and cloned with the Topo TA cloning kit

(Invitrogen) into the pCR 4-TOPO vector. Vectors were transformed into DH5*α*-T1 competent

cells. Sense and antisense probes were prepared by linearizing plasmid with the restriction

endonucleases *Not1* or *Pme1* and then transcribing the linearized product with T7 or T3

polymerase, respectively. Probes are labeled with digoxigenin (Sigma-Aldrich) and hybridized

with the alligator embryos at 68°C. Methods for *in situ* hybridization followed GEISHA Project

miRNA Detection Protocol Version 1.1 (http://geisha.arizona.edu/).

Illustrations. The authors modified existing images of the mouse forelimb skeleton[58], alligator

forelimb skeleton[59], chicken limb skeleton[22], and chicken and alligator silhouettes[22]. *Anolis*

silhouette by Sarah Werning.

References

1   Coates, M. & Clack, J. Polydactyly in the earliest known tetrapod limbs. *Nature* **347**, 66-
69 (1990).
2   Lebedev, O. A. & Coates, M. I. The postcranial skeleton of the Devonian tetrapod
Tulerpeton curtum Lebedev. *Zoological Journal of the Linnean Society* **114**, 307-348,
doi:10.1111/j.1096-3642.1995.tb00119.x (1995).
3   Owen, R. *On the nature of limbs*. 132 (John Van Voorst, 1849).
4   Coates, M. Ancestors and homology. *Acta Biotheoretica* **41**, 411-424,
doi:10.1007/BF00709374 (1993).
5   Tabin, C. J. Why we have (only) five fingers per hand: hox genes and the evolution of
paired limbs. *Development* **116**, 289-296 (1992).
6   Zhu, J. & Mackem, S. John Saunders' ZPA, Sonic hedgehog and digit identity – How
does it really all work? *Developmental Biology* **429**, 391-400,
doi:http://dx.doi.org/10.1016/j.ydbio.2017.02.001 (2017).
7   Wang, Z., Young, R. L., Xue, H. & Wagner, G. P. Transcriptomic analysis of avian digits
reveals conserved and derived digit identities in birds. *Nature* **477**, 583–586,
doi:10.1038/nature10391 (2011).
8   Montavon, T., Garrec, J.-F. L., Kerzberg, M. & Duboule, D. Modelling HOX genes
regulation in digits: reverse collinearity and the molecular origin of thumbness. *Genes
Dev* **22**, 346-359, doi:10.1101/gad.1631708 (2008).
9   Chiang, C. *et al.* Manifestation of the Limb Prepattern: Limb Development in the Absence
of Sonic Hedgehog Function. *Developmental Biology* **236**, 421-435,
doi:http://dx.doi.org/10.1006/dbio.2001.0346 (2001).
10   Harfe, B. D. *et al.* Evidence for an expansion-based temporal Shh gradient in specifying
vertebrate digit identities. *Cell* **118**, 517-528, doi:10.1016/j.cell.2004.07.024 (2004).
11   Scherz, P. J., McGlinn, E., Nissim, S. & Tabin, C. J. Extended exposure to sonic
hedgehog is required for patterning the posterior digits of the vertebrate limb. *Dev Biol*
308, 343-354, doi:10.1016/j.ydbio.2007.05.030 (2007).
12   Reno, P. L. *et al.* Patterns of correlation and covariation of anthropoid distal forelimb
segments correspond to Hoxd expression territories. *Journal of Experimental Zoology
Part B: Molecular and Developmental Evolution* **310B**, 240-258, doi:10.1002/jez.b.21207
(2008).
13   Dahn, R. D. & Fallon, J. F. Interdigital regulation of digit identity and homeotic
transformation by modulated BMP signaling. *Science* **289**, 438-441,
doi:10.1126/science.289.5478.438 (2000).
14   Suzuki, T., Hasso, S. M. & Fallon, J. F. Unique SMAD1/5/8 activity at the phalanx –
forming region determines digit identity. *Proc Natl Acad Sci U S A* **105**, 4185-4190
doi:10.1073/pnas.0707899105 (2008).
15   Suzuki, T., Takeuchi, J., Koshiba-Takeuchi, K. & Ogura, T. Tbx Genes Specify Posterior
Digit Identity through Shh and BMP Signaling. *Developmental Cell* **6**, 43-53,
doi:http://dx.doi.org/10.1016/S1534-5807(03)00401-5 (2004).
16   Wagner, G. P. & Gauthier, J. A. 1,2,3=2,3,4: A solution to the problem of the homology
of the digits in the avian hand. *Proc Natl Acad Sci USA* **96**, 5111-5116, doi:10.1073/
pnas.96.9.5111 (1999).
17   Larsson, H. C. E. & Wagner, G. P. Pentadactyl ground state of the avian wing. *Journal of
Experimental Zoology* **294**, 146-151, doi:10.1002/jez.10153 (2002).
18   Feduccia, A. & Nowicki, J. The hand of birds revealed by early ostrich embryos.
Naturweissenschaften **89**, 391-393 (2002).
19   KundráT, M., Seichert, V., Russell, A. P. & Smetana, K. Pentadactyl pattern of the avian
wing autopodium and pyramid reduction hypothesis. *Journal of Experimental Zoology*
294, 152-159, doi:10.1002/jez.10140 (2002).
20   Burke, A. C. & Feduccia, A. Developmental Patterns and the Identification of
Homologies in the Avian Hand. *Science* **278**, 666-668, doi:10.1126/
          science.278.5338.666 (1997).

21 Welten, M. C. M., Verbeek, F. J., Meijer, A. H. & Richardson, M. K. Gene expression and
digit homology in the chicken embryo wing. *Evolution & Development* **7**, 18-28,
doi:10.1111/j.1525-142X.2005.05003.x (2005).
22 de Bakker, M. A. G. *et al.* Digit loss in archosaur evolution and the interplay between
selection and constraints. *Nature* **500**, 445-448, doi:10.1038/nature12336
http://www.nature.com/nature/journal/v500/n7463/abs/nature12336.html - supplementary-
information (2013).
23 Vargas, A. & Fallon, J. F. Birds have dinosaur wings: the molecular evidence. *J Exp*
Zoolog B Mol Dev Evol **304**, 86-90, doi:10.1002/jez.b.21023 (2005).
24 Carkett, M. D. *Comparative analysis of Sonic Hedgehog signalling and the response to*
Sonic Hedeghog signalling in vertebrate forelimbs and hindlimbs. PhD thesis, University
College London, (2015).
25 Ravasi, T. *et al.* An Atlas of Combinatorial Transcriptional Regulation in Mouse and Man.
Cell **140**, 744-752, doi:http://dx.doi.org/10.1016/j.cell.2010.01.044 (2010).
26 Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
(Methodological) **57**, 289-300, doi:citeulike-article-id:1042553
doi: 10.2307/2346101 (1995).
27 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics (Oxford,*
England) **26**, 139-40, doi:10.1093/bioinformatics/btp616 (2010).
28 McCarthy, J. D., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor
RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research* **40**, 9
(2012).
29 Warren, W. C. *et al.* A New Chicken Genome Assembly Provides Insight into Avian
Genome Structure. *G3: Genes|Genomes|Genetics* **7**, 109 (2017).
30 Towers, M., Signolet, J., Sherman, A., Sang, H. & Tickle, C. Insights into bird wing
evolution and digit specification from polarizing region fate maps. *Nat Commun* **2**, 1-7,
doi:10.1038/ncomms1437 (2011).
31 Newman, S. A. & Frisch, H. L. Dynamics of skeletal pattern formation in developing chick
limb. *Science* **205**, 662-668 (1979).
32 Raspopovic, J., Marcon, L., Russo, L. & Sharpe, J. Digit patterning is controlled by a
Bmp-Sox9-Wnt Turing network modulated by morphogen gradients. *Science* **345**,
566-570, doi:10.1126/science.1252960 (2014).
33 Wagner, G. P. *Homology, Genes and Evolutionary Innovation*. (University Press, 2014).
34 Bateson, W. *Problems of Genetics*. (Oxford University Press, 1913).
35 Stewart, T. A. The origin of a new fin skeleton through tinkering. *Biol. Letts* **11**, 1-5,
doi:10.1098/rsbl.2015.0415 (2015).
36 Salinas-Saavedra, M. *et al.* New developmental evidence supports a homeotic frameshift
of digit identity in the evolution of the bird wing. *Frontiers in Zoology* **11**, 33, 1-10,
doi:10.1186/1742-9994-11-33 (2014).
37 Yoon, S. & Nam, D. Gene dispersion is the key determinant of the read count bias in
differential expression analysis of RNA-seq data. *BMC Genomics* **18**, 408,
doi:10.1186/s12864-017-3809-0 (2017).
38 Rose, K. D. *The beginning of the age of mammals*. 431 (The Johns Hopkins University
Press, 2006).
39 Carroll, R. *Vertebrate Paleontology and Evolution* (Freeman, 1988).
40 Xu, X. & Mackem, S. Tracing the evolution of Avian Wing Digits. *Curr Biol* **23**, R538-
R544, doi:10.1016/j.cub.2013.04.071 (2013).
41 Xu, X. *et al.* A Jurassic ceratosaur from China helps clarify avian digital homologies.
Nature **459**, 940–944, doi:10.1038/nature08124
https://www.nature.com/articles/nature08124 - supplementary-information (2009).
42 Vargas, A., Wagner, G. & Gauthier, J. Limusaurus and bird digit identity. *Nature*
Proceedings **713** (2009).
43 Belon, P. *L'histoire de la natvre des oyseavx : avec levrs descriptions, & naïfs portraicts*
retirez du natvrel, escrite en sept livres. (... Gilles Corrozet ... 1555).

44   Ferguson, M. W. J. in *Biology of the Reptilia* Vol. 14 (ed Carl Gans) Ch. 5, 329-481
705   (1985).
706 45   Davis, L. M., Glenn, T. C., Elsey, R. M., Dessauer, H. C. & Sawyer, R. H. Multiple
707   paternity and mating patterns in the American alligator, Alligator mississippiensis.
708   *Molecular Ecology* **10**, 1011-1024, doi:10.1046/j.1365-294X.2001.01241.x (2001).
709 46   Chomczynski, P. & Mackey, K. Short Technical Report - Modification of the TRIZOL
710   reagent procedure for isolation of RNA from Polysaccharaide and proteoglycan rich
711   sources. *Biotechniques* **19**, 942-945 (1995).
712 47   Green, R. E. *et al.* Three crocodilian genomes reveal ancestral patterns of evolution
713   among archosaurs. *Science* **346**, doi:10.1126/science.1254449 (2014).
714 48   Anders, S., Pyl, P. T. & Huber, W. HTSeq - A Python framework to work with high-
715   throughput sequencing data. *Bioinformatics*, doi:10.1093/bioinformatics/btu638
716   (2014).
717 49   Sanger, T. J., Hime, P. M., Johnson, M. A., Diani, J. & Lsos, J. B. Laboratory protocols
718   for husbandry and embryo collection of *Anolis* lizards. *Herpetological Review* **39**, 58-63
719   (2008).
720 50   Sanger, T., Losos, J. & Gibson-Brown, J. A developmental staging series for the lizard
721   genus *Anolis*: a reptilian model system for the integration of evolution, development, and
722   ecology. *J. Morph.* **269**, 129-137 (2008).
723 51   O'Rahilly, R. Early human development and the chief sources of information on staged
724   human embryos. *Europ. J. Obstet. Gynec. Reprod. Biol.* **9**, 273-280 (1979).
725 52   Cotney, J. *et al.* The Evolution of Lineage-Specific Regulatory Activities in the Human
726   Embryonic Limb. *Cell* **154**, 185-196, doi:http://dx.doi.org/10.1016/j.cell.2013.05.056
727   (2013).
728 53   Wagner, G. P., Kin, K. & Lynch, V. J. Measurement of mRNA abundance using RNA-seq
729   data: RPKM measure is inconsistent among samples. *Theory in Biosciences* **131**, 281-
730   285, doi:10.1007/s12064-012-0162-3 (2012).
731 54   Suzuki, R. & Shimodaira, H. Pvclust: an R package for assessing the uncertainty in
732   hierarchical clustering. *Bioinformatics* **22**, 1540-1542, doi:10.1093/bioinformatics/btl117
733   (2006).
734 55   Fisher, A., Caffo, B., Schwartz, B. & Zipunnikov, V. Fast, Exact Bootstrap Principal
735   Component Analysis for p > 1 million. *Journal of the American Statistical Association* **111**,
736   846-860, doi:10.1080/01621459.2015.1062383 (2016).
737 56   Storey, J. D. The positive false discovery rate: a Bayesian interpretation and the q-value.
738   *Ann. Statist.* **31**, 2013-2035, doi:10.1214/aos/1074290335 (2003).
739 57   Untergasser, A. *et al.* Primer3 - new capabilities and interfaces. *Nucleic Acids Research*
740   **40**, e115 (2012).
741 58   Krebs, O. *et al.* Replicated anterior zeugopod (raz): a polydactylous mouse mutant with
742   lowered Shh signaling in the limb bud. *Development* **130**, 6037-6047,
743   doi:10.1242/dev.00861 (2003).
744 59   Muller, G. B. & Alberch, P. Ontogeny of the Limb Skeleton in Alligator mississipiensis:
745   Developmental Invariance and Change in the Evolution of Archosaur Limbs *Journal of*
746   *Morphology* **203**, 151-164 (1990).
747

Data availability

The RNA-sequencing data for mouse, alligator and *Anolis* (including counts of mapped reads), is

available on Gene Expression Omnibus (GEO) repository (*https://www.ncbi.nlm.nih.gov/geo/*)

under accession number GSE108337. Sequencing data for human limb samples is available

through the database of Genotypes and Phenotypes (dbGaP) under study accession number

phs001226.v1.p1. Supplementary data files include (1) unique gene IDs corresponding to all

gene lists described, (2) median gene lengths, (3) bootstrap values for all PCA plots, (4) mapped

reads for chicken. All code used for analyses is available on GitHub

(https://github.com/ThomasAStewart/digit_identity_project) and available upon request.

Acknowledgements

We thank the Wagner lab members and SE Newman for discussions on data interpretation. The

human embryonic material was provided by the Joint MRC (G0700089)/Wellcome Trust

(GR082557) Human Developmental Biology Resource. Work in the Wagner lab was supported by

the John Templeton Foundation (Integrating Generic and Genetic Explanations of Biological

Phenomena; ID 46919), by National Institutes of Health grant GM094780 (to J.P.N.), and by

National Science Foundation grant 1353691.

Author Contributions

T.A.S. and G.P.W. contributed project design and paper writing. T.A.S., T.J.S., J.L.C., and J.P.N.

contributed data for analysis. T.A.S., C.L., and G.P.W. contributed analyses of data.

Correspondence and requests for materials should be addressed to tomstewart@uchicago.edu

and gunter.wagner@yale.edu

Additional Information

Extended Data accompanies this paper.

Figure 1 | Pentadactyl amniote limbs have disparate patterns of genetic differentiation of digits. (a) Photographs of right forelimbs at the stages sampled, dorsal perspective. Analyses of limb patterning genes show that in (b) mouse and (c) alligator, replicates of each digit form clusters, indicating that the digits have distinct gene expression profiles. By contrast, (c) *Anolis* digits do not show clear differentiation of gene expression profiles.

Figure 2 | Differential expression analyses suggest homogeneity among *Anolis* digits. Histograms show the distribution of *p* values from expression analyses of adjacent digits. In *Anolis, p* value distributions that are close to uniform, indicating very weak genetic differentiation of adjacent fingers. Mouse and alligator, on the other hand, generally show strongly biased *p* value distributions.The number of genes that are identified as differentially expressed at a FDR threshold of 0.05 are noted in each panel as "genes," and the number of transcription factors among these are noted in each panel as "TF."

Figure 3 | Few genes are differentially expressed at the same position between pentadactyl limbs. Venn diagrams of one-to-one orthologous transcription factors genes for mouse, alligator, and *Anolis* that were identified as differentially expressed between adjacent digits with a FDR threshold of 0.05.

Figure 4 | D1 has a unique, conserved gene expression profile across amniotes. (a) Venn diagrams of one-to-one orthologous transcription factors genes of mouse, alligator, and chicken that are differentially expressed between adjacent digits (FDR threshold of 0.05). (b) Heatmap showing relative expression of genes in D2 and D1. Human transcriptomic data provides additional support for the hypothesis that D1 has a conserved developmental identity across amniotes. (c) Expression levels of *T-box* family genes across the autopod. Transcript per million (TPM) values presented in panels b and c calculated from the gene list of one-to-one orthologous genes between mouse, alligator, *Anolis*, chicken, and human.

Figure 5 | A conserved core set of digit patterning genes in amniotes. (a) ANOVA comparing all combination of digits within limbs. The number of genes identified as differentially expressed at a FDR threshold of 0.05 are noted in each panel as 'genes.' The number of transcription factors among these are noted as 'TF.' (b) Venn diagram showing one-to-one orthologous transcription factor genes that are differentially expressed across mouse, alligator, and chicken limbs. (c) HCA and PCA show these genes can recover patterns of digit correspondence similar to analyses of limb patterning genes and transcription factors.

Figure 6 | Digits D2, D3, D4 of the chicken forelimb correspond to D1, D3, and D4 of other limbs. This pattern is observed by (a) HCA and PCA of CDEGs for digits of chicken wing and hindlimb. (b) Expression levels of individual transcription factor genes in chicken. (c) *In situ* hybridization of alligator embryos confirms that the position of differential expression of transcription factors can be conserved between fore- and hindlimbs that differ in digit number. Arrowheads indicate distal-most expression along a developing digit. (d) Projection of chicken digit data upon PCA of CDEGs for mouse, alligator, and *Anolis*. TPM values shown in panels b and c calculated from the gene list of one-to-one orthologous genes between mouse, alligator, *Anolis*, chicken, and human.

Figure 7 | The evolution of digit gene expression in amniotes is highly dynamic. (a) A phylogeny of the taxa sampled by this study and schematic graphs summarizing the relative similarity of digits within limbs, where connections and line thickness reflect degree of similarity in gene expression profiles. (b) Schematic of a limited frame shift model for evolutionary origin of the avian wing in which the developmental identity of D1 was translocated to position D2.

Extended Data Figure 1 | Analyses of full transcriptomes do not show clustering of samples by digit. (a) In mouse, only D1 forms a cluster in HCA, indicating that similarity can be diagnosed when the expression of all genes is considered; more-posterior digits do not form clusters of replicates. (b) In alligator, a cluster with low bootstrap support is observed for D4 samples. Stage 18 and stage 19.5 samples are differentiated as filled or outlined points in PCA plot. (c) In *Anolis,* samples do not reveal stable clustering of digit replicates.

Extended Data Figure 2 | Analyses of all transcription factors show patterns of sample clustering consistent with analyses of limb patterning genes. (a) In mouse, D1 is markedly distinct from the posterior digits. (b) In alligator, two major clusters of digits are observed: (D1, D2, D3)(D4, D5). Stage 18 and stage 19.5 samples are differentiated as filled or outlined points in PCA plot. (c) In *Anolis*, digits do not show gene expression differentiation when all transcription factors are considered. Broadly, adjusted uncertainty values recovered by HCA are lower for a given cluster (*e.g.*, alligator D4) than when limb patterning genes are analyzed.

Extended Data Figure 3 | Clustering analyses of chicken hindlimb digit transcriptomes. PCA, heatmap of Pearson's correlations and HCA of (a) the full transcriptome and (b) transcription factor genes, (c) and known limb patterning genes. Each digit is represented by two data points, which correspond to one sample from stage 28 and another from stage 31. Clustering analyses show stable gene expression profiles for individual digits over this developmental window, even after phalangeal number has been established.

Extended Data Figure 4 | Differential expression analyses of adjacent chicken hindlimb digits. (a) *p* value distrubitions for pairwise tests of differential expression. The number of genes that are identified as differentially expressed at a FDR threshold of 0.05 are noted in each panel as "genes" and of these the number of transcription factors are noted in each panel as "TF." (b) Venn diagrams showing the genes that are one-to-one orthologous transcription factors and differentially expressed in each species to a FDR threshold of 0.05.

Extended Data Figure 5 | Comparing fold change of differentially expressed genes to random genes. Fold change of genes differentially expressed between adjacent digits for one species was compared to those of orthologous genes in other species and also to randomly selected genes of similar expression level for the other species. Comparisons were made among the pentadactyl limbs (a-d), and also considering chicken, rather than *Anolis* (e-g). Broadly, genes differentially expressed between D1 and D2 behave consistently between species. Among more posterior digits there is limited evidence for conserved behavior. The genes number of genes recovered as differentially expressed at each position for each species are reported in Fig 3. and Extended Data Fig. 4 b. Vertical lines in each plot represent the mean values of correlation among comparisons between the sets of orthologous or random genes.

Extended Data Figure 6 | Correspondence of chicken forelimb and hindlimb digits. (a) PCA and HCA of full transcriptomes does not reveal correspondence between digits of the fore- and hindlimb. However, analyses of (b) all transcription factors and (c) limb patterning genes show that the three digits in the avian wing correspond to hindlimb digits D1, D3, and D4. Stage 28 and 31 samples are differentiated as filled or outlined points in PCA plot.

Extended Data Figure 7 | Evaluating how differential expression analyses are impacted by variance of replicates. (a) Pearson's correlation values of sample replicates showing that in *Anolis* replicates are less highly correlated than the replicates of mouse and alligator; however, for the digits D1, D2, D4 and D5 two of the three replicates are correlated with values comparable to the other species (>0.99). Therefore, to assess whether variance in *Anolis* was biasing our analyses, tests of differential expression were replicated using only the two most-highly correlated replicates of each digit. (b) Venn diagrams of one-to-one orthologous transcription factors genes identified as differentially expressed between adjacent digits with a FDR threshold of 0.05 when differential expression analyses considered only the two most-highly correlated samples of each digit.

Extended Data Figure 8 | Mean dispersion values of the digit transcriptomes. Values calculated by edgeR showing that although pedigree likely impacts tests of differential gene expression, it does not explain the homogeneity of the *Anolis* digits.

Chicken forelimb and hindlimb

- Published data set: Wang et al. 2011
- Sampled at st.28 and 31
- One sample per digit for each time point
- Each sample contained
  20 pooled digits

Alligator forelimb

- Sampled at st.18 and 19.5
- 3 replicates per digit
- Each replicate contained
  5 pooled digits

Anolis **forelimb**

- Sampled at st.10
- 3 replicates per digit
- Each replicate contained
  5 pooled digits

Mouse forelimb

- Sampled at st.18 (E13.5)
- 3 replicates per digit
- Each replicate
  contained 5 pooled digits

Human forelimb and hindlimb

- Sampled at st.18
- 2 replicates per sample type, designated a and b when plotted on Fig. 2 b.
- One sample of each type from Individual 1; one sample of each forelimb type from Individual 2; one sample of each hindlimb type from Individual 3

Archosauria

Sauropsida

Mammalia

Extended Data Figure 9 | Summary of the data sampled. Illustrations are of the adult skeletons from the dorsal perspective, anterior is left. 'Replicates' refer to biological replicates.

Extended Data Figure 10 | Differential expression analyses of adjacent digits without PC1 correction of *Anolis* data. Venn diagrams of one-to-one orthologous transcription factors genes for mouse, alligator, and *Anolis* that were identified as differentially expressed between adjacent digits with a FDR threshold of 0.05.

Extended Data Table 1 | Human and mouse phenotypes have been described for 28 of the 49 CDEGs

gene	human phenptype or syndrome	mouse phenotypes
Alx1	Camptodactyly [2]	polydatyly [4]
Dlx5	Split hand/foot malformation [1,3]	ectrodactyly, monodactyly, syndactyly [4]
Dlx6		brachydactyly, ectrodactyly, syndactyly [4]
En1		25 phenotypes including adactyly, ectopic digits, polydactyly, syndactyly, truncation of digits, fused phalanges [4]
Ets2	Chitayat syndrome [3]	
Hand1		hypoplastic limb buds [4]
Hand2		19 phenotypes including oligodactyly, polydactyly, abnormal pollex morphology [4]
Hic1	Hand-foot-genital syndrome [1]	abnormal fore- and hindlimb morphology [4]
Hoxa13	Guttmacher syndrome [1,2], Hand-foot-genital syndrome [1,3], Postaxial hand polydactyly [2]	21 phenotypes including brachydactyly, clinodactyly, syndactyly [4]
Hoxd11		7 phenotypes including abnormal phalanx, abnormal and fused carpals [4]
Hoxd12		16 phenotypes including brachyphalangia, brachydactyly, clinodactyly, oligodactyly [4]
Hoxd13	Brachydactyly-syndactyly syndrome [1], VACTERL association, Brachydactyly type E [2]	29 phenotypes including brachydactyly, ectrodactyly, polydactyly, polysyndactyly [4]
Nkx3-2	Spondylo-megaepiphyseal-metaphyseal dysplasia [3]	
Pax9		polydactyly, polysyndactyly [4]
Pitx1	Clubfoot and Lower limb malformations [1], Liebenberg syndrome [3]	11 phenotypes including brachydactyly, oligodactyly, and clubfoot [4]
Runx2	Brachydactyly [1], MDMHB [3], cleidocranial dysplasia (which includes abnormal thumbs and brachydactyly) [4]	10 phenotypes including abnormal phalanx morphology [4]
Sall1	Townes-Brocks Syndrome [1,2,3] , Lenz microphthalmia syndrome [2]	10 phenotypes including oligodactyly, preaxial polydactyly, syndactyly and triphalangia [4]
Satb2	chromosome 2q32-q33 deletion syndrome (which includes clinodactyly archnodactyly, and Talpes equinovarus [4]	
Sox10	kallmann-syndrome [3], Klein-Waardenburg syndrome [3], PCWH syndrome (which includes Pes cavus) [4]	
Tbx2	17q23.1q23.2 microdeletion syndrome [2]	Polydactyly, postaxial polydactyly [4]
Tbx3	Ulnar-mammary syndrome [1], Limb-mammary syndrome [2], post-axial polydactyly [2]	16 phenotypes including oligodactyly [4]

Tbx5	Holt-Oram syndrome [1,2]	10 phenotypes including abnormal phalanx morphology [4]
Tbx15	Cousin syndrome (which includes brachydactyly) [1,2]	9 phenotypes including abnormal phalanx morphology [4]
Tfap2b	Char syndrome [1,2]	Polydactyly, postaxial polydactyly [4]
Trps1	Trichorhinophalangeal syndrome type 1 [1,2], type 2 [2], and type 3 [2]	
Twist2	Ablepharon macrostomia syndrome [2]	
Zic2		6 phenotypes mostly restricted to carpals [4]
Zic3	VACTERL association [2], Radial abnormalities [2], aplasia/Hypoplasia of the radius [2]	

References

[1] Stenson *et al*. The Human Gene Mutation Database (HGMD®): 2003 Update. Hum. Mutat. **21**, 577-581 (2003)

[2] NIH Genetic and Rare Disease Information Center. http://rarediseases.info.nih.gov.

[3] National Library of Medicine (US). Genetics Home Reference [Internet]. https://ghr.nlm.nih.gov (2013).

[4] Blake JA *et al*. Mouse Genome Database (MGD)-2017: community knowledge resource for the laboratory mouse. Nucl. Acids Res. 4, D723-D729 (2017).