1   **_E. coli_ translation strategies differ across nutrient conditions**

4   **Authors:** Sophia Hsin-Jung Li[1], Zhiyuan Li[2], Junyoung O. Park[3,4], Christopher G. King[5], Joshua
5   D. Rabinowitz[3,6], Ned S. Wingreen*[1,3], and Zemer Gitai*[1]

Affiliations:

[1]Department of Molecular Biology,

[2]Princeton Center for Theoretical Science,

[3]Lewis-Sigler Institute for Integrative Genomics,

[4]Department of Chemical and Biological Engineering,

[5]Department of Physics, and

[6]Department of Chemistry, Princeton University, Princeton, NJ, USA.

*Correspondence to: wingreen@princeton.edu, zgitai@princeton.edu

Abstract

For cells to grow faster they must increase their protein production rate. Microorganisms have traditionally been thought to accomplish this increase by producing more ribosomes to enhance protein synthesis capacity, leading to the linear relationship between ribosome level and growth rate observed under most growth conditions previously examined. Past studies have suggested that this linear relationship represents an optimal resource allocation strategy for each growth rate, independent of any specific nutrient state. Here we investigate protein production strategies in continuous cultures limited for carbon, nitrogen, and phosphate, which differentially impact substrate supply for protein versus nucleic acid metabolism. Unexpectedly, we find that at slow growth rates, *E. coli* achieves the same protein production rate using three different strategies under the three different nutrient limitations. Upon phosphate (P) limitation, translation is slow due to a particularly low abundance of ribosomes, which are RNA-rich and thus particularly costly for phosphorous-limited cells. In nitrogen (N) limitation, translation is slowed by limited glutamine and stalling at glutamine codons, resulting is slow elongation. In carbon (C) limitation, translation is slowed by accumulation of inactive ribosomes not bound to mRNA. These extra ribosomes enable rapid growth acceleration upon nutrient upshift. Thus, bacteria tune ribosome usage across different limiting nutrients to enable balanced nutrient-limited growth while also preparing for future nutrient upshifts.

Introduction

Resource allocation during growth is a fundamental challenge faced by all cells[1-4]. For example, with a fixed resource budget, cells must balance production of the machinery that makes proteins (ribosomes, tRNAs, translation factors) with the production of the proteins themselves. This balance is generally represented by the RNA:protein ratio (R/P ratio)[5]. The R/P ratio captures protein production capacity, as >95% of total RNA is devoted to translation (rRNAs and tRNAs[5,6]). In single-cell organisms like *E. coli*, previous studies demonstrated that

there is a linear relationship between R/P ratio and growth rate, with faster growth rates

requiring more protein production capacity and therefore higher R/P ratios[5,7,8]. Production of

ribosomes is costly as each contains 52 protein subunits and three large rRNAs[9,10]; hence, it is

advantageous for the cell to saturate ribosomes with substrates. In this efficient ribosome

scenario, the ribosome level should be fixed and independent of nutrient conditions for any

growth rate, with the only way to increase protein synthesis rate being to increase the number of

ribosomes[1,5,11]. One surprise for such a seemingly optimized system is that multiple studies

have demonstrated that at slow growth rates *E. coli* accumulates inactive ribosomes[12]. There

are two possible explanations for the presence of inactive ribosomes. First, it is possible that *E.*

coli translation is constrained in such a way that it cannot function when ribosome levels drop

too low[12]. Alternatively, *E. coli* could regulate ribosome production independently of growth rate.

Here we settle this debate by showing that *E. coli* ribosome production and usage differ across

nutrient conditions.

Results

Phosphate-limited cells achieve the same growth rate with fewer ribosomes than Carbon-

or Nitrogen-limited cells

To determine the generality of the relationship between growth rate and ribosome

content, we examined how the R/P ratio changes as a function of growth upon different nutrient

limitations. We measured R/P ratios in *E. coli* under glucose- (C, carbon), ammonia- (N,

nitrogen), and phosphate- (P, phosphorous) limitations over a range of different growth rates in

chemostats (Figure 1A). Surprisingly, P-limited cells consistently exhibited lower R/P ratios than

C-limited or N-limited cells, with a roughly 2-fold difference at the slowest growth rate tested (0.1

h^{-1}, Figure 1B). Measured protein levels were similar in all cells regardless of growth rate or

nutrient limitation (Figure S1A-B), indicating that the changes in the R/P ratios reflect changes in

ribosome availability.

The finding that *E. coli* can grow at the same rate with a lower R/P ratio implies that P-

limited cells make the same amount of protein with fewer ribosomes, i.e. use ribosomes more

efficiently. Thus, C/N-limited *E. coli* cells do not use ribosomes with optimal efficiency and their

supply of "extra" ribosomes cannot be ascribed to requirements for productive translation. Since

RNA accounts for two-thirds of the mass of bacterial ribosomes[9,14], producing fewer ribosomes

upon phosphate-limitation makes sense as a way for cells to deal with a limitation that

preferentially reduces an elemental substrate needed to make RNA but not protein. This effect

may be a direct consequence of low phosphate resulting in limited nucleotide pools and thus

slow RNA synthesis, as deletion of phosphate sensing or storage systems, phoB[15] or ppk[16], did

not alter the R/P ratio (Figure S1C-D).

N-limited ribosomes translate slowly while C-limited cells accumulate more mRNA-free

ribosomes

Why do C/N-limited cells accumulate so many ribosomes if P-limited cells can achieve

the same protein synthesis rates with fewer ribosomes? One possibility is that the ribosomes in

these cells translate slowly. We thus used a *lacZ* induction assay to compare the translation

elongation rates of slow-growing C-, N-, and P-limited cells (0.1 h^{-1})[17,18]. We observed a reduced

elongation rate in N-limited cells compared to C- and P-limited cells but no difference between

C- and P-limited cells (Figure 1C). Thus, N-limited cells may need higher ribosome numbers to

compensate for their slow translation elongation, but something else must explain the elevated

ribosome numbers in C-limited cells.

To characterize ribosome pools we performed polysome profiling, which separates

ribosome species using a sucrose gradient[19]. Regardless of the growth condition, all cells

exhibited similar fractional pools of dissociated 30S and 50S subunits (Figure 1D). In contrast,

the fraction of 70S monosomes was significantly larger in C/N-limited cells than in P-limited cells

(Figure 1D). Since growth rate is proportional to protein synthesis rate, growth rate can be

estimated by the product of the number of active ribosomes and the translation elongation rate.

However, using the assumption that all 70S monosomes are active yielded very different growth

rate estimates for C-, N-, and P-limited cells (Figure 1F and SI), which is inconsistent with the

fact that these cells are growing at the same rate and have the same protein content. This

inconsistency suggested that a fraction of the 70S ribosomes may not be active.

The mass of a single mRNA is small relative to the mass of a ribosome, such that 70S

monosomes could represent either mRNAs with only one ribosome per transcript or inactive

"free" ribosomes that are not associated with an mRNA. To distinguish free and mRNA-bound

70S monosomes, we utilized their differential sensitivity to high potassium levels (170 mM)[20].

High potassium causes free ribosomes to shift to a lower density but does not shift the density

of mRNA-bound monosomes[20]. We thus designed a high-resolution "free-ribosome profiling"

method to resolve this density shift (Figure S2). As controls, we confirmed that our assay

detects the potassium-dependent shift of 70S monosomes induced by puromycin (Figure S2),

which releases elongating ribosomes from their associated mRNAs[21]. There was no potassium-

dependent shift detected in fast-growing cells that lack free ribosomes (Figure S2).

By combining traditional and free-ribosome polysome profiling, we quantified the relative

fractions of all ribosome species in slow-growing C-, N-, and P-limited *E. coli*. The fraction of

free monosomes was roughly 3-fold greater in the C-limited cells than in the P-limited cells,

while the fraction of mRNA-bound monosomes remained relatively constant across nutrient

limitations (Figure 1E). The accumulation of free 70S monosomes in C- and N-limited cells

appears to be independent of a previously-described RaiA-dependent mechanism for ribosome

storage as deletion of *raiA* had no impact on R/P ratios or polysome profiles (Figure S3).

Importantly, revising our protein synthesis rate estimates to account for the fraction of inactive

70S monosomes yielded similar values for all cells, regardless of nutrient limitation (Figure 1F).

These results both validate our experimental measurements and suggest that in different

nutrient states, *E. coli* differentially tune ribosome number, translation rate, and active fraction to

produce proteins at the same rates (Figure 2A).

Quantitative modeling describes three different strategies of ribosome dynamics to

achieve the same protein production rate

To better understand nutrient-dependent ribosome dynamics, we probed translation in

slow-growing C-, N-, and P-limited cells by ribosome profiling. Analysis of ribosome densities as

a function of distance from the start and stop codons revealed higher ribosome occupancy near

the start codon (Figure 2B). Ribosome density thereafter was similar, with no decrease in

ribosome density between the first and second halves of genes (Figures 2B and S4A),

suggesting that there is little to no aborted translation after the first few codons. We quantified

ribosome dynamics by building a simple macroscopic model (Figure 2C, S4B-E and details in

SI). In this model, unbound ribosomes (R_u) can bind to an mRNA with a free ribosomal binding

site (with rate constant k_f) to become initiating ribosomes (R_i). The initiating ribosomes can

proceed to elongation (with rate constant k_p) to become working ribosomes (R_w), or can abort

translation (with rate constant k_r). Working ribosomes elongate to finish translation and become

unbound (with rate constant k_{el}/N_{aa}), where N_{aa} is the length of an average protein in amino

acids. We defined the fraction of ribosomes bound to the first 10 codons as initiating ribosomes

since the ribosome footprint size is ~10 codons[22,23]. To validate our model we used it to

calculate elongation rates, which closely agreed with those we measured experimentally (Figure

S5A).

Analysis of our model indicated that the system can be characterized by two main

dimensionless parameters: the "relative proceeding rate", defined as the ratio between the rate

of ribosomes proceeding from initiation to elongation (k_p, s^{-1}) and the rate of elongation (k_{el}/N_{aa},

s^{-1}); and the "saturation parameter" ($R_t/(K_m+R_t)$), reflecting the degree of saturation of ribosome

binding sites on mRNAs, where R_t is the total ribosome number and $K_m = (k_r + k_p)/k_f$ (Figures

2D-E and S5B-C). Fitting the measured ribosome densities, translation elongation rates, and

pool sizes of ribosome species to the macroscopic model revealed that P-limited cells have the

highest relative proceeding rate while C-limited cells have the lowest saturation parameter, and

N-limited cells have intermediate values for both parameters (Figures 2D-E and S5B-H). Thus,

C-, N-, and P-limited cells produce proteins at the same rate using three different strategies: P-

limited cells have few ribosomes that are mostly active and elongate rapidly, N-limited cells

have more ribosomes but fewer are active and they elongate slowly, and C-limited cells have

many ribosomes, which elongate rapidly, but even fewer are bound to mRNA (Figure 2F and

S5I).

N-limited ribosomal regulation is mediated by RelA, the ppGpp alarmone synthase

Insight into the molecular basis of nutrient-specific ribosome regulation came from

analysis of the codon occupancies in our ribosome profiling data. Codon-specific ribosome

stalling leads to increased codon occupancy and is a hallmark of insufficient pools of the

corresponding charged tRNAs. P-limited cells exhibited no elevated codon frequencies,

consistent with these cells' efficient ribosome usage (Figure 3A). In contrast, both C- and N-

limited cells exhibited significant codon-specific stalling. Under N-limitation, ribosomes stalled at

both of the two glutamine-encoding codons, which together account for 4.4% of all predicted

ORF codons in *E. coli*. This result is consistent with previous studies indicating that glutamine is

the most strongly-depleted amino acid pool upon N-limitation and serves as an intracellular

sensor for extracellular nitrogen levels[24-26]. Upon C-limitation, we observed elevated occupancy

of the Leu-CUA codon, which was surprising as there is no known intracellular carbon sensor

that controls translation. *E. coli* has six leucine codons decoded by five leucine tRNA

species[27,28]. Leu-CUA is the rarest Leu codon, accounting for only 0.4% of all predicted ORF

codons.

In addition to clarifying the link between metabolism and translation for N-limitation, the

observation that ribosomes stall at specific codons upon C/N-limitation suggested a molecular

mechanism for nutrient-specific translation regulation (Figure 3A). In bacteria, insufficient

charged tRNA pools activate the stringent response to induce accumulation of the cellular

alarmone, ppGpp[30,31]. ppGpp is known to regulate rRNA transcription *in vivo*. Its role in

translation is less well-understood but ppGpp has been shown *in vitro* to inhibit translation

factors such as EF-Tu and IF2 by competing with GTP[32,33]. To test how ppGpp accumulation

might affect translation *in vivo*, we induced ppGpp synthesis by treating batch-grown *E. coli* with

serine hydroxamate (SHX). SHX is a serine analog that competitively inhibits serine tRNA

synthetase to yield uncharged serine-tRNA and thereby activate the RelA ppGpp synthase[34].

SHX treatment increased the pool of mRNA-free 70S monosomes, and this effect was

completely dependent on *relA* (Figures S6A-B). Thus, inducing ppGpp by activating RelA alters

translation by increasing the fraction of inactive free 70S ribosomes.

RelA is primarily required for the accumulation of ppGpp upon N-limitation but not C-

limitation[31]. Consistently, deletion of *relA* had little impact on the accumulation of free ribosomes

upon C-limitation, but significantly reduced free ribosome pools upon N-limitation to levels

similar to those observed upon P-limitation (Figures 3B and S6C). We could not probe the role

of ppGpp in C-limited ribosome accumulation because unlike N-limitation, C-limitation elevates

ppGpp through *spoT*, which is essential[35]. N-limited *ΔrelA* cells also contained more polysomes

than wild type (Figure 3B), suggesting that these cells had a higher fraction of elongating

ribosomes. This result was initially confusing because the wild type and *ΔrelA* N-limited cells

were grown at the same growth rate and maintained the same R/P ratio (Figure S6D). We thus

measured the rate of translation elongation and found that while N-limited *ΔrelA* cells have

higher fractions of translating ribosomes, their ribosomes elongate more slowly (Figure 3C),

resulting in the same rate of protein production as wild type.

To understand how RelA influences ribosome dynamics we performed ribosome

profiling. N-limited *ΔrelA* cells exhibited more pronounced ribosome stalling at both glutamine

codons than wild type N-limited cells, while *relA* deletion had no effect on P- or C-limited cells

(Figures 3D and S7). Fitting the *ΔrelA* cells measurements to our ribosome dynamics model

revealed that *ΔrelA* cells specifically increase the relative proceeding rate under N-limitation, but

display no effect on relative proceeding rate under P- or C-limitation and no effect on the

saturation parameter in any condition (Figures 3E and S5B-C). These results suggest that,

consistent with its known effects on IF-2 *in vitro*, in N-limited cells RelA functions to restrict the

transition from translational initiation to elongation. Thus, in the absence of RelA, more

ribosomes attempt to elongate, which exacerbates the depletion of charged tRNA pools, leading

to increased stalling and a slower translational elongation rate.

Extra ribosomes may facilitate growth acceleration upon nutrient upshift

While cells can modulate different aspects of ribosome dynamics to achieve the same

protein production rate, what benefits might be served by the inefficient translation system used

by C-limited cells where many ribosomes are inactive? Since free ribosomes accumulate the

most at the slowest growth rates, we hypothesized that our findings could reflect a trade-off

between steady-state growth rate and the ability to respond to a fluctuating environment. Such

transitions could include rapidly and safely slowing growth when nutrients become depleted and

rapidly increasing growth rate when nutrients are replenished. In this scenario, cells may benefit

more by optimizing their ability to rapidly utilize new nutrients, for example to outcompete their

neighbors or maximally utilize a transient pulse of nutrients, than by optimizing steady-state

growth rate when nutrient levels are low.

We built a mathematical model of cellular resource allocation to predict the growth

dynamics upon upshift. This model supported the hypothesis that the larger free ribosome pools

of C/N-limited cells should enable them to increase their growth rates more quickly than P-

limited cells upon nutrient upshift (Figures 4A, S8 and details in SI)[4,36]. We experimentally tested

this prediction by measuring the growth rates of slow-growing (0.1 h[-1]) *E. coli* immediately after

being shifted to rich media (LB + 0.4% glucose) [37]. As predicted, C/N-limited cells increased

their growth rates significantly faster than P-limited cells (Figure 4B). Thus, the distinct

translation strategies employed under different nutrient conditions may represent nutrient-

specific adaptations, with P-limited cells optimizing for current steady-state growth under slow

growth conditions, and C- and N-limited cells favoring the ability to rapidly recover growth

(Figure 4C).

Discussion

In previous studies, *E. coli* were found to vary R/P ratio with growth rate independently of

the specific nutrient limitation used to produce a given growth rate[5,8]. Meanwhile, transcriptional

analysis in *Saccharomyces cerevisiae* suggested that the primary determinant of the response

to a wide range of stresses was the cellular growth rate rather than the specific stressor[38].

Together, these studies suggested that the primary regulator of microbial physiology is growth

rate. However, our findings demonstrate that at the same growth rate, *E. coli* exhibit significantly

different translation strategies across nutrient limitations; at the lowest growth rate tested, P-

limited cells produced the same amount of protein with roughly half as many ribosomes as C/N-

limited cells. P-limited cells also exhibited smaller inactive 70S monosome pools and higher

relative proceeding rates than C/N-limited cells. Furthermore, while C/N-limited cells have

similar ribosome numbers, they also display differences in free ribosome pools, translational

elongation rates, and sensitivity to the loss of the ppGpp synthase RelA. Thus, our results

suggest that the extra ribosomes of C/N-limited cells do not reflect essential constraints, but

rather reflect a selectively beneficial adaptation.

Our findings also implicate ppGpp as a key mediator of ribosome activity that specifically

modulates the transition from translation initiation to elongation. ppGpp is a well-characterized

cellular alarmone that senses translational activity and inhibits rRNA transcription[31,39]. However,

ppGpp can competitively inhibit GTP-dependent enzymes, including the translation initiation

factor IF-2 that is required for the transition to elongation[40]. Our data thus provide *in vivo* support

for a previous *in vitro* study demonstrating that ppGpp inhibits IF-2 function[33]. Since ppGpp can

also affect other GTP-binding proteins involved in translation[32,41], future studies on bacterial

translation will confirm exactly how ppGpp interferes with ribosome function.

In native environments such as the mammalian gut, bacteria such as *E. coli* are faced

with feast and famine cycles induced by feeding cycles[42]. Here we show that the linear

relationship between R/P ratio and growth rate in *E. coli* does not reflect the optimization of

steady-state growth rate but, instead, reflects the ability of cells with higher inactive ribosome

pools to rapidly accelerate growth upon nutrient repletion. These findings suggest that *E. coli*

may improve their fitness by sacrificing their maximal growth rate in nutrient-poor periods in

return for the ability to respond to a changing environment, including rapidly accelerating growth

in nutrient-rich periods. Our macroscopic model highlights how cells can tune total ribosome

number, the fraction of working ribosomes, and the rate of translational elongation to achieve

the same total protein production rate while balancing other constraints such as reduced amino

acid availability or the need to rapidly accelerate growth. Future studies will address the

consequences of adaptation strategies in dynamic conditions, for example the generality of

bacteria optimizing growth rate transitions at the expense of steady-state growth[43], but this

strategy could explain recent reports of the sub-optimality of protein allocation for *E. coli* in the

presence of poor carbon sources[44], as well as the sub-optimal expression levels of essential

genes in *B. subtilis*[45].

Author Contributions
S.H.L., J.O.P., J.D.R., N.S.W, and Z.G. designed the experiments. S.H.L., J.O.P., and C.K.
performed experiments. Z.L. and N.S.W. constructed the mathematical models. S.H.L.

conducted computational analysis of sequencing data. S.H.L, Z.L., N.S.W, and Z.G. wrote the
paper with assistance from J.D.R.

Acknowledgments:

We thank the members of the Gitai, Wingreen, and Rabinowitz labs for helpful discussions. We
thank Gene-Wei Li for his support for the ribosome profiling experiments. We thank the
Microarray Core Facility at the Lewis-Sigler Institute (Daniel Sanchez, Jennifer M. Miller, Jessica
Wiggins, and Wei Wang) for RNA-Seq sample processing and sequencing and the Princeton
Proteomics Core Facility (Henry Shwe and Tharan Srikumar) for ribosome profiling sample
processing. Lance Parsons provided helpful technical support for bioinformatics analysis of the
sequencing data. We thank the Botstein lab, particularly Sandy Silverman, for chemostat
operation support and all former members for discussions relating to microbial growth.

The ribosome sequencing data that support the findings of this paper will be deposited in
GenBank with the accession code at publication.

This work was supported by grants from the NIH (DP1AI124669 and R01GM082938).

References and Notes:

1.   You, C. *et al.* Coordination of bacterial proteome with metabolism by cyclic AMP
signalling. **500,** 301–306 (2013).
2.   Li, G.-W., Burkhardt, D., Gross, C. & Weissman, J. S. Quantifying absolute protein
synthesis rates reveals principles underlying allocation of cellular resources. *Cell*
157, 624–635 (2014).
3.   Scott, M., Klumpp, S., Mateescu, E. M. & Hwa, T. Emergence of robust growth laws
from optimal regulation of ribosome synthesis. *Mol. Syst. Biol.* **10,** 747–747 (2014).
4.   Giordano, N., Mairet, F., Gouzé, J.-L., Geiselmann, J. & de Jong, H. Dynamical
Allocation of Cellular Resources as an Optimal Control Problem: Novel Insights into
Microbial Growth Strategies. *PLoS Comput. Biol.* **12,** e1004802 (2016).
5.   Bremer, H. & Dennis, P. P. Modulation of Chemical Composition and Other
Parameters of the Cell at Different Exponential Growth Rates. *EcoSal* 1–56 (2012).
6.   Baracchini, E. & Bremer, H. Determination of synthesis rate and lifetime of bacterial
mRNAs. *Anal. Biochem.* **167,** 245–260 (1987).
7.   Schaechter, M., Maaløe, O. & Kjeldgaard, N. O. Dependency on Medium and
Temperature of Cell Size and Chemical Composition during Balanced Growth of
Salmonella typhimurium. *Microbiology* **19,** 592–606 (1958).
8.   Scott, M., Gunderson, C. W., Mateescu, E. M., Zhang, Z. & Hwa, T.
Interdependence of cell growth and gene expression: origins and consequences.
Science **330,** 1099–1102 (2010).
9.   Yamamoto, T., Izumi, S. & Gekko, K. Mass spectrometry of hydrogen/deuterium
exchange in 70S ribosomal proteins from E. coli. *FEBS Lett.* **580,** 3638–3642
(2006).
10.   Nomura, M. & Gourse, R. Regulation of the synthesis of ribosomes and ribosomal
components. *Annual review of …* (1984).
11.   Maaløe, O. & Kjeldgaard, N. O. *Control of Macromolecular Synthesis, WA*
Beniamin. (Inc., 1966).

12. Dai, X. *et al.* Reduction of translating ribosomes enables Escherichia coli to maintain elongation rates during slow growth. *Nat Microbiol* **2,** 16231 (2016).

13. Klumpp, S., Scott, M., Pedersen, S. & Hwa, T. Molecular crowding limits translation and cell growth. *Proc. Natl. Acad. Sci. U.S.A.* **110,** 16754–16759 (2013).

14. Lewin, B. *Genes VIII*. (Pearson education, Inc, 2003).

15. Wanner, B. L. Gene regulation by phosphate in enteric bacteria. *Journal of Cellular Biochemistry* **51,** 47–54 (1993).

16. Ahn, K. & Kornberg, A. Polyphosphate kinase from Escherichia coli. Purification and demonstration of a phosphoenzyme intermediate. *J. Biol. Chem.* **265,** 11734–11739 (1990).

17. Schleif, R., Hess, W., Finkelstein, S. & Ellis, D. Induction kinetics of the L-arabinose operon of Escherichia coli. *J. Bacteriol.* **115,** 9–14 (1973).

18. Dalbow, D. G. & Young, R. Synthesis time of β-galactosidase in Escherichia coli B/r as a function of growth rate. *Biochemical Journal* **150,** 13–20 (1975).

19. Qin, D. & Fredrick, K. Analysis of polysomes from bacteria. *Meth. Enzymol.* **530,** 159–172 (2013).

20. Beller, R. J. & Lubsen, N. H. Effect of polypeptide chain length on dissociation of ribosomal complexes. *Biochemistry* **11,** 3271–3276 (1972).

21. Nathans, D. Puromycin inhibition of protein synthesis: incorporation of puromycin into peptide chains. *Proc. Natl. Acad. Sci. U.S.A.* **51,** 585–592 (1964).

22. Steitz, J. A. Polypeptide Chain Initiation: Nucleotide Sequences of the Three Ribosomal Binding Sites in Bacteriophage R17 RNA. **224,** 957–964 (1969).

23. Ingolia, N. T., Ghaemmaghami, S., Newman, J. R. S. & Weissman, J. S. Genome-wide analysis in vivo of translation with nucleotide resolution using ribosome profiling. *Science* **324,** 218–223 (2009).

24. Doucette, C. D., Schwab, D. J., Wingreen, N. S. & Rabinowitz, J. D. α-Ketoglutarate coordinates carbon and nitrogen utilization via enzyme I inhibition. *Nature Chemical Biology* **7,** 894–901 (2011).

25. Yuan, J. *et al.* Metabolomics-driven quantitative analysis of ammonia assimilation in E. coli. *Mol. Syst. Biol.* **5,** 302 (2009).

26. Ikeda, T. P., Shauger, A. E. & Kustu, S. Salmonella typhimuriumApparently Perceives External Nitrogen Limitation as Internal Glutamine Limitation. *Journal of Molecular Biology* **259,** 589–607 (1996).

27. Subramaniam, A. R., Pan, T. & Cluzel, P. Environmental perturbations lift the degeneracy of the genetic code to regulate protein levels in bacteria. *Proc. Natl. Acad. Sci. U.S.A.* **110,** 2419–2424 (2013).

28. Sørensen, M. A. *et al.* Over Expression of a tRNALeu Isoacceptor Changes Charging Pattern of Leucine tRNAs and Reveals New Codon Reading. *Journal of Molecular Biology* **354,** 16–24 (2005).

29. Elf, J., Nilsson, D., Tenson, T. & Ehrenberg, M. Selective charging of tRNA isoacceptors explains patterns of codon usage. *Science* **300,** 1718–1722 (2003).

30. Hauryliuk, V., Atkinson, G. C., Murakami, K. S., Tenson, T. & Gerdes, K. Recent functional insights into the role of (p)ppGpp in bacterial physiology. *Nat Rev Micro* **13,** 298–309 (2015).

31. Potrykus, K. & Cashel, M. (p)ppGpp: Still Magical? *. *Annu. Rev. Microbiol.* **62,** 35–51 (2008).

32. Mitkevich, V. A. *et al.* Thermodynamic characterization of ppGpp binding to EF-G or IF2 and of initiator tRNA binding to free IF2 in the presence of GDP, GTP, or ppGpp. *Journal of Molecular Biology* **402,** 838–846 (2010).

33. Milon, P. *et al.* The nucleotide-binding site of bacterial translation initiation factor 2 (IF2) as a metabolic sensor. *Proc. Natl. Acad. Sci. U.S.A.* **103,** 13962–13967

363   (2006).

34.   Jiang, M., Sullivan, S. M., Wout, P. K. & Maddock, J. R. G-protein control of the ribosome-associated stress response protein SpoT. *J. Bacteriol.* **189,** 6140–6147 (2007).

35.   Gentry, D. R. & Cashel, M. Mutational analysis of the Escherichia coli spoT gene identifies distinct but overlapping regions involved in ppGpp synthesis and degradation. *Mol. Microbiol.* **19,** 1373–1384 (1996).

36.   Pavlov, M. Y. & Ehrenberg, M. Optimal control of gene expression for fast proteome adaptation to environmental change. *Proc. Natl. Acad. Sci. U.S.A.* **110,** 20527–20532 (2013).

37.   Kjeldgaard, N. O., Maaløe, O. & Schaechter, M. The Transition Between Different Physiological States During Balanced Growth of Salmonella typhimurium. *Microbiology* **19,** 607–616 (1958).

38.   Brauer, M. J. *et al.* Coordination of growth rate, cell cycle, stress response, and metabolic activity in yeast. *Mol. Biol. Cell* **19,** 352–367 (2008).

39.   Paul, B. J., Ross, W., Gaal, T. & Gourse, R. L. rRNA Transcription in Escherichia coli. *Annu. Rev. Genet.* **38,** 749–770 (2004).

40.   Laursen, B. S., Sørensen, H. P., Mortensen, K. K. & Sperling-Petersen, H. U. Initiation of protein synthesis in bacteria. *Microbiology and Molecular Biology Reviews* **69,** 101–123 (2005).

41.   Rojas, A.-M., Ehrenberg, M. N., Andersson, S. G. E. & Kurland, C. G. ppGpp inhibition of elongation factors Tu, G and Ts during polypeptide synthesis. *Mol. Gen. Genet.* **197,** 36–45 (1984).

42.   Zarrinpar, A., Chaix, A., Yooseph, S. & Panda, S. Diet and Feeding Pattern Affect the Diurnal Dynamics of the Gut Microbiome. *Cell Metabolism* **20,** 1006–1017 (2014).

43.   Fischer, E. & Sauer, U. Large-scale in vivo flux analysis shows rigidity and suboptimal performance of Bacillus subtilis metabolism. *Nat. Genet.* **37,** 636–640 (2005).

44.   Towbin, B. D. *et al.* Optimality and sub-optimality in a bacterial growth law. *Nat Commun* **8,** 14123 (2017).

45.   Peters, J. M. *et al.* A Comprehensive, CRISPR-based Functional Analysis of Essential Genes in Bacteria. *Cell* **165,** 1493–1506 (2016).

46.   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2,** 2006.0008 (2006).

47.   Zhu, M., Dai, X. & Wang, Y.-P. Real time determination of bacterial in vivoribosome translation elongation speed based on LacZα complementation system. *Nucleic Acids Res.* gkw698 (2016). doi:10.1093/nar/gkw698

48.   Dunn, J. G. & Weissman, J. S. Plastid: nucleotide-resolution analysis of next-generation sequencing and genomics data. *BMC Genomics* **17,** 958 (2016).

49.   Woolstenhulme, C. J., Guydosh, N. R., Green, R. & Buskirk, A. R. High-precision analysis of translational pausing by ribosome profiling in bacteria lacking EFP. *Cell Rep* **11,** 13–21 (2015).

Materials and Methods

Cell strains and growth conditions

    Escherichia coli strain NCM3722 was grown in batch or continuous cultures. To achieve different growth rates, different carbon or nitrogen sources were provided in batch culture, whereas dilution rates ranging from 0.1 h^{-1} to 0.7 h^{-1} were used in continuous (chemostat)

cultures. The chemostat (Sixfors, HT) volume was 300mL with oxygen and pH probes to
monitor the culture. pH was maintained at 7.2 +/- 0.1 and the aeration rate was set at 4.5 l/h. 40
mM MOPS media (M2120, Teknova) was used with glucose (0.4%, Sigma G8270), ammonia
(9.5 mM NH_4Cl, Sigma A9434) and phosphate (1.32 mM K_2HPO_4, Sigma P3786) added
separately. For carbon- and nitrogen- limiting media, glucose and ammonia concentrations were
reduced by 5-fold (0.08% and 1.9mM respectively). Phosphorus-limiting medium contains 0.132
mM K_2HPO_4. Δ*relA* mutant was generated by P1 transduction from the KEIO collection[46] into
Escherichia coli strain NCM3722
420
Nutrient upshift growth measurement
Cells from chemosats were mixed with 4X volumes of fresh pre-warmed media and grown
in flasks in a 37°C water bath. Cell growth was monitored every 5 minutes by checking
absorbance at 600nm in a quartz cuvette (Starna, 16.160-Q-10/Z8.5) using a
spectrophotometer (GENESYS™ 20, Thermo Scientific). The LB media (244610, BD) used for
upshift was supplemented with 0.4% glucose.
427
Total RNA measurement
The method for RNA measurement was adapted from You *et al.*[1]. 1.5 mL of cultures were
pelleted by centrifugation for 1 min at 13,000 X g. The pellet was frozen on dry ice and the
supernatant was taken to measure absorbance at 600 nm for cell loss. The pellet was then
washed twice with 0.6M $HClO_4$ and digested with 0.3M KOH for 1 hour at 37°C. The solution
was then precipitated with 3M $HClO_4$ and the supernatant was collected. The pellet was re-
extracted again with 0.5M $HClO_4$. The supernatants were combined and absorbance measured
at 260nm using NanoDrop (ND-1000, NanoDrop). Total RNA concentration was determined by
multiplying the A260 absorbance with 31 (µg RNA/mL) as the extinction coefficient.
437
Total protein measurement
The method for protein measurement is adapted from You *et al.*[1]. 1.5 mL of cell cultures were
pelleted by centrifugation for 1 min at 13,000 X g. Cells were washed with 1 mL MOPS buffer
once, re-suspended in 200 uL water and left on dry ice. All the supernatant was collected and
OD600 was measured for cell loss. To measure the protein content, samples were thawed,
100µL 3M NaOH were added, and the sample was heated for 5 min at 98°C. The samples were
cooled down to RT for 5 min before 300µL 0.1% $CuSO_4$ were added for biuret assay. The
samples were incubated at RT for 5 min and centrifuged at 13,000 X g for 1 min. The
supernatant was collected and absorbance was measured at 555nm for a 200µL sample
volume in a microplate reader (Synergy HT, BioTek) with software Gen 5.0. Proper dilution of
albumin (23209, Thermo) with known concentrations was used to infer the total protein
concentration in the cell.
450
Polysome profiling and quantification of ribosome fraction
200mL of cells were collected from cultures by filtration through 90mm cellulose acetate
membranes with a 0.2 µm pore size (CA029025, Strelitech) at 37°C, scratched with a clean and
pre-warmed stainless steel spatula, and snap-frozen in liquid nitrogen. The whole filtration
procedure did not take more than 2 minutes in order to maintain the original physiological state.
Cell pellets were mixed with 650µL lysis buffer frozen nuggets (20mM Tris-HCl pH 8.0, 10mM
$MgCl_2$, 100mM NH_4Cl, 0.4% Triton X-100, 0.1% NP-40, 1 mM Chloramphenicol, 100 U/mL
RNase-free DNase I (04716728001 Roche)) in a pre-chilled 10mL jar (014620331, Retsch).
Pulverization was done by cryomill (Retsch) at 15 Hz for 15 minutes. The thawed cell lysates
were quantified by NanoDrop and 200µL of lysates with RNA concentration ranging from 80µg
to 500µg were used. For overall polysome quantification, lysates were loaded to 10%-55%
linear sucrose gradients (20mM Tris-HCl pH 8.0, 10mM $MgCl_2$, 100mM NH_4Cl, 300 µM

Chloramphenicol) made by GradientMaster (BioComp). The gradients were placed in a SW41Ti
bucket and centrifuged in Optima XE-100 Ultracentrifuge (Beckman Coulter) at 35,000 rpm for 2
hours at 4°C. Gradients were fractionated by BioComp Gradient Fractionator and the absorption
curves at 254nm were recorded by a UV monitor (EM-1, BioRad).
Quantification of the polysome profiles was done using customized MATLAB codes. First,
baselines were estimated using the readings where no peaks existed and the background was
subtracted. The background from the free nucleotides and tRNA was removed by fitting an
exponential decay function to the first peak representing the source of non-ribosome signals.
Each ribosome peak was picked and quantified by integrating the area underneath the curve.
To quantify different species of ribosomes in the 70S peak, 100mM NH_4Cl was replaced with
170mM KCl. Cell lysates were loaded onto 10%-30% linear gradient and centrifuged at 35,000
rpm at 4°C for 5 h. Because the two peaks for the free and mRNA-bound 70S ribosomes are
very close in mass and clean separation was not possible, the MATLAB file-exchange package,
peakfit, was used to fit the two overlapped peaks as two Gaussian distributions. For the free
ribosome control, cells were treated with 100μM puromycin for 5 minutes and collected. For
serine hydroxamate (SHX, Sigma S4503) treatment, cells were grown in MOPS glucose
minimal media until OD ~0.3 and treated with SHX with final concentration of 1 mg/mL for ten
minutes before collection.
481
lacZ induction and translational elongation rate measurement
The method was adapted from Zhu et al[47]. A final concentration of 5 mM IPTG (I2481C-25,
Gold Biotechnology) was added to cultures. At every 15 seconds, 1 mL of culture was collected
in a tube containing 10μL 100mM chloramphenicol, snap frozen in liquid nitrogen and stored at -
20°C before subsequent measurement. After cells were thawed, 400μL of the sample was
added to 100 μL 5x Z-buffer (0.3M $Na_2HPO_4.7H_2O$, 0.2M $NaH_2PO_4.H_2O$, 50mM KCl, 5mM
$MgSO_4$, 20mM β-mercaptoethanol) and incubated at 37°C for 10 minutes. 100μL of 4mg/mL
MUG (337210010, ACROS Organics) in DMSO was then added to each sample every 10
seconds to accurately control the reaction time. The samples were incubated at 37°C in a
thermomixer at 1,400 rpm mixing rate for 30 minutes to 2 hours, depending on the enzyme
expression level. The reaction was stopped by addition of 300μL of 1M Na_2CO_3. The tubes were
spun down at 16,000 X g for 3 minutes to sediment the cell debris. 200 μL of supernatant were
taken and measured fluorescence by a microplate reader (365 nm excitation and 450 nm
emission filter).
To infer translational elongation rate, the square root of the signal in excess of the signal at
time zero was plotted. A linear fit was performed on the points after signal began to increase.
The lag time is the x-intercept of the line. A previous study has shown that the initiation time
remains constant for about 10 seconds across a wide range of conditions tested[47]. Therefore,
we corrected the elongation times measured by subtracting 10 seconds in the lag times
measured.
502
Ribosome footprinting and total RNA extraction for RNA-Seq
The cell collection step was the same as for polysome profiling except that 1mM
chloramphenicol was used in the sucrose solution. The footprinting and library preparation steps
were adapted from Li *et al.*[2] After quantification of RNA concentration with NanoDrop, samples
with 500μg RNA were digested with 750U MNase (10107921001, Roche) for 1 hour at 25°C
before being quenched with 6mM EGTA. The lysates were then layered onto a 10%-55%
sucrose gradient and centrifuged. The monosome fraction was collected and snap frozen in
liquid nitrogen. No polysome peaks were observed, indicating a thorough digestion. The RNA
was isolated using hot phenol and size selected on 15% TBE-Urea PAGE gels run for 1 hour at
210V. Gels were stained with SYBR Gold and visualized using Dark Reader (Clare Chemical
Research). RNA fragments with size between 25-40 nt were extracted using isopropanol

precipitation. Total RNA was extracted with TRIZOL from the same pulverized cells used for
footprinting. After DNaseI (04716728001,Roche) treatment and cleanup using RNA clean &
concentrator 5 (R1016, Zymo Research), ribosomal RNA was subtracted using MicrobeExpress
(AM1905, Ambion). The recovered RNA was fragmented and size selected. RNA fragments
with size between 25-40nt were extracted using isopropanol precipitation.
519
Library preparation and sequencing
Fragments from footprints and total RNA were dephosphorylated at the 3' end by PNK
(M0201, NEB). The repaired fragments were linked to the Universal miRNA Cloning Linker
(S1315S, NEB), reverse transcribed (18080044, Thermo)and circularized (CL4111K, Epicentre).
The circularized samples were PCR amplified (M0531L, NEB) and size selected. High quality
PCR samples checked by Bioanalyzer high sensitive DNA chip. Deep sequencing was done by
Illumina HiSeq 2500 on Rapid flowcells with settings of single end and 75 nt-long read length.
527
Mapping and sequencing data analysis
Data manipulation including barcode splitting, linker trimming and mapping were done
using Galaxy. The processed reads were mapped to *Escherichia coli* genome
escherichia_coli_k12_nc_000913_3 from the NCBI database with the BWA short read mapping
algorithm. Only the reads between 20-45 nt that aligned to the coding region were used for
further analysis.
To infer the ribosome A-site position, python package Plastid[48] was used to align the 3' end
of reads to the stop and start codons[49], which are known to have higher ribosome densities. We
found that the offsets were 12 nt for stop codon and 15 nt for start codon. Therefore, we used
11nt for A site position and 14nt for P site. The counts were normalized to the total counts in the
coding region as reads per million (RPM) and transcripts per million (TPM). Further analysis
was done using customized Python and R codes with packages including Plastid, dplyr, tidyr
and ggplot2.
541
Analysis of ribosome profiling data
Transcripts per million (TPM) was used to identify highly expressed genes for codon
occupancy analysis. After assigning each mapped read to the A-site nucleotide, the raw counts
were first normalized by the length of their mapped coding region, to yield count density for each
gene. These count densities were then globally normalized to one million counts across all
genes within one sample. To determine codon occupancy, transcripts having total counts over
100 TPM and containing more than 200 codons were considered. Ribosome footprint counts for
the first and last 40 codons were removed to avoid possible effects from initiation and
termination, and counts per codon were recorded for the remaining counts. For each gene and
each codon type, the codon occupancy ratio is defined as the ratio of measured counts to
expected counts. Expected counts are simply proportional to the frequency of that particular
codon in the gene. The final reported codon occupancy ratio is the average of codon occupancy
ratios from all genes considered weighted equally.
We calculated ribosome counts along a transcript using reads per million (RPM). Only
transcripts having total counts over 10 TPM and containing more than 100 codons were
considered. Total counts after filtering were normalized to one million reads. The selected genes
account for more than 85% of the total counts. The ribosome counts are the sum of RPM at
each position.
To do the first and second halves comparison of ribosome counts of genes, the same set
of genes and trimming processing for codon occupancy were used. The RPM from positions
were summed up based on its assigned location (first or second half) on the transcript and plot
against each other.
564

565 Code availability

566 All the codes used for data analysis in this paper are available upon request.

567 Data availability

568 All the data used to reach the conclusion of this paper will be made available upon publication
569 on SRA BioProject with submission number SUB3256640.