Allosteric effects in catalytic impaired variants of the enzyme cyclophilin A may be explained by changes in nano-microsecond time scale motions

Pattama Wapeesittipan,[a] Antonia S. J. S. Mey,[a] Malcolm D. Walkinshaw,[b] and Julien Michel[a]*

[a] EaStCHEM School of Chemistry, David Brewster road, Joseph Black Building, The King's Buildings, Edinburgh, EH9 3FJ, UK.

[b] School of Biological Sciences, Michael Swann Building, Max Born Crescent, Edinburgh, Scotland, EH9 3BF, UK

E-mail: mail@julienmichel.net

Keywords: cyclophilins, catalysis, protein dynamics, molecular dynamics simulations

Abstract:

There is much debate about the mechanisms by which molecular motions influence catalysis in enzymes. This work investigates the connection between stochastic protein dynamics and function for the enzyme cyclophilin A (CypA) in wild-type (WT) form, and three variants that features several mutations that are distal from the active site. Previous biophysical studies have suggested that conformational exchange between a 'major' active and a 'minor' inactive state on millisecond timescales plays a key role in catalysis for CypA. Here this hypothesis was addressed by a variety of molecular dynamic (MD) simulation techniques. The simulations reproduce X-ray crystallography derived evidence for a shift in populations of major and minor active site conformations between the wild-type and mutant forms. Strikingly, exchange between these active site conformations occurs at a rate that is 5 to 6 orders of magnitude faster than previously proposed. Further analyses indicate that the minor active site conformation is catalytically impaired, and that decreased catalytic activity of the mutants may be explained by changes in Phe113 motions on a ns-µs timescale. Therefore previously described millisecond time scale motions may not be necessary to explain allosteric effects in CypA mutants.

A major goal of modern molecular biophysics is to clarify the connection between protein motions and enzymatic catalysis.[1–3] A wide range of experimental methods, e.g. neutron scattering, X-ray crystallography, NMR, or vibrational spectroscopy have been used to characterize internal protein motions occurring from femtosecond to second timescales.[4,5] While there is broad consensus that protein motions are implicated in catalysis, there is much debate around the role of conformational changes occurring on a millisecond timescale, and several studies have linked changes in millisecond protein motions with changes in enzymatic function.[6–9] However, it remains unclear whether such motions have a causal link to catalysis, or are merely a manifestation of the inherent flexibility of proteins over a broad range of timescales.

There have been vigorous debates about the meaning of dynamics in the context of enzymatic catalysis.[10–12] In the framework of transition state theory, the reaction rate is given by equation 1:

$$k = A(T)e^{-\Delta G^{\ddagger}(T)/RT} \quad (1)$$

where T is the temperature and R the gas constant. The pre-exponential term $A(T)$ includes contributions from non-statistical motions such as re-crossing or tunnelling, The exponential term involves the activation free energy of the chemical step $\Delta G^{\ddagger}(T)$. If transitions between reactant states are fast compared to the time scale of the chemical reaction, $\Delta G^{\ddagger}(T)$ is the free energy difference between the thermally equilibrated ensembles describing the reactant and transition states.[13,14] Non-statistical motions described by $A(T)$ have typically been found to make a small contribution to rate constants with respect to the exponential term that involves equilibrium fluctuations of the protein and solvent degrees of freedom.[15]

The current work is concerned with the connection between rates of thermally equilibrated motions, and catalysis in enzymes. Specifically, the focus is on clarifying the nature of protein motions implicated in catalysis for the well-studied enzyme cyclophilin A (CypA). CypA is a

member of the cyclophilin family of peptidyl-prolyl isomerases which catalyzes the *cis/trans* isomerization of amide groups in proline residues.[16] CypA plays an essential role in protein folding and regulation, gene expression, cellular signaling and the immune system. Notably, CypA is involved in the infectious activity and the viral replication of HIV-1.[17] Accordingly, CypA has been the subject of structure-based drug design efforts for decades.[18–20] Because of its significance as a medical target, the catalytic mechanism of CypA has been the subject of extensive studies.[2,3,21–30] Computational studies have shown that the speedup of the rate of *cis/trans* isomerization rate of the prolyl peptide bond is a result of preferential transition-state stabilization through selective hydrogen bonding interactions in the active site of CypA.[26,30] Figure 1(**a**) depicts key interactions between the substrate and active site residues, whereas Figure 1(**b**) highlights the relevant ω angle of the substrate used to track the *cis/trans* isomerization reaction.

Elegant NMR relaxation experiments by Eisenmesser et al. have also characterized the existence of intrinsic motions in apo CypA that couple a 'major' state M with a 'minor' conformational state m with a rate constant $k_{M \to m} = 60$ s^{-1}.[27] Fraser et al. later used ambient temperature X-ray crystallographic data to determine a high-resolution structure of this CypA state m, revealing an interconversion pathway with the 'major' state M that involves coupled rotations of a network of side-chains involving residues Ser99, Phe113, Met61, and Arg55. To establish the relevance of this 'minor' state m to catalysis, the distal residue Ser99 was mutated to Thr99 (now only referred to as ST). Further X-ray and NMR measurements on the free enzyme confirmed that the ST mutant increased the population of the m state, while decreasing the conversion rate $k_{M \to m}$ to 1 s^{-1}.[31] Remarkably, additional NMR experiments established that this 60-fold decrease in conversion rate between M and m states in the ST mutant correlates with

a ca. 70-fold decrease in bidirectional isomerization rate $(k_{iso} = k_{cis \to trans} + k_{trans \to cis})$ of a model substrate with respect to wild-type (WT). The effect is comparable to rate decreases observed for mutations of key active site residues such as Arg55.[31] More recently, two further mutants were reported in an effort to rescue the lost enzymatic activity of ST. These mutations were S99T and C115S (now only referred to as STCS), or S99T, C115S, and I97V (now only referred to as STCSIV). The two newly introduced mutants recover the enzyme activity to some extent, which correlates with an increase in $k_{M \to m}$ values.[32]

While this body of work suggested a link between millisecond time scale motions and catalysis in enzymes, there is currently no detailed mechanistic explanation for the decreased catalytic activity of the mutants. The present study uses a variety of extensive equilibrium and biased molecular dynamics (MD) simulations to clarify the link between catalytic activity and rates of molecular motions of CypA in wild-type and the three mutant variants.

Figure 1. The active site of cyclophilin A. (**a**) Key residues in the active site of cyclophilin A that form hydrogen bonds (cyan sticks, dashed lines) or are in contact (purple sticks) with the transition-state form of the peptide Ace-AAPF-Nme (orange sticks). For clarity, the Phe side-chain of the substrate is represented as transparent sticks. The distal residues Ser99, Cys115 and Ile97 are depicted in green. (**b**) The ω torsional angle of -Ala2-Pro3- is used to track the progression of the isomerization reaction between *cis* and *trans* forms.

Results

The proposed 'major' and 'minor' conformations exchange on timescales of nanoseconds in apo CypA

Fraser et al. have described the proposed 'major' and 'minor' states according to sets of values of χ_1 (Phe113, Ser/Thr99), χ_2 (Met61) and χ_3 (Arg55) angles.[31,33] These dihedrals as well as the side-chain dihedrals χ_1 of Ile97 and Cys115 were used to construct a Markov state model (MSM) to obtain quantitative information on thermodynamic and kinetic properties of the WT protein and the three experimentally studied mutants. The consistency of the MSMs was evaluated using standard protocols (See Supplementary Figures S1, S2). In the case of WT the accuracy of the MSM was additionally evaluated by back-calculation of previously reported NMR observables.[34] The MSM yields predictions of observables that show broadly similar accuracy to that of the NMR ensembles of Chi et al. and Otter et al.(see Supplementary Figure S3).[35,36] Thus the simulations were deemed sufficiently consistent with experimental data to warrant further analyses.

The X-ray structures of the key active site dihedrals in their dominantly populated states (if multiple occupancy is observed) are shown in Figure 2(**a**) for WT and ST, 2(**b**) for WT and STCS, and 2(**c**) for WT and STCSIV mutants. The most striking feature of the 'major' and 'minor' conformations are the rotameric states of χ_1 of Phe113 from the crystal structures, which in the 'minor' conformation is $\chi_1 \approx -60°$. This will be referred to as the '*out*' conformation. In contrast, the 'major' state $\chi_1 \approx 60°$, takes an '*in*' conformation. In Figure 2(**d**), (**e**), and (**f**) crystal structure occupancies for Phe113 χ_1 are compared to the MSM-derived dihedral distributions comparing WT and ST, WT and STCS, and WT and STCSIV respectively. The simulations suggest that in apo WT the Phe113 '*in*' and '*out*' orientations are equally likely, which is

consistent with the relatively similar occupancies of the two rotamers in the X-ray structure (occupancies = 0.63 and 0.37 respectively).[31] In apo ST there is a significant population shift towards the *'out'* orientation (χ_1=-60º), and the *'in'* orientation has a marginal population (ca. 1%), see Figure 2(**d**). This agrees with the X-ray structure of ST where only the Phe113 *'out'* rotamer is observed (occupancy = 1.0). In the STCS and STCSIV mutants the *'in'* rotamer is also destabilized with respect to wild-type but to a lesser extent (populations of ca. 16% and 17% respectively). Though only one *'out'* rotamer was resolved in the X-ray structure of STCS (Figure 2(**e**)), a major *'out'* and a minor distorted *'in'* rotamer (χ_1=+31º, occupancy 0.21) are observed in the X-ray structure of STCSIV (Figure 2(**f**)). Rotamers of other side-chain dihedrals of the key residues for all WT and mutants are found in Supplementary figures S4 and S5.

Figure 2. Comparison of X-ray and MSM-derived conformational preferences of Phe113. The X-ray structures of WT (blue and cyan) are compared with ST (red) (**a**), STCS (purple) (**b**) and STCSIV (dark and light green) (**c**).[31,32] The MSM-derived probability distributions of χ_1 in Phe113 for WT and ST (**d**), STCS (**e**) and STCSIV (**f**) are depicted as solid lines. The X-ray crystallography χ_1 values are depicted for WT and mutants with their respective occupancies as dashed lines.[31,32]

Surprisingly the Phe113 χ_1 dihedral was observed to flip frequently in MD trajectories of 200 ns duration (Supplementary Figure S6), suggesting faster motions than what NMR experiments have suggested. Therefore the MSM was used to obtain quantitative information on transition rates between *'in'* and *'out'* states as defined by the Phe113 χ_1 rotamer. Figure 3 summarises the MSM results. WT shows the fastest overall dynamics and ST the slowest, while the STCS and

STCSIV mutants show an intermediate behaviour. The exchange rates vary from $238 \pm 18 \ \mu s^{-1}$ (ST) to $47 \pm 4 \ \mu s^{-1}$ (STCS). Remarkably these values are five orders of magnitude faster than the exchange rates that have been determined by NMR measurements for motions involving Phe113.

	WT	S99T	S99TCS	S99TCSIV
P(out)	0.54 ± 0.00	0.99 ± 0.01	0.84 ± 0.01	0.83 ± 0.01
P(in)	0.46 ± 0.01	0.01 ± 0.00	0.16 ± 0.01	0.17 ± 0.01
k_A	65.7 ± 0.7	3.8 ± 0.2	7.3 ± 0.3	9.5 ± 0.5
k_B	79.2 ± 2.0	234.7 ± 17.6	39.2 ± 3.6	47.6 ± 4.9

Figure 3. Thermodynamics and kinetics of the Phe113 in/out flip. The diagram illustrates the flip of the Phe113 χ_1 dihedral between 'in' or 'out' states. Fractional populations and transition rates between 'in' and 'out' states in μs^{-1} for the four CypA variants are given in the table below the diagram. Error bars represent one standard deviation of the mean.

No differential catalytic activity is observed in the 'in' conformations; the 'out' conformation is catalytically inactive

Given that the timescales of rotations of Phe113 in the four CypA variants appear much faster than previously suggested, attention turned next to substrate bound CypA simulations. Results from umbrella sampling (US) simulations were used to quantify the isomerization free energy profile for WT and the ST mutant and investigate the role of Phe113 motions in catalysis.

The isomerization free energy profiles for WT and ST mutant with the side-chain of the Phe113 in an *'in'* and *'out'* conformations are shown in Figure 4(**a**) and 4(**b**) respectively. Ladani and Hamelberg have previously shown that fixed-charge classical force fields reproduce the energetics of amide bond rotation due to relatively small changes in intramolecular polarization during this process reasonably well.[28] The calculated activation free energy for the uncatalyzed *cis*→*trans* isomerization process in water is consistent with experimental data (20.1 ± 0.1 kcal•mol^{-1} vs ca. 19.3 kcal•mol^{-1} for the related substrate Suc-AAPF-pNA at 283 K).[37,38] The free energy profile for the substrate bound to CypA WT and ST in the *'in'* conformation shows that the enzyme catalyzes the isomerization reaction in both directions via a transition state with a positive ω value (ca. 90-100º) equally well (Figure 4(**a**)). There is a more significant decrease in activation free energy for *trans*→*cis* (ca. -6 kcal•mol^{-1}) than for *cis*→*trans* with less than 1 kcal•mol^{-1} difference between WT and ST, because the *cis* form is more tightly bound to CypA than the *trans* form. According to Figure 4(**b**), there is no catalytic benefit from the *'out'* conformation of the enzyme since the activation free energy of the isomerization reaction in CypA is similar to that of the substrate in water. The calculated free energy profiles for isomerization reactions in STCS and STCSIV show a similar trend (Supplementary Figure S8).

Systems	$\Delta G^{\ddagger}(cis{\rightarrow}trans)$	$\Delta G^{\ddagger}(trans{\rightarrow}cis)$
AAPF	20.1 ± 0.1	20.9 ± 0.1
WT-AAPF	19.7 ± 1.3	15.2 ± 0.8
S99T-AAPF	19.2 ± 1.4	15.1 ± 1.0

Systems	$\Delta G^{\ddagger}(cis{\rightarrow}trans)$	$\Delta G^{\ddagger}(trans{\rightarrow}cis)$
AAPF	20.1 ± 0.1	20.9 ± 0.1
WT-AAPF	20.4 ± 0.6	20.2 ± 0.5
S99T-AAPF	20.6 ± 1.5	19.9 ± 0.8

Figure 4. Catalytic energy profiles in the Phe113 'in' and 'out' conformations. **(a)** Isomerization free energy profiles (in kcal.mol^{-1}) for the substrate AAPF in water (black), and bound to WT (blue) or ST mutant (red) forms of CypA with starting structures in the 'in' conformation. The free energies of the *trans* conformation were set to zero at ω = 180°. Error bars represent one standard error of the mean. The table shows the activation free energies for both directions of the isomerization reaction. **(b)** same as **(a)**, but with stimulations starting in a 'out' configuration.

Decreased hydrogen-bonding interactions with multiple active site residues explain transition state destabilization in the minor conformation

Further analysis of the umbrella sampling trajectories shows that for the simulations started in the *'in'* configuration in both WT and ST the transition state region (ω ca. 90-100º) is electrostatically stabilized by more negative Coulombic interactions between substrate and binding site atoms as show in Figure 5(**a**). Figure 5(**b**) breaks down the different contribution of active site residues, showing that Arg55, Trp121, Asn102, His126, and Gln63 are important for the stabilization of the transition state ensemble via hydrogen bonding interactions as shown in Figure 5(**e**). In contrast, Figure 5(**c**) shows that for simulations in the *'out'* configuration no transition state stabilization through electrostatic interactions is observed, this is further reflected by the per-residue split of interaction energy contribution at the transition state in Figure 5(**d**) and the lack of hydrogen bond formation in Figure 5(**f**). Hydrogen bonding probabilities for simulations from the *'in'* and *'out'* starting conformations are shown in Supplementary Figures S9-S11. A similar picture holds for the STCS and STCSIV mutants (Supplementary Figure S12).

Figure 5. Structural basis for the differential catalytic activity of the Phe113 in and out conformations. (**a**) Electrostatic energies between the substrate and active site residues as a function of the ω angle in WT (blue) and the ST systems (red) with simulations started from the *'in'* configuration. (**b**) Average electrostatic energies per-active site residues at the transition state region from (**a**). (**c**) same as (**a**) with simulations started from the *'out'* configuration. (**d**) same as (**b**) with simulations started in the *'out'* conformation. Error bars denote the standard

error of the mean. (**e**) and (**f**) typical hydrogen bonding pattern at the transition state for simulations started in the *'in'* and *'out'* conformation respectively.

Preorganization can explain observed decreases in catalytic activity of the mutants

Taken together the MSM and US data suggest a mechanistic explanation for the effect of distal mutations on the catalytic activity of cyclophilin A. In WT free form the enzyme rapidly interconverts between a catalytically active F113 *'in'* form and a catalytically inactive F113 *'out'* form. Because the interconversion rate between in and out forms (ca. 7.10^7 s^{-1}) is faster than the substrate binding rate as suggested by NMR experiments (ca. 2.10^4 s^{-1}, based on k_{on} rate ca. 2.10^7 s^{-1}.M^{-1} and substrate concentration ca. 1 mM.)[39] the free enzyme rapidly equilibrates between catalytically active and inactive forms before substrate binding (Figure 6(**a**)). In the mutants form, the interconversion rates between catalytically active and inactive forms are still within the μs^{-1} timescale, but the equilibrium is shifted towards the catalytically inactive form (Figure 6(**b**)), thus the mutants are less pre-organized than WT and the overall catalytic activity is decreased.

In the case of the ST mutant and WT forms, Fraser et al. have reported bi-directional on-enzyme isomerization rates ($k_{cis \rightarrow trans} + k_{trans \rightarrow cis}$) by NMR spectroscopy, and found a ratio of 68±13 between WT and ST.[31] According to the model proposed in Figure 6 and by combining the MSM-derived populations and the US-derived activation free energies, a ratio of $12 < 46 <$ 176 can be derived from the simulations (see SI for details). The uncertainty from the simulations is larger than that of the measurements because small variations in activation free energies contribute large change in catalytic rates. Thus the model described in Figure 6 appears consistent with experimental data for WT and ST. No bidirectional isomerization rates have been reported for the STCS and STCSIV mutants.[32] However, the STCS and STCSIV mutants show

populations of the catalytically active Phe113 '*in*' conformation that are intermediate between WT and ST, which is consistent with their increased catalytic activity with respect to ST.

A defining feature of this model is that the χ1 rotamers of a number of active-site side-chains such as Gln63, Ile/Val97, Phe113, Cys/Ser115 flip in WT and mutants on ns-μs timescales. Back-calculation of C_β-C_γ order parameters shows that this effect is captured by a decrease in S^2 values upon increasing the averaging window from 10 ns to 100 ns (Supplementary Figure S13). Motions on these timescales are too rapid to be detected by CPMG or CEST experiments that have been used extensively to study μs-ms processes in cyclophilin A.[3,25,31,40,41] Likewise relaxation experiments cannot detect motions on this timescale as they are limited to processes occurring faster than the tumbling time τ_c of cyclophilin A (ca. 10 ns).[42] Residual Dipolar Couplings (RDCs) can, however, provide information about dynamic orientation of inter-nuclear vectors on the supra-τ_c time scale.[43] Such experiments have been reported for backbone and methyl-RDCs in ubiquitin.[43,44] Therefore the model predictions can be experimentally tested with combined nuclear spin relaxation and RDC based model-free analyses coupled with a labelling scheme that resolves χ1 side-chain motions.[43,45]

Figure 6. Proposed mechanism for allosteric inhibition of cyclophilin A function. **(a)** Catalysis in WT, with the favored route being the *'in'* conformations, which are similarly populated to the *'out'* conformations. **(b)** Catalysis in the mutants still occurs through the '*in*' conformations, which have a lower population than the 'out' conformations.

Discussion

This work highlights the potential of detailed molecular simulation studies to guide the interpretation of biophysical measurements for the elucidation of protein allosteric mechanisms.[46] Previous work has suggested that exchange on millisecond timescales between conformational states in CypA are linked to its catalytic cycle,[27] leading to a proposal for a slow exchange between a 'major' and a 'minor' state of a set of side chain rotamers linking distal residue Ser99 to active-site residues.[27,31] The present results do not support or reject this hypothesis because the MD simulations used here do not resolve motional processes occurring on timescales slower than microseconds. However a major finding of this study is that transitions between *'in'* and *'out'* rotamers of Phe 113 in WT and mutants occur on a time scale of ns-μs, thus five to six orders of magnitude faster than suggested by earlier NMR relaxation dispersion

measurements.[31] Nevertheless the simulations reproduce well the population shifts in Phe113 rotamers observed in room-temperature X-ray crystallography experiments. This suggests that the X-ray structures may have resolved motional processes occurring on a distinct timescales from the processes resolved by CPMG experiments. Indeed in reported CPMG experiments the millisecond motions of Phe113 are coupled to a large network of ca. 30 residues,[31] whereas the $\chi 1$ rotameric flip observed in the simulations, is a largely local motion.

Nevertheless, the simulations suggest that a local '*in*' to '*out*' rotation of Phe113 is sufficient to abrogate catalysis in cyclophilin A, and variations of exchange parameters on the ns-μs timescale between these two conformational states appear sufficient to explain the decreased catalytic activity of the ST, STCS, STCSIV mutants with respect to WT. Therefore it is advisable to carry out additional experiments to confirm the existence of Phe113 $\chi 1$ rotations on the ns-μs timescale before causally linking catalysis to millisecond time scale motions. On the computational side, efforts should focus on advancing MD methodologies such that millisecond timescale processes observed in experiments can be resolved in atomistic details.

The contribution of protein flexibility on the ps-ns and μs-ms timescales to enzymatic catalysis has been the focus of several computational and experimental studies.[3,8,10,13,15,25,27,31,32,47] Our work suggests that more efforts should be directed at resolving conformational processes on the ns-μs timescale. This has important conceptual implications for enzyme design and optimization strategies.

Methods

Systems preparation

Models for apo/substrate bound human CypA of the WT and ST were prepared for MD

simulations from PDB structures 3K0N (R=1.39 Å) and 3K0O (R= 1.55 Å) respectively. For apo STCS and STCSIV two structures were prepared from PDB structures 6BTA (R=1.5 Å) and 5WC7 (R=1.43Å) and also by mutating residues in WT using Schrödinger's Maestro.[48] For WT the major conformation of 3K0N (altloc A, occupancy 0.63) was retained. For STCS and STCSIV the residues with higher occupancy were chosen for initial structures. Supplementary Tables S1-S2 summarise all simulations conducted in this study. The proteins were solvated in a rhombic dodecahedron box of TIP3P water molecules with edges extending 1 nm away from the proteins and chloride counter-ions were added to neutralise the overall net-charge. The Charmm22* forcefield[49] was used to describe protein atoms in the apo simulations. Steepest descent minimized was used for 50,000 steps followed by equilibration for 100 ps in an NVT ensemble, and 100 ps NPT ensemble, with heavy protein atoms restraint using a harmonic force constant of 1000 kJ·mol^{-1}·nm^{-2}.

Models of CypA WT and other mutants in complex with the Ace-AAPF-Nme substrate were prepared, using the amber99sb forcefield with optimised ω angle parameters for amides as reported by Doshi and co-workers was used.[50] The crystal structure of the CypA-*cis* AAPF peptide complex (PDB ID: 1RMH)[51] was used to obtain a suitable orientation for the substrate in the active site of WT and other mutants. PDB structure 1RMH was aligned to the structure of WT and all mutants, and the N-terminal and C-terminal ends of the proteins and substrate were capped using Schrödinger's Maestro.[48] In order to generate starting structures of *'in'* and *'out'* CypA-substrate complexes, MD simulations of CypA-substrate complexes (*cis*-conformation) were performed for 10 ns. For ST, STCS and STCSIV mutants the χ_1 values of Phe113 were measured to monitor transitions between *'out'* and *'in'* rotamers. The last snapshot structure of *'in'* and *'out'* complexes structures were used as input umbrella sampling calculations. For WT

complexes, only the *'in'* rotamer was observed in a 10-ns MD simulation. Thus, Umbrella samplings (US) simulations of χ_1 (Phe113) were performed serially to generate the *'out'* ($\chi_1\approx$-60°) rotamer starting from the *'in'* rotamer ($\chi_1\approx$60°) using the software PLUMED2.[52] Also, in order to retain the substrate in the active site, the distance between the proline ring of substrate and the phenyl rings of Phe113 and Phe60 were restrained using a force constant of 300 kJ·mol⁻¹·rad⁻². Each US simulation was performed for 5 ns. The bias parameters and the restrained variables for the US of χ_1 (Phe113) are summarised in Supplementary Table S5.

apo WT and mutant MD simulations

80 independent 200 ns MD trajectories of the apo WT, ST, STCS, and STCSIV proteins (20 each) were generated using Gromacs 5.0.[53] For apo STCS and STCSIV the MD simulations were split between both structures prepared independently. A 2 fs time step was used, and the first 5 ns discarded for equilibration. Temperature was maintained at 300 K with a stochastic Berendsen thermostat.[54] The Parrinello-Rahman barostat was used for pressure coupling at 1 bar.[55] The Particle Mesh Ewald scheme was used for long-range electrostatic interactions with a Fourier grid spacing of 0.16 nm, and fourth-order cubic interpolation.[56] Short-range van der Waals and electrostatic interaction were cutoff at 1 nm. The LINCS algorithm was used to constrain all bonds.[57]

Markov state models

All MSM analysis was carried out with the software package pyemma version 2.3.2.[58] The focus was on the side-chain motion of binding site residues. Details on which dihedral angles were used for TICA[59] is given in Supplementary Table S6. A more detailed description of the Markov State model in particular with respect to best model selection is given in the SI. Clustering was done using all trajectory data from the WT and mutant trajectories using a set of

24 input coordinates, with selecting dominant coordinates using a 90% variance in TICA for the subsequent k-means clustering. It was found that 200 clusters was necessary to separate in and out rotamers of Phe113 in ST. With the same cluster assignment for all trajectories, MSM transition matrices were estimated, using the Bayesian MSM option, and choosing lagtimes 0.6 ns for WT, S99T, C115S, and I97V. Means and errors of observables (e.g. populations and MFPT) were estimated from the Bayesian MSM using the provided functions in pyEMMA. Membership assignments were based on the MSM microstate dihedral probabilities of being in the *'in'* or *'out'* state respectively. The microstate definition used for the MSMs is the same across the WT and all mutants. The MFPTs are estimated between the manually grouped two states depending on whether the Phe113 rotamer is *'in'* or *'out'* in the microstate. MSM validation and further details on the MSM can be found in the SI and in particular Supplementary Figures S1-S5.[60]

Umbrella sampling simulations

Series of Umbrella Sampling (US) simulations[61–63] of the *'in'* and *'out'* conformers were performed to compute free energy profiles along ω.[26,28,64,65] For substrate in solution, the initial structure of US was in a *trans* conformation taken from 10-ns equilibration run, while all protein-substrate complexes were in a *cis* conformation. For both of *'in'* and *'out'* US calculations, a standard harmonic potential was used to bias the ω angle towards a series of target values ω_k spanning the interval [-180°,180°]. The force constants of the biasing potential and the spacing between ω_k values were adjusted by trial and error in order to obtain a good overlap between probability distributions of neighbouring ω_k values (Supplementary Tables S3-S4 and Figure S7). For *'out'* US calculation the distances between the proline ring of substrate and the phenyl rings of Phe113 and Phe60 were restrained using a flat-bottom harmonic restraint with force

constants of 200 kJ·mol^{-1}·rad^{-2} and 300 kJ·mol^{-1}·rad^{-2}, respectively. Simulations were performed serially initially for 7 ns, with the starting conformation for a given target angle ω_k taken from the preceding run performed at the neighbouring $\omega_{k+\Delta\omega}$ value. Each US was then extended to 20 ns. A total of 22 (substrate in solution) or 24 (substrate bound to protein) umbrellas were used. In order to estimate uncertainties of free energy profiles six repeats of the entire procedure were performed for *'in'* and *'out'* US. All simulations were carried out using a PLUMED2 patched version of Gromacs 5.0 with simulation parameters identical to the previously described apo MD simulation protocols unless otherwise mentioned. The weighted histogram analysis method (WHAM) was used to produce a free energy profile from the pool of US simulations.[66]

Other trajectory analyses

Average proton-proton distances were derived as $r_{ij}^{avg} = \langle r_{ij}^{-6} \rangle^{\frac{-1}{6}}$ from snapshots sampled from the MSM of WT for comparison with NOEs and eNOEs-derived distance intervals.[67] $^{3}J(H^{N},H^{\alpha})$, $^{3}J(H^{N},C')$, and $^{3}J(H^{N},C^{\beta})$ were also computed using Karplus equations and backbone dihedral angle values $\langle\phi\rangle$ and $\langle\psi\rangle$ sampled from the MSM.[68]

Interaction energies between binding site residues (Arg55, Ile57, Phe60, Met61, Gln63, Asn102, Gln111, Phe113, Trp121, Leu122 and His126) and all atoms of the substrate were analysed with the Gromacs g_energy module, using snapshots from the US simulations. The probability distribution of distances between key residues and substrate atoms during the simulations were computed using the MDAnalysis library.[69]

ASSOCIATED CONTENT. Detailed computational protocols, supplementary figures, supplementary tables, input files for the MD calculations.

AUTHOR INFORMATION

Corresponding Author

* mail@julienmichel.net

Present Addresses

Author Contributions

PW, ASJSM: Carried out experiments, analysed data, wrote manuscript. MDW: Analysed data, wrote manuscript. JM: Designed study, analysed data, wrote manuscript.

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Data availability. All input files and scripts used for the preparation, execution and analysis of the MD, MSM and US calculations are freely available at https://github.com/michellab/CypAcatalysis_input and are also provided as Supplementary dataset S1. Other data that support the findings of this study are available from the corresponding author upon reasonable request.

Acknowledgments

Gratitude is expressed to Fernanda Duarte for thoughtful discussions about this work.

Funding Sources

Julien Michel is supported by a University Research Fellowship from the Royal Society. The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement No. 336289. Pattama Wapeesittipan is supported by The Development and Promotion of Science

and Technology Talents Project (DPST) Scholarship, Royal Thai Government. This project made use of time on ARCHER granted via the UK High-End Computing Consortium for Biomolecular Simulation, HECBioSim (http://hecbiosim.ac.uk), supported by EPSRC (grant no. EP/L000253/1)

REFERENCES

1. Warshel, A. Dynamics of Enzymatic Reactions. *Proc. Natl. Acad. Sci.* **81,** 444–448 (1984).

2. McGowan, L. C. & Hamelberg, D. Conformational plasticity of an enzyme during catalysis: Intricate coupling between cyclophilin a dynamics and substrate turnover. *Biophys. J.* **104,** 216–226 (2013).

3. Eisenmesser, E. Z., Bosco, D. A., Akke, M. & Kern, D. Enzyme Dynamics During Catalysis. *Science.* **295,** 1520–1523 (2002).

4. Rader, S. D. & Agard, D. a. Conformational substates in enzyme mechanism: the 120 K structure of alpha-lytic protease at 1.5 A resolution. *Protein Sci.* **6,** 1375–86 (1997).

5. Bu, Z. *et al.* A view of dynamics changes in the molten globule native folding step by quasielastic neutron scattering. *J. Mol. Biol.* **301,** 525–536 (2000).

6. Henzler-Wildman, K. & Kern, D. Dynamic personalities of proteins. *Nature* **450,** 964–972 (2007).

7. Wolf-Watz, M. *et al.* Linkage between dynamics and catalysis in a thermophilic-mesophilic enzyme pair. *Nat. Struct. Mol. Biol.* **11,** 945–9 (2004).

8. Palmer, A. G., Kroenke, C. D. & Loria, J. P. NMR methods for quantifying microsecond-to-millisecond motions in biological macromolecules. *Methods Enzym.* **339,** 204–238

(2001).

9.   Schlegel, J., Armstrong, G. S., Redzic, J. S., Zhang, F. & Eisenmesser, E. Z. Characterizing and controlling the inherent dynamics of cyclophilin-A. *Protein Sci.* **18,** 811–824 (2009).

10.   Pisliakov, A. V, Cao, J., Kamerlin, S. C. L. & Warshel, A. Enzyme millisecond conformational dynamics do not catalyze the chemical step. *Proc. Natl. Acad. Sci. U. S. A.* **106,** 17359–17364 (2009).

11.   Kamerlin, S. C. L. & Warshel, A. At the Dawn of the 21st Century : Is Dynamics the Missing Link. *Proteins Struct. Funct. Bioinforma.* **78,** 1339–1375 (2010).

12.   Warshel, A. & Bora, R. P. Perspective: Defining and quantifying the role of dynamics in enzyme catalysis. *J. Chem. Phys.* **144,** (2016).

13.   Kohen, A. Role of dynamics in enzyme catalysis: Substantial versus semantic controversies. *Acc. Chem. Res.* **48,** 466–473 (2015).

14.   Wolfenden, R. Transition state analogues for enzyme catalysis. *Nature* **223,** 704–705 (1969).

15.   Delgado, M. *et al.* Convergence of Theory and Experiment on the Role of Preorganization, Quantum Tunneling, and Enzyme Motions into Flavoenzyme-Catalyzed Hydride Transfer. *ACS Catal.* **7,** 3190–3198 (2017).

16.   Go, S. F., Marahiel, M. A. & Gothel, S. F. Peptidyl-prolyl cis-trans isomerases , a superfamily of. *Cell Mol Life Sci* **55,** 423–436 (1999).

17.   Li, J. *et al.* Discovery of dual inhibitors targeting both HIV-1 capsid and human cyclophilin A to inhibit the assembly and uncoating of the viral capsid. *Bioorganic Med. Chem.* **17,** 3177–3188 (2009).

18. Dornan, J., Taylor, P. & Walkinshaw, M. Structures of Immunophilins and their Ligand Complexes. *Curr. Top. Med. Chem.* **3,** 1392–1409 (2003).

19. Yang, Y. *et al.* Structure-based discovery of a family of synthetic cyclophilin inhibitors showing a cyclosporin-A phenotype in Caenorhabditis elegans. *Biochem. Biophys. Res. Commun.* **363,** 1013–1019 (2007).

20. Georgiou, C. *et al.* Pushing The Limits Of Detection Of Weak Binding Using Fragment Based Drug Discovery: Identification Of New Cyclophilin Binders. *bioRxiv* (2017).

21. Nagaraju, M., McGowan, L. C. & Hamelberg, D. Cyclophilin a inhibition: Targeting transition-state-bound enzyme conformations for structure-based drug design. *J. Chem. Inf. Model.* **53,** 403–410 (2013).

22. Kofron, J. L., Kuzmic, P., Kishore, V., Colón-Bonilla, E. & Rich, D. H. Determination of kinetic constants for peptidyl prolyl cis-trans isomerases by an improved spectrophotometric assay. *Biochemistry* **30,** 6127–6134 (1991).

23. Doshi, U., Holliday, M. J., Eisenmesser, E. Z. & Hamelberg, D. Dynamical network of residue-residue contacts reveals coupled allosteric effects in recognition, catalysis, and mutation. *Proc. Natl. Acad. Sci. U. S. A.* **113,** 4735–4740 (2016).

24. Camilloni, C. *et al.* Cyclophilin A catalyzes proline isomerization by an electrostatic handle mechanism. *Proc. Natl. Acad. Sci. U. S. A.* **111,** 10203–8 (2014).

25. Holliday, M. J., Camilloni, C., Armstrong, G. S., Vendruscolo, M. & Eisenmesser, E. Z. Networks of Dynamic Allostery Regulate Enzyme Function. *Structure* **25,** 276–286 (2017).

26. Hamelberg, D. & McCammon, J. A. Mechanistic insight into the role of transition-state stabilization in cyclophilin A. *J. Am. Chem. Soc.* **131,** 147–152 (2009).

27. Eisenmesser, E. Z. *et al*. Intrinsic dynamics of an enzyme underlies catalysis. *Nature* **438,** 117–21 (2005).

28. Ladani, S. T. & Hamelberg, D. Entropic and surprisingly small intramolecular polarization effects in the mechanism of cyclophilin A. *J. Phys. Chem. B* **116,** 10771–10778 (2012).

29. Trzesniak, D. & van Gunsteren, W. F. Catalytic mechanism of cyclophilin as observed in molecular dynamics simulations: pathway prediction and reconciliation of X-ray crystallographic and NMR solution data. *Protein Sci.* **15,** 2544–51 (2006).

30. Ladani, S. T. & Hamelberg, D. Intricacies of interactions, dynamics and solvent effects in enzyme catalysis: a computational perspective on cyclophilin A. *Mol. Simul.* **40,** 765–776 (2014).

31. Fraser, J. S. *et al*. Hidden alternative structures of proline isomerase essential for catalysis. *Nature* **462,** 669–673 (2009).

32. Otten, R. *et al*. Rescue of conformational dynamics in enzyme catalysis by directed evolution. *Nat. Commun.* **9,** (2018).

33. Papaleo, E., Sutto, L., Gervasio, F. L. & Lindorff-Larsen, K. Conformational changes and free energies in a proline isomerase. *J. Chem. Theory Comput.* **10,** 4169–4174 (2014).

34. Vögeli, B., Kazemi, S., Güntert, P. & Riek, R. Spatial elucidation of motion in proteins by ensemble-based structure calculation using exact NOEs. *Nat. Struct. Mol. Biol.* **19,** 1053–1058 (2012).

35. Chi, C. N. *et al*. A Structural Ensemble for the Enzyme Cyclophilin Reveals an Orchestrated Mode of Action at Atomic Resolution. *Angew. Chemie - Int. Ed.* **54,** 11657–11661 (2015).

36. Ottiger, M., Zerbe, O., Güntert, P. & Wüthrich, K. The NMR solution conformation of

unligated human cyclophilin A. *J. Mol. Biol.* **272,** 64–81 (1997).

37.   Harrison, R. K. & Stein, R. L. Mechanistic studies of enzymic and nonenzymic prolyl cis-trans isomerization. *J. Am. Chem. Soc.* **114,** 3464–3471 (1992).

38.   Dugave, C. & Demange, L. Cis-trans isomerization of organic molecules and biomolecules: Implications and applications. *Chem. Rev.* **103,** 2475–2532 (2003).

39.   Kern, D., Kern, G., Scherer, G., Fischer, G. & Drakenberg, T. Kinetic analysis of cyclophilin-catalyzed prolyl cis/trans isomerization by dynamic NMR spectroscopy. *Biochemistry* **34,** 13594–602 (1995).

40.   Holliday, M. J., Armstrong, G. S. & Eisenmesser, E. Z. Determination of the Full Catalytic Cycle among Multiple Cyclophilin Family Members and Limitations on the Application of CPMG-RD in Reversible Catalytic Systems. *Biochemistry* **54,** 5815–5827 (2015).

41.   Holliday, M. J. *et al.* Structure and dynamics of GeoCyp: A thermophilic cyclophilin with a novel substrate binding mechanism that functions efficiently at low temperatures. *Biochemistry* **54,** 3207–3217 (2015).

42.   Gu, Y., Li, D. W. & Brüschweiler, R. NMR order parameter determination from long molecular dynamics trajectories for objective comparison with experiment. *J. Chem. Theory Comput.* **10,** 2599–2607 (2014).

43.   Lakomek, N. A. *et al.* Self-consistent residual dipolar coupling based model-free analysis for the robust determination of nanosecond to microsecond protein dynamics. *J. Biomol. NMR* **41,** 139–155 (2008).

44.   Farès, C. *et al.* Accessing ns-μs side chain dynamics in ubiquitin with methyl RDCs. *J. Biomol. NMR* **45,** 23–44 (2009).

45. Kasinath, V., Valentine, K. G. & Wand, A. J. A 13C labeling strategy reveals a range of aromatic side chain motion in calmodulin. *J. Am. Chem. Soc.* (2013). doi:10.1021/ja4001129

46. Michel, J. Current and emerging opportunities for molecular simulations in structure-based drug design. *Phys. Chem. Chem. Phys. Phys. Chem. Chem. Phys* **16,** 4465–4477 (2014).

47. Liu, C. T., Wang, L., Goodey, N. M., Hanoian, P. & Benkovic, S. J. Temporally overlapped but uncoupled motions in dihydrofolate reductase catalysis. *Biochemistry* **52,** 5332–5334 (2013).

48. Schrödinger Release 2017-4: Maestro, Schrödinger, LLC, New York, NY, 2017. .

49. Piana, S., Lindorff-Larsen, K. & Shaw, D. E. How robust are protein folding simulations with respect to force field parameterization? *Biophys. J.* **100,** (2011).

50. Doshi, U. & Hamelberg, D. Reoptimization of the AMBER force field parameters for peptide bond (Omega) torsions using accelerated molecular dynamics. *J. Phys. Chem. B* **113,** 16590–16595 (2009).

51. Zhao, Y. & Ke, H. Crystal structure implies that cyclophilin predominantly catalyzes the Trans to Cis isomerization. *Biochemistry* **35,** 7356–7361 (1996).

52. Tribello, G. A., Bonomi, M., Branduardi, D., Camilloni, C. & Bussi, G. PLUMED 2: New feathers for an old bird. *Comput. Phys. Commun.* **185,** 604–613 (2014).

53. Abraham, M. J. *et al.* Gromacs: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **1–2,** 19–25 (2015).

54. Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity rescaling. *J. Chem. Phys.* **126,** (2007).

55. Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new molecular dynamics method. *J. Appl. Phys.* **52,** 7182–7190 (1981).

56. Essmann, U. *et al.* A smooth particle mesh Ewald method. *J. Chem. Phys.* **103,** 8577–8593 (1995).

57. Hess, B., Bekker, H., Berendsen, H. J. C. & Fraaije, J. G. E. M. LINCS: A linear constraint solver for molecular simulations. *J. Comput. Chem.* **18,** 1463–1472 (1997).

58. Scherer, M. K. *et al.* PyEMMA 2: A Software Package for Estimation, Validation, and Analysis of Markov Models. *J. Chem. Theory Comput.* **11,** 5525–5542 (2015).

59. Schwantes, C. R. & Pande, V. S. Improvements in Markov State Model construction reveal many non-native interactions in the folding of NTL9. *J. Chem. Theory Comput.* **9,** 2000–2009 (2013).

60. Prinz, J. H. *et al.* Markov models of molecular kinetics: Generation and validation. *J. Chem. Phys.* **134,** (2011).

61. Torrie, G. M. & Valleau, J. P. Nonphysical sampling distributions in Monte Carlo free-energy estimation: Umbrella sampling. *J. Comput. Phys.* **23,** 187–199 (1977).

62. Hooft, R. W. W., van Eijck, B. P. & Kroon, J. An adaptive umbrella sampling procedure in conformational analysis using molecular dynamics and its application to glycol. *J Chem Phys* **97,** 6690–6694 (1992).

63. Northrup, S. H., Pear, M. R., Lee, C. Y., McCammon, J. A. & Karplus, M. Dynamical theory of activated processes in globular proteins. *Proc. Natl. Acad. Sci.* **79,** 4035–4039 (1982).

64. Agarwal, P. K. Cis/trans isomerization in HIV-1 capsid protein catalyzed by cyclophilin A: Insights from computational and theoretical studies. *Proteins Struct. Funct. Genet.* **56,**

449–463 (2004).

65.   Hamelberg, D., Shen, T. & McCammon, J. A. Phosphorylation effects on cis/trans isomerization and the backbone conformation of serine-proline motifs: Accelerated molecular dynamics analysis. *J. Am. Chem. Soc.* **127,** 1969–1974 (2005).

66.   Kumar, S., Rosenberg, J. M., Bouzida, D., Swendsen, R. H. & Kollman, P. A. The weighted histogram analysis method for free energy calculations on biomolecules. I. The method. *J. Comput. Chem.* **13,** 1011–1021 (1992).

67.   Zagrovic, B. & Van Gunsteren, W. F. Comparing atomistic simulation data with the NMR experiment: How much can NOEs actually tell us? *Proteins Struct. Funct. Genet.* **63,** 210–218 (2006).

68.   Salvador, P. Dependencies of J-Couplings upon Dihedral Angles on Proteins. *Annu. Reports NMR Spectrosc.* **81,** 185–227 (2014).

69.   Michaud-Agrawal, N., Denning, E. J., Woolf, T. B. & Beckstein, O. MDAnalysis: A toolkit for the analysis of molecular dynamics simulations. *J. Comput. Chem.* **32,** 2319–2327 (2011).

Graphical TOC