Mapping the phenotypic repertoire of the cytoplasmic 2-Cys peroxiredoxin – thioredoxin system. 1. Understanding commonalities and differences among cell types.

Gianluca Selvaggio[1,2], Pedro M. B. M. Coelho[1] and Armindo Salvador[1,3,*]

[1] CNC – Center for Neuroscience and Cell Biology, University of Coimbra

[2] MIT-Portugal Program Bioengineering Systems Doctoral Program

[3] CQC, Department of Chemistry, University of Coimbra, Portugal

*Corresponding author: salvador@cnc.uc.pt

Abbreviations

ASK1, apoptosis signal-regulating kinase 1; Cat, catalase; GSH, glutathione; GPx1, glutathione peroxidase 1; Grx, glutaredoxin; KEAP1, Kelch-like ECH-associated protein 1; NRF2, nuclear factor erythroid 2-related factor 2; Prx, typical 2-Cys peroxiredoxin; PTTRS, peroxiredoxin / thioredoxin / thioredoxin reductase system; Srx, sulfiredoxin; Trx, thioredoxin; TrxR, thioredoxin reductase.

Abstract

The system (PTTRS) formed by typical 2-Cys peroxiredoxins (Prx), thioredoxin (Trx), Trx

reductase (TrxR), and sulfiredoxin (Srx) is central in antioxidant protection and redox signaling

in the cytoplasm of eukaryotic cells. Understanding how the PTTRS integrates these functions

requires tracing phenotypes to molecular properties, which is non-trivial. Here we analyze this

problem based on a model that captures the PTTRS' conserved features. We have mapped the

conditions that generate each distinct response to H_2O_2 supply rates (v_{sup}), and estimated the

parameters for thirteen human cell types and for *Saccharomyces cerevisiae*. The resulting

composition-to-phenotype map yielded the following experimentally testable predictions. The

PTTRS permits many distinct responses including ultra-sensitivity and hysteresis. However,

nearly all tumor cell lines showed a similar response characterized by limited Trx-S⁻ depletion

and a substantial but self-limited gradual accumulation of hyperoxidized Prx at high v_{sup}. This

similarity ensues from strong correlations between the TrxR, Srx and Prx activities over cell lines,

which contribute to maintain the Prx-SS reduction capacity in slight excess over the maximal

steady state Prx-SS production. In turn, in erythrocytes, hepatocytes and HepG2 cells high v_{sup}

depletes Trx-S⁻ and oxidizes Prx mainly to Prx-SS. In all nucleated human cells the Prx-SS

reduction capacity defined a threshold separating two different regimes. At sub-threshold v_{sup}

cytoplasmic H_2O_2 is determined by Prx, nM-range and spatially localized, whereas at

supra-threshold v_{sup} it is determined by much less active alternative sinks and μM-range

throughout the cytoplasm. The yeast shows a distinct response where the Prx Tsa1 accumulates

in sulfenate form at high v_{sup}. This is mainly due to an exceptional stability of Tsa1's sulfenate.

The implications of these findings for thiol redox regulation and cell physiology are discussed.

All estimates were thoroughly documented and provided, together with analytical

approximations for system properties, as a resource for quantitative redox biology.

Keywords

Redox relays; Redox signaling; Thiol redox regulation; Quantitative redox biology; Systems

design space methodology

Introduction

The PTTRS (Figure 1) plays key roles in antioxidant protection and redox signaling in the cytoplasm of eukaryotic cells. This system controls cytoplasmic hydrogen peroxide (H_2O_2) concentrations at low oxidative loads [1–3], and plays prominent signaling roles in vascular adaptation [4], mitogenesis [5], inflammation [6], tumorigenesis [7] and apoptosis [8,9]. But despite the numerous studies associating the PTTRS to redox signaling, a consensus about the mechanisms conveying H_2O_2 signals to redox-regulated targets, and how this system integrates signaling and antioxidant protection is yet to emerge.[10–13] Clarifying how the dynamics of the PTTRS in cells relates to the properties and abundances of these proteins is a critical step towards understanding these problems.

This dynamics is largely determined by the redox behavior of the cytoplasmic Prx (Figure 1). These are pentamers of dimers in antiparallel orientation. Each monomer carries a very H_2O_2-reactive thiolate (peroxidatic cysteine, C_P) and a less reactive thiol (resolving cysteine, C_R). H_2O_2 oxidizes the peroxidatic cysteine to a sulfenate (Prx-SO$^-$), which then condenses with the resolving cysteine of the opposing monomer to form a disulfide (Prx-SS). In eukaryotic Prx the rate of this last step is limited by the local unfolding (LU) around the active site that is required to bring the sulfenate close enough to the resolving cysteine.[12] This delay prompts the accumulation of the sulfenate form, and thereby its further oxidation (called "hyperoxidation") by additional H_2O_2 molecules to sulfinate (Prx-SO$_2^-$) and sulfonate (Prx-SO$_3^-$). The conversion to sulfonate irreversibly inactivates the peroxidatic activity. However, Prx-SO$_2^-$ can be slowly reduced to Prx-SO$^-$ at the expense of ATP and reducing equivalents, under catalysis by sulfiredoxin (Srx) [14]. In turn, Prx-SS is reduced by thioredoxin (Trx-S$^-$) eventually returning Prx to its fully folded (FF) thiolate form. Thioredoxin becomes oxidized to a disulfide (Trx-SS) in the process, and this disulfide is reduced by NADPH under catalysis by thioredoxin reductase (TrxR).

The characterization of the responses of the PTTRS to H_2O_2 in a variety of conditions, cell types and organisms highlighted both commonalities and differences, as the following examples illustrate. Treatment of human erythrocytes (5×10^6 cells/mL) with H_2O_2 boluses up to 200 μM caused both PrxII and Trx1 to accumulate in disulfide form, with virtually no Prx hyperoxidation.[1] Hyperoxidation was detectable only in erythrocytes treated with ≥100 μM H_2O_2 boluses after Cat inhibition.[1] In contrast, treatment of Jurkat T cells (10^6 cells/mL) with ≥100 μM H_2O_2 caused extensive hyperoxidation even in absence of Cat inhibition.[1] Remarkably, this happens despite the predominant Prx in Jurkat T cells being PrxI [15], which is more resistant to hyperoxidation than the dominant Prx in erythrocytes, PrxII.[5,16] Extensive

hyperoxidation of PrxI and PrxII was also observed when confluent cultures of human umbilical

vein endothelial cells (HUVEC) were exposed to $\geq$30 μM H_2O_2 boluses.[17]   In these cells

hyperoxidation of PrxII is already extensive, and that of PrxI clearly observable, by 2 min after a

100 μM H_2O_2 bolus.[17]

Sobotta et al. [18] exposed 10^6 HEK293 cells/mL to either 0.2 – 3.7 μM H_2O_2 steady states for

1 h or 2.5 μM – 5 mM H_2O_2 boluses for 5 min. In the former case the fraction of PrxII in disulfide

form increased in a dose-dependent manner from $\approx$5% to 100%. In the latter case that fraction

peaked at $\approx$80% for a 25 μM bolus and progressively decreased for increasing boluses,

presumably due to increasing double hyperoxidation of the dimers preventing disulfide

formation. Tomalin et al. [19] reported that treatment of 2$\times$$10^6$ HEK293 cells/mL with 10 –

80 μM H_2O_2 boluses for 10 min caused a progressive increase in total hyperoxidation only after

a $\approx$20 μM H_2O_2 bolus threshold.

These commonalities and differences prompt important questions with practical relevance for

research in thiol redox regulation and for therapy. To what extent are results obtained in one

organism or cell type generalizable? What among the many factors that may change form cell

type to cell type explain the observed differences in the responses of the PTTRS? What are the

determinants of the observed response thresholds? How can oxidative stress and apoptosis be

most effectively induced in tumor cells? Will different types of tumor cells react in different

ways? Again, understanding the answers to these questions requires clarifying how the

dynamics of the PTTRS relates to the properties and cellular abundances of these proteins.

Mathematical modeling has consistently yielded useful insights about the operation of

antioxidant and thiol redox systems [20–27] and is recognized as an important tool for the

progress of redox biology [28]. Most previous computational studies have focused on accurately

modeling specific cells. Instead, the present work seeks to identify generic principles connecting

design and function in redox signaling and antioxidant protection by the PTTRS, and on

understanding the underpinnings of differences among cell types. Further, we focus on overall

dynamic properties, and not yet on details that hinge on further experimental characterization

of the components. These distinct goals required a distinct modeling approach. Thus, we

proceeded as follows. First, we extensively reviewed the literature and databases to identify (a)

the features of the PTTRS in the cytoplasm of eukaryotic cells that are conserved and most

relevant for its dynamics, and (b) the typical ranges of the kinetic and composition parameters.

This preliminary analysis revealed that most current uncertainties — such as the contribution of

GSH for Prx reduction, or a contribution of generic protein thiols for buffering H_2O_2 and oxidizing

Trx1 — have a minor impact on the overall dynamics of the PTTRS and can be neglected in a first

approach. The quantitative analyses supporting this conclusion are documented in the

Supplementary Information Section 3 (SI3).

Second, we set up a simple coarse grained model of the PTTRS and analyzed it through a

mathematical framework [29,30] that provides an approximated but intelligible comprehensive

description of the relationship between system and molecular properties. The usefulness of this

approach to clarify the functional significance of biological variability has been

demonstrated.[31] This analysis permitted enumerating the qualitatively distinct states and

responses available to the PTTRS and determine closed-form analytical relationships among

H_2O_2 supply rates, protein concentrations and kinetic parameters that take the system to each

state. Importantly, these results do not depend on numerical parameter values, but just on the

order-of-magnitude considerations that informed model set up.

Third, based on selected data in the literature and quantitative proteomics datasets we

estimated the kinetic parameters and cytoplasmic concentrations for human erythrocytes,

hepatocytes, eleven human cell lines, and *S. cerevisiae*. (These estimates are thoroughly

documented in the SI.) We validated the quantitative models by comparing computational

predictions to the most comprehensive quantitative observations of the PTTRS' responses to

H_2O_2 available. In light of the analysis mentioned in the previous paragraph, we then examined

the functional implications of the variation in protein composition and properties among cell

types, and we dissected the underpinnings of commonalities and differences among the

predicted responses.

These analyses show that the PTTRS can *in principle* respond to H_2O_2 supply in many distinct

ways. Nevertheless, once the actual parameter values and composition are considered distinct

tumor cell lines are predicted to show a surprisingly similar response that (a) prevents strong

Trx-S⁻ depletion, (b) favors a gradual moderate accumulation of hyperoxidized Prx at high v_{sup},

and (c) avoids a run-away hyperoxidation of all the Prx. This response hinges on the Prx-SS

reduction capacity just slightly exceeding the maximal steady state Prx-SS production. Its near-

universality over cell lines with quite heterogeneous protein composition is due in part to

previously undocumented strong correlations between the concentrations of TrxR, Srx and Prx.

In turn, erythrocytes, hepatocytes and hepatoma cells are predicted to show a distinct response

where at high v_{sup} Trx-S⁻ is depleted and Prx accumulates mainly as Prx-SS.

In all *nucleated* human cells examined the Prx-SS reduction capacity, which is in most cases

determined by the TrxR activity, defines a threshold separating two different H_2O_2 signaling

regimes. At sub-threshold v_{sup} cytoplasmic H_2O_2 concentrations are determined by Prx, very low

(nM-range) and spatially localized, whereas at supra-threshold v_{sup} H_2O_2 concentrations are

determined by much less active alternative sinks and in the μM range throughout the cytoplasm.

The PTTRS in *S. cerevisiae* is predicted to show a distinct response where at high v_{sup} Prx

accumulates in sulfenate form. This is due to the exceptional stability of TSA1's sulfenate.

The computational predictions are experimentally testable and have important implications for

understanding how the PTTRS integrates redox signaling and antioxidant protection, which are

examined in the Discussion.

Model Formulation

We set up a minimal model that captures the basic features of the PTTRS common to

most cells where it occurs (Figure 1A). It provides a reasonable description of system behavior

in absence of stresses that might deplete NADPH over an extended period. The latter should

only occur under strong stresses or in cells with a compromised pentose phosphate pathway.

Most healthy cells have glucose 6-phosphate (G6P) dehydrogenase (G6PD) activity in large

excess of their capacity to supply G6P, which ensures a fast and effective NADPH supply-demand

coupling and avoids strong depletion [32,33] (SI3.2.9). Indeed, even a 500 μM H_2O_2 bolus caused

just 30% NADPH depletion to human fibroblasts.[34]

Figure 1. A. A simple model of the peroxiredoxin / thioredoxin / thioredoxin reductase system. *The kinetic parameters for each process are indicated near the respective arrow. B. Notation used to designate each phenotypic region.*

In the model, v_{sup} stands for the overall supply of H_2O_2 to the cytoplasm. H_2O_2 can be reduced

by Prx through both Prx-S$^-$ (rate constant k_{Ox}) and Prx-SO$^-$ (k_{Sulf}). The rate constant k_{Ox} can

be regarded as an *effective* rate constant for H_2O_2 reduction by Prx-S$^-$, which allows to consider

a diminished effective peroxidatic activity such as postulated [21] for PrxII in human

erythrocytes. The H_2O_2 can also be cleared by efflux from the cytoplasm, by the activities of

catalase, peroxidases, 1-Cys Prx, and by reaction with other thiolates. These alternative sinks

were aggregated into a single process with first-order kinetics (rate constant k_{Alt}), as

peroxidases should only saturate at very high (>17 μM) intracellular H_2O_2 concentrations

(SI3.2.2.1). With the exception of erythrocytes [21], under low v_{sup} these alternative sinks

contribute little for H_2O_2 clearance in all other cells analyzed (SI3.2.2, Supplementary Table, ST,

6).

For simplicity, we consider each Prx's proximate C_P-C_R pair as an independent functional unit.

The Tpx1 peroxiredoxin of *Schizosaccharomyces pombe* [19] and the human mitochondrial Prx3

[35] may violate this assumption. However, there is no evidence that the same happens to the

same extent for other eukaryotic Prx in the cytoplasm [35]. Further, even in the noted

exceptions, the PTTRS shows qualitatively the same overall dynamics.

The FF → LU transition plus condensation sequence, as well as the Prx-SS reduction plus LU → FF

transition were each aggregated into single steps, with apparent first- and second-order rate

constants k_{Cond} and k_{Red}, respectively.

The reduction of Prx-SO$_2^-$ to Prx-SO$^-$ is treated as a pseudo-first-order process (k_{Srx}). Srx has

relatively low Michaelis constants for ATP and Trx1-S$^-$ [36], which supports this approximation.

Its Michaelis constant for Prx-SO$_2^-$ has not been determined. However, from results in ref. [37]

it can be inferred that the K_M of *Saccharomyces cerevisiae* Srx for Tsa1 is ≈20 μM. This suggests

that Prx-SO$_2^-$ reduction by Srx follows pseudo-first order kinetics with respect to this substrate

in vivo, except under high oxidative stress.

The reduction of Trx-SS was treated as a one-substrate Michaelis-Menten process. This process

is catalyzed by TrxR and follows a ping-pong mechanism that uses NADPH as a second substrate

[38]. However, the $K_{\text{M,TrxR,NADPH}}$ of human TrxR1 is 6 μM [39] and the apparent

$K_{\text{M,TrxR,NADPH}}$ is substantially lower when the enzyme is far from saturation with Trx-SS.

Therefore, physiological concentrations of NADPH should be saturating for TrxR, which justifies

neglecting the influence of this substrate on the reaction rate. Below we write the rate

expression for TrxR as function of a V_{Max}^{App} to highlight that the maximal rate of TrxR may

decrease at very high v_{sup} due to NADPH depletion or to TrxR inactivation by electrophilic lipid

peroxidation products [40].

PrxI can be glutathionylated at C52, C173 and C83, and the extent of glutathionylation was

40%-60% increased (*vs.* untreated controls) 10 min after treatment of A549 or HeLa cells with a

10 μM H_2O_2 bolus.[41] Both Srx and glutaredoxin 1 (Grx1) can catalyze the deglutathionylation

of these residues.[41] Altogether, these findings suggest that glutathione (GSH) might reduce

PrxI-SS and/or PrxI-SO⁻. This hypothesis was recently supported by the observation that GSH plus

Grx1 can reduce both PrxII-SS and PrxII-SO⁻.[42] However, as discussed in SI3.2.4 these reactions

do not dominate the dynamics of the PTTRS and can thus be neglected in a coarse-grained model

for cells where PrxI is the dominant cytoplasmic Prx. They can play a significant role in increasing

resistance to hyperoxidation and reducing Prx-SS in erythrocytes, where PrxII is the dominant

Prx and the TrxR activity is very low.[42] But even here only a very minor fraction of PrxII is

glutathionylated.[42]

Trx-SS can also be reduced by GSH + Grx1, but similarly to the case for Prx this is not the

dominant reductive process [43], and can thus be neglected in a coarse-grained model of the

PTTRS.

Although Trx1-S⁻ can be oxidized by numerous protein disulfides and several enzyme-catalyzed

processes, our estimates (SI3.2.7) indicate that under oxidative stress Prx-SS reduction is the

dominant process oxidizing Trx1-S⁻. In turn, at lower v_{sup} cells' TrxR activity is sufficient to keep

Trx1 strongly reduced despite all the oxidizing processes. Because the objective for the model is

to provide a simple description of the PTTRS dynamics and not a quantitative estimation of the

Trx redox status we neglected the other processes oxidizing Trx-S⁻. For similar reasons we also

neglected the oxidation of Prx-S⁻ by Trx-SS.

Note that the overall reaction for the chemical system represented in this model couples the

reduction of H_2O_2 to water to NADPH oxidation. The overall process is extremely

thermodynamically favorable owing largely to the high redox potential of the first half reaction

[+1.32 V at 25 °C, pH 7 [44]]. Therefore, the PTTRS should operate irreversibly.

The assumptions discussed above and the scheme in Figure 1A translate into the following

system of algebraic differential equations to describe the dynamics of the PTTRS:

234
$$\begin{cases} \dfrac{d\,H_2O_2}{dt} = v_{\sup} - k_{Alt} \times H_2O_2 - k_{Ox} \times Prx\text{-}S^- \times H_2O_2 - k_{Sulf} \times Prx\text{-}SO^- \times H_2O_2 \\[2mm] \dfrac{d\,Prx\text{-}SO^-}{dt} = k_{Ox} \times Prx\text{-}S^- \times H_2O_2 + k_{Srx} \times Prx\text{-}SO_2^- - k_{Sulf} \times Prx\text{-}SO^- \times H_2O_2 - k_{Cond} \times Prx\text{-}SO^- \\[2mm] \dfrac{d\,Prx\text{-}SO_2^-}{dt} = k_{Sulf} \times Prx\text{-}SO^- \times H_2O_2 - k_{Srx} \times Prx\text{-}SO_2^- \\[2mm] \dfrac{d\,Prx\text{-}SS}{dt} = k_{Cond} \times Prx\text{-}SO^- - k_{Red} \times Trx\text{-}SH \times Prx\text{-}SS \\[2mm] \dfrac{d\,Trx\text{-}SS}{dt} = k_{Red} \times Trx\text{-}SH \times Prx\text{-}SS - \dfrac{V_{Max}^{App} \times Trx\text{-}SS}{K_M + Trx\text{-}SS} \\[2mm] Prx_T = Prx\text{-}S^- + Prx\text{-}SS + Prx\text{-}SO^- + Prx\text{-}SO_2^- \\[1mm] Trx_T = Trx\text{-}SH + Trx\text{-}SS \end{cases}$$

(1)

Henceforth, we will denote this model by "Model 1".

For physiological Prx concentrations there are substantial concentration gradients of H_2O_2,

Prx-SO⁻, and Prx-SS over the cytoplasm [13,22,45], which this model does not account for.

However, recent reaction-diffusion simulations [13] confirm that the model provides a good

overall description of the behavior of the average concentrations of these species.

Methods

The systems design space analysis was performed using the algorithms described in SI1,

which extends the applicability of the approach published in ref. [30]. Sensitivity analyses were

performed according to the methodology in refs. [46,47]. Except where otherwise stated, all

these analyses and the numerical simulations were performed in *Mathematica*™ 11 [48].

Parameters for human erythrocytes were estimated as described in ref. [21].   Those for

Saccharomyces cerevisiae and for human Jurkat T, A549, GAMG, HEK293, HeLa, HepG2, K562,

LnCap, MCF-7, RKO, and U-2 OS and hepatocyte cells were estimated from the literature and

databases as described in SI3.

Numerical simulations for a quantitative comparison between computational and experimental

results on PrxII were based in the two-Prx model described in SI4.

Results

A phenotypic map of the PTTRS

We first seek to map the properties of the system as a function of kinetic parameters and protein concentrations. As a starting point, this requires analyzing the steady state solutions of Model 1. However, these solutions cannot be expressed in closed analytical form, and the large number of parameters prevents an effective numerical exploration. We therefore applied the system design space methodology [29,30] to obtain an intelligible approximate description. This methodology subdivides the parameters space into a set of regions. The dynamics in each region is described by a distinct combination of alternatively dominant production and consumption fluxes for each chemical species, and of alternatively dominant concentrations among the forms included in each moiety-conservation cycle. Whenever a region contains a steady state solution, this is guaranteed to be unique and analytically described by a simple product of power laws of the parameters. By the construction of the approximation, these regions represent qualitatively distinct behaviors of the system, and are accordingly denoted by "*phenotypic regions*". The parameters space partitioned into the set of phenotypic regions is called the system's "*design space*".

The construction of the design space for Model 1 is explained in SI1,2. This design space contains 13 regions with positive steady state solutions, but not all of these are biologically plausible. In order to select these, one has to consider the ranges of kinetic parameters and protein concentrations found in real cells. We consider the following three plausibility criteria cumulatively (see SI2 for mathematical definitions).

First, the maximum flux of Prx-SO$_2^-$ reduction is the lowest maximum flux of the system. This is because Srx is an inefficient enzyme [37,49–51] and is much less abundant in cells than the other proteins considered in the model (ST6).

Second, the pseudo-first-order rate constant for H$_2$O$_2$ reduction by Prx-S$^-$ exceeds the rate constant for Prx-SO$^-$ condensation. This follows from the high reactivity and abundance of typical 2-Cys Prx in the cytoplasm, contrasting with the kinetic limitation imposed by the local unfolding to condensation step.

Third, Prx sulfinylation is the slowest among all (aggregated) H$_2$O$_2$-consuming processes in Model 1. This holds with a large margin for all cells we examined (SI3).

Only the 8 phenotypic regions that we describe below satisfy the three plausibility criteria above. We designate each region by a four-character code as explained in Figure 1B. A pictorial

representation of system steady state properties in each of these regions is presented in Table
1. For the corresponding mathematical descriptions of the steady state properties and of the
boundaries of each region in the parameters space please see ST1 and ST2, respectively, in SI2.1.

Phenotypic regions TTPU and TTAU are characterized by the thiol(ate) forms of Prx and Trx being
the dominant ones and differ on whether most of the H_2O_2 is consumed by Prx (TTPU) or by
alternative sinks (TTAU) (Table 1). These regions occur where, cumulatively, the H_2O_2 supply is
low and the TrxR activity is not too low (ST2, Figure 2). In these regions, the concentrations of
Prx-SO⁻, Prx-SS and Trx-SS show an approximately linear response to changes in v_{sup} (ST1, Figure
2).

Region HTAU is characterized by extensive Prx sulfinylation and low Trx oxidation (ST1, Figure
2). This region occurs at very high v_{sup} and not very low TrxR activity (ST2, Figure 2).

Under some conditions, regions TTPU and HTAU partly overlap (Figure 2C). When this occurs,
there is also a region (HTPU) of unstable steady states that coincides with the overlap between
TTPU and HTAU. This feature reveals the possibility of bi-stability and hysteresis in the PTTRS,
which had not been appreciated before. That is, under these conditions as v_{sup} increases up to
the critical value (v_{sup}^{*}) at the right hand side of region TTPU, entering region HTAU, the
concentrations of Prx-SO₂⁻ and H_2O_2 abruptly increase, whereas those of Prx-S⁻, Prx-SO⁻, Prx-SS,
and Trx-SS abruptly decrease. However, as v_{sup} decreases from high values in region HTAU the
opposite transition occurs only at a lower critical value ($v_{sup}^{**} < v_{sup}^{*}$) at the left hand side of
region HTAU, entering region TTPU. The abrupt changes are driven by the following positive
feedback. As v_{sup}, and hence H_2O_2 concentration, increases Prx becomes more hyperoxidized,
depleting Prx-S⁻. As reduction by Prx-S⁻ is the main H_2O_2 sink, this depletion causes a sharp
increase in the concentration of H_2O_2, which accelerates hyperoxidation even further. Biological
systems use toggle switches like this to avoid random back-and-forth switching between
discrete physiological or developmental states, driven by gene expression noise and
environmental fluctuations.[52] On the other hand, the positive feedback just described will
cause near-complete Prx hyperoxidation, from which cells can recover only very slowly due to
the usually very low Srx activity. Therefore, the findings above raise the intriguing question of
whether cells use the PTTRS' hysteretic behavior as part of a stress switch, or avoid it due to its
potentially very deleterious consequences.

Table 1. System states in the physiologically plausible phenotypic regions. *Each phenotypic region is characterized by a hierarchy of concentrations of alternative protein forms (represented by symbols of different sizes) and one or more flux hierarchies (arrows of different widths). Distinct flux hierarchies belong to the same phenotypic region when they originate the same steady state solution. A bar at a reactant tip of an arrow indicates that the reactant in point is saturating for the corresponding process.*

Region	Phenotype
HTPU	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2
TTPU	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2
STAU	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2
HTAU	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2
TTAU	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2
DTAU	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2
DDAU	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2
DDAS	H_2O_2, Prx-SS, Trx-S$^-$, Prx-SO$_2^-$ ⇌ Prx-SO$^-$, Prx-S$^-$, Trx-SS, H_2O_2

Regions STAU and DTAU are characterized by Trx being predominantly in thiol form and the dominant Prx forms being Prx-SO$^-$ or Prx-SS, respectively (Table 1, Figure 2B,D, ST1). Both regions occur at intermediate H_2O_2 supplies and high TrxR activities, only under conditions where regions TTPU and HTAU do not overlap (ST2, Figure 2B,D).

Figure 2. Slices of the design space of the PTTRS over (v_{sup}, V_{Max}^{App}) planes, showing the relative locations of the various phenotypic regions for distinct cell compositions. *The second and third rows show the responses of the fractions of Prx and Trx in each form, TrxR saturation and cytoplasmic H_2O_2 concentration to v_{sup} for the V_{Max} values marked by the dashed horizontal lines in panels A-D. The vertical dotted lines mark region boundaries. H_2O_2 concentration is scaled by maximal value achieved in each plot. Each region is characterized by distinct concentration hierarchies as well as by distinct dependencies on H_2O_2 supply. Boundaries among phenotypic regions correspond approximately to crossover points where these concentration hierarchies or TrxR saturation qualitatively change. The composition ranges yielding each type of section are as follows:*

(A) $k_{Alt} k_{Cond}^{-1} k_{Sulf}^{-1} k_{Srx} < Prx_T < k_{Alt} k_{Ox}^{-1} \wedge \sqrt{k_{Ox} k_{Cond} k_{Sulf}^{-1} k_{Srx}} \, k_{Red}^{-1} < Trx_T < K_M$,

(B) $k_{Alt} k_{Ox}^{-1} < Prx_T < k_{Alt} k_{Cond}^{-1} k_{Sulf}^{-1} k_{Srx} \wedge Trx_T > \max(K_M, k_{Cond} k_{Red}^{-1})$,

(C) $Prx_T > k_{Alt} \max\left(\left(k_{Ox}^{-1}, k_{Cond}^{-1} k_{Sulf}^{-1} k_{Srx}, k_{Cond} k_{Sulf}^{-1} k_{Srx} k_{Red}^{-2} Trx_T^{-2} \right) \right) \wedge Trx_T > K_M$,

(D) $k_{Alt} k_{Ox}^{-1} < Prx_T < k_{Alt} k_{Cond} k_{Sulf}^{-1} k_{Srx} k_{Red}^{-2} Trx_T^{-2} \wedge K_M < Trx_T < k_{Cond} k_{Red}^{-1}$. *Delogaritmized landmark coordinates are as follows:*

$$p_1 : \left(k_{Alt} \sqrt{ \frac{k_{Cond}}{k_{Sulf}} \frac{k_{Srx}}{k_{Ox}} }, Prx_T \sqrt{ \frac{k_{Cond}}{k_{Sulf}} k_{Srx} k_{Ox} } \right), \quad p_2 : \left(k_{Cond} Prx_T, k_{Cond} Prx_T \right), \quad p_3 : \left(k_{Cond} Prx_T, \frac{k_{Alt} k_{Srx}}{k_{Sulf}} \right),$$

$$p_4 : \left(\sqrt{ \frac{k_{Cond}}{k_{Sulf}} k_{Srx} k_{Alt} Prx_T }, \sqrt{ \frac{k_{Cond}}{k_{Sulf}} k_{Srx} k_{Alt} Prx_T } \right),$$

$$p_5 : (V', V'), \quad \left(V' = Prx_T \min \left(k_{Cond}, \sqrt{ \frac{k_{Cond}}{k_{Sulf}} k_{Srx} k_{Ox} }, k_{Red} Trx_T \right) \right) \qquad p_6 : \left(k_{Red} Prx_T Trx_T, k_{Red} Prx_T Trx_T \right),$$

$$p_7 : \left(\frac{k_{Cond} k_{Alt} k_{Srx}}{k_{Sulf} k_{Red} Trx_T}, k_{Red} Prx_T Trx_T \right), \qquad l_1 : V_{Max}^{App} = \frac{k_{Ox} Prx_T}{k_{Alt}} \frac{K_M}{Trx_T} v_{sup} ,$$

$$l_2 : V_{Max}^{App} = \frac{k_{Cond}}{k_{Sulf}} \frac{K_M}{Trx_T} \frac{k_{Alt} k_{Srx} Prx_T}{v_{sup}} , \quad l_3 : V_{Max}^{App} = v_{sup} , \quad l_4 : V_{Max}^{App} = \frac{k_{Cond}}{k_{Sulf}} \frac{k_{Alt} k_{Srx} Prx_T}{v_{sup}} .$$

For multistationarity to occur the ratio between the highest and the lowest value of v_{sup} in the overlap region in (C) must exceed 2.

Finally, regions DDAU and DDAS are characterized by the dominance of the disulfide forms of

both Prx and Trx, and differ on whether TrxR is saturated (DDAS) or not (DDAU) (Table 1, ST1).

They occur at intermediate H_2O_2 supplies and low TrxR activities (Table 1, Figure 2, ST2).

The analysis of the design space permits the following generalizations. First, the system can

always be driven to either TTPU or TTAU by making $v_{\sup}$ sufficiently low, and to HTAU by making

$v_{\sup}$ sufficiently high. However, the latter high $v_{\sup}$ values are not necessarily physiological.

Second, the system can always be driven to regions DDAU and DDAS through a strong enough

inhibition/under-expression of TrxR or Trx, though DDAS becomes unreachable where the

concentration of Trx is too low to saturate TrxR. Third, not all dominance configurations are

feasible or biologically plausible. For instance, whenever Trx-SS is the dominant Trx form Prx-SS

is the dominant Prx form, and whenever Prx-SO$_2^-$ is the dominant Prx form Trx-S$^-$ is the dominant

Trx form. Fourth, only regions TTPU, HTPU and HTAU can overlap. In particular, regions TTAU

and HTAU cannot overlap, which means that the hysteretic behavior described above can only

occur when Prx are the main H_2O_2 consumers under low oxidative loads.

The PTTRS permits many distinct responses to H_2O_2 supply

As illustrated in Figure 2 the eight phenotypic regions can be arranged in several

alternative ways, yielding potentially many qualitatively distinct responses of the various

chemical species to changes in $v_{\sup}$ and V_{Max}^{App}. We have systematically inventoried the

qualitatively different arrangements of the eight phenotypic regions in the $(v_{\sup}, V_{Max}^{App})$ plane

that are possible under physiologically plausible conditions (SI2.2). This analysis reveals that 12

qualitatively distinct generic configurations can occur (SF2). Furthermore, 10 qualitatively

distinct generic types of responses to $v_{\sup}$ are possible, corresponding to as many different

sequences of phenotypic regions. ("Generic" here means that it does not hinge on specific

"pointwise" relationships between parameters.) Six of these generic types of responses are

illustrated in Figure 2, and the conditions where each of the 10 responses hold are listed in ST3.

With relevance for signal transduction, these results demonstrate that the PTTRS can exhibit

many distinct types of responses to $v_{\sup}$. These include: (i) saturation, as for *Trx-SS* and

Prx-SO$^-$ transitioning from TTAU to DDAU (Figure 2A); (ii) ultrasensitivity, as for *Prx-SS*, *Trx-SS*

and *Trx-S$^-$* transitioning from TTPU to DDAS and from DDAS to HTAU (Figure 2B); (iii) hysteresis

(toggle-switch behavior), as for all variables transitioning from DTAU to HTAU over the overlap

region (Figure 2C); (iv) non-monotonic behavior of all variables except *H_2O_2*, *Prx-S$^-$* and *Prx-SO$_2^-$*.

Composition and kinetic parameters of the PTTRS show high variability across cell types

349
350

All the results obtained above depend on just a few generic assumptions about
parameter values and protein concentrations. They suggest a potentially high diversity of
responses of the PTTRS to H_2O_2 supply under the biological plausibility conditions specified
above. But do cells manifest all this diversity, or do they orchestrate protein concentrations and
kinetic parameters to generate a common phenotype? That is the problem we address over the
next sections.

In order to address the previous question and to quantitatively define region boundaries and
their locations relative to physiological operating ranges, parameter values must be known. Until
recently, this requirement would pose an insurmountable obstacle. However, a consistent effort
by several laboratories has provided reliable determinations of most of the necessary kinetic
parameters, and quantitative proteomic methodologies now allow reliable estimates of the
concentrations of all the relevant PTTRS proteins. These developments permitted estimating the
relevant parameters for human erythrocytes (as per ref. [21]) and hepatocytes, one non-cancer
human cell line (HEK293), ten human tumor cell lines, and *S. cerevisiae*. Protein abundances for
the eleven human cell lines were determined in a single laboratory through the same state-of-
the-art methods [53], which adds confidence on the comparability of the results. Furthermore,
a separate quantitative proteomics study [54] obtained protein abundances for human
hepatocytes and for the HepG2 cell line under the same conditions.   The estimates are
documented in SI3, and parameters for all cell types are presented in Table 2. A more detailed
account of the estimated protein concentrations and activities contributing for these aggregated
parameters is presented in ST5. The latter results suggest that at low oxidative loads the 2-Cys
peroxiredoxins consume ~99% of the H_2O_2 in the cytoplasm of human cells. In turn, glutathione
peroxidase 1 (GPx1) is much less abundant and typically consumes <0.2% of the H_2O_2, and Cat
has a negligible contribution because it is located in peroxisomes. The main contributor for
cytoplasmic H_2O_2 clearance other than the typical 2-Cys peroxiredoxins is PrxVI. Despite their
very high PrxII concentrations, erythrocytes are an exception to this pattern, as experimental
evidence [2,55–57] indicates that Cat plays a major role in H_2O_2 clearance in these cells even at
modest H_2O_2 supply rates. This is due to a very high cytoplasmic concentration of Cat and
presumably to a postulated [21] strong and quickly reversible PrxII inhibition Cat.

 Table 2. Summary of estimated parameters for the cell types addressed in this work.

	Prx_T	Trx_T	k_{Alt}	k_{Ox}	k_{Cond}	k_{Sulf}	k_{Red}	k_{Srx}	V_{Max}	K_M
Units:	µM	µM	s^{-1}	$M^{-1}s^{-1}$	s^{-1}	$M^{-1}s^{-1}$	$M^{-1}s^{-1}$	$10^{-3}\ s^{-1}$	mMs^{-1}	µM
A549	49.	29.	44.	4.0×10^7	8.7	1.7×10^3	2.1×10^5	4.5	0.59	1.8
GAMG	76.	44.	65.	4.0×10^7	8.5	2.0×10^3	2.1×10^5	2.6	0.58	1.8
HEK293	1.4×10^2	46.	1.6×10^2	4.0×10^7	7.3	3.7×10^3	2.1×10^5	0.41	0.19	1.8
HeLa	65.	24.	1.8×10^2	4.0×10^7	7.3	3.8×10^3	2.1×10^5	1.1	0.23	1.8
HepG2	93.	27.	1.3×10^2	4.0×10^7	7.1	4.1×10^3	2.1×10^5	2.5	0.12	1.8
Jurkat T	1.7×10^2	36.	85.	4.0×10^7	7.0	4.3×10^3	2.1×10^5	0.52	0.18	1.8
K562	1.0×10^2	28.	1.8×10^2	4.0×10^7	6.9	4.4×10^3	2.1×10^5	0.34	0.13	1.8
LnCap	86.	17.	1.3×10^2	4.0×10^7	5.9	5.8×10^3	2.1×10^5	0.93	0.31	1.8
MCF-7	92.	23.	79.	4.0×10^7	6.4	5.1×10^3	2.1×10^5	3.3	0.23	1.8
RKO	87.	69.	1.4×10^2	4.0×10^7	6.6	4.7×10^3	2.1×10^5	6.6	0.27	1.8
U-2 OS	84.	18.	97.	4.0×10^7	8.0	2.7×10^3	2.1×10^5	2.4	0.34	1.8
HepG2*	1.1×10^2	24.	74.	4.0×10^7	7.4	3.6×10^3	2.1×10^5	1.5	0.047	1.8
Hepatocytes	86.	63.	2.1×10^2	4.0×10^7	7.4	3.7×10^3	2.1×10^5	0.29	0.050	1.8
Erythrocytes	5.8×10^2	0.56	2.6×10^2	3.8×10^5**	1.8	1.2×10^4	2.1×10^5	0.1	0.010	1.8
S. cerevisiae	31.	1.5	34.	4.7×10^7	0.58	7.	1.2×10^6	0.059	1.0	0.8
Max/Min***	3.5	4.1	4.8	1.0	1.5	3.4	1.0	23.	12.	1.0
Max/Min	19.	95.	7.6	1.2×10^2	15.	8.3×10^2	5.7	1.1×10^2	59.	2.2

*Dataset from ref. [54]. **Value reflects the PrxII inhibition postulated in ref. [21]. ***For nucleated human cells.

The last two rows in Table 2 highlight that protein concentrations and kinetic parameters of the

PTTRS can vary by several orders of magnitude over the various cell types and organisms.

Remarkably, both Srx and TrxR activities can vary by over one order of magnitude among the

nucleated human cells in our dataset. This variability makes the question about phenotypic

diversity especially pertinent.

387 The underlying model is consistent with and yields insight on experimental
388 observations

Before analyzing the predicted phenotypes for the various cells one must test if the

underlying model and parameter estimates yield predictions that are consistent with known

experimental observations. Few studies so far have investigated the responses of the PTTRS

quantitatively under steady state conditions. However, experimental observations of cells'

responses to H_2O_2 boluses can also be used to validate model and estimates.

Sobotta *et al*. [18] examined the response of PrxII redox status of HEK293 cells exposed to

$0.2 - 3.7\ \mu M\ H_2O_2$ steady states and to 5 min $2.5 - 5000\ \mu M\ H_2O_2$ boluses. Because PrxI is more

abundant than PrxII in these cells (ST6) the *Prx-SS* variable in Model 1 reflects mainly the

concentration of PrxI-SS. Seeking a more direct quantitative comparison between

computational predictions and these experimental observations we set up the model presented

in SI4. This model ("Model 2") differs from Model 1 by considering the redox cycles of PrxI and

PrxII separately and by considering H_2O_2 exchange between the medium and the cytoplasm

explicitly. Parameterized with the data from Table 2 and ST6 for HEK293 cells, Model 2 shows

qualitative agreement with the experimental observations (SF9). Namely, a near-linear increase

in PrxII-SS with extracellular H_2O_2 concentration up to near-maximal values and the biphasic

response of PrxII-SS to H_2O_2 boluses.

However, the maximal fraction of PrxII disulfide attained is much lower than observed. The

design space analysis above helps tracing the likely source of this discrepancy. The Prx-SS

response curves in Figure 2 show that for high fractions of Prx crosslinking to be achieved with

a near-linear response to v_{sup} as observed, the protein composition should allow attainment of

region DTAU (*e.g.* Figure 2D). The analysis shows that this will happen if the total concentration

of Trx is lower than assumed. Indeed, numerical simulations of Model 2 with the cytoplasmic

concentration of Trx1 set to $\approx 1.5~\mu M$ ($\approx 3\%$ of the previously estimated value) show remarkable

agreement with the observations (SF10A).

Model 2 with $Trx_T = 1.5~\mu M$ also agrees with the experimental observations [19] of a threshold

H_2O_2 bolus beyond which Prx hyperoxidation starts increasing, and intracellular H_2O_2 increases

more steeply (SF10B,C). This threshold behavior has been attributed to the saturation of

alternative H_2O_2 sinks and a buffering effect by generic protein thiols.[19] However, it is

predicted even by Model 1, which includes neither of these effects. A comparison of the

simulated progress curves of the various species for boluses around the threshold value (SF11)

points to the following alternative explanation. The steeper increase in hyperoxidation starts

once the bolus becomes sufficient to fully oxidize Prx to Prx-SS, thereby sharply decreasing the

cytoplasmic H_2O_2 clearance rate. In turn, the ensuing sharp increase in cytoplasmic H_2O_2

concentration causes a strong increase in sulfinylation and accumulation of Prx-SO$_2^-$. Noting that

the full oxidation of the Prx-S$^-$ pool to Prx-SS can only happen when the rate of Prx-S$^-$ oxidation

exceeds the rate of Prx-SS reduction, one can make several experimentally testable predictions

about the factors that influence the threshold value of extracellular H_2O_2. First, the threshold

must be inversely proportional to the membrane permeability (SF12A,B), as for the same

extracellular H_2O_2 the H_2O_2 influx rate is proportional to membrane permeability. There is

growing evidence that cells' permeability to H_2O_2 is largely determined by some aquaporins

[58,59], and the previous prediction can thus be tested through the use of aquaporin inhibitors.

Second, the threshold should be higher the higher the concentration of Trx available in the

cytoplasm (SF12C,D), as Prx-SS reduction in these cells is rate limited by Trx availability. Third,

increasing the TrxR activity should have qualitatively the same effect as increasing the Trx

concentration, but the effect should be much less pronounced in these cells than the latter

(SF12C,D) because here the TrxR activity is not rate-limiting for Trx-SS reduction. In most other

cells examined TrxR activity is rate-limiting for Trx-SS reduction in absence of extensive Trx1

sequestration and should there have a stronger effect on the threshold. Fourth, the activity of

alternative H_2O_2 sinks should have little effect on the threshold in these cells (SF12E,F). This

because while there is Prx-S$^-$ available those sinks contribute little for H_2O_2 elimination and to

determine the cytoplasmic H_2O_2 concentration. Fifth, the total concentration of Prx should have

virtually no effect on the threshold (SF12E,F). This because as Prx-S$^-$ consumes the overwhelming

majority of the H_2O_2, the rate of Prx-SS formation is determined by the rate of H_2O_2 supply and

not influenced by Prx_T. The threshold's dependence on the last four factors was not tested for

HEK293 cells. However, the observations reported in ref. [19] for *Schizosaccharomyces pombe*

agree with all these predictions, which further supports our interpretation of the threshold's

underpinnings.

Further experimental data for model validation can be found in Low *et al.* [1], where the

response of PrxII redox status in Jurkat T cells and erythrocytes to various H_2O_2 boluses is

examined. Model 2, parameterized with the data from Table 2 and ST6 for Jurkat T cells yields

results in very good agreement with the experimental observations for these cells (SF14). A

much sharper threshold response of PrxII-SO$_2$$^-$ than for HEK293 cells is predicted, which the

experimental observations do indeed support. Remarkably, in the case of Jurkat T cells setting

the Trx concentration to 3% or even 30% of the value estimated from the quantitative

proteomics data yields a poor agreement with the experimental observations, which indicates

that Trx1 is not much sequestered in these cells.

Because in erythrocytes PrxII is by far the dominant Prx, one can use Model 1 to simulate the

redox state of PrxII. Parameterizing Model 1 with the data from Table 2 and ST6 for these cells

again yields results in very good agreement with the experimental observations (SF15).

Although some of the simulations above were carried out with Model 2, the single-Prx Model 1

provides a good approximation of the overall Prx redox state (SF16). Thus, altogether, the

results above validate the model and parameter estimates underlying the subsequent analysis,

though the possibility that in some cells just a small fraction of the Trx1 is available to reduce

Prx-SS needs to be considered. To obtain a more general perspective of the predicted

phenotypes of the cell types under consideration we now turn again to the design space analysis,

now informed by the quantitative estimates.

Most human cell lines share a common PTTRS phenotype

When all the Trx1 is considered available to reduce Prx-SS the slices of the quantitative design spaces in the plane (v_{sup}, V_{Max}^{App}) (Figure 3, SF17), and the responses to v_{sup} (Figure 4, SF18) for all cell types reveal the following remarkable patterns.

First, the design spaces based on two quantitative proteomic datasets obtained in distinct laboratories [53,54] for the HepG2 cell line are remarkably consistent (Figure 3A,B). Namely, they both predict that at relatively wide range of intermediate values of v_{sup} ($\approx 40 - 450$ μM s^{-1}) both Prx and Trx accumulate mainly in disulfide form (Figure 4A,B). This accumulation sets in at a sharp v_{sup} threshold and is accompanied by a sharp ultrasensitive increase in the H_2O_2 concentration. Prx-SO$_2^-$ only becomes the predominant Prx form at even higher v_{sup}. This behavior matches response PDS in ST3.

Second, the design space (Figure 3C) and response to v_{sup} (Figure 4C) for hepatocytes are very similar to those for the hepatoma-derived HepG2 cell line. Thus, the distinctive compositional pattern of hepatocytes (Table 2, ST6) does not translate into qualitative differences from HepG2 cells in the response of the PTTRS to H_2O_2.

Third, the responses of human erythrocytes to H_2O_2 supply (Figure 3D, Figure 4D) are predicted to show some unique features, characteristic of Response A in ST3. Namely, at low v_{sup} the PTTRS of erythrocytes operates in phenotypic region TTAU, and not in TTPU as all the other cells examined. Also unlike all the other cells, in erythrocytes the concentration of Trx1 is insufficient to saturate TrxR, and therefore at higher v_{sup} the PTTRS operates in region DDAU and not in region DDAS. Consequently, the Trx-SS concentration should not show the sharp ultrasensitive behavior predicted for hepatocytes and HepG2 cells, and the H_2O_2 and Prx-S$^-$ concentrations also should not show the strongly ultrasensitive behavior predicted for all other human cells in this study. The response of the PTTRS in erythrocytes is otherwise most similar to that of hepatocytes and HepG2 cells. Henceforth, for simplicity, we will denote all the responses that involve substantial Trx-S$^-$ depletion over a wide interval of intermediate v_{sup} (*i.e.*, responses A, ADU, ADS, PDU and PDS from ST3) as "Response D".

Fourth, the remaining 10 human cell lines are predicted to show broadly similar responses that are transitional between the generic responses P, PD, and PDS in ST3. Namely, they share the following features. (i) Little Trx-S$^-$ depletion at medium-high values of v_{sup}. (ii) High Prx-SS

Figure 3. Slices of the design space of the PTTRS over the physiological (v_{sup},V_{Max}^{App}) plane for human cell types and S. cerevisiae as computed for the parameters estimated in ref. [21] and SI3. The black scales inside the plots mark the apparent V_{Max} of TrxR and the values of v_{sup} corresponding to 1 µM, 10 µM and 100 µM extracellular H_2O_2. These values of v_{sup} were estimated based on the known cell permeability and morphology (HeLa, MCF-7, Jurkat T cells, erythrocytes and S. cerevisiae) or assuming $k_{Inf} = 10 s^{-1}$ (all other cells). Note the logarithmic scales. Color codes are as for Figure 2. A, HepG2, ref. [53]; B, HepG2, ref. [54]; C, hepatocytes; D, erythrocytes; E, HEK293; F, Jurkat T; G, HeLa; H, K562; I, MCF-7; J, A549; K, LnCap; L, S. cerevisiae. Design space slices for GaMG, RKO and U-2 OS cells are presented in SF17.

496   accumulation at only a narrow range of v_{sup}, followed by progressively decreasing

497   concentrations with increasing v_{sup}. (iii) An ultrasensitive increase in the cytoplasmic

Figure 4. Responses of the PTTRS to H₂O₂ supply rates for human cell types and S. cerevisiae *as computed for the parameters estimated in ref.* [21] *and SI3. Note the logarithmic scales. The plots were obtained by numerical integration of equations (1) with the parameters in Table 2. Predictions of the responses at $v_{sup} > \approx 0.5$ mM s⁻¹ may be inaccurate due to neglect of NADPH depletion. Color codes are as for Figure 2, except that cytoplasmic H₂O₂ concentrations are scaled by 100 μM. A, HepG2, ref.* [53]; *B, HepG2, ref.* [54]; *C, hepatocytes; D, erythrocytes; E, HEK293; F, Jurkat T; G, HeLa; H, K562; I,MCF-7; J, A549; K, LnCap; L, S. cerevisiae. Responses of GaMG, RKO and U-2 OS cells are presented in SF18.*

concentration of H₂O₂ and Prx-SO₂⁻, and decrease of that of Prx-S⁻ by several orders of magnitude

over this narrow v_{sup} range. Henceforth we will denote this overall behavior as "Response H".

If only 3% of the Trx1 is available to reduce Prx-SS, all the nucleated human cells are predicted

to exhibit Response PD from ST3, where region DTPU appears between regions TTPU and HTAU

(SF19). Accordingly, Prx-SS becomes the dominant Prx form over a range of v_{sup} and the increase

of Prx-SO$_2^-$ with v_{sup} becomes more gradual (SF20). A comparison of Figure 3 and Figure 4 to

SF19 and SF20 (respectively) highlights the following three points. First, a strong Trx oxidation

can be prevented by decreasing the concentration of Trx available to reduce Prx-SS. Second, the

decreased Trx availability shifts the v_{sup} at which Prx-SS becomes dominant (v_{sup}^d) to lower

values. Third, although the strongly decreased Trx availability makes the phenotype of all

nucleated cells qualitatively similar, it increases the variability in cells tolerance to Prx oxidation.

Tracing phenotype to molecular properties

What specific differences and similarities in protein composition underlie the

differences and similarities in the predicted PTTRS phenotypes of human cells?

Response D and the high resistance of PrxII in human erythrocytes to hyperoxidation were

attributed to the low TrxR activity in these cells [1]. The analysis below supports this view and

provides a more complete understanding of the multiple factors underlying this outcome. It

follows from rearranging the boundary conditions in ST2 that for the PTTRS to enter the regions

where both Prx and Trx are mostly oxidized to disulfides (DDAS or DDAU) at some range of v_{sup}

the following condition must hold:

518
$$V_{Max}^{App} \min(1, \frac{Trx_T}{K_M}) <$$
$$< \min\left(\sqrt{\min(k_{Alt}, k_{Ox} Prx_T) \frac{k_{Cond} k_{Srx} Prx_T}{k_{Sulf}}}, \frac{k_{Alt} k_{Cond}}{k_{Sulf}}, k_{Cond} Prx_T, V_{Prx-SS}^- \right).$$
(2)

519 Here

520
$$V_{Prx-SS}^- = k_{Red} Prx_T Trx_T$$
(3)

521 stands for the maximum rate of Prx-SS reduction when not limited by the TrxR activity. For the

522 kinetic parameters and concentration ranges of Prx and Trx in most human cells in this study,

523 $\sqrt{\min(k_{Alt}, k_{Ox} Prx_T) \frac{k_{Cond} k_{Srx} Prx_T}{k_{Sulf}}} < V_{Prx-SS}^- < k_{Cond} Prx_T < \frac{k_{Alt} k_{Cond}}{k_{Sulf}}$ holds, and therefore

524 condition (2) simplifies to:

$$525 \qquad V_{Max}^{App} \min\left(1, \frac{Trx_\mathrm{T}}{K_M}\right) < \sqrt{\min(k_\mathrm{Alt}, k_\mathrm{Ox} Prx_\mathrm{T}) \frac{k_\mathrm{Cond} k_{Srx} Prx_\mathrm{T}}{k_\mathrm{Sulf}}}. \qquad (4)$$

The left hand side of this inequality approximates the maximal possible rate of Trx-SS reduction
in the system, and the right hand side approximates the maximal rate of Prx-SS production (and
thus Trx oxidation) taking into account the balance between hyperoxidation and
Prx-SO$_2^-$ reduction. Therefore, this inequality shows that a maximal Trx reduction rate lower than
the maximal Trx oxidation rate warrants the accumulation of both Prx-SS and Trx-SS at some
range of v_sup.

In cells showing Response D the value of v_sup (v_sup^h) at which Prx-SO$_2^-$ becomes the dominant Prx
form (*i.e.*, the PTTRS crosses from regions DDAU or DDAS into region HTAU) is approximately:

$$534 \qquad v_\mathrm{sup}^h \approx \frac{k_\mathrm{Cond}}{k_\mathrm{Sulf}} k_{Srx} k_\mathrm{Alt} \frac{Prx_\mathrm{T}}{V_{Max}^{App} \min\left(1, \frac{Trx_\mathrm{T}}{K_M}\right)}, \qquad (5)$$

as can be derived from the expressions in ST2. Therefore, a high resistance of Prx to
hyperoxidation in these cells depends on the following factors. (i) The intrinsic resistance of the
dominant Prx to hyperoxidation ($k_\mathrm{Cond} / k_\mathrm{Sulf}$); (ii) a high Srx activity; (iii) the availability of
alternative H$_2$O$_2$ sinks (k_Alt); and (iv) a low Prx reduction turnover, as expressed by the last factor
in expression (5). (The value of v_sup^h for erythrocytes inferred from Figure 3E is underestimated
because Srx reduces PrxII more rapidly than PrxI [60] and because the PrxII-SO$^-$ glutathionylation
rate approaches the condensation rate,[42] thereby inhibiting hyperoxidation to some extent.)

In turn, the value of v_sup (v_sup^d) at which Prx-SS and Trx-SS become the dominant forms (*i.e.*, the
PTTRS crosses from regions TTPU or TTAU into regions DDAU or DDAS) is approximately:

$$544 \qquad v_\mathrm{sup}^d \approx V_{Max}^{App} \max\left(1, \frac{Trx_T}{K_M}\right) \max\left(1, \frac{k_\mathrm{Alt}}{k_\mathrm{Ox} Prx_T}\right) < v_\mathrm{sup}^h, \qquad (6)$$

545 which for cells transitioning from TTPU into DDAS simplifies to:

$$546 \qquad v_\mathrm{sup}^d \approx V_{Max}^{App}. \qquad (7)$$

548 Response H emerges in cells where at some intermediate range of v_{sup} the PTTRS crosses the

549 overlap between regions TTPU, HTPU and HTAU. According to the design space analysis the

550 range of v_{sup} where the overlap occurs is determined by:

$$
\max\left(k_{\text{Srx}} Prx_{\text{T}}, V_{\text{Prx-SS}}^{+}\right) < v_{\text{sup}} <
$$

551
$$
< \min\left(k_{\text{Cond}} Prx_{\text{T}}, \sqrt{\frac{k_{\text{Cond}} k_{\text{Ox}} k_{\text{Srx}}}{k_{\text{Sulf}}}} Prx_{\text{T}}, V_{\text{Prx-SS}}^{-}, V_{Max}^{App}, \frac{V_{Max}^{App}}{K_M} Trx_{\text{T}}\right),
$$
(8)

552 with

$$
V_{\text{Prx-SS}}^{+} = \sqrt{\frac{k_{\text{Cond}}}{k_{\text{Sulf}}} k_{\text{Srx}} k_{\text{Alt}} Prx_T}
$$
(9)

554 representing the maximal steady state rate of Prx oxidation to Prx-SS. This is a particular case of

555 the expression in the right hand side of Expression (4) reflecting the fact that the overlap only

556 occurs where $k_{\text{Ox}} Prx_T > k_{\text{Alt}}$. Taking the kinetic parameters and protein concentration ranges

557 in the human cell lines into account, the overlap range simplifies to:

$$
V_{\text{Prx-SS}}^{+} < v_{\text{sup}} < \min\left(V_{Max}^{App}, V_{\text{Prx-SS}}^{-}\right),
$$
(10)

559 which yields the following condition for occurrence of the overlap:

$$
R = \frac{V_{\text{Prx-SS}}^{+}}{\min\left(V_{Max}^{App}, V_{\text{Prx-SS}}^{-}\right)} < 1.
$$
(11)

Note that the denominator approximates the maximal rate of Prx-SS reduction, and therefore

this expression means that the overlap occurs when this maximal rate exceeds the maximal rate

of Prx-SS production. With A549 cells as the only significant exception, in the examined

nucleated human cells it is the TrxR activity that limits the rate of Prx-SS reduction, in absence

of substantial Trx1 sequestration.

The overlap between regions indicates that multistability might occur, but the approximations

in the design space analysis tend to underestimate the minimum and overestimate the

maximum values of v_{sup} for overlap (Figure 2C). As result, only Jurkat T cells are numerically

predicted to show hysteresis over a very narrow v_{sup} range (Figure 4F). But remarkably all

Response H cells with known morphometry have R values in the range 0.61 to 0.76 (except

$R=0.34$ for LnCap cells as an outlier), indicating that their PTTRS operates at the margins of the

overlap region. For progressively lower values of this ratio the switch from low to high

hyperoxidation becomes increasingly ultrasensitive and then hysteretic. Importantly, the

maximum fraction of Prx in Prx-SS form and the maximum fractions of oxidized Trx attained at

the switch get progressively lower.

For cells predicted to show Response H, $Prx\text{-}SO_2^-$ becomes the dominant Prx form at

$$v_{\sup}^{h'} = \min\left(V_{Max}^{App}, k_{\text{Cond}} Prx_{\text{T}}, \sqrt{\frac{k_{\text{Cond}}}{k_{\text{Sulf}}} k_{\text{Srx}} k_{\text{Ox}}} \, Prx_{\text{T}}, V_{\text{Prx-SS}}^- \right),$$ (12)

which in most cells corresponds to

$$v_{\sup}^{h'} \approx V_{Max}^{App}.$$ (13)

R is higher than 1 when V_{Max}^{App} or Trx_T are too low to cope with the maximum Prx-SS production

rate. When the low V_{Max}^{App} is the culprit, Response D ensues, as discussed above. In turn, when

V_{Max}^{App} is high enough but Trx_T is not, Prx can accumulate as Prx-SS without substantial Trx

oxidation over a range of $v_{\sup}$ (Region DTAU). This will happen in virtually all nucleated human

cells in our set if only 3% of the Trx_T estimated from the proteomic datasets is available to

reduce Prx-SS as estimated above for HEK293 cells (SF19,20).

More precise conditions for occurrence of this behavior can be derived by rearranging the

boundary conditions for region DTAU in ST2:

$$V_{\text{Prx-SS}}^- < \min\left(\sqrt{\min(k_{\text{Alt}}, k_{\text{Ox}} Prx_{\text{T}}) \frac{k_{\text{Cond}} k_{\text{Srx}} Prx_{\text{T}}}{k_{\text{Sulf}}}}, k_{\text{Alt}} \frac{k_{\text{Cond}}}{k_{\text{Sulf}}}, k_{\text{Cond}} Prx_{\text{T}}, V_{Max}^{App} \right),$$ (14)

which for the kinetic parameters and protein concentration ranges in the human cell lines in this

study simplifies to

$$V_{\text{Prx-SS}}^- < \min\left(V_{\text{Prx-SS}}^+, V_{Max}^{App} \right).$$ (15)

Under these conditions, $Prx\text{-}SO_2^-$ becomes the dominant Prx at [compare to (5)]

$$v_{\sup}^{h'} \approx \frac{k_{\text{Cond}}}{k_{\text{Sulf}}} k_{\text{Srx}} k_{\text{Alt}} \frac{1}{k_{\text{Red}} Trx_{\text{T}}},$$ (16)

reflecting the fact that the Prx reduction turnover is then determined by Trx availability. In turn,

Prx-SS becomes the dominant form at

596
$$v_{\text{sup}}^{d'} \approx V_{\text{Prx-SS}}^{-} < v_{\text{sup}}^{h'}. \tag{17}$$

597 As R approaches 1, v_{sup}^{d}, v_{sup}^{h}, $v_{\text{sup}}^{d'}$ and $v_{\text{sup}}^{h'}$ converge at

598
$$v_{\text{sup}}^{h} \approx V_{\text{Prx-SS}}^{+}, \tag{18}$$

599 which is the minimum value of v_{sup} at which Prx-SO$_2^-$ can become the dominant Prx form (Figure

600 2C).

Enzyme activities in the PTTRS are correlated over cell lines

Is the similarity of the responses among human cell lines due to similar concentrations

of the PTTRS proteins in all cells, or to correlations among the protein concentrations over cells?

The following two observations support the latter possibility. First, Table 2 shows that there is

substantial compositional heterogeneity among cell lines. For instance, k_{Srx} varies over a

$\approx$20-fold range over cells showing Response H. Second, the values of k_{Srx} over these cells are

very strongly correlated with the k_{Srx} threshold value,

608
$$k_{\text{Srx}}^{*} = \frac{k_{\text{Sulf}}}{k_{\text{Cond}}} \frac{\min\left(V_{\text{Max}}^{\text{App}}, V_{\text{Prx-SS}}^{-}\right)^{2}}{k_{\text{Alt}} Prx_{T}}, \tag{19}$$

for occurrence of the overlap (Figure 5A).

A statistical analysis reveals three prominent features of the orchestration among the

parameters of the PTTRS over cells predicted to show Response H. First, a positive correlation

between $V_{\text{Max}}^{\text{App}}$ and k_{Srx} (Figure 5B). Remarkably, $V_{\text{Max}}^{\text{App}}$ scales approximately as $\sqrt{k_{\text{Srx}}}$ over

these cell lines (Figure 5B), suggesting that the relative concentrations of TrxR and Srx are

balanced in such way as to make the ratio between the maximal Prx-SS production rate and the

maximal Trx-SS reduction rate approximately invariant over cell lines.

Second, a negative correlation between k_{Srx} and Prx_{T} (Figure 5D). In Region TTPU, where most

cells operate in absence of stress, the concentration of Prx-SO$_2^-$ is inversely proportional to the

product $k_{\text{Srx}} Prx_{T}$ (ST2). Furthermore, the value of v_{sup} at which Prx-SO$_2^-$ becomes the

dominant Prx form in Response H cells is approximately proportional to $\sqrt{k_{\text{Srx}} Prx_{T}}$. Therefore,

the negative correlation between k_{Srx} and Prx_{T} attenuates the heterogeneity of

Prx-SO$_2^-$ concentration and hyperoxidation resistance among cell lines.

Figure 5. Correlations between parameters over cell lines. *(A)* k_{Srx} *strongly correlates with the minimum value* (k_{Srx}^*) *necessary to avoid the overlap region: Spearman rank correlation* (σ_S) *0.89, (p=0.0068) for all cell lines with Response H and known morphometry. Black dashed line indicates* $k_{Srx} = k_{Srx}^*$. *Red dots, cells showing Response D; green, cells with unknown morphometry showing response H; blue, cells with known morphometry showing response H. The following parameters are strongly correlated over cells with known morphometry showing Response H. (B) The activities of TrxR and Srx:* σ_S= 0.81, p= 0.015. *Dashed line: best log-log fit over Response H cells with known morphometry, yielding a scaling exponent 0.38±0.13 (R^2= 0.66). (C) The activity of TrxR and the total concentration of PrxI+PrxII:* σ_S= -0.86, p= 0.0065. *Dashed line: best log-log fit over Response H cells with known morphometry, yielding a scaling exponent -0.87±0.31 (R^2= 0.57). (D) The activity of Srx and the total concentration of PrxI+PrxII:* σ_S= -0.74, p= 0.037. *Dashed line: best log-log fit over Response H cells with known morphometry, yielding a scaling exponent -0.29±0.11 (R^2= 0.51). Note the logarithmic scales.*

Third, a strong negative correlation between V_{Max}^{App} and Prx_T (Figure 5C). This correlation may

be just a consequence of the previous ones, as it lacks an obvious direct functional relevance.

Interestingly, V_{Max}^{App} correlates with $V_{Prx\text{-}SS}^+$ over Response H cells (σ_S= 0.65, p= 0.043), but Trx_T

and $V_{Prx\text{-}SS}^-$ do not (σ_S= 0.28, p= 0.42; σ_S= 0.018, p= 0.96, respectively). This is consistent with

the estimates from the data in Table 2 indicating that Prx-SS reduction is rate limited by TrxR

activity and not by Trx1 availability in most human cells. It also consistent with the notions that

Prx-SS reduction is the process with the greatest capacity to drive Trx1 oxidation (see SI3.2.7)

and that Trx1 is not strongly sequestered in most cells.

Finally, Figure 5 also highlights that Response D cells in our dataset have Srx and Prx

concentrations in the range of the other cells but have disproportionally low TrxR.

The PTTRS in yeast has distinctive properties

Does the dynamics of the PTTRS in other eukaryotic organisms differ from that in human

cells? In order to address this question we estimated the parameters for *S. cerevisiae* (SI3.4).

The design space analysis (Figure 3L) and numerical simulations of Model 1 (Figure 4L) based on

these parameters predict that the responses of the PTTRS in the yeast depart from those

described above for human cells in at least the following three aspects. First,

Prx-SO⁻ concentrations are ≈7-fold higher than those of Prx-SS over all the range of v_{sup} , whereas

in human cells Prx-SO⁻ concentrations are roughly similar to those of Prx-SS. Second, at

intermediate values of v_{sup} Prx accumulates mainly as Prx-SO⁻, and this remains the dominant

Prx species over a wide range of v_{sup} values. Third, the concentration of Prx-SO₂⁻ increases more

gradually with v_{sup} than in human cells.

Table 2 shows many differences in protein composition and kinetic parameters between yeast

and human cells. Which of these determine the differences between the behavior of the PTTRS

in the yeast and in human cells? The high Prx-SO⁻/Prx-SS ratio in the yeast is explained as follows.

The results in ST1 show that in all the phenotypic regions that can be attained in cells with high

TrxR activity (*i.e.,* the *T*U regions) the ratio between the concentrations of Prx-SO⁻ and Prx-SS

is approximated by $k_{\text{Red}} Trx_T / k_{\text{Cond}}$. The about one order of magnitude lower cytoplasmic Trx

concentration in the yeast is compensated by a ≈6-fold higher k_{Red} (SI3.4.3), yielding a similar

value of $k_{\text{Red}} Trx_T$. However, the value of k_{Cond} is >10-fold lower for yeast's Tsa1 than for

human PrxI, explaining the higher ratio.

In turn, the following conditions for Prx-SO⁻ to become the dominant Prx species over a range of

v_{sup} can be derived from the results in ST2 for region STAU:

654
$$k_{\text{Cond}} Prx_T < \min\left(V_{\text{Prx-SS}}^-, \frac{k_{\text{Srx}}}{k_{\text{Sulf}}} k_{\text{Alt}}, \frac{k_{\text{Srx}}}{k_{\text{Sulf}}} k_{\text{Ox}} Prx_T, V_{Max}^{App}, V_{Max}^{App} \frac{Trx_T}{K_M} \right) \wedge k_{\text{Sulf}} Prx_T < k_{\text{Alt}} \qquad (20)$$

Both yeast and human cells fulfill the condition $k_{Sulf} Prx_T < k_{Alt}$. However, only yeast cells fulfill

the condition $k_{Cond} Prx_T < \dfrac{k_{Srx}}{k_{Sulf}} k_{Alt}$. This is due to a combination of factors: (a) both k_{Cond}

and Prx_T are lower in the yeast; (b) although both k_{Srx} and k_{Alt} are also lower in the yeast,

k_{Sulf} is much lower than in the human cells, making $\dfrac{k_{Srx}}{k_{Sulf}} k_{Alt}$ higher in the yeast. Therefore

this behavior can be ascribed to the much higher stability of Tsa1's sulfenate relative to human

PrxI/II-SO⁻, and to the low Tsa1+Tsa2 concentration in the yeast.

The more gradual increase in Trx hyperoxidation in the yeast is explained by the following

considerations. In the intermediate region STAU the concentration of Prx-SO⁻ is practically

constant with v_{sup}, as most of the Prx was accumulated in this form. In turn, the concentration

of H_2O_2 increases linearly with v_{sup}. As a consequence, the rate of hyperoxidation increases

linearly with v_{sup}, rather than quadratically as in region TTPU.

Discussion

The PTTRS can respond to H_2O_2 supply in multiple ways

The results provide an approximated but simple description of the responses of the

PTTRS in the cytoplasm of eukaryotic cells to physiological H_2O_2 supply rates (v_{sup}), and of how

these responses depend on protein concentration and kinetic parameters. The approximations

were thoroughly justified in the previous sections and in SI3, and are valid for v_{sup} values that

are insufficient to strongly deplete NADPH. For simplicity the model neglected the strong

cytoplasmic concentration gradients of H_2O_2, Prx-SO⁻ and Prx-SS expected under low v_{sup} [13].

The concentrations discussed below are thus spatially-averaged cytoplasmic concentrations.

The analysis highlights the versatility of the PTTRS. For physiologically plausible parameters the

PTTRS can exhibit eight qualitatively distinct types of steady states ("phenotypes"),

corresponding to distinct hierarchies among fluxes and concentrations of the various Prx and

Trx forms (Table 1, Figure 2, ST1). Two of these phenotypes allow the maintenance of Prx and

Trx mostly in thiolate form and of strong concentration gradients. The remaining six are stress

phenotypes where Prx is predominantly oxidized and gradients collapse.

The PTTRS can transition across 10 qualitatively distinct sequences of phenotypes as v_{sup}

increases from very low to very high values, defining as many distinct responses. In the process,

the PTTRS has the potential to generate, proportional, saturable, ultrasensitive, non-monotonic,

and even hysteretic behaviors. Cells may in principle take advantage of this diversity of behaviors

to achieve distinct modes of regulation of downstream processes by H_2O_2 supply rate signals.

We mapped systematically the relationships among kinetic parameters and protein

concentrations that make each of these phenotypes and responses emerge (ST2,3), and we

derived simple approximations for the main PTTRS response thresholds. These phenotypes and

thresholds usually depend on an interplay among multiple factors, as is well illustrated by the

sensitivity to hyperoxidation *in vivo*. As shown by expressions (5) and (16) this property depends

not only on the Prx's intrinsic susceptibility of the Prx and on the Srx activity but also on the

activity of non-Prx H_2O_2 sinks, on the Prx concentration and on the Prx-SS reduction capacity of

the cytoplasm. Proper interpretation of the functional significance of gene expression, protein

abundance or metabolic changes must take this interplay into account, which is difficult to

accomplish without the help of modeling approaches such as illustrated here.

The quantitative model is consistent with experimental observations and yields

novel insights

In order to examine what of the potential behaviors discussed above can be realized in

real cells, we estimated the necessary protein concentrations and kinetic parameters for ten

human cancer cell lines, one non-cancer human cell line (HEK293), two differentiated human

cell types (hepatocytes and erythrocytes), and *S. cerevisiae*. These estimates are based on

quantitative proteomics data and published kinetic data. They are also thoroughly documented

in SI3 and summarized in Table 2 and ST6, which we hope will provide a useful reference for

quantitative redox biology.

With these data we first tested if simulations based on the parameterized models yielded results

in agreement with available quantitative experimental data for human erythrocytes, Jurkat T

cells and HEK293 cells. Computational predictions for the first two cell types were in very good

agreement with the experimental observations (SF14,15).[1] Of note, they fully capture the

observations that whereas treatment of 5×10^6 erythrocytes/mL with H_2O_2 boluses up to 200 μM

caused PrxII to accumulate in disulfide form and little Prx hyperoxidation, treatment of 10^6

Jurkat T cells/mL with 100 μM or 200 μM H_2O_2 boluses caused extensive hyperoxidation [1].

In turn, the simulations for HEK293 cells agree qualitatively with the observations in ref. [18],

but predict much lower Prx-SS fractions for the same extracellular H_2O_2 concentrations. This

discrepancy is eliminated if only $\approx$3% (1.5 μM) of the Trx1 concentration estimated from the

proteomic dataset is available to reduce Prx-SS (SF10A). Because the Trx1 concentrations Table

2 are in the range determined by other methods (see SI3.2.6) [61,62], the most likely explanation

for the discrepancy is the sequestration of Trx1 in the nucleus [63,64] and/or in complexes with

other proteins [65].

The following observations suggest that a larger fraction of Trx1 is available for Prx-SS reduction

in other cell lines. First, the small fraction of PrxII oxidized to PrxII-SS in Jurkat T cells exposed to

various H_2O_2 boluses in ref. [1] is not consistent with extensive Trx1 sequestration (SF14).

Second, the TrxR activity, and not the Trx1 concentration, correlates with the maximum steady

state rate of Prx-SS production over Response H cells, which suggests that in most cells Prx-SS

reduction is limited by TrxR activity and not by Trx1 availability.

Threshold behavior of Prx hyperoxidation and cytoplasmic H_2O_2 is intrinsic to the

PTTRS dynamics

Surprisingly, this simple model for HEK293 cells (with $Trx_T = 1.5\ \mu M$) shows very good

agreement with the experimental observations [19] of a threshold H_2O_2 bolus beyond which Prx

hyperoxidation starts increasing, and intracellular H_2O_2 increases more steeply (SF10B,C).

Because the model considers neither a buffering effect by generic protein thiols nor saturable

alternative H_2O_2 sinks, the observed agreement strongly argues against inferring the occurrence

of such phenomena from the observed threshold behavior. Moreover, the balance of

experimental evidence does not support the notion that generic (non-redoxin) protein thiols can

provide a significant buffering capacity (see SI3.2.2.3).　Instead, our results (SF11) indicate that

the threshold behavior described in ref. [19] is intrinsic to the dynamics of the PTTRS. Concretely,

the steeper increase in cytoplasmic H_2O_2 concentration and hyperoxidation starts once the bolus

becomes sufficient to fully oxidize Prx to Prx-SS, thereby sharply decreasing the cytoplasmic H_2O_2

clearance rate. The cytoplasmic H_2O_2 concentration then becomes sufficient to cause

accumulation of Prx-SO$_2^-$. Further supporting this interpretation, simulations using Model 1 to

analyze the influence of the Trx concentration, TrxR activity, activity of alternative H_2O_2 sinks

and Prx concentration on the threshold (SF12) agree with the observations in ref. [19]. There is

a close correspondence between these thresholds and the v_{sup} threshold marking the border of

the TTPU region in the design space, whose physiological significance we will discuss below.

Namely, they all reflect the saturation of cells' Prx-SS reduction capacity.

Human cell lines are predicted to show similar responses to H_2O_2 supply

Once applied to the quantitative estimates, the phenotypic map we derived allowed to

make quantitative predictions about the responses of the PTTRS to H_2O_2 supply and TrxR

modulation and to classify distinct cell types according to the predicted behavior of their PTTRS.

This analysis revealed several intriguing commonalities and differences, which we discuss below.

When all the Trx1 is available to reduce Prx-SS in the cytoplasm, the 13 human cell types in our

sample are predicted show just two major types of responses to v_{sup} despite the diversity of cell

types (ST4), and the variability in protein concentrations (ST6). Namely, in "Response D"

intermediate values of v_{sup} lead to oxidation of both Prx and Trx to disulfide forms, as well as to

Trx-S⁻ depletion. In this response substantial hyperoxidation occurs only at very high v_{sup} . In

turn, in "Response H" there is modest Trx-S⁻ depletion at any v_{sup}, and the steady state fraction

of Prx-SS only reaches high values at a very narrow range of v_{sup} . In contrast, responses including

an extensive accumulation of Prx-SS or Prx-SO⁻ without Trx-S⁻ depletion over a wide range of

v_{sup}, or an abrupt bi-stable switch between a low and a high hyperoxidation state are possible

(Figure 2, ST3) but not predicted to occur in any of the cells considered. (Jurkat T cells may show

bistability over a tiny v_{sup} range but this is strongly dependent on parameter uncertainties.)

Remarkably, all but one of the cell lines are predicted to exhibit Response H, and the two

differentiated human cell types are predicted to exhibit Response D. The exception is the

hepatoma HepG2 cell line. The model parameterizations based on both independent proteomic

datasets for HepG2 [53,54] consistently lead to the prediction that these cells exhibit the same

type of response as hepatocytes: Response D. This observation raises the question of whether

cancer cell lines tend to respond to v_{sup} similarly to the differentiated cell types they derive

from, but unfortunately we are unaware of other data sets that would permit such a comparison

for other cell types.

Our analysis revealed the underpinnings of the predicted similarities and distinctions among the

PTTRS responses of human cells. Response D occurs in cells where the maximal Trx reduction

rate is substantially lower than the maximal Trx-SS production rate [Expression (4)]. Response H

occurs where the TrxR activity and/or the maximal Prx-SS reduction slightly exceed the maximal

Prx-SS production rate [$0.3 \leq R \leq 1$ in expression (11)].

In turn, when just ≈3% of the Trx1 is considered available to reduce Prx-SS, all *nucleated* human

cells in our sample are predicted to show Response PD (ST3), which differs from Response H in

that Prx-SS becomes the predominant Prx form over a wide range of v_{sup} (SF19,20). The

threshold value of v_{sup} at which the Prx-S⁻ pool collapses also decreases. While for Response D

and for most cells showing Response H this threshold occurs at $v_{\sup} \approx V_{Max}^{App} < V_{\text{Prx-SS}}^{-}$

[expressions (7) and (13)], for Response PD it occurs at $v_{\sup} \approx V_{\text{Prx-SS}}^{-} < V_{Max}^{App}$ [expression (17)].

While the conditions above for occurrence of each type of response are intuitive and nearly

trivial when stated in terms of maximum steady state fluxes, these fluxes have a more complex

dependence on the kinetic parameters and protein concentrations. However, the design space

approach permitted deriving relatively simple approximations that can be straightforwardly

used to predict responses and threshold values.

Landmarks of the quantitative design space have a physiological correspondence

Irrespective of the predicted Response type, under physiological, non-stress conditions

all human cells analyzed here operate far within regions TTAU (erythrocytes) or TTPU (all others).

As $v_{\sup}$ increases, the fraction of Prx in oxidized forms gradually increases. Then, as $v_{\sup}$

approaches 50–500 μM s^{-1}, corresponding to 5–50 μM extracellular H_2O_2, nucleated human cells

transition into regions DTAU, DDAS or HTAU, where Prx-SS or Prx-SO$_2^-$ predominate (Figure

3,Figure 4,SF17,18). This transition is accompanied by an ultrasensitive increase of the

intracellular H_2O_2 concentration from the 1–10 nM to the μM range over a relatively narrow

range of $v_{\sup}$ (Figure 4, SF18). The sharp increase is due to cellular H_2O_2 sinks other than PrxI and

PrxII being less active towards H_2O_2 (PrxVI), much less abundant (GPx1) and/or not directly

accessible from the cytoplasm (Cat). The inability of these sinks to maintain a strong

transmembrane H_2O_2 gradient is experimentally demonstrated by the observation that partial

Cat inhibition has a substantial effect on the pseudo-first order rate constant (k_{cells}) for the

consumption of extracellular H_2O_2 boluses that are sufficient to extensively oxidize PrxI and PrxII

[66,67]: otherwise H_2O_2 consumption would be strongly limited by the membrane permeation

step and virtually insensitive to Cat inhibition (see SI3.2.1). It is also supported by recent

experiments in living zebra fish.[59] In turn, a strong transmembrane gradient when Prx-S$^-$ is

predominant is predicted by reaction-diffusion models [13,68] and was experimentally

estimated as $\approx$650 for HeLa cells [68].

Could the Prx-S$^-$ collapse and ensuing sharp increase in cytoplasmic H_2O_2 associated to the

transition from TTPU to the stress regions mark the threshold for H_2O_2 toxicity? Antunes &

Cadenas [69] found that steady state H_2O_2 induces apoptosis of Jurkat T cells only above a

threshold extracellular concentration. Moreover, this threshold is very sharp, such that doubling

the H_2O_2 concentration from the threshold value causes the fraction of apoptosing cells to

increase from control to saturation values. The threshold is also dependent on exposure

duration, suggesting that H_2O_2 toxicity manifests only beyond a given dosage

(concentration×time) threshold.[69] Huang & Sikes [68] recently reported similar findings for

HeLa cells with respect to intracellularly produced H_2O_2 and further elaborated this concept.

Interestingly, the extracellular H_2O_2 concentration where the TTPU→HTAU transition in Jurkat

T cells is expected (Figure 3F, Figure 4F) corresponds approximately to the steady state

concentration observed to cause apoptosis of 50% of these cells [69]. Furthermore, steady state

extracellular H_2O_2 concentrations that are just enough to fully oxidize PrxII to PrxII-SS, but not

lower concentrations, cause apoptosis of HEK239 cells.[18] According to our simulations of the

PTTRS in these cells (SF13), under these circumstances PrxI should also become fully oxidized

and intracellular H_2O_2 should attain the μM range. Therefore, these observations support the

notion that the onset of H_2O_2 toxicity is associated to the collapse of the Prx-S⁻ pool. Accordingly,

it should be modulated by the TrxR activity in cells showing Responses H or D, or Trx1 availability

in cells where extensive Trx1 sequestration dictates Response PD. These predictions are

experimentally testable.

The considerations above question the assertion that perturbations of the intracellular H_2O_2

concentration in the order of 10 nM can induce apoptosis. This assertion was based on a

re-evaluation of the transmembrane H_2O_2 gradient to account for the activity of the cytoplasmic

Prx.[68] This estimated ≈650-fold transmembrane gradient applies where the H_2O_2 supply rate

to the cytoplasm is insufficient to strongly oxidize Prx (*i.e.*, well within region TTPU). However,

apoptosis seems to be induced only upon the collapse of the Prx-S⁻ pool. Under these conditions

the transmembrane gradient should be no higher than the ≈7-fold determined in ref. [70], and

intracellular H_2O_2 concentrations should then approach the μM range [69].

Could the changes in Prx and/or Trx redox state that occur as the PTTRS transitions from region

TTPU to the stress regions directly regulate apoptosis? Apoptosis induction by H_2O_2 is mediated

by activation of apoptosis signal-regulating kinase 1 (ASK1). In the most consensual redox

regulation model, this protein is inhibited by non-covalent binding of Trx1-S⁻ to its Trx1-binding

domain, and released from inhibition by Trx1-S⁻ oxidation to Trx-SS.[71] However, the

dissociation constant for the complexes of Trx1-S⁻ and Trx1-SS with ASK1's Trx1-binding domain

— 0.3±0.1 μM [72] and 4±2 μM [73], respectively — are *both* much lower than typical

cytoplasmic Trx1 concentrations (Table 2). Therefore, not even the full oxidation of Trx can

dissociate the inhibitory Trx1:ASK1 complex unless the dissociation from Trx1-SS is coadjuvated

by other factors. These might include destabilizing interactions with other proteins (e.g. tumor

necrosis factor receptor-associated factors 2 and 6), extensive Trx1 sequestration and/or

oxidation of ASK1's Cys residues.[74] The latter might be driven by the recently characterized

redox relay mediated by Prx1-SS.[9] It is thus tempting to speculate that the PrxI-SS

accumulation as cells cross from phenotypic region TTPU to DTAU, DDAS or HTAU might thereby

trigger apoptosis. However, ASK1 regulation is complex and still poorly understood [75], and

none of the proposed models explains, for instance, the iron dependence of H_2O_2-induced

apoptosis [69].

Although if sustained over a long time a strong oxidation of the Prx-S$^-$ pool may trigger cell death,

temporary strong Prx oxidation does occur in the normal physiology of higher organisms. The

events associated to zebra fish wounding illustrate this point: 2–5 µM extracellular H_2O_2

concentrations are attained within ≈30 nm of wound margins during tens of minutes, which

strongly oxidize the cells' Prx-S$^-$ pool.[59] Studying the steady state response to H_2O_2 supply rates

beyond region TTPU yields insight on cells' responses to such temporary H_2O_2 surges. In all

human cells examined in this work, sudden exposure to v_{sup} beyond the threshold initially

causes a short-lived surge in Prx-SO$^-$ and then extensively oxidizes Prx-S$^-$ to Prx-SS, dramatically

increasing the cytoplasmic H_2O_2 concentration within seconds (SF21). In Response H and

Response D cells, but not in Response PD cells, Trx-S$^-$ is also rapidly and extensively oxidized. In

turn, Prx-SO$_2^-$ accumulates much slower. In Response H cells, Prx-SO$_2^-$ may eventually

accumulate as the dominant Prx form thereby relieving the load on the Trx-S$^-$ pool. However, in

other cells this would only happen at v_{sup} beyond the thresholds defined by expressions (5), (16)

which cells are unlikely to survive. Recovery of the Prx-S$^-$ pool from Prx-SS once the stimulus

stops can occur in seconds, but recovery from Prx-SO$_2^-$ can take hours.

Dynamics of the PTTRS defines two distinct H_2O_2 signaling regimes

The considerations above have important implications for redox signaling in the

cytoplasm of eukaryotic cells. The much higher activity of PrxI and PrxII as H_2O_2 reductants

relative to the alternative H_2O_2 sinks determines the existence of two very distinct H_2O_2 signaling

regimes with physiological relevance. For v_{sup} below the cytoplasmic capacity to reduce Prx-SS,

cytoplasmic H_2O_2 concentrations are determined by the Prx-S$^-$ pool. Under these conditions

there are strong H_2O_2 concentration gradients both across the cell membrane and over the

cytoplasm.[13,68] And even though cytoplasmic H_2O_2 concentrations are much higher near H_2O_2

supply sites than elsewhere, they are far too low to directly oxidize thiolates other than those in

the active sites of peroxiredoxins and peroxidases in a signaling time frame even there.[13]

Therefore, in this regime Prx must act as primary H_2O_2 sensors, which then actuate regulatory

targets through localized[13] redox relays [9,10,76–78] or non-covalent binding/release of the

oxidized Prx forms [11]. H_2O_2 chemotaxis by leukocytes far away from wound borders may be

an example of H_2O_2 signaling in this regime.[59]

In turn, at v_{sup} above the cytoplasmic capacity to reduce Prx-SS, cytoplasmic H_2O_2

concentrations are determined by the activities of peroxiredoxin VI, GPx1 and Cat. These

activities are, collectively, too low to impose a significant H_2O_2 gradient over the cytoplasm or

even a very large transmembrane gradient. Thus, H_2O_2 concentrations are expected to be in the

μM range throughout the cytoplasm. However, as per the previous section, cells are unlikely to

survive such H_2O_2 concentrations for more than a few tens of minutes, which constrains the

redox targets that can be directly actuated by H_2O_2 even in this regime: For 50% of a target's

molecules to be oxidized within 1h by 1 μM H_2O_2 the oxidation rate constant must exceed

$\ln(2)/(3600 \text{ s} \times 10^{-6} \text{ M}) = 193$ $M^{-1}s^{-1}$. Indeed, several regulatory targets that are involved in

adaptation to oxidative stress and thus should be actuated in this regime and not in the low-v_{sup}

one, have H_2O_2 reactivities in this range. These include glyceraldehyde 3-phosphate

dehydrogenase (GAPDH, k= 500 $M^{-1}s^{-1}$ [79,80]), Kelch-like ECH-associated protein 1 (KEAP1,

k= 140 $M^{-1}s^{-1}$ [81]), and a still unidentified target that regulates nuclear factor erythroid 2-related

factor 2 (NRF2) protein synthesis ($k \geq 1300$ $M^{-1}s^{-1}$ [81]). Of these, the oxidative inhibition of

GAPDH by H_2O_2 contributes for adaptation by redirecting the metabolic flux from glycolysis to

the oxidative part of the pentose phosphate pathway flux, thereby increasing NADPH

regeneration.[82] The reactions with KEAP1 and the NRF2 regulator induce multiple antioxidant

defenses whose promoters carry the antioxidant response element. On the other hand, various

protein tyrosine phosphatases that are rapidly inactivated upon cells' stimulation with mitogenic

factors have substantially lower rate constants for direct oxidation by H_2O_2.[83,84] Therefore,

even in this regime their inactivation requires other mechanisms (e.g. peroximonocarbonate-

mediated [85,86]).

The analysis of a simpler reaction-diffusion model of H_2O_2 signaling also supports the notion of

two distinct regimes,[13] with the transition here occurring through the hysteretic switch to the

high-hyperoxidation regime (as in Figure 2C, bottom). This hysteretic switch is unlikely to occur

in the cells examined in the present work, except under glutathione depletion (discussed below).

However, this does not change the remaining conclusions in ref. [13]. In particular, the estimated

times required to actuate redox targets by direct oxidation by H_2O_2 remain pertinent. It has been

argued [86] that by neglecting target reduction ref. [13] overestimated the signaling response

times. However, the analysis in ref. [86] refers to the time required for the target's redox state

to approach a new steady state, not to the time required to oxidize a given fraction of the

target's molecules as ref. [13] does. Target reduction cannot shorten the latter time. And it only

accelerates the response time by decreasing the fraction of the target that is oxidized at the new

steady state — that is, fewer target molecules need to be oxidized for approaching the new

steady state. The experimental observations that most targets are substantially oxidized upon

the stimuli indicate that reduction could accelerate the responses perhaps by up to 3-fold, but

not by orders of magnitude.

Is Response H an optimal trade-off between signaling and protection?

Why is Response H so prevalent among human cell lines? Could this be explained by any

functional advantages? A main distinguishing feature of Response H over Responses D and PD is

that at high v_{sup} Prx-SO$_2^-$ gradually accumulates, eventually replacing Prx-SS as the dominant Prx

form (SF21A). This behavior may have the following advantages.

First, it may better protect nascent proteins in proliferating cells against aggregation. The strong

increase in cytoplasmic H$_2$O$_2$ to μM concentrations accompanying the collapse of the PTTRS'

redox capacity should dramatically accelerate oxidative damage by Fenton-reaction-derived

hydroxyl radicals, lipid-peroxidation-derived electrophiles, etc. Recently synthesized proteins

that have not yet folded [87] and misfolded proteins [88] are particularly vulnerable to this

damage and to aggregation. Interestingly, hyperoxidation converts Prx to efficient holdases [89–

91], which protect damaged and misfolded proteins against aggregation [92]. The following

recent findings [93] support an even more central role of Prx-SO$_2^-$ in protection against

H$_2$O$_2$-induced protein aggregation/inclusions. In *S. cerevisiae*, hyperoxidized Tsa1 recruits Hsp70

chaperones to damaged or unfolded proteins, thereby promoting their folding or destruction.

Additionally, it recruits Hsp70/Hsp104 to H$_2$O$_2$-induced protein inclusions, promoting their

disaggregation. Something similar may happen in human cells. Accordingly, PrxII hyperoxidation

protected HeLa cells from H$_2$O$_2$-induced apoptosis, whereas hyperoxidation-resistant,

peroxidase-competent PrxII mutants were less protective [90]. Supporting a connection with cell

proliferation, PrxII hyperoxidation correlated with reversible H$_2$O$_2$-induced cell cycle arrest of

C10 mouse lung epithelial cells [94].

Second, the higher hyperoxidation decreases Prx-SS production and the ensuing load on the

Trx-S$^-$ pool (compare SF21A to SF21G), which has at least three advantages. (A) Leaving this pool

available for other protective processes such as the reduction of methionine sulfoxides.[95]

(B) Improving resistance to apoptosis by delaying ASK1 activation. (C) Decreasing the excretion

of pro-inflammatory glutathionylated PrxI-SS and PrxII-SS [96–98]. Excretion of these Prx forms

is proportional to the intracellular Prx-SS concentration [97], and it may signal the imminent
collapse of the cell's ability to sustain oxidative stress.

Also important, in cells lacking sufficient TrxR activity or Trx1 to avoid Responses D and PD the
Prx-S$^-$ pool collapses at lower v_{sup} than in cells with higher TrxR and Trx concentrations and
otherwise identical protein concentrations and activities. Those cells may thus be less tolerant
of H_2O_2 supply.

The discussion in the previous paragraphs raises the question of why shouldn't cells carry more
Trx1 and TrxR, as this might increase their resistance to H_2O_2. Perhaps the physiological H_2O_2
supply rates are rarely high enough for the advantages of increased resistance to compensate
the costs of increased protein expression. But there may be two other reasons. First, higher Trx1
and TrxR would cause bi-stability, such that transient increases in v_{sup} beyond v_{sup}^h in Expression
(18) could then trigger the positive feedback causing near-complete Prx hyperoxidation (Figure
2C, bottom). This positive feedback is intensified, making the switch more likely, if the GSH pool
is depleted. This is both because the decrease in H_2O_2 clearance due to hyperoxidation then has
a stronger impact on H_2O_2 concentrations and because Srx-catalyzed Prx-SO$_2^-$ reduction is itself
GSH-dependent. Recovery from the high-hyperoxidation state can start only after v_{sup} decreases
below v_{sup}^h and is very slow, limited by the low Srx activity. Therefore, cells would be left with a
diminished H_2O_2 clearance capacity for a long time, which would eventually trigger apoptosis.
Intriguingly, the Trx1 and TrxR concentrations in almost all cell lines in our sample are just low
enough to avoid this bi-stable behavior, which suggests that precisely controlling the extent of
hyperoxidation is critical for optimal antioxidant protection.

The second reason to avoid very high Trx1 and TrxR concentrations is that this would prevent
the attainment of high Prx-SS and/or Trx-SS concentrations, thereby disengaging redox relays
mediated by these species.

Altogether, the considerations above suggest that Response H embodies an optimal trade-off
between effective signal transduction and effective antioxidant protection. Together with the
prediction that both differentiated human cell types in our sample show Response D they also
raise the question of whether Response H is associated to cell proliferation and helps tumor cells
survive.

The hypothesized functional advantages of Response H are experimentally testable through
genetic or pharmacological manipulation of the PTTRS.

Strong correlations among protein concentrations over cell lines ensure the maintenance of Response H

The predicted similarity among the responses of the PTTRS is not simply due to a similar
protein composition of the cell lines. Instead, it is due to strong correlations among the
concentrations of several proteins over cell lines in such a manner as to keep the maximum rate
of Prx-SS production and the capacity for Prx-SS reduction approximately balanced. A positive
correlation between V_{Max}^{App} and k_{Srx} (Figure 5B), and negative correlations between V_{Max}^{App} and
Prx_T (Figure 5C) as well as between k_{Srx} and Prx_T (Figure 5D) are strong and statistically
significant. However, these three correlations are insufficient to explain the low variability of R
over these cell lines, as can be verified by replacing the scalings described in Figure 5B-D into
Expression (11). The additional dependencies among protein concentrations and activities that
are required to explain the invariance are likely embedded in various other correlations that
border the significance threshold. Thus, larger cell type samples and/or more precise
determinations may reveal additional aspects of the orchestration among the concentrations of
the PTTRS' proteins.

This orchestration should be at least in part the result of developmental processes associated to
cell differentiation and determining, for instance, that mean Srx concentrations are >100-fold
higher in some human cell types than in others. However, it likely also reflects faster
(hours-scale) regulatory mechanisms ensuring that each cell keeps a proper balance among the
PTTRS' proteins despite substantial fluctuations ("noise") of each protein's concentration.
Microscopy techniques allowing to follow the abundances of, say, fluorescently labeled Srx and
TrxR in single cells over time may clarify this point in the near future.

Given the strong involvement of the PTTRS in cancer and vascular processes [99–101], the
characterization of these regulatory mechanisms may have important implications for human
health. For instance, therapies that interfere with these mechanisms thereby unbalancing the
PTTRS may be more effective against cancer than therapies modulating the activity of any single
PTTRS protein.

The orchestration discussed above also implies that the functional consequences of up- or
down-regulation of any single PTTRS protein must be evaluated in light of likely compensatory
regulation of the other proteins. Quantifying multiple selected proteins will thus be important
for understanding potential disturbances of the PTTRS in disease. However, even these
multiplexed approaches may yield little insight in lack of a proper theoretical framework to help
interpret the results.

 ### Yeast show a distinctive response to H_2O_2 supply

The strong similarities in the response of the PTTRS across human cell lines prompted

to question if the PTTRS from phylogenetically distant eukaryotes shows similar responses. This

turns out not to be the case. *S. cerevisiae* was predicted to exhibit a distinctive response to H_2O_2

supply: its main Prx (Tsa1p) accumulates predominantly as Prx-SO$^-$ over a wide range of

intermediate v_{sup} (Figure 3L, Figure 4L). Furthermore, over all values of v_{sup}

Prx-SO$^-$ concentrations are $\approx$7-fold higher than those of Prx-SS, whereas in human cells they are

similar or lower than those of Prx-SS. That is mainly a consequence of the low $k_{Cond} = 0.58\,s^{-1}$

estimated for Tsa1 (SI3.4.3) from the data in ref. [102]. The low k_{Cond} and consequent high

Prx-SO$^-$ concentrations might make the yeast very sensitive to hyperoxidation. But remarkably

the same data indicate that Tsa1's sulfenate group is exceptionally stable, with a

$k_{Sulf} = 7.\,M^{-1}s^{-1}$ (estimated in SI3.4.3), as compared to $1.2 \times 10^4\,M^{-1}s^{-1}$ for human PrxII and

PrxIII [103] and $1.3 \times 10^3\,M^{-1}s^{-1}$ for PrxI (estimated in SI3.2.3). Consequently, the yeast is

predicted (Figure 4L) to be more resistant to Prx hyperoxidation than most human cell lines,

which is indeed experimentally observed [91].

How these "special" properties of Tsa1 relate to its recently characterized [93] roles in

proteostasis and aging is an interesting matter for future research.

Concluding remarks

The relationship between the PTTRS' responses to H_2O_2 and its proteins' concentrations

and properties is complex, but it can be expressed in terms of simple analytical approximations.

The concentrations-to-phenotype map presented here can guide the interpretation of protein

concentration or gene expression datasets as well as the design of new redox biology

experiments and therapies.

The analysis identifies the Prx-SS reduction capacity as a critical factor separating two distinct

H_2O_2 signaling regimes in all nucleated human cells. At H_2O_2 supply rates below Prx-SS reduction

capacity, H_2O_2 concentrations are very low throughout the cytoplasm and highly localized.

Signaling in this regime should thus be mediated by localized Prx-mediated redox relays. In turn,

at H_2O_2 supply rates above Prx-SS reduction capacity μM-range H_2O_2 concentrations are attained

uniformly throughout the cytoplasm. Signaling in this regime can occur through direct oxidation

of some targets by H_2O_2. But as cells are unlikely to tolerate such high cytoplasmic H_2O_2 for more

than a few tens of minutes this signaling mechanism is only effective for targets with a H_2O_2

reactivity $>10^2$ $M^{-1}s^{-1}$.

Nearly all human cell lines analyzed are predicted to show a similar response to H_2O_2 supply that

may optimize a trade-off between effective redox-relay signaling and effective antioxidant

protection. This response favors a gradual and moderate accumulation of hyperoxidized Prx

after the H_2O_2 supply exceeds the above-mentioned threshold. It hinges on the Prx-SS reduction

capacity exceeding the maximal steady state Prx-SS production rate by a small margin that is

insufficient to enable a hysteretic switch to a full-hyperoxidation state. This balance between

maximal Prx-SS production and reduction capacity is maintained through strong correlations

between the concentrations of TrxR, Srx, and Prx over cell lines.

The Tsa1-based PTTRS in *S. cerevisiae* is predicted to show a distinctive response to H_2O_2 supply,

largely by virtue of the exceptional stability of Tsa1-SO$^-$.

The phenotypic map can be gradually refined as the kinetic properties of further relevant redox

interactions are determined, allowing to better define the responses to high H_2O_2 supply rates,

the functional complementarity between PrxI and PrxII, and the interactions between the PTTRS

and the Grx/GSH/GSSG reductase system.

Acknowledgements

This work was funded by fellowship SFRH/BD/51576/2011 and grants UID/NEU/04539

COMPETE (POCI-01-0145-FEDER-007440), PEst-OE/QUI/UI0612/2013, and FCOMP-01-0124-

FEDER-020978 financed by FEDER through the "Programa Operacional Factores de

Competitividade, COMPETE" and by national funds through "FCT, Fundação para a Ciência e a

Tecnologia" (project PTDC/QUI-BIQ/119657/2010). We thank Dr. Michael Savageau and Dr.

Jason Lomnitz (University of California – Davis) for illuminating discussions, and Ms. Lisa Susi for

careful review of the manuscript.

Competing financial interests

1060 The authors declare no competing financial interests.

References

[1] F.M. Low, M.B. Hampton, A. V Peskin, C.C. Winterbourn, Peroxiredoxin 2 functions as a

noncatalytic scavenger of low-level hydrogen peroxide in the erythrocyte, Blood. 109

(2007) 2611–2617. doi:10.1182/blood-2006-09-048728.

[2]   R.M. Johnson, J. r Goyettejr G, Y. Ravindranath, Y.S. Ho, Hemoglobin autoxidation and
regulation of endogenous H2O2 levels in erythrocytes, Free Radical Biology and
Medicine. 39 (2005) 1407–1417. doi:10.1016/j.freeradbiomed.2005.07.002.

[3]   P.A. Karplus, A primer on peroxiredoxin biochemistry, Free Radical Biology and Medicine.
80 (2015) 183–190. doi:10.1016/j.freeradbiomed.2014.10.009.

[4]   M.H. Choi, I.K. Lee, G.W. Kim, B.U. Kim, Y.H. Han, D.Y. Yu, H.S. Park, K.Y. Kim, J.S. Lee, C.
Choi, Y.S. Bae, B.I. Lee, S.G. Rhee, S.W. Kang, Regulation of PDGF signalling and vascular
remodelling by peroxiredoxin II, Nature. 435 (2005) 347–353. doi:10.1038/nature03587.

[5]   H.A. Woo, S.H. Yim, D.H. Shin, D. Kang, D.Y. Yu, S.G. Rhee, Inactivation of Peroxiredoxin I
by Phosphorylation Allows Localized H2O2 Accumulation for Cell Signaling, Cell. 140
(2010) 517–528. doi:10.1016/j.cell.2010.01.009.

[6]   D.Y. Jin, H.Z. Chae, S.G. Rhee, K.T. Jeang, Regulatory Role for a Novel Human Thioredoxin
Peroxidase in NF-κB Activation, The Journal of Biological Chemistry. 272 (1997) 30952–
30961. doi:10.1074/jbc.272.49.30952.

[7]   J. Cao, J. Schulte, A. Knight, N.R. Leslie, A. Zagozdzon, R. Bronson, Y. Manevich, C. Beeson,
C.A. Neumann, Prdx1 inhibits tumorigenesis via regulating PTEN/AKT activity, The EMBO
Journal. 28 (2009) 1505–1517. doi:10.1038/emboj.2009.101.

[8]   S.Y. Kim, T.J. Kim, K.Y. Lee, A novel function of peroxiredoxin 1 (Prx-1) in apoptosis signal-
regulating kinase 1 (ASK1)-mediated signaling pathway, FEBS Letters. 582 (2008) 1913–
1918. doi:10.1016/j.febslet.2008.05.015.

[9]   R.M. Jarvis, S.M. Hughes, E.C. Ledgerwood, Peroxiredoxin 1 functions as a signal
peroxidase to receive, transduce, and transmit peroxide signals in mammalian cells, Free
Radical Biology and Medicine. 53 (2012) 1522–1530.
doi:10.1016/j.freeradbiomed.2012.08.001.

[10]   C.C. Winterbourn, Reconciling the chemistry and biology of reactive oxygen species,
Nature Chemical Biology. 4 (2008) 278–286. doi:10.1038/nchembio.85.

[11]   C.A. Neumann, J. Cao, Y. Manevich, Peroxiredoxin 1 and its role in cell signaling, Cell
Cycle. 8 (2009) 4072–4078. doi:10.4161/cc.8.24.10242.

[12]   Z.A. Wood, Peroxiredoxin Evolution and the Regulation of Hydrogen Peroxide Signaling,
Science. 300 (2003) 650–653. doi:10.1126/science.1080405.

[13] R.D.M. Travasso, F. Sampaio dos Aidos, A. Bayani, P. Abranches, A. Salvador, Localized
Redox Relays as a Privileged Mode of Cytoplasmic Hydrogen Peroxide Signaling, Redox
Biology. 12 (2017) 233–245. doi:10.1016/j.redox.2017.01.003.

[14] H.A. Woo, H.Z. Chae, S.C. Hwang, K.S. Yang, S.W. Kang, K. Kim, S.G. Rhee, Reversing the
Inactivation of Peroxiredoxins Caused by Cysteine Sulfinic Acid Formation, Science. 300
(2003) 653–656. doi:10.1126/science.1080273.

[15] T. Geiger, A. Wehner, C. Schaab, J. Cox, M. Mann, Comparative proteomic analysis of
eleven common cell lines reveals ubiquitous but varying expression of most proteins,
Molecular   &   Cellular   Proteomics: MCP.   11   (2012)   M111.014050.
doi:10.1074/mcp.M111.014050.

[16] J.H. Seo, J.C. Lim, D.Y. Lee, K.S. Kim, G. Piszczek, H.W. Nam, Y.S. Kim, T. Ahn, C.H. Yun, K.
Kim, P.B. Chock, H.Z. Chae, Novel Protective Mechanism against Irreversible
Hyperoxidation of Peroxiredoxin N-alpha-terminal acetylation of human peroxiredoxin,
Journal Of Biological Chemistry. 284 (2009) 13455–13465. doi:10.1074/jbc.M900641200.

[17] A. Mitsumoto, Y. Takanezawa, K. Okawa, A. Iwamatsu, Y. Nakagawa, Variants of
peroxiredoxins expression in response to hydroperoxide stress, Free Radical Biology and
Medicine. 30 (2001) 625–635. doi:10.1016/S0891-5849(00)00503-7.

[18] M.C. Sobotta, A.G. Barata, U. Schmidt, S. Mueller, G. Millonig, T.P. Dick, Exposing cells to
H2O2: A quantitative comparison between continuous low-dose and one-time high-dose
treatments,   Free   Radical   Biology   and   Medicine.   60   (2013)   325–335.
doi:10.1016/j.freeradbiomed.2013.02.017.

[19] L.E. Tomalin, A.M. Day, Z.E. Underwood, G.R. Smith, P. Dalle Pezze, C. Rallis, W. Patel,
B.C. Dickinson, J. Bähler, T.F. Brewer, C.J.-L. Chang, D.P. Shanley, E.A. Veal, Increasing
extracellular H2O2 produces a bi-phasic response in intracellular H2O2, with
peroxiredoxin hyperoxidation only triggered once the cellular H2O2-buffering capacity is
overwhelmed,   Free   Radical   Biology   and   Medicine.   95   (2016)   333–348.
doi:10.1016/j.freeradbiomed.2016.02.035.

[20] C.S. Pillay, J.H. Hofmeyr, L.N. Mashamaite, J.M. Rohwer, From Top-Down to Bottom-Up:
Computational Modeling Approaches for Cellular Redoxin Networks, Antioxidants &
Redox Signaling. 18 (2013) 2075–2086. doi:10.1089/ars.2012.4771.

[21] R. Benfeitas, G. Selvaggio, F. Antunes, P.M.B.M. Coelho, A. Salvador, Hydrogen peroxide

metabolism and sensing in human erythrocytes: A validated kinetic model and

reappraisal of the role of peroxiredoxin II, Free Radical Biology and Medicine. 74 (2014)

35–49. doi:10.1016/j.freeradbiomed.2014.06.007.

[22] J.B. Lim, T.F. Langford, B.K. Huang, W.M. Deen, H.D. Sikes, A reaction-diffusion model of

cytosolic hydrogen peroxide, Free Radical Biology and Medicine. 90 (2016) 85–90.

doi:10.1016/j.freeradbiomed.2015.11.005.

[23] C.S. Pillay, B.D. Eagling, S.R. Driscoll, J.M. Rohwer, Quantitative measures for redox

signaling, Free Radical Biology and Medicine. 96 (2016) 290–303.

doi:10.1016/j.freeradbiomed.2016.04.199.

[24] A. Salvador, F. Antunes, R.E. Pinto, Kinetic modelling of in vitro lipid peroxidation

experiments--'low level' validation of a model of in vivo lipid peroxidation, Free Radical

Research. 23 (1995) 151–172. doi:0.3109/10715769509064029.

[25] F. Antunes, A. Salvador, H.S. Marinho, R. Alves, R.E. Pinto, A.S. Marinho, Lipid

peroxidation in mitochondrial inner membranes. I. An integrative kinetic model, Free

Radical Biology and Medicine. 21 (1996) 21(7):917-943. doi:10.1016/S0891-

5849(96)00185-2.

[26] F. Antunes, A. Salvador, R.E. Pinto, PHGPx and phospholipase A2/GPx: comparative

importance on the reduction of hydroperoxides in rat liver mitochondria, Free Radical

Biology and Medicine. 19 (1995) 669–677. doi:10.1016/0891-5849(95)00040-5.

[27] A. Salvador, J. Sousa, R.E. Pinto, Hydroperoxyl, superoxide and pH gradients in the

mitochondrial matrix: a theoretical assessment, Free Radical Biology and Medicine. 31

(2001) 1208–1215.

[28] G.R. Buettner, Moving free radical and redox biology ahead in the next decade(s), Free

Radical Biology and Medicine. 78 (2015) 236–238.

doi:10.1016/j.freeradbiomed.2014.10.578.

[29] P.M.B.M. Coelho, A. Salvador, M.A. Savageau, Quantifying Global Tolerance of

Biochemical Systems: Design Implications for Moiety-Transfer Cycles, PLoS

Computational Biology. 5 (2009) e1000319. doi:10.1371/journal.pcbi.1000319.

[30] M.A. Savageau, P.M.B.M. Coelho, R.A. Fasani, D.A. Tolla, A. Salvador, Phenotypes and

tolerances in the design space of biochemical systems, Proceedings of the National

Academy of Sciences. 106 (2009) 6435–6440. doi:10.1073/pnas.0809869106.

[31] P.M.B.M. Coelho, A. Salvador, M.A. Savageau, Relating Mutant Genotype to Phenotype
via Quantitative Behavior of the NADPH Redox Cycle in Human Erythrocytes, PLoS ONE.
5 (2010) e13031. doi:10.1371/journal.pone.0013031.

[32] A. Salvador, M.A. Savageau, Quantitative evolutionary design of glucose 6-phosphate
dehydrogenase expression in human erythrocytes, Proceedings of the National Academy
of Sciences. 100 (2003) 14463–14468. doi:10.1073/pnas.2335687100.

[33] A. Salvador, M.A. Savageau, Evolution of enzymes in a series is driven by dissimilar
functional demands, Proceedings of the National Academy of Sciences. 103 (2006) 2226–
2231. doi:10.1073/pnas.0510776103.

[34] A. Kuehne, H. Emmert, J. Soehle, M. Winnefeld, F. Fischer, H. Wenck, S. Gallinat, L.
Terstegen, R. Lucius, J. Hildebrand, N. Zamboni, Acute Activation of Oxidative Pentose
Phosphate Pathway as First-Line Response to Oxidative Stress in Human Skin Cells.,
Molecular Cell. 59 (2015) 359–371. doi:10.1016/j.molcel.2015.06.017.

[35] R.A. Poynton, A. V Peskin, A.C. Haynes, W.T. Lowther, M.B. Hampton, C.C. Winterbourn,
Kinetic analysis of structural influences on the susceptibility of peroxiredoxins 2 and 3 to
hyperoxidation, Biochemical Journal. 473 (2016) 411–421. doi:10.1042/BJ20150572.

[36] T.S. Chang, W. Jeong, H.A. Woo, S.M. Lee, S. Park, S.G. Rhee, Characterization of
Mammalian Sulfiredoxin and Its Reactivation of Hyperoxidized Peroxiredoxin through
Reduction of Cysteine Sulfinic Acid in the Active Site to Cysteine, Journal of Biological
Chemistry. 279 (2004) 50994–51001. doi:10.1074/jbc.M409482200.

[37] X. Roussel, S. Boukhenouna, S. Rahuel-Clermont, G. Branlant, The rate-limiting step of
sulfiredoxin is associated with the transfer of the γ-phosphate of ATP to the sulfinic acid
of overoxidized typical 2-Cys peroxiredoxins, FEBS Letters. 585 (2011) 574–578.
doi:10.1016/j.febslet.2011.01.012.

[38] E.S. Arnér, A. Holmgren, Physiological functions of thioredoxin and thioredoxin
reductase, European Journal of Biochemistry. 267 (2000) 6102–6109.
doi:10.1046/j.1432-1327.2000.01701.x.

[39] S. Urig, J. Lieske, K. Fritz-Wolf, A. Irmler, K. Becker, Truncated mutants of human
thioredoxin reductase 1 do not exhibit glutathione reductase activity, FEBS Letters. 580
(2006) 3595–3600. doi:10.1016/j.febslet.2006.05.038.

[40] M. Cebula, E.E. Schmidt, E.S. Arnér, TrxR1 as a Potent Regulator of the Nrf2-Keap1

Response System, Antioxidants & Redox Signaling. 23 (2015) 823–853.
doi:10.1089/ars.2015.6378.

[41]   J.W. Park, J.J. Mieyal, S.G. Rhee, P.B. Chock, Deglutathionylation of 2-Cys Peroxiredoxin
Is Specifically Catalyzed by Sulfiredoxin, The Journal of Biological Chemistry. 284 (2009)
23364–23374. doi:10.1074/jbc.M109.021394.

[42]   A. V. Peskin, P.E. Pace, J.B. Behring, L.N. Paton, M. Soethoudt, M.M. Bachschmid, C.C.
Winterbourn, Glutathionylation of the Active Site Cysteines of Peroxiredoxin 2 and
Recycling by Glutaredoxin, The Journal of Biological Chemistry. 291 (2016) 3053–3062.
doi:10.1074/jbc.M115.692798.

[43]   Y. Du, H. Zhang, J. Lu, A. Holmgren, Glutathione and glutaredoxin act as a backup of
human thioredoxin reductase 1 to reduce thioredoxin 1 preventing cell death by
aurothioglucose, Journal of Biological Chemistry. 287 (2012) 38210–38219.
doi:10.1074/jbc.M112.392225.

[44]   W.H. Koppenol, Thermodynamics of reactions involving nitrogen-oxygen compounds,
Methods in Enzymology. 268 (1996) 7–12. doi:10.1016/S0076-6879(96)68005-7.

[45]   J.B. Lim, B.K. Huang, W.M. Deen, H.D. Sikes, Analysis of the lifetime and spatial
localization of Hydrogen peroxide generated in the cytosol using a reduced kinetic
model., Free Radical Biology and Medicine. 89 (2015) 47–53.
doi:10.1016/j.freeradbiomed.2015.07.009.

[46]   A. Salvador, Synergism analysis of biochemical systems. I. Conceptual framework,
Mathematical Biosciences. 163 (2000) 105–129. doi:10.1016/S0025-5564(99)00056-5.

[47]   A. Salvador, Synergism analysis of biochemical systems. II. Tensor formulation and
treatment of stoichiometric constraints, Mathematical Biosciences. 163 (2000) 131–158.
doi:10.1016/S0025-5564(99)00057-7.

[48]   I. Wolfram Research, Mathematica, Version 7., Wolfram Research, Inc., 2008.

[49]   X. Roussel, A. Kriznik, C. Richard, S. Rahuel-Clermont, G. Branlant, Catalytic Mechanism
of Sulfiredoxin from Saccharomyces cerevisiae Passes through an Oxidized Disulfide
Sulfiredoxin Intermediate That Is Reduced by Thioredoxin, Journal of Biological
Chemistry. 284 (2009) 33048–33055. doi:10.1074/jbc.M109.035352.

[50]   S.G. Rhee, W. Jeong, T.S. Chang, H.A. Woo, Sulfiredoxin, the cysteine sulfinic acid
reductase specific to 2-Cys peroxiredoxin: its discovery, mechanism of action, and

biological significance, Kidney International. 72 (2007) S3–S8. doi:10.1038/sj.ki.5002380.

[51] X. Roussel, G. Béchade, A. Kriznik, A. Van Dorsselaer, S. Sanglier-Cianferani, G. Branlant,
S. Rahuel-Clermont, Evidence for the Formation of a Covalent Thiosulfinate Intermediate
with Peroxiredoxin in the Catalytic Mechanism of Sulfiredoxin, Journal of Biological
Chemistry. 283 (2008) 22371–22382. doi:10.1074/jbc.M800493200.

[52] J.J. Tyson, K.C. Chen, B. Novak, Sniffers, buzzers, toggles and blinkers: dynamics of
regulatory and signaling pathways in the cell, Current Opinion In Cell Biology. 15 (2003)
221–231. doi:10.1016/s0955-0674(03)00017-6.

[53] T. Geiger, A. Wehner, C. Schaab, J. Cox, M. Mann, Comparative Proteomic Analysis of
Eleven Common Cell Lines Reveals Ubiquitous but Varying Expression of Most Proteins,
Molecular & Cellular Proteomics. 11 (2012) M111.014050-M111.014050.
doi:10.1074/mcp.M111.014050.

[54] J.R. Wiśniewski, A. Vildhede, A. Norén, P. Artursson, In-depth quantitative analysis and
comparison of the human hepatocyte and hepatoma cell line HepG2 proteomes, Journal
of Proteomics. 136 (2016) 234–247. doi:10.1016/j.jprot.2016.01.016.

[55] R.M. Johnson, Y.-S. Ho, D.-Y. Yu, F.A. Kuypers, Y. Ravindranath, G.W. Goyette, The effects
of disruption of genes for peroxiredoxin-2, glutathione peroxidase-1, and catalase on
erythrocyte oxidative metabolism, Free Radical Biology and Medicine. 48 (2010) 519–
525. doi:10.1016/j.freeradbiomed.2009.11.021.

[56] H.S. Jacob, S.H. Ingbar, J.H. Jandl, Oxidative hemolysis and erythrocyte metabolism in
hereditary acatalasia, Journal of Clinical Investigation. 44 (1965) 1187–1199.
doi:10.1172/JCI105225.

[57] C.S. Cho, G.J. Kato, S.H. Yang, S.W. Bae, J.S. Lee, M.T. Gladwin, S.G. Rhee, Hydroxyurea-
Induced Expression of Glutathione Peroxidase 1 in Red Blood Cells of Individuals with
Sickle Cell Anemia, Antioxidants and Redox Signaling. 13 (2010) 1–11.
doi:10.1089/ars.2009.2978.

[58] G.P. Bienert, A.L.B. Møller, K.A. Kristiansen, A. Schulz, I.M. Møller, J.K. Schjoerring, T.P.
Jahn, Specific aquaporins facilitate the diffusion of hydrogen peroxide across
membranes., The Journal of Biological Chemistry. 282 (2007) 1183–92.
doi:10.1074/jbc.M603761200.

[59] M. Jelcic, B. Enyedi, J.B. Xavier, P. Niethammer, Image-Based Measurement of H_2O_2

Reaction-Diffusion in Wounded Zebrafish Larvae, Biophysical Journal. 112 (2017) 2011–
2018. doi:10.1016/j.bpj.2017.03.021.

[60] M. Chevallet, E. Wagner, S. Luche, A. van Dorsselaer, E. Leize-Wagner, T. Rabilloud,
Regeneration of peroxiredoxins during recovery after oxidative stress - Only some
overoxidized peroxiredoxins can be reduced during recovery after oxidative stress,
Journal Of Biological Chemistry. 278 (2003) 37146–37153. doi:10.1074/jbc.M305161200.

[61] S.J. Montano, J. Lu, T.N. Gustafsson, A. Holmgren, Activity assays of mammalian
thioredoxin and thioredoxin reductase: Fluorescent disulfide substrates, mechanisms,
and use with tissue samples, Analytical Biochemistry. 449 (2014) 139–146.
doi:10.1016/J.AB.2013.12.025.

[62] M. Kontou, C. Adelfalk, M. Hirsch-Kauffmann, M. Schweiger, Suboptimal Action of NF-κB
in Fanconi Anemia Cells Results from Low Levels of Thioredoxin, Biological Chemistry.
384 (2003). doi:10.1515/BC.2003.166.

[63] S.J. Wei, A. Botero, K. Hirota, C.M. Bradbury, S. Markovina, A. Laszlo, D.R. Spitz, P.C.
Goswami, J. Yodoi, D. Gius, Thioredoxin nuclear translocation and interaction with redox
factor-1 activates the activator protein-1 transcription factor in response to ionizing
radiation., Cancer Research. 60 (2000) 6688–95.

[64] P. Schroeder, R. Popp, B. Wiegand, J. Altschmied, J. Haendeler, Nuclear redox-signaling is
essential for apoptosis inhibition in endothelial cells--important role for nuclear
thioredoxin-1., Arteriosclerosis, Thrombosis, and Vascular Biology. 27 (2007) 2325–31.
doi:10.1161/ATVBAHA.107.149419.

[65] S. Lee, S.M. Kim, R.T. Lee, Thioredoxin and Thioredoxin Target Proteins: From Molecular
Mechanisms to Functional Significance, Antioxidants and Redox Signaling. 18 (2012)
1165–1207. doi:10.1089/ars.2011.4322.

[66] B.A. Wagner, J.R. Witmer, T.J. van 't Erve, G.R. Buettner, An Assay for the Rate of Removal
of Extracellular Hydrogen Peroxide by Cells., Redox Biology. 1 (2013) 210–217.
doi:10.1016/j.redox.2013.01.011.

[67] C.M. Doskey, V. Buranasudja, B.A. Wagner, J.G. Wilkes, J. Du, J.J. Cullen, G.R. Buettner,
Tumor cells have decreased ability to metabolize H2O2: Implications for pharmacological
ascorbate in cancer therapy, Redox Biology. 10 (2016) 274–284.
doi:10.1016/j.redox.2016.10.010.

[68] B.K. Huang, H.D. Sikes, Quantifying intracellular hydrogen peroxide perturbations in terms of concentration, Redox Biology. 2 (2014) 955–962. doi:10.1016/j.redox.2014.08.001.

[69] F. Antunes, E. Cadenas, Cellular titration of apoptosis with steady state concentrations of H2O2: submicromolar levels of H2O2 induce apoptosis through Fenton chemistry independent of the cellular thiol state, Free Radical Biology and Medicine. 30 (2001) 1008–1018. doi:10.1016/S0891-5849(01)00493-2.

[70] F. Antunes, E. Cadenas, Estimation of H2O2 gradients across biomembranes, FEBS Letters. 475 (2000) 121–126. doi:10.1016/S0003-9861(02)00049-8.

[71] M. Saitoh, H. Nishitoh, M. Fujii, K. Takeda, K. Tobiume, Y. Sawada, M. Kawabata, K. Miyazono, H. Ichijo, Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1, The EMBO Journal. 17 (1998) 2596–2606. doi:10.1093/emboj/17.9.2596.

[72] D. Kosek, S. Kylarova, K. Psenakova, L. Rezabkova, P. Herman, J. Vecer, V. Obsilova, T. Obsil, Biophysical and structural characterization of the thioredoxin-binding domain of protein kinase ASK1 and its interaction with reduced thioredoxin., The Journal of Biological Chemistry. 289 (2014) 24463–74. doi:10.1074/jbc.M114.583807.

[73] S. Kylarova, D. Kosek, O. Petrvalska, K. Psenakova, P. Man, J. Vecer, P. Herman, V. Obsilova, T. Obsil, Cysteine residues mediate high-affinity binding of thioredoxin to ASK1, The FEBS Journal. 283 (2016) 3821–3838. doi:10.1111/febs.13893.

[74] T. Obsil, V. Obsilova, Structural aspects of protein kinase ASK1 regulation, Advances in Biological Regulation. (2017) 0–1. doi:10.1016/j.jbior.2017.10.002.

[75] T. Nishida, K. Hattori, K. Watanabe, The regulatory and signaling mechanisms of the ASK family, Advances in Biological Regulation. (2017) 1–21. doi:10.1016/j.jbior.2017.05.004.

[76] L. Flohé, Changing paradigms in thiology from antioxidant defense toward redox regulation, in: Methods Enzymol, 2010: pp. 1–39. doi:10.1016/S0076-6879(10)73001-9.

[77] H.J. Forman, M. Maiorino, F. Ursini, Signaling Functions of Reactive Oxygen Species, Biochemistry. 49 (2010) 835–842. doi:10.1021/bi9020378.

[78] M.C. Sobotta, W. Liou, S. Stöcker, D. Talwar, M. Oehler, T. Ruppert, A.N. Scharf, T.P. Dick, Peroxiredoxin-2 and STAT3 form a redox relay for H2O2 signaling, Nature Chemical Biology. 11 (2014) 64–70. doi:10.1038/nchembio.1695.

[79] C. Little, P.J. O'brien, Mechanism of Peroxide-Inactivation of the Sulphydryl Enzyme
Glyceraldehyde-3-Phosphate Dehydrogenase, European Journal of Biochemistry. 10
(1969) 533–538. doi:10.1111/j.1432-1033.1969.tb00721.x.

[80] C.C. Winterbourn, M.B. Hampton, Thiol chemistry and specificity in redox signaling, Free
Radical   Biology   and   Medicine.   45   (2008)   549–561.
doi:10.1016/j.freeradbiomed.2008.05.004.

[81] H.S. Marinho, C. Real, L. Cyrne, H. Soares, F. Antunes, Hydrogen peroxide sensing,
signaling and regulation of transcription factors, Redox Biology. 2 (2014) 535–562.
doi:10.1016/j.redox.2014.02.006.

[82] D. Peralta, A.K. Bronowska, B. Morgan, É. Dóka, K. Van Laer, P. Nagy, F. Gräter, T.P. Dick,
A proton relay enhances H2O2 sensitivity of GAPDH to facilitate metabolic adaptation.,
Nature Chemical Biology. 11 (2015) 156–63. doi:10.1038/nchembio.1720.

[83] U. Schwertassek, A. Haque, N. Krishnan, R. Greiner, L. Weingarten, T.P. Dick, N.K. Tonks,
Reactivation of oxidized PTP1B and PTEN by thioredoxin 1, FEBS Journal. 281 (2014)
3545–3558. doi:10.1111/febs.12898.

[84] J.M. Denu, K.G. Tanner, Specific and Reversible Inactivation of Protein Tyrosine
Phosphatases by Hydrogen Peroxide: Evidence for a Sulfenic Acid Intermediate and
Implications   for   Redox   Regulation,   Biochemistry.   37   (1998)   5633–5642.
doi:10.1021/bi973035t.

[85] H. Zhou, H. Singh, Z.D. Parsons, S.M. Lewis, S. Bhattacharya, D.R. Seiner, J.N. LaButti, T.J.
Reilly, J.J. Tanner, K.S. Gates, The Biological Buffer Bicarbonate/CO2 Potentiates H2O2 -
Mediated Inactivation of Protein Tyrosine Phosphatases, Journal of the American
Chemical Society. 133 (2011) 15803–15805. doi:10.1021/ja2077137.

[86] F. Antunes, P.M. Brito, Quantitative biology of hydrogen peroxide signaling, Redox
Biology. (2017). doi:10.1016/j.redox.2017.04.039.

[87] B. Medicherla, A.L. Goldberg, Heat shock and oxygen radicals stimulate ubiquitin-
dependent degradation mainly of newly synthesized proteins., The Journal of Cell
Biology. 182 (2008) 663–73. doi:10.1083/jcb.200803022.

[88] S. Dukan, A. Farewell, M. Ballesteros, F. Taddei, M. Radman, T. Nystrom, Protein
oxidation in response to increased transcriptional or translational errors, Proceedings Of
The National Academy Of Sciences Of The United States Of America. 97 (2000) 5746–

5749. doi:10.1073/pnas.100422497.

[89]   H.H. Jang, K.O. Lee, Y.H. Chi, B.G. Jung, S.K. Park, J.H. Park, J.R. Lee, S.S. Lee, J.C. Moon, J.W. Yun, Y.O. Choi, W.Y. Kim, J.S. Kang, G.-W. Cheong, D.-J. Yun, S.G. Rhee, M.J. Cho, S.Y. Lee, Two enzymes in one. Two yeast peroxiredoxins display oxidative stress-dependent switching from a peroxidase to a molecular chaperone function., Cell. 117 (2004) 625–35. doi:10.1016/j.cell.2004.05.002.

[90]   J.C. Moon, Y.-S. Hah, W.Y. Kim, B.G. Jung, H.H. Jang, J.R. Lee, S.Y. Kim, Y.M. Lee, M.G. Jeon, C.W. Kim, M.J. Cho, S.Y. Lee, Oxidative stress-dependent structural and functional switching of a human 2-Cys peroxiredoxin isotype II that enhances HeLa cell resistance to H2O2-induced cell death, Journal of Biological Chemistry. 280 (2005) 28775–28784. doi:10.1074/jbc.M505362200.

[91]   J.C. Lim, H.-I. Choi, Y.S. Park, H.W. Nam, H.A. Woo, K.-S. Kwon, Y.S. Kim, S.G. Rhee, K. Kim, H.Z. Chae, Irreversible oxidation of the active-site cysteine of peroxiredoxin to cysteine sulfonic acid for enhanced molecular chaperone activity., The Journal of Biological Chemistry. 283 (2008) 28873–80. doi:10.1074/jbc.M804087200.

[92]   J.H. Hoffmann, K. Linke, P.C.F. Graf, H. Lilie, U. Jakob, Identification of a redox-regulated chaperone network, The EMBO Journal. 23 (2003) 160 LP-168. doi:10.1038/sj.emboj.7600016.

[93]   S. Hanzén, K. Vielfort, J. Yang, F. Roger, V. Andersson, S. Zamarbide-Forés, R. Andersson, L. Malm, G. Palais, B. Biteau, B. Liu, M.B.B. Toledano, M. Molin, T. Nyström, Lifespan Control by Redox-Dependent Recruitment of Chaperones to Misfolded Proteins, Cell. 166 (2016) 140–151. doi:10.1016/j.cell.2016.05.006.

[94]   T.J. Phalen, K. Weirather, P.B. Deming, V. Anathy, A.K. Howe, A. van der Vliet, T.J. Jönsson, L.B. Poole, N.H. Heintz, Oxidation state governs structural transitions in peroxiredoxin II that correlate with cell cycle arrest and recovery, Journal of Cell Biology. 175 (2006) 779–789. doi:10.1083/jcb.200606005.

[95]   A.M. Day, J.D. Brown, S.R. Taylor, J.D. Rand, B.A. Morgan, E.A. Veal, Inactivation of a Peroxiredoxin by Hydrogen Peroxide Is Critical for Thioredoxin-Mediated Repair of Oxidized Proteins and Cell Survival, Molecular Cell. 45 (2012) 398–408. doi:10.1016/j.molcel.2011.11.027.

[96]   S. Salzano, P. Checconi, E.-M. Hanschmann, C.H. Lillig, L.D. Bowler, P. Chan, D. Vaudry, M.

Mengozzi, L. Coppo, S. Sacre, K.R. Atkuri, B. Sahaf, L.A. Herzenberg, L.A. Herzenberg, L.

Mullen, P. Ghezzi, Linkage of inflammation and oxidative stress via release of

glutathionylated peroxiredoxin-2, which acts as a danger signal., Proceedings of the

National Academy of Sciences of the United States of America. 111 (2014) 12157–62.

doi:10.1073/pnas.1401712111.

[97] L. Mullen, E.-M. Hanschmann, C.H. Lillig, L.A. Herzenberg, P. Ghezzi, Cysteine Oxidation

Targets Peroxiredoxins 1 and 2 for Exosomal Release through a Novel Mechanism of

Redox-Dependent Secretion., Molecular Medicine (Cambridge, Mass.). 21 (2015) 98–

108. doi:10.2119/molmed.2015.00033.

[98] P. Checconi, S. Salzano, L. Bowler, L. Mullen, M. Mengozzi, E.-M. Hanschmann, C.H. Lillig,

R. Sgarbanti, S. Panella, L. Nencioni, A.T. Palamara, P. Ghezzi, Redox Proteomics of the

Inflammatory Secretome Identifies a Common Set of Redoxins and Other

Glutathionylated Proteins Released in Inflammation, Influenza Virus Infection and

Oxidative Stress, PLOS ONE. 10 (2015) e0127086. doi:10.1371/journal.pone.0127086.

[99] M. Mishra, H. Jiang, L. Wu, H.A. Chawsheen, Q. Wei, The sulfiredoxin-peroxiredoxin (Srx-

Prx) axis in cell signal transduction and cancer development., Cancer Letters. 366 (2015)

150–9. doi:10.1016/j.canlet.2015.07.002.

[100] A. Mukherjee, S.G. Martin, The thioredoxin system: a key target in tumour and

endothelial cells, The British Journal of Radiology. 81 (2008) S57–S68.

doi:10.1259/bjr/34180435.

[101] R. Bretón-Romero, S. Lamas, Hydrogen peroxide signaling in vascular endothelial cells.,

Redox Biology. 2 (2014) 529–534. doi:10.1016/j.redox.2014.02.005.

[102] C.A. Tairum Jr., M.A. de Oliveira, B.B. Horta, F.J. Zara, L.E.S. Netto, Disulfide Biochemistry

in 2-Cys Peroxiredoxin: Requirement of Glu50 and Arg146 for the Reduction of Yeast Tsa1

by Thioredoxin, Journal of Molecular Biology. 424 (2012) 28–41.

doi:10.1016/j.jmb.2012.09.008.

[103] A. V Peskin, N. Dickerhof, R.A. Poynton, L.N. Paton, P.E. Pace, M.B. Hampton, C.C.

Winterbourn, Hyperoxidation of peroxiredoxins 2 and 3: Rate constants for the reactions

of the sulfenic acid of the peroxidatic cysteine, The Journal of Biological Chemistry. 288

(2013) 14170–14177. doi:10.1074/jbc.M113.460881.

1404