1 **Title:** *AtRsgA* from *Arabidopsis thaliana* controls maturation of the small subunit of
2 the chloroplast ribosome
3

4 **Authors:** Marcin Janowski[1], Reimo Zoschke[1], Lars Scharff[2], Silvia Martinez Jaime[1],
5 Camilla Ferrari[1], Sebastian Proost[1], Nooshin Omranian[1], Magdalena Musialak-
6 Lange[1], Zoran Nikoloski[1,3], Alexander Graf[1], Mark A. Schöttler[1], Arun
7 Sampathkumar[1], Neha Vaid[1]*, Marek Mutwil[1,4]*
8
[1]Max Planck Institute of Molecular Plant Physiology, Am Muehlenberg 1, 14476
Potsdam, Germany
[2]Copenhagen Plant Science Centre, Department of Plant and Environmental
Sciences, University of Copenhagen, Thorvaldsensvej 40, DK-1871 Frederiksberg C,
Denmark
[3]Bioinformatics Group, Institute of Biochemistry and Biology, University of Potsdam,
Karl- Liebknecht-Str. 24-25, 14476 Potsdam-Golm, Germany
[4]School of Biological Sciences, Nanyang Technological University, 60 Nanyang
Drive, Singapore 637551, Singapore

*Corresponding authors:

Marek Mutwil

Max Planck Institute of Molecular Plant Physiology,

Am Muehlenberg 1,

14476 Potsdam,

Germany

Email: mutwil@mpimp-golm.mpg.de

Neha Vaid

Max Planck Institute of Molecular Plant Physiology,

Am Muehlenberg 1,

14476 Potsdam,

Germany

Email: vaid@mpimp-golm.mpg.de

Summary: Plastid ribosomes are very similar in structure and function to ribosomes of their bacterial ancestors. Since ribosome biogenesis is not thermodynamically favourable at biological conditions, it requires activity of many assembly factors. Here, we have characterized a homolog of bacterial *rsgA* in *Arabidopsis thaliana* and show that it can complement the bacterial homolog. Functional characterization of a strong mutant in Arabidopsis revealed that the protein is essential for plant viability, while a weak mutant produced dwarf, chlorotic plants that incorporated immature pre-16S ribosomal RNA into translating ribosomes. Physiological analysis of the mutant plants revealed smaller, but more numerous chloroplasts in the mesophyll cells, reduction of chlorophyll *a* and *b*, depletion of proplastids from the rib meristem and decreased photosynthetic electron transport rate and efficiency. Comparative RNA-sequencing and proteomic analysis of the weak mutant and wild-type plants revealed that various biotic stress-related, transcriptional regulation and post-transcriptional modification pathways were repressed in the mutant. Intriguingly, while nuclear- and chloroplast-encoded photosynthesis-related proteins were less abundant in the mutant, the corresponding transcripts were upregulated, suggesting an elaborate compensatory mechanism, potentially via differentially active retrograde signalling pathways. To conclude, this study reveals a new chloroplast ribosome assembly factor and outlines the transcriptomic and proteomic responses of the compensatory mechanism activated during decreased chloroplast function.

Significance statement: AtRsgA is an assembly factor necessary for maturation of the small subunit of the chloroplast ribosome. Depletion of AtRsgA leads to dwarfed, chlorotic plants and smaller, but more numerous chloroplasts. Large-scale transcriptomic and proteomic analysis revealed that chloroplast-encoded and -targeted proteins were less abundant, while the corresponding transcripts were upregulated in the mutant. We analyse the transcriptional responses of several retrograde signalling pathways to suggest a mechanism underlying this compensatory response.

Introduction

Plastids have retained many prokaryotic features from its bacterial ancestor, such as the presence of operons and polycistronic RNAs for most of the chloroplast-encoded genes. These genes are transcribed by a bacteria-type, chloroplast-encoded RNA polymerase and nucleus-encoded sigma factors of the s70 type that recognize promoters resembling bacterial promoters (Liere and Börner, 2007). Chloroplast ribosomes share many features with bacterial ribosomes as ribosomal RNAs (rRNAs) and 21 ribosomal proteins from the small subunit (30S) have clear orthologs in *Escherichia coli*. Similarly, the large ribosomal subunit (50S) composed of three rRNAs (4.5S rRNA, 5S rRNA and 23S rRNA) and 31 ribosomal proteins have orthologs in *E.coli*. Interestingly, 12 out of 21 proteins associated with 30S and 22 out of 31 proteins associated with 50S ribosomal subunits are encoded by the nucleus rather than the chloroplast genome (Yamaguchi and Subramanian, 2000). Many of these proteins are essential in *E.coli* and plants (Rogalski *et al.*, 2006; Fleischmann *et al.*, 2011; Tiller and Bock, 2014). Also, several ribosome assembly factors, tRNAs, initiation, elongation, and termination factors are conserved between plastids and *E.coli* (Britton, 2009; Alkatib *et al.*, 2012; Meurer *et al.*, 2002; Albrecht *et al.*, 2006; Motohashi *et al.*, 2007; Shen *et al.*, 2013).

Ribosomal biogenesis is among the most energy consuming processes of the cell, and involves stepwise processing of the primary rRNA transcript and the correct assembly of ribosomal proteins and rRNAs into functional ribosomal subunits. This highly coordinated process is facilitated by many trans-acting factors, including accessory proteins and small RNAs (Fromont-Racine *et al.*, 2003; Kaczanowska and Rydén-Aulin, 2007). *In-vitro* reconstitution of ribosomes without trans-acting factors requires high ionic strength, heating and prolonged incubation time (Nierhaus and Dohme, 1974; Traub and Nomura, 1969), indicating the essential role of these factors under physiological conditions. Not surprisingly, more than half of these trans-acting factors are ATP- and GTP-binding proteins (rATPases and rGTPases) that provide energy for processes such as breaks and inter/intramolecular rearrangements, thus allowing association and dissociation of other assembly factors or ribosomal proteins for sequential progress of the ribosome assembly (Shajani *et al.*, 2011).

Since many plastid ribosomal proteins and assembly factors are conserved
between chloroplast and *E.coli*, and many of these proteins can be integrated into
E.coli ribosomes, a great deal of information regarding bacterial ribosome biogenesis
can be conveyed to the chloroplast ribosomes (Bubunenko and Subramanian, 1994;
Weglohner *et al.*, 1997). The detailed assembly maps for both ribosomal subunits
and the necessary assembly factors needed for rRNA processing and/or ribosome
maturation from *E.coli* are conserved in chloroplasts, despite more than 1.6 billion
years since the endosymbiosis of the cyanobacterial ancestor of chloroplast (Dabbs,
1991; Talkington *et al.*, 2005; Kaczanowska and Rydén-Aulin, 2007; Woodson,
2008; Connolly and Culver, 2013; Fournier *et al.*, 2010; Davies *et al.*, 2010; Jacob *et*
al., 2013). However, since more than 90 % of endosymbiont bacterial genes
migrated to host cell nuclear genome (Sato, 1999; Timmis *et al.*, 2004), all of the
ribosome biogenesis factors are nuclear-encoded. Consequently, their site of action
in a plant cell (chloroplast or mitochondrion or both) remains to be determined.

RsgA is among the most extensively studied *E.coli* rGTPases. The null
mutants of RsgA in *E.coli* and *Bacillus subtillis* show decreased growth rates,
decreased levels of mature 70S ribosomes, and accumulation of 17S RNA
(precursor of 16S RNA), indicating its role in the biogenesis of the 30S ribosomal
subunit (Campbell *et al.*, 2005; Himeno *et al.*, 2004). However, its *Arabidopsis*
thaliana homolog null mutant shows embryo lethality (Tzafrir *et al.*, 2004), indicating
that the assembly pathways of 70S ribosomes in bacteria and chloroplasts might
have some subtle differences. In general, the rGTPases associate with the immature
ribosomal subunits in GTP-bound form, resulting in enhanced GTPase activity,
followed by dissociation of GDP-bound rGTPase from the subunit (Goto *et al.*, 2011).
Structurally, most rGTPases possess a nucleotide binding site and a GTPase
domain. The latter consists of five conserved motifs; G1 to G5 (Leipe *et al.*, 2002).
The GTPase domain of several rGTPases is organized in a non-canonical order of
G4-G5-G1-G2-G3, often referred to as circular permutated GTPase domain (Leipe *et*
al., 2002; Elias and Novotny, 2008). Structural studies on immature 30S subunit from
ΔrsgA E.coli revealed severe distortion at the decoding center of the 30S subunit,
which makes it unable to associate with the 50S subunit to carry out mRNA
translation (Jomaa *et al.*, 2011). A recent study reports that *ΔrsgA* in *E.coli*, the
immature 30S subunit can still assemble with a mature 50S subunit, although at a
slower rate, explaining the slower growth rate phenotype (Thurlow *et al.*, 2016).

However, it's function and site of action in plants remains unclear, which prompted
us to study Arabidopsis thaliana RsgA (AtRsgA) gene in detail.

Results

AtRsgA can complement the small ribosomal subunit maturation deficiency in
an *E. coli ΔrsgA* mutant

E.coli RsgA is involved in biogenesis of the small ribosomal subunit (30S). Its null
mutant is characterized by slow growth under favourable growth conditions (Kimura
et al., 2008; Goto *et al.*, 2011). We identified homologs of *E.coli* RsgA in a
cyanobacterium, a green alga, and a plant, demonstrating that RsgA belongs to a
conserved protein family (Figure S1). The multiple protein sequence alignment
revealed a conserved oligonucleotide/oligosaccharide binding motif (OB) and a
GTPase domain. We also observed a plant-specific sequence insertion within the
GTPase domain (Figure S1, marked in red box).

To investigate if the *Arabidopsis thaliana* RsgA homolog (AtRsgA) protein is
also involved in 30S maturation, we transformed *E.coli ΔrsgA* null mutant with full-
length *AtRsgA* cDNA. We observed that *AtRsgA* can partially rescue the growth
phenotype in liquid (Figure 1A) and solid media (Figure 1B), when compared to
ΔrsgA and *ΔrsgA* transformed with empty vector (EV). To confirm if the rescue of
slow growth phenotype is due to improved 30S subunit biogenesis, we quantified
ribosomal RNAs (rRNAs) which are indicative of the amount of their respective
ribosomal subunits (Tiller *et al.*, 2012)(Figure S2). We observed an increased ratio of
16S rRNA (30S subunit) to 23S rRNA (50S subunit) in *AtRsgA* complemented *ΔrsgA*
as compared to the controls (Figure 1C). This demonstrated that AtRsgA can
partially rescue the 16S maturation deficiency and confirmed that AtRsgA is a 30S
assembly factor.

Arabidopsis RsgA is essential for morphogenesis and viability

To decipher the potential role of *AtRsgA* in *Arabidopsis thaliana*, we analysed two
transfer DNA (T-DNA) insertion lines, *rsgA-i* (insertion in the fourth intron) and *rsgA-e*
(insertion in the third exon, Figure 2A). We could not obtain homozygous plants for
the *rsgA-e* allele, and heterozygous plants produced wild-type (WT):heterozygous
plants in 1:2 ratio (31:62, as defined by PCR), indicating that the *rsgA-e* allele is
recessive and embryo or seedling lethal in homozygous state. Real-time PCR

analysis showed significantly reduced *AtRsgA* transcript in both lines, with

homozygous *rsgA-i* showing a stronger decrease (Figure 2B). While the

heterozygous *rsgA-e* plants showed a wild-type like morphology, the homozygous

rsgA-i plants were severely dwarfed and chlorotic at the rosette stage (Figure 2C)

and as mature plants (Figure 2D). The dwarfed growth of *rsgA-i* is marked with both

reduced growth rate (Figure S3A), rosette area (Figure S3B), and a significant

reduction in seed yield (Table S1, Figure 2E). Interestingly, although heterozygous

rsgA-e plants showed no net reduction in seed yield, 24.9±2.7 % of immature seeds

were white (Figure 2D-E, Table S1), suggesting that the white seeds were

homozygous for the *rsgA-e* allele and not viable.

To show that the T-DNA insertions in *AtRsgA* are causing the observed

phenotypes, we analyzed progeny of heterozygous *rsgA-e* pollinated with

homozygous *rsgA-i*, and observed the expected wild-type (WT/*rsgA-i*):dwarf (*rsgA-*

e/*rsgA-i)* chlorotic ratio of phenotypes and genotypes (Table S2, Figure S4A, X^2 P =

0.097 testing significantly different phenotype distribution). This showed that *AtRsgA*

mutations are responsible for the observed phenotypes in both mutant alleles.

Arabidopsis RsgA is involved in 30S ribosome maturation in chloroplasts

The chlorotic phenotype of *rsgA-i,* together with its ability to rescue a 30S assembly

factor mutant in *E.coli* indicated that AtRsgA was involved in the maturation of the

small subunit of chloroplasts. To verify this in plants, we compared rRNA profiles of

rsgA-i and WT plants using a bioanalyzer electropherogram. The cytosolic plant

rRNAs appear in form of two major peaks corresponding to 25S (60S subunit) and

18S (40S subunit) rRNA molecules, while the chloroplast rRNA profile is presented

as a single peak for 16S rRNA (30S subunit) and three smaller peaks for 23S (23-1,

23-2, 23-3, 50S subunit) rRNA, which are generated post-transcriptionally by two

hidden breaks (Figure S5, Figure 3A). The unchanged ratio of cytosolic subunits

(25S/18S) in *rsgA-i* and WT samples indicated that the cytosolic ribosome

biogenesis was not affected in *rsgA-i.* However, the 16S rRNA was significantly

decreased as noted by decrease in 16S/23S-1 (Figure 3B, small and large

chloroplast subunit) and 16S/18S (Figure 3C, small subunits from chloroplast and

cytosol, respectively) amount in *rsgA-i* as compared to WT. The ratio of large

chloroplast subunit was decreased when compared with the small cytosolic subunit

(23S-1/18S, Figure 3C), but the change was less pronounced than for 16S.

To corroborate the finding that 16S rRNA maturation was specifically affected
and to identify the steps where the maturation pathway was perturbed, we performed
Northern blot analysis using a specific probe against mature 16S rRNA (Figure 3A).
As positive controls for small chloroplast subunit defects, we included two known
chloroplast 30S assembly factor mutants *rif1* (resistant to inhibition by FSM)(Flores-
Pérez *et al.*, 2008) and *bpg2* (Brz-insensitive-pale green 2)(Kim *et al.*, 2012). While
in WT, the mature 16S rRNA was the most abundant rRNA, *rsgA-i*, *rif1* and *bpg2*
mutants showed accumulation of 17S rRNA, a precursor of the 16S rRNA (Figure
3D)(Fristedt *et al.*, 2014). Northern blot analysis showed normal processing for
mitochondrial 18S (Figure S6A), mitochondrial 26S (Figure S6B), cytosolic 18S
(Figure S6C) and chloroplast 23S rRNAs (Figure S6D), indicating that the *rsgA-i*
defect is causing a specific maturation deficiency of chloroplast 16S rRNA. To
ascertain if 17S rRNA precursor accumulation in *rsgA-i* was due to processing defect
at 5' and/or 3' termini of 16S, we performed northern blot analysis using specific
probes for the respective termini. The results showed that both 5' and 3' termini were
not efficiently processed in the three mutants (Figure 3D).

Mutants of several *E.coli* ribosome assembly factors show temperature
sensitivity (Charollais *et al.*, 2003; Charollais *et al.*, 2004; Connolly *et al.*, 2008).
Moreover, elevated temperatures can partially rescue lethality of chloroplast
ribosome biogenesis defects in tobacco (Ehrnthaler *et al.*, 2014). Indeed, *rsgA-i*
mutants showed a more severe phenotype at lower temperature (18 °C) as
compared to normal growth temperatures (23 °C, Figure S7). However, elevated
temperature (28 °C) could not rescue the chlorotic phenotype observed at 23 °C
(Figure S7A). Interestingly, although at lowered temperature (18 °C) *rsgA-i* seedlings
exhibited more severe phenotype, this was not accompanied with visibly increased
accumulation of 17S precursor, as compared to higher temperatures (23 °C and 28
°C, Figure S7B). Similar results were obtained for *rif1* and *bpg2* (Figure S7). Taken
together, *rsgA-i*, *rif1* and *bpg2* are potentially active at a similar step of 30S subunit
maturation.

To test if 17S rRNA can be integrated into mature ribosomes, we performed a
polysome loading experiment. Polysomes consist of translating ribosomes in
complex with mRNA, which can be separated from single ribosomes in a sucrose
gradient upon centrifugation. Low sucrose fractions contain free subunits,
monosomes and disomes, while high sucrose fractions contain polysomes, multiple

ribosomes associated with one mRNA. While WT produced a single 16S rRNA in
monosomes and polysomes (Figure 4A), the *rsgA-i* showed presence of 16S and
17S rRNA precursor in low (fractions 3-5) and high (fractions 11-12) density fractions
(Figure 4B). The presence of 17S in high density fractions indicated that the
immature 17S rRNA was incorporated into translating ribosomes. Furthermore, the
lower ratio of 17S/16S in the high density fractions (Figure 4C), when compared to
the low density fractions, indicated that ribosomes containing 17S rRNA tend to be
less associated with mRNA.

RsgA mutation leads to deficiency of the photosynthesis machinery
To corroborate the function of *AtRsgA* in chloroplast ribosome biogenesis, we
performed *AtRsgA* expression analysis using Real-time PCR, and observed higher
expression of the gene in green photosynthesizing organs as compared to non-
photosynthesizing organs (roots and inflorescence), with additionally highest
expression in mature seeds (Figure 5A). The expression of *AtRsgA* in green tissues
was confirmed by high GUS (β-glucuronidase) activity in transgenics expressing
GUS driven by the *AtRsgA* promoter (Figure 5B).

To estimate the influence of the ribosome assembly defect on chloroplast
function, we first estimated the levels of tetrapyrroles. Tetrapyrrole biosynthesis
takes mainly place in chloroplast stroma, and through branching leads to production
of chlorophyll *a*/*b*, hemes, siroheme, phytochromobilin and xanthophyll molecules
(Mochizuki *et al.*, 2010). Xanthophylls are most abundant in the light-harvesting
complexes and their main function is photoprotection by channelling excess energy
away from chlorophylls (reviewed in Ruiz-Sola and Rodríguez-Concepción, 2012).
Interestingly, while the amount of chlorophylls was decreased in the *rsgA-i* mutant,
xanthophylls remained unchanged, suggesting a differential regulation of the two
compounds (Figure 5C).

Next, we measured several chlorophyll-a fluorescence parameters, such as
rhe Fv/Fm ratio (maximum quantum efficiency of photosystem II photochemistry),
linear electron transport rate (ETRII) and non-photochemical quenching (qN,
dissipation of excess energy as heat). The results showed a decreased Fv/Fm ratio
(Figure 5D) and linear electron transport capacity (Figure 5E), while the induction of
non-photochemical quenching was strongly shifted to lower light intensities, and

maximum non-photochemical quenching capacity was increased (Figure 5F),
indicating reduced photosynthesis efficiency in *rsgA-i* plants.
265

RsgA is important for proper plant and chloroplast morphology
To investigate the effect of potentially decreased chloroplast protein synthesis on
chloroplast biogenesis, we investigated mesophyll cells of *rsgA-i* and WT seedlings
by microscopy. The analysis revealed that chloroplasts in mesophyll cells of *rsgA-i*
mutant plants were smaller than in WT (Figure 6A). We also observed that while the
average cell size was not significantly different to WT plants in *rsgA-i*, chloroplast
number per cell was significantly higher in *rsgA-i* (Figure 6B, $P < 0.0001$).
Furthermore, despite the higher number per cell, the chloroplasts in *rsgA-i* line
occupy less area in the cell due to their smaller size (Figure 6B).

Despite the similarity in mesophyll cell size, *rsgA-i* leaves were smaller
compared to WT (Figure 2C, D), which could be caused by a decreased cell number
and/or altered anatomy of the leaves. Therefore, we analyzed leaf cross-sections by
microscopy, and in agreement with the previous result (Figure 6B), we observed no
differences in cell sizes between *rsgA-i* and WT, and decreased chloroplast area
occupancy in *rsgA-i* (Figure 6C). However, the mutant had visibly less intercellular
air space between the mesophyll cells (Figure 6C), suggesting that increased cell
packing might also play a role in the decreased leaf size.

In higher plants, most of the aboveground cells originate from a small group of
undifferentiated, dividing cells called shoot apical meristem (SAM). The SAM
consists of multiple cells layers (L_{1-3}) and central rib meristem part. The shoot
meristematic cells contain 10 to 20 proplastids (Possingham and Lawrence, 1983),
which multiply to more than 100 chloroplasts in mesophyll cells of fully expanding
leaves (Lopez-Juez and Pyke, 2005). To investigate if SAM morphology and
chloroplast biogenesis were affected in *rsgA-i*, SAM of 4 weeks old plants was
analyzed by confocal microscopy. We observed a severely reduced diameter of SAM
in *rsgA-i* (Figure 6D). Furthermore, chlorophyll fluorescence, which is indicative of
chloroplasts, was mostly located in rib meristem region in WT, while in *rsgA-i* the
signal was largely absent (Figure 6D). Interestingly, proplastids accumulated more in
some cells of L_1 in *rsgA-i,* compared to WT plants (Figure 6D). These results indicate
a connection between plastidial protein synthesis, chloroplast biogenesis and SAM
formation.

Impact of defective chloroplast protein synthesis on the transcriptome and proteome

A defect in plastid ribosome biogenesis machinery is expected to cause dramatic changes to the transcriptome and proteome of mutant plants, as it needs to adjust to the severely perturbed chloroplast function. To understand the molecular basis of this adjustment, we carried out transcriptomic (RNA sequencing) and proteomic (LC-MS/MS) analysis of *rsgA-i* and WT plants.

We first investigated how mRNA and protein expression of gene products targeted to different cellular compartments was perturbed in *rsgA-i*. The analysis revealed that most of the plastid-encoded transcripts tended to be up-regulated, while the corresponding proteins accumulated less (Figure 7A, Figure S8A for all compartments). This effect was also seen to a weaker degree for chloroplast-targeted nuclear-encoded genes. We also determined which gene products show significantly ($P < 0.05$) different expression in the mutant, and identified 1676 up-regulated and 1678 down-regulated transcripts, respectively. The proteomic analysis revealed 42 and 114 up- and down-regulated proteins, respectively (Table S3 for the transcriptome, and Table S4 for the proteome). Similarly to the previous result, the compartment with the largest number of differentially regulated elements was the chloroplast, where the majority of chloroplast-encoded and -targeted transcripts showed up-regulation. Conversely, the corresponding proteins showed significant decrease (Figure 7B, Figure S8B for all compartments). Indeed, this opposite behaviour of mRNAs and proteins corresponded to the largest fraction of differentially expressed proteins (Figure 7C, 64/156 proteins).

To investigate which biological pathways were affected by the decreased protein synthesis in chloroplast, we performed a MapMan enrichment analysis (Thimm *et al.*, 2004). Transcripts significantly upregulated in *rsgA-i* were involved in photosynthesis (light reaction, photorespiration, Calvin-Benson cycle) and chloroplast protein synthesis (Figure 7D, blue cells, P<0.05). Conversely, significantly downregulated transcripts belonged to pathways involved in biotic stresses, acid phosphatases, different transcription factor families, post-translational modification, and other signalling pathways (Figure 7D, red cells). MapMan analysis of the proteome revealed that photosynthesis was the only significantly enriched MapMan bin and most of these proteins were decreased (Figure 7D, P < 0.05). We

confirmed these results by carrying out qRT-PCR and western blot analysis for some

of the differentially expressed genes and proteins involved in photosynthesis.

Indeed, transcripts of photosystem I (*psaA*, *psaB*), cytochrome b6 (*petA*), ATP

synthase (*atpA*) were upregulated in *rsgA-i* (Figure S9A), while AtpA, PsaA and

RbcL (RuBisCO) proteins were decreased (Figure S9B). Taken together, these

results indicate that the decreased protein synthesis in chloroplast is met with an

upregulation of photosynthesis- and protein synthesis-related transcripts.

Analysis of chloroplast retrograde signalling pathways

To better understand the mechanism that induced higher expression of nuclear-

encoded chloroplast-targeted transcripts and increased chloroplast number, we

studied the retrograde signalling pathways. These pathways can control expression

of nuclear genes by metabolic and protein signals to preserve plastid homeostasis in

changing environment (operational signalling) and during development (biogenic

signalling, reviewed in Chan *et al.*, 2016). Still unknown signals from chloroplasts are

conveyed to the nucleus, where they modulate the expression of photosynthesis-

associated nuclear genes (PhANGs), plastid redox–associated nuclear genes

(PRANGs), singlet oxygen–responsive genes (SORGs) and defense response

genes. The proposed metabolic signals (orange boxes) and their biosynthesis

pathways and genes they regulate (green boxes) are shown on Figure 7E.

Metabolic signals which are not likely candidates behind the observed

phenotypes include 3-phosphoadenosine 5-phosphate (PAP), methylerythritol

cyclodiphosphate (MEcPP), Heme and Dihydroxyacetone phosphate (DHAP). PAP

alters gene expression by inhibiting the RNA-degrading activity of XRNs which target

APX2 (Estavillo *et al.*, 2011), whose transcript accumulation was unchanged in *rsgA-

i* (Figure 7E). MEcPP induces wounding and high light stress-related nuclear gene

expression of the chloroplast-targeted hydroxyperoxide lyase (HPL1) protein (Xiao *et

al.*, 2012), which was unchanged in *rsgA-i* (Figure 7E). Heme acts as a positive

retrograde signal from plastids, and overexpression of Ferrochelatase 1 (FC1)

transcript increases expression of PhANGs (Woodson *et al.*, 2011). However, *rsgA-i*

showed downregulation of FC1, while expression of PhANGS was increased (Figure

7E). DHAP activates MPK6, which subsequently induces the nuclear transcription of

ERF-TFs (such as *ERF6*, Vogel *et al.*, 2014), which remained unchanged in *rsgA-i.*

The metabolic signals that could cause the observed phenotypes in the *rsgA-i*

mutant include Mg–protoporphyrin IX (Mg-ProtoIX), chloroplast biogenesis 5 (CLB5)

and beta-cyclocitral (β-cyc). Mg-ProtoIX is an intermediate in chlorophyll

biosynthesis (Kim *et al.*, 2013) and might act by suppressing PhANG expression by

binding to Long Hypocotyl 5 (HY) via Heat Shock 90 (HSP90)(Kindgren *et al.*, 2011).

Chlorophyll (and likely Mg-ProtoIX) is decreased in *rsgA-i*, while expression of

multiple PhANGs is increased (Figure 7E). Loss of CLB5 causes a decrease in

carotenoid biosynthesis, and repression of genes involved in the biosynthesis of

isoprenoid precursors (DXS1 and HDR)(Avendano-Vazquez *et al.*, 2014). The latter

showed increased mRNA levels in *rsgA-i*. Finally, β-cyc expression increases at

elevated $^1O_2$ and induces expression of multiple genes, such as GST12 and LTI30

(Ramel *et al.*, 2012), of which LTI30 was downregulated on the transcript level.

The protein signals are shown on Figure S10, together with expression

analysis of 121 PhANGs (Zones *et al.*, 2015), 418 SORGs (Gonzalez-Perez *et al.*,

2011), 27 defense genes (Desveaux *et al.*, 2004), 2 PRANGs (Chan *et al.*, 2016)

and 120 chloroplast-encoded genes (Table S3 for transcripts, Table S4 for proteins).

The majority of the plastid-targeted nuclear-encoded genes were upregulated on the

transcript level, while no changes were observed on the level of proteins (e.g.

PRIN2, SG1, HDS). PhANGs and chloroplast-encoded genes were upregulated on

the level of transcripts, but down-regulated on the level of proteins (39 and 59

proteins for PhANGs and chloroplast-encoded proteins analysed, respectively).

SORGs and defense genes were down-regulated on the level of transcripts, with no

observable differences in protein levels (397 and 17 proteins analysed for SORGs

and defense genes, respectively). PRANG marker genes (APX2 and ERF6) showed

no changes on the level of transcripts, indicating that the PRANG signalling pathway

is not activated in *rsgA-i*. Of the nucleus-localized signals, only GLK transcription

factors showed differential transcript expression. GLK1 and GLK2 induce expression

of PhANGs and are down-regulated during chemically-inhibited plastid protein

synthesis (Waters *et al.*, 2009). Surprisingly, GLK2 was upregulated on the

transcriptional level in *rsgA-i*, suggesting that GLK2 might be responsible for higher

expression of PhANGs.

Several genetic perturbations can alter chloroplast numbers. These include

RSH3, a guanosine penta- and tetraphosphate synthase, whose overexpression

causes an increased number of chloroplasts, but decreased chlorophyll level and

chloroplast diameter (Sugliani *et al.*, 2016). Plastid division (PDV) mutants display
lowered number of large chloroplasts, while the over-expression line shows a higher
number of smaller chloroplasts (Okazaki *et al.*, 2009). Finally, gibberellin (GA)
deficient mutants have increased number of chloroplasts and chlorophyll, and
similarly to *rsgA-i*, less air space between cells. However, the expression level of
RSH3, PDV and GA-regulated genes, such as GNC and GNL (Figure S10)(Park *et*
al., 2013), remained unaltered in *rsgA-i* suggesting that increased chloroplast
number is potentially regulated by another, yet unknown mechanism.

Discussion
This work elucidates the function of the Arabidopsis AtRsgA protein, as a novel
chloroplast ribosome assembly factor. Despite more than 1.6 billion of years since
the endosymbiotic event that gave rise to chloroplasts, AtRsgA could partially
complement the slow growth and defect in 30S maturation of *E.coli ΔrsgA* mutant
(Figure 1). The incomplete complementation could be attributed to the fact that
AtRsgA was not codon-optimized for *E.coli*, due to minor structural differences
between the *E.coli* and chloroplast ribosomes, or differences in pH, GTP availability
and temperature between the cytosol of bacteria and plants. While *E.coli ΔrsgA* is
viable (Figure 1)(Thurlow *et al.*, 2016), the Arabidopsis *AtRsgA* is essential, as we
could not obtain homozygous T-DNA insertion mutants of *rsgA-e* (Figure 2E). This
suggests that the ribosome assembly (or the requirement of having a certain pool of
functional ribosomes) is in some aspects different between *E.coli* and chloroplasts.
Indeed, RbfA, an interaction partner of RsgA, shows processing defects in the 5' end
region of the 16S rRNA in *E.coli* (Xia *et al.*, 2003; Datta *et al.*, 2007), while the
chloroplast counterpart shows defects in both 5' and 3' end processing (Fristedt *et*
al., 2014). This suggests that ribosome assembly can be modified, and the assembly
factors can gain or lose certain functions. The presence of a plant-specific region in
the GTPase domain is especially interesting in this regard (Figure S1), and
identification of its function warrants further investigation.

The *rsgA-i* intron mutant showed a typical chloroplast deficiency phenotype,
such as chlorotic leaves with reduced rosette area (Figure 2), depleted
photosynthetic pigments and reduced photosynthesis (Figure 5). These phenotypes
have been described for numerous mutants with disturbed chloroplast biogenesis,
chlorophyll production (Liu *et al.*, 2015; Motohashi *et al.*, 2007; Maekawa *et al.*,

2015) and chloroplast ribosome assembly (Fristedt *et al.*, 2014; Liu *et al.*, 2010;
Komatsu *et al.*, 2010). These defects ultimately resulted in a massive growth
retardation and decrease in rosette area of the mutant (Figure S3). Mechanistically,
the rsgA-i mutant displayed a specific defect of the 16S rRNA accumulation that was
accompanied by an over accumulation of 17S precursor in chloroplasts, whereas
rRNA accumulation of cytosolic and mitochondrial ribosomes was unaffected.
Furthermore, AtRsgA was shown to interact with RbfA, which is absent in
mitochondria (Goto *et al.*, 2011; Fristedt *et al.*, 2014). Together, these evidences
strongly suggest that AtRsgA is important for maturation of the small chloroplast
subunit (Figure 3). While *rsgA-i* also shows a modest, but significant decrease in 23S
rRNA (50S subunit, Figure 3C), we have demonstrated that no precursors
accumulated in processing pathways of rRNAs from chloroplast 23S rRNA. This
suggests that the free 50S subunit can be degraded either (i) due to a lower level of
the 30S subunit or (ii) by decreased translation of chloroplast-encoded components
of 50S (such as L2 and L22). Interestingly, *rbf1* which acts at a similar step of 30S
subunit maturation (Figure 3D)*,* showed no differences in 50S subunit accumulation
(Fristedt *et al.*, 2014), suggesting some dissimilarity in the role of these two proteins.

The negative correlation between chloroplast number and chloroplast size has
been well known for over a decade (Figure 5 A,B), but the sensing and actuating
mechanism underlying this phenomenon is largely unknown (Pyke, 1997). Similarly
to *rsgA-i,* the RSH3 over-expression (controlling homeostasis of penta- and tetra-
phosphate)(Sugliani *et al.*, 2016) and PDV over-expression (Plastid Division
Protein)(Okazaki *et al.*, 2009) result in a larger number of smaller chloroplasts.
However, transcripts of these genes did not show a significantly altered expression
(Figure 7), suggesting that the herein observed increased chloroplast number is
controlled by another mechanism.

Nuclear- and chloroplast-encoded photosynthetic proteins were significantly
less abundant in *rsgA-i* (Figure 6C, D). While decrease of chloroplast-encoded
proteins could be explained by the decreased protein synthesis capacity in
chloroplast, the decrease of the nuclear-encoded photosynthesis-related proteins
took place despite the increase of the corresponding transcripts. This degradation
most likely takes place in chloroplasts rather than cytosol, and we suggest that it is
caused by the inability to assemble chloroplast complexes that consist of both
nuclear- and chloroplast-encoded subunits. Indeed, damaged and unassembled

protein complexes have been shown to be rapidly degraded in chloroplasts (Adam
and Ostersetzer, 2001).

Transcriptomic analysis of *rsgA-i* showed a potential compensatory response
of protein translation deficiency in chloroplast, by upregulation of chloroplast- and
photosynthesis-associated nuclear genes (PhANGs, Figure 7E, Figure S10). To
better understand which signalling pathway induced the expression of
photosynthesis-related transcripts in the nucleus, we analysed the metabolic and
protein components of the retrograde signalling pathways. The analysis of
differentially expressed genes revealed that the altered metabolic signal candidates
are Mg–protoporphyrin IX (precursor of chlorophylls), CLB5 or β-cyc. Mg-
protoporphyrin signal IX is a likely candidate, as the metabolite acts as a
suppressant of PhANG expression (Kindgren *et al.*, 2011), and the chlorophyll (and
most likely Mg-protoporphyrin IX) levels are decreased in *rsgA-i* (Figure 5C), while
PhANGs are upregulated (Figure 7E, Figure S10). The *clb5* mutant shows
downregulaton of photosynthesis-related genes (Avendano-Vazquez *et al.*, 2014),
while we found that *CLB5* transcript and PhANGs are upregulated in *rsgA-i* (Figure
S10). Treatment of *rsgA-i* with products and inhibitors specific to these signalling
pathways (e.g. feeding the plants with Mg-protoporphyrin IX to suppress PhANG
expression or treatment of *clb5* with fluridone to suppress its chlorotic
phenotype)(Avendano-Vazquez *et al.*, 2014), could reveal which of these signals
plays a role in the observed phenotypes.

Taken together, our work has identified a novel assembly factor needed for
maturation of 30S ribosomal subunit in chloroplasts, while the transcriptomic and
proteomic analysis revealed the responses underpinning the adaptation to a
decreased chloroplast function.

Materials and methods

Cloning of *E.coli* complementation vector and Arabidopsis GFP constructs.
AtRsgA cDNA fragment was amplified by PCR from ABRC clone (S80886) using a
high-fidelity polymerase (Phusion; New England Biolabs) and primers rsgA_fwd and
rsgA_rv/rsgA_rv (Table S5). The product was inserted into pENTR4 vector using
restriction based cloning (NcoI/NotI restriction sites) strategy and shuttled to pET60
(for *E.coli* complementation) or pUBC/pUBN destination vector (for GFP fusion

constructs) using Gateway cloning strategy. *AtRsgA* promoter region was amplified

from *Arabidopsis thaliana* Col-0 genomic DNA using primers promotor_fwd and

promotor_rv (Table S5), digested with BamHI and NcoI and ligated with T4 ligase

(Roche) into pCambia 1305 vector.

E.coli ΔrsgA mutant complementation

ΔrsgA mutant was transformed with pET60 vector with either pT7::GST-AtRsgA-His

or pT7::GST-HIS (empty vector without ccdB cassette) expression cassette via

electroporation. Three colonies from transformed/ non-transformed *ΔrsgA* and WT

E.coli were inoculated in 5mL LB medium and grown overnight at 28□C and 180

RPM shaking. 50 µl of overnight cultures were used to inoculate 5 ml fresh LB

medium without antibiotics to achieve starting culture OD_{600} = 0.005. Cultures were

grown for 8 h at 28□C with 180 rpm shaking, and OD_{600} was measured every hour.

For plate growth experiment, 10 µl of each starting culture (OD_{600} = 0.005) was

plated on LB agar plates without antibiotics and grown overnight at 28□C.

Plant growth conditions and genotyping

Two independent T-DNA lines, *rsgA-e* (GABI_538D07) and *rsgA-i* (GK-716H11)

were used for characterization of *RsgA* (*At1g67440*). Mutant plants were genotyped

with primers *rsgA-e* fwd and *rsgA-e* rv for GABI_538D07, primers *rsgA-i* fwd and

rsgA-i rv for GK-716H11, and T-DNA left border primer GABI LB for GABI lines

(Table S1), with PCR parameters as recommended for GABI lines. For growth on

sucrose-containing medium, seeds were surface sterilized and plated on Murashige

and Skoog agar medium containing 1 % Suc (Murashige and Skoog, 1962) and

grown in growth chamber under 16-h day (140 µmol m^{-2} s^{-1}) / 8-h night regime. For

growth in soil, two weeks old seedlings were transplanted into soil and grown in the

greenhouse at 16-h day (250 µmol m^{-2} s^{-1}) / 8-h night regime at 20 /16 °C

(day/night).

rRNA analysis with Bioanalyzer

Total RNA from *E.coli* was isolated using TRizol reagent (Thermo Fisher Scientific).

RNA from seedlings was harvested using Spectrum Plant Total RNA Kit (Sigma-

Aldrich). rRNA ratios were analyzed with an Agilent 2100 Bioanalyzer using the

Agilent RNA 6000 nano kit following manufacturer's instructions. rRNA ratios were

determined as described previously (Tiller *et al.*, 2012). Bacterial electropherogram
curves for 16S and 17S rRNA overlap, due to which peak area calculation using
Bioanalyzer software could not be used. Therefore, peak area was calculated with
normalmixEM function from the R package mixtools, which was applied to the
sampled data in order to deconvolute the peak as a mixture of two Gaussian
distributions. Pairwise Student t-test was used to calculate the significance of the
differences in the ratio of the areas.

Northern blotting
Northern blot analysis was essentially performed as previously described (Beick *et*
al., 2008). In bried, total RNA was separated on formaldehyde-containing 1.2 % (w/v)
agarose gels and blotted onto Hybond N membranes (GE Healthcare). Hybridization
probes (Table S6) were generated by PCR amplification and purified by agarose gel
electrophoresis, or commercially synthesized and then labeled with [α-32P]dCTP by
random priming (Multiprime DNA labelling system; GE Healthcare) or with [γ-
32P]dATP by 5' end phosphate exchange (T4 PNK, New England Biolabs),
respectively. Hybridizations were performed at 65 °C.

Polysome loading assay
The analysis was performed as in (Barkan, 1993), with gradient concentrations of 15
%, 30 %, 45 % and 60 %. Samples were ultracentrifuged in 9 ml tubes for 4 hours at
37000 rpm in 4□C.

Quantitative real-time PCR
cDNA used for qRT-PCR was obtained via Reverse transcriptase polymerase chain
reaction using SuperScript™ IV Reverse Transcriptase (Invitrogen). qRT-PCR was
carried out using the Maxima SYBR® Green/ROX qPCR Master Mix (Thermo
Scientific, UK). The expression levels of three technical replicates for each three
biological replicates were measured on One Step Plus real time PCR system
(Applied Biosystems, Life Technologies, Germany). Equipment setup, data collection
and analysis were done using ABI SDS 2.3 software (Applied Biosystems, Life
Technologies, Germany). The relative expression of target genes was normalized
separately to housekeeping genes *Ubiquitin10* (*UBQ10*) and *Actin*. The samples

which gave similar relative expression to both reference genes were considered for
analysis. Data were analyzed by Livak method (Livak and Schmittgen, 2001).

568 Protein Isolation, Gel Electrophoretic Separation and Immunodetection

100 mg of plant tissue was ground to fine powder and 150 µl of extraction buffer (40
mM Tris-HCl pH 7.0, 4 % SDS, 5 % glycerol, 2x Protein inhibitor cocktail) was
added. Samples were vortexed for 5 min and spun down at 13000g for 30 min at
12□C. Supernatant containing total protein extract was collected and used for further
experiments. The protein content was measured using BCA assay kit (Thermo
Fisher Scientific). Total protein extract was separated by SDS-PAGE (6 % stacking
gel, 12 % separation gel), and the proteins were subsequently transferred to
polyvinylidene difluoride membranes (Immobilone; Millipore). Membranes were
blocked with 5 % milk for 1 hour at room temperature with gentle shaking and then
incubated with the primary antibody overnight at 4 °C. Blotted membranes were
washed five times for 5 minutes using Tris-buffered saline with 1 % (v/v) Tween 20
(TTBS) buffer and incubated at room temperature with horseradish peroxidase-
conjugated secondary antibody for 1 hour. The membranes were then washed five
times for 5 minutes with TTBS buffer, and the immunoreactive proteins were
visualized following 5 minutes incubation with detection reagents from the
SuperSignal WestPico horseradish peroxidase detection kit (Thermo Scientific).
Quantification of the immunoblots was done using the Fujifilm LAS-1000 software.

587 Measuring plant growth kinetics and photosynthesis parameters

Total rosette leaf area was measured with the IMAG3D Maxi-Version of the Imaging
PAM M-series, which was equipped with a 3D scanner based on a structured
illumination approach (Heinz Walz GmbH, Effeltrich). Rosette leaf area was
determined 21, 24, 27 and 30 days after sowing. To determine chlorophyll *a*
fluorescence parameters on intact rosettes, the Maxi-version of the Imaging-PAM M-
series was used. Light response curves between 0 and 1000 µE $m^{-2}s^{-1}$ actinic light
intensity were measured with measuring times per intensity ranging between 150
seconds under light-limited and 60 seconds under light-saturated conditions. Plants
were dark-adapted for 30 min prior to all measurements. Linear electron transport
(ETRII) was corrected for leaf absorbance.

Leaf cross-section analysis

For leaf cross section, two weeks old seedlings were fixed with 4 % paraformaldehyde and 0.2 % glutaraldehyde by vacuum infiltration and incubated at 4 °C overnight. The samples were washed in 0.1M phosphate buffer (pH 7.4), and dehydrated in an ethanol series (30, 50, 70, 80, 90 and 100 %) with incubation times of 1 hour in each solution. Samples were infiltrated with Technovit 7100 resin (Heraeus Kulzer) for 24 hours, followed by polymerization at room temperature (Beeckman and Viane, 2000). Microscopic analysis was performed on 5 µm cross-sections cut using a rotary microtome (RM 2265; Leica) and placed on poly-L-lysine-coated glass slides (Sigma). The slides were dried at 37 °C for 2 hours on a heating plate, and stained with toluidine blue O (0.05 %). Finally, the sections were examined with motorized epi-fluorescence microscope (Olympus BX 61) using cellP software (Olympus).

.

Shoot apical meristem imaging and transient expression in protoplasts

Cut shoot apical meristems from *rsgA-i* line and wild type were transferred to a fresh 2 % agarose in a small round petri dish and used for imaging. 0.1 % Propidium iodide (PI) solution was used to stain cell walls. To image the meristem, a 40X water-dipping lens was used. The stack of images was obtained with a resolution of 0.5µm along the Z-axis. 561nm laser was used to excite PI and emission wavelengths between 605 to 630nm were recorded on a Leica SP8 confocal microscope. Arabidopsis protoplast were isolated and transformed with 10 µg of plasmid (Wu *et al.*, 2009), and observed with Leica DM6000B/SP5 confocal laser scanning microscope (Leica Microsystems). The GFP was excited with 488 nm wavelength and emission of GFP was collected between 500 and 520 nm.

Pigment content estimation

For chlorophyll extraction, leaf discs from 6 weeks old plants were cut and crushed in 80 % acetone solution followed by 12-16 hour incubation at room temperature. The readings were taken in triplicates at 470 nm, 645 nm and 662 nm using ELISA plate reader (Molecular devices, Versamax microplate reader). Chlorophyll a, chlorophyll b and xanthophyll + carotenoid amounts were calculated as in (Fan, 1997).

RNA-Seq analysis

Samples were processed using LSTrAP (Proost *et al.*, 2017), using default settings. All samples passed LSTrAP's quality control. Log_2 fold changes, *p*-values and corrected *p*-values (BH) for genes differentially expressed when comparing mutants with WT plants were calculated using DESeq2 (package version 1.10.1)(Love *et al.*, 2014). Significantly differentially expressed genes ($P < 0.05$ after correction) were selected and used for a MapMan pathway analysis (MapMan v 3.6.0RC1)(Thimm *et al.*, 2004). MapMan bins significantly enriched for genes ($P < 0.05$ after Benjamini and Hochberg (BH) correction) in this subset were determined and stored (Table S7).

Proteomic analysis

50 µg of total protein extract was digest using trypsin. Peptides were re-suspended in 40 µl 5 % v/v acetonitrile, 2 % v/v trifluoroacetic acid. Measurements were performed on a Q Exactive HF coupled with an Easy nLC1000 HPLC (Thermo Scientific). Samples were loaded onto Acclaim PepMap RSLC reversed-phase column (75 µm inner diameter, 25 cm long Thermo Scientific) at flow 0.4 µl min^{-1} in 3 % v/v acetonitrile, 0.5 % v/v acetic acid, and eluted with a 3 % to 30 % v/v acetonitrile concentration gradient over 200 min, 30 % to 40 % in 20 min and from 40 % to 60 % in 20 min and followed by a washing step with 90 % v/v acetonitrile for 5 min, at 0.3 µl/min. Peptide ions were detected in a full scan from mass-to-charge ratio 200 to 2000. MS/MS scans were performed for the 15 peptides with the highest MS signal. Quantitative analysis of MS/MS measurements was performed with the Progenesis QI software (Nonlinear Dynamics). Proteins were identified from spectra using Mascot (Matrix Science). A t-test was performed for each peptide and the resulting p-values were corrected for multiple testing using FDR (BH). For the MapMan plots, only significantly changed P-values (corrected $P < 0.05$) were included in the plots (Table S8).

Acknowledgments:

We would like to thank Profs. Staffan Persson, Mark Stitt and Ralph Bock for the

helpful discussions and feedback. We would like to thank the Max Planck Society for

funding. The authors declare no conflict of interest.

Supplemental figures

Figure S1. Multiple sequence alignment of RsgA protein sequence.

Figure S2. Bioanalyzer analysis of rRNA from *E.coli.*

Figure S3. Rosette growth dynamics.

Figure S4. Crosses of *rsgA-e* and *rsgA-i.*

Figure S5. rRNA Electropherogram of *rsgA-i* and wild type.

Figure S6. Northern blot against mitochondrial, cytosolic and chloroplast rRNAs.

Figure S7. Phenotypes of mutant and WT lines at different growth temperatures.

Figure S8. Differential expression analysis of *rsgA-i* and WT.

Figure S9. qRT-PCR and western blot analysis of the selected genes.

Figure S10. Chloroplast retrograde signalling responses.

Supplemental tables

Table S1. Number and phenotypes of seeds in WT and the mutant lines.

Table S2. Phenotypic segregation of the *rsgA-e/rsgA-i* cross.

Table S3. Transcriptomic analysis of rsgA-i and WT.

Table S4. Proteomic analysis of rsgA-i and WT.

Table S5. Primer sequences.

Table S6. Probes used for Northern blot analyses.

Table S7. MapMan term enrichment analysis of the transcriptome.

Table S8. MapMan term enrichment analysis of the proteome.

References:

Adam, Z. and Ostersetzer, O. (2001) Degradation of unassembled and damaged

thylakoid proteins. *Biochem. Soc. Trans.*, **29**, 427–30. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/11498002.

Albrecht, V., Ingenfeld, A. and Apel, K. (2006) Characterization of the snowy
cotyledon 1 mutant of Arabidopsis thaliana: The impact of chloroplast elongation
factor G on chloroplast development and plant vitality. *Plant Mol. Biol.*, **60**, 507–
518.

Alkatib, S., Fleischmann, T.T., Scharff, L.B. and Bock, R. (2012) Evolutionary
constraints on the plastid tRNA set decoding methionine and isoleucine. *Nucleic*
Acids Res., **40**, 6713–6724.

Avendano-Vazquez, A.-O., Cordoba, E., Llamas, E., et al. (2014) An
Uncharacterized Apocarotenoid-Derived Signal Generated in -Carotene
Desaturase Mutants Regulates Leaf Development and the Expression of
Chloroplast and Nuclear Genes in Arabidopsis. *Plant Cell*, **26**, 2524–2537.
Available at: http://www.plantcell.org/cgi/doi/10.1105/tpc.114.123349.

Barkan, A. (1993) Nuclear mutants of maize with defects in chloroplast polysome
assembly have altered chloroplast RNA metabolism. *Plant Cell*, **5**, 389–402.

Beeckman, T. and Viane, R. (2000) Embedding thin plant specimens for oriented
sectioning.   *Biotech.   Histochem.*,   **75**,   23–6.   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/10810979.

Beick, S., Schmitz-Linneweber, C., Williams-Carrier, R., Jensen, B. and Barkan,
A. (2008) The Pentatricopeptide Repeat Protein PPR5 Stabilizes a Specific
tRNA Precursor in Maize Chloroplasts. *Mol. Cell. Biol.*, **28**, 5337–5347.
Available at: http://mcb.asm.org/cgi/doi/10.1128/MCB.00563-08.

Britton, R.A. (2009) Role of GTPases in Bacterial Ribosome Assembly. *Annu. Rev.*
Microbiol.,   **63**,   155–176.   Available   at:
http://www.annualreviews.org/doi/10.1146/annurev.micro.091208.073225.

Bubunenko, M.G. and Subramanian, A.R. (1994) Recognition of novel and
divergent higher plant chloroplast ribosomal proteins by Escherichia coli
ribosome during in vivo assembly. *J. Biol. Chem.*, **269**, 18223–18231.

Campbell, T.L., Daigle, D.M. and Brown, E.D. (2005) Characterization of the
Bacillus subtilis GTPase YloQ and its role in ribosome function. *Biochem. J.*,
389, 843–52. Available at: http://www.biochemj.org/content/389/3/843.abstract.

Chan, K.X., Phua, S.Y., Crisp, P., McQuinn, R. and Pogson, B.J. (2016) Learning
the Languages of the Chloroplast: Retrograde Signaling and Beyond. *Annu.*
Rev.   Plant   Biol.,   **67**,   25–53.   Available   at:
http://www.annualreviews.org/doi/10.1146/annurev-arplant-043015-111854.

Charollais, J., Dreyfus, M. and Iost, I. (2004) CsdA, a cold-shock RNA helicase
from Escherichia coli, is involved in the biogenesis of 50S ribosomal subunit.
Nucleic Acids Res., **32**, 2751–2759.

Charollais, J., Pflieger, D., Vinh, J., Dreyfus, M. and Iost, I. (2003) The DEAD-box
RNA helicase SrmB is involved in the assembly of 50S ribosomal subunits in
Escherichia coli. *Mol. Microbiol.*, **48**, 1253–1265.

Connolly, K. and Culver, G. (2013) Overexpression of RbfA in the absence of the
KsgA checkpoint results in impaired translation initiation. *Mol. Microbiol.*, **87**,
968–981.

Connolly, K., Rife, J.P. and Culver, G. (2008) Mechanistic insight into the ribosome
biogenesis functions of the ancient protein KsgA. *Mol. Microbiol.*, **70**, 1062–
1075.

Dabbs, E.R. (1991) Mutants lacking individual ribosomal proteins as a tool to
investigate ribosomal properties. *Biochimie*, **73**, 639–645.

Datta, P.P., Wilson, D.N., Kawazoe, M., et al. (2007) Structural Aspects of RbfA
Action during Small Ribosomal Subunit Assembly. *Mol. Cell*, **28**, 434–445.

Davies, B.W., Köhrer, C., Jacob, A.I., Simmons, L.A., Zhu, J., Aleman, L.M.,
RajBhandary, U.L. and Walker, G.C. (2010) Role of escherichia coli YbeY, a
highly conserved protein, in rRNA processing. *Mol. Microbiol.*, **78**, 506–518.

Desveaux, D., Subramaniam, R., Després, C., Mess, J.N., Lévesque, C., Fobert,
P.R., Dangl, J.L. and Brisson, N. (2004) A "Whirly" transcription factor is
required for salicylic acid-dependent disease resistance in Arabidopsis. *Dev.*
Cell, **6**, 229–240.

Ehrnthaler, M., Scharff, L.B., Fleischmann, T.T., Hasse, C., Ruf, S. and Bock, R.
(2014) Synthetic lethality in the tobacco plastid ribosome and its rescue at
elevated growth temperatures. *Plant Cell*, **26**, 765–76. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/24563204%5Cnhttp://www.pubmedcentral.

nih.gov/articlerender.fcgi?artid=PMC3967039.

Elias, M. and Novotny, M. (2008) cpRAS: a novel circularly permuted RAS-like

GTPase domain with a highly scattered phylogenetic distribution. *Biol. Direct*, **3**,

21.   Available   at:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2430557&tool=pmcen

trez&rendertype=abstract.

Estavillo, G.M., Crisp, P.A., Pornsiriwong, W., et al. (2011) Evidence for a SAL1-

PAP Chloroplast Retrograde Pathway That Functions in Drought and High Light

Signaling in *Arabidopsis*. *Plant Cell*, **23**, 3992–4012. Available at:

http://www.plantcell.org/lookup/doi/10.1105/tpc.111.091033.

Fan, L. (1997) Antisense Suppression of Phospholipase D[alpha] Retards Abscisic

Acid[mdash] and Ethylene-Promoted Senescence of Postharvest Arabidopsis

Leaves. *PLANT CELL ONLINE*, **9**, 2183–2196. Available at:

http://www.plantcell.org/cgi/doi/10.1105/tpc.9.12.2183.

Fleischmann, T.T., Scharff, L.B., Alkatib, S., Hasdorf, S., Schöttler, M.A. and

Bock, R. (2011) Nonessential plastid-encoded ribosomal proteins in tobacco: a

developmental role for plastid translation and implications for reductive genome

evolution. *Plant Cell*, **23**, 3137–55. Available at:

http://www.ncbi.nlm.nih.gov/pubmed/21934145%5Cnhttp://www.pubmedcentral.

nih.gov/articlerender.fcgi?artid=PMC3203423.

Flores-Pérez, U., Sauret-Güeto, S., Gas, E., et al. (2008) A mutant impaired in the

production of plastome-encoded proteins uncovers a mechanism for the

homeostasis of isoprenoid biosynthetic enzymes in Arabidopsis plastids. *Plant*

Cell, **20**, 1303–15. Available at:

http://www.plantcell.org/cgi/doi/10.1105/tpc.108.058768%5Cnhttp://www.pubme

dcentral.nih.gov/articlerender.fcgi?artid=2438453&tool=pmcentrez&rendertype=

abstract%5Cnhttp://www.plantcell.org/content/20/5/1303.long.

Fournier, G.P., Neumann, J.E. and Gogarten, J.P. (2010) Inferring the ancient

history of the translation machinery and genetic code via recapitulation of

ribosomal subunit assembly orders. *PLoS One*, **5**.

Fristedt, R., Scharff, L.B., Clarke, C.A., Wang, Q., Lin, C., Merchant, S.S. and
Bock, R. (2014) RBF1, a Plant Homolog of the Bacterial Ribosome-Binding
Factor RbfA, Acts in Processing of the Chloroplast 16S Ribosomal RNA. *PLANT*
Physiol.,   **164**,   201–215.   Available   at:
http://www.plantphysiol.org/cgi/doi/10.1104/pp.113.228338.

Fromont-Racine, M., Senger, B., Saveanu, C. and Fasiolo, F. (2003) Ribosome
assembly in eukaryotes. *Gene*, **313**, 17–42.

Gonzalez-Perez, S., Gutierrez, J., Garcia-Garcia, F., Osuna, D., Dopazo, J.,
Lorenzo, O., Revuelta, J.L. and Arellano, J.B. (2011) Early Transcriptional
Defense Responses in Arabidopsis Cell Suspension Culture under High-Light
Conditions. *PLANT Physiol.*, **156**, 1439–1456. Available at:
http://www.plantphysiol.org/cgi/doi/10.1104/pp.111.177766.

Goto, S., Kato, S., Kimura, T., Muto, A. and Himeno, H. (2011) RsgA releases
RbfA from 30S ribosome during a late stage of ribosome biosynthesis. *EMBO J.*,
30,   104–14.   Available   at:
http://www.ncbi.nlm.nih.gov/pubmed/21102555%5Cnhttp://www.pubmedcentral.
nih.gov/articlerender.fcgi?artid=PMC3020115.

Himeno, H., Hanawa-Suetsugu, K., Kimura, T., et al. (2004) A novel GTPase
activated by the small subunit of ribosome. *Nucleic Acids Res.*, **32**, 5303–5309.

Jacob, A.I., Köhrer, C., Davies, B.W., RajBhandary, U.L. and Walker, G.C. (2013)
Conserved Bacterial RNase YbeY Plays Key Roles in 70S Ribosome Quality
Control and 16S rRNA Maturation. *Mol. Cell*, **49**, 427–438.

Jomaa, A., Stewart, G., Martin-Benito, J., Zielke, R., Campbell, T.L., Maddock,
J.R., Brown, E.D. and Ortega, J. (2011) Understanding ribosome assembly:
the structure of in vivo assembled immature 30S subunits revealed by cryo-
electron microscopy. *RNA*, **17**, 697–709. Available at:
http://rnajournal.cshlp.org/cgi/doi/10.1261/rna.2509811.

Kaczanowska, M. and Rydén-Aulin, M. (2007) Ribosome biogenesis and the
translation process in Escherichia coli. *Microbiol. Mol. Biol. Rev.*, **71**, 477–494.

Kim, B.H., Malec, P., Waloszek, A. and Arnim, A.G. von (2012) Arabidopsis
BPG2: A phytochrome-regulated gene whose protein product binds to plastid

ribosomal RNAs. *Planta*, **236**, 677–690.

Kim, S., Schlicke, H., Ree, K. Van, Karvonen, K., Subramaniam, A., Richter, A., Grimm, B. and Braam, J. (2013) Arabidopsis Chlorophyll Biosynthesis: An Essential Balance between the Methylerythritol Phosphate and Tetrapyrrole Pathways. *Plant Cell Online*, **25**, 4984–4993. Available at: http://www.plantcell.org/content/25/12/4984.abstract%5Cnhttp://www.plantcell.org/content/25/12/4984.full.pdf.

Kimura, T., Takagi, K., Hirata, Y., Hase, Y., Muto, A. and Himeno, H. (2008) Ribosome-Small-Subunit-Dependent GTPase Interacts with tRNA-Binding Sites on the Ribosome. *J. Mol. Biol.*, **381**, 467–477.

Kindgren, P., Eriksson, M.-J., Benedict, C., Mohapatra, A., Gough, S.P., Hansson, M., Kieselbach, T. and Strand, Å. (2011) A novel proteomic approach reveals a role for Mg-protoporphyrin IX in response to oxidative stress. *Physiol. Plant.*, **141**, 310–320. Available at: http://doi.wiley.com/10.1111/j.1399-3054.2010.01440.x.

Komatsu, T., Kawaide, H., Saito, C., et al. (2010) The chloroplast protein BPG2 functions in brassinosteroid-mediated post-transcriptional accumulation of chloroplast rRNA. *Plant J.*, **61**, 409–422.

Leipe, D.D., Wolf, Y.I., Koonin, E. V and Aravind, L. (2002) Classification and evolution of P-loop GTPases and related ATPases. *J. Mol. Biol.*, **317**, 41–72. Available at: http://linkinghub.elsevier.com/retrieve/pii/S0022283601953781.

Liere, K. and Börner, T. (2007) Transcription of Plastid Genes. In *Regulation of Transcription in Plants*. pp. 184–224.

Liu, H., Lau, E., Lam, M.P.Y., et al. (2010) OsNOA1/RIF1 is a functional homolog of AtNOA1/RIF1: Implication for a highly conserved plant cGTPase essential for chloroplast function. *New Phytol.*, **187**, 83–105.

Liu, J., Zhou, W., Liu, G., et al. (2015) The Conserved Endoribonuclease YbeY Is Required for Chloroplast Ribosomal RNA Processing in Arabidopsis. *Plant Physiol.*, **168**, 205–221. Available at: http://www.plantphysiol.org/lookup/doi/10.1104/pp.114.255000.

Livak, K.J. and Schmittgen, T.D. (2001) Analysis of Relative Gene Expression
Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods*, **25**,
402–408. Available at:
http://linkinghub.elsevier.com/retrieve/pii/S1046202301912629.

Lopez-Juez, E. and Pyke, K.A. (2005) Plastids unleashed: Their development and
their integration in plant development. *Int. J. Dev. Biol.*, **49**, 557–577.

Love, M.I., Anders, S. and Huber, W. (2014) *Differential analysis of count data - the*
DESeq2 package, Available at:
http://biorxiv.org/lookup/doi/10.1101/002832%5Cnhttp://dx.doi.org/10.1186/s130
59-014-0550-8.

Maekawa, S., Takabayashi, A., Huarancca Reyes, T., Yamamoto, H., Tanaka, A.,
Sato, T. and Yamaguchi, J. (2015) Pale-green phenotype of atl31atl6 double
mutant leaves is caused by disruption of 5-aminolevulinic acid biosynthesis in
Arabidopsis thaliana. *PLoS One*, **10**, e0117662. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/25706562%5Cnhttp://www.pubmedcentral.
nih.gov/articlerender.fcgi?artid=PMC4338271.

Meurer, J., Lezhneva, L., Amann, K., Godel, M., Bezhani, S., Sherameti, I. and
Oelmuller, R. (2002) A peptide chain release factor 2 affects the stability of
UGA-containing transcripts in Arabidopsis chloroplasts. *Plant Cell*, **14**, 3255–
3269. Available at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt
=Citation&list_uids=12468741.

Mochizuki, N., Tanaka, R., Grimm, B., Masuda, T., Moulin, M., Smith, A.G.,
Tanaka, A. and Terry, M.J. (2010) The cell biology of tetrapyrroles: A life and
death struggle. *Trends Plant Sci.*, **15**, 488–498.

Motohashi, R., Yamazaki, T., Myouga, F., et al. (2007) Chloroplast ribosome
release factor 1 (AtcpRF1) is essential for chloroplast development. *Plant Mol.*
Biol., **64**, 481–497.

Murashige, T. and Skoog, F. (1962) A Revised Medium for Rapid Growth and Bio
Assays with Tobacco Tissue Cultures. *Physiol. Plant.*, **15**, 473–497.

Nierhaus, K.H. and Dohme, F. (1974) Total reconstitution of functionally active 50S

ribosomal subunits from Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.*, **71**,
4713–7.   Available   at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=433966&tool=pmcent
rez&rendertype=abstract.

Okazaki, K., Kabeya, Y., Suzuki, K., Mori, T., Ichikawa, T., Matsui, M., Nakanishi,
H. and Miyagishima, S. -y. (2009) The PLASTID DIVISION1 and 2
Components of the Chloroplast Division Machinery Determine the Rate of
Chloroplast Division in Land Plant Cell Differentiation. *PLANT CELL ONLINE*,
21,   1769–1780.   Available   at:
http://www.plantcell.org/cgi/doi/10.1105/tpc.109.067785.

Park, J., Nguyen, K.T., Park, E., Jeon, J.-S. and Choi, G. (2013) DELLA Proteins
and Their Interacting RING Finger Proteins Repress Gibberellin Responses by
Binding to the Promoters of a Subset of Gibberellin-Responsive Genes in
Arabidopsis. *Plant Cell*, **25**, 927–943. Available at:
http://www.plantcell.org/cgi/doi/10.1105/tpc.112.108951.

Possingham, J. V. and Lawrence, M.E. (1983) Controls to Plastid Division. *Int.*
Rev. Cytol., **84**, 1–56.

Proost, S., Krawczyk, A. and Mutwil, M. (2017) LSTrAP: Efficiently combining RNA
sequencing data into co-expression networks. *BMC Bioinformatics*, **18**.

Pyke, K.A. (1997) The genetic control of plastid division in higher plants. *Am. J. Bot.*,
84, 1017–1027.

Ramel, F., Birtic, S., Ginies, C., Soubigou-Taconnat, L., Triantaphylides, C. and
Havaux, M. (2012) Carotenoid oxidation products are stress signals that
mediate gene responses to singlet oxygen in plants. *Proc. Natl. Acad. Sci.*, **109**,
5535–5540.   Available   at:
http://www.pnas.org/cgi/doi/10.1073/pnas.1115982109.

Rogalski, M., Ruf, S. and Bock, R. (2006) Tobacco plastid ribosomal protein S18 is
essential for cell survival. *Nucleic Acids Res.*, **34**, 4537–4545.

Ruiz-Sola, M.Á. and Rodríguez-Concepción, M. (2012) Carotenoid Biosynthesis in
Arabidopsis: A Colorful Pathway. *Arab. B.*, **10**, e0158. Available at:
http://www.bioone.org/doi/abs/10.1199/tab.0158.

Sato, S. (1999) Complete Structure of the Chloroplast Genome of Arabidopsis
thaliana. *DNA Res.*, **6**, 283–290. Available at:
https://academic.oup.com/dnaresearch/article-
lookup/doi/10.1093/dnares/6.5.283.

Shajani, Z., Sykes, M.T. and Williamson, J.R. (2011) Assembly of Bacterial
Ribosomes. *Annu. Rev. Biochem.*, **80**, 501–526. Available at:
http://www.annualreviews.org/doi/10.1146/annurev-biochem-062608-160432.

Shen, Y., Li, C., McCarty, D.R., Meeley, R. and Tan, B.C. (2013) Embryo
defective12 encodes the plastid initiation factor 3 and is essential for
embryogenesis in maize. *Plant J.*, **74**, 792–804.

**Sugliani, M., Abdelkefi, H., Ke, H., Bouveret, E., Robaglia, C., Caffarri, S. and
Field, B.** (2016) An Ancient Bacterial Signaling Pathway Regulates Chloroplast
Function to Influence Growth and Development in Arabidopsis. *Plant Cell*, **28**,
661–679. Available at: http://www.plantcell.org/lookup/doi/10.1105/tpc.16.00045.

Talkington, M.W.T., Siuzdak, G. and Williamson, J.R. (2005) An assembly
landscape for the 30S ribosomal subunit. *Nature*, **438**, 628–632. Available at:
http://www.nature.com/doifinder/10.1038/nature04261.

Thimm, O., Bläsing, O., Gibon, Y., et al. (2004) MAPMAN: A user-driven tool to
display genomics data sets onto diagrams of metabolic pathways and other
biological processes. *Plant J.*, **37**, 914–939.

**Thurlow, B., Davis, J.H., Leong, V., Moraes, T.F., Williamson, J.R. and Ortega,
J.** (2016) Binding properties of YjeQ (RsgA), RbfA, RimM and Era to assembly
intermediates of the 30S subunit. *Nucleic Acids Res.*, **44**, 9918–9932.

Tiller, N. and Bock, R. (2014) The translational apparatus of plastids and its role in
plant development. *Mol. Plant*, **7**, 1105–1120.

**Tiller, N., Weingartner, M., Thiele, W., Maximova, E., Schöttler, M.A. and Bock,
R.** (2012) The plastid-specific ribosomal proteins of Arabidopsis thaliana can be
divided into non-essential proteins and genuine ribosomal proteins. *Plant J.*, **69**,
302–316.

Timmis, J.N., Ayliffe, M.A., Huang, C.Y. and Martin, W. (2004) Endosymbiotic

gene transfer: organelle genomes forge eukaryotic chromosomes. *Nat. Rev.*
Genet., **5**, 123–135. Available at:
http://www.nature.com/doifinder/10.1038/nrg1271.

Traub, P. and Nomura, M. (1969) Structure and function of Escherichia coli
ribosomes: VI. Mechanism of assembly of 30 s ribosomes studied in vitro. *J.*
Mol. Biol., **40**, 391–413.

Tzafrir, I., Pena-Muralla, R., Dickerman, A., et al. (2004) Identification of genes
required for embryo development in Arabidopsis. *Plant Physiol.*, **135**, 1206–20.
Available at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=519041&tool=pmcent
rez&rendertype=abstract.

Vogel, M.O., Moore, M., Konig, K., Pecher, P., Alsharafa, K., Lee, J. and Dietz,
K.-J. (2014) Fast Retrograde Signaling in Response to High Light Involves
Metabolite Export, MITOGEN-ACTIVATED PROTEIN KINASE6, and AP2/ERF
Transcription Factors in Arabidopsis. *Plant Cell*, **26**, 1151–1165. Available at:
http://www.plantcell.org/cgi/doi/10.1105/tpc.113.121061.

Waters, M.T., Wang, P., Korkaric, M., Capper, R.G., Saunders, N.J. and
Langdale, J.A. (2009) GLK Transcription Factors Coordinate Expression of the
Photosynthetic Apparatus in Arabidopsis. *PLANT CELL ONLINE*, **21**, 1109–
1128. Available at: http://www.plantcell.org/cgi/doi/10.1105/tpc.108.065250.

Weglohner, W., Junemann, R., Knoblauch, K. von and Subramanian, A.R.
(1997) Different consequences of incorporating chloroplast ribosomal proteins
L12 and S18 into the bacterial ribosomes of Escherichia coli. *Eur. J. Biochem.*,
249, 383–392.

Woodson, J.D., Perez-Ruiz, J.M. and Chory, J. (2011) Heme synthesis by plastid
ferrochelatase i regulates nuclear gene expression in plants. *Curr. Biol.*, **21**,
897–903.

Woodson, S.A. (2008) RNA folding and ribosome assembly. *Curr. Opin. Chem.*
Biol., **12**, 667–673.

Wu, F.-H., Shen, S.-C., Lee, L.-Y., Lee, S.-H., Chan, M.-T. and Lin, C.-S. (2009)
Tape-Arabidopsis Sandwich - a simpler Arabidopsis protoplast isolation method.

Plant Methods, **5**, 16. Available at:
http://plantmethods.biomedcentral.com/articles/10.1186/1746-4811-5-16.

Xia, B., Ke, H., Shinde, U. and Inouye, M. (2003) The role of RbfA in 16S rRNA
processing and cell growth at low temperature in Escherichia coli. *J Mol Biol*,
332, 575–584. Available at:
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt
=Citation&list_uids=12963368%5Cnhttp://www.ncbi.nlm.nih.gov/pubmed/12963
368.

Xiao, Y., Savchenko, T., Baidoo, E.E.K., et al. (2012) Retrograde signaling by the
plastidial metabolite MEcPP regulates expression of nuclear stress-response
genes. *Cell*, **149**, 1525–1535.

Yamaguchi, K. and Subramanian, A.R. (2000) The plastid ribosomal proteins.
Identification of all the proteins in the 50 S subunit of an organelle ribosome
(chloroplast). *J. Biol. Chem.*, **275**, 28466–28482.

Zones, J.M., Blaby, I.K., Merchant, S.S. and Umen, J.G. (2015) *High-Resolution*
Profiling of a Synchronized Diurnal Transcriptome from Chlamydomonas
reinhardtii Reveals Continuous Cell and Metabolic Differentiation, Available at:
http://www.plantcell.org/content/early/2015/10/02/tpc.15.00498.abstract?papeto
c.

Figures:

Figure 1. AtRsgA from *Arabidopsis thaliana* partially complements *ΔrsgA* deletion strain from *E.coli*. A) Growth curve of wild type (WT), mutant (*ΔrsgA*), complemented (*ΔrsgA+AtRsgA*) and control (*ΔrsgA+empty vector, EV*) strains in liquid media. The error bars indicate standard deviation (n = 3). B) Growth of transformed and non-transformed cultures on solid LB medium. C) 16S/23S rRNA ratios (as calculated from bioanalyzer electropherogram) of transformed and non-transformed cultures grown in liquid cultures. The asterisks indicate ratios significantly different from the complemented strain (*ΔrsgA+AtRsgA*, P < 0.05).

Figure 2. Phenotypes of At*RsgA* T-DNA insertion lines in *Arabidopsis thaliana*. A) Exon-intron structure of *AtRsgA*, together with the position of the two T-DNA lines. Untranslated regions, exons and introns are indicated by white boxes, gray boxes and lines, respectively. B) *AtRsgA* expression analysis in heterozygous *rsgA-e* and homozygous *rsgA-i* lines. The expression of *AtRsgA* in *rsgA-e* and *rsgA-i* lines was normalized to its expression in WT sample. Error bars represent standard deviations (n=3, asterisk = P < 0.01). C) Rosette phenotype after four weeks of growth (scale bar = 1 cm). D) Phenotypes of eight week old plants (scale bar = 5 cm). E) Opened siliques from WT, *rsgA-e* and *rsgA-i* (scale bar = 1 mm). F) Exposed immature seeds from WT, *rsgA-e* and *rsgA-I* (scale bar = 500 μm).

Figure 3. Northern blot analysis of *rsgA-i* and chloroplast 30S assembly mutants. A) Physical map of the plastid rDNA operon and maturation pattern of the plastid rRNA. The 7.3-kb primary transcript, processing intermediates, hidden breaks (HB, black triangles) and the mature forms of the 16S and 23S rRNAs are shown. Positions of hybridization probes are indicated by green lines. B) Bioanalyzer estimation of ratio of peak area of chloroplast rRNAs. Error bars indicate standard deviation (n = 3), while asterisk show significant difference between WT and *rsgA-i* (P < 0.01). C) Close-up of 16S/18S and 23S-1/18S ratios as estimated in (B). D) Northern blot analysis of mature 16S and 5' and 3' termini of 17S rRNA in WT, *rsgA-i*, *rif1* and *bpg2*. As a loading control, methylene blue-stained nylon membranes are shown. The bands on methylene blue membrane correspond to (from highest nt

1017 size): 25S, 18S rRNA from cytosolic ribosome, plastid 16S and two HB fragments of
1018 plastid 23S rRNA.

1019

1020 **Figure 4. Polysome analysis of A) WT and B) *rsgA-i*.** 254nm absorbance spectra
1021 were used to indicate the presence of ribosomal subunits, monosomes and
1022 polysomes. The collected fractions were subjected to 16S rRNA northern blot
1023 analysis, while methylene blue stain was used as loading control. The 17S and 16S
1024 rRNAs are indicated. C) Ratio of 17S/16S rRNA in the collected fractions.

1025

Figure 5. Expression analysis of *AtRsgA* and comparison of photosynthetic
parameters of wild type and mutant lines. A) qRT-PCR of *AtRsgA* in different
tissues, normalized to the expression in roots. B) *AtRsgA* promoter-GUS activity in
one week old seedling showing high expression in photosynthetically active green
tissue. Scale bar = 5mm. C) Photosynthetic pigment content measurements in 2-
weeks old seedlings of WT, heterozygous *rsgA-e* and homozygous *rsgA-i* (n = 3,
asterisk = P < 0.05). D-F) Photosynthesis efficiency parameters of WT and *rsgA-i*
plants showing reduced photosynthetic capacity of *rsgA-i* plants. The measurements
were taken at 21 and 27 days after germination (DAG). D) Maximum quantum
efficiency of PSII. E) Light response curve of linear electron transport II (ETRII). F)
Light response curve of non-photochemical quenching (qN).

1037

Figure 6. Physiological analysis of *rsgA-i*. A) Mesophyll cells from first true leaves
of two weeks old WT and *rsgA-i* plants, fixed in 10 % formalin. Scale bar = 20 µm. B)
Comparison of chloroplast number, mesophyll cell size and chloroplast area per cell
area in WT (white bar) and *rsgA-i* (gray bar). Error bars represent standard deviation
(n = 50, asterisk = P<0.01). C) Leaf cross-section with chloroplast stained with
toluidine blue O (scale bar = 100 µm). D) Confocal image of shoot apical meristem
and leaf primordia. Red color corresponds to plastid autofluorescence. The L1, L2
and L3 layers and rib meristem (RB) are indicated (scale bar = 50 µm).

1046

Figure 7. Transcriptome and proteome analysis of wild type and *rsgA-i*. A)
Violin plots showing the log2-fold-change between *rsgA-i* and WT, for transcripts
(blue side of the violin plot) and proteins (yellow side of the violin plot) targeted to, or
encoded in the different compartments. The solid lines within the violin plot indicate

median, while dashed lines show quartiles. Values above 0 indicate expression

higher in *rsgA-i* than WT, while values lower than 0 indicate expression higher in WT

than *rsgA-i*. B) Distribution of significantly (P < 0.05) differentially upregulated (blue

bars), downregulated (red bars) or unchanged (white bars) transcripts (left bars) and

proteins (right bars) in the different compartments. C) The numbers indicate how

many proteins and transcripts are commonly up- or downregulated in *rsgA-i*, as

compared to WT. D) MapMan analysis of the differentially expressed transcripts and

proteins. Blue color indicates genes (cells in the heatmap) that are upregulated

(blue) or downregulated (red) in *rsgA-i*. E) Biochemical pathways of the different

metabolite retrograde signals. Known retrograde signals are shown in orange, solid

arrows indicate a single biochemical step, and dash-dotted arrows are simplified

representations of more than one biochemical step. Transcriptomic and proteomic

responses are indicated by colored boxes, where blue, red, white and striked field

indicates up-regulated, down-regulated, unchanged and not detectable, respectively.

A

Normalized absorbance 254 nm

Ribosomal subunits
Monosomes

WT - 16S probe

Polysomes

16S

Methylene blue

Sucrose gradient

B

Normalized absorbance 254 nm

Ribosomal subunits
Monosomes

rsgA-i - 16S probe

Polysomes

17S
16S

Methylene blue

Sucrose gradient

C

17S/16S ratio

fractions