Metabolic retroconversion of trimethylamine *N*-oxide and the gut microbiota

Lesley Hoyles[1]*, Maria L. Jiménez-Pranteda[2]*, Julien Chilloux[1]*, Francois Brial[3], Antonis

Myridakis[1], Thomas Aranias[3], Christophe Magnan[4], Glenn R. Gibson[2], Jeremy D. Sanderson[5], Jeremy

K. Nicholson[1], Dominique Gauguier[1,3], Anne L. McCartney[2]† and Marc-Emmanuel Dumas[1]†

[1]Integrative Systems Medicine and Digestive Disease, Department of Surgery and Cancer, Faculty of

Medicine, Imperial College London, Exhibition Road, London SW7 2AZ, UK

[2]Food Microbial Sciences Unit, Department of Food and Nutritional Sciences, School of Chemistry,

Food and Pharmacy, Faculty of Life Sciences, The University of Reading, Whiteknights Campus,

Reading RG6 6UR, UK

[3]Sorbonne Universities, University Pierre & Marie Curie, University Paris Descartes, Sorbonne Paris

Cité, INSERM UMR_S 1138, Cordeliers Research Centre, Paris, France

[4]Sorbonne Paris Cité, Université Denis Diderot, Unité de Biologie Fonctionnelle et Adaptative, CNRS

UMR 8251, 75205 Paris, France

[5]Department of Gastroenterology, Guy's and St Thomas' NHS Foundation Trust and King's College

London, London, UK

*These authors made equal contributions to this work; shared first authorship.

†Corresponding authors: Anne L. McCartney; Marc-Emmanuel Dumas

Keywords: co-metabolic axis; gut–liver axis; metabolomics; *Enterobacteriaceae*; lactic acid bacteria.

Email addresses: L. Hoyles, lesley.hoyles11@imperial.ac.uk; M.L. Jiménez-Pranteda, jimepra@ugr.es; J.

Chilloux, j.chilloux@imperial.ac.uk; F. Brial, francois.brial@crc.jussieu.fr; A. Myridakis,

a.myridakis@imperial.ac.uk; T. Aranias, thomas.aranias@free.fr; C. Magnan, christophe.magnan@univ-paris-

diderot.fr; G.R. Gibson, g.r.gibson@reading.ac.uk; J.D. Sanderson, jeremy.sanderson@kcl.ac.uk; J.K.

Nicholson, j.nicholson@imperial.ac.uk; D. Gauguier, dominique.gauguier@upmc.fr; A.L. McCartney,

a.l.mccartney@reading.ac.uk; M.-E. Dumas, m.dumas@imperial.ac.uk

ABSTRACT

BACKGROUND: The dietary methylamines choline, carnitine and phosphatidylcholine are used by

the gut microbiota to produce a range of metabolites, including trimethylamine (TMA). However,

little is known about the use of trimethylamine *N*-oxide (TMAO) by this consortium of microbes.

RESULTS: A feeding study using deuterated TMAO in C57BL6/J mice demonstrated microbial

conversion of TMAO to TMA, with uptake of TMA into the bloodstream and its conversion to

TMAO. Microbial activity necessary to convert TMAO to TMA was suppressed in antibiotic-treated

mice, with deuterated TMAO being taken up directly into the bloodstream. In batch-culture

fermentation systems inoculated with human faeces, growth of *Enterobacteriaceae* was stimulated in

the presence of TMAO. Human-derived faecal and caecal bacteria (n = 66 isolates) were screened on

solid and liquid media for their ability to use TMAO, with metabolites in spent media analysed by ^{1}H-

NMR. As with the *in vitro* fermentation experiments, TMAO stimulated the growth of

Enterobacteriaceae; these bacteria produced most TMA from TMAO. Caecal/small intestinal isolates

of *Escherichia coli* produced more TMA from TMAO than their faecal counterparts. Lactic acid

bacteria produced increased amounts of lactate when grown in the presence of TMAO, but did not

produce large amounts of TMA. Clostridia (*sensu stricto*), bifidobacteria and coriobacteria were

significantly correlated with TMA production in the mixed fermentation system but did not produce

notable quantities of TMA from TMAO in pure culture.

CONCLUSIONS: Reduction of TMAO by the gut microbiota (predominantly *Enterobacteriaceae*)

to TMA followed by host uptake of TMA into the bloodstream from the intestine and its conversion

back to TMAO by host hepatic enzymes is an example of metabolic retroconversion. TMAO

influences microbial metabolism depending on isolation source and taxon of gut bacterium.

Correlation of metabolomic and abundance data from mixed microbiota fermentation systems did not

give a true picture of which members of the gut microbiota were responsible for converting TMAO to

TMA; only by supplementing the study with pure culture work and additional metabolomics was it

possible to increase our understanding of TMAO bioconversions by the human gut microbiota.

BACKGROUND

Dietary methylamines such as choline, trimethylamine *N*-oxide (TMAO),

phosphatidylcholine (PC) and carnitine are present in a number of foodstuffs, including meat, fish,

nuts and eggs. It has long been known that gut bacteria are able to use choline in a fermentation-like

process, with trimethylamine (TMA), ethanol, acetate and ATP among the known main end-products

[1–3]. TMA can be used by members of the order *Methanomassiliicoccales* (*Archaea*) present in the

human gut to produce methane [4], or taken up by the host. Microbially produced TMA derived from

PC is absorbed in the small intestine [5]. TMA diffuses into the bloodstream from the intestine via the

hepatic vein to hepatocytes, where it is converted to trimethylamine *N*-oxide (TMAO) by hepatic

flavin-containing mono-oxygenases (FMOs) [6]. The bulk of TMAO, and lesser amounts of TMA,

derived from dietary methylamines can be detected in urine within 6 h of ingestion, and both

compounds are excreted in urine but not faeces [7,8]. TMAO can also be detected in human skeletal

muscle within 6 h of an oral dose of TMAO [8].

TMAO present in urine and plasma is considered a biomarker for non-alcoholic fatty liver

disease (NAFLD), insulin resistance and cardiovascular disease (CVD) [9–12]. Feeding TMAO to

high-fat-fed C57BL/6 mice exacerbates impaired glucose tolerance, though the effect on the gut

microbiota is unknown [13]. Low plasma PC and high urinary methylamines (including TMAO) were

observed in insulin-resistant mice on a high-fat diet, leading to the suggestion microbial methylamine

metabolism directly influences choline bioavailability; however, the microbiota of the mice was not

studied nor was labeled choline used to determine the metabolic fate of TMA/TMAO [9]. Choline

bioavailability is thought to contribute to hepatic steatosis in patients [11]. Spencer *et al.* [11]

manipulated dietary choline levels in 15 females for 2 weeks and monitored changes in the fecal

microbiota during the intervention. They found the level of choline supplementation affected

Gammaproteobacteria (inhibited by high levels of dietary choline, negatively correlated with changes

in liver fat/spleen fat ratios) and Erysipelotrichi (positively correlated with changes in liver fat/spleen

fat ratios), suggesting baseline levels of these taxa may predict the susceptibility of an individual to

liver disease. However, the study was under-powered and requires replication with a much larger

cohort to fully examine the link between choline bioavailability and liver disease. High levels of

circulating TMAO are associated with CVD [12], potentially via a platelet-mediated mechanism [14].

Circulating TMAO has been suggested to play a role in protection from hyperammonemia in mice,

acting as an osmoprotectant, and from glutamate neurotoxicity in mice and primary cultures of

neurons [15,16]. Recently, chronic exposure to TMAO was shown to attenuate diet-associated

impaired glucose tolerance in mice, and reduce endoplasmic reticulum stress and adipogenesis in

adipocytes [10]. The relevance of these findings to human health remains to be determined. There is a

basal level of TMA and TMAO detected in human urine even in the absence of dietary

supplementation [17], suggesting use of (microbial and/or host) cell-derived choline or PC in the

intestinal tract by the gut microbiota.

While it is well established that choline [2,11,18], PC [2,12] and carnitine [19–22] are used

by the human and rodent gut microbiotas to produce TMA, little is known about the reduction of

TMAO (predominantly from fish) to TMA (or other compounds) by members of these consortia.

TMAO is found at concentrations of 20–120 mg per 100 g fish fillet [23]. Ingestion of fish by humans

leads to increased urinary excretion of dimethylamine (DMA) and TMA, from 5.6 to 24.1 and from

0.2 to 1.6 µmol/24 h/kg of body weight, respectively [24]. Individuals with trimethylaminuria (fish

odour syndrome), in which TMA is not detoxified to TMAO by hepatic FMOs, have been shown to

reduce 40–60 % of an oral dose of TMAO to TMA [25]. It was suggested the gut microbiota was

responsible for reducing TMAO in these individuals, and in those individuals without

trimethylaminuria the TMA was re-oxidized in the liver before being excreted in the urine. The

process was termed '*metabolic retroversion*' "*(i.e., a cycle of reductive followed by oxidative*

reactions to regenerate TMAO)" [25]. The reduction of TMAO to TMA is most commonly associated

with the gut microbiota of marine fish, with the TMA generated by these bacteria contributing to the

characteristic odour of rotting fish [26]. Members of the class *Gammaproteobacteria*, which includes

a number of food spoilage organisms, are known to reduce TMAO to TMA quantitatively and are

unable to reduce TMA further [26,27]. The conversion of TMAO to TMA by bacteria represents a

unique form of anaerobic respiration, in which TMAO reductase acts as a terminal electron acceptor

(from NADH or formate) for respiratory electron flow [27].

Mining of metagenomic data (*torA*) suggests *Proteobacteria* (particularly *Escherichia* and

Klebsiella spp.) are likely to contribute greatest to the production of TMA from TMAO in the human

gut via the TMAO reductase pathway, with *Actinobacteria* (*Eggerthellaceae*) becoming more

important under stress [28]. Other microbial genes are associated with production of TMA from

choline (*cutC*), glycine-betaine (*grdH*), L-carnitine (*cntA*) and γ-butyrobetaine (*cntA*) [4]. A search of

the NCBI nucleotide database with the phrase 'TMAO reductase' also suggests many other bacteria of

human intestinal origin (namely, *Salmonella*, *Helicobacter*, *Prevotella*, *Bacillus* and *Bacteroides* spp.)

should be able to reduce TMAO to TMA. However, this trait has not been examined *in vitro* for

isolates of intestinal origin.

Consequently, we carried out an *in vivo* study in mice to confirm use of TMAO by the mouse

gut microbiota and to allow us to examine metabolic retroconversion of TMAO in a murine model.

We then used an *in vitro* fermentation system to highlight the effect of TMAO on the human faecal

microbiota. Finally, we screened a panel of human faecal and caecal/small intestinal isolates to

determine which members of the human gut microbiota were able to reduce TMAO to TMA, and

whether their metabolism was affected by being grown in the presence of TMAO.

METHODS

Animal work

Groups of six-week-old C57BL6/J mice (Janvier Labs, Courtaboeuf, France) were received

and acclimated in specific pathogen-free (SPF) maintenance conditions. They were fed with a

standard chow diet (R04-40, Safe, Augy, France) and were either given free access to tap water or tap

water containing the antibiotic cocktail (0.5 g/L vancomycin hydrochloride, 1 g/L neomycin trisulfate,

1 g/L metronidazole, 1 g/L ampicillin sodium; all antibiotics were purchased from Sigma-Aldrich) for

14 days. Mice were given in the morning by gavage either a solution of d_9-TMAO at 1×10^{-4} M

(Cambridge Isotope Laboratories Inc., DLM-4779-0, UK) or saline and euthanized 6 h later by

cervical dislocation. Blood samples were collected by tail tipping every 2 h in Microvette® CB 300

Lithium Heparin (Sarstedt, Marnay, France). Plasma was separated by centrifugation (10 min, 5000 $\boldsymbol{g}$,

4 °C) and stored at -80 °C until analysed by ultra-performance liquid chromatography–tandem mass

spectrometry (UPLC–MS/MS).

UPLC–MS/MS determination of plasma TMA, d_9-TMA, TMAO and d_9-TMAO

UPLC–MS/MS was employed for the determination of TMA, d_9-TMA, TMAO and d_9-

TMAO. Samples (10 µL) were spiked with 10 µL Internal Standard solution (d_9-choline and $^{13}C_3/^{15}N$-

TMA in water; 1 mg/L, Sigma-Aldrich). Ethyl 2-bromoacetate solution (45 µL) (15 g/L ethyl 2-

bromoacetate, 1 % NH_4OH in acetonitrile; ChromaSolv grade, Sigma-Aldrich) was added and

derivatization of TMAs (TMA, d_9-TMA and $^{13}C_3/^{15}N$-TMA) to their ethoxy-analogues was completed

after 30 min at room temperature. Protein/lipid precipitation solution (935 µL) (94 % acetonitrile/5

%water/1 % formic acid; ChromaSolv grade, Sigma-Aldrich) was added, samples were centrifuged

(20 min, 20,000 $\boldsymbol{g}$, 4 °C) and were transferred to UPLC-autosampler vials. Sample injections (5 µL

loop) were performed with a Waters Acquity UPLC-Xevo TQ-S UPLC–MS/MS system equipped

with an Acquity BEH HILIC (2.1×100 mm, 1.7 µm) chromatographic column. An isocratic elution

was applied with 10 mM ammonium formate (Sigma-Aldrich) in 95:5 (v/v) acetonitrile:water for 7

min at 750 µL/min and 50 °C. Positive electrospray (ESI^+) was used as ionization source and mass

spectrometer parameters were set as follows: capillary, cone and sources voltages at -700, -18 and

50 V, respectively, desolvation temperature at 600 °C, desolvation/cone/nebuliser gases were high

purity nitrogen (BOC) at 1000 L/h, 150 L/h and 7 bar, respectively. Collision gas was high-purity

argon (BOC). Mass spectrometer was operated in multiple reaction monitoring mode. The monitored

transitions were the following: for TMA, +146 → +118/59 m/z (23/27 V); for d_9-TMA, +155 →

+127/68 m/z (21/23 V); for $^{13}C_3/^{15}N$-TMA, +150 → +63/122 m/z (27/22 V); for TMAO, +76 →

+59/58 m/z (12/13 V); for d_9-TMAO, +85 → +68/66 m/z (18/20 V); and for d_9-choline, +108 →

+60/45 m/z (20/22 V). The system was controlled by MassLynx software, also used for data

acquisition and analysis.

[1]H-NMR spectroscopy and data analysis

Medium samples were randomized and centrifuged (5 min, 16000 g). Aliquots (50 µL) of the supernatants were diluted in 550 µL D_2O containing 1 mM trimethylsilyl-(2,2,3,3-2H$_4$)-1-propionate. Samples were transferred to 5 mm NMR tubes and measured on a NMR spectrometer (Bruker) operating at 600.22 MHz [1]H frequency as described [29]. [1]H-NMR spectra were pre-processed and analysed as described [9] using the Statistical Recoupling of Variables-algorithm [30]. Structural assignment was performed as described [31], using in-house and publicly available databases.

In vitro fermentation systems

Freshly voided faecal samples were obtained from three healthy human volunteers (one male and two females; age range 29–31 years), none of whom had been prescribed antibiotics 6 months prior to the study, eaten fish 3 days prior to sample collection or had any history of gastrointestinal disease. The University of Reading's Research Ethics Committee (UREC) does not require that specific ethical review and approval be given by UREC for the collection of faecal samples from healthy human volunteers to inoculate *in vitro* fermentation systems. Samples were processed immediately by diluting them 1 in 10 (w/w) in pre-reduced minimal broth and homogenizing them in a filter bag in a stomacher (Stomacher 400 Lab System; Seward) for 2 min at 'high' speed. For each donor, a batch culture vessel containing 175 mL of pre-reduced minimal broth was inoculated with 20 mL of the faecal homogenate and 5 mL of sterile H_2O containing 2 g TMAO dihydrate (Sigma-Aldrich). Control vessels were inoculated with 20 mL of the faecal homogenate and 5 mL of sterile H_2O. The final working volume of each batch culture vessel was 200 mL. The pH of each vessel (pH 6.5) was controlled automatically by the addition of 2 M HCl or 2 M NaOH. pH controllers were supplied by Electrolab. The contents of each vessel were stirred constantly. An anaerobic environment was maintained by constantly sparging the vessels with O_2-free N_2. The temperature of each vessel was maintained at 37 °C by use of a circulating waterbath connected to each fermentation vessel. The experiment was run for 9 h, with samples taken at 0, 1, 2, 3, 4, 5, 6 and 9 h.

Fluorescence *in situ* hybridization (FISH) analysis

Aliquots (2× 375 μL) of sample were fixed in ice-cold 4 % paraformaldehyde for 4 h, washed

in sterile phosphate-buffered saline and stored for FISH analysis as described [32]. **Supplementary**

Table 1 gives details for probes used in this study. Probes were synthesized commercially (MWG-

Biotech) and labelled with the fluorescent dye cyanine 3 (Cy3; excitation λ, 514 nm; emission λ, 566

nm; fluorescence colour, orange–red). FISH was done as described [32].

Slides were viewed under a Nikon E400 Eclipse microscope. DAPI slides were visualized

with the aid of a DM400 filter; probe slides were visualized with the aid of a DM575 filter. Cells

(between 15 and 50 per field of view) were counted for 15 fields of view, and the numbers of bacteria

were determined by using the following equation:

$DF \times ACC \times 6{,}732.42 \times 50 \times DF_{sample}$

Where the DF (dilution factor) was calculated by taking into account the concentration of the

original sample (375 μL to 300 μL = 0.8×). ACC (average cell count) was determined by counting 15

fields of view and assumes that a normal distribution was observed for the counts. The figure

6,732.42 refers to the area of the well divided by the area of the field of view. DF_{sample} refers to the

dilution of sample used with a particular probe (e.g. 5× for t_0 Lab158 counts). The detection limit of

this method was 89,766 bacteria/mL of sample (= $\log_{10}$ 4.95).

Screening of bacteria for ability to reduce TMAO

Part of an in-house collection (University of Reading) of bacteria isolated from human

caecal/small intestinal and faecal samples (**Table 1**) was screened on minimal agar [g/L: glucose

(Fisher Scientific), 4; bacteriological peptone (Oxoid), 20; NaCl (Fisher Scientific), 5; neutral red

(Sigma-Aldrich), 0.03; agar technical no. 3 (Oxoid), 15; pH 7] with and without 1 % (w/v) TMAO

dihydrate (Sigma-Aldrich). The composition of the agar was based on [27], who used the medium

with crystal violet and bile salts to screen members of the *Enterobacteriaceae* for TMAO reductase

activity. Colonies of isolates able to ferment glucose were red, whereas those able to reduce TMAO

were white. Growth curves (OD_{600} measured hourly for 10–12 h, then at 24 h) were determined for

selected isolates grown anaerobically at 37 °C in anaerobic minimal broth [g/L: glucose, 4;

bacteriological peptone, 20; NaCl, 5; L-cysteine HCl (Sigma-Aldrich), 0.5; resazurin solution (0.25

mg/mL; Sigma-Aldrich), 4 mL; pH 7] with and without 1 % (w/v) TMAO dihydrate. Glucose was

substituted by raffinose (Sigma-Aldrich) [33] in the minimal broth when working with bifidobacteria

because of the poor growth of these bacteria on the glucose-based medium. pH and metabolite

profiles of culture medium were examined at the end of the growth experiment (i.e. at t_{24}).

Statistical analyses

Differences between metabolites produced by faecal and caecal/small intestinal isolates of

Escherichia coli and lactic acid bacteria in the presence and absence of TMAO were analysed using

Student's *t* test. FISH data from the *in vitro* fermentation systems were analysed using the

Kolmogorov–Smirnov test with statistical significance, after correction for multiple testing, taken at P

< 0.05. Because of the presence of ties in the metabolite data from *in vitro* fermentation systems, the

bootstrapped Kolmogorov–Smirnov test (10,000 replications) was used with these data. Data from *in*

vitro fermentation systems (FISH and NMR) were correlated using Spearman's rank correlation

(corrected for ties) with results corrected for multiple testing using the method of Benjamini and

Hochberg [34].

RESULTS

***In vivo* confirmation of metabolic retroconversion**

First, to confirm *in vivo* metabolic retroconversion (i.e. microbial conversion of TMAO to

TMA, followed by host conversion of TMA to TMAO), we administered isotopically-labelled d_9-

TMAO or saline to mice that had or had not been treated with a broad-spectrum antibiotic cocktail for

14 days to suppress the gut microbiota [12]. Reduction of d_9-TMAO to d_9-TMA was quantified in

urine by UPLC–MS/MS up to 6 h after d_9-TMAO gavage, together with unlabelled TMA and TMAO

potentially produced from dietary sources (**Fig. 1**).

In the absence of antibiotics, d_9-TMAO was converted to d_9-TMA within 2 h of gavage. This

conversion was dramatically reduced, leading to a significantly lower concentration of d_9-TMA and a

higher concentration of d_9-TMAO, when the gut microbiota was suppressed by antibiotics (**Fig. 1A–**

D). We noted that in the absence of antibiotics all animals excreted approximately three times higher

levels of urinary unlabelled TMA than TMAO (**Fig. 1E–H**), corresponding to constitutively low

FMO3 activity in mice [35]. Levels of unlabelled TMAO/TMA were not significantly different from

one another in the control and d_9-TMAO-fed animals, while antibiotic treatment significantly reduced

the amount of both TMAO and TMA (**Fig. 1E**, **F**). Bioavailability of unlabelled TMAO/TMA, as

assessed by area under the curve (AUC), was significantly reduced by antibiotic treatment in both

saline and d_9-TMAO-gavaged animals, with almost no TMA detected in either experimental group

(**Fig. 1G**, **H**).

Effect of TMAO on human gut bacteria within a mixed system

After *in vivo* validation of the role of the gut microbiota in metabolic retroconversion, we

analysed the effect of TMAO on the faecal microbiota in an anaerobic batch-culture fermentation

system. Fermenter vessels filled with the glucose-containing medium supplemented or not with 1 %

(w/v) TMAO were inoculated with faecal slurries from three healthy donors and monitored for 9 h.

With the exception of enhanced growth of the *Enterobacteriaceae* (probe Ent), the presence of

TMAO in the medium had no statistically significant effect on the growth of bacteria within the

fermentation systems at 9 h (**Fig. 2A**, **Supplementary Figure 1**).

Huge variability, as measured by ^{1}H-NMR, was observed in the amount of TMA and DMA

produced by gut bacteria in the TMAO-containing fermentation systems, with the concentrations of

both metabolites increasing steadily from 0 to 9 h and differing significantly ($P < 0.05$) from the

control systems (**Fig. 2B**). The amount of TMAO in the systems was seen to decrease at 8 h.

Correlation of metabolite and FISH data demonstrated *Clostridium* clusters I and II

(Chis150), *Enterobacteriaceae* (Ent), bifidobacteria (Bif164) and coriobacteriia (Ato291) were

associated with TMA, acetate, ethanol and lactate (**Fig. 2C**). The *Betaproteobacteria* (Bet42a) were

anti-correlated with TMA, acetate, ethanol and lactate, which is unsurprising given this was the only

group of bacteria whose representation decreased in the fermentation systems over the course of the

experiment (**Supplementary Figure 1**, **Fig. 2C**). The *Enterobacteriaceae* and clostridia were

positively correlated with DMA production. The lactic acid bacteria (Lab158) were not significantly

correlated with any of the metabolites in the mixed culture.

Growth of pure cultures of gut bacteria in the presence of TMAO

Using an in-house collection of bacteria, we initially screened isolates on a modified version

of the agar of Takahi & Ishimoto [27] as a rapid means of screening bacteria for TMAO reductase

activity, and thereby their ability to reduce TMAO to TMA. While members of the

Enterobacteriaceae produced expected results (i.e. white colonies when grown in the presence of

TMAO and glucose, rather than red colonies on the glucose only control), we observed a number of

unexpected outcomes depending on the species under study. For example, colonies of lactic acid

bacteria were larger (almost twice their usual size) on TMAO-containing agar than on the glucose

control but remained red in colour, suggesting they had not reduced TMAO to TMA at detectable

levels but TMAO was influencing their growth. The clostridia examined produced mixed results on

the control and TMAO-containing media (i.e. white colonies on both plates or on the control plate

only, larger colonies on the TMAO-containing medium but without a colour change of the medium).

To determine whether these isolates were converting TMAO to TMA but a low level, we

examined the growth of all isolates in liquid culture and metabolites in the spent medium using NMR.

The growth of the *Enterobacteriaceae* was most greatly affected by the presence of TMAO in the

medium, with a faster, longer-lasting exponential phase than for the same isolates grown in the

control medium (**Fig. 3A**). pH of the spent medium (after 24 h) when *Enterobacteriaceae* were grown

in the presence of TMAO increased from a mean of 4.7 ± 0.3 (for the control) to 7.6 ± 0.3 ($n = 20$)

(the change in pH is what causes the colonies to appear white on TMAO-containing agar). The

growth of lactic acid bacteria, including *Enterococcus* and *Streptococcus* (**Fig. 3A**) spp., was

enhanced in the presence of TMAO, but not to the same extent as seen for the *Enterobacteriaceae*.

There was no significant difference ($P = 0.27$, t test) in the pH of the spent medium for these bacteria

after 24 h (mean 4.67 ± 0.9 compared with 4.33 ± 0.27 for the control). The growth of members of

Clostridium cluster I (e.g. *Clostridium perfringens*, **Fig. 3A**) was not enhanced in the presence of

TMAO, though some of these bacteria changed the colour of both the control and TMAO-containing

plates yellow during their growth. The pH of the spent liquid medium confirmed this observation to

be due to the alkalinity of the media in the control and TMAO-containing media after 24 h incubation:

e.g. *Clostridium sporogenes* D1(9) (pH 6.25 compared with 6.76 in the control medium), *Clostridium*

paraputrificum L16-FAA6 (pH 6.45 vs 5.38) and *Clostridium perfringens* L20-BSM1 (pH 5.56 vs

4.64).

[1]H-NMR analysis of spent medium from the TMAO-containing and control samples

demonstrated that, as expected, the *Enterobacteriaceae* produced the greatest amount of TMA from

TMAO (mean 38.79 ± 11.08 mM compared with 0.03 ± 0.01 mM, $n = 20$) (**Fig. 3B**; **Supplementary**

Table 2). Members of the families *Peptostreptococcaceae* (3.72 mM, $n = 1$), *Clostridiaceae* (Cluster

I) (2.62 ± 1.83 mM, $n = 3$), *Porphyromonadaceae* (1.42 mM, $n = 1$), *Bacteroidaceae* (1.40 ± 0.31

mM, $n = 3$), *Enterococcaceae* (1.19 ± 0.05 mM, $n = 5$), *Erysipelotrichaceae* (0.94 mM, $n = 1$;

[*Clostridium*] *ramosum*), *Staphylococcaceae* (0.34 mM, $n = 1$), *Streptococcaceae* (0.30 ± 0.16 mM, n

$= 5$), *Lactobacillaceae* (0.17 ± 0.07 mM, $n = 2$), *Pseudomonadaceae* (0.12 mM, $n = 1$) and

Bifidobacteriaceae (0.13 ± 0.1 mM, $n = 17$) produced low levels of TMA from TMAO (**Fig. 3B**;

Supplementary Table 2). There was great variability in the ability of the bifidobacteria to produce

TMA from TMAO, with several isolates and [*Clostridium*] *innocuum*, *Actinomyces odontolyticus*,

Fusobacterium ulcerans and *Actinomyces viscosus* not producing TMA from TMAO (**Fig. 3B**;

Supplementary Table 2).

Differences in the metabolic capabilities of faecal and caecal *Escherichia coli* isolates

Comparison of the amounts of TMA, and co-metabolites, produced by the faecal ($n = 7$) and

caecal ($n = 9$) isolates of *Escherichia coli* demonstrated significantly higher amounts of TMA were

produced by the caecal isolates compared with the faecal isolates in the TMAO-containing medium

(**Fig. 3C**). *Escherichia coli* of caecal origin produced more TMA than faecal isolates of the same

bacterium or other enterobacteria (*Hafnia*, *Citrobacter* and *Klebsiella* spp.) (**Supplementary Table**

2). The faecal isolates produced more acetate and lactate than the caecal isolates when grown in the

control medium. Taken together, these results demonstrate the different metabolic capabilities of

isolates of *Escherichia coli* recovered from different regions of the human gut.

Lactic acid bacteria produce more lactate in the presence of TMAO

Differences were also seen in the amount of lactic acid produced by lactic acid bacteria in the presence and absence of TMAO (**Fig. 3B**, **D**). In raffinose-containing medium, bifidobacteria produced increased of amounts of lactate when grown in the presence of TMAO (the bifidobacteria grew poorly, if at all, in glucose-containing media). Unlike the bifidobacteria, the *Streptococcaceae* and *Enterococcaceae* grew well in the glucose-containing medium and produced over 25 mM lactic acid in the TMAO-containing samples compared with <5 mM in the control samples (**Fig. 3D**). To the best of our knowledge, this is the first time TMAO has been shown to influence the metabolism of gut bacteria – specifically lactic acid bacteria – without producing appreciable amounts of TMA.

DISCUSSION

TMAO is a circulating metabolite produced as a direct result of microbial degradation of dietary methylamines in the intestinal tract, and can be readily detected along with its precursor TMA in human urine, blood and skeletal muscle [8,12,18,22]. It worsens atherosclerosis in some mouse models of CVD, and is positively correlated with CVD severity in humans. Beneficial effects associated with TMAO include potential protection from hyperammonia and glutamate neurotoxicity, alleviation of endoplasmic reticulum stress and improved glucose homeostasis by stimulating insulin secretion by pancreatic cells [10,15,16].

Over many decades it has been established that choline, PC and carnitine are dietary methylamines that contribute directly to microbiome-associated circulating levels of TMAO found in humans and other animals – e.g. [2,12,18,19,21,22]. However, TMAO itself is a water-soluble osmolyte found in high abundance in fish. Based on their observations that individuals with trimethylaminuria could reduce an oral dose of TMAO to TMA could not detoxify it to TMAO, al-Waiz *et al.* [25] suggested the gut microbiota could use TMAO as a substrate, and in those individuals without trimethylaminuria the TMA was re-oxidized in the liver before being excreted in the urine. This led these authors to propose the process of '*metabolic retroversion*'. Reference to the literature

associated with fish spoilage and recent mining of metagenomic data have predicted members of the

Enterobacteriaceae (particularly *Escherichia coli* and *Klebsiella pneumoniae*) have the potential to

convert TMAO to TMA in the intestinal tract, but this has not been tested *in vitro* or *in vivo* to date

[4,8,26–28]. Consequently, we instigated this study to demonstrate metabolic retroconversion of

TMAO, and to determine the effect of TMAO on the growth and metabolism of human-derived

intestinal bacteria in pure and mixed cultures.

Through *in vivo* administration of deuterated TMAO to mice via oral gavage in mice, we

unambiguously demonstrated that TMAO is converted to TMA, with this TMA detectable in plasma

within 2 h of administration. This conversion was highly dependent on the gut microbiota, as

conversion of TMAO to TMA was dramatically reduced when the microbiota was suppressed by

treatment of animals with broad-spectrum antibiotics. Even in the presence of antibiotics, there was

low-level conversion of TMAO to TMA, suggesting a subpopulation of the microbiota was resistant

to the antibiotics used in our experiment. However, administration of a broad-spectrum antibiotic

cocktail for 14 days has been demonstrated to be an effective means of suppressing the gut microbiota

in studies associated with gut microbial use of dietary methylamines [12].

Our *in vivo* experiment clearly demonstrates the gut microbiota converts TMAO to TMA, and

that this TMA is re-oxidized to TMAO, in line with the process of '*metabolic retroversion*' [25]. Gut-

associated microbial conversion of the majority of TMAO to TMA is at odds with recent findings [8],

in which it was suggested TMAO is taken up intact and not metabolised by the gut microbiota of

humans. The ratio of TMAO-to-TMA is around 10:1 in humans and 1:10 in mice, meaning TMAO

was *N*-oxidized before Taesuwan *et al.* [8] were able to observe it in circulating blood of their human

subjects. In a human system with high FMO3 activity, ^{17}O-labelled TMAO would need to be used in

any study evaluating O turnover to allow calculation of the true rate of retroconversion.

Based on work done in mice [10,36], metabolic retroconversion of TMAO may be protective,

and may even go some way to explaining why TMA and TMAO are detected at low levels in urine in

the absence of dietary methylamines [17]. Chronic exposure of high-fat-fed mice to TMAO reduced

diet-associated endoplasmic stress and adipogenesis, and improved glucose tolerance [10]. Exposure

of high-fat-fed mice to TMA reduced low-grade inflammation and insulin resistance via inhibition of

interleukin-1 receptor-associated kinase 4 (IRAK-4) [36]. Concentrations of the microbial signalling

metabolite TMA used by Chilloux *et al.* [36] were comparable to those found in normal human

plasma. Continual exposure of the human system to TMA/TMAO may bring similar benefits to host

metabolic and inflammatory responses; these remain to be studied.

Having conducted our *in vivo* experiment in mice, we examined the ability of a range of

human-derived gut bacteria to convert TMAO to TMA. In the mixed microbiota system and in pure

cultures, the growth of the *Enterobacteriaceae* – the main TMA-producers – was quickly affected by

the presence of TMAO. This is likely to happen in the human gut also. Consequently, we believe

dietary TMAO undergoes metabolic retroconversion in mammals, with the TMA produced as a result

of bacterial activity in the gut available to the host for conversion back to TMAO by FMOs in

hepatocytes. It should be noted that [8] did not suppress or monitor the intestinal/faecal microbiota

when they administered isotopically labeled TMAO to humans, nor did they measure d_9-TMA and d_9-

TMAO in the portal vein, bypassing subsequent hepatic *N*-oxidation of d_9-TMA, so it is not possible

to interpret their results in the context of presence/absence of microbial activity.

Of note is the finding that caecal isolates of enterobacteria produce more TMA from TMAO

than faecal isolates of the same bacterium. The speed with which TMAO was reduced to TMA by the

Enterobacteriaceae in the present study suggests bacterial conversion of TMAO to TMA takes place

in the small intestine/proximal colon of humans and small intestine/caecum of mice. It is, therefore,

unsurprising that caecal bacteria – representing the microbiota present at the intersection of the small

and large intestine – are metabolically more active than their faecal counterparts with respect to

TMAO metabolism. This finding is relevant to functional studies of the gut microbiota where gene

annotations are based largely on faecal isolates, whose functionalities may be greatly different from

those of bacteria in other regions of the human intestinal tract [37,38]. It has already been

demonstrated that the microbiota of the small intestine is enriched for functions associated with rapid

uptake and fermentation of simple carbohydrates compared with the faecal microbiota, and that

streptococci isolated from this niche are functionally very different from the same bacteria isolated

from different habitats [37,38]. It is, therefore, important we characterize the functions and genomes

of bacteria isolated from all regions of the intestinal tract, not just those of faecal bacteria, to gain a

true picture of how microbial activity influences host health.

Enterobacteriaceae made the greatest contribution to the conversion of TMAO to TMA, both

in pure culture and in a mixed microbiota. These Gram-negative bacteria are a source of the virulence

factor lipopolysaccharide (LPS), which is associated with low-grade inflammation in high-fat-fed

mice and elevated plasma levels that define metabolic endotoxaemia [39,40]. High-fat feeding has

been shown to increase the representation of *Enterobacteriaceae* in the caecal microbiota of obesity-

prone Sprague–Dawley rats [41], though this mode of feeding is known to modulate the microbiota of

mice independent of obesity [42]. Non-LPS-associated virulence of *Enterobacteriaceae*, *Vibrio*

cholerae and *Helicobacter pylori* is increased when these bacteria are grown anaerobically or

microaerophilically in the presence of as little as 5–10 mM TMAO [43–46], and may be an additional

means by which the gut microbiota contributes to CVD and other diseases in which increased

representation of *Gammaproteobacteria* is observed. This warrants attention in future animal studies.

Lactic acid bacteria clearly grow better in the presence of TMAO. The relatively high

concentration (1 %) of TMAO used in this study may have contributed to this improved growth, as

TMAO is an osmolyte that stabilizes proteins. Future work will involve growing lactic acid bacteria in

a range of TMAO concentrations to determine how this compound affects their growth and gene

expression, and comparing faecal and caecal isolates. Similar to the *Enterobacteriaceae*, a large

number of these bacteria are facultative anaerobes able to grow over a range of conditions, and whose

representation is increased in obese, and cirrhotic patients [47,48]. *Streptococcus* and *Enterococcus*

spp. are commensal lactic acid bacteria of the gut microbiota known to modulate immune function;

but little is known about their metabolic activities in mixed microbial populations [49]. Understanding

how commensal lactic acid bacteria influence the host in dysbiosis in mixed microbial communities

may allow the development of approaches to modulate their activity and influence host health.

With respect to the lactic acid bacteria, it is important to note that our mixed culture work did

not highlight these as being relevant to TMAO metabolism. This is unsurprising given these bacteria

do not produce large quantities of TMA from TMAO. However, we have shown their metabolism is

affected by presence of TMAO in growth medium, and the increased lactate they produce in its

presence may contribute to cross-feeding associated with short-chain fatty acid production [50]. It is

difficult to determine relevance of correlations from mixed microbial ecosystems in the absence of

isotope labelling or pure culture work: i.e. correlation does not equate with causation. As an example,

three (*Clostridium* clusters I and II, bifidobacteria, coriobacteriia) of the four groups of bacteria

correlated with TMA production in our fermentation study did not produce notable quantities of TMA

from TMAO based on our pure culture work. Therefore, correlating microbiota and metabolite data

derived from complex systems will not give a true picture of which members of microbiota contribute

to specific metabolic processes, and work with pure cultures is required to supplement functional

studies to increase our understanding of poorly understood microbially driven metabolic processes

within the human gut.

CONCLUSIONS

We have demonstrated metabolic retroconversion – another example of host–microbial co-

metabolism – occurs in the mammalian system with respect to TMAO, whereby TMAO is reduced by

the gut microbiota to TMA and regenerated by host hepatic enzymes. We have also demonstrated that

growth and metabolism of members of the gut microbiota are affected by TMAO in a source- and

taxon-dependent manner, with the family *Enterobacteriaceae* making the greatest contribution to

production of TMA in the gut.

ABBREVIATIONS

AUC, area under the curve; DMA, dimethylamine; FISH, fluorescence *in situ* hybridization; FMO,

flavin mono-oxygenase; MMA, monomethylamine; PC, phosphatidylcholine; TMA, trimethylamine;

TMAO, trimethylamine *N*-oxide; UPLC–MS/MS, ultra-performance liquid chromatography–tandem

mass spectrometry.

DECLARATIONS

Ethics approval: All animal procedures were authorized following review by the institutional ethics

committee (Sorbonne Universities) and carried out under national license conditions. Ethical approval

to collect caecal effluent from patients was obtained from St Thomas' Hospital Research Ethics

Committee (06/Q0702/74) covering Guy's and St Thomas' Hospitals, and transferred by agreement to

London Bridge Hospital. Patients provided written consent to provide samples.

Consent for publication: Not applicable.

Availability of data and material: The datasets used and/or analysed during the current study are

available from the corresponding authors upon reasonable request.

Competing interests: The authors have no competing interests.

Funding: MLJP was sponsored by Fundación Alfonso Martin Escudero (Spain). Work was supported

by a grant from the EU-FP7 (METACARDIS, HEALTH-F4-2012-305312) to DG and MED. LH is in

receipt of an MRC Intermediate Research Fellowship in Data Science (grant number MR/L01632X/1,

UK Med-Bio).

Authors' contributions: MLJP, LH and ALM performed the microbiological work, JC was responsible

for NMR analysis and interpretation of the NMR data, TA and FB did the animal work, MED

supervised the NMR work, ALM supervised the microbiological work, CM provided invaluable

advice for the labelling study and DG supervised the animal work. All authors contributed to the

writing of the manuscript.

Acknowledgements: Not applicable.

REFERENCES

[1] Wünsche R. Uber die wirkung von darmbakterien auf trimethylammoniumbasen. Ber Ges
Physiol Exp Pharm. 1940;118:303.
[2] Zeisel SH, Wishnok JS, Blusztajn JK. Formation of methylamines from ingested choline and
lecithin. J Pharmacol Exp Ther. 1983;225:320-24.
[3] Bradbeer C. The clostridial fermentations of choline and ethanolamine. 1. Preparation and
properties of cell-free extracts. J Biol Chem. 1965;240:4669-74.

[4] Borrel G, McCann A, Deane J, Neto MC, Lynch DB, Brugère JF, et al. Genomics and
metagenomics of trimethylamine-utilizing *Archaea* in the human gut microbiome. ISME J.
2017;11:2059-74.
[5] Stremmel W, Schmidt KV, Schuhmann V, Kratzer F, Garbade SF, Langhans CD, et al. Blood
trimethylamine-*N*-oxide originates from microbiota mediated breakdown of phosphatidylcholine
and absorption from small intestine. PLoS One. 2017;12:e0170742.
[6] Bennett BJ, de Aguiar Vallim TQ, Wang Z, Shih DM, Meng Y, Gregory J, et al.
Trimethylamine-*N*-oxide, a metabolite associated with atherosclerosis, exhibits complex genetic
and dietary regulation. Cell Metab. 2013;17:49-60.
[7] de la Huerga J, Popper H. Urinary excretion of choline metabolites following choline
administration in normals and patients with hepatobiliary diseases. J Clin Invest. 1951;30:463-
70.
[8] Taesuwan S, Cho CE, Malysheva OV, Bender E, King JH, Yan J, et al. The metabolic fate of
isotopically labeled trimethylamine-*N*-oxide (TMAO) in humans. J Nutr Biochem. 2017;45:77-
82.
[9] Dumas ME, Barton RH, Toye A, Cloarec O, Blancher C, Rothwell A, et al. Metabolic profiling
reveals a contribution of gut microbiota to fatty liver phenotype in insulin-resistant mice. Proc
Natl Acad Sci U S A. 2006;103:12511-6.
[10] Dumas ME, Rothwell AR, Hoyles L, Aranias T, Chilloux J, Calderari S, et al. Microbial-host
co-metabolites are prodromal markers predicting phenotypic heterogeneity in behavior, obesity,
and impaired glucose tolerance. Cell Rep. 2017;20:136-48.
[11] Spencer MD, Hamp TJ, Reid RW, Fischer LM, Zeisel SH, Fodor AA. Association between
composition of the human gastrointestinal microbiome and development of fatty liver with
choline deficiency. Gastroenterology. 2011;140:976-86.
[12] Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, et al. Intestinal microbial
metabolism of phosphatidylcholine and cardiovascular risk. N Engl J Med. 2013;368:1575-84.
[13] Gao X, Liu X, Xu J, Xue C, Xue Y, Wang Y. Dietary trimethylamine *N*-oxide exacerbates
impaired glucose tolerance in mice fed a high fat diet. J Biosci Bioeng. 2014;118:476-81.
[14] Zhu W, Wang Z, Tang WHW, Hazen SL. Gut microbe-generated trimethylamine *N*-oxide from
dietary choline is prothrombotic in subjects. Circulation. 2017;135:1671-3.
[15] Kloiber O, Banjac B, Drewes LR. Protection against acute hyperammonemia: the role of
quaternary amines. Toxicology. 1988;49:83-90.
[16] Miñana MD, Hermenegildo C, Llsansola M, Montoliu C, Grisolía S, Felipo V. Carnitine and
choline derivatives containing a trimethylamine group prevent ammonia toxicity in mice and
glutamate toxicity in primary cultures of neurons. J Pharmacol Exp Ther. 1996;279:194-9.
[17] de la Huerga J, Gyorgy P, Waldstein S, Katz R, Popper H. The effects of antimicrobial agents
upon choline degradation in the intestinal tract. J Clin Invest. 1953;32:1117-20.

[18] Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, et al. Gut flora metabolism of
phosphatidylcholine promotes cardiovascular disease. Nature. 2011;472:57-63.

[19] Rebouche CJ, Mack DL, Edmonson PF. L-Carnitine dissimilation in the gastrointestinal tract of
the rat. Biochemistry. 1984;23:6422-6.

[20] Rebouche CJ, Chenard CA. Metabolic fate of dietary carnitine in human adults: identification
and quantification of urinary and fecal metabolites. J Nutr. 1991;121:539-46.

[21] Seim H, Schulze J, Strack E. Catabolic pathways for high-dosed L(-)- or D(+)-carnitine in germ-
free rats? Biol Chem Hoppe Seyler. 1985;366:1017-21.

[22] Koeth RA, Wang Z, Levison BS, Buffa JA, Org E, Sheehy BT, et al. Intestinal microbiota
metabolism of L-carnitine, a nutrient in red meat, promotes atherosclerosis. Nat Med. 2013;19,
576-85.

[23] Jebsen JW, Riaz M. Breakdown products of trimethylamine oxide in airdried stockfish. Means
of enhancing the formation of formaldehyde and dimethylamine. Fish Dir Skr Ser Ernoering.
1977;1:145-53.

[24] Zeisel SH, Da Costa KA. Increase in human exposure to methylamine precursors of *N*-
nitrosamine after eating fish. Cancer Res. 1986;46:6136-8.

[25] al-Waiz M, Ayesh R, Mitchell SC, Idle JR, Smith RL. Disclosure of the metabolic retroversion
of trimethylamine *N*-oxide in humans: a pharmacogenetic approach. Clin Pharmacol Ther.
1987;42:608-12.

[26] Barrett EL, Kwan HS. Bacterial reduction of trimethylamine oxide. Ann Rev Microbiol.
1985;39:131-49.

[27] Takagi M, Ishimoto M. *Escherichia coli* mutants defective in trimethylamine *N*-oxide reductase.
FEMS Microbiol Lett. 1983;17:247-50.

[28] Jameson E, Doxey AC, Airs R, Purdy KJ, Murrell JC, Chen Y. Metagenomic data-mining
reveals contrasting microbial populations responsible for trimethylamine formation in human
gut and marine ecosystems. Microb Genom. 2016;2:e000080.

[29] Dona AC, Jiménez B, Schäfer H, Humpfer E, Spraul M, Lewis MR., et al. Precision high-
throughput proton NMR spectroscopy of human urine, serum, and plasma for large-scale
metabolic phenotyping. Anal Chem. 2014;86:9887-94.

[30] Blaise BJ, Shintu L, Elena B, Emsley L, Dumas ME, Toulhoat P. Statistical recoupling prior to
significance testing in nuclear magnetic resonance based metabonomics. Anal Chem.
2009;81:6242-51.

[31] Dona AC, Kyriakides M, Scott F, Shephard EA, Varshavi D, Veselkov K, et al. A guide to the
identification of metabolites in NMR-based metabonomics/metabolomics experiments. Comput
Struct Biotechnol J. 2016;14:135-53.

[32] Martín-Peláez S, Gibson GR, Martín-Orúe SM, Klinder A, Rastall RA, La Ragione RM, et al. *In vitro* fermentation of carbohydrates by porcine faecal inocula and their influence on *Salmonella* Typhimurium growth in batch culture systems. FEMS Microbiol Ecol. 2008;66:608-19.

[33] Trojanová I, Vlková E, Rada V, Marounek M. Different utilization of glucose and raffinose in *Bifidobacterium breve* and *Bifidobacterium animalis*. Folia Microbiol (Praha). 2006;51:320-4.

[34] Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Stat Soc Ser B. 1995;57:289-300.

[35] Zhang J, Cerny MA, Lawson M, Mosadeghi R, Cashman JR. Functional activity of the mouse flavin-containing monooxygenase forms 1, 3, and 5. J Biochem Mol Toxicol. 2007;21:206-15.

[36] Chilloux J, Everard A, Zhang L, Blancher C, Fuchs J, Hoyles L, et al. Gut microbial trimethylamine inhibits IRAK-4 and chronic inflammation in metabolic syndrome. bioRxiv. 2018;doi:10.1101/277434.

[37] Zoetendal EG, Raes J, van den Bogert B, Arumugam M, Booijink CC, Troost FJ, et al. The human small intestinal microbiota is driven by rapid uptake and conversion of simple carbohydrates. ISME J. 2012;6:1415-26.

[38] van den Bogert B, Boekhorst J, Herrmann R, Smid EJ, Zoetendal EG, Kleerebezem M. Comparative genomics analysis of *Streptococcus* isolates from the human small intestine reveals their adaptation to a highly dynamic ecosystem. PLoS One. 2013;8:e83418.

[39] Cani PD, Amar J, Iglesias MA, Poggi M, Knauf C, Bastelica D, et al. Metabolic endotoxemia initiates obesity and insulin resistance. Diabetes. 2007;56:1761-72.

[40] Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM, Delzenne NM, et al. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. Diabetes. 2008;57:1470-81.

[41] de La Serre CB, Ellis CL, Lee J, Hartman AL, Rutledge JC, Raybould HE. Propensity to high-fat diet-induced obesity in rats is associated with changes in the gut microbiota and gut inflammation. Am J Physiol Gastrointest Liver Physiol. 2010;299:G440-8.

[42] Hildebrandt MA, Hoffmann C, Sherrill-Mix SA, Keilbaugh SA, Hamady M, Chen YY, et al. High-fat diet determines the composition of the murine gut microbiome independently of obesity. Gastroenterology. 2009;137:1716-24.

[43] Ando H, Abe H, Sugimoto N, Tobe T. Maturation of functional type III secretion machinery by activation of anaerobic respiration in enterohaemorrhagic *Escherichia coli*. Microbiology. 2007;153:464-73.

[44] Lee KM, Park Y, Bari W, Yoon MY, Go J, Kim SC, Lee HI, et al. Activation of cholera toxin production by anaerobic respiration of trimethylamine *N*-oxide in *Vibrio cholerae*. J Biol Chem. 2012;287:39742-52.

[45] Paiva JB, Penha Filho RA, Pereira EA, Lemos, MV, Barrow PA, Lovell MA, et al. The

609   contribution of genes required for anaerobic respiration to the virulence of *Salmonella enterica*

610   serovar Gallinarum for chickens. Braz J Microbiol. 2009;40:994–1001.

611   [46] Wu D, Cao M, Peng J, Li N, Yi S, Song L, et al. The effect of trimethylamine *N*-oxide on

612   *Helicobacter pylori*-induced changes of immunoinflammatory genes expression in gastric

613   epithelial cells. Int Immunopharmacol. 2017;43:172-8.

614   [47] Qin N, Yang F, Li A, Prifti E, Chen Y, Shao L, et al. Alterations of the human gut microbiome

615   in liver cirrhosis. Nature. 2014;513:59-64.

616   [48] Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, et al. Richness of human gut

617   microbiome correlates with metabolic markers. Nature. 2013;500:541-6.

618   [49] van den Bogert B, Meijerink M, Zoetendal EG, Wells JM, Kleerebezem M. Immunomodulatory

619   properties of *Streptococcus* and *Veillonella* isolates from the human small intestine microbiota.

620   PLoS One. 2014;9:e114277.

621   [50] Hoyles L, Wallace RJ. Gastrointestinal tract: intestinal fatty acid metabolism and implications

622   for health. In *Handbook of Hydrocarbon and Lipid Microbiology*, vol. 4, part 7, *Human-Animal-*

623   *Plant Health and Physiology Consequences of Microbial Interactions with Hydrocarbons and*

624   *Lipids*. 2010, 3120-32. Edited by T. McGenity, K. N. Timmis, J. R. van der Meer & V. de

625   Lorenzo. New York: Springer.

626   [51] Roger LC, Costabile A, Holland DT, Hoyles L, McCartney AL. Examination of faecal

627   *Bifidobacterium* populations in breast- and formula-fed infants during the first 18 months of life.

628   Microbiology. 2010;156:3329-41.

629   [52] Hoyles L, Murphy J, Neve H, Heller KJ, Turton JF, Mahony J, et al. *Klebsiella pneumoniae*

630   subsp. *pneumoniae*–bacteriophage combination from the caecal effluent of a healthy woman.

631   PeerJ. 2015;doi:10.7717/peerj.1061.

 FIGURE LEGENDS

Fig. 1. *In vivo* confirmation of metabolic retroconversion of TMAO. Reduction of d_9-TMAO to d_9-

TMA was quantified by UPLC–MS/MS up to 6 h after d_9-TMAO gavage and antibiotic treatment,

together with unlabelled TMA and TMAO levels. Plasma quantification of post-gavage (A) d_9-TMA

and (B) d_9-TMAO. *, Significantly ($P < 0.05$; *t* test and corrected for multiple comparison using the

Holm–Sidak method) different from the respective groups not treated with antibiotics. (C) d_9-TMA

bioavailability (AUC). (D) d_9-TMAO bioavailability (AUC). Plasma quantification of post-gavage

unlabelled/endogenous (E) TMA and (F) TMAO. *, Significant between d_9 and d_9 antibiotic

treatment; $, significant between TMAO and TMAO antibiotic treatment. (G) Unlabelled/endogenous

TMA bioavailability (AUC). (H) Unlabelled/endogenous TMAO bioavailability (AUC). Data ($n = 6$

per group) are shown as mean ± SEM. (A, B, E and F). Differences between the bioavailabilities (C,

D, G and H) were assessed using one-way analysis of variance (ANOVA), followed by Holm–Sidak

post hoc tests. Data with different superscript letters are significantly different ($P < 0.05$).

Fig. 2. Effect of TMAO on mixed faecal microbial population *in vitro*. (A) Enumeration of selected

bacteria in fermentation vessels by FISH analysis. Red lines, TMAO-containing systems; blue lines,

negative controls. Data are shown as mean + SD ($n = 3$). Eub338, total bacteria; Ent,

Enterobacteriaceae; Bif164, *Bifidobacterium* spp.; Lab158, lactic acid bacteria. *, Statistically

significantly different (adjusted $P < 0.05$) from the control at the same time point. Full data are shown

in **Supplementary Figure 1**. (B) ^{1}H-NMR data for batch culture samples. Data are shown mean ± SD

($n = 3$). Red lines, TMAO-containing systems; blue lines, negative controls. *, Statistically

significantly different ($P < 0.05$) from the negative control at the same time point. (C) Bidirectional

clustering of correlation matrix of FISH data and data for the six metabolites found in highest

amounts in the NMR spectra from the batch-culture samples. +, Adjusted P value (Benjamini–

Hochberg) statistically significant ($P < 0.05$). FISH and metabolite data and a table of correlations and

adjusted P values (Benjamini–Hochberg) for the batch-culture samples are available in

Supplementary Tables 3–5.

Fig. 3. Influence of TMAO on growth and metabolism of pure cultures of gut bacteria. (A)

Representative growth curves for isolates grown in the presence and absence of TMAO. Red lines,

TMAO-supplemented cultures; blue lines, negative controls. Data are shown as mean ± SD ($n = 3$).

(B) Biplot showing production of various metabolites when isolates were grown in the presence of

TMAO. Summary of data from **Supplementary Table 2**. The larger a circle, the more of the

metabolite produced by an isolate. (C) Differences in metabolites produced when caecal and faecal

isolates of *Escherichia coli* were grown in the presence (+) and absence (−) of 1 % TMAO. Adjusted

(Benjamini–Hochberg) *P* values indicate the caecal isolates were significantly different from the

faecal isolates for a particular metabolite. (D) Lactate production by lactic acid bacteria was increased

in the presence of TMAO. *Enterobacteriaceae*, $n = 20$; *Bifidobacteriaceae*, $n = 17$; *Streptococcaceae*,

$n = 7$; *Enterococcaceae*, $n = 5$. Members of the *Enterococcaceae* and *Streptococcaceae* are

homofermenters (produce only lactic acid from glucose fermentation), whereas the

Bifidobacteriaceae are heterofermenters (produce ethanol, CO_2 and lactic acid from glucose

fermentation), though it should be noted the bifidobacteria included in this study were grown on

raffinose-containing media. Red, TMAO-containing medium; blue, negative control. *, Statistically

significantly different from its negative control (adjusted *P* value < 0.05).

Fig. 1

Fig. 2

681   **Fig. 3**

683 **Table 1.** Details for human-derived gut bacteria screened for their ability to reduce or utilize TMAO

Isolate*	Identified as	Source	Reference	Facultative anaerobe?
D2(9)	[*Clostridium*] *innocuum*	Human faeces	M.L. Jiménez-Pranteda, unpublished	No
L16-FAA1	[*Clostridium*] *ramosum*	Human caecum	L. Hoyles, unpublished	No
L12-BSM1	*Actinomyces odontolyticus*	Human caecum	L. Hoyles, unpublished	No
L6-BSM10	*Actinomyces viscosus*	Human caecum	L. Hoyles, unpublished	No
L6-FAA7	*Bacteroides fragilis*	Human caecum	L. Hoyles, unpublished	No
D1(4)	*Bacteroides vulgatus*	Human faeces	M.L. Jiménez-Pranteda, unpublished	No
L9-FAA7	*Bacteroides vulgatus*	Human caecum	L. Hoyles, unpublished	No
DSM 20083^T	*Bifidobacterium adolescentis*	Human intestine	DSMZ†	No
LCR26	*Bifidobacterium animalis* subsp. *lactis*	Infant faeces	[51]	No
LCR11	*Bifidobacterium bidifum*	Infant faeces	[51]	No
L25-MRS2	*Bifidobacterium bifidum*	Human caecum	L. Hoyles, unpublished	No
LCR5	*Bifidobacterium breve*	Infant faeces	[51]	No
LCR8	*Bifidobacterium breve*	Infant faeces	[51]	No
LCR1	*Bifidobacterium dentium*	Infant faeces	[51]	No
DSM 20093^T	*Bifidobacterium gallicum*	Human intestine	DSMZ	No
LCR6	*Bifidobacterium longum*	Infant faeces	[51]	No
DSM 20088^T	*Bifidobacterium longum* subsp. *infantis*	Infant intestine	DSMZ	No
LCR2	*Bifidobacterium longum* subsp. *infantis*	Infant faeces	[51]	No
DSM 20219^T	*Bifidobacterium longum* subsp. *longum*	Human intestine	DSMZ	No
LCR3	*Bifidobacterium pseudocatenulatum*	Infant faeces	[51]	No
L19-MRS1	*Bifidobacteroum longum* subsp. *longum*	Human caecum	L. Hoyles, unpublished	No
L25-MRS8	*Bifidobacteroum longum* subsp. *longum*	Human caecum	L. Hoyles, unpublished	No
L26-MRS4	*Bifidobactrium animalis* subsp. *lactis*	Human caecum	L. Hoyles, unpublished	No
L26-FAA1	*Citrobacter gillenii*	Human caecum	L. Hoyles, unpublished	Yes
L8-FAA3	*Citrobacter koseri*	Human caecum	L. Hoyles, unpublished	Yes
L16-FAA6	*Clostridium paraputrificum*	Human caecum	L. Hoyles, unpublished	No
L20-BSM1	*Clostridium perfringens*	Human caecum	L. Hoyles, unpublished	No
D1(9)	*Clostridium sporogenes*	Human faeces	M.L. Jiménez-Pranteda, unpublished	No
D2(14)	*Enterococcus faecalis*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D3(1)	*Enterococcus faecalis*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D5(2)	*Enterococcus faecalis*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D6(1)	*Enterococcus faecium*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D6(5)	*Enterococcus gallinarum*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D1(2)	*Escherichia coli*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D2(1)	*Escherichia coli*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D2(2)	*Escherichia coli*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D2(8)	*Escherichia coli*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D3(8)	*Escherichia coli*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D4(15)	*Escherichia coli*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D5(1)	*Escherichia coli*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
L1-FAA5	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L13-FAA2	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L16-FAA5	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L19-FAA2	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L20-FAA3	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L24-FAA5	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L5-FAA2	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L6-FAA1	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L9-MRS1	*Escherichia coli*	Human caecum	L. Hoyles, unpublished	Yes
L9-FAA5	*Fusobacterium ulcerans*	Human caecum	L. Hoyles, unpublished	Yes
L15-FAA9	*Hafnia paralvei*	Human caecum	L. Hoyles, unpublished	No
L4-FAA5	*Klebsiella pneumoniae* subsp. *pneumoniae*	Human caecum	[52]	Yes
L26-MRS5	*Lactobacillus fermentum*	Human caecum	L. Hoyles, unpublished	No

L26-FAA6	*Lactobacillus rhamnosus*	Human caecum	L. Hoyles, unpublished	No
L13-FAA10	*Parabacteroides johnsonii*	Human caecum	L. Hoyles, unpublished	No
D4(1)	*Paraclostridium bifermentans*	Human faeces	M.L. Jiménez-Pranteda, unpublished	No
L1-FAA6	*Pseudomonas aeruginosa*	Human caecum	L. Hoyles, unpublished	Yes
D2(4)	*Staphyloccoccus hominis*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D1(5)	*Streptococcus anginosus*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
D5(12)	*Streptococcus anginosus*	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
L25-MRS1	*Streptococcus gallolyticus*	Human caecum	L. Hoyles, unpublished	Yes
L4-MRS5	*Streptococcus oralis*	Human caecum	L. Hoyles, unpublished	Yes
L4-MRS1	*Streptococcus sanguinis*	Human caecum	L. Hoyles, unpublished	Yes
D4(3)	*Streptococcus* sp.	Human faeces	M.L. Jiménez-Pranteda, unpublished	Yes
L26-MRS7	*Streptococcus vestibularis*	Human caecum	L. Hoyles, unpublished	Yes
LCR4	Unknown *Bifidobacterium*	Infant faeces	[51]	No

685   *L*n*- prefix, different numbers indicate isolates recovered from different individuals.

686   †DSMZ, Leibniz Institute DSMZ – German Collection of Microorganisms and Cell Cultures.