Cost evaluation during decision making in patients at early stages of psychosis

Anna O. Ermakova[1], Nimrod Gileadi, Franziska Knolle[2,3], Azucena Justicia Diaz[2,4], Rachel Anderson[2], Paul C. Fletcher[2,3,4,6], Michael Moutoussis[7], Graham K. Murray[2,3,4]

Unit for Social & Community Psychiatry, WHO Collaborating Centre for Mental Health Services Development, East London NHS Foundation Trust

Department of Psychiatry, University of Cambridge

Behavioural and Clinical Neuroscience Institute, University of Cambridge

Cambridgeshire and Peterborough NHS Foundation Trust

Department of Psychology, University of Cambridge

Institute of Metabolic Science, University of Cambridge

Faculty of Brain Sciences, University College London

Correspondence to Dr Graham Murray gm285@cam.ac.uk

ABSTRACT

Background: Jumping to conclusions during probabilistic reasoning is a cognitive bias reliably observed in psychosis, and linked to delusion formation. Although the reasons why people jump to conclusions are unknown, one suggestion is that they view sampling information as costly. However, previous computational modelling (Moutoussis *et al.* 2011) has provided evidence against it, suggesting that patients with chronic schizophrenia jump to conclusion because of noisy decision making. We therefore developed a novel version of the classical beads-task where the cost of information gathering is systematically manipulated. **Methods:** For 31 individuals with early psychosis, and 31 healthy volunteers we examined the numbers of 'draws to decision' when information sampling had no, a fixed, or an escalating cost. Computational modelling was identical to previous work, and involved estimating a cost of information sampling and a cognitive noise parameters. **Results:** Overall patients sampled less information than controls. However, group differences in numbers of draws became less prominent at higher cost trials, as less information was sampled by both groups. The attenuation of group difference was not due to floor effects, as in the most costly condition participants sampled more information than an ideal Bayesian agent. Computational modelling showed that patients attributed higher costs to information sampling than controls, but groups did not differ in the noise parameter. **Conclusion:** Using a psychological manipulation and computational modelling, we provide evidence that early psychosis patients jump to conclusions because of attributing higher costs to sampling information, and not because of being noisy decision makers.

INTRODUCTION

A consistent psychological finding in schizophrenia research is that patients, especially those with delusions, gather less information before reaching a decision. This tendency to draw a conclusion on the basis of little evidence has been called a jumping to conclusions (JTC) bias (Garety & Freeman 2013). According to cognitive theories of psychosis, a JTC bias is a trait representing liability to delusions. People who jump to conclusions easily accept implausible ideas, discounting alternative explanations, thus ensuring the persistence of delusions (Garety & Freeman 1999; Garety et al. 2007; Garety et al. 2013). Reviews and meta-analyses confirm the specificity, strength and reliability of the association of JTC bias and psychotic symptoms (Fine *et al.* 2007; So *et al.* 2010, 2015; Garety & Freeman 2013; Ross *et al.* 2015; Dudley *et al.* 2016).

The JTC bias is commonly measured in psychosis research using variants of the "beads in the jar" task (Huq *et al.* 1988). A person is presented with beads drawn one at time from one of two jars containing beads of two colours mixed in opposite ratios (Huq *et al.* 1988; Garety *et al.* 1991). Jumping to conclusions has been operationally defined in the 'beads' tasks as making decisions after just one or two beads (Garety *et al.* 2005; Warman *et al.* 2007). The commonly used outcome in this task is the number of beads seen before choosing the jar, known as draws to decision (DTD). The task requires the participant to decide of how much information to sample before making a final decision. This behaviour can be compared to the behaviour of an ideal Bayesian reasoning agent (Huq *et al.* 1988). However, when people experience evidence seeking as costly, it is thought that gathering less information could be seen as optimal leading to monetary gains at the expense of accuracy (Furl & Averbeck 2011).

Computational models allow to consider important factors influencing decisions: The "costed Bayesian model" (Moutoussis *et al.* 2011) incorporates a Bayesian consideration of future outcomes with the subjective benefits or penalty (cost) for gathering additional information on each trial and the noise during the decision making

Although the JTC bias has been well replicated, the neurocognitive mechanisms underlying it are unknown; many possible psychological explanations have been put forward, not all of which are mutually exclusive (Evans *et al.* 2015). Motivational factors, such as intolerance of uncertainty (Bentall & Swarbrick 2003; Broome *et al.* 2007), a "need for closure" (Colbert & Peters 2002), a cost to self-esteem of seeming to need more information (Bentall & Swarbrick 2003) or an abnormal

"hypersalience of evidence" (Menon *et al.* 2008; Speechley *et al.* 2010; Esslinger *et al.* 2013) have been posited as potentially underlying the JTC bias. A common theme emerging from the intolerance of uncertainty, the need for closure, and the cost to self-esteem hypotheses is that patients experience an excessive cost of sampling information.

Moutoussis and colleagues (2011) applied a costed Bayesian model to the information sampling behaviour of a sample of chronic schizophrenia patients undertaking the beads task. Using computational modelling, they found, contrary to their expectations, that a higher perceived cost of the information gathering did not underpin the JTC bias. Therefore, they concluded that differences in the 'noise' of decision making were more useful in explaining the differences between patients and controls than the perceived cost of the information sampling. However, here we reason that a rejection of the "increased cost of information sampling" account of the JTC bias based on this finding is premature, as patients with chronic schizophrenia may not be representative for all psychosis patients, especially not for those at early stages of psychosis. Additionally, different cognitive factors may contribute to the JTC bias; the balance of contributory factors may differ in different patient populations, with noisy decision making relating to executive cognitive impairments predominating in chronic, but perhaps not in early stages of psychosis. Furthermore, Moutoussis and colleagues (2011) applied their model to an existing dataset (Corcoran *et al.* 2008) which used the classic beads task. In this task, no explicit value was assigned to getting an answer correct or incorrect, and no explicit cost was assigned to gathering information. The authors themselves concluded that their work required replication, including incorporation of experimental manipulation of rewards and penalties.

This study investigated the hypothesis that patients with early psychosis attribute higher costs to information sampling using a novel version of the traditional beads task and computational modelling. Focusing on patients at early stages of psychosis allowed us to investigate the JTC bias before the onset of a potential neuropsychological decline seen in patients with chronic schizophrenia, and to study a largely unmedicated sample of psychosis patients. Specifically, we were interested in testing whether patients adapted their decision strategies when there is an explicit cost of information sampling. We, therefore, developed a variation of the beads task in which there were blocks with and without an explicit cost of information sampling, and we gave feedback for correct and incorrect answers. This manipulation allowed the comparison between groups on different cost schedules. However, it would also allow us to test the competing hypothesis which is that psychosis patients jump to conclusions because of "noisy" decision making behaviour.

Under this account, patients should be insensitive to a cost manipulation in the novel setup of the paradigm and apply random decision making.

This study presents a novel investigation of the processes that lead to reduced information sampling in psychosis. We hypothesised that (1) psychosis patients would gather less information than controls when gathering information is cheap, and that (2) psychosis patients and control would adjust their information sampling according to experimental cost manipulations and that the adjustments would mitigate, but not abolish, the difference between the groups. Finally, we hypothesised that (3) the costed-Bayesian model applied to this paradigm and largely unmedicated early psychosis group would provide explanatory evidence for JTC bias in favour of less information because of higher perceived costs rather than noisy decision making.

METHODS

Study participants

The study was approved by the Cambridgeshire 3 National Health Service research ethics committee. An early psychosis group (N=31) was recruited, consisting of individuals with first episode psychotic illness (N=14) or with At Risk Mental States (ARMS; N=17), from the Cambridge early intervention service in psychosis, CAMEO. Inclusion criteria were as follows: age 16–35 years, current psychotic symptoms meeting either ARMS or first episode of psychosis criteria according to the Comprehensive Assessment of At-Risk Mental States (CAARMS) (Yung *et al.* 2005; Morrison *et al.* 2012). Healthy volunteers (N=35) without a history of psychiatric illness or brain injury were recruited as control subjects. None of the healthy controls had drug or alcohol dependence. Healthy volunteers have not reported any personal or family history of neurological or psychotic illnesses, and were matched with regard to age, gender, handedness, level of education, and maternal level of education. None of the patients with ARMS were taking antipsychotic medication, and four patients with FEP were on antipsychotic medication at the time of testing. All of the experiments were completed with the participant's written informed consent.

Jumping to conclusions task

This was a novel task (Figure 1), based on previously published tasks (Huq *et al.* 1988; Garety *et al.* 1991) of reasoning bias in psychosis, but amended in the light of decision-making theory, according to which the amount of evidence sought is inversely proportional to the costs of information sampling. These costs include the high subjective cost of uncertainty, and the cost to self-esteem or other factors (Moutoussis *et al.* 2011). Participants were told that there are two lakes, each containing black and gold fish in two different ratios (60:40). The ratios were explicitly stated and displayed on the introductory slide. A series of fish was drawn from one of the lakes; all the previously 'caught' fish were visible, in order to reduce the working memory load. The participants were informed that fish were being 'caught' randomly from either of the two lakes and then allowed to 'swim away.' We used a pseudorandomised order for each trial which was the same for all of the participants. The lake where the fish were drawn from was also pseudorandomised.

Participants could ask for a maximum of 20 fish being shown. After each fish shown they either indicated whether the fish came from Lake G or B, or asked to see t more fish. The outcome measure was draws to decision (DTD), and the trial terminated when the subject chose the lake.

There were four blocks, each with the 10 trials of the predetermined sequences, in order to increase reliability. The first block was similar to the classical beads task, and was included to provide a reference point. The only difference was that feedback ('correct' or 'incorrect') was provided after each trial. In the second block, a win was assigned to a correct decision (100 points) and a loss (–100 points) to an incorrect decision. In the third block, the cost of each extra fish after the first one was introduced (–5 points) and was subtracted from the possible win or loss of 100 points for making a correct or incorrect decision respectively. Block 4 was similar to Block 3, but the information sampling cost was incrementally increased. The first fish would cost 0 points, the second –5 points, the third –15 points, etc.). Thus, the more fish would be sampled, the more points would be lost. Subjects performed the task at their own pace. Whether Lake G or B was correct was randomised. The task consisted of four blocks. Within each block there were 10 trials of predetermined sequences of fish, in order to increase reliability. All of the fish that were 'caught' during one trial were visible on the screen in order to minimise the working memory load. Block order was not randomised because the task increased in complexity.

The main outcome variable was the number of fish sampled (draws to decision, DTD). Secondary outcomes were the accuracy of the decision, calculated according to Bayes' theorem based on the probability of the chosen lake given the colour and number of fish seen (Everitt & Skrondal 2010)

(see supplementary material for details), and the dichotomous JTC variable (whether or not the decision is made after two fish or less).

Rating scales and questionnaires

The participants underwent a general psychiatric interview and assessment (Table 1) using the Comprehensive Assessment of At Risk Mental States (CAARMS) (Yung *et al.* 2005; Morrison *et al.* 2012), the Positive and Negative Symptom Scale (PANSS) (Kay *et al.* 1987), the Scale for the Assessment of Negative Symptoms (SANS) (Andreasen 1989) and the Global Assessment of Functioning (GAF) (Hall 1995). The Beck's Depression Inventory (BDI) (Beck *et al.* 1996) was used to assess depressive symptoms during the last 2 weeks. IQ was estimated using the Culture Fair Intelligence Test (Cattell & Cattell 1973). Schizotypy was measured with the 21-item Peters et al. Delusions Inventory (PDI-21) (Peters *et al.* 2004).

Statistical analysis of the behavioural data

The effect of manipulations of wins, losses and costs on the DTD and accuracy was assessed by repeated-measures analysis of variance (ANOVA) using SPSS 21 (IBM Corp.). We looked at the trend of reducing DTD with increasing costs of decision making of the last three blocks only in order to better interpret the group × block interactions (the last three blocks are matched in terms of an explicit value of a correct response, whereas the first block has no explicit value of a correct response). Although DTD was not normally distributed, repeated-measures ANOVA is robust to violations of normality, and was therefore an appropriate test to run. IQ and depressive symptoms scores were initially included as covariates and dropped from the subsequent analysis where non-significant. We report two-tailed p-values, which were significant at $p < 0.05$. When the assumption of sphericity was violated we applied Greenhouse–Geisser corrections. We also examined whether DTD was correlated with the severity of psychotic symptoms in the patient group (CAARMS positive symptoms) and with schizotypy scores on the PDI in controls using Spearman's correlation coefficients. The intra-class correlation coefficients (ICCs) were used to estimate the consistency of decision making. We calculated the ICCs of the mean number of choices in each condition separately for patients and controls.

For completeness, we report data on the dichotomous variable JTC, defined as making a decision after receiving one or two pieces of information (see supplementary materials).

Computational Modelling

In addition to the statistical analysis above, we conducted further analysis using a computational model used for a similar experiment (Moutoussis *et al.* 2011). This model is based on the behaviour of an ideal Bayesian agent operating on the beads problem. The ideal Bayesian agent has a known cost associated with sampling more data, and known reward values for guessing the colour correctly. The agent makes choices that maximise its expected reward. Behaviour under these assumptions is explored. To adapt this model to human behaviour, a noise parameter is introduced, which allows the ideal Bayesian decisions (which are deterministic) to be corrupted with probability governed by the noise parameter (Moutoussis *et al.* 2011).

Thus there are two parameters that could describe a participant's behaviour:
- CS: the cost of sampling, is a measure of how much cost the participant assigns to getting more information, compared to the final reward. High values of CS mean that decisions are made early.
- T: the noise parameter. As this value increases, the probability of the participant following the ideal behaviour (for their given value of CS) decreases, and their actions become more uniformly random.

Given sufficient data, one could estimate these two parameters for each participant, and make statistical claims about different participant groups given those values. Unfortunately, deriving these parameters from our experimental data is difficult: there are different combinations of the cost and noise parameters that could explain one participant's behaviour, and the available data are not sufficient to estimate a pair of parameters for each participant.

Instead, we used what is known as a random effects model: We assumed that a participant's model parameters are drawn from a group distribution of parameters. Instead of estimating each individual participant's parameters, we estimate the group's distribution, characterised by a mean and a variance. So instead of fitting CS and T for each participant, we try to estimate the mean and variance for CS and for T for the whole group. For a group of, for example, 30 participants with 10 data points each, we use 300 data points to estimate 4 values (i.e. mean and variance for each of the two parameter distributions), rather than 2 different parameters for each of the 30 participants (i.e. 60 parameters in total). This allows us to get meaningful and statistically correct results from a smaller data set.

To do this parameter estimation, we used a technique called EM (expectation maximization), using code adapted from Moutoussis et al. (2011). This is an optimization method that tries to find maximum-likelihood (ML) parameters for a model, given experimental data.

For our experiments, we ran between 25 iterations of the EM algorithm. Runs took about 30s per iteration per participant on a single core of an Intel Core2 Duo CPU and 4GB of RAM.

Once the ML parameters for a given group are estimated, they can be interpreted using the integrated Bayesian Information Criterion (iBIC (Huys *et al.* 2012, 2015)) uses the likelihood values of best-fit parameters for two different models to decide which model best represents the data. It contains a penalty term for models which have more parameters and can be used as a form of hypothesis testing: If one model has all participant parameters coming from one distribution (4 parameters), and a second model separates the control and unhealthy groups (8 parameters), a comparison of IBIC values can be used to decide if splitting the groups is justified.

RESULTS

Demographical characteristics of the participants

In Table 1 the socio-demographic and cognitive parameters as well as clinical measures of the participants are presented. We have excluded 4 health volunteers with the IQ score > 140 from our original sample, leaving 31 controls. Although the remaining healthy volunteers had higher IQ, compared to the psychosis group, the difference was not significant. The groups were matched with regard to level of maternal education and number of years of education. The control and patient groups did not differ with regard to gender or age. There were significantly more smokers in the patient groups compared with the control group, and the patient group used more drugs. For all of the other measures (e.g. alcohol or cannabis) there were no significant group differences. As expected there are significant differences between the healthy volunteers and the patients in all subscales of CAARMS (see Supplementary Table S1), on the self-reported depression questionnaire (BDI) and self-reported schizotypy scores (PDI). In patient group only we have performed additional clinical assessments. The mean (±SD) score for PANSS positive symptoms was 13.68 (±3.99); for PANSS negative symptoms 9.87 (±4.88); SANS 0.35 (±0.75) and GAF 55.00 (±18.54).

Group differences in the number of DTD and points

Inspection of Figure 2A reveals that in all of the blocks the controls took more draws to decision than the patients. Mauchly's test indicated that the assumption of sphericity was not violated (W(5)=0.211, p<0.001). On mixed-model ANOVA, there were significant main effects of block (F(1)=94.49 , p<0.001 and of group (F(1)=5.99, p=0.017) on the number of DTD. The interaction between the group and the block was also significant (F(3)=39.37, p=0.006). This indicates that, depending on the group, block change had different effects on DTD. Post-hoc Bonferroni-corrected tests demonstrated that group differences were significant in the first two blocks (block 1: p=0.007; block2: p=0.028), whereas in the third and fourth block there were no group differences (block3: p=0.059; block 4: p=0.419). The group differences became increasingly reduced as the cost of decision-making became increasingly high. Controls behaviour was more similar to the ideal Bayesian reasoner than patients on the first three blocks but not block 4 (see Supplementary Table S2). Group differences in the probability of being correct were very similar to the results in the number of DTD (Figure 2B and Supplementary materials).

Analysing the point won/lost in block 2-4 (Figure 3), we identified four outliers in each groups that significantly exceeded ±2SD threshold. After excluding these subjects, the mixed-model ANOVA revealed a significant effect for block (F(2)=93.73, p<0.001), a marginally significant interaction between block x group (F(2)=2.52, p=0.086) and a significant groups effect (F(1)=4.14, p=0.047). Bonferroni corrected post-hoc analysis showed that patients won significantly fewer points in block 2 (F(2)=4.65, p=0.035), but did not differ from controls in block 3 and 4 (both p>0.3).

Intra-class correlations of DTD and SD of the mean DTD

Intra-class correlation coefficients (ICCs) of the mean number of DTD within each block were calculated separately for patients and controls. Within all blocks in both groups the correlations were very high, indicating that behaviour was consistent within each block (ICC values: block 1: patients 0.965, controls 0.943; block 2: patients 0.982, controls 0.973; block 3: patients 0.972, controls 0.976; and block 4: patients 0.973, controls 0.978).

Correlation of symptoms and IQ with DTD

We calculated two-tailed Spearman's correlations with positive psychotic symptoms in patients and with PDI scores in controls. In the patient group, we used a summary score of the three CAARMS

subscales that quantify positive psychotic symptoms, namely Unusual Thought Content, Non-Bizzare Ideas (mainly persecutory ideas) and Perceptual Aberrations. An additional advantage of the summary measure is that it provides one measure to reduce the number of correlations that need to be performed. We also ran correlations BDI scores because the groups differed on this measure. In the group with psychotic symptoms the correlations with the overall CAARMS score were significant in the first three blocks (block 1: rho=–0.515, p=0.003; block 2: rho= –0.489, p=0.005; block 3: rho= –0.491, p=0.005). There were no correlations in the psychotic symptoms group between DTD and BDI score (for all p>0.1). Negative symptoms were not correlated with DTD.

In controls, we found that there were significant correlations of DTD in block 1 with the Distress (rho=-394, p=0.035) and Preoccupation (rho = –0.462, p = 0.012) subscales of the PDI. DTD in block 2 correlated with the Preoccupation subscale of PDI (rho=–0.387, p=0.038). There were no significant correlations with BDI scores (e.g. for block 1 rho=-0.023, p=0.905).

In the patients we furthermore found a positive correlation between IQ and the first three blocks (block 1: rho=–0.481, ρ = 0.006; block 2: rho=–0.362, p=0.045; block 3: rho=–0.364, p=0.044). There was no such correlation in the controls.

Computational Modelling Results

Our computational analysis found that in block 1-3, patients with early psychosis assigned a higher cost to sampling more data than healthy controls (Table 2). For example, in Block 1, where there was no explicit cost, the modelled mean cost of sampling in the controls was very low 1.9×10^{-3}, and in patients 1.7. The modelled variance was also higher for the patient group (13) compared to the controls (2.0×10^{-6}). The estimated noise parameters were similar in each group, with overlapping confidence intervals. For example, in block 1, the mean noise parameter for controls was 3.5, and 4.0 in patients. The logliklihoods (LLs) of the models were closer to zero when fitting the models as two different groups compared to a single group, consistent with the theory that the two groups do differ from each other in their behaviour.

In block 3, where an explicit cost of 5 points per fish sampled was assigned, the model finds that patients very slightly overestimated the cost of sampling compared to the assigned cost (estimated mean cost of sample for patients 5.2), whereas the controls underestimated it (estimated mean cost of sample 1.3).

In block 4, we explicitly told participants that the cost for additional information is not fixed, but increases with amount of information requested. We did not fit the computational model to block 4 because the model does not take into account the increasing cost structure set up.

To test the null hypothesis that both groups are drawn from the same distribution of behaviour parameters, we used iBIC. Table 3 shows the results of that calculation, where the null hypothesis can be rejected with strong evidence. For example, in Block 1, despite the fact the iBIC penalises the use of extra parameters substantially, when fitting CSm, CSv, Tm and Tv, separately for each group iBIC was 3450.2, compared to a iBIC of 3489.7 for fitting them as one combined group. Taken together, the iBIC can be used to compare the hypothesis about the models of interest. In our case those are the following two: 1) Unhealthy and healthy groups have independent CS and T parameter distributions, characterised by 8 numbers (mean and variance for CS and for T, for each group). 2) All participants come from the same pool of noise and cost parameters, characterised by 4 numbers. The iBIC suggests that hypothesis 1 is better explained by the separate model in blocks 1 and 2, and by the combined model in block 3.

DISCUSSION

Our study shows that early psychosis patients generally gather less information before coming to a conclusion compared to healthy controls. Both groups slightly increased their DTD when rewarded for a correct answer (block 2), and significantly decreased their DTD when there was an explicit cost for the sampling of information (block 3 and 4). The decrease was strongest in block 4 including the incremental cost increase for each 'extra fish.' These effects were especially strong in controls, as they gathered significantly more information during block 1 and 2 compared to patients. Thus, patients had a smaller range to exhibit less DTD with increasing cost in patients. However, we want to point out that an ideal Bayesian decision agent would still sample fewer information in than controls or patients in block 3 and 4. This indicates that potential floor effects may not be responsible for the decreased reduction of DTD in patients compared to control. Together with our modelling results, this finding rather supports the hypothesis that independent of the true cost-value of information, patients with early psychosis experience information sampling as more costly than controls.

In this novel version of the beads task using conditions with an increasing cost scheme, jumping to conclusions is actually an advantageous strategy, because sampling a large amount of very costly information would cancel out the potential win of correctly identifying the lake. Consistent with this, group differences were especially strong in the first two blocks, where information sampling was free. Despite being rewarded for a correct decision in block 2, patients did not significantly adapt their behaviour compared to block 1. They still sample significantly fewer information resulting in lower accuracy and few points won in block 1 and 2. In the third block, when there was only a small cost of information, both groups responded to that change by lowering the number fish seen. The difference between the groups was marginally significant, that patients still applied fewer DTD. However, further increasing the information sampling cost completely abolished the group differences. Our data furthermore show that patients were significantly worse in overall accuracy (i.e. probability of being correct at the time of making the decision) and total points won indicating the application of an unsuccessful strategy. In general, these results show that healthy controls were more flexible in adapting their information sampling to the changed task conditions. They increased the number of fish sampled when there was a reward for the correct decision, and decreased it when information sampling had a cost. Patients also decreased the number of DTD when information sampling became more costly, but not as much as the controls, showing that the patients view information sampling generally as costly, independent the actual value and the feedback. The results on block 1 and 2, furthermore, indicate that psychosis patients have difficulties integrating feedback appropriately to update their future decisions. This is similar to what has been reported in a recent study on the win-stay/lose-shift behaviour in an early psychosis group (Ermakova *et al.* 2017).

The variance across the two groups was similar, indicating consistent decision making behaviour across the 10 trials in each block and within each group. If patients were acting more randomly, they would apply a more noisy decision making behaviour. This is in contrast to Moutoussis et al. (2011). Concluding from their Bayesian modelling data, they proposed that patients had more noise in their responses, leading to the reduced number of DTD. When modelling our data in a similar way, we found a significant difference between the groups in the perceived cost of information sampling, but not in the noise of the decision behaviour. We argue that this is mainly due to the differences in the patient groups used in the two studies. Whereas Moutoussis et al. (2011) used chronic schizophrenia patients with a potential neuropsychological decline, our study used patients at early stages with preserved cognitive functioning. Although we altered also our task to specifically investigate information sampling in relation to cost evaluation, block 1 was very similar to the previous study. It

only differed in providing feedback (i.e. 'correct' or 'incorrect') after each decision, which potentially led the patients to stick to the same strategy throughout, which might be why we also did not see differences in the noise parameter in block 1. However, as argued above it rather seems that patients are unable to integrate the feedback appropriately. When analysing true cost of information sampling, we found that patients overestimated the cost, while the controls underestimated it. We, furthermore, noticed that key contributors to the best fit parameters are those who jump to conclusions most: participants who consistently made a decision after viewing only one fish. The proportion of those individuals was significantly higher among the patients.

The percentage of people with psychosis who demonstrated JTC reasoning style in our study was relatively low, at 26%, compared to previous studies that report 40% of individuals (Dudley *et al.* 2011) or, half to two-thirds of the individuals with delusions (So *et al.* 2010). Likewise, our DTD measure was slightly higher than that in most other studies. This might be due to the use of early stage psychosis patients and presence of feedback in our task compared to previous version. Feedback has been shown to increase information sampling and accuracy both in patients with delusions and in controls (Lincoln *et al.* 2010).

Additionally, we found a correlation between IQ and DTD in the first three blocks in the patient group, so we cannot completely exclude the contribution of intelligence to the information-gathering bias. Some argue that bad executive functions or working memory deficits contribute to the JTC bias (Garety *et al.* 2013; Falcone *et al.* 2015). Alternatively, a low IQ could lead to a low tolerance for uncertainty and an equivalent high cost of the information sampling, as well as the inability to integrate feedback in order to update future decisions. If more information is unpleasant, because it exceeds one's capacity to utilise it, it could be viewed as costly. However, the fact that the correlation does not appear in controls argues against it. Therefore, low IQ is likely to contribute to the JTC bias, but cannot solely explain its existence in psychosis.

Rather than focusing on ICD-10 or DSM-V schizophrenia patients, we studied a group of patients early in their course of psychosis. All patients in our study suffered from current psychotic symptoms. The distinction between ARMS and FEP patients is useful for clinical purposes, but it is potentially not very informative when investigating the underlying mechanisms of psychosis. This was motivated by the hypothesis that positive symptoms vary as a continuous variables throughout the population. Psychosis is often viewed as an upper part of the continuum, ranging from rare occurrences of delusions or hallucinations at one end, through individuals with many 'schizotypal'

traits (van Os *et al.* 2009). Consistent with this approach, we found that the more severe their positive symptoms were, the less information they sampled. To investigate the idea of the psychosis continuum further, we looked at the correlations between information sampling and schizotypy characteristics in healthy volunteers. In this group, we found a negative correlation between the number of DTD and the scores on the distress and preoccupation subscales of the PDI, indicating that less information sampling is associated with higher scores. This is consistent with studies by Colbert and Peters (2002) and Lee et al. (2011), as well as the recent meta-analysis by Ross et al. (2015) and fits with a continuum model of psychosis.

Summary

In summary, we have found that early psychosis patients demonstrate a hasty decision-making style compared with healthy volunteers, sampling significantly fewer information. Moreover, psychosis patients were less able to adapt to the changing demands of the task. Patients performed worse with increased total score for positive symptoms.

Figures and Tables

Figure 1: Experimental design of a single trial. In 50% of the trials fish were coming from the black lake, and in 50% they were coming from the gold lake. The order was pseudorandomised, so that the same sequences were used for all participants. Feedback, depending on the block, was either the words 'Correct' or 'Incorrect' in block 1 or the number of points won or lost)during the trial in all subsequent blocks.

Figure 2: A) Mean number of draws to decision in the four blocks of the task. B) Probability of being correct (Accuracy) at the time of making the decision in four task blocks.

Figure 3: Mean number of point won or lost in block 2-3 across all 10 trials.

Table 1: Sample characteristics for healthy controls and people with psychotic symptoms.

Variable	Controls (n = 31)		Early psychosis (n = 31)		Statistics
	Mean	SD	Mean	SD	Value (df), Significance p < 0.05
Age (years)	21.58	2.41	22.52	4.66	t(60) = 0.993, p = 0.325
Gender (male/female)	18/13		18/13		χ(1) = 0.0, p = 1
IQ	110.52	15.79	102.26	17.91	t(60) = −1.926, p = 0.059
Level of education	2.35	0.79	2.00	1.17	U(2) = 373.5, p = 0.117
Mother's level of education	2.19	0.91	2.11	1.37	U(2) = 440.0, p = 0.884
Smoking (yes/no)*	6/25		18/13		χ(1) = 9.79, p = 0.004
Alcohol	2.42	0.85	1.78	1.37	U(2) = 419.5, p = 0.368
Cannabis	0.90	0.79	1.26	1.23	U(2) = 430.0, p = 0.459
Other drugs*	0.49	1.02	1.11	1.36	U(2) = 355.5 , p=0.039
PDI-21*	4.45	2.34	7.83	4.58	t(45) = 3.359, p = 0.002
Distress*	9.52	6.94	24.00	15.32	t(45) = 4.430, p < 0.001
Preoccupation*	10.28	6.48	24.78	16.11	t(45) = 4.341, p < 0.001
Conviction*	13.31	8.13	26.56	17.11	t(45) = 3.58, p = 0.001
BDI*	3.40	3.90	25.34	14.12	t(57) = 8.197, p < 0.001
CAARMS Summary Score*	0.52	1.15	18.00	7.53	t(60) = 12.776 , p < 0.001

Education was measured on a 5-point scale (from no education to higher university degree). Intelligence was measured with a Culture Fair Intelligence Test. Smoking: 0 = non-smoker, 1 = smoker. Substance use was measured on a 5-point scale (from 0 = never used to 5 = daily user). Other drugs: hallucinogens, stimulants or sedatives. PDI, Peters Delusion Inventory. BDI, Beck Depression Inventory, CAARMS, Comprehensive Assessment of At Risk Mental States summary score is a summary score of Unusual Thought Content, Non-Bizzare Ideas and Perseptual Abnormalities intensity and frequency subscales (individual subscales of CAARMS, and other clinical assessment measures can be found in the supplementary table x). χ, Pearson's chi-square; t, t-test, SD, standard deviation. U, Mann–Whitney test. *Significant differences.

Table 2: Best-fit distribution parameters for all groups in all experiments.

Group	CSM	CSV	TM	TV	LL
Block 1: No cost					
Control	1.9×10^{-3}	2.0×10^{-6}	3.4	13	-943.747
Psychosis	1.7	13	4.2	13	-762.993
Combined	0.7	2.2	3.6	13	-1735.65
Block 2: No cost, but win or loss (±100)					
Control	5.0×10^{-3}	1.3×10^{-5}	3.6	21	-841.202
Psychosis	3.0	44	2.9	7.5	-690.567
Combined	1.5	13	2.5	7.0	-1549.28
Block 3: Fixed cost, plus win or loss (±100)					
Control	1.6	9.9	4.1	2.2	-707.604
Psychosis	5.2	100	4.4	2.2	-545.704
Combined	3.1	38	4.2	1.0	-1260.45

Best fit parameters were calculated for each participant group separately, and for the group formed by combining both participant groups. CS_M is the mean cost per sample for a particular group. CS_V is the variance of cost per sample within a particular group. T_M is the mean noise parameter for a particular group. Tv is the variance of the noise parameter within a particular group. LL is the log likelihood of the data given the parameters

Table 3: iBIC values for the model where all participants are drawn from the same distribution, vs the model where the healthy and unhealthy groups differ in their distributions.

Block	iBIC for combined	iBIC for separate	Difference
1 no cost	3489.7	3450.2	39.5 (Very strong)
2 no cost, but win or loss (±100)	3116.9	3100.3	16.7 (Very strong)
3 fixed cost, plus win or loss (±100)	2539.3	2543.4	-4.1 (Preference for combined model)

REFERENCES

Andreasen NC (1989). *Scale for the Assessment of Negative Symptoms (SANS). British Journal of Psychiatry* **155**, 53–58.

Beck AT, Steer RA, Brown GK (1996). Manual for the Beck depression inventory-II. *San Antonio, TX: Psychological Corporation*, 1–82.

Bentall RP, Swarbrick R (2003). The best laid schemas of paranoid patients: autonomy, sociotropy and need for closure. *Psychology and psychotherapy* **76**, 163–71.

Broome MR, Johns LC, Valli I, Woolley JB, Tabraham P, Brett C, Valmaggia L, Peters E, Garety PA, McGuire PK (2007). Delusion formation and reasoning biases in those at clinical high risk for psychosis. *British Journal of Psychiatry* **191**

Cattell RB, Cattell AKS (1973). *Measuring intelligence with the culture fair tests*. Institute for Personality and Ability Testing.

Colbert SM, Peters ER (2002). Need for Closure and Jumping-To-Conclusions in Delusion-Prone Individuals. *The Journal of Nervous and Mental Disease* **190**, 27–31.

Corcoran R, Rowse G, Moore R, Blackwood N, Kinderman P, Howard R, Cummins S, Bentall RP (2008). A transdiagnostic investigation of 'theory of mind' and 'jumping to conclusions' in patients with persecutory delusions. *Psychological Medicine* **38**, 1577.

Dudley R, Shaftoe D, Cavanagh K, Spencer H, Ormrod J, Turkington D, Freeston M (2011). 'Jumping to conclusions' in first-episode psychosis. . Wiley Online Library *Early intervention in psychiatry* **5**, 50–56.

Dudley R, Taylor P, Wickham S, Hutton P (2016). Psychosis, delusions and the 'Jumping to Conclusions' reasoning bias: A systematic review and meta-analysis. *Schizophrenia Bulletin* **42**, 652–665.

Ermakova AO, Knolle F, Justicia A, Bullmore ET, Jones PJ, Robbins TW, Fletcher PC, Murray GK (2017). Abnormal reward prediction error signalling in antipsychotic naïve individuals with first episode psychosis or clinical risk for psychosis. *bioRxiv*

Esslinger C, Braun U, Schirmbeck F, Santos A, Meyer-Lindenberg A, Zink M, Kirsch P (2013). Activation of Midbrain and Ventral Striatal Regions Implicates Salience Processing during a Modified Beads Task. *PLoS ONE* **8**

Evans SL, Averbeck BB, Furl N (2015). *Jumping to conclusions in schizophrenia. Neuropsychiatric Disease and Treatment* **11**, 1615–1624.

Everitt BS, Skrondal A (2010). *The Cambridge Dictionary of Statistics. Journal of Chemical Information and Modeling* vol 53

Falcone MA, Murray RM, O'Connor JA, Hockey LN, Gardner-Sood P, Di Forti M, Freeman D, Jolley S (2015). Jumping to conclusions and the persistence of delusional beliefs in first episode psychosis. . Elsevier *Schizophrenia research* **165**, 243–246.

Fine C, Gardner M, Craigie J, Gold I (2007). Hopping, skipping or jumping to conclusions? Clarifying the role of the JTC bias in delusions. *Cognitive Neuropsychiatry* **12**, 46–77.

Furl N, Averbeck BB (2011). Parietal Cortex and Insula Relate to Evidence Seeking Relevant to Reward-Related Decisions. *Journal of Neuroscience* **31**, 17572–17582.

Garety PA, Bebbington P, Fowler D, Freeman D, Kuipers E (2007). Implications for neurobiological research of cognitive models of psychosis: a theoretical paper. *Psychological Medicine* **37**

Garety PA, Freeman D (1999). Cognitive approaches to delusions: a critical review of theories and evidence. *The British journal of clinical psychology / the British Psychological Society* **38 (Pt 2)**, 113–154.

Garety PA, Freeman D (2013). *The past and future of delusions research: From the inexplicable to the treatable. British Journal of Psychiatry* **203**, 327–333.

Garety PA, Freeman D, Jolley S, Dunn G, Bebbington PE, Fowler DG, Kuipers E, Dudley R (2005). Reasoning, Emotions, and Delusional Conviction in Psychosis. *Journal of Abnormal Psychology* **114**, 373–384.

Garety PA, Hemsley DR, Wessely S (1991). Reasoning in Deluded Schizophrenic and Paranoid Patients. *The Journal of Nervous and Mental Disease* **179**, 194–201.

Garety P, Joyce E, Jolley S, Emsley R, Waller H, Kuipers E, Bebbington P, Fowler D, Dunn G, Freeman D (2013). Neuropsychological functioning and jumping to conclusions in delusions. *Schizophrenia Research* **150**, 570–574.

Hall RCW (1995). Global Assessment of Functioning: A Modified Scale. *Psychosomatics* **36**, 267–275.

Huq SF, Garety PA, Hemsley DR (1988). Probabilistic Judgements in Deluded and Non-Deluded Subjects. *The Quarterly Journal of Experimental Psychology Section A* **40**, 801–812.

Huys QJM, Eshel N, O'Nions E, Sheridan L, Dayan P, Roiser JP (2012). Bonsai trees in your head: How the pavlovian system sculpts goal-directed choices by pruning decision trees. *PLoS Computational Biology* **8**

Huys QJM, Lally N, Faulkner P, Eshel N, Seifritz E, Gershman SJ, Dayan P, Roiser JP (2015). Interplay of approximate planning strategies. *Proceedings of the National Academy of Sciences* **112**, 3098–3103.

Kay SR, Fiszbein A, Opler LA (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia bulletin* **13**, 261–76.

Lee G, Barrowclough C, Lobban F (2011). The influence of positive affect on jumping to conclusions in delusional thinking. *Personality and Individual Differences* **50**, 717–722.

Lincoln TM, Ziegler M, Mehl S, Rief W (2010). The jumping to conclusions bias in delusions: Specificity and changeability. *Journal of Abnormal Psychology* **119**, 40–49.

Menon M, Mizrahi R, Kapur S (2008). "Jumping to conclusions" and delusions in psychosis: Relationship and response to treatment. *Schizophrenia Research* **98**, 225–231.

Morrison AP, French P, Stewart SLK, Birchwood M, Fowler D, Gumley AI, Jones PB, Bentall RP, Lewis SW, Murray GK, Patterson P, Brunet K, Conroy J, Parker S, Reilly T, Byrne R, Davies LM, Dunn G (2012). Early detection and intervention evaluation for people at risk of psychosis: multisite randomised controlled trial. *British Medical Journal* **344**, e2233.

Moutoussis M, Bentall RP, El-Deredy W, Dayan P (2011). Bayesian modelling of Jumping-to-Conclusions bias in delusional patients. *Cognitive Neuropsychiatry* **16**, 422–447.

van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L (2009). A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness–persistence–impairment model of psychotic disorder. *Psychological Medicine* **39**, 179.

Peters E, Joseph S, Day S, Garety P (2004). Measuring Delusional Ideation: The 21-Item Peters et al. Delusions Inventory (PDI). *Schizophrenia Bulletin* **30**, 1005–1022.

Ross RM, McKay R, Coltheart M, Langdon R (2015). Jumping to conclusions about the beads task? A meta-analysis of delusional ideation and data-gathering. *Schizophrenia Bulletin* **41**, 1183–1191.

So SH, Garety PA, Peters ER, Kapur S (2010). Do Antipsychotics Improve Reasoning Biases? A Review. *Psychosomatic Medicine* **72**, 681–693.

So SH wai, Siu NY fan, Wong H lam, Chan W, Garety PA (2015). *'Jumping to conclusions' data-gathering bias in psychosis and other psychiatric disorders — Two meta-analyses of comparisons between patients and healthy individuals*. *Clinical Psychology Review* **46**, 151–167.

Speechley WJ, Whitman JC, Woodward TS (2010). The contribution of hypersalience to the 'jumping to conclusions' bias associated with delusions in schizophrenia. *Journal of Psychiatry and Neuroscience* **35**, 7–17.

Warman DM, Lysaker PH, Martin JM, Davis L, Haudenschield SL (2007). Jumping to conclusions and the continuum of delusional beliefs. *Behaviour Research and Therapy* **45**, 1255–1269.

Yung AR, Yuen HP, McGorry PD, Phillips LJ, Kelly D, Dell'Olio M, Francey SM, Cosgrave EM, Killackey E, Stanford C, Godfrey K, Buckby J (2005). Mapping the onset of psychosis: The comprehensive assessment of at risk mental states. *Schizophrenia Research* **39**, 964–971.