Title: Tracking the dynamic functional connectivity structure of the human brain across the adult lifespan

Yunman Xia[a,b], Qunlin Chen[a,b], MengZe Li [a,b], Jiang Qiu[a,b]*

Affiliation:

[a]Key Laboratory of Cognition and Personality (Ministry of Education), Chongqing 400715, China

[b]School of Psychology, Southwest University, Chongqing 400715, China

Corresponding author: J. Qiu, School of Psychology, Southwest University, Beibei, Chongqing 400715, China.

E-mail address: qiuj318@swu.edu.cn.

Phone: +86-023-68367942

Abstract

The transition from early adulthood to older is marked by pronounced functional and structural brain transformations that impact cognition and behaviour. Here, we use dynamic functional network connectivity method to examine resting state functional network changes over aging process. In general, the features of dynamic functional states are generally varying across ages, such as the frequency of expression and the amount of time spent in the certain state. Increasing age is associated with less variability of functional state across time at rest period. From age point of view, examining the age-related difference of topology index revealed 19-30 age range has the significant largest global efficiency, largest local efficiency of default-mode network (DMN), cognitive control network (CCN) and salience network (SN). As for functional states, one state displayed the whole positive connectivity, in the meantime, it has the largest global efficiency and local efficiency of three subnetworks. Besides, the frequency of another state was negatively correlated to the box block (The Wechsler Adult Intelligence Scale subset, which is thought to evaluate fine motor skills, processing speed, and visuospatial ability), while positively correlated with age, and the box block was inversely correlated to age. The results suggested that cognitive aging may be characterized by the dynamic functional network connectivity. Taken together, these findings suggested the importance of a dynamic approach to understanding cognitive aging in lifespan.

Key words: dynamic functional network connectivity; aging; graph theory; Wechsler Adult Intelligence Scale

Introduction

With the age of the increasing, our brain undergoes structural and functional changes (Betzel, et al., 2014). Structural studies revealed a tendency toward cortical 'disconnection', a rapid decline that levels off in adult life, suggestive of reduced within-network covariance (DuPre & Spreng, 2016). The 'disconnection' is accompany with dynamic changes of resting state functional connectivity. For example, the default mode or the execute control network, which is involved in attention, memory and executive control functions, has less interregional BOLD correlations within network in the cognitive aging process. (Vidal-Piñeiro, et al., 2014; Chan, et al., 2014). It is great importance to investigate changes across lifespan based integrating multi-networks, which provided a whole brain level understanding of aging process. (Zuo et al., 2017)

Functional connectivity (FC) describes different brain regions activity how to interaction over time, which is usually measured by the Pearson correlation between the fMRI time series (Biswal et al., 1995). Previous studies examined age-related differences in functional connectivity based at a static, whole-brain level using neuroimaging techniques, and are linked to cognitive ability and behavioural changes. For example, interactions between intrinsic connectivity networks (ICNs) alter over time across age (Fair et al., 2007; Power et al., 2010; Thomason et al., 2008). Although static FC is widely used in previous literature (Greicius, 2008; Van den Heuvel and Hulshoff Pol, 2010), it may not be enough to fully characterise the human brain. This depiction of development neglects the dynamic characteristic of the brain's functional connectivity, with the potential assumption of FC remains constant throughout the

resting period (Biswal et al., 1995; Fox et al., 2005). Recent research shows that aging not only affects static FC but also its spontaneous reconfiguration over time.

Recently, a growing body of research has investigated the dynamic FC in healthy subjects and patients with neuropsychiatric diseases (Calhoun et al., 2014). For example, Allen and colleagues (Allen et al., 2014) examined resting-state FC dynamics in healthy young adults. And in a follow-up study, researchers found dynamic rs-FC alterations in schizophrenia (Damaraju et al., 2014). Besides resting state FC, researchers examined task FC and confirmed a direct link between cognitive performance and the dynamic reorganisation of the network structure of the brain (James, et al., 2016). Furthermore, researchers found a link between dynamic connectivity in fMRI data and concurrently collected EEG data, and the result revealed the stationarity of connectivity cannot be assumed (Allen et al., 2017). Another study evaluated replicability of dynamic connectivity patterns in 7500 resting fMRI datasets, and results showed distinct connectivity states that are similar across groups during rest (Abrol et al., 2017). These collective findings imply that dynamic FC is a promising avenue for clinical neuroimaging and can enrich our knowledge of the functional organisation of the human brain.

In this study, we provided a method to examine age-related differences in dynamic FC. We began to discuss the correlation between dynamic FC and aging. Based on the rapid increase in journal papers focused on dynamic connectivity (Allen et al., 2014; Calhoun et al., 2014; Damaraju et al., 2014; James, et al., 2016; Abrol et al., 2017), we compared the dynamic functional network connectivity (FNC) in different age cohorts.

Briefly, group independent component analysis (ICA) was first used to extract resting state networks (RSNs), and dynamic FNC matrices were then created using sliding window correlation approach. Subsequently, K-means algorithm was employed to cluster these matrices into different dynamic states (these can be thought of as average patterns that subjects tend to return to during the course of the experiment), and state analysis (based on the patterns of connectivity within each state as well as high-level summaries such as the dwell time each individual subject spends in each state) was finally carried out to compare the dynamic FNC in different age cohorts. We described the states by calculating their network graphic properties including: local efficiency, global efficiency, as well as hub areas. Finally, we discussed some of the issues associated with dynamic FNC and intelligence. We focused on recently developed dynamic FNC and provided a special perspective to examine the correlation between dynamic FNC and aging.

Materials and methods

Subjects

The large sample was drawn from an ongoing project exploring the associations among individual development in brain structure and function, cognitive ability and mental health (Qunlin, et al., 2016; Wei, et al., 2017). A total of 494 healthy volunteers were recruited from SouthWest University (SWU) by means of the campus network, advertisements on bulletin boards and through face-to-face communications, but 60 participants were excluded due to large head motion ($FD > 0.2$mm) (Laumann et al., 2016). Thus, the final sample was composed of 434 subjects (165 males; mean age =

44.44, *SD* = 17.28; age range = 19-80). All participants were required to be healthy and none had a history of psychiatric disorder or substance abuse (including illicit drugs and alcohol), and MRI contraindications. The project was approved by the SWU Brain Imaging Center Institutional Review Board, and written informed consent was obtained from each subject prior to the study. Participants received payment depending on time and tasks completed.

Image acquisition

All functional images were obtained from a 3-T Siemens Magnetom Trio scanner (Siemens Medical, Erlangen, Germany) at the Brain Imaging Research Central in Southwest University, Chongqing, China. The whole-brain resting-state functional images were acquired using T2-weighted gradient echo planar imaging (EPI) sequence: slices = 32, repetition time (TR)/echo time (TE) = 2000/30 ms, flip angle = 90 degrees, field of view (FOV) = 220 mm × 220 mm, thickness = 3 mm, slice gap = 1 mm, matrix = 64 × 64, resulting in a voxel with $3.4 \times 3.4 \times 4$ mm^3. During the functional images acquisition, participants were asked to close eyes, keep still and not to fall asleep (confirmed by all participants immediately after the experiment). The scan lasted for 484 s and acquired 242 volumes in total for each subject. Additionally, high-resolution T1-weighted anatomical images were acquired for each participant (TR = 1900 ms; TE = 2.52 ms; inversion time = 900 ms; flip angle = 9 degrees; resolution matrix = 256 × 256; slices = 176; thickness = 1.0 mm; voxel size = $1 \times 1 \times 1$ mm^3).

Data preprocessing

The sMRI ($1 \times 1 \times 1$ mm^3) data was processed by using SPM8 (Welcome Department

of Cognitive Neurology, London, UK; www.fil.ion.ucl.ac.uk/spm) implemented in MATLAB 2012a (MathWorks Inc., Natick, MA, USA). Each sMRI was first displayed in SPM8 to check quality. Firstly, the reorientation of the images was manually set to the anterior commissure. Then, the images were segmented into gray matter, white matter, and cerebrospinal fluid by using the segmentation tool in SPM8.

The resting-State fMRI data were preprocessed using Data Processing Assistant for Resting-State fMRI (DPARSF, http://resting-fmri.sourceforge.net/) implemented in the MATLAB 2012a (Math Works, Natick, MA, USA) platform. Resting-State fMRI preprocessing steps included the following: eliminate first 10 time points of each subject, slice timing correction, realignment, registration functional images (MNI pace), normalization ($3\times3\times4$ mm^3), smoothing (FWHM = 6 mm), band pass filtering (0.023-0.18 Hz) (Gonzalez-Castillo, et al., 2015; Leonardi & Van De Ville, 2015), nuisance regressors included white matter and cerebrospinal fluid signals in addition to 24 movement regressors derived by expansion, frames with frame wise displacement (FD) > 0.2mm were censored. The residual effects of motion was regressed out in group statistical analysis by including mean frame wise displacement (FD) derived with Jenkinson's relative root mean square (RMS) algorithm as a regressor of no interest. These preprocessing steps were followed by the standard protocol published (Yan et al., 2016).

Head motion correction

Recent studies have demonstrated that head motion has a substantial impact on dynamic FC (Laumann et al., 2016；Siegel et al., 2016). So we use the following steps

to further minimize the effects of head motion. Artifacts were reduced using excluding subjects with high head-motion, nuisance regression (excluding censored frames), interpolation (Power et al., 2014), and band pass-filtering ($0.023 < f < 0.18$ Hz) (Gonzalez-Castillo, et al., 2015; Leonardi & Van De Ville, 2015). Nuisance regressors included the cerebrospinal fluid signals, white matter, and their derivatives, in addition to 24 movement regressors derived by expansion (Friston et al. 1996; Yan et al. 2013). Subjects with max head motion > 0.2mm and 2.0 degrees were censored (Laumann et al., 2016),

Group independent component analysis

Group ICA was performed using the GIFT toolbox (http://mialab.mrn.org/software/gift). We followed the Allen et al. (2014), so we used a relatively high model order (number of components, C= 100) to achieve a 'functional parcellation' of refined cortical and subcortical components corresponding to known anatomical and functional segmentations. Principal component analysis (PCA) was used to reduce data at two levels. First, participant-specific data were reduced to 150 principal components (PCs). Then the reduced data of all participants were further decomposed into 100 PCs using the expectation-maximization algorithm to avoid otherwise prohibitive memory requirements (Roweis 1998). The Infomax ICA algorithm was repeated 20 times in ICASSO (http://www.cis.hut. fi/projects/ica/icasso) and aggregate spatial maps (SMs) were estimated as the modes of the component clusters, and then a total of 59 components were identified as intrinsic connectivity networks (ICNs) finally. The algorithmic and statistical reliability/stability is

investigated by running the algorithm many times with different initial values or with differently bootstrapped data sets, respectively. Furthermore, following a previous study (Allen et al., 2014), we performed additional post processing steps on time courses of the 59 ICNs, including 1) detrend linear, quadratic, and cubic trends, 2) multiple regression of the 6 realignment parameters and their temporal derivatives, 3) low-pass filtering 0.15 Hz, and 4) spike removal to ensure that artifactual spikes do not negatively impact the signal analysis.

Dynamic FNC computation

Before dynamic FNC computation, the time courses of RSNs were temporally bandpass filtered (0.023–0.18 Hz) (Gonzalez-Castillo, et al., 2015; Leonardi & Van De Ville, 2015) to reduce the effects of low-frequency drift and high-frequency physiological noise. The dynamic FNC was computed using a sliding window correlation approach. Since there was currently no formal consensus regarding the window length, we selected the length (22TR) according to former study with a Gaussian of $\sigma = 3$ TR (Allen et al., 2014). The window was shifted with a step of 1 TR, resulting in 210 windows. In each window, the time courses of each pair of the 59 RSNs were used to calculate FNC (Pearson's correlation coefficient) and a 59 * 59 correlation matrix was obtained. A Fisher's r-to-z transformation was then applied to all FNC matrices to improve the normality of the correlation distribution as where r is the Pearson correlation coefficient and z is approximately normally distributed.

K-means clustering

For the dFNC patterns reoccur within subjects across time and across subjects, we

applied the *k*-means algorithm to divide the dFNC windows into separate clusters. The clustering algorithm was applied to a subset of all subject data corresponding to windows that showed greater variance in FNC, and was repeated 150 times to increase the likelihood of escaping local minima, with random initial cluster centroid positions (Allen et al., 2014; Liu et al., 2016). Subsampling was chosen both to reduce redundancy between windows (the chosen time step of 1 TR induces high autocorrelation in FC time series) and to reduce computational demands. The optimal number of clusters was estimated using the elbow criterion, calculating as the ratio of within-cluster distance to between-cluster distance. A *k* of 5 was determined in a search window of *k* ranging from 2 to 10. Finally, the resulting centroids of subsample were used as starting points to cluster all data into 5 clusters. A *k* of 4 and 6 was used respectively to validate the robustness of our results (see SI.B). It has been suggested that the L1-norm is a more effective similarity measure for high dimensional data (Lloyd, 1982).

State analysis

In addition, we performed an exploratory experiment in which we calculated and compared temporal metrics derived from each subject's state vector (Allen et al., 2014). Specifically, we computed four measures in each subject, including: (1) frequency of each state, measured as the number of consecutive windows in each state; (2) dwell time of state, measured as the frequency of unchanged between time window and time+1 window, while the current window and the next window have the same state, the dwell time of the state plus one; (3) total number of transitions, measured as the

number of state transitions; and (4) state transition frequency, measured as the frequency of transitioning from one state to another state.

Parametric tests (Pearson product-moment correlation) were utilized to evaluate the correlation between age and these measures.

Three dimensions of age $*$ time $*$ state

Before the analysis, each connection matrix of cluster centroids was converted to a z-value connection matrix by using Fisher's *r*-to-*z* transform to improve the normality. Threshold RSFC matrixes is critical for obtaining a sparse adjacency matrix, which should be optimal, contains not too many functional connections, and may dilute group difference. Using absolute thresholding method may ultimately change the properties of the original global and local functional connectivity, which may bias the comparisons of graph-theoretic metrics between different groups of subjects (Song, et al., 2014). Therefore, we applied a proportional network threshold of 15% (Whitfield-Gabrieli, &Nieto-Castanon, 2012). We globally thresholded the RSFC matrix at a fixed threshold (K= 0.15), and then measured local and global efficiency based all weighted adjacency matrix by using Brain Connectivity Toolbox (https://sites.google.com/site/bctnet/) (Whitfield-Gabrieli, &Nieto-Castanon, 2012).

For each time window, we can get a matrix and its graphic indexes (global and local efficiency). We averaged the graphic indexes of all time windows of each subject, and then put the average value as the graphic indexes of each subject. Then we divided the subjects into 6 groups according to age, each age span of 10 years (19-30; 31-40; 41-50; 51-60; 61-70; 71-80). This way we can investigate the correlation between age and

graphic indexes. Previous literatures indicted that DMN (Damoiseaux et al., 2008; Biswal et al., 2010; Evers et al., 2012), CCN (Elizabeth DuPre & R. Nathan Spreng, 2016), and SN (Kamen Tsvetanov, et al., 2016) are more susceptible to the effects of aging, therefore, we calculated the correlation between local efficiency of these subnetworks and age.

Besides, we averaged the graphic indexes of the certain time window of the certain age range, and then put the average value as the graphic indexes of the certain time window of the certain age range. Then we can observe the varying curve of different age range in the time course.

Topological properties of discrete functional connectivity states

To characterize the state topological properties, we applied a proportional network threshold of 15%, calculated global efficiency, subnetwork local efficiency, and the degree of every node of every matrixes, and then listed the top three of degree (The Brain Connectivity Toolbox, http://www.brain-connectivity-toolbox.net/) as the hub nodes of matrices. With global efficiency, local efficiency and hub regions of each state, we can depict the characteristic of each state.

The correlation between WAIS and state

The Wechsler Adult Intelligence Scale (WAIS) is an IQ test designed to measure intelligence and cognitive ability in adults and older adolescents (Kaufman & Lichtenberger, 2005). The WAIS-R, a revised form of the WAIS, and consisted of six verbal and five performance subtests (Wechsler, 1981). Firstly, we investigated the relationship between age, state indexes (frequency of each state; dwell time of state;

numbers of transitions), and total score of WAIS. Secondly, we chose one of verbal subsets, called similarity, which measures abstract verbal reasoning as well as semantic knowledge. In this section, participants are given two words or concepts and have to describe how they are similar. We examined the relationship between similarity (the subtest of the Wechsler Adult Intelligence Scale-Revised Chinese revision (WAIS-RC; Gong, 1992)), state indexes, and age. Lastly, we tested the relationship between block design test (the subtest of the Wechsler Adult Intelligence Scale-Revised Chinese revision (WAIS-RC; Gong, 1992), state indexes, and age. The block design test, which is thought to evaluate fine motor skills, processing speed, and visuospatial ability, is most affected by age (Hoogendam, et al., 2014). We hypothesized that age affects the brain, and then the brain affects the behavior.

Results

Static FNC

Figure 1 displays the ICNs identified with group ICA. Network components are provided in Figure 1.Based on their anatomical and presumed functional properties, ICs are arranged into groups of sub-cortical (SC), auditory (AUD), somatomotor (SM), visual (VIS), cognitive control (CC), default-mode (DM), and salience (SN) networks by spatial correlation and visual recognition. The manual arrangement of ICNs is very similar to a previous study (Allen et al., 2014). ICNs are similar to those observed in previous high model order ICA decompositions (Smith et al. 2009; Allen et al. 2011) and cover the majority of subcortical and cortical grey matter. Figure 2 displayed the static FC between ICNs, computed over the entire scan length and averaged over

subjects.

INSERT Figure 1 in HERE

INSERT Figure 2 in HERE

The duration and transition of connectivity states as a function of age

In addition to the describing the static FNC, we can also examine the frequency of states as a function of time and the transitions between them. Figure 3 shows the state assignments as a function of time for the 4 example subjects. As we would see, FC tends to sustain single state during a long period, while transition times are rarely less. We can characterize transition behavior by calculating the frequency of changing from one state to another.

INSERT Figure 3 in HERE

The total dwell time (the sum of dwell time of all states) of states was positively correlated with age ($r = 0.20$, $p = 0.000$, n=434), while the total transition of states was negatively correlated with age ($r = -0.20$, $p = 0.000$, n=434). The states indexes at rest were correlated with age (See Table 1), the distribution of each state in each age ranges is shown in Figure 4.

Besides, recent studies have demonstrated that head motion has a substantial impact on dynamic FC (Laumann et al., 2016；Siegel et al., 2016), therefore, we examined the correlation between head motion and state indexes. The results indicated head motion is negatively correlated with the frequency of state 3 ($r = -0.15$, $p = 0.002$) and the times of transition of state 3-5 ($r = -0.10$, $p = 0.034$).

INSERT Table 1 in HERE

INSERT Figure 4 in HERE

Three dimensions of age ∗ time ∗ state

To characterize the age effects on the global network topological properties, two key graphic metrics were employed, global efficiency and local efficiency, which were all calculated based on weighted networks. Firstly, we found significant age-related differences in network global and local efficiency (see in Table 2). Notably, the age-related difference is observed for global efficiency (global efficiency: $r = -0.24$, $p = 0.000$), 19-30 years old age range has significant greater global efficiency than 51-60, 61-70, and 71-80 years old age range. The correlation between subnetwork local efficiency and age (DMN Elocal: $p = 0.055$, $r = -0.092$; CC_Elocal: $p = 0.019$, $r = -0.11$; SN_Elocal: $p = 0.000$, $r = -0.19$) behaved differently. Intriguingly, no matter which network, its local efficiency of 19-30 age range is the best.

Secondly, we divided the subjects into 6 groups according to age, each age span of 10 years (19-30; 31-40; 41-50; 51-60; 61-70; 71-80). The sample was not equally distributed amongst the age groups. The each age groups have 149, 32, 63, 103, 59, and 28 samples respectively. We considered the distribution may impact differences of global efficient and local efficient among age groups. So we randomly sampled 28 subjects from each age groups and made them a new sample. Then we compared the global efficiency and local efficiency among different age groups in the new sample. We ran the above-mentioned analysis 100 times and reported the ratio of significant result ($p < 0.05$) in Table 2. The results indicated that the age-related difference in global efficiency is reliable and stable, and that in local efficiency of different subnetwork is

unstable.

Besides, we have plotted the three dimensions of age $*$ time $*$ graphic indexes to observed varying curve of different age range in the time course (see Figure 5).

<div align="center">INSERT Table 2 in HERE</div>

<div align="center">INSERT Figure 5 in HERE</div>

Characteristic of dynamic functional connectivity states

We used sliding window approach to estimate dFNC network for each subject. For the dFNC patterns reoccur within subjects across time and across subjects, we then applied the k-means algorithm to divide the dFNC windows into separate clusters. Figure 6 shows the centroids of the 5 dFNC states. In state 1, the whole network displayed slight and moderate negative connectivity. State 2 showed the high positive correlation among AUD, SMN, and VIS, while the negative correlation between SCN and other networks. In state 3, there are negative connections between DMN and other networks as well as SCN and others. In state 4, the whole network displayed slight and moderate positive connectivity, and the high positive coupling among AUD, SMN, and VIS appeared again. In state 5, AUD, SMN, and SN were negatively correlated with DMN and it displayed a whole negative pattern.

To characterize the state topological properties, we applied a proportional network threshold of 15% and calculated global efficiency, subnetwork local efficiency and degree based on thresholded weighted networks (See Table 3). It's worth noting that state 4 has the largest global efficiency, DMN local efficiency, CC local efficiency and SN local efficiency. The hub nodes (the node of the top three degree) are located in

visual and auditory cortex, DMN (posterior cingulate cortex) and CCN (supplementary motor area).

INSERT Table 3 in HERE

INSERT Figure 6 in HERE

The correlation between WAIS and age

Firstly, the total score of WAIS and similarity were both negatively correlated with age (total score: r = -0.51, p = 0.000, n = 82; similarity: r = -0.49, p = 0.000, n = 93), and similarity was negatively correlated with transition of state 3-1 (r = -0.22, p = 0.031, n = 93). However, the box block design is inversely correlated to age (r = -0.53, p = 0.000, n = 82) and frequency of state 5 (r = -0.26, p = 0.019, n = 82), while the correlation between age and the frequency is not significant (r = 0.15, p = 0.18, n = 82). However, we think age is positively correlated with the frequency of state 5, since the total samples (n = 434) suggested the two factors are positively correlated (r = 0.11, p = 0.022). The size and age range of the subsample (n = 82, age range: 23-65) may be the cause of that statistical significance failed to reach borderline in subsample.

Robust Analysis

Some literature (Gonzalez-Castillo, et al., 2015; Leonardi & Van De Ville, 2015) recommend removing frequency components below 1/w (where w is the length of the window), so we used a band pass filtering (0.023-0.18 Hz). However, others (Abrol, et al., 2016; Damaraju, et al., 2014) low-pass filtered time series with a high frequency cutoff of 0.15 Hz by Gift software (http://mialab.mrn.org/software/). To examine our main results, we use the second filtering parameter to re-analysis the correlation

between aging and dynamic FC.

We examined the frequency of states as a function of time and the transitions between them. As we previous results, FC tends to sustain single state during a long period, while transition times are rarely less. The total dwell time (the sum of dwell time of all states) of states was positively correlated with age ($r = 0.14$, $p = 0.003$), while the total transition of states was negatively correlated with age ($r = -0.14$, $p = 0.003$). The results verified the previous finding. As aging process occurring, the pattern of functional connectivity tended to keep stable in resting period. Some states indexes were correlated with age (See SI.D), and the distribution of each state in each age range was shown in SI.E.

Then we compared the states of two analyzes (states identified in first analysis and robust analysis). Firstly, we calculated the Pearson correlation coefficient among states. Considering the correlation coefficient and visual pattern, we think state 5_second is similarity with state 4_first (See SI.F). Secondly, we applied a proportional network threshold of 15% and calculated the global efficiency, subnetwork local efficiency, and degree of each state. The state 5_second, just like state 4_first, had greatest global efficiency and subnetwork local efficiency (See SI.G). Lastly, SI.H showed the centroids of the intrinsic states.

Discussion

Brain's dynamics functional network connectivity over aging

The present study examined the effect of age on dynamic whole-brain FC. The analysis revealed 5 recurring FC states departed with substantial internetwork

correlation variability. In resting state period, the non - random distribution of states in different age ranges suggested that dynamic changes of large-scale brain network may be a fundamental feature of aging process.

Firstly, the number of transitions occurring between multi-connectivity states and the rate of transition between states were all higher among younger than older participants. We can assume that the thinking is more active for young people than older people, or the speed of mind change is faster in resting period.

Secondly, the frequency of occurrence state 1 and 5 increases over aging, while the same parameter for state 3 and 4 decrease over aging. This suggests that the patterns of state 1 and 5 become more active in the older life span, while the patterns of state 3 and 4 become more active in the young life span. State1 and state 5 display a whole negative connectivity network, while state 3 and 4 have relative more positive connectivity. That is mean young participations may experience more network cooperation in the rest period.

Thirdly, a number of studies have proposed that head motion during a scan can substantially affect functional connectivity estimates (Power et al., 2012; Yan et al., 2013; Laumann et al., 2016；Siegel et al., 2016), therefore, we examined the correlation between head motion and state indexes. On the one hand, the relationship between head motion and state dwell time is not significant, which verified the dynamic FC over time is not artifacts. On the other hand, head motion is negatively correlated with the frequency of state 3 while positively correlated with age, meantime the frequency of state 3 is negatively correlated with age. Maybe the state 3 reflects the psychological

characteristics associated with head movement, such as control ability, which decreases in the aging process. That is to say, head motion also changes systematically with age, which may reflect true neurobiological effects of aging. (Geerligs, Tsvetanov., & Henson, 2017).

Lastly, our resulted suggested that dwell time is positively correlated with age, while previous literature (Hutchison & Morton, 2015) reported the dwell time is negatively correlated with age. One possible reason is that previous sample mainly focused 9-32 age range; however, the age range of our subjects included 19-80. After 32 years old, dwell time is still increasing, and 40 years old is a turning point, twists and turns down. To some extent, our work can enlarge and depth this issue. Another is the results are not necessarily comparable as we use a different number of states.

Age-related dynamic changes in network topology

In this study, we examined the age-related difference of global efficiency, different network local efficiency. 19-30 age range has significant greater global efficiency and local efficiency of subnetworks than older age range. The random sampling results indicated that the age-related difference in global efficiency is reliable and stable, however, the local efficiency of subnetwork didn't show the stable difference. Young have greater global efficiency than older maybe because they have more amounts of state 4, which has largest global efficiency in all states. In state 4, the whole network displayed slight and moderate positive connectivity, and the high positive coupling among AUD, SMN, and VIS. The whole positive connectivity may reflect individual experiences more network cooperation in the rest period.

Static FC research found that the absolute global efficiency of brain functional networks showed no significant relationship with age (Cao, et al., 2014), increasing age is accompanied by increasing global efficiency (Chan, et al., 2014; Sala-Llonch 2014), or young have greater global efficiency than older (Achard & Bullmore, 2007). It is not necessarily to compare dynamic FC study and static FC results

Characterize each state

Recently study also revealed discrete functional connectivity states appear to be quite a resemblance in different groups (Abrol et al., 2017), the same as, states identified here also repeat previous states to a certain degree (Allen et al., 2014; Abrol et al., 2017). In state 1, the whole network displayed slight and moderate negative connectivity. State 2 showed the high positive correlation among AUD, SMN, and VIS, while the negative correlation between SCN and other networks. In state 3, there are negative connections between DMN and other networks as well as SCN and others. In state 4, the whole network displayed slight and moderate positive connectivity, and the high positive coupling among AUD, SMN, and VIS appeared again. In state 5, it rendered whole brain negative correlation.

The previous study (Abrol, et al., 2017) and this results evidenced that the patterns of the certain states are recurrent during resting state period, and the occurrence doesn't change with the research methods, sample source or data quality, and thus can as a measure of functional brain aging.

The correlation of cognitive ability and state

Firstly, the total score of WAIS and similarity were both negatively correlated with

age, and similarity was negatively correlated with transition of state 3-1. Secondly, the box block design is inversely correlated to age and frequency of state 5, while the correlation between age and the frequency is not significant. However, we think age is positively correlated with the frequency of state 5, since the total samples (n = 434, age range: 19-80) suggested the two factors are positively correlated. The size and age range of the subsample (n = 82, age range: 23-65) may be the cause of that statistical significance failed to reach borderline in subsample. The block design test evaluates fine motor skills, processing speed, and visuospatial ability, which are decreasing accompanied by age increasing (Hoogendam, et al., 2014). In state 5, AUD, SMN, and SN were negatively correlated with DMN and it displayed a whole negative pattern. The pattern of state 5 in older may reflect a true decreasing in certain cognitive abilities.

Conclusion

The findings motivate a reconceptualization of the link between aging and FC. The previous model assumed that FC remains static throughout the resting period, neglecting the dynamic feature of the brain's functional connectivity. Our results instead suggest that these networks are, like aging process themselves, transient and dynamic. There are also some limitations to note in our study. One potential problem is that although we provided robust analysis, the direct relationship between aging and brain state requires further confirmation, perhaps utilizing the independent sample to verify the reliability of the results. Secondly, using fMRI data alone, it is not possible to determine whether network changed as aging increased structure connectivity

between brain regions, or whether the topological changes were merely a necessary temporary state. Besides, using functional imaging across development are age-associated motion artifacts and physiological signals (Power et al., 2012; Lahmann et al., 2017; Geerligs, et al., 2017), however, the impact of these factors on dynamic FC can be minimized by larger sampling (Zuo & Xing, 2014; Zuo et al., 2014), rigorous head motion control (Laumann et al., 2016; Siegel et al., 2017) or simultaneous psychological recordings. Finally, we plan to investigate the underlying significance of states in more detail using methods such as time-varying analysis. Future work would focus on depicting the cognitive symbolize of connectivity state. There is a lot of useful information that we can learn from characterizing the network properties of each state individually; however, we are left with properties across multiple states.

Acknowledgments

This research was supported by the National Natural Science Foundation of China (31470981; 31571137; 31500885), National Outstanding young people plan, the Program for the Top Young Talents by Chongqing, the Fundamental Research Funds for the Central Universities (SWU1509383,SWU1509451,SWU1609177), Natural Science Foundation of Chongqing (cstc2015jcyjA10106), Fok Ying Tung Education Foundation (151023) , General Financial Grant from the China Postdoctoral Science Foundation (2015M572423, 2015M580767), Special Funds from the Chongqing Postdoctoral Science Foundation (Xm2015037, Xm2016044), Key research for Humanities and social sciences of Ministry of Education (14JJD880009).

Reference

Abrol, A., Chaze, C., Damaraju, E., & Calhoun, V. D. (2016). The chronnectome: Evaluating replicability of dynamic connectivity patterns in 7500 resting fMRI datasets. In *Engineering in Medicine and Biology Society (EMBC), 2016 IEEE 38th Annual International Conference of the* (pp. 5571-5574). IEEE.

Achard, S., & Bullmore, E. (2007). Efficiency and cost of economical brain functional networks. *PLoS computational biology*, 3(2), e17.

Aggarwal CC, Hinneburg A, Keim DA. On the Surprising Behavior of Distance Metrics in High Dimensional Space. In: Bussche J, Vianu V, eds. *Database Theory* — ICDT (2001): 8th International Conference London, UK, January 4–6, 2001 Proceedings. Berlin, Heidelberg: Springer Berlin Heidelberg; 2001:420-434.

Alexander-Bloch, A.F., Vertes, P.E., Stidd, R., Lalonde, F., Clasen, L., Rapoport, J., Giedd, J., Bullmore, E.T., Gogtay, N., (2013). The anatomical distance of functional connections predicts brain network topology in health and schizophrenia. *Cereb. Cortex* 23, 127–138..

Allen EA, Erhardt EB, Damaraju E, Gruner W, Segall JM, Silva RF, Havlicek M, Rachakonda S, Fries J, Kalyanam R et al. (2011). A baseline for the multivariate comparison of resting-state networks. *Front Syst Neurosci*. 5.

Allen EA, Damaraju E, Plis SM, Erhardt EB, Eichele T, Calhoun VD., (2014). Tracking whole-brain connectivity dynamics in the resting state. *Cereb Cortex* 24:663–676.

Allen EA, Damaraju E, Eichele T, Wu L, Calhoun VD., (2017). EEG Signatures of Dynamic Functional Network Connectivity States. *Brain Topogr*.

Betzel, R. F., Byrge, L., He, Y., Goñi, J., Zuo, X. N., & Sporns, O. (2014). Changes in structural and functional connectivity among resting-state networks across the human lifespan. *Neuroimage, 102*

Pt 2(2), 345.

Biswal B, Yetkin FZ, Haughton VM, Hyde JS, (1995): Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magn Reson Me*d 34:537–541.

Biswal,B.B.,Mennes,M.,Zuo,X.-N.,Gohel,S.,Kelly,C.,Smith,S.M.,et al., (2010). Toward discovery science of human brain function. *Proc.Natl.Acad.Sci.USA* 107, 4734–4739.doi:10.1073 /pnas.0911855107

Calhoun VD, Miller R, Pearlson G, Adali T., (2014): The chronnectome: Time-varying connectivity networks as the next frontier in fMRI data discovery. *Neuron* 84:262–274.

Cao, M., Wang, J. H., Dai, Z. J., Cao, X. Y., Jiang, L. L., Fan, F. M., ... & Milham, M. P. (2014). Topological organization of the human brain functional connectome across the lifespan. *Developmental cognitive neuroscience*, 7, 76-93.

Chan, M. Y., Park, D. C., Savalia, N. K., Petersen, S. E., & Wig, G. S. (2014). Decreased segregation of brain systems across the healthy adult lifespan. Proceedings of the National Academy of Sciences, 111(46), E4997-E5006.

DA, Dosenbach NU, Church JA, Cohen AL, Brahmbhatt S, Miezin FM, Barch DM, Raichle ME, Petersen SE, Schlaggar BL (2007) Development of distinct control networks through segregation and integration. *Proc Natl Acad Sci* U S A 104:13507–13512. CrossRef Medline

Damoiseaux, J.S., Beckmann, C.F., Arigita, E.J.S., Barkhof, F., Scheltens, P., Stam, C. J., et al. (2008). Reduced resting-state brain activity in the "default network" in normal aging. *Cereb.Cortex* 18, 1856–1864.doi:10.1093/cercor/ bhm207

DuPre, E., & Spreng, R. N. (2016). Structural covariance networks across the lifespan, from 6-94 years of age. *bioRxiv*, 090233.

Evers, E.A.T., Klaassen, E.B., Rombouts, S.A., Backes, W.H., & Jolles, J. (2012). The effects of sustained cognitive task performance on subsequent resting state functional connectivity in healthy young and middle-aged male school teachers. *Brain Connect*. 2, 102–112.doi:10.1089/brain.2011.0060Friston KJ, Williams S, Howard R, Frackowiak RS, Turner R. (1996). Movement-related effects in fMRI time-series. *Magn Reson Med*. 35(3):346–355.

Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME (2005): The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci* USA 102:9673–9678.

Geerligs, L., Tsvetanov, K. A., & Henson, R. N. (2017). Challenges in measuring individual differences in functional connectivity using fMRI: The case of healthy aging. *Human Brain Mapping*.

Gong, Y. X. (1992). Chinese version of the Wechsler Adult Intelligence Scale (WAIS-RC). *Changsha: Hunan Map Publishing House*, *35150*.

Gonzalez-Castillo, J., Hoy, C. W., Handwerker, D. A., Robinson, M. E., Buchanan, L. C., Saad, Z. S., & Bandettini, P. A. (2015). Tracking ongoing cognition in individuals using brief, whole-brain functional connectivity patterns. *Proceedings of the National Academy of Sciences*, 112(28), 8762-8767.

Hutchison RM, Womelsdorf T, Allen EA, Bandettini PA, Calhoun VD, Corbetta M, Della Penna S, Duyn JH, Glover GH, Gonzalez-Castillo J, Handwerker DA, Keilholz S, Kiviniemi V, Leopold DA, de Pasquale F, Sporns O, Walter M, Chang C. , (2013). Dynamic functional connectivity: promise, issues, and interpretations. *Neuroimage* 80:360 –378.

Hoogendam, Y. Y., Hofman, A., van der Geest, J. N., van der Lugt, A., & Ikram, M. A. (2014).

Patterns of cognitive function in aging: the Rotterdam Study. European journal of epidemiology, 29(2), 133-140.

Kamen A. Tsvetanov, Richard N.A. Henson, Lorraine K. Tyler, Adeel Razi, Linda Geerligs, Timothy E. Ham, &James B. Rowe, (2016). Extrinsic and Intrinsic Brain Network Connectivity Maintains Cognition across the Lifespan Despite Accelerated Decay of Regional Brain Activation. *J. Neurosci.*, March 16, 2016 • 36(11):3115–3126

Kaufman, A. S., & Lichtenberger, E. O. (2005). Assessing adolescent and adult intelligence. *John Wiley & Sons.*

Laumann, T. O., Snyder, A. Z., Mitra, A., Gordon, E. M., Gratton, C., Adeyemo, B., & McCarthy, J. E. (2016). On the stability of bold fmri correlations. *Cerebral Cortex.*

Lehmann, B. C., White, S. R., Henson, R. N., & Geerligs, L. (2017). Assessing dynamic functional connectivity in heterogeneous samples. *NeuroImage*.

Leonardi, N., & Van De Ville, D. (2015). On spurious and real fluctuations of dynamic functional connectivity during rest. *Neuroimage*, 104, 430-436.

Liu, F., Wang, Y., Li, M., Wang, W., Li, R., Zhang, Z., & Chen, H. (2016). Dynamic functional network connectivity in idiopathic generalized epilepsy with generalized tonic-clonic seizure. *Human brain mapping*.

Liu, W., Wei, D., Chen, Q., Yang, W., Meng, J., Wu, G., ... & Qiu, J. (2017). Longitudinal test-retest neuroimaging data from healthy young adults in southwest China. *Scientific Data*, *4*.

Liu X, Chang C, Duyn JH. (2013). Decomposition of spontaneous brain activity into distinct fMRI co-activation patterns. *Front Syst Neurosci* 7:101. CrossRef Medline

Lloyd S. (1982). Least squares quantization in PCM. *IEEE Transaactions Information Theory*, vol

IT-28, 1982; 28:129-137.

Power JD, Fair DA, Schlaggar BL, Petersen SE., (2010). The development of human functional brain networks. *Neuron* 67:735–748. CrossRef Medline

Power JD, Mitra A, Laumann TO, Snyder AZ, Schlaggar BL, Petersen SE. (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage*. 84:320–341.

Roser Sala-Llonch, David Bartrés-Faz and Carme Junqué., (2015). Reorganization of brain networks in aging: a review of functional connectivity studies. *Front. Psychol.*, 21 May 2015 | https://doi.org/10.3389/fpsyg.2015.00663

Roweis S. (1998). EM algorithms for PCA and SPCA. *Neural Inform Process Syst*. 10:626–632.

Sala-Llonch, R., Junqué, C., Arenaza-Urquijo, E. M., Vidal-Piñeiro, D., Valls-Pedret, C., Palacios, E. M., ... & Bartrés-Faz, D. (2014). Changes in whole-brain functional networks and memory performance in aging. *Neurobiology of aging*, *35*(10), 2193-2202.

Siegel, J. S., Mitra, A., Laumann, T. O., Seitzman, B. A., Raichle, M., Corbetta, M., & Snyder, A. Z. (2016). Data quality influences observed links between functional connectivity and behavior. *Cerebral Cortex*.

Smith SM, Fox PT, Miller KL, Glahn DC, Fox PM, Mackay CE, Filippini N, Watkins KE, Toro R, Laird AR. (2009). Correspondence of the brain's functional architecture during activation and rest. *Proc Natl Acad Sci*. 106:13040–13045.

Song, J., Birn, R. M., Boly, M., Meier, T. B., Nair, V. A., Meyerand, M. E., & Prabhakaran, V. (2014). Age-related reorganizational changes in modularity and functional connectivity of human brain networks. *Brain connectivity*, 4(9), 662-676.

Thomason ME, Chang CE, Glover GH, Gabrieli JD, Greicius MD, Gotlib IH (2008). Default-mode

function and task-induced deactivation have overlapping brain substrates in children. *Neuroimage* 41:1493–1503. CrossRef Medline

Vidal-Piñeiro, D., Valls-Pedret, C., Fernández-Cabello, S., Arenaza-Urquijo, E. M., Sala-Llonch, R., Solana, E., & Bartrés-Faz, D. (2014). Decreased default mode network connectivity correlates with age-associated structural and cognitive changes. Frontiers in aging neuroscience, 6, 256.Fair

Wechsler, D., (1981). Scale-Revised, W. A. I. The Psychological Corporation. New York, 1, 309

Whitfield-Gabrieli, S., &Nieto-Castanon, A. (2012). Conn: A functional connectivity toolbox for correlated and anticorrelated brain networks. *Brain Connect*, 2, 125-141

Yan CG, Cheung B, Kelly C, Colcombe S, Craddock RC, Di Martino A, Li Q, Zuo XN, Castellanos FX, Milham MP. (2013). A comprehensive assessment of regional variation in the impact of head micromovements on functional connectomics. *Neuroimage*. 76:183–201.

Yan, C.G., Wang, X.D., Zuo, X.N., Zang, Y.F., (2016). DPABI: Data Processing & Analysis for (Resting-State) Brain Imaging. *Neuroinformatics* 14, 339-351. doi:10.1007/s12021-016-9299-4.

Zuo, X. N., Anderson, J. S., Bellec, P., Birn, R. M., Biswal, B. B., & Blautzik, J., et al. (2014). An open science resource for establishing reliability and reproducibility in functional connectomics. *Scientific Data,1*, 140049.

Zuo, X. N., He, Y., Betzel, R. F., Colcombe, S., Sporns, O., & Milham, M. P. (2017). Human Connectomics across the Life Span. *Trends in Cognitive Sciences*.

Zuo, X. N., & Xing, X. X. (2014). Test-retest reliabilities of resting-state fmri measurements in human brain functional connectomics: a systems neuroscience perspective. *Neuroscience & Biobehavioral Reviews, 45*, 100.

Tables & Figures

Table 1. The state index expressed at rest was correlated with age

State index	r	p
Total dwell time	0.20	0.000
Total transition time	-0.20	0.000
Frequency of state 1	0.32	0.000
Frequency of state 3	-0.238	0.000
Frequency of state 4	-0.19	0.000
Frequency of state 5	0.11	0.022
Transition times of 1-5	0.19	0.000
Transition times of 2-3	-0.15	0.002
Transition times of 3-2	-0.12	0.015
Transition times of 3-4	-0.20	0.000
Transition times of 3-5	-0.17	0.000
Transition times of 4-3	-0.21	0.000
Transition times of 5-1	0.16	0.001
Transition times of 5-3	-0.14	0.003
Head motion	0.41	0.000

Table 2. Global efficiency and local efficiency of each age group

Age range	N	Global efficiency	DMN Local efficiency	CC Local efficiency	SN Local efficiency
19-30	149	0.351	0.601	0.603	0.616
31-40	32	0.347	0.598	0.580	0.599
41-50	63	0.341	0.588	0.577	0.607
51-60	103	0.334	0.576	0.575	0.571
61-70	59	0.328	0.566	0.552	0.553

71-80	28	0.325 94	0.578 11	0.562 16	0.534 79
Numbers of Significant results (in 100 times)					

Table 3. Graphic indexes of each state

State Indexes	State 1	State 2	State 3	State 4	State 5
Global efficiency	0.1938	0.2069	0.2388	0.3049	0.2005
DMN local efficiency	0.3442	0.3025	0.3830	0.5334	0.4610
CCN local efficiency	0.2932	0.2316	0.3929	0.4190	0.3625
SN local efficiency	0.3847	0.4412	0.5327	0.6566	0.4454
Hub node	VIS(71,28), DMN(52)	VIS(71,55, 28,54)	VIS(71,28), AUD(51,63)	VIS(28,71, 55)	AUD(63), VIS(28), CCN(47)
N	25879	10111	28300	10641	16209

IC labels: 28, 47, 51, 52, 54, 55, 63, 71

Figure 1

Figure 1. Composite maps of the 59 identified intrinsic connectivity networks (ICNs),

sorted into seven sub-networks. Each color in the composite maps corresponds to a

different ICN.

Figure 2

Figure 2. Static functional network connectivity matrix of ICNs during rest computed over the entire scan length and averaged over subjects.

Figure 3

Figure 3. The state assignments as a function of time for the 4 example subjects. Examples of FC dynamics for subject 34, subject 85, subject 185, subject 318.

Figure 4

Figure 4. The distribution of each state in each age range.

Figure 5

Figure 5. The varying curve of different age range in the time course. The global efficiency of

young age range was always higher than it of the older age range in whole time course. The DMN

local efficiency of elderly reached its peak at the beginning, while it of young reached its peak in

the intermediate process. The CCN local efficiency of elderly reached its peak at the beginning,

while it of young reached its peak in the second half. The SN local efficiency of elderly maintained

a relative low level, while it of young had been growing up.

Figure 6.

Figure 6. The centroids of the 5 dFNC states. In state 1, the whole network displayed slight and

moderate negative connectivity. State 2 showed the high positive correlation among AUD, SMN,

and VIS, while the negative correlation between SCN and other networks. In state 3, there are

negative connections between DMN and other networks as well as SCN and others. In state 4, the

whole network displayed slight and moderate positive connectivity, and the high positive coupling

among AUD, SMN, and VIS appeared again. In state 5, AUD, SMN, and SN were negatively

correlated with DMN and it displayed a whole negative pattern.