Short report

Host shifts result in parallel genetic changes when viruses adapt to closely related species

Ben Longdon[*1,2]**, Jonathan P Day**[2]**, Joel M Alves**[2,3]**, Sophia CL Smith**[2]**, Thomas M Houslay**[1]**, John E McGonigle**[2]**, Lucia Tagliaferri**[2] **and Francis M Jiggins**[2]

[1] Biosciences, College of Life & Environmental Sciences, University of Exeter, Penryn Campus, TR10 9FE, UK

[2] Department of Genetics, University of Cambridge, Cambridge, CB2 3EH, UK

[3] CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Universidade do Porto, 4485-661 Vairão, Portugal

*b.longdon2@exeter.ac.uk

Abstract

Host shifts, where a pathogen jumps into a new host species, are a major source of emerging infectious diseases. They frequently occur between related host species and often rely on the pathogen evolving adaptations that increase their fitness in the novel host species. To investigate genetic adaptation to novel hosts, we experimentally evolved replicate lineages of an RNA virus (Drosophila C Virus) in 19 different species of Drosophilidae and deep sequenced the viral genomes. We found a strong pattern of parallel evolution, where viral lineages from the same host were genetically more similar to each other than to lineages from other host species, indicating species-specific adaptation. When we compared viruses that had adapted to different host species, we found that parallel genetic changes were more likely to occur if the two host species were closely related. This suggests that when a virus adapts to one host it might also become better adapted to closely related host species. This may explain in part why host shifts tend to occur between related species, and may mean that when a new pathogen appears in a given species, closely related species may become vulnerable to the new disease.

Introduction

Host shifts – where a pathogen jumps into a new host species – are a major source of emerging infectious diseases. RNA viruses seem particularly prone to host shift [1-4], with HIV, Ebola virus and SARS coronavirus all having been acquired by humans from other host species [5-7]. Whilst some pathogens may be pre-adapted to a novel host, there is increasing evidence that adaptation to the new host is frequently essential for a host shift to succeed [8]. These adaptations may allow a pathogen to enter host cells, increase replication rates, avoid or suppress the host immune response, or optimise virulence or transmission [9, 10]. For example, in the 2013-2016 Ebola virus epidemic in West Africa, a mutation in the viral glycoprotein gene that arose early in the outbreak and rose to high frequency was found to increase infectivity in human cells, and decrease infectivity in bats, which are thought to be the source of Ebola virus [11, 12]. Likewise, a switch of a parvovirus from cats to dogs resulted in mutations in the virus capsid that allowed the virus to bind to cell receptors in dogs, but resulted in the virus losing its ability to infect cats [13].

In some instances adaptation to a novel host relies on specific mutations that arise repeatedly whenever a pathogen switches to a given host. For example, in the jump of HIV-1 from chimps to humans, codon 30 of the *gag* gene has undergone a change that increases virus replication in humans, and this has occurred independently in all three HIV-1 lineages [5, 14]. Similarly, five parallel mutations have been observed in the two independent epidemics of SARS coronavirus following its jump from palm civets into humans [15]. Similar patterns have been seen in experimental evolution studies, where parallel genetic changes occur repeatedly when replicate viral lineages adapt to a new host species in the lab. For example, when Vesicular Stomatitis Virus was passaged in human or dog cells, the virus evolved parallel mutations when evolved on the same cell type [16]. Likewise, a study passaging Tobacco Etch Potyvirus on four plant species found parallel mutations occurred only when the virus infected the same host species [17]. These parallel mutations provide compelling evidence that these genetic changes are adaptive, with the same mutations evolving independently in response to natural selection [18].

The host phylogeny is important for determining a pathogens ability to infect a novel host, with pathogens tending to replicate most efficiently when they infect a novel host that is closely related to their original host. This pattern is seen in nature, where host shifts tend to occur most frequently between closely related hosts, and in laboratory cross-infection studies, where viruses tend to replicate more rapidly when the new host is related to the pathogens natural host [2, 19-32].

These phylogenetic patterns suggest that mutations that adapt a virus to one host may also adapt it to closely related host species. Therefore, there may be a knock-on effect, where a host shift leaves closely related species vulnerable to the new disease. To investigate this, we asked whether viruses acquire the same genetic adaptations when adapting to closely related host species. We experimentally evolved replicate lineages of an RNA virus called Drosophila C Virus (DCV; Discistroviridae) in 19 species of Drosophilidae that vary in their relatedness and shared a common ancestor approximately 40 million years ago [33, 34]. We then sequenced the genomes of the evolved viral lineages and tested whether the same genetic changes arose when the virus was evolved in closely related host species.

91
92 **Results**
93
94 **Parallel genetic changes occur in DCV lineages that have evolved in the same host species**
95
To examine how viruses adapt to different host species we experimentally evolved DCV in 19 species
of Drosophilidae by serially passaging the virus. In total we infected 22,095 adult flies and generated
173 independent replicate lineages (6-10 per host species). We deep sequenced the evolved virus
genomes to generate over 740,000 300bp sequence reads from each viral lineage. Out of 8989 sites,
584 contained a SNP with a derived allele frequency >0.05 in at least one viral lineage, and 84 of
these were tri-allelic. None of these variants were found at an appreciable frequency in five
sequencing libraries produced from the ancestral virus, indicating that they had spread though
populations during the experiment (Figure 1). In multiple cases these variants had nearly reached
fixation (Figure 1).
105
A powerful way to detect adaptation is to test whether the same genetic changes occur in parallel
when different populations encounter the same environment. Of the 584 SNPs, 102 had derived
allele frequencies >0.05 in at least two viral lineages, and some had risen to high frequencies in
multiple lineages (Figure 1). We estimated the genetic differentiation between viral lineages by
calculating F_{ST}. We found that viral lineages that had evolved within the same host were genetically
more similar to each other than to lineages from other host species (Figure 2; $P<0.001$). The
observation of parallel evolution within the same host species indicates that the viruses are
genetically adapting to the different host species.
114
115 To examine the genetic basis of parallel evolution, we individually tested whether each SNP in the
116 DCV genome showed a signature of parallel evolution among viral lineages passaged in the same
117 host species (i.e. we repeated the analysis in Figures 2 for each SNP). We identified 56 SNPs with a
118 significant signal of parallel evolution within the same host species ($P<0.05$; significantly parallel
119 SNPs are shown with a red asterisk in Figure 1; the false discovery rate is estimated to be 17% [35]).
120

Ancestral virus
D. prosaltans
D. saltans
S.pattersoni
S.lebanonesis
D. euronotus
D. paramelanica
D. mojavensis
D. arizonae
D. montana
D. americana
D. lummei
D. teissieri
D. santomea
D. sechellia
D. simulans
D. mauritiana
D. melanogater
D. pseudoobscura
D. obscura

ORF1   ORF2

Position in DCV genome

136
137 **Figure 2. Viral lineages from the same host species were genetically more similar to each other**
138 **than to lineages from different host species.** The mean pairwise F_{ST} between all possible pairs of
139 viral lineages from the same host species was calculated. The red line shows the observed value. The
140 grey bars are the null distribution of this statistic obtained by permuting the viral lineages across
141 host species 1000 times.
142
143 **Viruses in closely related hosts are genetically more similar**
144
We investigated if viruses passaged through closely related hosts showed evidence of parallel
molecular adaptations. We calculated F_{ST} between all possible pairs of viral lineages that had evolved
in different host species. We found that viral lineages from closely related hosts were more similar
to each other than viral lineages from more distantly related hosts (Figure 3A). This is reflected in a
significant positive relationship between virus F_{ST} and host genetic distance (Figure 3B, Permutation
test: *r*=0.15, *P*=0.002). We lacked the statistical power to identify the specific SNPs that are causing
the signature of parallel evolution in Figure 3 (false discovery rate >0.49 for all SNPs).
152

Figure 3. Viral lineages from more closely related host species are genetically more similar. (A) The
correlation between the genetic differentiation of viral lineages and the genetic distance between
the species they have evolved in. Linear regression line is shown in red. This plot includes
comparisons between viral lineages that evolved in the same host species (genetic distance = 0).
Genetic distances were scaled so that the distance from the root to the tip of the tree was one. (B)
Pearson's correlation coefficient (r) of F_{ST} between pairs of viral lineage and the genetic distance
between the host species they evolved in. The observed value is in red and the grey bars are the null
distribution obtained by permutation. This analysis did not include comparisons between viral
lineages from the same host species

Discussion

When a pathogen infects a novel host species, it finds itself in a new environment to which it must
adapt [4, 9, 41, 42]. When DCV was passaged through different species of Drosophilidae, we found
the same genetic changes arose repeatedly in replicate viral lineages in the same host species. Such
repeatable parallel genetic changes to the same host environment are compelling evidence that
these changes are adaptive [18]. We then examined whether these same genetic changes might
adapt the virus to closely related host species, as these are likely to present a similar environment
for the virus. We found that viruses evolved in closely related hosts were more similar to each other
than viruses that evolved in more distantly related species. Therefore, mutations that are adaptive in
one host species frequently arise when the virus infects closely related hosts. This finding of parallel
genetic changes in closely related host species suggests that when a virus adapts to one host it might
also become better adapted to closely related host species.

Such phylogenetic patterns of host adaptation may in part explain why pathogens tend to be more
likely to jump between closely related host species. At a phenotypic level, similar patterns have been
reported in nature [19, 27]. Likewise, in a large cross-infection experiment Drosophila sigma viruses
(Rhabdoviridae) isolated from different species of *Drosophila*, the viruses tended to replicate most
efficiently in species closely related to their natural hosts [32]. This suggests that these viruses had
acquired adaptations to their host species that benefitted them when they infected closely related
species. Our results demonstrate that this pattern is apparent at the level of specific nucleotides,
and can arise very shortly after a host shift.

While the susceptibility of a novel host is strongly correlated to its relatedness to the pathogens'
original host, it is also common to find exceptions to this pattern. This is seen both in nature when
pathogens shift between very distant hosts [43, 44], and in laboratory cross-infection experiments
[31, 32]. This pattern is also seen in our data where we also observe parallel genetic changes
occurring between more distantly related hosts. For example, a mutation at position 8072 was not
only near fixation in most of the lineages infecting two closely related species, but also occurred at a
high frequency in replicate lineages in a phylogenetically distant host (Figure 1).

In conclusion, we have found that host relatedness can be important in determining how viruses
evolve when they find themselves in a new host. This study suggests that while some genetic
changes will be found only in specific hosts, we frequently see the same changes occurring in closely

related host species. This could mean that mutations that adapt a virus to one host might also adapt
it to closely related species.
201
202
203 **Methods**
204
205 *Virus production*
206
DCV is a positive sense RNA virus in the family Discistroviridae that was isolated from *D.*
melanogaster, which it naturally infects in the wild [45, 46]. To minimise the amount of genetic
variation in the DCV isolate we used to initiate the experimental evolution study, we aimed to isolate
single infectious clones of DCV using a serial dilution procedure. DCV was produced in Schneider's
Drosophila line 2 (DL2) cells [47] as described in [48]. Cells were cultured at 25°C in Schneider's
Drosophila Medium with 10% Fetal Bovine Serum, 100 U/ml penicillin and 100 µg/ml streptomycin
(all Invitrogen, UK). The DCV strain used was isolated from *D. melanogaster* collected in Charolles,
France [49]. DL2 cells were seeded into two 96-well tissue culture plates at approximately 10^4 cells in
100 µl of media per well. Cells were allowed to adhere to the plates by incubating at 25°C for five
hours or over-night. Serial 1:1 dilutions of DCV were made in complete Schneider's media, giving a
range of final dilutions from $1:10^8 - 1:4 \times 10^{14}$. 100 µl of these dilutions were then added to the cells
and incubated for 7 days, 8 replicates were made for each DCV dilution. Each well was then
examined for DCV infection of the DL2 cells, and a well was scored as positive for DCV infection if
clear cytopathic effects were present in the majority of the cells. The media was taken from the
wells with the greatest dilution factor that were scored as infected with DCV and stored at -80°C.
This processes was then repeated using the DCV samples from the first dilution series. One clone,
B6A, was selected for amplification and grown in cell culture as described above. Media containing
DCV was removed and centrifuged at 3000 x g for 5 minutes at 4°C to pellet any remaining cell
debris, before being aliquoted and stored at -80°C. The Tissue Culture Infective Dose 50 ($TCID_{50}$) of
the DCV was 6.32×10^9 infectious particles per ml using the Reed-Muench end-point method [50].
227
228 *Inoculating fly species*
229
We passaged the virus through 19 species of Drosophilidae, with 6-10 independent replicate
passages for each species. We selected species from across the phylogeny (that shared a common
ancestor approximately 40 million years ago [33, 34]), but included clades of closely related species
that recently shared common ancestors less than 5 million years ago (Figure 1). All fly stocks were
reared at 22°C. Stocks of each fly species were kept in 250ml bottles at staggered ages. Flies were
collected and sexed, and males were placed on cornmeal medium for 4 days before inoculation.
Details of the fly stocks used can be found in the supplementary materials.
237
4-11 day old males were infected with DCV using a 0.0125 mm diameter stainless steel needle
(26002–10, Fine Science Tools, CA, USA) dipped in DCV solution. For the first passage this was the
cloned DCV isolate in cell culture supernatant (described above), and then subsequently was the
virus extracted from the previous passage (described below). The needle was pricked into the
pleural suture on the thorax of flies, towards the midcoxa. Each replicate was infected using a new
needle and strict general cleaning procedures were used to minimise any risk of cross-contamination

between replicates. Species were collected and inoculated in a randomised order each passage. Flies
were then placed into vials of cornmeal medium and kept at 22°C and 70% relative humidity. Flies
were snap frozen in liquid nitrogen 3 days post-infection, homogenised in Ringer's solution (2.5μl
per fly) and then centrifuged at 12,000g for 10 mins at 4°C. The resulting supernatant was removed
and frozen at -80°C to be used for infecting flies in the subsequent passage. The remaining
homogenate was preserved in Trizol reagent (Invitrogen) and stored at -80°C for RNA extraction.
The 3 day viral incubation period was chosen based on time course and pilot data showing that viral
load reaches a maximum at approximately 3 days post-infection. This process was repeated for 10
passages for all species, except *D. montana* where only 8 passages were carried out due to the fly
stocks failing to reproduce. Each lineage was injected into a mean of 11 flies at each passage (range
4-18). Experimental evolution studies in different tissue types have seen clear signals of adaptation
in 100 virus generations [16]. Based on $\log_2$ change in RNA viral load we estimate that we have
passaged DCV for approximately 100-200 generations.

Sequencing

After passaging the virus, we sequenced evolved viral lineages from 19 host species, with a mean of
9 independent replicate lineages of the virus per species (range 6-10 replicates). cDNA was
synthesised using Invitrogen Superscript III reverse-transcriptase with random hexamer primers
(25°C 5mins, 50°C 50mins, 70°C 15mins). The genome of the evolved viruses, along with the initial
DCV ancestor (x5) were then amplified using Q5 high fidelity polymerase (NEB) in nine overlapping
PCR reactions (see supplementary Table S2 for PCR primers and cycle conditions). Primers covered
position 62-9050bp (8989bp) of the Genbank refseq (NC_001834.1) giving 97% coverage of the
genome. PCRs of individual genomes were pooled and purified with Ampure XP beads (Agencourt).
Individual Nextera XT libraries (Illumina) were prepared for each viral lineage. In total we sequenced
173 DCV pooled amplicon libraries on an Illumina MiSeq (Cambridge Genomic Service) v3 for 600
cycles to give 300bp paired-end reads.

Bioinformatics and variant calling

FastQC, version 0.11.2 [51] was used to assess read quality and primer contamination. Trimmomatic,
version 0.32 [52] was used to removed low quality bases and adaptor sequences, using the following
options: MINLEN=30 (Drop the read if it is below 30 base pairs), TRAILING=15 (cut bases of the end
of the read if below a threshold quality of 15), SLIDINGWINDOW=4:20 (perform a sliding window
trimming, cutting once the average quality within a 4bp window falls below a threshold of 20), and
ILLUMINACLIP=TruSeq3-PE.fa:2:20:10:1:true (remove adapter contamination; the values correspond
in order to: input fasta file with adapter sequences to be matched, seed mismatches, palindrome clip
threshold, simple clip threshold, minimum adapter length and logical value to keep both reads in
case of read-through being detected in paired reads by palindrome mode).

To generate a reference ancestral Drosophila C Virus sequence we amplified the ancestral starting
virus by PCR as above. PCR products were treated with exonuclease 1 and Antarctic phosphatase to
remove unused PCR primers and dNTPs and then sequenced directly using BigDye reagents (ABI) on
an ABI 3730 capillary sequencer in both directions (Source Bioscience, Cambridge, UK). Sequences
were edited in Sequencer (version 4.8; Gene Codes), and were manually checked for errors. Fastq

reads were independently aligned to this reference sequence (Genbank accession: MG570143) using
BWA-MEM, version 0.7.10 {Li, 2009 #1605} with default options with exception of the parameter –
M, which marks shorter split hits as secondary. 99.5% of reads had mapping phred quality scores of
>60. The generated SAM files were converted to their binary format (BAM) and sorted by their
leftmost coordinates with SAMtools, version 0.1.19 (website: http://samtools.sourceforge.net/) [53].
Read Group information (RG) was added to the BAM files using the module
AddOrReplaceReadGroups from Picard Tools, version 1.126 (https://broadinstitute.github.io/picard).

The variant calling was then performed for each individual BAM using UnifiedGenotyper tool from
GATK, version 3.3.0. As we were interested in calling low frequency variants in our viruses, we
assumed a ploidy level of 100 (-sample_ploidy:100). The other parameters were set to their defaults
except --stand_call_conf:30 (minimum phred-scaled confidence threshold at which variants should
be called) and --downsample_to_coverage:1000 (down-sample each sample to 1000X coverage)

Host phylogeny

We used a trimmed version of an ultrametric (time-based) phylogeny produced previously [31]. This
phylogeny was inferred using seven genes with a relaxed molecular clock model in BEAST (v1.8.0)
[40, 54]. The tree was pruned to the 19 species used using the Ape package in R [55, 56].

Statistical Analysis
304
We examined the frequency of alternate alleles (single nucleotide polymorphisms: SNPs) in five
ancestral virus replicates (aliquots of the same virus stock that was used to found the evolved
lineages). SNPs in these ancestral viruses may represent pre-standing genetic variation, or may be
sequencing errors. We found the mean SNP frequency was 0.000923 and the highest frequency of
any SNP was 0.043 across the ancestral viruses. We therefore included a SNP in our analyses if its
frequency was >0.05 in any of the evolved viral lineages. For all analyses we included all three alleles
at triallelic sites.
312
313 *Parallel evolution within species*
314
As a measure of genetic differentiation we estimated F_{ST} between all the virus lineages based on the
heterozygosity (*H*) of the SNPs we called [57]:

$$F_{ST} = \frac{H_b - H_w}{H_b} \qquad \text{(Equation 1)}$$

where H_b is the mean number of differences between pairs of sequence reads sampled from the two
different lineages. H_w is mean number of differences between sequence reads sampled from within
each lineage. H_b and H_w were calculated separately for each polymorphic site, and the mean across
sites used in equation (1). H_w was calculated separately for the two lineages being compared, and
the unweighted mean used in equation (1).
323
To examine whether there had been parallel evolution among viral lineages that had evolved within
the same fly species, we calculated the mean F_{ST} between lineages that had evolved in the same fly
species, and compared this to the mean F_{ST} between lineages that had evolved in different fly

species. We tested whether this difference was statistically significant using a permutation test. The
fly species labels were randomly reassigned to the viral lineages, and we calculated the mean F_{ST}
between lineages that had evolved in the same fly species. This was repeated 1000 times to
generate a null distribution of the test statistic, and this was then compared to the observed value.
331
332   To identify individual SNPs with a signature of parallel evolution within species, we repeated this
333   procedure separately for each SNP.
334
335   *Parallel evolution between species*
336
We next examined whether viral lineages that had evolved in different fly species tended to be more
similar if the fly species were more closely related. Considering all pairs of viral lineages from
different host species, we correlated pairwise F_{ST} with the genetic distance between the fly species.
To test the significance of this correlation, we permuted the fly species over the *Drosophila*
phylogeny and recalculated the Pearson correlation coefficient. This was repeated 1000 times to
generate a null distribution of the test statistic, and this was then compared to the observed value.
To identify individual SNPs whose frequencies were correlated with the genetic distance between
hosts we repeated this procedure separately for each SNP.
345
346   We confirmed there was no relationship between rates of molecular evolution (SNP frequency) and
347   either genetic distance from the host DCV was isolated from (*D. melanogaster*) or estimated viral
348   population size (see supplementary Figures S1 and S2).
349
350   Sequence data (fastq files) are available in the NCBI SRA (Accession: SRP119720). BAM files, data and
351   R scripts for analysis are available from the NERC data repository (funding requirement - awaiting
352   doi, temporary link to data and scripts https://figshare.com/s/b119ba86def8bca58782).
353
354

355   **Author contributions**
356   Designing experiment: BL, FMJ. Lab work: JPD, SCLS, TMH, LT, BL. Bioinformatics analysis: JMA, JEM,
357   FMJ, BL. Statistical analysis: BL, FMJ. Manuscript written by BL and FMJ with input from all authors.
358

359   **Funding**
360
361   BL and FMJ are supported by a NERC grant (NE/L004232/1 http://www.nerc.ac.uk/) and by an ERC
362   grant (281668, DrosophilaInfection, http://erc.europa.eu/). BL is supported by a Sir Henry Dale
363   Fellowship jointly funded by the Wellcome Trust and the Royal Society (Grant Number
364   109356/Z/15/Z).
365

366   **References**
367
1.   Cleaveland S, Laurenson MK, Taylor LH. Diseases of humans and their domestic mammals:
pathogen characteristics, host range and the risk of emergence. Philosophical Transactions of the
Royal Society of London Series B-Biological Sciences. 2001;356(1411):991-9. PubMed PMID:
WOS:000170315900003.

2.   Davies TJ, Pedersen AB. Phylogeny and geography predict pathogen community similarity in
wild primates and humans. Proceedings of the Royal Society B-Biological Sciences.
2008;275(1643):1695-701. doi: 10.1098/rspb.2008.0284. PubMed PMID: ISI:000256387500014.
3.   Taylor LH, Latham SM, Woolhouse ME. Risk factors for human disease emergence. Philos
Trans R Soc Lond B Biol Sci. 2001;356(1411):983-9. Epub 2001/08/23. doi: 10.1098/rstb.2001.0888.
PubMed PMID: 11516376; PubMed Central PMCID: PMC1088493.
4.   Woolhouse ME, Haydon DT, Antia R. Emerging pathogens: the epidemiology and evolution
of species jumps. Trends Ecol Evol. 2005;20(5):238-44. Epub 2006/05/17. doi: S0169-5347(05)00038-
8 [pii]
10.1016/j.tree.2005.02.009. PubMed PMID: 16701375.
5.   Sharp PM, Hahn BH. The evolution of HIV-1 and the origin of AIDS. Philosophical
Transactions of the Royal Society B-Biological Sciences. 2010;365(1552):2487-94. doi:
10.1098/rstb.2010.0031. PubMed PMID: WOS:000280097000008.
6.   Leroy EM, Kumulungui B, Pourrut X, Rouquet P, Hassanin A, Yaba P, et al. Fruit bats as
reservoirs of Ebola virus. Nature. 2005;438(7068):575-6. doi: 10.1038/438575a. PubMed PMID:
WOS:000233593100030.
7.   Li WD, Shi ZL, Yu M, Ren WZ, Smith C, Epstein JH, et al. Bats are natural reservoirs of SARS-
like coronaviruses. Science. 2005;310(5748):676-9. doi: 10.1126/science.1118391. PubMed PMID:
WOS:000232997700042.
8.   Woolhouse M, Adair K, Brierley L. RNA Viruses: A Case Study of the Biology of Emerging
Infectious Diseases. Microbiol Spectr. 2013;1(1). doi: 10.1128/microbiolspec.OH-0001-2012.
PubMed PMID: 26184815.
9.   Longdon B, Brockhurst MA, Russell CA, Welch JJ, Jiggins FM. The Evolution and Genetics of
Virus Host Shifts. PLoS Pathog. 2014;10(11):e1004395. Epub 2014/11/07. doi:
10.1371/journal.ppat.1004395. PubMed PMID: 25375777; PubMed Central PMCID: PMC4223060.
10.   Parrish CR, Kawaoka Y. The origins of new pandemic viruses: the acquisition of new host
ranges by canine parvovirus and influenza A viruses. Annual review of microbiology. 2005;59:553-86.
Epub 2005/09/13. doi: 10.1146/annurev.micro.59.030804.121059. PubMed PMID: 16153179.
11.   Diehl WE, Lin AE, Grubaugh ND, Carvalho LM, Kim K, Kyawe PP, et al. Ebola Virus
Glycoprotein with Increased Infectivity Dominated the 2013-2016 Epidemic. Cell. 2016;167(4):1088-
98 e6. doi: 10.1016/j.cell.2016.10.014. PubMed PMID: 27814506; PubMed Central PMCID:
PMCPMC5115602.
12.   Urbanowicz RA, McClure CP, Sakuntabhai A, Sall AA, Kobinger G, Muller MA, et al. Human
Adaptation of Ebola Virus during the West African Outbreak. Cell. 2016;167(4):1079-87 e5. doi:
10.1016/j.cell.2016.10.013. PubMed PMID: 27814505; PubMed Central PMCID: PMCPMC5101188.
13.   Truyen U, Evermann JF, Vieler E, Parrish CR. Evolution of canine parvovirus involved loss and
gain of feline host range. Virology. 1996;215(2):186-9. Epub 1996/01/15. doi:
10.1006/viro.1996.0021. PubMed PMID: 8560765.
14.   Wain LV, Bailes E, Bibollet-Ruche F, Decker JM, Keele BF, Van Heuverswyn F, et al.
Adaptation of HIV-1 to its human host. Mol Biol Evol. 2007;24(8):1853-60. Epub 2007/06/05. doi:
10.1093/molbev/msm110. PubMed PMID: 17545188.
15.   Liu W, Tang F, Fontanet A, Zhan L, Wang TB, Zhang PH, et al. Molecular epidemiology of
SARS-associated coronavirus, Beijing. Emerg Infect Dis. 2005;11(9):1420-4. Epub 2005/10/19. doi:
10.3201/eid1109.040773. PubMed PMID: 16229772; PubMed Central PMCID: PMC3310602.
16.   Remold SK, Rambaut A, Turner PE. Evolutionary genomics of host adaptation in vesicular
stomatitis virus. Mol Biol Evol. 2008;25(6):1138-47. Epub 2008/03/21. doi: 10.1093/molbev/msn059.
PubMed PMID: 18353798.
17.   Bedhomme S, Lafforgue G, Elena SF. Multihost experimental evolution of a plant RNA virus
reveals local adaptation and host-specific mutations. Mol Biol Evol. 2012;29(5):1481-92. Epub
2012/02/10. doi: 10.1093/molbev/msr314. PubMed PMID: 22319146.

18.   Bollback JP, Huelsenbeck JP. Parallel Genetic Evolution Within and Between Bacteriophage
Species of Varying Degrees of Divergence. Genetics. 2009;181(1):225-34. doi:
10.1534/genetics.107.085225. PubMed PMID: WOS:000262595500021.
19.   Streicker DG, Turmelle AS, Vonhof MJ, Kuzmin IV, McCracken GF, Rupprecht CE. Host
Phylogeny Constrains Cross-Species Emergence and Establishment of Rabies Virus in Bats. Science.
2010;329(5992):676-9. doi: 10.1126/science.1188836. PubMed PMID: WOS:000280602700037.
20.   Faria NR, Suchard MA, Rambaut A, Streicker DG, Lemey P. Simultaneously reconstructing
viral cross-species transmission history and identifying the underlying constraints. Philos Trans R Soc
Lond B Biol Sci. 2013;368(1614):20120196. Epub 2013/02/06. doi: 10.1098/rstb.2012.0196. PubMed
PMID: 23382420; PubMed Central PMCID: PMC3678322.
21.   Cooper N, Griffin R, Franz M, Omotayo M, Nunn CL, Fryxell J. Phylogenetic host specificity
and understanding parasite sharing in primates. Ecol Lett. 2012;15(12):1370-7. Epub 2012/08/24.
doi: 10.1111/j.1461-0248.2012.01858.x. PubMed PMID: 22913776.
22.   Waxman D, Weinert LA, Welch JJ. Inferring host range dynamics from comparative data: the
protozoan parasites of new world monkeys. Am Nat. 2014;184(1):65-74. Epub 2014/06/13. doi:
10.1086/676589. PubMed PMID: 24921601.
23.   Huang S, Bininda-Emonds ORP, Stephens PR, Gittleman JL, Altizer S. Phylogenetically related
and ecologically similar carnivores harbor similar parasite assemblages. Journal of Animal Ecology.
2013:n/a-n/a. doi: 10.1111/1365-2656.12160.
24.   Hadfield JD, Krasnov BR, Poulin R, Nakagawa S. A Tale of Two Phylogenies: Comparative
Analyses of Ecological Interactions. The American Naturalist. 2014;0(0):000. doi: 10.1086/674445.
25.   Ramsden C, Holmes EC, Charleston MA. Hantavirus evolution in relation to its rodent and
insectivore hosts: no evidence for codivergence. Mol Biol Evol. 2009;26(1):143-53. Epub 2008/10/17.
doi: msn234 [pii]
10.1093/molbev/msn234. PubMed PMID: 18922760.
26.   de Vienne DM, Hood ME, Giraud T. Phylogenetic determinants of potential host shifts in
fungal pathogens. Journal of Evolutionary Biology. 2009;22(12):2532-41. doi: 10.1111/j.1420-
9101.2009.01878.x. PubMed PMID: WOS:000271785800019.
27.   Gilbert GS, Webb CO. Phylogenetic signal in plant pathogen-host range. Proceedings of the
National Academy of Sciences of the United States of America. 2007;104(12):4979-83. doi:
10.1073/pnas.0607968104. PubMed PMID: WOS:000245256700040.
28.   Tinsley MC, Majerus MEN. Small steps or giant leaps for male-killers? Phylogenetic
constraints to male-killer host shifts. Bmc Evolutionary Biology. 2007;7. doi: 10.1186/1471-2148-7-
238. PubMed PMID: WOS:000252786000001.
29.   Russell JA, Goldman-Huertas B, Moreau CS, Baldo L, Stahlhut JK, Werren JH, et al.
Specialization and geographic isolation among Wolbachia symbionts from ants and lycaenid
butterflies. Evolution. 2009;63(3):624-40. Epub 2008/12/05. doi: 10.1111/j.1558-5646.2008.00579.x.
PubMed PMID: 19054050.
30.   Perlman SJ, Jaenike J. Infection success in novel hosts: An experimental and phylogenetic
study of Drosophila-parasitic nematodes. Evolution. 2003;57(3):544-57. PubMed PMID:
WOS:000182193800010.
31.   Longdon B, Hadfield JD, Day JP, Smith SC, McGonigle JE, Cogni R, et al. The Causes and
Consequences of Changes in Virulence following Pathogen Host Shifts. PLoS Pathog.
2015;11(3):e1004728. Epub 2015/03/17. doi: 10.1371/journal.ppat.1004728. PubMed PMID:
25774803.
32.   Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM. Host phylogeny determines viral
persistence and replication in novel hosts. PLoS Pathogens. 2011;7((9)):e1002260. doi:
10.1371/journal.ppat.1002260.
33.   Obbard DJ, Maclennan J, Kim K-W, Rambaut A, O'Grady PM, Jiggins FM. Estimating
divergence dates and substitution rates in the Drosophila phylogeny. Molecular Biology and
Evolution. 2012;29(11):3459-73.

34.   Tamura K, Subramanian S, Kumar S. Temporal patterns of fruit fly (Drosophila) evolution
revealed by mutation clocks. Mol Biol Evol. 2004;21(1):36-44. Epub 2003/09/02. doi:
10.1093/molbev/msg236
msg236 [pii]. PubMed PMID: 12949132.
35.   Storey JD. A direct approach to false discovery rates. J Roy Stat Soc B. 2002;64:479-98. doi:
Unsp 1369-7412/02/64479
Doi 10.1111/1467-9868.00346. PubMed PMID: WOS:000177425500009.
36.   Jan E. Divergent IRES elements in invertebrates. Virus Res. 2006;119(1):16-28. doi:
10.1016/j.virusres.2005.10.011. PubMed PMID: 16307820.
37.   Johnson KN, Christian PD. The novel genome organization of the insect picorna-like virus
Drosophila C virus suggests this virus belongs to a previously undescribed virus family. J Gen Virol.
1998;79 (Pt 1):191-203. Epub 1998/02/14. PubMed PMID: 9460942.
38.   Nakashima N, Nakamura Y. Cleavage sites of the "P3 region" in the nonstructural polyprotein
precursor of a dicistrovirus. Arch Virol. 2008;153(10):1955-60. doi: 10.1007/s00705-008-0208-5.
PubMed PMID: 18810573.
39.   Nakashima N, Uchiumi T. Functional analysis of structural motifs in dicistroviruses. Virus Res.
2009;139(2):137-47. doi: 10.1016/j.virusres.2008.06.006. PubMed PMID: 18621089.
40.   Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and the
BEAST 1.7. Mol Biol Evol. 2012;29(8):1969-73. Epub 2012/03/01. doi: 10.1093/molbev/mss075.
PubMed PMID: 22367748; PubMed Central PMCID: PMC3408070.
41.   Parrish CR, Holmes EC, Morens DM, Park EC, Burke DS, Calisher CH, et al. Cross-species virus
transmission and the emergence of new epidemic diseases. Microbiology and Molecular Biology
Reviews. 2008;72(3):457-70. doi: 10.1128/mmbr.00004-08. PubMed PMID: WOS:000258951200004.
42.   Woolhouse ME, Gowtage-Sequeria S. Host range and emerging and reemerging pathogens.
Emerg Infect Dis. 2005;11(12):1842-7. Epub 2006/02/21. PubMed PMID: 16485468.
43.   Webby RJ, Webster RG. Emergence of influenza A viruses. Philos Trans R Soc Lond B Biol Sci.
2001;356(1416):1817-28. Epub 2002/01/10. doi: 10.1098/rstb.2001.0997. PubMed PMID: 11779380;
PubMed Central PMCID: PMC1088557.
44.   Weinert LA, Welch JJ, Suchard MA, Lemey P, Rambaut A, Fitzgerald JR. Molecular dating of
human-to-bovid host jumps by Staphylococcus aureus reveals an association with the spread of
domestication. Biol Lett. 2012;8(5):829-32. Epub 2012/05/26. doi: 10.1098/rsbl.2012.0290. PubMed
PMID: 22628096; PubMed Central PMCID: PMC3440972.
45.   Christian PD. Studies of Drosophila C and A viruses in Australian populations of Drosophila
melanogaster: Australian National University; 1987.
46.   Webster CL, Waldron FM, Robertson S, Crowson D, Ferrai G, Quintana JF, et al. The
discovery, distribution and evolution of viruses associated with Drosophila melanogaster. PLOS
Biology. 2015;13(7): e1002210.
47.   Teixeira L, Ferreira A, Ashburner M. The Bacterial Symbiont Wolbachia Induces Resistance to
RNA Viral Infections in Drosophila melanogaster. Plos Biology. 2008;6(12):2753-63. doi: DOI
10.1371/journal.pbio.1000002. PubMed PMID: ISI:000261913700016.
48.   Longdon B, Cao C, Martinez J, Jiggins FM. Previous Exposure to an RNA Virus Does Not
Protect against Subsequent Infection in Drosophila melanogaster. Plos One. 2013;8(9):e73833. doi:
10.1371/journal.pone.0073833.
49.   Jousset FX, Plus N, Croizier G, Thomas M. [Existence in Drosophila of 2 groups of picornavirus
with different biological and serological properties]. C R Acad Sci Hebd Seances Acad Sci D.
1972;275(25):3043-6. Epub 1972/12/18. PubMed PMID: 4631976.
50.   Reed LJ, Muench H. A simple method of estimating fifty per cent endpoints. The American
Journal of Hygiene. 1938;27:493-7.
51.   Andrews S. FastQC: a quality control tool for high throughput sequence data. Available at:
http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc.2010.

52.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. 2014;30(15):2114-20. doi: 10.1093/bioinformatics/btu170. PubMed PMID:
24695404; PubMed Central PMCID: PMCPMC4103590.
53.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. doi:
10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed Central PMCID:
PMCPMC2723002.
54.   Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees. BMC
Evolutionary Biology. 2007;7:214. doi: Artn 214
Doi 10.1186/1471-2148-7-214. PubMed PMID: ISI:000253468300001.
55.   Team RDC. R: a language and environment for statistical computing. V 2.4. 2006.
56.   Paradis E, Claude J, Strimmer K. APE: Analyses of Phylogenetics and Evolution in R language.
Bioinformatics. 2004;20(2):289-90. PubMed PMID: 14734327.
57.   Hudson RR, Slatkin M, Maddison WP. Estimation of levels of gene flow from DNA sequence
data. Genetics. 1992;132(2):583-9. Epub 1992/10/01. PubMed PMID: 1427045; PubMed Central
PMCID: PMC1205159.
539