Rapid, noninvasive, and unsupervised detection of sleep/wake using piezoelectric monitoring for pharmacological studies in narcoleptic mice

Sarah Wurts Black[a,*], Jessica D. Sun[a], Alex Laihsu[a], Nikki Kimura[a], Pamela Santiago[a], Kevin D. Donohue[b,c], Bruce F. O'Hara[b,d], Ross Bersot[a], Paul S. Humphries[a]

a. Reset Therapeutics, 260 Littlefield Ave, Suite 200, South San Francisco, CA 94080, United States of America

b. Signal Solutions, LLC, 145 Graham Ave., Lexington, KY 40506, United States of America

c. Department of Electrical and Computer Engineering, University of Kentucky, Lexington, KY 40506, United States of America

d. Department of Biology, University of Kentucky, Lexington, KY 40506, United States of America

* Corresponding author

Email: sallyb@resettherapeutics.com

Phone: 650-235-4136

Fax: 650-235-4151

Address: 260 Littlefield Ave, Suite 200, South San Francisco, CA 94080, USA

Type of Article: Methodology article

Running Title: PiezoSleep assay for murine narcolepsy

Pages:29   **Figures:** 4   **Tables:** 2

Abstract

Background: Assessment of sleep/wake by electroencephalography (EEG) and electromyography (EMG) is invasive, resource intensive, and not amenable to rapid screening at scale for drug discovery. In the preclinical development of therapeutics for narcolepsy, efficacy tests are hindered by the lack of a non-EEG/EMG based translational test of symptom severity. A new method of noninvasive, unsupervised piezoelectric monitoring of gross body movement, together with respiration patterns during behavioral quiescence, was used to determine sleep/wake. In the PiezoSleep narcolepsy screen, these measurements were sampled during the dark period, and the ratio of time spent in long wake bouts to short sleep bouts was used to determine a translational wake-maintenance score by which to rank narcoleptic and wild type mice by sleepiness severity. The wake-maintenance score was used to identify the most highly symptomatic mice for resource-intensive EEG/EMG studies for further study. PiezoSleep was validated for high-throughput screening of wake-promoting compounds using modafinil in narcoleptic and wild type mice.

Results: The PiezoSleep narcolepsy screen revealed a 15-fold difference in wake-maintenance scores between wild type mice and the most severely affected narcoleptic mice. Compared to historical methods, the PiezoSleep narcolepsy screen increased by 16% the number of mice that could be identified that met the cataplexy inclusion criteria for more intensive studies. The ED_{50} for wake-promotion by modafinil was approximately 50 mg/kg in both genotypes. Using unsupervised piezoelectric monitoring, the efficacy of wake-promoting compounds can be determined in a 5-arm study with 60 mice in less than one week—a fraction of the time compared to EEG/EMG studies.

Conclusions: PiezoSleep offers rapid, scalable assessment of sleep/wake for high-throughput screening in drug discovery.

Keywords

modafinil, mouse, narcolepsy, non-invasive sleep scoring, orexin, piezoelectric, transgenic

Background

Pharmacological studies with arousal state endpoints are typically invasive, resource intensive, and time consuming. Rodent polysomnography requires surgical implantation of electrodes, recovery time, adaptation to sleep implant hardware, and manual scoring of arousal states for the highly accurate detection of sleep microstructure. Using standard electroencephalography (EEG) and electromyography (EMG), wakefulness is defined as low amplitude, desynchronous activity in the EEG across a broad frequency range with variable EMG activity. Sleep in rodents is classified as either rapid-eye-movement (REM) sleep, with pronounced activity in the EEG theta band (4.5-9.5 Hz) and atonia in the EMG, or non-REM (NREM) sleep, with slow, synchronous activity in the EEG delta band (0.5 to 4.5 Hz) and low, steady EMG tone. Some of the burden of rodent polysomnography has been relieved by the development of automated methods of scoring EEG/EMG records. However, the expediency and objectivity value of these methods is diminished for those that require a subset of manually scored samples from individual animals to train the algorithm [1-4]. Unsupervised, automated scoring algorithms have addressed these limitations of machine learning with varying degrees of accuracy (72-96% agreement) compared to standard manual scoring [5-7]. Nevertheless, all EEG/EMG based systems of arousal state monitoring are hindered by the time required for surgery, recovery, and in many cases, the repeated-measures experimental designs used for studies in which a small number of animals are used. There is a need in drug discovery for a fully automated method of sleep/wake staging that can scale to a sample size that is amenable to high-powered, between-subject designs for the rapid screening of compounds.

The use of non-invasive proxy-measures of sleep/wake, such as actigraphy or

cardiovascular correlates of arousal states instead of traditional polysomnography, can expedite

sleep/wake determination in rodents [8-11]. In particular, the use of a highly sensitive motion

detector (piezoelectric sensor) on the cage floor to monitor gross motor activity, together with

breathing patterns during periods of behavioral quiescence, yields unsupervised sleep/wake

detection with 90% accuracy vs. manual EEG scoring [12]. This piezoelectric sleep monitoring

system (PiezoSleep) has been used to characterize daily sleep/wake amounts and patterns in a

transgenic mouse model of Alzheimer's disease [13] and in 180 knockout lines as part of a

large-scale phenotyping program [14].

Transgenic *orexin/ataxin-3* (Atax) mice model the human sleep disorder narcolepsy with

high fidelity. Like their human counterparts, Atax mice experience degeneration of wake-

promoting orexin (also known as hypocretin) neurons which leads to cataplexy (a sudden,

reversible loss of muscle tone and REM-sleep like EEG) and/or increased sleep propensity at

times of day when wakefulness would otherwise predominate [15]. While cataplexy is

pathognomonic of narcolepsy, excessive daytime sleepiness (EDS) is the most problematic

symptom in patients with narcolepsy [16]. The EDS leads to an inability of patients with

narcolepsy to sustain long periods of wakefulness during the Maintenance of Wakefulness Test

(MWT) and to fall asleep with short latency on the Multiple Sleep Latency Test (MSLT). Both

tests are indicators of symptom severity; the MSLT is used in the diagnosis of narcolepsy and

the MWT is used to assess treatment efficacy [17]. Modafinil (MOD) is a first-line choice of

treatment for EDS in patients, and promotes EEG/EMG-defined wakefulness in narcoleptic

mice, but is ineffective in cataplexy reduction [17, 18]. Although Atax mice exhibit cataplexy that

is readily identifiable according to EEG/EMG-dependent consensus criteria [19], 34% of Atax

mice have fewer than 3 cataplexy episodes per day, thus presenting a challenge to sort mice for

inclusion in studies with cataplexy reduction as an endpoint [16]. Currently, there is no

prescreening method prior to EEG/EMG studies for sorting mice that are highly symptomatic for

narcolepsy.

108   Drug discovery efforts aimed at finding improved therapeutics for narcolepsy would

benefit from a translational assay in mice similar to the MSLT and MWT that is rapid, scalable,

and not dependent upon polysomnography nor sleep deprivation protocols [20]. In the current

report, we introduce the "wake-maintenance score" as a translational measure of sleep/wake

fragmentation as determined by the PiezoSleep narcolepsy screen. This assay may help

prescreen Atax mice that are highly symptomatic and prone to cataplexy, as patients with

severe cataplexy are the least able to stay awake on the MWT test [21]. In a previous study,

Atax mice were monitored with piezoelectric detection of movement and heartrate for a few

hours during the light period, but the algorithm for autoscoring was only 73% accurate against

manual EEG scoring [22] and missed the time of day in which cataplexy and sleep/wake

fragmentation occurs most frequently [16]. Here, we demonstrate how the PiezoSleep system

can be used to determine the severity of narcoleptic symptoms in Atax mice for subject

stratification and to quantify sleep time and consolidation in pharmacological studies at a scale

amenable to screening wake-promoting therapeutics in drug discovery.

Methods

124   All experimental procedures were approved by the Institutional Animal Care and Use

125   Committee at Reset Therapeutics and were conducted in accordance with the principles set

126   forth in the *Guide for Care and Use of Laboratory Animals of the National Institute of Health*.

Animals

128   Male and female hemizygous transgenic B6.Cg-Tg(HCRT-MJD)1Stak/J mice (Atax) and

wild type (WT) littermates (JAX stock #023418, Jackson Labs, Bar Harbor, ME) were used

(2001 Hara Neuron). Separate cohorts were used for immunohistochemistry, development of

the narcolepsy phenotyping assay and PiezoSleep narcolepsy screen, EEG/EMG and video

monitoring, and pharmacology. Male C57BL/6J mice (JAX stock #000664, Jackson Labs, Bar

Harbor, ME) were also used in pharmacological studies. Male CD-1 mice (Charles River

Laboratories, Hollister, CA), 7-8 weeks old, were used in pharmacokinetic studies. Animals were

individually housed, except for CD-1 mice which were housed 3 per cage, at 22-25 ^{o}C, 30-70%

relative humidity, and LD12:12 with *ad libitum* access to food, water and nesting material.

Immunohistochemistry

All histological procedures and cell counts were performed at Jackson Laboratories (Bar

Harbor, ME). Male and female Atax (*N* = 8 and 5, respectively) and WT littermates (*N* = 8 males

and 6 females), aged 16.9 ± 0.1 and 28.5 ± 0.8 g were euthanized by pentobarbitol overdose

and transcardially perfused with 0.4 % paraformaldehyde. Brains were harvested, postfixed

overnight, and cryo sectioned at 40 µm from approximately -1.23 to -1.91 relative to Bregma in

coronal sections. After blocking in normal donkey serum (5% in 0.3% triton in PBS), free-floating

brain sections were incubated overnight at room temperature with goat anti-donkey orexin A

antibody (SC-8070, Santa Cruz Biotechnology, CA) at 1:2000 dilution. After washing with

phosphate-buffered saline, sections were incubated for 2 hours at room temperature with

biotinylated donkey anti-goat (Jackson Immunoresearch, West Grove, PA) at 1:500 dilution in

blocking solution. After further washes, sections were developed with the avidin-biotin-

peroxidase system using diaminobenzidine as the chromagen (Vector Laboratories,

Burlingame, CA). Sections were mounted onto slides, and anatomically matched sections

between brains that represented the anterior, mid, and posterior regions of the orexin field were

identified for manual cell counting. Counts were performed bilaterally on these sections at 20x

magnification under light microscopy. Only somata with complete and intact cell membranes

were included in the analysis.

Piezoelectric monitoring

The narcoleptic phenotype of Atax mice and their sleep/wake responses in

pharmacological studies were determined by piezoelectric recording of gross body movement

and breath rate (PiezoSleep version 2.11, Signal Solutions, Lexington, KY). The PiezoSleep

apparatus consisted of 60 individual polycarbonate cages, each of which contained a

polyvinylidene difluoride piezo sensor beneath a thin (0.025 cm) plastic shield. The sensor

covered the entire area of the cage floor (232.3 cm^2) and was connected to a single channel

tube amplifier (RoMo version 1.2, Signal Solutions, Lexington, KY). Each amplifier fed into a

digital acquisition device connected to a computer for data collection and sleep/wake

determination as previously described [8, 12, 23]. Briefly, signals were amplified and filtered

between 0.5 and 10 Hz and analog-to-digital converted at 120 Hz sampling rate. The frequency,

amplitude, and peak spectral energy of the signal were evaluated in 2 second increments over a

tapered, 8 second window to automatically calculate a sleep/wake decision statistic by simple

linear discriminant analysis. Signals were classified as sleep if they exhibited a periodicity with a

consistent, low relative amplitude at a frequency in the typical breathing range (1-4 Hz).

Wakefulness was associated with high amplitude signals with variable frequency and broad

spectral energy. Signal features related to these properties were extracted and combined to

obtain a decision statistic for each interval. Over time, the decision statistics per increment

accumulated in a bimodal distribution. The saddle point of the distribution was determined using

an adaptive threshold for the classification of sleep and wake. Bouts of contiguous intervals of

sleep or wake were defined to terminate when an interval, equal to the minimum bout length of

30 s, included less than 50% of the targeted state.

Piezoelectric experiments began by loading mice into individual PiezoSleep cages that

contained standard rodent bedding (diamond dry cellulose approximately 1 cm thick) with food

and nesting material from each mouse's home cage. Mice were permitted at least 24 h of

acclimation before experimental data were sampled. For development of the narcolepsy

phenotype assay, after the acclimation period, Atax mice (N = 31, aged 15.7 ± 0.3 weeks, 29.4

± 0.4 g) and their WT littermates (N = 17, aged 14.8 ± 0.5, 29.5 ± 0.4 g) were recorded

undisturbed for 36 h beginning at zeitgeber time (ZT)12 (21:00). Hourly sleep percentages,

mean sleep bout durations (SBD) and wake bout durations (WBD) per h, and histograms of the

percent time spent asleep or awake in bouts from 0.5 to ≥ 64 min were calculated using

SleepStats version 2.11 (Signal Solutions, Lexington, KY). Based on the circadian distribution of

mean SBD and WBD per h, and on the increased propensity for cataplexy during the early dark

period [16], sleep and wake data from ZT12-18 were used to differentiate mice based on their

ability to maintain long bouts of wakefulness at that time of day. Scatterplots of time spent in

short SBD (< 8 min) vs. long WBD (≥ 32 min) were used to screen Atax mice that were highly

symptomatic for murine narcolepsy (i.e., compared with other Atax mice, they spent more than

the mean amount of time in short SBD and less than the mean amount of time in long WBD.) All

mice were assigned a wake-maintenance score based on their ratio of time in long WBD:short

SBD during ZT12-18 for the PiezoSleep narcolepsy screen.

EEG/EMG & video monitoring

A second cohort of male Atax mice (N = 37, aged 11.6 weeks, 26.6 ± 0.4 g) was

subjected to the PiezoSleep narcolepsy screen to identify the individuals that were the most

highly symptomatic for narcolepsy and thus the best candidates for EEG and EMG studies. The

16 Atax mice with the lowest wake-maintenance scores were surgically implanted at 16.9 ± 0.2

weeks of age (30.2 ± 0.7 g) with transmitters (F20-EET, Data Sciences International, St. Paul,

MN) for EEG and EMG recording as previously described [24]. Mice were anesthetized with

isoflurane and transmitters were placed IP along the midline. Biopotential leads were routed

subcutaneously to the head, and EMG leads were positioned bilaterally through the nuchal

muscles. Cranial holes were drilled 1 mm anterior to bregma and 1 mm lateral to midline, and

contralaterally, 2 mm posterior to bregma and 2 mm lateral to midline. EEG leads were placed

subchranially over the dura and attached to the skull with cyanoacrylate and dental acrylic.

Analgesia was managed with prophylactic ketoprofen (5 mg/kg, SC) followed by buprenorphine

(0.1 mg/kg, SC) upon emergence from anesthesia. Ketoprofen (5 mg/kg, SC, QD) was

continued for 3 days postsurgery. Mice received postoperative thermal support for 24 h and

hydration (0.7 mL lactated Ringer's solution, SC and dietary recovery gel) for 3 days. Mice were

permitted 3 weeks postsurgical recovery, which included 2 weeks adaptation to running wheels

and the light-controlled (LD12:12 with ZT0 = 03:00), sound-attenuated and ventilated recording

chamber.

Physiological and video-recorded behavioral data were simultaneously acquired with

Ponemah version 5.2 (Data Sciences International, St. Paul, MN). Electrophysiological data

were sampled at 500 Hz and digital videos were recorded at 15 frames per second (640 x 480

pixels) using Noldus Media Recorder (Leesburg, VA). Recordings from mice in their home

cages (with nesting material and running wheels) began at ZT6 and continued for the following

30 h. Data were manually scored using NeuroScore version 3.2.1 (Data Sciences International,

St. Paul, MN) by scorers trained with ≥ 96% interrater reliability by an expert scorer. Data were

classified in 10-second epochs as wakefulness, wheel-running during wakefulness, NREM

sleep, REM sleep, or cataplexy. Criteria for cataplexy were ≥ 10 seconds of EMG atonia, theta-

dominated EEG, and video-confirmed behavioral immobility preceded by ≥ 40 seconds of

wakefulness [19]. Bouts of each arousal state were defined to terminate with the occurrence of

a single epoch of another state. Of the 16 implanted mice, 11 were included in the analysis

(aged 19.9 ± 0.3 weeks at recording), 2 did not survive surgical recovery, and 3 had corrupted

data files or unscorable EEG signals.

Compound formulation

Modafinil (MOD, 2-[(diphenylmethyl)sulfinyl]acetamide) was purchased from Focus

Synthesis (San Diego, CA) for dosing in CD-1 and WT and from Advanced ChemBlocks, Inc.

(Burlingame, CA) for dosing in Atax mice because MOD was no longer available from Focus

Synthesis. For pharmacokinetic studies, MOD was formulated in 1.25% Hydroxypropyl

Methylcellulose / 0.25% Methylcellulose in Water for Injection, which formed a homogenous

suspension. For efficacy studies, after MOD was weighed, premade vehicle (20% propylene

glycol, 25% polyethylene glycol 400 in water for injection) was added at a volume appropriate

for the highest concentration, followed by overnight sonication, which produced a clear solution.

Serial dilutions were created and re-sonicated approximately 6 h prior to dosing. All dosing

solutions were formulated within 24 h prior to dosing.

Pharmacokinetic studies

Male CD-1 mice were acclimated for a minimum of 3 days prior to drug administration.

Modafinil was administered as 10, 30, 50, or 100 mg/kg via oral gavage (PO) to CD-1 mice at

ZT0 to ZT1. Blood and tissue samples were collected from three mice per time point at pre-

dose, 15, 30, 60, 90, 180, 360, 720, and 1440 min following oral gavage from the CD-1 mice.

Blood (500 µL) was obtained via cardiac puncture into Microtainer Blood Collection Tubes with

K_2EDTA (BD Biosciences, San Jose, CA), and gently mixed. Blood was immediately

centrifuged, and the resulting plasma was frozen on dry ice then stored at -80 °C until it was

analyzed. Entire brain from each animal was harvested, weighed, frozen on dry ice, and stored

at -80 °C until it was analyzed. The concentrations of MOD were determined by a liquid

chromatography – mass tandem spectrometry (LC-MS/MS) method (Quintara Discovery,

Hayward, CA) following the standard protocol. Briefly, Brain samples were homogenized in two

volumes of ice cold water. Then 20 µL of each plasma or brain homogenate sample were

processed by protein precipitation with methanol: acetonitrile (5:95 v:v) containing internal

standard, verapamil. Extracts were analyzed by LC-MS/MS using positive electrospray

ionization under the multiple-reaction-monitoring mode for transitions m/z 274->167 (MOD) and

m/z 455-> 165 (verapamil). The lower limit of quantitation was 1.8 nM for plasma and 5.5 nM

for tissues. Pharmacokinetic parameters of MOD were generated by non-compartmental

analysis with Phoenix™ WinNonlin® software (Pharsight Corporation, St. Louis, MO).

Efficacy studies

In a between-subjects design, male C57BL/6 mice (WT, 15.7 ± 0.1 weeks, 28.5 ± 0.2 g,

N = 12/group) and Atax mice (16.4 ± 0.7 weeks, 30.0 ± 0.7 g, *N* = 9-10/group) were dosed

separately in two pharmacological studies. Mice were loaded into PiezoSleep cages for 42 h

acclimation, followed by 29 h baseline recording prior to dosing at ZT5 (14:00), the time of day

in which wake-promoting/sleep-inhibiting effects would be expected to be most pronounced.

Syringes were filled immediately prior to dosing, which was completed within a 15-minute

window. After pharmacokinetic studies (Table 1) and based on efficacy in previous reports [18,

25], mice received either MOD (10, 30, 50, 100 mg/kg, 10 mL/kg) or vehicle orally and were

recorded for 44 h post dosing. Mice were undisturbed during the recording period except for

dosing. Groups were balanced across dosing order, dosing personnel, and cage location per

shelf and rack. Atax mice had previously been screened in the narcolepsy phenotype assay,

and highly symptomatic mice were balanced across dosing groups.

Table 1. Pharmacokinetic data after oral modafinil at ZT0 in CD-1 mice

Dose (mg/kg)	$T_{1/2}$ (hrs)		MRT (hrs)		Cmax (µM)		AUC (µM.h)	
	Plasma	Brain	Plasma	Brain	Plasma	Brain	Plasma	Brain
10	1.1	1.4	1.8	1.5	14.2	4.1	22.5	7.3
30	0.9	1.2	1.7	1.5	76.1	15.4	131.3	34.5
50	0.8	1.2	2.1	1.9	80.1	33.7	185.5	72.4
100	1.0	1.1	2.4	2.7	306.2	76.5	958.5	294.9

273    $T_{1/2}$: half-life
274   MRT: mean residence time
275   AUC: area under the curve

Data analysis and statistics

Statistics were performed using GraphPad Prism version 7.03 (La Jolla, CA). Unless otherwise noted, all data are presented as the mean ± S.E.M. Orexin cell counts were compared between male and female Atax and WT mice using two-way analysis of variance (ANOVA) and between genotypes for combined sexes using two-tailed, unpaired t test. In phenotyping studies, sleep time and mean SBD and WBD were compared between Atax and WT mice using two-way mixed-model ANOVA on factors "genotype" (between subjects) and "hour" (within subjects). Histograms representing the distribution of time spent in various SBD and WBD were evaluated using two-way mixed-model ANOVA on factors "genotype" (between subjects) and "bout duration bin" (within subjects). Ratios of time spent in long wake bouts and short sleep bouts were compared between genotypes using two-tailed, unpaired t test. Comparisons of wake time measurements via piezoelectric monitoring vs. EEG/EMG with video recording over 24 h were made within subjects using two-way repeated-measures (RM) ANOVA, and for 24 h totals using two-tailed, paired t test. Pharmacological data were assessed within genotypes by two-way mixed-model ANOVA on between-subjects factor "drug condition" and within-subjects factor "hour" or "bout duration bin." The maximum sleep deficit was calculated as the minimal value in the cumulative change from baseline in sleep % time during the 12 h post dosing. Consolidation of wakefulness after dosing was measured as the change from baseline in the % time spent in long wake bouts ≥ 8 min. Sleep deficits and consolidated wakefulness were compared between genotypes after dosing using two-way ANOVA. For each analysis, when ANOVA indicated significance (α = 0.05), contrasts between relevant factor levels (vs. WT or vehicle, as appropriate) were detected *post hoc* with Bonferroni's multiple comparisons test. The total sample size ($N = 97$) for the pharmacological study permitted

300　sufficient power (0.8) for the detection of an effect size of 36% at $P < 0.05$ (computed using

301　WebPower https://webpower.psychstat.org/wiki/).

Results

Histology

To confirm orexin cell loss from Atax mice taken out of cryopreservation at JAX, orexin

cell counts were compared between the transgenics and their WT littermates at an adult age in

which they would be used for sleep studies. Representative images (Figure 1) of the tuberal

region of the hypothalamus and cell counts (Table 2) indicated that, as expected, Atax mice had

fewer cells that stained positive for orexin A. Total numbers of orexin-positive cells counted from

WT males vs. Atax males were 511 ± 21 vs. 12.9 ± 3.0, while from WT females vs. Atax females

were 399 ± 54 vs. 19.0 ± 12. There were no sex differences in the number of orexin neurons

that were sampled, $F_{(1, 23)} = 3.58$, *n.s.* The extent of orexin neurodegeneration was consistent

with that observed in a different colony of Atax mice [24].

Table 2. Normalized orexin cell counts in wild type (WT) and *orexin/ataxin-3* (Atax) mice

	N/group	WT	Atax	statistic
Males	8	100 ± 4.1	**2.5 ± 0.6**	**$F_{(1, 23)} = 207.5$**
Females	5-6	100 ± 13.6	**5.1 ± 3.8**	
Both sexes	13-14	100 ± 6.0	3.4 ± 1.2	*t*(25) = 15.3

Data are the mean ± S.E.M. expressed as the % of WT mean per sex

Bold = main effect for genotype; no interaction or main effects for sex

Figure 1. Loss of orexin neurons in orexin/ataxin-3 (Atax) mice. Representative images of
Orexin A+ cells in the posterior lateral hypothalamus of male **(top)** and female **(bottom)** wild
type littermates **(left)** and Atax mice **(right)**. Images taken at 10x magnification (bar = 100 µM),
approximately -1.5 mm from Bregma; fornix (f), internal capsule (ic), optic tract (opt), third
ventricle (3V).

Phenotyping of narcolepsy model

Piezoelectric monitoring of sleep/wake was used to evaluate the phenotype of Atax mice

and to sort mice by the severity of murine narcolepsy symptoms. Compared to WT littermates,

Atax mice slept more at a few time points during the dark period, particularly during the last hour

prior to lights-on (Figure 2A), $F(36, 1656) = 4.81$, $P < 0.0001$. However, the total amount of

sleep per 24 h was the same between genotypes (WT: 45.5 ± 0.8 % vs. Atax: 45.8 ± 1.0 %),

$t(46) = 0.03$, *n.s.*, as expected. Genotype differences were observed in sleep and wake bout

duration measures. Atax mice exhibited shorter mean SBD per h across the day (Figure 2B,

$F(36, 1649) = 4.65$, $P < 0.0001$) and shorter mean WBD during the dark period (Figure 2C,

$F(36, 1656) = 8.62$, $P < 0.0001$).

Figure 2. The PiezoSleep narcolepsy phenotyping assay. Determination of narcoleptic
phenotype in male orexin/ataxin-3 (Atax, *N* = 31) mice **(black)** vs. wild-type (WT, *N* = 17)
littermates **(gray dashed lines)** using PiezoSleep. Sleep % per h **(A)** and mean bout durations
per h of sleep **(B)** and wake **(C)** over the 36 h recording period. Percentage of time during the
first half of the dark period (zeitgeber time (ZT)12-18) spent asleep **(D)** and awake **(E)** in bout
durations lasting from 0.5 to ≥ 64 min. Percentage of time spent in short sleep bouts (< 8 min)
vs. long wake bouts (≥ 32 min) during the first half of the dark period per individual mouse **(F)**.
Two-way mixed-model analysis of variance with Bonferroni's multiple comparisons: * P < 0.05
vs. WT.

Because differences in both mean SBD and WBD were particularly robust during the first

half of the dark period, and because that time of day is associated with an increased probability

of cataplexy occurrence [16], ZT12-18 was chosen as the focus for determination of narcolepsy

symptom severity. The percentage of time spent in sleep bouts that were 0.5 to 4 min long was

greater in Atax than for WT mice, while WT mice spent more time than Atax mice in sleep bouts

that were 8 to 32 min long (Figure 2D), $F(7, 368) = 24.31$, $P < 0.0001$. Compared to WT

littermates, Atax mice spent more time in wake bouts that were 8-32 min long and less time in

the longest wake bouts of ≥ 64 min duration (Figure 2E), $F(7, 368) = 25.74$, $P < 0.0001$. The

percentage of time spent in short SBD (< 8 min) vs. time spent in long WBD (≥ 32 min) were

plotted in Figure 2E. Atax mice clustered on the scatterplot in a manner indicative of more time

spent in short sleep bouts and less time in long wake bouts than their WT littermates. The ratio

of time spent in long WBD to short SBD determined the "wake-maintenance score" and was

smaller in Atax mice (1.5 ± 0.2) than in WT (9.9 ± 2.7), $t(46) = 4.19$, $P < 0.0001$.

The wake-maintenance score was used to sort individual Atax mice by murine

narcolepsy symptom severity to prioritize EEG/EMG instrumentation on the mice most likely to

demonstrate cataplexy. Using the PiezoSleep narcolepsy screen on a separate cohort of Atax

mice, 16 mice were identified as "highly symptomatic" for murine narcolepsy, with lower wake-

maintenance scores (0.68 ± 0.1) than the remaining Atax littermates (3.15 ± 0.4), $t(35) = 5.17$, P

< 0.0001, and were implanted for EEG/EMG monitoring. The wake-maintenance scores of the

11 mice that were ultimately included in the EEG/EMG analysis (0.67 ± 0.1) were representative

of the original 16 animals.

The percentage of time spent awake per hour across 24 h was compared between

piezoelectric recordings from the phenotype screen and EEG/EMG with video monitoring that

was performed 8 weeks later (Figure 3A). This comparison of non-simultaneous recordings from

different environments was not intended to validate the accuracy of the piezoelectric sleep/wake

detection against polysomnography, as has previously been done [8, 12], but rather was

intended to assess the predictive consistency between the two methods under different use

cases. The pattern of the daily distribution of wakefulness appeared to be similar between

sleep/wake recording methods, but piezoelectric determination yielded slightly more

wakefulness per 24 h (53.3 ± 0.9%) than EEG/EMG with video monitoring (49.7 ± 1.3%), $t(20) =$

2.31, $P = 0.03$). However, the only time point in which the PiezoSleep recordings differed from

the EEG/EMG recordings was in the hour immediately prior to lights on, $F(23, 230) = 29.3$, $P <$

0.0001.

Time spent in NREM and REM sleep (Figure 3B) and during cataplexy and wheel-

running (Figure 3C) were quantified in these mice from the EEG/EMG with video recordings.

During the dark period, 9 of the 11 mice (82%) exhibited ≥ 3 bouts of cataplexy, the typical

criteria for inclusion in pharmacological studies on cataplexy [16]. These mice exhibited 5.4 ±

0.9 bouts of cataplexy that each lasted 1.2 ± 0.2 min long for 4.7 ± 1.2 cumulative minutes

during the dark period.

Figure 3. Quantification of arousal states in narcoleptic mice by EEG/EMG recording.
Comparison of wake % per h in male orexin/ataxin-3 (Atax, $N = 11$) mice as measured by
piezoelectric monitoring **(A, black line)** vs. manual scoring **(A, gray dashed line)** of
EEG/EMG/video recordings from the same mice 8 weeks later. The % per h of NREM **(B, black**
line), REM **(B, gray dashed line)**, cataplexy **(C, black line)**, and wheel running **(C, gray**
dashed line) as measured by manual scoring of EEG/EMG/video recordings. Two-way
repeated-measures analysis of variance with Bonferroni's multiple comparisons: * P < 0.05 vs.
EEG/EMG/video.

Sleep/wake pharmacology using PiezoSleep

To validate piezoelectric monitoring of sleep/wake for the screening of wake-promoting compounds for narcolepsy, male Atax mice ($N = 46$) and WT mice ($N = 60$) were dosed with MOD (10, 30, 50, 100 mg/kg, 10 mL/kg) or vehicle orally at ZT5 when sleep propensity is high. As expected, MOD induced a dose-related inhibition of sleep for 3 h (30 and 50 mg/kg) and 6 h (100 mg/kg) post dosing vs. vehicle in WT (Figure 4A), $F(24, 330) = 11.4$, $P < 0.0001$. In Atax mice, MOD inhibited sleep for 2, 4, and at least 7 h post dosing at 30, 50, and 100 mg/kg, respectively, compared to vehicle (Figure 4B), $F(24, 246) = 8.25$, $P < 0.0001$. Although a dose-related response to MOD (10-100 mg/kg) was evident for the maximum sleep deficit in both Atax and WT mice (Figure 4C), $F(4, 96) = 96.0$, $P < 0.0001$ (main effect for dose), there was no interaction with genotype $F(4, 96) = 1.63$, *n.s.*

Modafinil fragmented sleep bouts in both mouse models. Under baseline and vehicle conditions during the last 7 h of the light period, WT mice spent most of their sleep time (26-31% of total time) in bouts that were 32-64 min long (Figure 4D). After MOD (10-100 mg/kg), the percentage of time spent in these long sleep bouts was reduced in dose-related manner, $F(28, 385) = 5.38$, $P < 0.0001$, and the distribution of time spent in the different SBD bins shifted such that more time was spent in shorter bouts than longer bouts (i.e., leftward shift of the peak), especially at the highest dose. Unlike WT mice, Atax mice in baseline and vehicle conditions spent 5-6% of the time in long sleep bouts ≥ 32 min, and most of their sleep time (22-25% of total time) was in bouts that were 8-32 min long (Figure 4E). Modafinil (10-100 mg/kg) further fragmented the sleep of Atax mice, as the percentage of time spent in sleep bouts 8-32 min long was reduced in a dose-related manner, $F(24, 246) = 8.25$, $P < 0.0001$. However, only the highest dose of MOD shifted the peak of the sleep bout distribution curve to the left toward shorter bouts. A dose-related increase in the percentage of time spent in wake bouts ≥ 8 min was evident after MOD (10-100 mg/kg) in WT and Atax mice (Figure 4F), $F(4, 96) = 98.6$, $P <$

0.0001 (main effect for dose), but the effect did not depend on genotype $F(4, 96) = 2.87$, $P =$

0.03, no significant contrasts per dose.

Figure 4. Piezoelectric monitoring of sleep/wake in pharmacological studies. Inhibition of
sleep and promotion of wakefulness by modafinil (MOD) in orexin/ataxin-3 (Atax, **filled symbols,**
B, E and **black lines C, F,** N = 9-10/group) vs. wild type (**open symbols, A, D** and **gray lines C,**
F, N = 12/group) male mice as detected by PiezoSleep. Mice were administered MOD at 0
(vehicle, **gray squares**), 10 (**circles**), 30 (**black squares**), 50 (**diamonds**), or 100 (**triangles**)
mg/kg (mpk) at 5 h after lights on (zeitgeber time (ZT)5, **arrow**) after 30 h undisturbed baseline.
Sleep % per h (**A-B**) and the percentage of time spent in sleep bout durations lasting from 0.5 to ≥
64 min (**D-E**) for 7 h post dosing were plotted against values taken from undisturbed baseline

(**gray dashed line**) for reference. The maximum sleep deficit (**C**) and the % of time spent in
consolidated wake bouts ≥ 8 min for 7 h post dosing, expressed as the change from baseline (**F**),
revealed dose-related wake-promotion by MOD (differences between genotypes, *n.s.*). Two-way
analysis of variance with Bonferroni's multiple comparisons: * P < 0.05 vs. vehicle for doses and at
time or bout duration bin ranges as indicated by bars at top of graphs.

Discussion

Piezoelectric monitoring of sleep/wake was used to identify key features of the

narcoleptic phenotype that distinguish Atax mice by symptom severity, thus enabling rapid, non-

invasive sorting of mice for EEG/EMG studies for cataplexy detection. The PiezoSleep

narcolepsy screen was used to assign a symptom severity score—the ratio of time spent in long

WBD (≥ 32 min) to time in short SBD (☐ 8 min) during the first half of the dark period—to

individual mice. These wake-maintenance scores spanned a 15-fold difference between the

most severely narcoleptic mice and their WT littermates. The PiezoSleep system was also

demonstrated to detect changes in sleep/wake time and consolidation after administration of the

wake-promoting therapeutic modafinil in normal and transgenic mice. Using this method, the

efficacy of wake-promoting compounds can be determined in a 5-arm study with up to 60 mice

in less than one week for high-throughput screening *in vivo*.

The PiezoSleep narcolepsy screen was designed to be a translational test of the ability

to sustain long periods of wakefulness as is assessed in humans by the MWT and the MSLT.

For the human tests, patients are provided 4-5 opportunities throughout the daytime to either

nap within 20 min (MSLT) or to try to stay awake for 20-40 min (MWT) in the absence of alerting

factors. The latency to polysomnographically defined sleep is measured for each test. A mean

sleep onset latency ≤ 8 min with ≥ 2 sleep-onset REM periods (abnormal transition into REM

sleep within 15 min of sleep onset) is a positive result on the MSLT and aids in the diagnosis of

narcolepsy [17]. A murine MSLT has been devised in which the latency to EEG/EMG-defined

sleep is measured during four 20-min nap opportunities presented after 20 min of enforced

wakefulness [20]. The PiezoSleep narcolepsy screen provides a rapid, automated alternative to

the murine MSLT and does not require the manipulation of sleep homeostatic processes to

obtain the measurements. The translational utility of the PiezoSleep phenotyping assay is

limited by the inability of the current version of the algorithm to distinguish between NREM and

REM sleep. Perhaps a new algorithm under development [26] will permit quantification of short

REM sleep latency into the assay as a murine equivalent of sleep-onset REM periods.

The PiezoSleep narcolepsy screen can be used to prescreen narcoleptic mice to identify

those with the most fragmented sleep/wake for inclusion in resource-intensive EEG/EMG

studies. The wake-maintenance score clearly differentiated narcoleptic mice with high sleep

propensity from WT littermates that were able to spontaneously sustain long bouts of

wakefulness from ZT12-18 (Figure 2). It is unknown if the wake bouts in Atax mice terminated

because of sleep onset or from the intrusion of cataplexy, which may register on the PiezoSleep

assay as sleep due to the similar motor and autonomic physiology between REM sleep and

cataplexy. Nevertheless, this distinction does not detract from the utility of the PiezoSleep

narcolepsy screen to differentiate Atax mice by severity of sleep/wake fragmentation. Out of the

37 Atax mice screened for inclusion in the EEG/EMG study, 16 mice with severe symptoms

were identified through rank ordering of the wake-maintenance scores. These mice scored 4.7x

lower on the PiezoSleep narcolepsy screen than the remainder of the Atax mice, and only 2 out

of the final 11 included in analysis (18%) had □ 3 cataplexy bouts. Compared to a previous

report that 34% of Atax mice fall below this study inclusion criteria [16], the current study

represents a 16% improvement in yield. However, it is difficult to determine if the yield

differences are due to the new sorting methodology or to differences between colonies and

recording environments. The reason for high interindividual differences in cataplexy expression

between Atax mice remains unclear, but may recapitulate the etiology of human narcolepsy type

2 (narcolepsy without cataplexy) which is also poorly understood [16]. Determination of whether

the wake-maintenance scores on the PiezoSleep narcolepsy screen correlate with the severity

of cataplexy would require EEG/EMG measurement from mice across a broader range of

symptom severity scores, and was beyond the scope of the current study. Comparison of

EEG/EMG-defined sleep/wake between mice with low and high wake-maintenance scores

identified by piezoelectric monitoring would be an important next step in the validation of the

PiezoSleep narcolepsy screen.

Some differences between studies in the comparison of piezoelectric monitoring of

sleep/wake and polysomnography are worth noting. Previous studies have validated the

PiezoSleep sleep/wake classifier against manually scored EEG/EMG signals that had been

simultaneously recorded from the same mice (across 4 strains) for direct epoch-to-epoch

comparison [8, 12]. The current analysis in Atax mice does not permit this level of scrutiny, as

the PiezoSleep and EEG recordings were staged in 2 and 10 second epochs, respectively, and

occurred 8 weeks apart in different environments (e.g., the EEG recordings occurred in

standard cages equipped with running wheels in a sound-attenuated recording chamber).

Wakefulness over 24 h was 3.6% higher after piezoelectric monitoring than after EEG/EMG

recording in the current study, vs. 2.3% lower as previously reported [12]. In addition to the

caveats between methods as noted above, the discrepancy could be due to the different mouse

strains studied (Atax vs. CFW Swiss Webster). Orexin knockout mice tend to breathe shallow

and fast compared to WT when both asleep and awake as determined by EEG/EMG [27]. This

respiration pattern during sleep could conceivably be misconstrued as quiet wakefulness in

orexin-deficient Atax mice in the PiezoSleep assay, thus slightly overinflating the measurement

of time awake compared to EEG/EMG in the current study. However, if this respiration-induced,

false detection of wake in narcoleptic mice were true, then piezoelectric monitoring would be

expected to detect more wakefulness in Atax mice than in WT littermates, which was not the

case (Figure 2A). The differences in sleep/wake as determined by EEG/EMG and piezoelectric

monitoring in the current study are likely due to the different environments used for recording.

Future studies are needed in which simultaneous piezoelectric EEG/EMG recordings are

collected during the dark period from narcoleptic mice to assess cataplexy detection using an

unsupervised algorithm that has been tuned to extract features of REM sleep [26] to determine

if cataplexy can be identified by piezoelectric monitoring.

Perhaps more noteworthy than the small discrepancies between polysomnography and

PiezoSleep monitoring in the current study was the consistency in the pattern of sleep/wake

between the two recording methods (Figure 3A). The hour-by-hour increase and decrease in

wake time after EEG/EMG recording paralleled the pattern observed after piezoelectric

monitoring, despite the 8 week difference in recording dates. The hour immediately prior to

lights-on was the only time point in which PiezoSleep registered more wakefulness in Atax mice

than polysomnography (i.e., At ZT23, Atax mice slept more as determined by polysomnography

than they did 8 weeks prior as measured with PiezoSleep). This time point was the same ZT in

which Atax mice slept more than WT mice after piezoelectric monitoring of both genotypes. The

increased sleep observed in Atax mice at the end of the dark period may reflect progressive

worsening of EDS over time in this model. The differences in sleep/wake measurement between

genotypes and between methods at ZT23 are less likely due to reduced accuracy of PiezoSleep

at that time point, as could be speculated, because the variance was similar at that hour

between the different runs and between other ZT hours. It is also not clear what physiological or

behavioral factors would differentially impact the detection of respiration rates and gross motor

activity at ZT23 vs. other times of day.

In the current study, piezoelectric measurement of sleep/wake was validated as a

method to screen wake-promoting compounds rapidly and at scale for the development of

therapeutics for narcolepsy. Dosing WT mice at ZT5 was chosen as the first pass of screening

using MOD because of the similarities between Atax and WT mice in the high amount of sleep

per hour (Figure 2A) and in the very short WBD (Figure 2C) during the light period. Modafinil (50

mg/kg) with total brain exposure of 33.7 µM approximated the ED_{50} in both WT and Atax mice in

terms of the time course of sleep inhibition, the maximum sleep deficit induced, and the

consolidation of wakefulness (Figure 4) and is recommended as the positive control for

screening the wake-promoting efficacy of compounds. Using the PiezoSleep assay, a total of

106 mice in two cohorts were screened across 4 concentrations of MOD vs. vehicle in two

weeks, representing substantial improvement in efficiency over conventional sleep/wake

bioassays [18, 25]. Future studies are needed to validate the PiezoSleep assay for

pharmacological studies at different times of day and with hypnotics. Of particular interest would

be to test the hypothesis that wake-promoting therapeutics administered at ZT12 would rescue

the narcoleptic phenotype of Atax mice as measured by the wake-maintenance score on the

PiezoSleep narcolepsy screen.

Piezoelectric monitoring of sleep/wake offers advantages over video-based methods,

even those that claim to discriminate NREM and REM sleep [28] for several reasons. Because

PiezoSleep does not depend on a visual signal, it permits mice to sleep while burrowed in

nesting material for assessment of more ecologically relevant behavior. The highly sensitive

piezoelectric sensor captures a detailed respiratory measure that enables it to distinguish quiet,

resting wakefulness from sleep using relevant physiology [12]. To the best of our knowledge,

sleep state discrimination based on postural changes in the aspect ratio of the body on video

recordings [28] has not been replicated nor validated in mouse models characterized by obesity,

such as Atax mice.

Conclusions

The non-invasive, unsupervised piezoelectric monitoring of sleep/wake is a new tool for

efficacy studies with arousal state endpoints. The PiezoSleep narcolepsy screen can be used in

mice to determine a wake-maintenance score that translates the ability to sustain long bouts of

wakefulness, as measured in humans with the MWT and MSLT, and can be used to improve

phenotypic sorting for resource intensive EEG/EMG studies. PiezoSleep has been validated as

a means to rapidly quantify sleep/wake time and consolidation for high-throughput screening *in*

vivo in drug discovery.

List of Abbreviations

ANOVA, analysis of variance

Atax, *orexin/ataxin-3*

AUC, area under the curve

Cmax, maximum concentration

EDS, excessive daytime sleepiness

EEG, electroencephalography

EMG, electromyography

f, fornix

ic, internal capsule

LD12:12, 12 h light period, 12 h dark period

MOD, modafinil

MSLT, multiple sleep latency test

MWT, maintenance of wakefulness test

NREM, non-rapid-eye-movement

opt, optic tract

PBS, phosphate-buffered saline

PO, *per os*

REM, rapid-eye-movement

RM ANOVA, repeated-measures analysis of variance

SBD, sleep bout duration

SC, subcutaneous

S.E.M., standard error of the mean

mpk, mg per kg

MRT, mean residence time

$T_{1/2}$, half life

WBD, wake bout duration

WT, wild type

ZT, zeitgeber time

3V, third ventricle

Declarations

Ethics approval and consent to participate: All experimental procedures were approved by the Institutional Animal Care and Use Committee at Reset Therapeutics and were conducted in accordance with the principles set forth in the *Guide for Care and Use of Laboratory Animals of the National Institute of Health*.

Consent for publication: Not applicable

Availability of data and material: The datasets generated and/or analyzed during the current study are not publicly available due to commercial ownership, but may be available upon request.

Competing interests: K.D. and B.O.H. own Signal Solutions, LLC, a company that produces and sells the PiezoSleep equipment and software used in this study, however no aspects of the research presented here were sponsored by Signal Solutions. S.W.B., J.D., A.L., N.K., P.S., R.B., and P.S.H. have been employed by Reset Therapeutics.

Funding: This work was supported by Reset Therapeutics and by the Michael J. Fox Foundation (grant number 11432 to P.S.H.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the Michael J. Fox Foundation.

Authors' contributions: S.W.B. designed and managed the efficacy studies, analyzed the data, prepared the figures and wrote the manuscript. J.D. designed and managed the PK study, prepared the table, formulated the compound, scored the data and reviewed the manuscript. A.L., N.K., and P.S. executed the studies, performed surgeries, scored the data and reviewed the manuscript. K.D. and B.O.H. provided the PiezoSleep system, created custom analysis tools, and reviewed the manuscript. R.B. and P.S.H. conceptualized the research goals, supervised the team, acquired funding and resources, and reviewed the manuscript.

Acknowledgements: We thank Drs. Thomas Scammell (Harvard University) and Takeshi Sakurai (University of Tsukuba) for the donation of the orexin/ataxin-3 mice to Jackson Labs,

621 Dr. Orsolya Kiraly (The Jackson Laboratory) for performing the histology, and Tod Steinfeld and

622 Dr. Bruce Clapham (Reset Therapeutics) for helpful discussions.

623 **Authors' information (optional):** Not applicable

References

1.   Van Gelder RN, Edgar DM, Dement WC: Real-time automated sleep scoring: validation

of a microcomputer-based system for mice. *Sleep* 1991, 14(1):48-55.

2.   Gross BA, Walsh CM, Turakhia AA, Booth V, Mashour GA, Poe GR: Open-source logic-

based automated sleep scoring software using electrophysiological recordings in rats. *J*

Neurosci Methods 2009, 184(1):10-18.

3.   Rytkönen KM, Zitting J, Porkka-Heiskanen T: Automated sleep scoring in rats and mice

using the naive Bayes classifier. *J Neurosci Methods* 2011, 202(1):60-64.

4.   Gross BA, Vanderheyden WM, Urpa LM, Davis DE, Fitzpatrick CJ, Prabhu K, Poe GR:

Stress-free automatic sleep deprivation using air puffs. *J Neurosci Methods* 2015,

251:83-91.

5.   Libourel PA, Corneyllie A, Luppi PH, Chouvet G, Gervasoni D: Unsupervised online

classifier in sleep scoring for sleep deprivation studies. *Sleep* 2015, 38(5):815-828.

6.   Stephenson R, Caron AM, Cassel DB, Kostela JC: Automated analysis of sleep-wake

state in rats. *J Neurosci Methods* 2009, 184(2):263-274.

7.   Sunagawa GA, Séi H, Shimba S, Urade Y, Ueda HR: FASTER: an unsupervised fully

automated sleep staging method for mice. *Genes Cells* 2013, 18(6):502-518.

8.   Flores AE, Flores JE, Deshpande H, Picazo JA, Xie XS, Franken P, Heller HC, Grahn

DA, O'Hara BF: Pattern recognition of sleep in rodents using piezoelectric signals

generated by gross body movements. *IEEE Trans Biomed Eng* 2007, 54(2):225-233.

9.   Pack AI, Galante RJ, Maislin G, Cater J, Metaxas D, Lu S, Zhang L, Von Smith R, Kay

T, Lian J *et al*: Novel method for high-throughput phenotyping of sleep in mice. *Physiol*

Genomics 2007, 28(2):232-238.

10.   Zeng T, Mott C, Mollicone D, Sanford LD: Automated determination of wakefulness and

sleep in rats based on non-invasively acquired measures of movement and respiratory

activity. *J Neurosci Methods* 2012, 204(2):276-287.

11.   Fisher SP, Godinho SI, Pothecary CA, Hankins MW, Foster RG, Peirson SN: Rapid

assessment of sleep-wake behavior in mice. *J Biol Rhythms* 2012, 27(1):48-58.

12.   Mang GM, Nicod J, Emmenegger Y, Donohue KD, O'Hara BF, Franken P: Evaluation of

a piezoelectric system as an alternative to electroencephalogram/ electromyogram

recordings in mouse sleep studies. *Sleep* 2014, 37(8):1383-1392.

13.   Sethi M, Joshi SS, Webb RL, Beckett TL, Donohue KD, Murphy MP, O'Hara BF, Duncan

MJ: Increased fragmentation of sleep-wake cycles in the 5XFAD mouse model of

Alzheimer's disease. *Neuroscience* 2015, 290:80-89.

14.   Sethi M, Joshi SS, Striz M, Cole N, Ryan J, Lhamon ME, Agarwal A, Sukoff Rizzo SJ,

Denegre JM, Braun RE *et al*: Analysis of sleep traits in knockout mice from the large-

scale KOMP2 population using a non-invasive, high-throughput piezoelectric system.

BMC Bioinformatics 2015, 16(15):P15.

15.   Hara J, Beuckmann CT, Nambu T, Willie JT, Chemelli RM, Sinton CM, Sugiyama F,

Yagami K, Goto K, Yanagisawa M *et al*: Genetic ablation of orexin neurons in mice

results in narcolepsy, hypophagia, and obesity. *Neuron* 2001, 30(2):345-354.

16.   Black SW, Yamanaka A, Kilduff TS: Challenges in the development of therapeutics for

narcolepsy. *Progress in neurobiology* 2017, 152:89-113.

17.   Cao MT, Guilleminault C: Narcolepsy: Diagnosis and management. In: *Principles and*

Practice of Sleep Medicine. 6th edn. Edited by Kryger MH, Roth T. Philadelphia,

Pennsylvania: Elsevier; 2017: 873-882.

18.   Black SW, Schwartz MD, Chen TM, Hoener MC, Kilduff TS: Trace Amine-Associated

Receptor 1 Agonists as Narcolepsy Therapeutics. *Biological psychiatry* 2017, 82(9):623-

633.

19.   Scammell TE, Willie JT, Guilleminault C, Siegel JM: A consensus definition of cataplexy

in mouse models of narcolepsy. *Sleep* 2009, 32(1):111-116.

20.   Veasey SC, Yeou-Jey H, Thayer P, Fenik P: Murine Multiple Sleep Latency Test:

phenotyping sleep propensity in mice. *Sleep* 2004, 27(3):388-393.

21.   Mitler MM, Walsleben J, Sangal RB, Hirshkowitz M: Sleep latency on the maintenance of

wakefulness test (MWT) for 530 patients with narcolepsy while free of psychoactive

drugs. *Electroencephalography and clinical neurophysiology* 1998, 107(1):33-38.

22.   Sato M, Sagawa Y, Hirai N, Sato S, Okuro M, Kumar S, Kanbayashi T, Shimizu T, Sakai

N, Nishino S: Noninvasive detection of sleep/wake changes and cataplexy-like behaviors

in orexin/ataxin-3 transgenic narcoleptic mice across the disease onset. *Experimental*

neurology 2014, 261:744-751.

23.   Donohue KD, Medonza DC, Crane ER, O'Hara BF: Assessment of a non-invasive high-

throughput classifier for behaviours associated with sleep and wake in mice. *Biomed*

Eng Online 2008, 7:14.

24.   Black SW, Morairty SR, Fisher SP, Chen TM, Warrier DR, Kilduff TS: Almorexant

promotes sleep and exacerbates cataplexy in a murine model of narcolepsy. *Sleep*

2013, 36(3):325-336.

25.   Fujiki N, Cheng T, Yoshino F, Nishino S: Specificity of direct transition from wake to

REM sleep in orexin/ataxin-3 transgenic narcoleptic mice. *Experimental neurology* 2009,

217(1):46-54.

26.   Yaghouby F, Donohue KD, O'Hara BF, Sunderam S: Noninvasive dissection of mouse

sleep using a piezoelectric motion sensor. *J Neurosci Methods* 2016, 259:90-100.

27.   Kuwaki T: Orexinergic modulation of breathing across vigilance states. *Respiratory*

physiology & neurobiology 2008, 164(1-2):204-212.

28.   McShane BB, Galante RJ, Biber M, Jensen ST, Wyner AJ, Pack AI: Assessing REM

sleep in mice using video data. *Sleep* 2012, 35(3):433-442.