Fe oxidation by a fused cytochrome-porin common to diverse Fe-oxidizing bacteria

Clara S. Chan[1,2]*, Sean M. McAllister[2]^, Arkadiy Garber[1]^, Beverly J. Hallahan[1], Sharon Rozovsky[3]

[1]Dept. of Geological Sciences, University of Delaware, Newark, DE

[2]School of Marine Science and Policy, University of Delaware, Newark, DE

[3]Dept. of Chemistry and Biochemistry, University of Delaware, Newark, DE

*Corresponding author (cschan@udel.edu, Dept. of Geological Sciences, University of Delaware, 103 Penny Hall, Newark, DE 19716)

^Equal contributions

Running title: Fe oxidation by a fused cytochrome-porin

Summary

Fe oxidation is one of Earth's major biogeochemical processes, key to weathering, soil

formation, water quality, and corrosion. However, our ability to track the contributions of Fe-

oxidizing microbes is limited by our relatively incomplete knowledge of microbial Fe oxidation

mechanisms, particularly in neutrophilic Fe-oxidizers. The genomes of many Fe-oxidizers

encode homologs to an outer-membrane cytochrome (Cyc2) that has been shown to oxidize Fe in

two acidophiles. Here, we demonstrate the Fe oxidase function of a heterologously expressed

Cyc2 homolog derived from a neutrophilic Fe oxidizer. Phylogenetic analyses show that Cyc2

from neutrophiles cluster together, suggesting a common function. Sequence analysis and

modeling reveal the entire Cyc2 family is defined by a unique structure, a fused cytochrome-

porin, consistent with Fe oxidation on the outer membrane, preventing internal Fe oxide

encrustation. Metatranscriptomes from Fe-oxidizing environments show exceptionally high

expression of *cyc2*, supporting its environmental role in Fe oxidation. Together, these results

provide evidence that *cyc2* encodes Fe oxidases in diverse Fe-oxidizers and therefore can be used

to recognize microbial Fe oxidation. The presence of *cyc2* in 897 genomes suggests that

microbial Fe oxidation may be a widespread metabolism.

Introduction

Fe oxidation occurs in virtually all near-surface environments, producing highly reactive Fe

oxyhydroxides that often control the fate of carbon, phosphorous, and other metals (Borch *et al.*,

2010). It is commonly assumed that abiotic mechanisms are sufficient to account for Fe

oxidation, particularly at near-neutral pH. However, Fe-oxidizing microbes are increasingly

observed in a wide range of environments (Emerson *et al.*, 2010; Kappler *et al.*, 2015), leading

us to ask to what extent microbes drive Fe oxidation. To address this, we need to confidently

identify the Fe oxidase. But unlike other major microbial metabolisms, we have relatively

incomplete knowledge of Fe oxidation pathways (Bird *et al.*, 2011; Hedrich *et al.*, 2011; Ilbert

and Bonnefoy, 2013), and no candidates for a broadly distributed Fe oxidasehave emerged until

now.

Rising interest in Fe-oxidizing microbes has resulted in a surge of Fe-oxidizer sequencing,

including isolate genomes, single cell genomes, metagenomes, and metatranscriptomes (refs. in

Table 1) (Moya-Beltrán *et al.*, 2014; Quaiser *et al.*, 2014; Fukushima *et al.*, 2015; Jewell *et al.*,

2016; Fullerton *et al.*, 2017), enabling us to search for the genes involved in microbial Fe

oxidation. Analyses of chemolithotrophic and phototrophic Fe-oxidizer isolate genomes have

revealed that most possess homologs to *cyc2* (Emerson *et al.*, 2013; Kato *et al.*, 2015; Chiu *et al.*,

2017; Mori *et al.*, 2017; Crowe *et al.*, 2017), which encodes an Fe-oxidizing outer membrane

cytochrome first characterized in *Acidithiobacillus ferrooxidans* (Castelle *et al.*, 2008). Fe

oxidation has also been shown for a distant Cyc2 homolog, Cyt_{572}, purified from an acid mine

drainage *Leptospirillum sp.* (Jeans *et al.*, 2008). In all, this suggests that Cyc2 may be an Fe

oxidase in a wide range of Fe-oxidizers.

To prove this, we need functional information on Cyc2 from neutrophilic chemolithotrophic Fe-

oxidizing bacteria (FeOB). Dark neutral pH environments are prevalent, and to date, these FeOB

have been found in a wide variety of marine, terrestrial, and engineered environments, including

aquifers, soils, sediments, hydrothermal vents, and water treatment systems (Kappler *et al.*, 2015;

Emerson and de Vet, 2015). In these environments, FeOB grow by coupling Fe oxidation to the

reduction of O_2 or nitrate, using this energy to fuel carbon fixation, thus serving as primary

producers (Emerson *et al.*, 2010). Known neutrophilic chemolithoautotrophic FeOB mostly fall

within the marine Zetaproteobacteria (*Mariprofundus spp.*, *Ghiorsea spp.*) and freshwater

Betaproteobacteria (Gallionellales genera *Gallionella*, *Sideroxydans,* and *Ferriphaselus*)

(Emerson *et al.*, 2010; Kato *et al.*, 2015; Mori *et al.*, 2017). All sequenced genomes of

Zetaproteobacteria and Gallionellales FeOB have *cyc2* homologs, and these sequences are

among the closest homologs to one another despite being from separate classes of Proteobacteria,

forming a cluster separate from the acidophilic *cyc2* (Kato *et al.*, 2015; He *et al.*, 2017). A

second potential Fe oxidase gene, *mtoA*, was found in the Gallionellales *Sideroxydans*

lithotrophicus ES-1 (Emerson *et al.*, 2013), and functional and genetic information supports the

role of MtoA and its homolog PioA in Fe oxidation (Jiao and Newman, 2007; Liu *et al.*, 2012).

However, few other FeOB genomes contain *mtoA*, suggesting that Cyc2 is potentially a more

widespread Fe oxidase.

Thus, we set out to demonstrate the function of a Cyc2 from a neutrophilic Fe-oxidizer. We first

analyzed the Cyc2 family phylogeny and then made structure-function predictions, which

informed the design of the gene constructs that we expressed in *E. coli*. To support the Fe

oxidase function, we performed whole cell Fe oxidation assays on Cyc2-expressing *E. coli*. To

determine the environmental relevance, we analyzed *cyc2* expression in a new marine Fe mat

metatranscriptome and reanalyzed a published Fe-rich aquifer metatranscriptome (Jewell *et al.*,

2016). Finally, we compare the genomic distribution and expression of *cyc2* and *mtoA*, to better

understand the relative significance of these two putative Fe oxidases.

Phylogeny of Cyc2

We started by producing a comprehensive phylogenetic tree of Cyc2 sequences acquired from

databases (National Center for Biotechnology Information (NCBI) and

Integrated Microbial Genomes (IMG)). To ensure that we were analyzing true homologs, we

screened the sequences for appropriate length (352 to 587 aa, average/median 446 aa), the

cytochrome c binding motif CXXCH, and beta barrel porin portion (see Cyc2 structure and

conservation below). Our search yielded 897 unique near-full length sequences, which were

reduced to 530 sequences when closely-related sequences were removed. The resulting Cyc2 fell

into three distinct clusters, with sequences distributed amongst various bacterial taxa, largely

Proteobacteria (**Fig. 1, Supplemental File 1**). Cyc2 homologs are present in all well-established

neutrophilic microaerophilic chemolithotrophic FeOB (**Table 1**), many of which are obligate

FeOB that lack other apparent Fe oxidase candidates. These microaerophilic FeOB Cyc2

sequences form a well-supported cluster (Cluster 1 in **Fig. 1**), with the marine Zetaproteobacteria

within one subcluster, and the freshwater Gallionellales forming a separate subcluster that also

includes the neutrophilic photoferrotrophic *Chlorobi*. This clustering suggests a common

function for the Cyc2 in these neutrophilic FeOB.

Beyond neutrophilic FeOB, *cyc2* is found in the genomes of various acidophilic FeOB, which

suggests a common adaptation to Fe oxidation. There are homologs in various acidophilic FeOB

genomes: *Ferrovum spp.*, *Thiomonas spp.*, and Burkholderiales GJ-E10, in addition to the

functionally-verified Cyc2 from *A. ferrooxidans* and *L. ferriphilum*. Unlike the neutrophiles, the

Cyc2 sequences from acidophiles do not form a single cluster, and instead are scattered across

the Cyc2 tree. Notably, the two functionally verified Cyc2 from *A. ferrooxidans* and *L.*

ferriphilum fall in different regions of the tree, Clusters 2 and 3 respectively. This presents the

intriguing possibility that many or all of the Cyc2 homologs are Fe oxidases.

At first glance, a common Fe oxidation pathway for both neutrophiles and acidophiles might not

be expected, due to the drastically different redox potential of Fe(II)/Fe(III) at acidic versus

neutral pH (770 mV at pH 2 vs. 24 mV at pH 7 (Bird *et al.*, 2011; Majzlan, 2013)). However, the

Cyc2 tree shows horizontal transfer between various lineages, and furthermore, certain

neutrophilic and acidophilic FeOB show signs of horizontal transfer of other electron transport

genes. Specifically, *S. lithotrophicus* ES-1, *Gallionella acididurans*, *M. ferrooxydans* PV-1,

Ferrovum spp., and Burkholderiales GJ-E10 all share a gene cassette that includes cbb_3-type

cytochrome c oxidase genes and the periplasmic cytochrome *cyc1* gene (**Supplemental Fig. 1A**).

The conservation in synteny and sequence homology signifies horizontal transfer, strongly

suggesting a common Fe oxidation pathway amongst diverse FeOB, involving Cyc2, Cyc1, and

cytochrome c oxidase (**Supplemental Fig. 1B**).

Cyc2 structure and conservation

To better understand the potential role of Cyc2 homologs and to prepare for functional studies,

we performed sequence and structure predictions, focusing on Cyc2 from FeOB. Despite the

great sequence diversity, the FeOB Cyc2 are all predicted to have a unique structure, a fused

cytochrome-porin. All contain a signal sequence, a c-type cytochrome, and a porin

(**Supplemental Fig. 2**). The signal sequence was predicted by SignalP, indicating that the protein

is exported to the periplasm. The cytochrome portion is identifiable by a single CXXCH heme-

binding motif, and is by far the most conserved part of the sequence (**Fig. 2; Supplemental Fig.**

2, 3). Conserved residues include a AXPXFAR[Q/K][T/Y] motif located 5 amino acids upstream

of the CXXCH heme binding site (AXPXFARQT in Clusters 1 and 2 sequences; AXPXFARKY

in Cluster 3). There is also a PXL motif 4 amino acids downstream of the CXXCH. This PXL

motif can be found in many other cytochromes, such as the structurally characterized MtoD

(Beckwith *et al.*, 2015) and Cyc1 gene in *Acidithiobacillus ferrooxidans* (CYC41 in (Abergel *et*

al., 2003)); the proline and lysine appear to help stabilize the heme (Abergel *et al.*, 2003). In

contrast, the AXPXFAR[Q/K][T/Y] motif is unique to Cyc2, and therefore could be used to

distinguish Cyc2-like outer membrane cytochromes.

The rest of the sequence corresponds to a porin, based on the presence of beta strands predicted

by PSIPRED (McGuffin *et al.*, 2000) (**Supplemental Fig. 4**) and homology matching by

HHpred (Söding *et al.*, 2005). This C-terminal section has low sequence homology, but high

structural homology to the outer membrane phosphate-selective porins OprO and OprP (PDB

structures 4RJW and 2O4V (Moraes *et al.*, 2006; Modi *et al.*, 2015); **Supplemental Table 1**).

Poor sequence conservation is typical of porins (Nikaido, 2003), and since the porin portion

constitutes most of the sequence, this explains why Cyc2 homologs tend to have low amino acid

identity (**Supplemental Fig. 3**). The porin structure is further supported by consistent 3D models

calculated by iTasser, MODELLER, and Phyre (Söding *et al.*, 2005; Zhang, 2008; Kelley *et al.*,

2015; Webb and Sali, 2016) (**Fig. 3**; **Supplemental Fig. 5**), which predict a barrel of 16 beta

strands, a common size porin (Schulz, 2004). Because porins are located in the outer membrane

(Hancock, 1987), Cyc2 is clearly an outer membrane cytochrome. This location is consistent

with our previous observations that Fe oxidation occurs at the cell surface, preventing internal Fe

oxide encrustation (Chan *et al.*, 2011; Comolli *et al.*, 2011).

Cyc2 heterologous expression and functional assay

Cyc2 of neutrophiles are in a distinct cluster from functionally-characterized Cyc2 homologs, so

its function still requires experimental evidence. Neutrophilic FeOB are challenging to grow in

quantities sufficient for protein assays, so we took a heterologous expression approach. The *cyc2*

gene sequence from *Mariprofundus ferrooxydans* PV-1 (*cyc2$_{PV-1}$*) was prepared for expression in

E. coli by (1) codon optimization, (2) replacing the signal sequence with that of the *E. coli* outer

membrane protein OmpA, and (3) adding a StrepII-tag at the C-terminus (**Supplemental Fig.**

6A). The resulting sequence was synthesized, cloned into pMAL-p4X, transformed into *E. coli*

C43(DE3), and co-expressed with the pEC86 plasmid containing the *ccm* cytochrome c

maturation genes under a constitutive promoter, to ensure proper cytochrome maturation under

aerobic conditions (Arslan *et al.*, 1998). The protein appeared to be somewhat toxic to *E. coli*, as

the yield of Cyc2-expressing cells (OD_{600} = 1.1) was much lower than that of cells with empty

vectors (2x-diluted OD_{600} = 1.9) even at low IPTG concentrations of 0.5 mM. Nevertheless,

expression was successful, as shown by western blot using Strep-Tactin antibody against the

protein N-terminal tag (**Fig. 4A**). This band runs close to the expected molecular weight of 43

kDa and contains heme, as established by heme-specific staining (**Fig. 4B**).

We tested the function by assaying Fe oxidation by Cyc2-expressing *E. coli* cells. Because of the

low expression levels, we tested relatively dense cell suspensions (OD=2), washed and

resuspended in fresh LB medium, buffered to pH 6 to help slow abiotic Fe oxidation. Fe(II) was

added from an anoxic stock solution of $FeCl_2$, to a concentration of 100 μM. The dense cell

suspension appeared to stabilize Fe(II), as cells with an empty vector (i.e. plasmid without *cyc2*)

showed considerably slower Fe oxidation relative to cell-free medium (**Supplemental Fig 7A**).

Cyc2-expressing cells did indeed oxidize Fe(II): cells oxidized 41% of the Fe(II) within 2

minutes, and 73% within 10 min (**Fig. 4C**). On further addition of Fe(II) at 45 min, the Cyc2-

expressing cells continued to oxidize Fe(II). In contrast, the empty vector control oxidized 14%

of the Fe(II) in 10 min (**Fig. 4C**). Consistent results were obtained in triplicate experiments from

cells taken from one specific expression stock (**Fig. 4**), as well as replicate assays using cells

from different expressions (**Supplemental Fig. 7B-D**). Azide (3 mM) reduced Fe oxidation by

50%. Azide inhibits cytochromes by binding Fe in heme (Yoshikawa *et al.*, 1998), suggesting

that Cyc2 and/or the terminal cytochrome c oxidase were partially inhibited, though 3 mM azide

may not have been sufficient to completely inhibit such a dense cell suspension. To confirm that

Fe oxidation is due to the Cyc2 cytochrome, we expressed and assayed the porin portion of

$Cyc2_{PV-1}$ ($Cyc2_{porin}$, 37 kDa as expected; **Fig. 4A**). These $Cyc2_{porin}$-expressing cells oxidized

Fe(II) very slowly (17% in 10 min; **Fig. 4C**), demonstrating that the cytochrome is required for

Fe oxidation. Taken together, the data show that Cyc2-expressing *E. coli* accelerate Fe oxidation,

so we conclude that Cyc2 confers the ability to oxidize Fe(II).

Expression of *cyc2* in the environment

If Cyc2 is an Fe oxidase, we would expect high expression of *cyc2* in Fe-oxidizing

environments. To investigate this, we analyzed metatranscriptomes from two ecosystems

dominated by Fe-oxidizing bacteria. We present a new dataset from the Loihi Seamount

(Hawaii) hydrothermal vent Fe microbial mat and reanalyze an existing dataset from an Fe-rich

alluvial aquifer in Rifle, Colorado (Jewell *et al.*, 2016).

At the Loihi seamount, Fe-oxidizing microbial mats thrive where hydrothermal vents emit

Fe(II)-rich fluids into oxygenated seawater (up to 700 µM Fe, <3 to 52 µM O_2 in the mats

(Glazer and Rouxel, 2009)). Here, Fe(II) is by far the most abundant electron donor, with

relatively low or localized sulfide (Glazer and Rouxel, 2009). Using a remotely operated vehicle

and syringe-based biomat sampler (Breier *et al.*, 2012) containing RNA Later, we obtained a 17

mL sample of a surface mat for metagenomic and metatranscriptomic analyses. The microbial

community was almost entirely Zetaproteobacteria (94.4% based on metagenome coverage), a

class in which all cultured representatives are microaerophilic FeOB (n=15 (Emerson and

Moyer, 2002; Emerson *et al.*, 2007; McAllister et al., 2011; McBeth et al., 2011; Field *et al.*,

2015; Makita *et al.*, 2016; Mumford *et al.*, 2016; Mori *et al.*, 2017; Chiu *et al.*, 2017; Laufer et

al., 2017; Barco et al., 2017; Beam *et al.,* in press)). One specific Zetaproteobacteria comprised

79.4% of the community. Overall, Zetaproteobacteria were also the most active, representing

90.5% of the mapped transcripts, with the dominant bin accounting for 78.9% of the transcripts.

For this dominant Zetaproteobacteria bin, *cyc2* is among the most highly expressed genes (91-

99% percentile; **Supplemental Fig. 8A**). We looked for genes that indicated other possible

respiratory metabolisms, but found none. This, combined with the high expression of *cyc2*,

strongly suggests that *cyc2* is a key gene in this marine Fe oxidation-based ecosystem.

In contrast to Zetaproteobacteria, freshwater Gallionellaceae genomes contain a wider range of

electron transport genes: *cyc2*, *mtoA*, and in some cases the sulfur oxidation *sox* and reverse *dsr*

genes. To see which were most highly expressed, we reanalyzed existing metatranscriptomes

from an oxidation experiment at the well-studied Rifle aquifer. Situated next to the Colorado

River, this aquifer was the subject of a long-term experimental study on uranium remediation by

Fe- and S-reducing microbes (Williams *et al.*, 2011). Acetate amendments resulted in a highly

reduced zone with a large reservoir of Fe(II) minerals (Williams *et al.*, 2009). Subsequently,

Jewell et al. (2016) re-oxidized some of this Fe(II) by injecting nitrate-amended oxic

groundwater, causing a bloom of Gallionellaceae. As the authors reported, *cyc2* was among the

most highly expressed genes, at 99.99-100th percentile in all three post-injection samples

(**Supplemental Fig. 8B**). However, the *cyc2* expression levels were not compared to *mtoA*, *sox*,

and *dsr* genes, so their relative importance was unclear. Our re-analysis shows that the *cyc2* gene

was expressed at much higher levels than *mtoA*, *sox* genes, and *dsr* genes (**Supplemental Table**

2). In particular, *cyc2* expression was approximately two orders of magnitude higher than *mtoA*

(**Table 2; Supplemental Fig. 9**). This was true in individual bins when both were expressed, and

also in total across all bins expressing these genes. Thus, while both putative Fe oxidases were

expressed when Fe oxidation was stimulated, *cyc2* was clearly preferentially expressed.

We can gain insight into the different niches of *cyc2* and *mtoA* by examining the temporal

expression patterns of different Gallionellaceae spp., (i.e. genomic bins; **Supplemental Fig. 9**).

Four species/bins expressed both *cyc2* and *mtoA* while >5 bins only expressed either *cyc2* or

mtoA. Overall, both putative Fe oxidase genes increased in expression level over time, but *mtoA*

slightly peaked at the 3rd time point, when the aquifer was largely anoxic (as indicated by a low

in terminal oxidase expression). The bin with the highest *mtoA* expression (22.6), showed peak

mtoA expression during this anoxic period. This suggests that *mtoA* may be more useful under

conditions of electron acceptor limitation.

The difference in *cyc2* and *mtoA* expression at the Rifle site led us to ask whether *cyc2* and

mto/mtr homologs commonly co-occur in genomes (*mtrABC* encodes for an Fe reductase system

homologous to the proposed MtoAB Fe oxidase system). We found that although *cyc2* and *mto*

genes co-occur in Gallionellaceae genomes, they are very rarely found in the same genome

(**Supplemental Fig. 10**), suggesting that *cyc2* and *mto/mtr* do indeed correspond to different

niches.

Cyc2 homologs in other Fe-cycling and extracellular electron-transporting organisms

Of the 897 Cyc2 homologs analyzed (represented in **Fig. 1**, fully labeled tree in **Supplemental**

File 1), very few are from genomes of well-established Fe-oxidizing taxa, in part because Fe

oxidation is not typically tested in isolates. This brings up the question of whether or not all

Cyc2 represent Fe oxidases. Although we do not know yet, we can gain insight from examining a

few homologs in other organisms known to cycle Fe and/or engage in extracellular electron

transport.

At least one of the organisms with *cyc2*, *Dechloromonas aromatica* RCB (Coates *et al.*, 2001), is

reported to be an anaerobic FeOB, coupling Fe oxidation with denitrification (Salinero *et al.*,

2009). Nitrate-dependent Fe oxidation is controversial because these organisms often require

organics, so it is not always clear if the Fe oxidation is enzymatic, or indirect via nitrite produced

by heterotrophic denitrification (reviewed by Kappler et al. (2015)). We can now hypothesize

that *cyc2* encodes an Fe oxidase in *D. aromatica*, as well as some other denitrifiers not yet tested

for Fe oxidation. If true, this would provide evidence in favor of enzymatic Fe oxidation in

heterotrophs, and give a mechanism for studying this metabolism in isolates and the

environment.

Another possibility is that Cyc2 functions more generally as a mechanism of extracellular

electron transport (EET), i.e. to transfer electrons to and from a cell via the cell surface. This is

not exclusive of Fe oxidation, as *M. ferrooxydans* PV-1 has been shown to oxidize a cathode

(Summers *et al.*, 2013) (though it is unknown whether $Cyc2_{PV-1}$ is involved). Homologs of *cyc2*

are also present in the genomes of organisms known to conduct EET, but not proven to oxidize

Fe. An example is the Gammaproteobacteria *Tenderia electrophaga*, which is the most active

organism in a stable cathode-oxidizing consortia (Eddie *et al.*, 2017). Curiously, *T. electrophaga*

also has the conserved gene cassette with *cyc1* and cytochrome c oxidases, found in FeOB

(**Supplemental Fig. 1**), but this organism has not yet been isolated or shown to oxidize Fe.

Homologs of *cyc2* are also found in *Geobacter spp.*, specifically *G. bemidjiensis*, *G.*

uraniireducens, and *Geobacter* M18, M21, and Rf64, members of a clade that predominates in

aquifers (Holmes *et al.*, 2007; Merkley *et al.*, 2015). These organisms reduce Fe(III), though it is

thought that this happens via the Omc system. A study on the *G. bemidjiensis* proteome showed

that the Cyc2 homolog Gbem_3353 was expressed at same levels during Fe(III) and fumarate

reduction (Merkley *et al.*, 2015), so function remained unclear, but could involve a more general

EET role.

Conclusions

This study has provided multiple lines of evidence that Cyc2 is an Fe oxidase in diverse FeOB,

including the first functional evidence that neutrophilic FeOB Cyc2 oxidizes Fe. Verifying the

function of neutrophilic FeOB Cyc2 was important because this cluster of Cyc2 sequences is

distinct and distant from the previously characterized Cyc2 and Cyt_{572} from acidophiles. It is still

not clear if *cyc2* is differentially expressed. If so, we would have a genetic marker of microbial

Fe oxidation activity, which is otherwise difficult to distinguish from abiotic Fe oxidation at

circumneutral pH. If not, the *cyc2* gene is still a valuable way of recognizing the Fe oxidation

potential in genomes and transcriptomes.

It is striking that there are so many Cyc2 homologs, including many from organisms not known

to oxidize Fe. Phylogenetic and genomic analyses show that *cyc2* has been horizontally

transferred between known FeOB and with other organisms. The addition of Cyc2 alone

conferred some Fe oxidation ability on *E. coli*, with Cyc2 likely interfacing with the existing

electron transport system. This suggests that acquisition of *cyc2* alone can allow an organism to

oxidize Fe. If this is generally true, the abundance of Cyc2 homologs suggests that microbial Fe

oxidation is more widespread than we currently recognize.

Experimental Procedures

Cyc2 phylogeny and amino acid identity calculations

Sequences with homology to Cyc2 were collected using blastp (Camacho *et al.*, 2009) against

the NCBI and IMG databases (maximum e-value $1*10^{-5}$). Query sequences were chosen to

represent all major groupings of the Cyc2 tree, including *Mariprofundus ferrooxydans* PV-1,

Gallionella capsiferriformans ES-2, *Acidithiobacillus ferrooxidans* ATCC 23270, *Tenderia*

electrophaga, and *Geobacter* sp. FRC-32. To remove identical sequences, the resulting 2,413

sequences were clustered at 100% identity using CD-HIT, reducing the database to 977 cluster

representatives (Li and Godzik, 2006). These sequences were imported into Geneious v.7.1.7,

where they were aligned using MUSCLE (Edgar, 2004). The resulting alignment was used to

filter out sequences of partial length, resulting in 897 sequences. Alignment columns with greater

than 30% gaps were removed, and a maximum likelihood phylogenetic tree was built using

RAxML (392 alignment columns, 100 bootstraps, CAT model of rate heterogeneity, JTT amino

acid substitution model (Stamatakis, 2014)). From the resulting tree, sequences that were closely

related and highly sampled (primarily from the *Burkholderia* and *Xanthomonas* genera) were

removed, resulting in 530 full-length sequences that were re-aligned, and a final phylogenetic

tree was built in RAxML (357 alignment columns, 300 bootstraps, CAT and JTT models). The

resulting phylogenetic tree was colored and names customized using the Iroki program (Moore *et*

al., 2017).

To calculate amino acid identities of the Cyc2 homologs, pairwise alignments between the 530

full-length Cyc2 sequences were constructed using Muscle (Edgar, 2004). Amino acid identity

(AAI) values were calculated from these pairwise alignments based on the full-length sequence,

as well as the cytochrome and porin domains separately. The cytochrome domain was defined as

the conserved region starting 14 residues upstream from the heme binding site, and ending 21

residues downstream. The rest of the sequence downstream of this was defined as the β-barrel

porin domain. The AAI values were imported into R and a histogram was plotted using the

ggplot2 package (Wickham, 2009).

Structural modeling

Signal peptides were predicted using SignalP(Petersen *et al.*, 2011) and secondary elements were

predicted using PSIPRED (McGuffin *et al.*, 2000). For identification of structural homologs to

Cyc2, we uploaded sequences to the HHPRED tool, available as part of the Max Planck Institute

Bioinformatics Toolkit (Söding *et al.*, 2005). Information for the structural homologs are

compiled in Supplementary Table 1. Structural modeling was carried out by

HHpred/MODELLER (Söding *et al.*, 2005; Webb and Sali, 2016), iTasser (Zhang, 2008), and

Phyre (Kelley *et al.*, 2015). The structural models from all three platforms were found to be in

close agreement.

Cloning and heterologous expression of Cyc2

The sequence of *cyc2* was optimized for expression in *Escherichia coli*, and the signal sequence

replaced with the signal sequence of a native *E. coli* gene, *ompA* (sequences in Supplementalry

Fig. 6). The gene was synthesized by Genscript (Piscataway, NJ, USA). The *cyc2* gene was

cloned into the EcoRI/ HindIII sites of the pMal-p4X plasmid (with the *malE* gene removed).

This *cyc2*-containing plasmid was co-transformed into *E. coli* C43(DE3) with pEC86, a plasmid

containing the cytochrome c maturation (*ccm*) genes under a constitutive promoter, to ensure

heme insertion into Cyc2 (Arslan *et al.*, 1998). We also co-transformed pEC86 and the pMalp4X

plasmid without the *cyc2* gene (and without *malE*) as an empty-vector control; cells with the

empty vector control received the same treatment throughout heterologous expression and Fe-

oxidation assays. For expression of *cyc2*, *E. coli* was grown aerobically at 37°C (with shaking at

200 RPM) in Lysogeny Broth (LB), buffered with 10 mM 2-(N-morpholino)ethanesulfonic acid

(MES), pH 6; in addition, the medium also contained ampicillin (100 µg/mL) for propagation of

the pMal-p4X plasmid, and chloramphenicol (30 µg/mL) for the pEC86 plasmid. After reaching

mid-log phase (2.5 h), cultures were amended with 0.5 mM isopropyl β-D-1-

thiogalactopyranoside (IPTG) for de-repression of the lac operon to induce *cyc2* expression;

induction proceeded for 20 hours at 18°C with shaking at ~200 RPM. Expression of the Cyc2$_{porin}$

was carried out in an identical manner, except for the absence of the pEC86 plasmid from cells

expressing this construct, and therefore the absence of chloramphenicol from the media.

SDS-PAGE, western and heme staining. Cells were harvested and from cultures before IPTG

induction and following 20 hours after induction. Biomass was adjusted to an optical density of

0.6 for the porin-only control and 1.2 for the empty-vector and Cyc2-expressing cells. Cells were

lysed by re-suspension in 100 µL of 5X sodium dodecyl sulfate (SDS) running buffer (125 mM

Tris, 1.25 M glycine, 0.5% SDS, pH 8.3) and passing the resuspension through a 27.5-gauge

needle 10 times. Cells were then combined with gel loading buffer (50 mM Tris-HCl, 12.5 mM

EDTA, 2% SDS, 10% glycerol, 0.02% bromophenol blue, pH 6.8), and centrifuged for 10 min at

15 000 x g. Fifteen µL of the sample was then loaded onto a 16% Tris-glycine SDS-PAGE gel,

and ran at 100 V for 30 min, then 160 V for 40 min. The gel was then either Coomasie-stained (1

g Coomasie Brilliant Blue Stain in 10% acetic acid, 40% ethanol), or transferred to a PVDF

membrane for heme stain and western blot. For heme and StrepII-tag detection, the SDS-PAGE

gel was transferred to a PVDF membrane at 30 V for 16 h (4°C) in transfer buffer (25 mM Tris,

192 mM glycine, 20% methanol, pH 7.2). Heme peroxidase activity was assessed by washing the

membrane with TBST buffer (20 mM Tris, 137 mM NaCl, 0.1% Tween-20, pH 7.6), and

incubating for 30 minutes with Pierce ECL luminol and substrate (Carlson *et al.*, 2013), before

imaging on the Typhoon FLA 9500 (GE Healthcare Life Sciences). For StrepII-tag detection, the

PVDF membrane was blocked for one hour with 5% bovine serum albumin (BSA) and 50

µg/mL avidin (from egg white) in TBST buffer, rinsed with TBST, and incubated for one hour

with Precision Protein Streptactin-HRP conjugate (1:60,000 dilution, Biorad). The membrane

was then washed with four 15-minute washes in TBST, then incubated with Pierce ECL luminol

and substrate for 5 minutes before imaging on the Typhoon FLA 9500.

Fe oxidation assay

After induction of Cyc2 in *E. coli*, triplicate cultures were centrifuged at 3200 x g, and washed

with sterile LB (supplemented with 10 mM MES, pH 6) before resuspension and incubation in

50 mL glass beakers in the same medium at an OD of 2 (final volume = 25 mL); these cell

suspensions were stirred to ensure homogeneity and aeration. In the case of azide experiments,

75 µL of a fresh 1M sodium azide stock solution was added to 25 mL cell suspension (to a final

concentration of 3 mM), and incubated for 5 min. In all experiments, $FeCl_2$ was added from an

anoxic, filter-sterilized 100 mM stock solution to a target concentration of 100 µM. Ferrozine

measurements were taken over a period of one hour (ferrozine method adapted from (Stookey,

1970)). At each time point, a sample was taken from each of the triplicate cultures, and

centrifuged at 15,000 x g for 20-30 s to remove cells. A 150 µL portion of the supernatant was

then combined with 40 µL of 1.225 mM ferrozine, 50 µL of 6.87 M acetate buffer, and 10 µL of

H_2O inside 200 µL 96-well plates, and incubated for 15 min in the dark before absorption was

measured at 562 nm using a plate reader (Perkin Elmer 1420 Multilabel Counter Victor[3]V). pH

of the cell suspensions was monitored before and after each assay, and was found to be stable at

pH 6 (within one-tenth of a pH unit).

Loihi Fe mat sampling and metatranscriptome analysis

Sampling. Sample J2-674-BM1-C123456 was collected from the Pohaku vents (Mkr 57) at Loihi

Seamount, Hawaii using the Jason II remotely operated vehicle in 2013. Using the mat sampler

designed by Breier et al. (2012), 60 mL of the top 1 cm of Fe mat was collected into each of six

syringes, each pre-loaded with 50 mL of RNALater (Ambion, United States) for immediate

DNA/RNA preservation. Upon recovery, samples were allowed to settle (~17 mL total mat

material), overlying fluids were decanted, and samples were stored at -80˚C until extraction.

Metagenome and metatranscriptome sequencing. Genomic DNA was extracted using the

FastDNA SPIN kit for soil (MP Bio, Solon, OH, USA) following the manufacturer's protocol,

with the addition of 250 μL 0.5 M sodium citrate pH 5.8 prior to cell lysis. RNA was extracted

using the NucleoSpin RNA kit (Macherey-Nagel, Bethlehem, PA, USA) following the

manufacturer's protocol. Metagenomic sequencing was performed on sample J2-674-BM1-C3.

An Illumina library with paired reads of 251 bp length with 450 bp insert size was prepared

following standard protocols and sequenced on a HiSeq 2500 (Illumina Inc., San Diego, CA).

Metatranscriptomic sequencing was performed on sample J2-674-BM1-C6. Ribosomal RNA-

depleted total RNA (Ribo-Zero, Epicenter, Madison, WI) was prepared for sequencing using

Illumina's NEXTflex Rapid RNA-Seq Library Prep Kit (Bioo Scientific, Austin, TX), and was

sequenced on a HiSeq 2500 producing single-end 51 bp reads.

Metagenome assembly and binning. Metagenomic reads were quality controlled (QC'ed) using

trimmomatic and merged using FLASH (see pipeline: https://github.com/mooreryan/qc). QC'ed

reads were then assembled using metaSPAdes (Nurk *et al.*, 2017), with a k-mer sweep from 21

to 127. In total 82,765 contigs were produced, from 128 bp to 79,748 bp (average 1,203 bp).

Contigs were then QC'ed so that only contigs with at least 1X coverage over 90% of their length

were used. Contigs over 2,000 bp in length (10,352 total) were binned using Binsanity (Graham

et al., 2017), CONCOCT (Alneberg *et al.*, 2014), MaxBin (Wu *et al.*, 2016), and MetaBAT

(Kang *et al.*, 2015). The best resulting bins were chosen using DAS Tool (Sieber *et al.*, 2017).

Bin completeness and redundancy were calculated using CheckM (Parks *et al.*, 2015). The

relative abundance of each bin was calculated using a length-normalized average of contig read

coverage.

Manual bin curation. Loihi metagenome sample 674-BM1-C3 was dominated by a single

Zetaproteobacteria OTU2 bin (S1_binsanity019), accounting for 79.4% of the total binned

average read coverage (50.3% including unbinned contigs). This bin was estimated by CheckM

to be 92.7% complete (3.03% redundancy, 0% strain heterogeneity), yet lacked any protein

BLAST hits to *cyc2*. Because previous research has identified *cyc2* within at least 9 genomic

representatives from ZOTU2 (3 SAGs, 6 MAGs (Field *et al.*, 2015; Fullerton *et al.*, 2017)), we

used two methods to find the cyc2 belonging to this bin: 1) subsampled reads for a simplified

assembly and 2) used information from the assembly graph to locate cyc2 connected with this

dominant bin. Quadruplicate, randomly-sampled read subsets at 10%, 2%, and 1% of the total

number of quality-controlled reads were independently assembled and binned to simplify the

assembly of the dominant ZOTU2 bin (starting at 1,352X coverage). Subsampling in this way

allowed us to increase the quality of the dominant ZOTU2 bin, with a maximum completeness of

98.1% (2.27% redundancy, 0% strain heterogeneity), though unfortunately cyc2 was again not

recovered from any of the binned contigs. However, the subsampled assembly was helpful in

recovering *cyc2* through use of the simplified assembly graph. Using blastp, 12 contigs were

identified that had homology to Cyc2. All twelve of these contigs were found within one section

of the assembly graph, connected through a minimal k-mer overlap of 21 bp to the assembly

network containing the binned contigs of interest. No other bins were contained within this

assembly graph network. Taking the section of the assembly graph containing these *cyc2* contigs,

overlap consensus assembly produced five unique contigs. Four of these contigs had sufficient

length to confirm their clustering within the dominant ZOTU2 bin using VizBin (Laczny *et al.*,

2015). These four contigs were combined with the dominant subsampled bin and used in further

analysis.

Metatranscriptome recruitment and analysis. Metatranscriptome reads were recruited to

metagenomic contigs using bowtie2 (Langmead and Salzberg, 2012). Expression estimates were

calculated by dividing the total read count by the gene length and total sequencing effort (reads

per thousand bp per million reads; RPKM). The relative abundance of the expression in each bin

was calculated as the percent of total reads mapping to that bin, normalized to bin length. RPKM

expression estimates from all expressed genes were then imported into R and plotted using the

ggplot2 package (Wickham, 2009).

Nucleotide submission. Raw sequence data were submitted to the sequence read archive (SRA) at

the National Center for Biotechnology Information (NCBI), with all appropriate metadata under

project accession number PRJNA412510.

Reanalysis of Rifle aquifer metatranscriptome

To quantify *cyc2* and other gene expression in an FeOB-dominated terrestrial ecosystem, we

extracted information from the supplementary dataset of Jewell et al.(2016), a

metatranscriptomic/metagenomic study of a nitrate/O_2-amended ferruginous Rifle aquifer. This

dataset consists of over 200,000 translated open reading frames (ORFs), grouped by bin and

associated with gene expression data in units of RPKM (reads per thousand base pairs per

million reads). From this dataset, amino acid sequences corresponding to ORFs in

Gallionellaceae bins 22.1-22.9 were used as a protein sequence database, which we used for

blastp to search for Cyc2, and Mto, Sox, Dsr protein sequences (**Supplementary table 2**).

RPKM expression values were then imported into R, log transformed, and plotted using the

ggplot2 package.

cyc2 **and** *mto/mtr* **gene distribution in genomes**

To look for *cyc2* and *mtoAB/mtrABC* gene distribution in genomes, we used blastp against the

non-redundant NCBI database to search for homologs of these genes from a representative set of

organisms: *Sideroxydans lithotrophicus* ES-1, *Shewanella oneidensis* MR-1, *Rhodoferax*

ferrireducens T118, *Magnetospirillum magneticum* AMB-1, and *Rhodopseudomonas palustris*

TIE-1. Blast results were analyzed using a custom Python script to identify which *cyc2* and

mto/*mtr* homologs co-occurred within the same genomes. Results were imported in R and plotted

using the VennDiagram package.

Conserved gene cassette identification

We used a custom Python script to identify organisms that encode the conserved gene cassette

described by Field et al. (2015). This cassette includes 2 subunits of the cbb3-type cytochrome c

oxidase, *cyc1*, spermidine synthase, ferredoxin, and several other c-type cytochromes; we blasted

these cassette genes against the non-redundant NCBI database (Release 84), and clustered close

homologs according to gene ID, which allowed us to detect genomes that included genes of

interest in close proximity, defined as within 20 genes of each other.

Acknowledgments

We thank Shawn Polson for assistance in bioinformatic analyses and the crews of R/V

Thompson and ROV Jason for support during sampling at the Loihi Seamount. This research

was funded by the National Science Foundation (EAR-1151682, OCE-1155290) and the Office

of Naval Research (N00014-17-1-2640). The authors declare no conflict of interest.

References

Abergel, C., Nitschke, W., Malarte, G., Bruschi, M., Claverie, J.-M., and Giudici-Orticoni, M.-T.
(2003) The Structure of *Acidithiobacillus ferrooxidans c4*-cytochrome. *Structure* **11**: 547–
555.

Alneberg, J., Bjarnason, B.S., de Bruijn, I., Schirmer, M., Quick, J., Ijaz, U.Z., et al. (2014)
Binning metagenomic contigs by coverage and composition. *Nat Methods* **11**: 1144–1146.

Arslan, E., Schulz, H., Zufferey, R., Künzler, P., and Thöny-Meyer, L. (1998) Overproduction of
the *Bradyrhizobium japonicum c*-type cytochrome subunits of the cbb3 oxidase in
Escherichia coli. Biochemical and Biophysical Research Communications **251**: 744–747.

Barco, R.A., Emerson, D., Sylvan, J.B., Orcutt, B.N., Meyers, M.E.J., Ramírez, G.A., et al.
(2015) The proteomic profile of an obligate iron-oxidizing chemolithoautotroph reveals new
insight into microbial iron oxidation. **81**: 5927–5937.

Barco, R.A., Hoffman, C.L., Ramírez, G.A., Toner, B.M., Edwards, K.J., and Sylvan, J.B.
(2017) *In-situ* incubation of iron-sulfur mineral reveals a diverse chemolithoautotrophic
community and a new biogeochemical role for *Thiomicrospira. Environ. Microbiol.* **19**:
1322–1337.

Beam, J., Scott, J.J., McAllister, S.M., Chan, C.S., McManus, J., Meysman, F., and Emerson, D.
Potential for biological rejuvenation of iron oxides in bioturbated marine sediments. *in press*.

Beckwith, C.R., Edwards, M.J., Lawes, M., Shi, L., Butt, J.N., Richardson, D.J., and Clarke,
T.A. (2015) Characterization of MtoD from *Sideroxydans lithotrophicus*: a cytochrome c

electron shuttle used in lithoautotrophic growth. *Front Microbiol* **6**:.

Bird, L.J., Bonnefoy, V., and Newman, D.K. (2011) Bioenergetic challenges of microbial iron
metabolisms. *Trends in Microbiology* **19**: 330–340.

Borch, T., Kretzschmar, R., Kappler, A., Cappellen, P.V., Ginder-Vogel, M., Voegelin, A., and
Campbell, K. (2010) Biogeochemical redox processes and their impact on contaminant
dynamics. *Environ Sci Technol* **44**: 15–23.

Breier, J.A., Gomez-Ibanez, D., Reddington, E., Huber, J.A., and Emerson, D. (2012) A
precision multi-sampler for deep-sea hydrothermal microbial mat studies. *Deep Sea Res Pt I*
70: 83–90.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden,
T.L. (2009) BLAST+: architecture and applications. *BMC Bioinformatics 2016 17:1* **10**: 421.

Cardenas, J.P., Lazcano, M., Ossandon, F.J., Corbett, M., Holmes, D.S., and Watkin, E. (2014)
Draft genome sequence of the iron-oxidizing acidophile *Leptospirillum ferriphilum* type
strain DSM 14647. *Genome Announc.* **2**: e01153–14–e01153–14.

Carlson, H.K., Clark, I.C., Blazewicz, S.J., Iavarone, A.T., and Coates, J.D. (2013) Fe(II)
oxidation is an innate capability of nitrate-reducing bacteria that involves abiotic and biotic
reactions. *Journal of Bacteriology* **195**: 3260–3268.

Castelle, C., Guiral, M., Malarte, G., Ledgham, F., Leroy, G., Brugna, M., and Giudici-Orticoni,
M.T. (2008) A new iron-oxidizing/O_2-reducing supercomplex spanning both inner and outer
membranes, isolated from the extreme acidophile *Acidithiobacillus ferrooxidans*. *Journal of
Biological Chemistry* **283**: 25803–25811.

Chan, C.S., Fakra, S.C., Emerson, D., Fleming, E.J., and Edwards, K.J. (2011) Lithotrophic iron-
oxidizing bacteria produce organic stalks to control mineral growth: implications for
biosignature formation. *ISME J* **5**: 717–727.

Chiu, B.K., Kato, S., McAllister, S.M., Field, E.K., and Chan, C.S. (2017) Novel pelagic iron-
oxidizing Zetaproteobacteria from the Chesapeake Bay oxic–anoxic transition zone. *Front
Microbiol* **8**: 4206.

Coates, J.D., Chakraborty, R., Lack, J.G., O'Connor, S.M., Cole, K.A., Bender, K.S., and
Achenbach, L.A. (2001) Anaerobic benzene oxidation coupled to nitrate reduction in pure
culture by two strains of *Dechloromonas*. *Nature* **411**: 1039–1043.

Comolli, L.R., Luef, B., and Chan, C.S. (2011) High-resolution 2D and 3D cryo-TEM reveals
structural adaptations of two stalk-forming bacteria to an Fe-oxidizing lifestyle. *Environ
Microbiol* **13**: 2915–2929.

Crowe, S.A., Hahn, A.S., Morgan-Lang, C., Thompson, K.J., Simister, R.L., Llirós, M., et al.
(2017) Draft genome sequence of the pelagic photoferrotroph *Chlorobium
phaeoferrooxidans*. *Genome Announc.* **5**: e01584–16.

Eddie, B.J., Wang, Z., Hervey, W.J., IV, Leary, D.H., Malanoski, A.P., Tender, L.M., et al.
(2017) Metatranscriptomics supports the mechanism for biocathode electroautotrophy by
"Candidatus *Tenderia electrophaga*." *mSystems* **2**: e00002–17.

Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Research* **32**: 1792–1797.

Emerson, D. and de Vet, W. (2015) The role of FeOB in engineered water ecosystems: a review.
J Am Water Works Ass **107**: E47–E57.

Emerson, D., Field, E.K., Chertkov, O., Davenport, K.W., Goodwin, L., Munk, C., et al. (2013)
Comparative genomics of freshwater Fe-oxidizing bacteria: implications for physiology,
ecology, and systematics. *Front Microbiol* **4**: 254.

Emerson, D., Fleming, E.J., and McBeth, J.M. (2010) Iron-oxidizing bacteria: an environmental
and genomic perspective. *Annu Rev Microbiol* **64**: 561–583.

Emerson, D. and Moyer, C.L. (2002) Neutrophilic Fe-oxidizing bacteria are abundant at the
Loihi Seamount hydrothermal vents and play a major role in Fe oxide deposition. *Appl.*
Environ. Microbiol. **68**: 3085–3093.

Emerson, D., Rentz, J.A., Lilburn, T.G., Davis, R.E., Aldrich, H., Chan, C., and Moyer, C.L.
(2007) A novel lineage of Proteobacteria involved in formation of marine Fe-oxidizing
microbial mat communities. *PLoS ONE* **2**: e667.

Fabisch, M., Beulig, F., Akob, D.M., and Kusel, K. (2011) New *Thiomonas* and *Bordetella*
strains involved in iron oxidation at a slightly acidic, heavy metal contaminated creek.
Mineralogical Magazine **75**: 825.

Field, E.K., Sczyrba, A., Lyman, A.E., Harris, C.C., Woyke, T., Stepanauskas, R., and Emerson,
D. (2015) Genomic insights into the uncultivated marine Zetaproteobacteria at Loihi
Seamount. *The ISME Journal* **9**: 857–870.

Fukushima, J., Tojo, F., Asano, R., Kobayashi, Y., Shimura, Y., Okano, K., and Miyata, N.
(2015) Complete genome sequence of the unclassified iron-oxidizing, chemolithoautotrophic
Burkholderiales bacterium GJ-E10, isolated from an acidic river. *Genome Announc.* **3**:
e01455–14.

Fullerton, H., Hager, K.W., McAllister, S.M., and Moyer, C.L. (2017) Hidden diversity revealed
by genome-resolved metagenomics of iron-oxidizing microbial mats from Loihi Seamount,
Hawaii. *The ISME Journal* **11**: 1900–1914.

Glazer, B.T. and Rouxel, O.J. (2009) Redox speciation and distribution within diverse iron-
dominated microbial habitats at Loihi Seamount. *Geomicrobiol J* **26**: 606–622.

Graham, E.D., Heidelberg, J.F., and Tully, B.J. (2017) BinSanity: unsupervised clustering of
environmental microbial assemblies using coverage and affinity propagation. *PeerJ* **5**:
e3035.

Hallberg, K.B., González-Toril, E., and Johnson, D.B. (2009) Acidithiobacillus ferrivorans, sp.
nov.; facultatively anaerobic, psychrotolerant iron-, and sulfur-oxidizing acidophiles isolated
from metal mine-impacted environments. *Extremophiles* **14**: 9–19.

Hancock, R.E. (1987) Role of porins in outer membrane permeability. *Journal of Bacteriology*
169: 929–933.

He, S., Barco, R.A., Emerson, D., and Roden, E.E. (2017) Comparative genomic analysis of
neutrophilic iron(II) oxidizer genomes for candidate genes in extracellular electron transfer.
Front Microbiol **8**: W400.

He, S., Tominski, C., Kappler, A., Behrens, S., and Roden, E.E. (2016) metagenomic analyses of
the autotrophic Fe(II)-oxidizing, nitrate-reducing enrichment culture KS. **82**: 2656–2668.

Hedrich, S., Schlömann, M., and Johnson, D.B. (2011) The iron-oxidizing proteobacteria.
Microbiology **157**: 1551–1564.

Heising, S., Richter, L., LUDWIG, W., and SCHINK, B. (1999) *Chlorobium ferrooxidans* sp.
nov., a phototrophic green sulfur bacterium that oxidizes ferrous iron in coculture with a
"*Geospirillum*" sp. strain. *Arch Microbiol* **172**: 116–124.

Holmes, D.E., O'Neil, R.A., Vrionis, H.A., N'Guessan, L.A., Ortiz-Bernad, I., Larrahondo, M.J.,
et al. (2007) Subsurface clade of Geobacteraceae that predominates in a diversity of Fe(III)-
reducing subsurface environments. *The ISME Journal* **1**: 663–677.

Ilbert, M. and Bonnefoy, V. (2013) Insight into the evolution of the iron oxidation pathways.
Biochim. Biophys. Acta **1827**: 161–175.

Jeans, C., Singer, S.W., Chan, C.S., VerBerkmoes, N.C., Shah, M., Hettich, R.L., et al. (2008)
Cytochrome 572 is a conspicuous membrane protein with iron oxidation activity purified
directly from a natural acidophilic microbial community. *The ISME Journal* **2**: 542–550.

Jewell, T.N.M., Karaoz, U., Brodie, E.L., Williams, K.H., and Beller, H.R. (2016)
Metatranscriptomic evidence of pervasive and diverse chemolithoautotrophy relevant to C,
S, N and Fe cycling in a shallow alluvial aquifer. *ISME J* **10**: 2106–2117.

Jiao, Y. and Newman, D.K. (2007) The pio operon is essential for phototrophic Fe(II) oxidation
in *Rhodopseudomonas palustris* TIE-1. *Journal of Bacteriology* **189**: 1765–1773.

Johnson, D.B., Hallberg, K.B., and Hedrich, S. (2014) Uncovering a microbial enigma: isolation
and characterization of the streamer-generating, iron-oxidizing, acidophilic bacterium
"Ferrovum myxofaciens". **80**: 672–680.

Kang, D.D., Froula, J., Egan, R., and Wang, Z. (2015) MetaBAT, an efficient tool for accurately
reconstructing single genomes from complex microbial communities. *PeerJ* **3**: e1165.

Kappler, A., Emerson, D., Gralnick, J.A., Roden, E.E., and Muehe, E.M. (2015)
Geomicrobiology of Iron. In, Ehrlich,H.L. and Newman,D.K. (eds), *Ehrlich's*

Geomicrobiology. Boca Raton, US, pp. 343–399.

Kato, S., Ohkuma, M., Powell, D.H., Krepski, S.T., Oshima, K., Hattori, M., et al. (2015)
Comparative genomic insights into ecophysiology of neutrophilic, microaerophilic iron
oxidizing bacteria. *Front Microbiol* **6**: 1265.

Kelley, L.A., Mezulis, S., Yates, C.M., Wass, M.N., and Sternberg, M.J.E. (2015) The Phyre2
web portal for protein modeling, prediction and analysis. *Nat Protoc* **10**: 845–858.

Kupka, D., Liljeqvist, M., Nurmi, P., Puhakka, J.A., Tuovinen, O.H., and Dopson, M. (2009)
Oxidation of elemental sulfur, tetrathionate and ferrous iron by the psychrotolerant
Acidithiobacillus strain SS3. *Research in Microbiology* **160**: 767–774.

Laczny, C.C., Sternal, T., Plugaru, V., Gawron, P., Atashpendar, A., Margossian, H.H., et al.
(2015) VizBin - an application for reference-independent visualization and human-
augmented binning of metagenomic data. *Microbiome 2017 5:1* **3**: 1.

Langmead, B. and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**: 357–359.

Laufer, K., Nordhoff, M., Halama, M., Martinez, R.E., Obst, M., Nowak, M., et al. (2017)
Microaerophilic Fe(II)-oxidizing Zetaproteobacteria isolated from low-Fe marine coastal
sediments: Physiology and characterization of their twisted stalks. *Appl. Environ.*
Microbiol. **83**: e03118-16.

Li, W. and Godzik, A. (2006) Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**: 1658–1659.

Liu, J., Wang, Z., Belchik, S.M., Edwards, M.J., Liu, C., Kennedy, D.W., et al. (2012)
Identification and characterization of MtoA: A decaheme c-type cytochrome of the
neutrophilic Fe(II)-oxidizing bacterium *Sideroxydans lithotrophicus* ES-1. *Front Microbiol*
3: 37.

Majzlan, J. (2013) Minerals and aqueous species of iron and manganese as reactants and
products of microbial metal respiration. In, Gescher,J. and Kappler,A. (eds), *Microbial Metal*
Respiration. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 1–28.

Makita, H., Tanaka, E., Mitsunobu, S., Miyazaki, M., Nunoura, T., Uematsu, K., et al. (2016)
Mariprofundus micogutta sp. nov., a novel iron-oxidizing zetaproteobacterium isolated from
a deep-sea hydrothermal field at the Bayonnaise knoll of the Izu-Ogasawara arc, and a
description of Mariprofundales ord. nov. and Zetaproteobacteria classis nov. *Arch Microbiol*
199: 335–346.

McAllister, S.M., Davis, R.E., McBeth, J.M., Tebo, B.M., Emerson, D., and Moyer, C.L. (2011)
Biodiversity and emerging biogeography of the neutrophilic iron-oxidizing
Zetaproteobacteria. *Appl. Environ. Microbiol.* **77**: 5445–5457.

McBeth, J.M., Little, B.J., Ray, R.I., Farrar, K.M., and Emerson, D. (2011) Neutrophilic iron-

oxidizing "Zetaproteobacteria" and mild steel corrosion in nearshore marine
environments. *Appl. Environ. Microbiol.* **77**: 1405–1412.

McGuffin, L.J., Bryson, K., and Jones, D.T. (2000) The PSIPRED protein structure prediction
server. *Bioinformatics* **16**: 404–405.

Merkley, E.D., Wrighton, K.C., Castelle, C.J., Anderson, B.J., Wilkins, M.J., Shah, V., et al.
(2015) Changes in protein expression across laboratory and field experiments in *Geobacter*
bemidjiensis. *J. Proteome Res.* **14**: 1361–1375.

Modi, N., Ganguly, S., Bárcena-Uribarri, I., Benz, R., van den Berg, B., and Kleinekathöfer, U.
(2015) Structure, dynamics, and substrate specificity of the OprO Porin from *Pseudomonas*
aeruginosa. *Biophys J* **109**: 1429–1438.

Moore, R.M., Harrison, A.O., McAllister, S.M., Marine, R.L., Chan, C.S., and Wommack, K.E.
(2017) Iroki: automatic customization for phylogenetic trees. *bioRxiv* 106138.

Moraes, T.F., Bains, M., Hancock, R.E.W., and Strynadka, N.C.J. (2006) An arginine ladder in
OprP mediates phosphate-specific transfer across the outer membrane. *Nat Struct Mol Biol*
14: 85–87.

Mori, J.F., Scott, J.J., Hager, K.W., Moyer, C.L., Küsel, K., and Emerson, D. (2017)
Physiological and ecological implications of an iron- or hydrogen-oxidizing member of the
Zetaproteobacteria, Ghiorsea bivora, gen. nov., sp. nov. *ISME J* **11**: 2624–2636.

Moya-Beltrán, A., Cárdenas, J.P., Covarrubias, P.C., Issotta, F., Ossandon, F.J., Grail, B.M., et
al. (2014) Draft genome sequence of the nominated type strain of "*Ferrovum myxofaciens,*"
an acidophilic, iron-oxidizing Betaproteobacterium. *Genome Announc.* **2**: e00834–14.

Mumford, A.C., Adaktylou, I.J., and Emerson, D. (2016) Peeking under the Iron Curtain:
Development of a microcosm for imaging colonization of steel surfaces by *Mariprofundus*
sp. DIS-1, an oxygen tolerant Fe-oxidizing bacterium. AEM.01990–16.

Nikaido, H. (2003) Molecular basis of bacterial outer membrane permeability revisited.
Microbiol Mol Biol R **67**: 593–656.

Nurk, S., Meleshko, D., Korobeynikov, A., and Pevzner, P.A. (2017) metaSPAdes: a new
versatile metagenomic assembler. *Genome Res.* **27**: 824–834.

Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., and Tyson, G.W. (2015) CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res.* **25**: 1043–1055.

Petersen, T.N., Brunak, S., Heijne, von, G., and Nielsen, H. (2011) SignalP 4.0: discriminating
signal peptides from transmembrane regions. *Nat Methods* **8**: 785–786.

Quaiser, A., Bodi, X., Dufresne, A., Naquin, D., Francez, A.-J., Dheilly, A., et al. (2014)
Unraveling the stratification of an iron-oxidizing microbial mat by metatranscriptomics.

PLoS ONE **9**: e102561.

Salinero, K., Keller, K., Feil, W.S., Feil, H., Trong, S., Di Bartolo, G., and Lapidus, A. (2009)
Metabolic analysis of the soil microbe Dechloromonas aromatica str. RCB: indications of a
surprisingly complex life-style and cryptic anaerobic pathways for aromatic degradation.
BMC Genomics **10**: 351.

Schulz, G.E. (2004) The structures of general porins. In *Bacterial and Eukaryotic Porins:*
Structure, Function, Mechanism. Benz, R. (ed). Weinheim, Germany: Wiley-VCH, pp. 25-
40.

Sieber, C.M.K., Probst, A.J., Sharrar, A., Thomas, B.C., Hess, M., Tringe, S.G., and Banfield,
J.F. (2017) Recovery of genomes from metagenomes via a dereplication, aggregation, and
scoring strategy. *bioarXiv* **107789**: 1–24.

Singer, E., Emerson, D., Webb, E.A., Barco, R.A., Kuenen, J.G., Nelson, W.C., et al. (2011)
Mariprofundus ferrooxydans PV-1 the first genome of a marine Fe(II) oxidizing
Zetaproteobacterium. *PLoS ONE* **6**: e25386.

Söding, J., Biegert, A., and Lupas, A.N. (2005) The HHpred interactive server for protein
homology detection and structure prediction. *Nucleic Acids Research* **33**: W244–W248.

Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**: 1312–1313.

Stookey, L.L. (1970) Ferrozine---a new spectrophotometric reagent for iron. *Anal Chem* **42**:
779–781.

Summers, Z.M., Gralnick, J.A., and Bond, D.R. (2013) Cultivation of an obligate Fe(II)-
oxidizing lithoautotrophic bacterium using electrodes. *MBio* **4**: e00420–12.

Ullrich, S.R., González, C., Poehlein, A., Tischler, J.S., Daniel, R., Schlömann, M., et al. (2016)
Gene loss and horizontal gene transfer contributed to the genome evolution of the extreme
acidophile "Ferrovum." *Front Microbiol* **7**: e78237.

Ullrich, S.R., Poehlein, A., Tischler, J.S., González, C., Ossandon, F.J., Daniel, R., et al. (2016)
Genome analysis of the biotechnologically relevant acidophilic iron oxidising strain JA12
indicates phylogenetic and metabolic diversity within the novel genus "Ferrovum." *PLoS*
ONE **11**: e0146832.

Webb, B. and Sali, A. (2016) Comparative protein structure modeling using MODELLER. *Curr*
Protoc Bioinformatics **54**: 5.6.1–5.6.37.

Wickham, H. (2009) ggplot2: Elegant Graphics for Data Analysis Springer, New York, NY.

Williams, K.H., Kemna, A., Wilkins, M.J., Druhan, J., Arntzen, E., N'Guessan, A.L., et al.
(2009) Geophysical monitoring of coupled microbial and geochemical processes during
stimulated subsurface bioremediation. *Environ Sci Technol* **43**: 6717–6723.

Williams, K.H., Long, P.E., Davis, J.A., Wilkins, M.J., N'Guessan, A.L., Steefel, C.I., et al.
(2011) Acetate availability and its influence on sustainable bioremediation of uranium-
contaminated groundwater. *Geomicrobiol J* **28**: 519–539.

Wu, Y.-W., Simmons, B.A., and Singer, S.W. (2016) MaxBin 2.0: an automated binning
algorithm to recover genomes from multiple metagenomic datasets. *Bioinformatics* **32**: 605–
607.

Yarzabal, A., Brasseur, G., Ratouchniak, J., Lund, K., Lemesle-Meunier, D., DeMoss, J., and
Bonnefoy, V. (2002) The high-molecular-weight cytochrome c Cyc2 of *Acidithiobacillus*
ferrooxidans is an outer membrane protein. *Journal of Bacteriology* **184**: 313–317.

Yoshikawa, S., Shinzawa-Itoh, K., Nakashima, R., Yaono, R., Yamashita, E., Inoue, N., et al.
(1998) Redox-coupled crystal structural changes in bovine heart cytochrome c oxidase.
Science **280**: 1723–1729.

Zhang, Y. (2008) I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics*
2016 17:1 **9**: 40.

Figure legends

Figure 1. Cyc2 protein phylogenetic tree, with all FeOB groups labeled, including the

functionally characterized *At. ferrooxidans* Cyc2 and *L. ferriphilum* Cyt$_{572}$, which have both

been shown to oxidize Fe(II) *in vitro*. Note that the neutrophilic FeOB (Zetaproteobacteria,

Gallionellaceae, *Chlorobi*) fall into a well-supported cluster (bootstrap value 93%). Also closely

related is the *Chlorobi* photoferrotroph Cyc2. Bootstrap values (%) corresponding to nodes

highlighted by circles: Zetaproteobacteria-most sequences (92), Gallionellaceae (98/75),

Chlorobi (100), Acidithiobacillus ferrooxidans (100), Ferrovum (100), Zetaproteobacteria-4

sequences (100), Leptospirillum spp. (100). See Supplemental File 1 for full tree with all taxa

and nodes labeled.

Figure 2. Alignment of cytochrome-containing section for Cyc2 from representative neutrophilic

and acidophilic FeOB. Red=CXXCH heme-binding motif. Blue=other motifs discussed in the

text. This section is preceded by a signal sequence, as detected by SignalP, and followed by a

beta strand predicted by PSIPRED. See Supplemental Figure 2 for full Cyc2 alignment.

Figure 3. Two views of the Cyc2 porin model, generated using iTasser. The Cyc2 cytochrome

(orange sphere=hypothesized location) is connected to the N terminal end of the porin (blue

strand).

Figure 4. Heterologous expression of Cyc2 and Fe oxidation assay results. (A) Western blot

showing expression of Cyc2 (lane 4) and Cyc2$_{porin}$ (lane 6); U=uninduced, I=induced with 0.5

mM IPTG. See corresponding Coomassie-stained gel in Supplemental Fig. 6B. (B) Heme stain

showing successful insertion of heme into Cyc2. (C) Fe oxidation assay results. Dissolved Fe^{2+}

monitored over time after Fe^{2+} addition to a suspension of *E. coli* cells expressing Cyc2,

Cyc2$_{porin}$, or neither (empty vector). Fe^{2+} was added to Cyc2-expressing cells a second time.

Error bars represent 1 SD of triplicate experimental runs.

Table 1. Distribution of putative Fe oxidation genes in FeOB genomes

Organism	*mtoAB*	*cyc2*	Reference
Freshwater neutrophilic microaerophilic FeOB (Gallionellaceae)			
Sideroxydans lithotrophicus ES-1	**X**	x	Liu *et al.*, 2012; Emerson *et al.*, 2013
Gallionella capsiferriformans ES-2*	x	x	Emerson *et al.*, 2013
Gallionellaceae *sp.* KS	x	x	He *et al.*, 2016
Ferriphaselus amnicola OYT1*		x	Kato *et al.*, 2015
Ferriphaselus globulitus R-1*		x	Kato *et al.*, 2015
Marine neutrophilic microaerophilic FeOB (Zetaproteobacteria)			
Mariprofundus ferrooxydans PV-1*, JV-1*, M34*, EKF-M39*		x	Singer *et al.*, 2011; Barco *et al.*, 2015; Field *et al.*, 2015
Mariprofundus aestuarium CP-5*		x	Chiu *et al.*, 2017
Mariprofundus ferrinatatus CP-8*		x	Chiu *et al.*, 2017
Mariprofundus micogutta ET2*		x	Makita *et al.*, 2016
Mariprofundus sp. DIS-1*		x	Mumford *et al.*, 2016
Ghiorsea bivora TAG-1, SV108		x	Mori *et al.*, 2017
Acidophilic FeOB			
Acidithiobacillus ferrooxidans ATCC23270		**X**	Yarzabal *et al.*, 2002; Castelle *et al.*, 2008
Acidithiobacillus ferrivorans CF27, SS3		x	Hallberg *et al.*, 2009; Kupka *et al.*, 2009
Leptospirillum sp., Iron Mountain Mine community		**X** (cyt_{572})	Jeans *et al.*, 2008
Leptospirillum ferriphilum DSM14647/P3A*		x	Cardenas *et al.*, 2014
Ferrovum myxofaciens P3G*		x	Johnson *et al.*, 2014
Ferrovum spp. JA12, PN-J185, Z-31		x	Ullrich, Poehlein, *et al.*, 2016; Ullrich, González, *et al.*, 2016
Thiomonas spp. FB-6, FB-Cd	x	x	Fabisch *et al.*, 2011
Phototrophic FeOB			
Chlorobium ferrooxidans KoFox		x	Heising *et al.*, 1999
Chlorobium phaeoferrooxidans KB01		x	Crowe *et al.*, 2017
Rhodopseudomonas palustris TIE-1	**X** (*pioAB*)		Jiao and Newman, 2007

x=present in genome

X=Fe oxidation function confirmed

*obligate FeOB, noted for pure cultures only

Table 2. *cyc2* and *mtoA* expression in Jewell et al. Rifle aquifer metatranscriptome

Gallionellaceae bin	*cyc2* max (RPKM)	*mtoA* max (RPKM)	*cyc2*:*mtoA* ratio
b22.1	136	0.714	190
b22.2	271	0.333	812
b22.3	173		
b22.5	1211		
b22.6	696	42.1	16.5
b22.7		5.09	
b22.8	176	0.166	1520
b22.9	171	0.317	540
all bins, median (n=11)	172	0.73	236

Figure 1. Cyc2 protein phylogenetic tree, with all FeOB groups labeled, including the functionally characterized *At. ferrooxidans* Cyc2 and *L. ferriphilum* Cyt₅₇₂, which have both been shown to oxidize Fe(II) *in vitro*. Note that the neutrophilic FeOB (Zetaproteobacteria, Gallionellaceae, *Chlorobi*) fall into a well-supported cluster (bootstrap value 93%). Also closely related is the *Chlorobi* photoferrotroph Cyc2. Bootstrap values (%) corresponding to nodes highlighted by circles: Zetaproteobacteria-most sequences (92), Gallionellaceae (98/75), Chlorobi (100), Acidithiobacillus ferrooxidans (100), Ferrovum (100), Zetaproteobacteria-4 sequences (100), Leptospirillum spp. (100). See Supplemental File 1 for full tree with all taxa and nodes labeled.

Figure 2. Alignment of cytochrome-containing section for Cyc2 from representative neutrophilic and acidophilic FeOB. Red=CXXCH heme-binding motif. Blue=other motifs discussed in the text. This section is preceded by a signal sequence, as detected by SignalP, and followed by a beta strand predicted by PSIPRED. See Supplemental Figure 2 for full Cyc2 alignment.

Figure 3. Two views of the Cyc2 porin model, generated using iTasser. The Cyc2 cytochrome (orange sphere=hypothesized location) is connected to the N terminal end of the porin (blue strand).

Figure 4. Heterologous expression of Cyc2 and Fe oxidation assay results. (A) Western blot showing expression of Cyc2 (lane 4) and Cyc2$_{porin}$ (lane 6); U=uninduced, I=induced with 0.5 mM IPTG. See corresponding Coomassie-stained gel in Supplemental Fig. 6B. (B) Heme stain showing successful insertion of heme into Cyc2. (C) Fe oxidation assay results. Dissolved Fe^{2+} monitored over time after Fe^{2+} addition to a suspension of *E. coli* cells expressing Cyc2, Cyc2$_{porin}$, or neither (empty vector). Fe^{2+} was added to Cyc2-expressing cells a second time. Error bars represent 1 SD of triplicate experimental runs.