Social and non-social autism symptom and trait domains are genetically dissociable

Varun Warrier[1,15], Roberto Toro[2,3], Hyejung Won[4], Claire S Leblond[2,3], Freddy Cliquet[2], Richard Delorme[2,5], Ward De Witte[6], Janita Bralten[6,7], Bhismadev Chakrabarti[1, 8], Anders D Børglum[9-11], Jakob Grove[9-12], Geert Poelmans[6], the 23andMe Research Team†, David A. Hinds[13,16] Thomas Bourgeron[2,3,14,16], and Simon Baron-Cohen[1,15,16]

[1] Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridgeshire, United Kingdom

[2] Institut Pasteur, Human Genetics and Cognitive Functions Unit, Paris, France

[3] CNRS UMR 3571: Genes, Synapses and Cognition, Institut Pasteur, Paris, France

[4] Department of Genetics, University of North Carolina, Chapel Hill, NC 27599, USA

[5] Child and Adolescent Psychiatry Department, Robert Debré Hospital, Paris, France

[6] Department of Human Genetics, Radboud University Medical Center, Nijmegen, The Netherlands

[7] Donders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

[8] Centre for Integrative Neuroscience and Neurodynamics, School of Psychology and Clinical Language Sciences, University of Reading, Reading, United Kingdom

[9] The Lundbeck Foundation Initiative for Integrative Psychiatric Research, iPSYCH, Aarhus, Denmark

[10] Centre for Integrative Sequencing, iSEQ, Aarhus University, Aarhus, Denmark

[11] Department of Biomedicine - Human Genetics, Aarhus University, Aarhus, Denmark

[12] Bioinformatics Research Centre, Aarhus University, Aarhus, Denmark

[13] 23andMe Inc., Mountain View, California 94043, USA

[14] Université Paris Diderot, Sorbonne Paris Cité, Human Genetics and Cognitive Functions, Paris, France

[15] **Corresponding authors**: Varun Warrier (vw260@medschl.cam.ac.uk) and Simon Baron-Cohen (sb205@cam.ac.uk). Autism Research Centre, Department of Psychiatry, Cambridge University, Douglas House, 18B Trumpington Road, Cambridge, CB2 8AH, United Kingdom. Telephone: 0044 (0) 1223 746057.

[16] Joint senior authors.

† A full list of consortium members appears at the end of the paper.

Abstract

The core diagnostic criteria for autism comprise two symptom domains – social and communication difficulties, and unusually repetitive and restricted behaviour, interests and activities. There is some evidence to suggest that these two domains are dissociable, yet, this hypothesis has not been tested using molecular genetics. We test this using a GWAS of a non-social autistic trait, systemizing (N = 51,564), defined as the drive to analyse and build systems. We demonstrate that systemizing is heritable and genetically correlated with autism. In contrast, we do not identify significant genetic correlations between social autistic traits and systemizing. Supporting this, polygenic scores for systemizing are significantly positively associated with restricted and repetitive behaviour but not with social difficulties in autistic individuals. These findings strongly suggest that the two core domains of autism are genetically dissociable, and point at how to fractionate the genetics of autism.

Introduction

The core diagnostic criteria of autism comprises two symptom domains: difficulties in social interactions and communication (the social domain) and unusually repetitive and restricted behaviour and stereotyped interests (the non-social domain)[1]. Multiple lines of evidence suggest that these two domains are dissociable[2,3]. First, factor and principal component analysis of autism and autistic traits have mostly identified two factors for autism – a social and a non-social factor[4–9]. Second, investigation of autistic traits in large cohorts have demonstrated a positive phenotypic correlation between different social traits and different non-social traits separately, but only a limited correlation between social and non-social traits[9–12]. Third, twin genetic correlations between social and non-social symptom domains in autism are low, though both social and non-social trait domains are highly heritable in neurotypical[13,14] or autistic twins[15]. Fourth, difficulties in social and non-social domains can occur independently of each other[16,17], which has been used to subgroup individuals on the spectrum based on the two domains[18]. This suggests that the genetic and phenotypic architecture of autism consists of at least two categories of broadly dissociable domains. This has implications for genetic, biological, and clinical studies of autism, since most studies have investigated autism as if it is a unitary condition[3]. The idea that social and non-social symptom domains are dissociable is unsurprising given their very different nature, and very different underlying cognitive processes one related to interpreting animate motion and mental states (theory of mind) and the other related to recognizing inanimate objects, events or patterns (systemizing), or their very different underlying neurology[3]. Nevertheless, the traditional view of autism is that it is a syndrome, meaning the diagnosis is only given when the social and non-social symptom domains cluster together.

However, to date, there has been limited molecular genetic evidence in support of this dissociability hypothesis, partly due to the limited large-scale research on the genetics of social and non-social domains. Most genetic research into the social and non-social domains has been primarily through linkage and genome-wide association studies (GWAS) in relatively small samples of autistic individuals and the general population ($N < 5K$)[19–25]. This has precluded a detailed molecular genetic investigation of the social and non-social domains associated with autism. Given currently available sample sizes with phenotypic information, investigating the genetics of the social and non-social domains in autistic individuals is difficult. However, several studies have demonstrated that the underlying liability for autism is normally distributed in the general population[26–29]. Factor analysis have failed to identify

discontinuities between clinical autism and autistic traits in the general population[30]. Autistic

traits are heritable[31–33], are elevated in family members of autistic individuals compared to the

general population[34,35], and are transmitted intergenerationally[36,37]. Factor analysis of autistic

traits measures have also identified two different factors in both the general population and

autistic individuals – one linked to the social domain, and another linked to the non-social

domain, mirroring the factor structure of clinical autism domains[6,9,30,38]. Studies have further

demonstrated moderate to high shared genetics between the extremes of the liability

distribution and the rest of the distribution[14,39–41]. One twin study investigated the bivariate

genetic correlation between research and clinical autism diagnosis and autistic traits and

identified high genetic correlations $(0.7 < r_g < 0.89)$[42]. Validating this, studies have identified

modest shared genetics between autism and autistic traits[43–45]. Taken together, there is

considerable evidence to suggest that autism represents the extreme end of the autistic traits

continuum.

While a few studies have investigated the genetics of traits contributing to the social

domains such as social and communications difficulties[19,44,45], empathy[46], and emotion

recognition[47], there have been limited studies investigating the genetics of the non-social

domain[25,48]. Neither of these studies have replicably identified significant variants associated

with the non-social domain, primarily because of the relatively modest sample sizes of the

GWAS. An alternate approach is to investigate the genetics of non-social autistic traits in the

typical population, maximizing the sample size. To better understand the genetics of a non-

social autistic trait, we investigate the genetics of systemizing measured using a 75-item well

validated self-report measure called the Systemizing Quotient-Revised (SQ-R) (**Methods**).

Systemizing involves identifying *input-operation-output* relationships in order to analyse and

build systems, and to understand the laws that govern specific systems[49]. The hyper-

systemizing theory of autism proposes that autistic individuals, on average, have superior

attention to detail, and a stronger drive to systemize[49]. This has been validated in several

studies[50,51] including a recent study in more 650,000 individuals including 36,000 autistic

individuals[12]. Several lines of evidence suggest that autistic individuals have intact or

superior systemizing. The idea was noted in the earliest papers describing autism by both

Hans Asperger[52] and Leo Kanner[53]. Further, autistic adults, on average, score higher on the

SQ-R compared to individuals in the general population[10,51], a pattern also observed in

autistic children[54]. Several items in the SQ-R specifically measures circumscribed interests

and insistence on sameness, two of the items mentioned in the DSM-5, and several of these

items map onto items on the Autism Spectrum Quotient (AQ), a well validated measure of

autistic traits[27] (**Supplementary Note**). Because systems follow rules, they repeat, such that

an operation on a given input produces the same output every time. A fascination with

systems may thus manifest as unusually repetitive behaviour. And because systems depend

on precise variables, a fascination with systems may also manifest as unusually narrow

interests in autism.

The present study has two aims: 1. To investigate the polygenic architecture of a non-

social trait linked to autism: *systemizing*, and 2. To investigate if social and non-social autistic

traits measured in the general population are genetically dissociable.

Results

We first conducted a GWAS of systemizing (N = 51,564) measured using the SQ-R.

Following this, and using data from GWAS of social traits genetically correlated with autism

(GWAS of self-reported empathy (N = 46,861)[46], and GWAS of social relationship

satisfaction[55] measured using friendship ($N_{effective}$ = 164,112) and family relationship ($N_{effective}$

=158,116) satisfaction scales) we investigated if the social and non-social domains autistic

traits are genetically dissociable in the general population. A flow-chart of the study design is

provided in **Figure 1**.

Insert Figure 1 here

Systemizing was measured in the 23andMe sample (N = 51,564) using scores from

the SQ-R[10]. Scores on SQ-R were normally distributed, with a mean of 71±21 out of 150. As

hypothesized based on previous research[10,12,51], males (76.5±20), on average, scored higher

than females (65.4±20.6) (P < 0.001, Cohen's d = 0.54, **Supplementary Figure 1**). Given the

significant sex differences in scores, we conducted a non-stratified and sex-stratified GWAS

for SQ-R. Genome-wide association analyses identified three significant loci (**Figure 2,**

Supplementary Table 1 and **Supplementary Figure 2**). Two of these were significant in the

non-stratified GWAS: rs4146336 on chromosome 3 (P = 2.58×10^{-8}) and rs1559586 on

chromosome 18 (P = 4.78×10^{-8}). The third significant locus was in the males-only GWAS

(rs8005092 on chromosome 14, P = 3.74×10^{-8}). rs8005092 and rs1559586 lie in regions of

high genetic recombination. Linkage-Disequilibrium Score Regression (LDSR) intercept

suggested that there was minimal inflation due to population stratification (**Figure 2**). Fine-

mapping of the three regions identified 14 credible SNPs (**Methods**). None of the SNPs overlapped with foetal brain eQTL. However, two of these SNPs mapped onto two genes - *LSAMP* and *PTMAP8*, both of chromosome 3 - using chromatin interaction data in the foetal brain. Of these, LSAMP is a neuronal adhesion molecule in the limbic system of the developing brain Additionally, gene-based analysis identified 4 significant genes *SDCCAG8*, *ZSWIM6*, *ZNF574* and *FUT8* (**Supplementary Table 2**). Of these, mutations in *ZSWIM6* cause a neurodevelopmental disorder with, in some cases, co-morbid autism and unusually repetitive movements and behaviour[56]. As supporting analyses, we investigated the direction of effect for all independent SNPs with $P < 1x10^{-6}$ in the non-stratified SQ-R GWAS in GWAS of autism[57], educational attainment[58], and cognitive aptitude[59]. Five out of six SNPs tested had concordant effect in the GWAS for educational attainment and GWAS for cognitive aptitude ($P = 0.21$, two-sided binomial sign test for each comparison). Similarly, four out of five SNPs tested had concordant effect direction in the GWAS for autism (**Supplementary Table 3a**) ($P = 0.37$, two-sided binomial sign test). For these three phenotypes, we additionally assessed effect direction concordance using binomial sign test at less stringent P-value thresholds in the SQ-R GWAS, after LD-based clumping ($P < 1, 0.5,$ 0.1 and $1x10^{-4}$). Binomial sign test was statistically significant at three of the four P-value thresholds ($P = 1, 0.5$ and 0.1) for all three phenotypes but not statistically significant at $P = 1E-4$, presumably due to the low statistical power (**Supplementary Table 3b**). Additionally, we tested effect direction concordance ($P < 1x10^{-6}$) in a GWAS ($N = 1,981$) of 'insistence on sameness' , a phenotype that's similar to systemizing (**Methods**). Four out of five SNPs had a concordant effect direction including the two SNPs with $P < 5x10^{-8}$ in the non-stratified SQ-R GWAS ($P = 0.37$, two-sided binomial sign test).

Insert Figure 2 here

Additive SNP-based heritability (h^2_{SNP}) calculated using LDSR was 0.12 ± 0.012 for the SQ-R ($P = 1.2x10^{-20}$). Despite small but significant sex-differences in the SQ-R scores, there was no significant difference in h^2_{SNP} between males and females ($P = 0.34$) (**Supplementary Figure 3 and Supplementary Table 4**), which was strengthened by the high genetic correlation between males and females (1 ± 0.17; $P = 3.91x10^{-10}$), suggesting a similar polygenic architecture between sexes. The per-SNP effect for the most significant SNPs was small, suggesting a highly polygenic architecture ($R^2 = 0.001 – 0.0002\%$, after correcting for winner's curse, **Supplementary Table 5**).

Partitioned heritability for functional categories identified significant enrichment for evolutionary conserved regions, transcription start sites, foetal DNase hyper-sensitivity sites, and H3 lysine 27 acetylation (H3K27ac), suggesting a prominent role for regulatory and conserved genomic regions in systemizing **(Supplementary Table 6)**. Partitioning heritability based on tissue specific active chromatin marks identified a significant enrichment for brain specific chromatin signatures highlighting the role of the brain for the SQ-R. Notably, this enrichment was significant in both adult and foetal brain specific active chromatin marks **(Supplementary Table 7 and Supplementary Figure 4)**. Enrichment for genes expressed in the brain was high but failed to reach statistical significance after correcting for the multiple tests conducted (**Supplementary Figure 5 and Supplementary Table 8**).

We identified a significant positive genetic correlation between the SQ-R and autism as well as measures of intelligence (cognitive aptitude and educational attainment) **(Supplementary Table 9 and Figure 3a)**. Of all the psychiatric conditions tested (**Methods**), SQ-R was only significantly genetically correlated with autism ($r_g = 0.26 \pm 0.06$; P $= 3.35 \times 10^{-5}$), demonstrating the relative specificity of the SQ-R to autism. Notably, the effect size of the genetic correlation between autism and the SQ-R is similar to the genetic correlation between autism and self-reported empathy (measured using the Empathy Quotient (EQ[60]): $r_g = -0.27 \pm 0.07$) and scores on the Social and Communication Disorders Checklist (SCDC[61]): $r_g = 0.27 \pm 0.13$). Controlling for the genetic effects of educational attainment on the SQ-R GWAS using genome-wide inferred statistics (GWIS) (**Methods**) attenuated the genetic correlation with autism only modestly, suggesting that the SQ-R scores are genetically correlated with autism independently of the genetic effects of education **(Figure 3b and Supplementary Table 10)**. We validated this using genomic structural equation modeling (GSEM) (**Methods**) using both educational attainment and cognitive aptitude **(Figure 3c)**. Further, the SQ-R was not genetically correlated with any of the social measures related to autism – friendship and family relationship satisfaction, scores on a self-report measure of empathy - the Empathy Quotient (EQ), and the scores on the Social and Communication Disorders Checklist (SCDC), which is a measure of social and communication difficulties (see **Supplementary Note** for how these traits map onto social domains in autism). Estimates of genetic correlations between SQ-R scores and the various social traits are also small, suggesting that there is limited shared genetics between social autism traits and the SQ-R.

To understand the genetic relationship between the SQ-R and autism in a broader

context, we evaluated the genetic correlations between multiple phenotypes with evidence of

significant genetic correlation with autism (15 phenotypes in total, see **Methods** for a list of

phenotypes included). Clustering highlighted three broad clusters: a social cluster, a

psychiatric cluster, and an intelligence cluster **(Figure 4a and Supplementary Tables 11**

and 12). The SQ-R clusters closely with measures of intelligence, but while educational

attainment and cognitive aptitude are significantly correlated with multiple social traits and

psychiatric conditions, the SQ-R is only genetically correlated with autism.

Given that the two major domains of autism as identified by the DSM-5 are persistent

difficulties in social interaction and communication and unusually restrictive, stereotyped,

and repetitive interests[1], we hypothesized that the combination of significant negative genetic

correlation with social traits (friendship satisfaction and empathy) and significant positive

genetic correlation with SQ-R is uniquely associated with autism (**Methods**). Indeed, across

the nine psychiatric conditions for which we had summary GWAS statistics, this combination

was uniquely seen for autism **(Figure 4b, Supplementary Table 13)**.

Given that our current analysis focussed on the general population, we sought to

investigate if polygenic scores from the SQ-R were associated with social and non-social

autism domains in 2,221 autistic individuals from the Simons Simplex Collection (**Methods**).

We hypothesized that SQ-R may be significantly associated with the non-social domain in

autism, but not associated with the social domain in autism. Polygenic scores for SQ-R were

significantly associated with scores on the Repetitive Behaviour Scale-Revised (Beta =

0.052 ± 0.02, P = 0.013), but not on the social and communication subscale of ADOS-G (Beta

= -0.00099 ± 0.018, P = 0.95) after adjusting for multiple test (Bonferroni alpha = 0.025). We

validated this in 426 additional individuals of which 401 had a diagnosis of autism with RBS-

R scores from the EU-AIMS LEAP, AGRE, and Paris cohorts. Here, we identified a

concordant effect direction for polygenic score of the SQ-R (Beta = 0.02 ± 0.05, P = 0.65),

though the results were not significant potentially due to the small sample size. Inverse-

variance meta-analysis of the discovery and the validation cohorts marginally improved the

significance of the association (Beta = 0.047 ± 0.018, P = 0.010), and the results remained

statistically significant (Bonferroni alpha = 0.025). In a separate sample of 475 autistic

individuals from the AGRE cohort, polygenic scores for the SQ-R were not associated with the social and communication subscale of ADOS-G (Beta = -0.046±0.04, P = 0.24). Meta-analysis of the two cohorts did not produce a statistically significant result (Beta = -0.008±0.016, P = 0.60) (see **Power calculations in the Supplementary Note)**. We note that the lack of association between the polygenic scores for the SQ-R and the ADOS-G social and communication subscale is not indicative of absence of shared genetics, but rather indicative of lower shared genetics than between the RBS-R and the SQ-R.

Finally, to further validate the results in autistic individuals, we conducted bivariate genetic correlations on scores on the RBS-R and the ADOS-G social and communication subscale in 2,989 individuals from the SSC, AGRE, EU-AIMS LEAP and Paris cohorts (2964 autistic individuals). Both the RBS-R (h^2_{SNP} = 0.11±0.11, P = 0.15) and the ADOS-G social and communication subscale (h^2_{SNP} = 0.26±0.10, P = 0.004) had modest h^2_{SNP}, though only the latter was statistically significant. We identified a small genetic correlation (r_g = 0.15±0.46, P = 0.74), which was not statistically different from 0. Given the small sample size, the genetic correlation is unlikely to be statistically significant. However, the effect was small and statistically less than 1 (P = 0.034, One-tailed T test).

Discussion

We present the largest GWAS of a non-social trait related to autism in the general population – systemizing, measured using the SQ-R. We demonstrate that systemizing is heritable and genetically correlated with autism. Associated loci are enriched in genomic regions containing brain chromatin signatures and we identify three genome-wide significant loci, but these must be replicated in an independent cohort. Despite the modest sample size, our GWAS is well-powered to investigate genetic correlations between various phenotypes including social traits related to autism, as the Z-score of the h^2_{SNP} is above the recommended threshold of four[62]. We identify high sign concordance of the top SNPs in genetically correlated traits, enrichment for active chromatin marks in foetal and adult brain, and significant polygenic score association with the RBS-R. Polygenic score analysis suggests that the shared genetics between systemizing and the non-social domain in autism is considerably higher than the shared genetics between systemizing and the social domain. In addition, using a smaller sample of autistic individuals, we provide preliminary evidence that the social and non-social domains in autistic individuals have low shared genetics. Our results

highlight the need to collect deeper clinical and cognitive information in autistic individuals
to better understand the phenotypic heterogeneity in autism.

Most studies model autism, and the underlying liability measured as autistic traits, as
a single domain. This has likely arisen because of the difficulties in recruiting and
phenotyping sufficient numbers of autistic people. Our study suggests that, both in the
general population and in autistic individuals, social and non-social autistic traits and
symptom domains are genetically dissociable. This may to some extent explain why,
compared to GWAS of other psychiatric conditions of roughly similar sample sizes[57,63–65],
GWAS of autism to date have identified fewer loci. One possible explanation is statistical
signal-attenuation because of the underlying heterogeneity. However, this does not
necessarily suggest that systemizing, or the other individual trait domains are less complex.
For instance, we observe similar h^2_{SNP} for SQ-R, self-reported empathy [46], and the largest and
most recent GWAS of autism[57]

It is important to investigate if these domains are dissociable in a larger cohort of
autistic individuals and identify potential convergence of the two domains in gene expression
networks in the developing brain. Our results confirm the need to rethink our understanding
of autism as existing along a single dimension[3,66]. We hypothesize that the dissociation of the
two domains will extend to these other research modalities in studies of autism and autistic
traits. It is important to note that, while our results demonstrate two broadly dissociable
autistic trait domains in the general population and in autistic individuals, more research is
needed to identify other potentially dissociable domains and to investigate if this
dissociability is driven by different designs of phenotypic instruments. For example, our
research does not make a distinction between communication vs. social interaction abilities,
or between sensory difficulties vs. repetitive behaviours, and future molecular genetic studies
may identify varying levels of overlap between these domains. The same principle applies to
other research modalities (neuroimaging, cognitive studies, hormonal assays, etc.,)
investigating the biology of autism and autistic traits. These different symptom domains of
autism may contribute to different co-morbidities. Our results identify shared genetics
between the social autistic traits and psychiatric conditions such as schizophrenia and
depression, but limited shared genetics between the SQ-R and these conditions. This needs to
be   evaluated   epidemiologically.

Methods

Participants: The current study included participants from 23andMe (Primary GWAS - SQ-R), from ALSPAC (GWAS of scores on the Social and Communication Disorders Checklist (SCDC)) and autistic individuals from the Simons Simplex Collection (SSC), the Autism Genetic Resource Exchange (AGRE), and the EU-AIMS LEAP and PARIS cohorts.

23andMe: Research participants in the GWAS of the SQ-R were from 23andMe and are described in detail elsewhere[67,68]. All participants provided informed consent and answered surveys online according to a human subjects' research protocol, which was reviewed and approved by Ethical & Independent Review Services, an external AAHRPP-accredited private institutional review board (http://www.eandireview.com). All participants completed the online version of the SQ-R on the 23andMe participant portal. Only participants who were primarily of European ancestry (97% European Ancestry) were selected for the analysis using existing methods[69]. Unrelated individuals were selected using a segmental identity-by-descent algorithm[70]. A total of 51,564 participants completed the SQ-R (males = 26,063, and females = 25,501).

ALSPAC: ALSPAC is a longitudinal cohort which recruited pregnant mothers in the Avon region of the UK. The ALSPAC cohort comprises 14,541 initial pregnancies from women in Avon resulting in a total of 13,988 children who were alive at 1 year of age. Children were enrolled in additional phases, described in greater detail elsewhere[71]. This study received ethical approval from the ALSPAC Law-and-Ethics Committee, and the Cambridge Human Biology Research Ethics Committee. Written informed consent was obtained from parent or a responsible legal guardian for the child to participate. Assent was obtained from the child participants where possible. We conducted a GWAS of scores on the SCDC in 5,421individuals from ALSPAC.

Other cohorts: We included data from four cohorts to conduct polygenic score and bivariate genetic correlation analysis. The SSC (n = 2,221 unrelated autistic individuals) consists of simplex autistic families, and are described elsewhere[72]. The AGRE cohort (n = 482 unrelated autistic individuals) consists of multiplex autism families, details of which are provided elsewhere[73]. Across all cohorts, all participants were of European ancestry as identified using multi-dimensional scaling. Additionally, we included 401 individuals (including 25 neurotypical individuals) from the EU-AIMS LEAP[74] and Paris[75] cohorts. Across all cohorts, we included only unrelated individuals, who were predominantly of

European Ancestry as defined by genetic principal components (5 SD deviations above or
below the mean European PC1).

Additionally, we also included data from 1,981 unrelated individuals (1000 males,
1981 females) from the Nijmegen Biomedical Study (NBS) to provide support for the
independent SNPs with P < 1x10-6 in the non-stratified GWAS. Participants were asked the
question: "It upsets me if my daily routine is disturbed", which is related to a non-social
domain of autism, and is similar to an item in the Autism Spectrum Quotient. Further
information including genotyping and quality control is provided elsewhere[43]. Genetic
association for the top SNPs were conducted using age, sex, and the first five genetic
principal components as covariates using linear regression.

Phenotypes: The primary phenotype for this study is the SQ-R, which was used to conduct a
GWAS in participants from 23andMe. The SQ-R is self-report measure of systemizing drive,
or interest in rule-based patterns[10]. The SQ-R taps a variety of domains of systemizing, such
as interest in mechanical (e.g., car engines), abstract (e.g., mathematics), natural (e.g., the
weather), motor (e.g., knitting), and collectible (e.g., stamp collecting) systems. There are 75
items on the SQ-R, with a maximum score of 150 and a minimum score of 0. Scores on the
test are normally distributed[10]. The SQ-R has good cross-cultural stability and good
psychometric properties with Cronbach's alpha ranging from 0.79 to 0.94 in different
studies[76]. Test-retest reliability available in a Dutch sample indicated high reliability of 0.79
(Pearson correlation)[76]. This was supported by another study in 4,058 individuals which
identified high internal cohesion[77]. Exploratory followed by confirmatory factor analysis
using Rasch modelling suggests that the SQ-R is unidimensional[77]. A sex difference has been
observed in multiple studies with males, on average, scoring significantly higher than
females[10,51]. Criterion validity shows that the SQ-R has a modest but significant correlation
with the Mental Rotation Test (r =.25, P =.013), as well as its subscales[78]. Autistic
individuals, on average, score higher on the SQ-R in multiple different studies[10,51,79]. Further,
the SQ-R also predicts autistic traits, with a combination of the SQ-R and the Empathy
Quotient predicting as much as 75% of the variance on the Autism Spectrum Quotient, a
measure of autistic traits[10]. The SQ-R has been validated using a short form in a very large
population of 600,000 controls and 36,000 autistic individuals (Greenberg et al, 2018).

In addition, we used the following secondary phenotypes: SCDC in ALSPAC,
ADOS-G social and communication scores and the RBS-R in the other cohorts. We also used
a single question which is a measure of 'insistence on sameness' in the NBS cohort.

The SCDC is a questionnaire that measures difficulties in verbal and nonverbal
communication, and social interaction including reciprocal social interaction[61]. The
questionnaire consists of 12 questions, with scores ranging from 0 – 24, with higher scores
reflecting difficulties in social interaction and communication. The SCDC has good internal
consistency (0.93) and good test-retest reliability (0.81)[61]. The SCDC has reasonable
specificity and sensitivity in distinguishing clinically diagnosed autism from control
individuals[80]. Previous research has demonstrated that the SCDC is genetically correlated
with autism[44,45,57]. We conducted a GWAS of SCDC to investigate if it is genetically
correlated with SQ-R in this study. We used mother-reported SCDC scores on children aged
8. While SCDC has been measured at different ages in the ALSPAC cohort, we chose SCDC
scores at age 8 as these had the largest sample size and have high h^2_{SNP} [19] ($h^2 = 0.24 \pm 0.07$).

We chose two measures of social and non-social traits. For the social trait, we used
the social and communication domain scores from the ADOS-G, a widely used instrument for
diagnosing and assessing autism in four cohorts (SSC, AGRE, EU-AIMS LEAP and Paris).
Participants completed one of the following ADOS-G modules[81]: 1 (used for children with
little or no phrase speech), 2 (for children with non-fluent speech), 3 (verbally fluent
children), and 4 (verbally fluent adolescents and adults). For this study, we used the raw
totals of the scores from the social domain and the communication domain, combined. Scores
for all 4 modules range from 0 – 24. The ADOS-G has high overall internal consistency, and
high test-retest reliability for the social and communication subscales[81]. The choice for
combining the social and communication domain scores were informed by factor analysis
which suggested that the two domains contribute to one underlying factor[82].

In contrast to the Social and Communication domain, the restricted and repetitive
behaviour domain of the ADOS-G has poor test retest reliability ($r < 0.6$) and a smaller range
of scores (0 – 8) as it captures fewer repetitive and restrictive behaviour[81]. Hence, for this
study, we used sores on the RBS-R[83]. The RBS-R is a measure developed to specifically
measure restricted and repetitive behaviours in autistic individuals and captures stereotyped,
self-injurious, sameness, compulsive, ritualistic, and restricted behaviour[84], and has high

inter-rater reliability and internal consistency[84]. The RBS-R comprises 43 questions with

scores ranging from 0 – 3 for each item based on a Likert scale.

'Insistence on sameness' in the NBS cohort was measured using a single item: "It

upsets me if my daily routine is disturbed". This is related to a non-social domain of autism,

and is again similar to an item in the Autism Spectrum Quotient. Participants were asked to

indicate on a 4-point Likert scale "definitely agree", "slightly agree", "slightly disagree",

"definitely disagree".

Genotyping, imputation, and quality control and genetic association in the 23andMe

cohort: Details of genotyping, imputation and quality control in the 23andMe cohort are

provided elsewhere[47]. Briefly, unrelated participants were included if they had a call rate of

greater than 98.5%, and were of primarily European ancestry (97% European ancestry). A

total of 1,030,430 SNPs (including InDels) were genotyped. SNPs were excluded if: they

failed the Hardy-Weinberg Equilibrium Test at $P < 10^{-20}$; had a genotype rate of less than

90%; they failed the parent-offspring transmission test using trio data in the larger 23andMe

research participant database; or if allele frequencies were significantly different from the

European 1000 Genomes reference data (chi-square test, $P < 10^{-20}$). Phasing was conducted

using Beagle (version 3.3.1)[85] in batches of 8000-9000 individuals. This was followed by

imputation against all-ethnicity 1000 Genomes haplotypes (excluding monomorphic and

singleton sites) using Minimac2[86]. Genetic association analyses were restricted to SNPs with

a minor allele frequency > 1%. After quality control, 9,955,952 SNPs (imputed and

genotyped) were included in the GWAS.

Our primary analysis was an additive model of genetic effects and was conducted

using a linear regression with age, sex, and the first five ancestry principal components

included as covariates. In addition, given the modest sex difference, we also conducted sex-

stratified analyses. SNPs were considered significant at a genome-wide threshold of P

$<5x10^{8}$. Leading SNPs were identified after LD-pruning using Plink ($r^2 > 0.8$). Winner's

curse correction was conducted using an FDR based shrinking[87].

We calculate variance explained by first standardizing the regression estimates and

then squaring the estimates. This is equivalent to:

$$R^2 = \hat{B}_j^2 \frac{2(MAF_j)(1 - MAF_j)}{\sigma^2{}_y}$$

Where R^2 is the proportion of variance explained for SNP j. $\hat{B}_j^2$ is the non-

standardized regression coefficient, MAF is the minor allele frequency for SNP j, and $\sigma^2{}_y$ is

the variance of SQ. Further details of this formula are provided in the **Supplementary Note.**

Genotyping, imputation, and quality control and genetic association in the ALSPAC:

The SCDC [61] scores were calculated from children of the 90s (ALSPAC cohort)[71], in children

aged 8. In total, SCDC scores were available on N = 7,825 children. From this, we removed

individuals for whom complete SCDC scores were not available. After excluding related

individuals and individuals with no genetic data, data was available on a total of N = 5,421

unrelated individuals.

Participants were genotyped using the Illumina® HumanHap550 quad chip by

Sample Logistics and Genotyping Facilities at the Wellcome Sanger Institute and LabCorp

(Laboratory Corportation of America) using support from 23andMe. Individuals were

excluded based on gender mismatches, high missingness (> 3%), and disproportionate

heterozygosity. We restricted subsequent analyses to individuals of European descent (CEU),

which were identified by multidimensional scaling analysis and compared with Hapmap II

(release 22). Individuals were also removed if cryptic relatedness, assessed using identity by

descent, was greater than 0.1. Genotyped SNPs were filtered out if they had more than 5%

missingness, violated Hardy-Weinberg equilibrium (P < 1×10^{-6}), and had a minor-allele

frequency less than 1%, resulting in a total of 526,688 genotyped SNPs. Haplotypes were

estimated using data from mothers and children using ShapeIT (v2.r644)[88]. Imputation was

performed using Impute2 V2.2.2[89] against the 1000 genomes reference panel (Phase 1,

Version 3). Imputed SNPs were excluded from all further analyses if they had a minor allele

frequency < 1% and info < 0.8. After quality control, there were 8,282,911 genotyped and

imputed SNPs that were included in subsequent analyses.

Dosage data from BGEN files were converted using hard-calls, with calls with

uncertainty > 0.1 treated as missing data. Post-imputation, we excluded SNPs that deviated

from Hardy-Weinberg Equilibrium (P < 1×10^{-6}), with minor allele frequency < 0.01 and

missing call rates > 2%. We further excluded individuals with genotype missing rates > 5%.

The SCDC score was not normally distributed so we log-transformed the scores and ran

regression analyses using the first two ancestry principal components and sex as the

covariates using Plink 2.0[90].

The log-transformed SCDC scores (henceforth, SCDC scores) had a modest but

significant h^2_{SNP} as quantified using LDSR (h^2_{SNP} = 0.12 ±0.05). LDSR intercept (0.99)

suggested that there was no inflation in GWAS estimates due to population stratification. The

λ_{GC} was 1.013. We replicated the previously identified genetic correlation with autism[57]

(constrained intercept) using our SCDC GWAS (r_g = 0.45±0.18, P = 0.01). In addition, we

also identified a negative genetic correlation between educational attainment[58] and SCDC (r_g

= -0.30± 0.11, P = 0.007).

Genomic inflation factor, heritability, and functional enrichment for the SQ-R GWAS:

LDSR[91,92] was used to calculate for inflation in test statistics due to unaccounted population

stratification. Heritability was calculated using LDSR using the north-west European LD

scores. Difference in heritability between males and females was quantified using:

$$Z = \frac{h^2_{males} - h^2_{females}}{\sqrt{SE^2_{males} + SE^2_{females}}}$$

where Z is the Z score for the difference in heritability for a trait, (h^2_{males} - $h^2_{females}$) is the

difference h^2_{SNP} estimate in males and females, and SE is the standard errors for heritability.

Two-tailed P-values were calculated and reported as significant if P < 0.05.

For the primary GWAS (non-stratified analyses), we conducted functional annotation

using FUMA[93]. We restricted our analyses to the non-stratified analyses due to the high

genetic correlation between the sexes and the low statistical power of the sex-stratified

GWAS. We conducted gene-based association analyses using MAGMA[94] within FUMA and

report significant genes after using a stringent Bonferroni corrected P < 0.05. In addition, we

conducted enrichment for tissue specific expression and pathway analyses within FUMA. For

the significant SNPs, we investigated enrichment for eQTLs using brain tissues in the

BRAINEAC and GTEx[95] database within FUMA. We further conducted partitioned

heritability for tissue specific active chromatin marks and baseline functional categories using

extended methods in LDSR[96].

Hi-C based annotations of fine mapped loci: We fine mapped three genome-wide

significant loci (index SNPs: rs4146336 and rs1559586 or SQ; rs8005092 for SQ-R males) to

obtain credible SNPs. First, we selected SNPs with P<0.01 that are located in the linkage

disequilibrium (LD) region (r^2>0.6) with an index SNP. LD structure within a locus was

constructed by calculating correlations between SNPs within a locus (1KG v20130502).

CAVIAR[97] was then applied to the summary association statistics and LD structure for each index SNP to generate potentially causal (credible) SNPs with a posterior probability of 0.95. In total, we identified 14 credible SNPs from the three GWS loci.

For each locus, candidate genes were identified by mapping credible SNPs based on physical interactions in foetal brain as previously described[98]. One locus (index SNP rs4146336) was mapped to two genes, *LSAMP* and *PTMAP8*, indicating that two credible SNPs (rs13066948 and rs11713893) located in this locus physically interact with these genes.

Genetic correlation: For all phenotypes we performed genetic correlation without constraining the intercept using LDSR. We identified significant genetic correlations using a Bonferroni adjusted P-value < 0.05. For the primary genetic correlation analysis with SQ-R, we included psychiatric conditions[57,63,99–102], personality traits[103–105], measures of intelligence[58,59,106,107], and social traits related to autism[46,55] including scores on the SCDC, as previous research has investigated the phenotypic correlation between these domains and systemizing[10,78,108–112].

To understand the correlation between systemizing and various phenotypes that have been genetically correlated with autism, we used GWAS data from 15 phenotypes including autism. 10 of these phenotypes (cognitive aptitude[59], educational attainment[58], tiredness[113], neuroticism[103], subjective wellbeing[103], schizophrenia[114], major depression[102], depressive symptoms[103], ADHD[63] and chronotype[115]), have been previously reported to be significantly genetically correlated with autism out of 234 phenotypes tested using LDHub[62] (P < 2.1x10^{-4}). We excluded college degree from this list, as previous work has identified near perfect genetic correlation between educational attainment and college dsegree[58]. In addition, we included data from friendship satisfaction[55], family satisfaction[55], systemizing, and self-reported empathy[46], all of which are also significantly genetically correlated with autism with P < 2.1x10^{-4}. These four additional phenotypes were not included in the previous paper which investigated genetic correlations with autism. Details of sample sizes with PMIDs/DOIs are provided in **Supplementary Table 12**. Cross trait genetic correlations were computed for all 15 phenotypes, and results were corrected for multiple testing using Bonferroni correction. A correlogram was created after using hierarchical clustering to cluster the phenotypes.

To investigate if the combination of negative genetic correlation social traits and positive genetic correlation for non-social traits is specific to autism, we conducted a genetic correlation between all psychiatric conditions for which we had access to summary GWAS

statistics (ADHD[63], Anxiety[116], Autism[57], Anorexia[101], Bipolar Disorder[99], Major Depressive

Disorder[102], OCD[117,118], PTSD[119], and Schizophrenia[114]) and SQ-R, self-reported empathy

measured using the EQ[46] and friendship satisfaction[55]. We chose friendship satisfaction and

self-reported empathy as representative of social traits as these are the most relevant to the

social domain of autism that we had access to GWAS summary statistics. The EQ is a short,

40-item self-report measure of empathy, which has been widely used and has good

psychometric properties[60,120]. For instance, in the DSM-5, one of the criteria for autism is

difficulties in making friends[1]. Additionally, differences in aspects of empathy compared to

the neurotypical population have been widely reported in autism[50,51,121], and is one of the

items in measures such as ADOS-G.

GWIS and GSEM: To investigate if the SQ-R is genetically correlated with autism

independent of the genetic effects of educational attainment, we constructed a unique SQ-R

phenotype after conditioning on the genetic effects of educational attainment using GWIS[122].

GWIS takes into account the genetic covariance between the two phenotypes to calculate the

unique component of the phenotypes as a function of the genetic covariance and the h^2_{SNP}.

Prior to performing GWIS, we standardized the beta coefficients for the SQ-R GWAS by

using the following formula:

$$\widehat{B_{std}} = \hat{B}\sqrt{\frac{2(MAF)(1 - MAF)}{\sigma^2_y}}$$

Where $\widehat{B_{std}}$ is the standardized regression coefficients, $\hat{B}$ is the regression coefficient

obtained from the non-standardized GWAS, MAF is the minor allele frequency, σ^2_y is the

variance of the SQ-R. This equation is explained in detail in the **Supplementary Note**. We

conducted GWIS using only educational attainment as we were unclear if the GWAS of

cognitive aptitude[59] was conducted on a standardized phenotype. Further, there is a high

genetic correlation between cognitive aptitude and educational attainment. In addition to

GWIS, to validate the findings, we conducted GSEM[123], a complementary but independent

method. GSEM uses the correlations and covariances calculated using LDSR after accounting

for sample overlap.

Polygenic scores in the SSC, AGRE, EU-AIMS LEAP and Paris cohorts: We

generated polygenic scores for SQ-R (mean weighted score of all the alleles that contribute to

higher systemizing) in 2,221 probands from the Simons Simplex Collection (Discovery

dataset). We downloaded genotype data from the SSC from SFARI base (https://www.sfari.org/resource/sfari-base/). Individuals were genotyped on three different platforms: Illumina Omni2.5, Illumina 1Mv3, or Illumina 1Mv1. Informed consent or assent was obtained from all participants. In addition, the research team obtained ethical approval from the Cambridge Human Biology Research Ethics Committee to access and analyse the de-identified data from the Simons Simplex collection. We conducted a stringent quality control and imputation to generate genotypes used in this analysis for each of the platforms separately. The full pipeline is available here: https://github.com/autism-research-centre/SSC_liftover_imputation. Briefly, individuals were excluded if they had: a genotyping rate < 95%, excessive or low heterozygosity (less or more than 3 SD from the mean), mismatched reported and genetic sex, and families with mendelian errors > 5%. We further removed SNPs that significantly deviated from Hardy-Weinberg Equilibrium ($P < 1x10^{-6}$), had mendelian errors in more than 10% of the families, and SNPs that were not genotyped in more than 10% of the families. We then conducted multidimensional scaling using the HapMap3 phase 3 population using the unrelated individuals CEU and TSI populations as representatives of the European population. This was conducted only in the parents to retain unrelated individuals for multidimensional scaling. Genetic principal components were calculated using only SNPs with minor allele frequency > 5%, and pruning the SNPs in Plink using an r^2 of 0.2. We excluded families from further downstream analyses if either one the parents were greater or less than 5 standard deviations from the means of the first two genetic principal components calculated using only the unrelated individuals in HapMap3 CEU and TSI populations. Quality control was done using Plink v 1.9 and R. Phasing and imputation was conducted using the Michigan Imputation Server (https://imputationserver.sph.umich.edu/start.html) using the 1000 genomes Phase 3 v5 as the reference panel.

Polygenic scores were generated using PRSice2 (https://choishingwan.github.io/PRSice/) for the SQ-R using the non-stratified GWAS data. We calculated the mean polygenic score for each of the 2,221 probands in the SSC, after clumping SNPs using an R^2 threshold of 0.1. Prior to generating polygenic scores, we confirmed that the probands were not related to each other using identity by descent PI-HAT > 0.15 as a relatedness cut-off. We used a P-value threshold of 1 as previous research on educational attainment, subjective wellbeing and social relationship satisfaction, all suggest that the maximum variance explained is at a threshold of 1[58,103]. This is expected for highly

polygenic traits where many SNPs incrementally contribute to the variance explained[124].
Polygenic scoring was done using standardized scores on two different phenotypes as the
dependent variable (RBS-R and the social and communication domain of the ADOS-G). We
included sex, platform, the first 15 genetic principal components and standardized full-scale
IQ as covariates. In addition, for the analysis of ADOS-G, we included the ADOS-G module
as a covariate. Linear regression was conducted in R. A total of 135,233 SNPs were included
in the polygenic score analyses after clumping and thresholding.

To validate the polygenic scores, we conducted additional polygenic score analysis
using data combined from the AGRE, EU-AIMS LEAP and Paris cohorts. We followed
similar quality control and imputation procedures to the SSC cohort. Given that this dataset
was a mix of related and unrelated individuals, we chose unrelated individuals using a
genomic relationship matrix (GRM) as provided in GCTA (--grm-cutoff 0.05)[125]. To
calculate GRMs, we included only SNPs with minor allele frequency > 1%. Scripts are
provided here: https://github.com/vwarrier/PARIS_LEAP_analysis. Polygenic scores were
calculated using PRSice2 as described for the SSC data. Given the differences in dataset,
polygenic scores were calculated separately for the AGRE dataset, and the EU-AIMS LEAP
and Paris datasets combined. For each regression, we included sex and the first 10 genetic
principal components (standardized). The dependent variables were standardized scores on
the RBS-R (N = 426) and the ADOS-G social and communication subscale (N = 475). IQ
information was unavailable for most individuals, and hence we did not include IQ as a
covariate. We combined the results of the EU-AIMS LEAP and Paris cohorts, and the AGRE
dataset using inverse variance weighted fixed-effect meta-analysis using the formula below:

$$w_i = 1/SE_i^2$$

$$SE_{meta} = \sqrt{1/\Sigma_i w_i}$$

$$Beta_{meta} = \Sigma_i \beta_i w_i / \Sigma_i w_i$$

Where β_i is the standardized regression coefficient of the polygenic scores, SE_i is the
associated standard error, and w_i is the weight.

Bivariate GREML: We conducted bivariate genetic correlation using GCTA GREML
to test the genetic correlation between the ADOS social and communication domains and the
RBS-R scores. We created a GRM after including autistic individuals from the SSC, AGRE,

EU-AIMS LEAP and Paris cohorts. We excluded SNPs and individuals using the same

quality control pipeline as applied to the SSC dataset outlined in the section above. We

further restricted our analysis only to SNPs with a minor allele frequency > 1%. We excluded

related individuals (--grm-cutoff 0.05) resulting in a total of 2,989 individuals. Of this, 2,652

individuals had scores for the ADOS social and communication domain and 2,550

individuals had scores on the RBS-R. We included sex and the first 10 genetic principal

components as covariates.

569   Data availability

The SQ-R GWAS results are available from 23andMe. The full set of summary statistics can

be made available to qualified investigators who enter into an agreement with 23andMe that

protects participant confidentiality. Interested investigators should email dataset-

request@23andme.com for more information. Top SNPs (n = 10,000) can be visualized here:

https://ghfc.pasteur.fr. Data for ALSPAC can be requested here:

http://www.bristol.ac.uk/alspac/researchers/access/. Data from the Simons Simplex

Collection can be requested here: https://www.sfari.org/resource/sfari-base/. Summary

GWAS statistics were downloaded from the PGC consortium:

http://www.med.unc.edu/pgc/results-and-downloads. Data for chronotype was downloaded

from http://www.t2diabetesgenes.org/data/. Data for self-reported tiredness was downloaded

from http://www.ccace.ed.ac.uk/node/335.

582   Software and code availability

Genomic-SEM: https://github.com/MichelNivard/GenomicSEM

GWIS: https://sites.google.com/site/mgnivard/gwis

Plink: https://www.cog-genomics.org/plink2/

PRSice2: https://choishingwan.github.io/PRSice/

CAVIAR: http://genetics.cs.ucla.edu/caviar/

Michigan Imputation Server: https://imputationserver.sph.umich.edu/index.html

Custom code for quality control of the SSC and the other cohorts can be downloaded from

https://github.com/autism-research-centre/SSC_liftover_imputation

(DOI: 10.5281/zenodo.3342561) and from

https://github.com/vwarrier/PARIS_LEAP_analysis

(DOI: 10.5281/zenodo.3342569)

References

1.   American Psychiatric Association. *Diagnostic and statistical manual of mental*
disorders (5th ed.). (2013).

2.   Happé, F. & Ronald, A. The 'Fractionable Autism Triad': A Review of Evidence from
Behavioural, Genetic, Cognitive and Neural Research. *Neuropsychol. Rev.* **18,** 287–
304 (2008).

3.   Happé, F., Ronald, A. & Plomin, R. Time to give up on a single explanation for
autism. *Nat. Neurosci.* **9,** 1218–1220 (2006).

4.   Shuster, J., Perry, A., Bebko, J. & Toplak, M. E. Review of Factor Analytic Studies
Examining Symptoms of Autism Spectrum Disorders. *J. Autism Dev. Disord.* **44,** 90–
110 (2014).

5.   Mandy, W. P. L. & Skuse, D. H. Research Review: What is the association between
the social-communication element of autism and repetitive interests, behaviours and
activities? *J. Child Psychol. Psychiatry* **49,** 795–808 (2008).

6.   Palmer, C. J., Paton, B., Enticott, P. G. & Hohwy, J. 'Subtypes' in the Presentation of
Autistic Traits in the General Adult Population. *J. Autism Dev. Disord.* **45,** 1291–1301
(2015).

7.   Guthrie, W., Swineford, L. B., Wetherby, A. M. & Lord, C. Comparison of DSM-IV
and DSM-5 Factor Structure Models for Toddlers With Autism Spectrum Disorder. *J.*
Am. Acad. Child Adolesc. Psychiatry **52,** 797–805.e2 (2013).

8.   Frazier, T. W., Youngstrom, E. A., Kubu, C. S., Sinclair, L. & Rezai, A. Exploratory
and Confirmatory Factor Analysis of the Autism Diagnostic Interview-Revised. *J.*
Autism Dev. Disord. **38,** 474–480 (2008).

9.   Grove, R., Baillie, A., Allison, C., Baron-Cohen, S. & Hoekstra, R. A. Empathizing,
systemizing, and autistic traits: Latent structure in individuals with autism, their
parents, and general population controls. *J. Abnorm. Psychol.* **122,** 600–609 (2013).

10.   Wheelwright, S. J. *et al.* Predicting Autism Spectrum Quotient (AQ) from the
Systemizing Quotient-Revised (SQ-R) and Empathy Quotient (EQ). *Brain Res.* **1079,**
47–56 (2006).

11.   Svedholm-Häkkinen, A. M., Halme, S. & Lindeman, M. Empathizing and systemizing
are differentially related to dimensions of autistic traits in the general population. *Int.*

629   *J. Clin. Heal. Psychol.* **18,** 35–42 (2018).

12.   Greenberg, D. M., Warrier, V., Allison, C. & Baron-Cohen, S. Testing the
Empathizing-Systemizing theory of sex differences and the Extreme Male Brain theory
of autism in half a million people. *Proc. Natl. Acad. Sci. U. S. A.* 201811032 (2018).
doi:10.1073/pnas.1811032115

13.   Ronald, A., Happe, F. & Plomin, R. The genetic relationship between individual
differences in social and nonsocial behaviours characteristic of autism. *Dev. Sci.* **8,**
444–458 (2005).

14.   Ronald, A. *et al.* Genetic Heterogeneity Between the Three Components of the Autism
Spectrum: A Twin Study. *J. Am. Acad. Child Adolesc. Psychiatry* **45,** 691–699 (2006).

15.   Dworzynski, K., Happé, F., Bolton, P. & Ronald, A. Relationship Between Symptom
Domains in Autism Spectrum Disorders: A Population Based Twin Study. *J. Autism*
Dev. Disord. **39,** 1197–1210 (2009).

16.   Norbury, C. F. Practitioner Review: Social (pragmatic) communication disorder
conceptualization, evidence and clinical implications. *J. Child Psychol. Psychiatry* **55,**
204–216 (2014).

17.   Uljarević, M., Evans, D. W., Alvares, G. A. & Whitehouse, A. J. O. Short report:
relationship between restricted and repetitive behaviours in children with autism
spectrum disorder and their parents. *Mol. Autism* **7,** 29 (2016).

18.   Georgiades, S. *et al.* Investigating phenotypic heterogeneity in children with autism
spectrum disorder: a factor mixture modeling approach. *J. Child Psychol. Psychiatry*
54, 206–215 (2013).

19.   St Pourcain, B. *et al.* Variability in the common genetic architecture of social-
communication spectrum phenotypes during childhood and adolescence. *Mol. Autism*
5, 18 (2014).

20.   Cantor, R. M. *et al.* ASD restricted and repetitive behaviors associated at 17q21.33:
genes prioritized by expression in fetal brains. *Mol. Psychiatry* (2017).
doi:10.1038/mp.2017.114

21.   Alarcón, M., Cantor, R. M., Liu, J., Gilliam, T. C. & Geschwind, D. H. Evidence for a
language quantitative trait locus on chromosome 7q in multiplex autism families. *Am.*
J. Hum. Genet. **70,** 60–71 (2002).

22.   Lowe, J. K., Werling, D. M., Constantino, J. N., Cantor, R. M. & Geschwind, D. H.
Social responsiveness, an autism endophenotype: genomewide significant linkage to
two regions on chromosome 8. *Am. J. Psychiatry* **172,** 266–75 (2015).

23.   Shao, Y. *et al.* Fine mapping of autistic disorder to chromosome 15q11-q13 by use of
phenotypic subtypes. *Am. J. Hum. Genet.* **72,** 539–48 (2003).

24.   Cannon, D. S. *et al.* Genome-wide linkage analyses of two repetitive behavior
phenotypes in Utah pedigrees with autism spectrum disorders. *Mol. Autism* **1,** 3 (2010).

25.   Tao, Y. *et al.* Evidence for contribution of common genetic variants within
chromosome 8p21.2-8p21.1 to restricted and repetitive behaviors in autism spectrum
disorders. *BMC Genomics* **17,** 163 (2016).

26.   Ruzich, E. *et al.* Measuring autistic traits in the general population: a systematic

review of the Autism-Spectrum Quotient (AQ) in a nonclinical population sample of
6,900 typical adult males and females. *Mol. Autism* **6,** 2 (2015).

27.   Baron-Cohen, S., Wheelwright, S. J., Skinner, R., Martin, J. & Clubley, E. The autism-
spectrum quotient (AQ): evidence from Asperger syndrome/high-functioning autism,
males and females, scientists and mathematicians. *J. Autism Dev. Disord.* **31,** 5–17
(2001).

28.   Posserud, M.-B., Lundervold, A. J. & Gillberg, C. Autistic features in a total
population of 7-9-year-old children assessed by the ASSQ (Autism Spectrum
Screening Questionnaire). *J. Child Psychol. Psychiatry* **47,** 167–175 (2006).

29.   Constantino, J. N. & Todd, R. D. Autistic Traits in the General Population. *Arch. Gen.*
Psychiatry **60,** 524 (2003).

30.   Constantino, J. N. *et al.* The factor structure of autistic traits. *J. Child Psychol.*
Psychiatry **45,** 719–726 (2004).

31.   Hoekstra, R. A. *et al.* Heritability of autistic traits in the general population. *Arch.*
Pediatr. Adolesc. Med. **161,** 372–7 (2007).

32.   Ronald, A. & Hoekstra, R. A. Autism spectrum disorders and autistic traits: A decade
of new twin studies. *Am. J. Med. Genet. Part B Neuropsychiatr. Genet.* **156,** 255–274
(2011).

33.   de Zeeuw, E. L., van Beijsterveldt, C. E. M., Hoekstra, R. A., Bartels, M. & Boomsma,
D. I. The etiology of autistic traits in preschoolers: a population-based twin study. *J.*
Child Psychol. Psychiatry **58,** 893–901 (2017).

34.   Wheelwright, S. J., Auyeung, B., Allison, C. & Baron-Cohen, S. Defining the broader,
medium and narrow autism phenotype among parents using the Autism Spectrum
Quotient (AQ). *Mol. Autism* **1,** 10 (2010).

35.   Sucksmith, E., Roth, I. & Hoekstra, R. A. Autistic Traits Below the Clinical
Threshold: Re-examining the Broader Autism Phenotype in the 21st Century.
Neuropsychol. Rev. **21,** 360–389 (2011).

36.   Sasson, N. J., Lam, K. S., Parlier, M., Daniels, J. L. & Piven, J. Autism and the broad
autism phenotype: familial patterns and intergenerational transmission. *J. Neurodev.*
Disord. **5,** 11 (2013).

37.   Constantino, J. N. & Todd, R. D. Intergenerational transmission of subthreshold
autistic traits in the general population. *Biol. Psychiatry* **57,** 655–660 (2005).

38.   Murray, A. L., McKenzie, K., Kuenssberg, R. & Booth, T. Do the Autism Spectrum
Quotient (AQ) and Autism Spectrum Quotient Short Form (AQ-S) Primarily Reflect
General ASD Traits or Specific ASD Traits? A Bi-Factor Analysis. *Assessment* **24,**
444–457 (2017).

39.   Lundström, S. *et al.* Autism Spectrum Disorders and Autisticlike Traits. *Arch. Gen.*
Psychiatry **69,** 46 (2012).

40.   Robinson, E. B. *et al.* Evidence that autistic traits show the same etiology in the
general population and at the quantitative extremes (5%, 2.5%, and 1%). *Arch. Gen.*
Psychiatry **68,** 1113–21 (2011).

41.   Ronald, A., Happé, F., Price, T. S., Baron-Cohen, S. & Plomin, R. Phenotypic and

genetic overlap between autistic traits at the extremes of the general population. *J. Am.*
Acad. Child Adolesc. Psychiatry **45,** 1206–14 (2006).

42. Colvert, E. *et al.* Heritability of autism spectrum disorder in a UK population-based
twin sample. *JAMA Psychiatry* **72,** 415–23 (2015).

43. Bralten, J. *et al.* Autism spectrum disorders and autistic traits share genetics and
biology. *Mol. Psychiatry* (2017). doi:10.1038/mp.2017.98

44. St Pourcain, B. *et al.* ASD and schizophrenia show distinct developmental profiles in
common genetic overlap with population-based social communication difficulties.
Mol. Psychiatry (2017). doi:10.1038/mp.2016.198

45. Robinson, E. B. *et al.* Genetic risk for autism spectrum disorders and neuropsychiatric
variation in the general population. *Nat. Genet.* **48,** 552–5 (2016).

46. Warrier, V. *et al.* Genome-wide analyses of self-reported empathy: Correlations with
autism, schizophrenia, and anorexia nervosa. *Transl. Psychiatry* **8,** (2018).

47. Warrier, V. *et al.* Genome-wide meta-analysis of cognitive empathy: heritability, and
correlates with sex, neuropsychiatric conditions and cognition. *Mol. Psychiatry* (2017).

48. Cantor, R. M. *et al.* ASD restricted and repetitive behaviors associated at 17q21.33:
genes prioritized by expression in fetal brains. *Mol. Psychiatry* **23,** 993–1000 (2018).

49. Baron-Cohen, S. The hyper-systemizing, assortative mating theory of autism. *Prog.*
Neuropsychopharmacol. Biol. Psychiatry **30,** 865–72 (2006).

50. Auyeung, B. *et al.* The Children's Empathy Quotient and Systemizing Quotient: Sex
Differences in Typical Development and in Autism Spectrum Conditions. *J. Autism*
Dev. Disord. **39,** 1509–1521 (2009).

51. Baron-Cohen, S. *et al.* Attenuation of typical sex differences in 800 adults with autism
vs. 3,900 controls. *PLoS One* **9,** e102251 (2014).

52. Asperger, H. 'Autistic psychopathy' in childhood. in *Autism and Asperger syndrome*
(ed. Frith, U.) 37–92 (Cambridge University Press, 1944).
doi:10.1017/CBO9780511526770.002

53. Kanner, L. Autistic disturbances of affective contact. *Nerv. Child J. Psychopathol.*
Psychother. Ment. Hyg. Guid. Child **2** 217–50 (1943).

54. Auyeung, B. *et al.* Foetal testosterone and the child systemizing quotient. *Eur. J.*
Endocrinol. **155,** S123–S130 (2006).

55. Warrier, V., the 23andMe Research team, Bourgeron, T. & Baron-Cohen, S. Genome-
wide association study of social relationship satisfaction: significant loci and
correlations with psychiatric conditions. *bioRxiv* 196071 (2017). doi:10.1101/196071

56. Palmer, E. E. *et al.* A Recurrent De Novo Nonsense Variant in ZSWIM6 Results in
Severe Intellectual Disability without Frontonasal or Limb Malformations. *Am. J.*
Hum. Genet. **101,** 995–1005 (2017).

57. Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum
disorder. *Nat. Genet.* **51,** 431–444 (2019).

58. Okbay, A. *et al.* Genome-wide association study identifies 74 loci associated with
educational attainment. *Nature* **533,** 539–542 (2016).

59.   Sniekers, S. *et al.* Genome-wide association meta-analysis of 78,308 individuals
755   identifies new loci and genes influencing human intelligence. *Nat. Genet.* (2017).
756   doi:10.1038/ng.3869

757   60.   Baron-Cohen, S. & Wheelwright, S. J. The Empathy Quotient: an investigation of
758   adults with Asperger syndrome or high functioning autism, and normal sex
759   differences. *J. Autism Dev. Disord.* **34,** 163–75 (2004).

760   61.   Skuse, D. H., Mandy, W. P. L. & Scourfield, J. Measuring autistic traits: heritability,
761   reliability and validity of the Social and Communication Disorders Checklist. *Br. J.*
762   *Psychiatry* **187,** 568–572 (2005).

763   62.   Zheng, J. *et al.* LD Hub: a centralized database and web interface to perform LD score
764   regression that maximizes the potential of summary level GWAS data for SNP
765   heritability and genetic correlation analysis. *Bioinformatics* **33,** 272–279 (2016).

766   63.   Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for
767   attention deficit/hyperactivity disorder. *Nat. Genet.* **51,** 63–75 (2019).

768   64.   Ripke, S. *et al.* Genome-wide association analysis identifies 13 new risk loci for
769   schizophrenia. *Nat. Genet.* **45,** 1150–1159 (2013).

770   65.   Stahl, E. *et al.* Genomewide association study identifies 30 loci associated with bipolar
771   disorder. *bioRxiv* 173062 (2018). doi:10.1101/173062

772   66.   Baron-Cohen, S. The extreme male brain theory of autism. *Trends Cogn. Sci.* **6,** 248–
773   254 (2002).

774   67.   Tung, J. Y. *et al.* Efficient replication of over 180 genetic associations with self-
775   reported medical data. *PLoS One* **6,** e23473 (2011).

776   68.   Do, C. B. *et al.* Web-based genome-wide association study identifies two novel loci
777   and a substantial genetic component for Parkinson's disease. *PLoS Genet.* **7,** e1002141
778   (2011).

779   69.   Eriksson, N. *et al.* Novel associations for hypothyroidism include known autoimmune
780   risk loci. *PLoS One* **7,** e34442 (2012).

781   70.   Henn, B. M. *et al.* Cryptic distant relatives are common in both isolated and
782   cosmopolitan genetic samples. *PLoS One* **7,** e34267 (2012).

783   71.   Boyd, A. *et al.* Cohort Profile: The 'Children of the 90s'—the index offspring of the
784   Avon Longitudinal Study of Parents and Children. *Int. J. Epidemiol.* **42,** 111–127
785   (2013).

786   72.   Fischbach, G. D. & Lord, C. The Simons Simplex Collection: A Resource for
787   Identification of Autism Genetic Risk Factors. *Neuron* **68,** 192–195 (2010).

788   73.   Geschwind, D. H. *et al.* The autism genetic resource exchange: a resource for the study
789   of autism and related neuropsychiatric conditions. *Am. J. Hum. Genet.* **69,** 463–6
790   (2001).

791   74.   Charman, T. *et al.* The EU-AIMS Longitudinal European Autism Project (LEAP):
792   clinical characterisation. *Mol. Autism* **8,** 27 (2017).

793   75.   Durand, C. M. *et al.* Mutations in the gene encoding the synaptic scaffolding protein
794   SHANK3 are associated with autism spectrum disorders. *Nat. Genet.* **39,** 25–27
795   (2007).

76.   Groen, Y., Fuermaier, A. B. M., Den Heijer, A. E., Tucha, O. & Althaus, M. The
Empathy and Systemizing Quotient: The psychometric properties of the Dutch version
and a review of the cross-cultural stability. *J. Autism Dev. Disord.* **45,** 2848–64 (2015).

77.   Allison, C., Baron-Cohen, S., Stone, M. H. & Muncer, S. J. Rasch modeling and
confirmatory factor analysis of the systemizing quotient-revised (SQ-R) scale. *Span. J.*
Psychol. **18,** E16 (2015).

78.   Ling, J., Burton, T. C., Salt, J. L. & Muncer, S. J. Psychometric analysis of the
systemizing quotient (SQ) scale. *Br. J. Psychol.* **100,** 539–552 (2009).

79.   Wakabayashi, A. *et al.* Empathizing and systemizing in adults with and without autism
spectrum conditions: Cross-cultural stability. *J. Autism Dev. Disord.* **37,** 1823–1832
(2007).

80.   Bölte, S., Westerwald, E., Holtmann, M., Freitag, C. M. & Poustka, F. Autistic traits
and autism spectrum disorders: the clinical validity of two measures presuming a
continuum of social communication skills. *J. Autism Dev. Disord.* **41,** 66–72 (2011).

81.   Lord, C. *et al.* The Autism Diagnostic Observation Schedule—Generic: A Standard
Measure of Social and Communication Deficits Associated with the Spectrum of
Autism. *J. Autism Dev. Disord.* **30,** 205–223 (2000).

82.   Gotham, K., Risi, S., Pickles, A. & Lord, C. The Autism Diagnostic Observation
Schedule: Revised Algorithms for Improved Diagnostic Validity. *J. Autism Dev.*
Disord. **37,** 613–627 (2007).

83.   Bodfish, J. W., Symons, F. J., Parker, D. E. & Lewis, M. H. Varieties of Repetitive
Behavior in Autism: Comparisons to Mental Retardation. *J. Autism Dev. Disord.* **30,**
237–243 (2000).

84.   Lam, K. S. L. & Aman, M. G. The Repetitive Behavior Scale-Revised: Independent
Validation in Individuals with Autism Spectrum Disorders. *J. Autism Dev. Disord.* **37,**
855–866 (2007).

85.   Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and
missing-data inference for whole-genome association studies By use of localized
haplotype clustering. *Am. J. Hum. Genet.* **81,** 1084–1097 (2007).

86.   Fuchsberger, C., Abecasis, G. R. & Hinds, D. A. minimac2: faster genotype
imputation. *Bioinformatics* **31,** 782–784 (2015).

87.   Bigdeli, T. B. *et al.* A simple yet accurate correction for winner's curse can predict
signals discovered in much larger genome scans. *Bioinformatics* **32,** 2598–2603
(2016).

88.   Delaneau, O., Marchini, J. & Zagury, J.-F. A linear complexity phasing method for
thousands of genomes. *Nat. Methods* **9,** 179–181 (2011).

89.   Howie, B. N., Donnelly, P. & Marchini, J. A flexible and accurate genotype
imputation method for the next generation of genome-wide association studies. *PLoS*
Genet. **5,** e1000529 (2009).

90.   Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based
linkage analyses. *Am. J. Hum. Genet.* **81,** 559–75 (2007).

91.   Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from

polygenicity in genome-wide association studies. *Nat. Genet.* **47,** 291–295 (2015).

92.   Bulik-Sullivan, B. K. *et al.* An atlas of genetic correlations across human diseases and
traits. *Nat. Genet.* **47,** 1236–41 (2015).

93.   Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping
and annotation of genetic associations with FUMA. *Nat. Commun.* **8,** 1826 (2017).

94.   de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: generalized
gene-set analysis of GWAS data. *PLoS Comput. Biol.* **11,** 1–19 (2015).

95.   Ardlie, K. G. *et al.* The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue
gene regulation in humans. *Science (80-.).* **348,** 648–60 (2015).

96.   Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-
wide association summary statistics. *Nat. Genet.* **47,** 1228–1235 (2015).

97.   Hormozdiari, F., Kostem, E., Kang, E. Y., Pasaniuc, B. & Eskin, E. Identifying Causal
Variants at Loci with Multiple Signals of Association. *Genetics* **198,** 497–508 (2014).

98.   Won, H. *et al.* Chromosome conformation elucidates regulatory relationships in
developing human brain. *Nature* **538,** 523–527 (2016).

99.   Sklar, P. *et al.* Large-scale genome-wide association analysis of bipolar disorder
identifies a new susceptibility locus near ODZ4. *Nat. Genet.* **43,** 977–983 (2011).

100.   Pardiñas, A. F. *et al.* Common schizophrenia alleles are enriched in mutation-
intolerant genes and in regions under strong background selection. *Nat. Genet.* 1
(2018). doi:10.1038/s41588-018-0059-2

101.   Duncan, L. E. *et al.* Significant locus and metabolic genetic correlations revealed in
genome-wide association study of anorexia nervosa. *Am. J. Psychiatry* appi.ajp.2017.1
(2017). doi:10.1176/appi.ajp.2017.16121402

102.   Wray, N. R. *et al.* Genome-wide association analyses identify 44 risk variants and
refine the genetic architecture of major depression. *Nat. Genet.* **50,** 668–681 (2018).

103.   Okbay, A. *et al.* Genetic variants associated with subjective well-being, depressive
symptoms, and neuroticism identified through genome-wide analyses. *Nat. Genet.* **48,**
624–633 (2016).

104.   de Moor, M. H. M. *et al.* Meta-analysis of genome-wide association studies for
personality. *Mol. Psychiatry* **17,** 337–49 (2012).

105.   van den Berg, S. M. *et al.* Meta-analysis of genome-wide association studies for
extraversion: findings from the genetics of personality consortium. *Behav. Genet.* **46,**
170–82 (2016).

106.   Rietveld, C. A. *et al.* Common genetic variants associated with cognitive performance
identified using the proxy-phenotype method. *Proc. Natl. Acad. Sci.* **111,** 13790–
13794 (2014).

107.   Rietveld, C. A. *et al.* GWAS of 126,559 individuals identifies genetic variants
associated with educational attainment. *Science* **340,** 1467–71 (2013).

108.   Hambrook, D., Tchanturia, K., Schmidt, U., Russell, T. & Treasure, J. Empathy,
systemizing, and autistic traits in anorexia nervosa: a pilot study. *Br. J. Clin. Psychol.*
47, 335–9 (2008).

109. Russell-Smith, S. N., Bayliss, D. M., Maybery, M. T. & Tomkinson, R. L. Are the
880   autism and positive schizotypy spectra diametrically opposed in empathizing and
881   systemizing? *J. Autism Dev. Disord.* **43,** 695–706 (2013).

882   110. Courty, A. *et al.* Levels of autistic traits in anorexia nervosa: a comparative
883   psychometric study. *BMC Psychiatry* **13,** 222 (2013).

884   111. Nettle, D. Empathizing and systemizing: what are they, and what do they contribute to
885   our understanding of psychological sex differences? *Br. J. Psychol.* **98,** 237–55 (2007).

886   112. Wakabayashi, A. & Kawashima, H. Is empathizing in the E–S theory similar to
887   agreeableness? The relationship between the EQ and SQ and major personality
888   domains. *Pers. Individ. Dif.* **76,** 88–93 (2015).

889   113. Deary, V. *et al.* Genetic contributions to self-reported tiredness. *Mol. Psychiatry* **23,**
890   609–620 (2018).

891   114. Ripke, S. *et al.* Biological insights from 108 schizophrenia-associated genetic loci.
892   *Nature* **511,** 421–7 (2014).

893   115. Jones, S. E. *et al.* Genome-Wide Association Analyses in 128,266 Individuals
894   Identifies New Morningness and Sleep Duration Loci. *PLOS Genet.* **12,** e1006125
895   (2016).

896   116. Otowa, T. *et al.* Meta-analysis of genome-wide association studies of anxiety
897   disorders. *Mol. Psychiatry* (2016). doi:10.1038/mp.2015.197

898   117. Mattheisen, M. *et al.* Genome-wide association study in obsessive-compulsive
899   disorder: results from the OCGAS. *Mol. Psychiatry* **20,** 337–344 (2015).

900   118. International Obsessive Compulsive Disorder Foundation Genetics Collaborative
901   (IOCDF-GC) and OCD Collaborative Genetics Association Studies (OCGAS), P. D. *et*
902   *al.* Revealing the complex genetic architecture of obsessive-compulsive disorder using
903   meta-analysis. *Mol. Psychiatry* **23,** 1181–1188 (2018).

904   119. Duncan, L. E. *et al.* Largest GWAS of PTSD (N=20 070) yields genetic overlap with
905   schizophrenia and sex differences in heritability. *Mol. Psychiatry* **23,** 666–673 (2017).

906   120. Allison, C., Baron-Cohen, S., Wheelwright, S. J., Stone, M. H. & Muncer, S. J.
907   Psychometric analysis of the Empathy Quotient (EQ). *Pers. Individ. Dif.* **51,** 829–835
908   (2011).

909   121. Kok, F. M., Groen, Y., Becke, M., Fuermaier, A. B. M. & Tucha, O. Self-Reported
910   Empathy in Adult Women with Autism Spectrum Disorders – A Systematic Mini
911   Review. *PLoS One* **11,** e0151568 (2016).

912   122. Nieuwboer, H. A., Pool, R., Dolan, C. V., Boomsma, D. I. & Nivard, M. G. GWIS:
913   Genome-Wide Inferred Statistics for functions of multiple phenotypes. *Am. J. Hum.*
914   *Genet.* **99,** 917–927 (2016).

915   123. Grotzinger, A. D. *et al.* Genomic structural equation modelling provides insights into
916   the multivariate genetic architecture of complex traits. *Nat. Hum. Behav.* 1 (2019).
917   doi:10.1038/s41562-019-0566-x

918   124. Boyle, E. A., Li, Y. I. & Pritchard, J. K. An Expanded View of Complex Traits: From
919   Polygenic to Omnigenic. *Cell* **169,** 1177–1186 (2017).

920   125. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide

921     complex trait analysis. *Am. J. Hum. Genet.* **88,** 76–82 (2011).

Acknowledgements

We are grateful to Michel Nivard and Beate St. Pourcain for their help with the analytical methods. We thank the research participants and employees of 23andMe for making this work possible. We are grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses. VW was funded by St. John's College, Cambridge, and the Cambridge Commonwealth Trust. This study was funded by grants to SBC from the Medical Research Council, the Wellcome Trust, the Autism Research Trust, the Templeton World Charity Foundation,, and to TB from the Institut Pasteur, the CNRS, the INSERM, The Fondamental Foundation, the APHP, the BioPsy Labex and the University Paris Diderot. The research was conducted in association with the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care East of England at Cambridgeshire and Peterborough NHS Foundation Trust. We also received support from the NIHR Cambridge Biomedical Research Centre. We acknowledge with gratitude the generous support of Drs Dennis and Mireille Gillings in strengthening the collaboration between SBC and TB, and between Cambridge University and the Institut Pasteur. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. Data obtained from 23andMe was supported by the National Human Genome Research Institute of the National Institutes of Health (grant number R44HG006981). The UK Medical Research Council and Wellcome (grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. GWAS data was generated by Sample Logistics and Genotyping Facilities at Wellcome Sanger Institute and LabCorp (Laboratory Corporation of America) using support from 23andMe. This publication is the work of the authors who will serve as guarantors for the content of this paper. The iPSYCH (The Lundbeck Foundation Initiative for Integrative Psychiatric Research) team acknowledges funding from The Lundbeck Foundation (grant no R102-A9118 and R155-2014-1724), the Stanley Medical Research Institute, the European Research Council (project no: 294838), the Novo Nordisk Foundation for supporting the Danish National Biobank resource, and grants from Aarhus and Copenhagen Universities and University Hospitals, including support to the iSEQ Center, the GenomeDK HPC facility, and the CIRRAU Center. The project leading to this application has received funding from the Innovative Medicines Initiative 2 Joint Undertaking (JU) under grant agreement No

777394. The JU receives support from the European Union's Horizon 2020 research and innovation programme and EFPIA and AUTISM SPEAKS, Autistica, SFARI. We thank the iPSCH-Broad Autism Group and the EU-AIMS LEAP group for sharing data. A full list of the authors and affiliations in the iPSYCH-Broad autism group and the EU-AIMS LEAP group is provided in the Supplementary Information.

Competing interests

DH and the 23andMe Research Team are employees of 23andMe, Inc. There is no conflict of interest for the other authors.

Author contributions

VW, TB, SBC, and DAH conceived and designed the analysis. CSL, FC, RD, WDW, JB, ADB, JG, GP, 23andMe Research Team, and DAH collected or contributed the data or analysis tools. VW, RT, HW, FC, and WDW performed the analysis. VW, RT, BC, DAH, TB, and SBC wrote the paper. DAH, TB, and SBC supervised the analysis.

†**23andMe Research Team**

Michelle Agee[1], Babak Alipanahi[1], Adam Auton[1], Robert K. Bell[1], Katarzyna Bryc[1], Sarah L. Elson[1], Pierre Fontanillas[1], Nicholas A. Furlotte[1], Karen E. Huber[1], Aaron Kleinman[1], Nadia K. Litterman[1], Jennifer C. McCreight[1], Matthew H. McIntyre[1], Joanna L. Mountain[1], Carrie A.M. Northover[1], Steven J. Pitts[1], J. Fah Sathirapongsasuti[1], Olga V. Sazonova[1], Janie F. Shelton[1], Suyash Shringarpure[1], Chao Tian[1], Joyce Y. Tung[1], Vladimir Vacic[1], and Catherine H. Wilson[1].

[17] 23andMe Inc., Mountain View, California 94043, USA

Figures and figure legends

Figure 1

Schematic diagram of the study. We conducted a GWAS of the SQ-R (N = 51,564) and
quantified SNP heritability (h^2_{SNP}), quantified genetic correlations with multiple phenotypes,
and conducted polygenic score analyses. Additionally, we conducted sex-stratified GWAS of
the SQ-R, and investigated h^2_{SNP} within sex and genetic correlation between sex. Finally, we
investigated the clustering of all phenotypes that are genetically correlated with autism, and
if the social and the non-social phenotypes associated with autism are genetically correlated.

Figure 2: Manhattan and QQ-plots for the three GWAS

Manhattan plot for the three SQ-R GWAS: non-stratified (A), males-only (C), females-only
(E). Significant SNPs are highlighted in red. QQ-plots for the three SQ-R GWAS: non-
stratified (B), males-only (D), females-only (F). SQ-R non-stratified (N = 51,564): λ_{GC} =
1.10, LDSR intercept = 0.99, SQ-R males only (N = 26,063): λ_{GC} = 1.06, LDSR intercept =
0.99, SQ-R females only (N =25,501): λ_{GC} = 1.05, LDSR intercept = 1.01.

Figure 3: Genetic correlation between the SQ-R and other phenotypes, and GWIS and GSEM estimates between SQ, educational attainment and cognitive aptitude

3A: Genetic correlations between the SQ-R and multiple other phenotypes provided. The bars represent 95% confidence intervals. Sample sizes and PMID are provided in Supplementary Table 9. The following genetic correlations were significant after Bonferroni correction: Autism (r_g = 0.26±0.06; P = 3.35x10^{-5}, N = 46,350), Years of Schooling 2016 (r_g = 0.13±0.03; P = 4.73x10^{-5}, N = 293,723), College completion (r_g = 0.18±0.05; P = 1.30x10^{-3}, N = 95427), and Cognitive aptitude (r_g = 0.19±0.04; P = 2.35x10^{-5}, N = 78,308). 3B: Results

of the GWIS analysis. Red lines represent genetic correlation with the SQ-R, blue lines represent genetic correlations with the SQ-R independent of the genetic effects of educational attainment. The bars represent 95% confidence intervals. 3C: Path diagrams providing the results of the standardized SEM models to investigate if the SQ-R is genetically correlated with autism independent of the genetic effects of cognitive aptitude (CA_g) and educational attainment (EA_g).GWIS is Genome-wide inferred statistics; GSEM is Genomic structural equation modeling.

Figure 4: Genetic correlogram of autism and related traits, and genetic correlations

between social and non-social traits and multiple psychiatric conditions

 4A: Correlogram of genetic correlations between all phenotypes that are genetically
 correlated with autism. Please note the upper and lower triangle are identical. Asterisk
 (provided only in the lower triangle) represent significant correlations after Bonferroni

correction. Genetic correlations have been clustered using hierarchical clustering. Colour as provides the magnitude of genetic correlation. 4B: Genetic correlation between empathy, friendship satisfaction, and systemizing with nine psychiatric conditions. Only autism was significantly genetically correlated with all three phenotypes.