Natural Variation of *OsLG3* Controls Drought Stress Tolerance in Rice by Inducing ROS Scavenging

Haiyan Xiong[1†], Jianping Yu[1†], Jinjie Li[1], Xin Wang[1], Pengli Liu[1], Hongliang Zhang[1], Yan Zhao[1], Zhigang Yin[1], Yang Li[1], Yan Guo[2], Binying Fu[3], Wensheng Wang[3], Zhikang Li[3], Jauhar Ali[4] and Zichao Li[1]*

[1]Key Lab of Crop Heterosis and Utilization of Ministry of Education / Beijing Key Lab of Crop Genetic Improvement, China Agricultural University, Beijing 100193, China

[2]State Key Laboratory of Plant Physiology and Biochemistry, College of Biological Sciences, China Agricultural University, Beijing 100193, China

[3]Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing, 100081, China

[4]International Rice Research Institute, DAPO Box 7777, Metro Manila 1301, Philippines

[†]These authors contributed equally to this work

*Corresponding author; E-mail: lizichao@cau.edu.cn; Tel: +86 10 62731414

Abstract

Background: Improving performance of rice under drought stress has potential to significant impact on rice productivity. Previously we reported that *OsLG3* positively control rice grain length and yield.

Results: In this study, we found that *OsLG3* was more strongly expressed in upland rice compared to lowland rice under drought stress condition. Candidate gene association analysis showed that the natural variation in *OsLG3* was associated with tolerance to water deficit stress in germinating rice seeds. Transgenic rice with enhanced *OsLG3* expression exhibited improved tolerance to drought and that is most likely due to enhanced ROS scavenging efficiency. Phylogenetic analysis and pedigree records indicated that the tolerant allele of *OsLG3* has potential to improve drought tolerance of *japonica* rice.

Conclusions: Collectively, our work revealed that the natural variation of *OsLG3* contributes to rice drought tolerance and the elite allele of *OsLG3* is a promising genetic resource for the development of drought-tolerant and high-yield rice varieties.

Keywords: rice; nature variation; *OsLG3*; drought tolerance; ROS

Background

Agriculture experience more than 50% of average yield losses worldwide due to abiotic stress, especially drought [1, 2]. Rice (*Oryza sativa* L.), one of the most important staple food for over half the world's population, requires high inputs of water during growth, which results in a number of production challenges due to water shortages and inadequate rainfall during the rice growing season [3]. In the face of these challenges, enhanced performance of rice under drought stress has the potential to significantly improve rice productivity.

Plants have evolved complex signaling pathways that enable them to respond and adapt to unfavorable environmental conditions through morphological, physiological, and biochemical changes [4-6]. The generation of reactive oxygen species (ROS) is a key process in plant responsiveness to various biotic and abiotic stresses. ROS are important signal transduction molecules, but also toxic by-products of stress metabolism, depending on their overall cellular amount [7]. At low to moderate concentrations, ROS are likely to function as secondary messengers in stress-signaling pathways, triggering stress defense/adaptation reactions. However, when ROS levels reach above a certain threshold, they can trigger progressive oxidative damage resulting in retarded growth and eventually cell death [8]. Plants have developed flexible ROS scavenging and regulation pathways to maintain homeostasis and avoid over-accumulation of ROS in cells. There are some evidences supporting that overproduction of ROS-scavenging related genes and synthesis of various functional proteins such as superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT) could increase tolerance to oxidative stress. For example, overexpression of

SNAC3 (Stress-responsive NAM, ATAF1/2, and CUC2 gene 3), increases rice drought and

heat tolerance by modulating ROS homeostasis through regulating the expression of genes

participated in ROS-scavenging [9]. Overexpression of an Ethylene-responsive factor (ERF)

family gene, *SERF1* (Salt-Responsive ERF1), improved salinity tolerance in rice, mainly

due to the regulation of ROS-dependent signaling during the initial phase of salt stress [10].

Overexpressing of *JERF3* (Jasmonate and Ethylene-Responsive Factor 3) improved the

expression of genes involved in ROS-scavenging and enhanced tolerance to drought,

freezing, and salt in tobacco [11]. Moreover, overexpression of the mitogen-activated

protein kinase kinase kinase (MAPKKK) gene *DSM1* (Drought-hypersensitive Mutant 1) in

rice, increased drought stress tolerance by regulating ROS scavenging. Conversely,

deficiency in *DSM1* resulted in a decrease in ROS scavenging and increased drought

hypersensitivity [12].

Transcription factors (TFs) play central roles in the regulation of gene expression in

the stress signaling and adaptation networks [13, 14]. One set of these is the

APETALA2/Ethylene Responsive Factor (AP2/ERF) superfamily. It has been implicated

that the ERF proteins play diverse roles in cellular processes involving flower development,

spikelet meristem determinacy, floral meristem, plant growth, pathogens and abiotic stress

tolerance [15-23]. There are emerging evidences to support that ERF proteins are involved

in response and adaptation to drought stress. For instance, transgenic rice lines

overexpression ERF TFs including SUB1A, *OsEREBP1*, *AP37*, *AP59*, *HYR* and *OsERF71*,

all showed strong resistance to drought stress [24-28]. Other ERF genes including *HARDY*,

T*RANSLUCENT GREEN* (*TG*) [29] and DREB genes [30, 31] from *Arabidopsis*, *TSRF1*

[32] from tomato, *TaERF3* [33] from wheat (*Triticum aestivum*), *GmERF3* [34] from

soybean, *JERF3* [11] from tobacco and *SodERF3* [35] from sugarcane (*Saccharum*

officinarum), have also been found to be involved in responses to water deficit stress

condition. Overall, these findings suggested that ERF TFs offer the potential for

engineering crops in a way that makes them more efficient under drought stress condition.

Linkage disequilibrium (LD)-based association mapping has been proven to be a

powerful tool for dissecting complex agronomic traits and identifying alleles that can

contribute to crop improvement [36-39]. Candidate gene association analysis, an effective

method to validate targets, has become easier and cheaper with the advances in

next-generation sequencing (NGS) technology facilitating the discovery and detection of

single nucleotide polymorphisms (SNPs) [40]. This strategy has been used successfully in

the genetic dissection of allelic diversity of genes controlling fatty acid content, kernel size,

ABA content, α-tocopherol and β-carotene content and aluminum tolerance in maize and

rice [41, 42]. There have also been reports of association studies on crop drought tolerance.

For instance, Y Lu, S Zhang, T Shah, C Xie, Z Hao, X Li, M Farkhari, JM Ribaut, M Cao,

T Rong, et al. [43] and Y Xue, ML Warburton, M Sawkins, X Zhang, T Setter, Y Xu, P

Grudloyma, J Gethi, J-M Ribaut, W Li, et al. [44] identified some QTLs underlying

drought tolerance in maize by genome-wide association analysis. S Liu, X Wang, H Wang,

H Xin, X Yang, J Yan, J Li, LS Tran, K Shinozaki, K Yamaguchi-Shinozaki, et al. [45]

found that DNA polymorphisms in the promoter region of *ZmDREB2.7* were associated

with maize drought tolerance. Analysis of the association found that an 82 bp insertion in

ZmNAC111 and 366 bp insertion in *ZmVPP1* affected drought tolerance in Maize [46, 47].

Recently, an association study of 136 wild and four cultivated rice accessions identified

three coding SNPs and one haplotype in a DREB (Dehydration Responsive Element

Binding) transcription factor (TF), *OsDREB1F*, that are potentially associated with drought

tolerance [48], and nine candidate SNPs were identified by association mapping of the ratio

of deep rooting in rice [49]. However, a role for these candidate genes or their causative

variations in enhanced drought tolerance remains to be verified experimentally.

Here, we characterize the role of an ERF family TF, *OsLG3* (LOC_Os03g08470), in

rice drought tolerance. *OsLG3* is located at the same locus as *OsERF62* [23] and *OsRAF* (a

Root Abundant Factor gene in *Oryza sativa*) [50]. In our previous work, we demonstrated

that *OsLG3* plays a positive role in rice grain length without affecting the grain quality [51].

In this study, we identified the natural variation in the promoter region of *OsLG3* was

associated with tolerance to drought stress among different rice accessions. Overexpression

of *OsLG3* in transgenic lines resulted in increased drought tolerance via regulating ROS

homeostasis. *OsLG3* function as a pleiotropic gene which can contribute to rice grain

length and drought stress tolerance together. These data provide insights that the elite allele

of *OsLG3* is a promising genetic resource for the genetic improvement of rice drought

tolerance and yield.

Results

***OsLG3* is associated with drought stress tolerance in rice.**

In our previous cDNA microarray experiment, comparison between upland rice (UR)

(IRAT109 and Haogelao, drought-resistant *japonica* rice) and lowland rice (LR)

(Nipponbare and Yuefu, drought-sensitive *japonica* rice) varieties under well-watered and

water deficit conditions showed that *OsLG3* expression was induced to a significantly

greater extent by water deficit stress in UR than LR [52]. To confirm this we analyzed the

expression level of *OsLG3* between IRAT109 and Nipponbare under increasingly severe

water deficit conditions using Quantitative real-time PCR (qRT-PCR). The data confirmed

that *OsLG3* is more highly expressed in IRAT109 than Nipponbare under well-watered

conditions and that *OsLG3* expression is strongly induced by drought in IRAT109 but not

Nipponbare (**Figure. 1a**). These results suggested that changes in *OsLG3* expression may

be involved in response to drought stresses.

We conducted a candidate gene association analysis to investigate if natural variation

in *OsLG3* is associated with rice drought tolerance. This approach utilized a mini core

collection (MCC) panel [53] (**Additional file 7: Table S1**) of 173 varieties that have

undergone deep sequencing to a 14.9× average depth (http://www.rmbreeding.cn/Index/).

Seeds from MCC lines were germinated on water or in the presence of 15% polyethylene

glycol (PEG) to simulate drought stress. The relative germination rate (RGR, ratio of

germination rates under 15%PEG condition to germination rates under water condition) of

each line was calculated after five days. We identified significant variation in water deficit

stress tolerance between the different varieties (**Additional file 1: Figure S1** and

Additional file 7: Table S1). A total of 97 SNPs within the *OsLG3* locus from these

accessions were identified. To reduce the incidence of false positives, we performed

general linear model (GLM) that controls population structure (Q matrix) to identify

significant genotypic and phenotypic associations. The association analysis detected three

significant SNPs ($P < 1.0 \times 10^{-3}$) (SNP_4352414, SNP_4352886, SNP_4352960) located

within the promoter region of *OsLG3* (**Figure 1b**). SNP_4352886, located 2449 bp

upstream from the start codon of *OsLG3*, showed greatest significant association with RGR

($P = 2.66 \times 10^{-6}$, **Figure 1b**), contributed 13.9% of the phenotypic variation in the MCC

population. SNP_4352886 was in strong LD with two other variations (SNP_4352414,

SNP_4352960) in the promoter ($r^2 \geq 0.8$), but not with the SNP_4348903, SNP_4352166,

SNP_4352793, SNP_4353076 and SNP_4353119, which were identified as marginally

significant ($P < 1.0 \times 10^{-2}$) (**Figure 1b**). SNPs identified within the coding region of

OsLG3 were not significantly associated with the RGR trait. Based on results above, we

conclude that the nucleotide polymorphisms in the promoter of *OsLG3* are associated with

differential germination rates under water deficit.

OsLG3 is an ERF family transcription activator that functions as a homodimer

OsLG3 encodes a putative protein with 334 amino acids. Amino acids 110-159 contain a

typical AP2 domain, including 11 putative DNA-binding sites, implying a strong DNA

binding capacity, and one putative nuclear localization signal (NLS) from amino acids

95-121 (**Additional file 2: Figure S2a, b**). Phylogenetic analysis comparing *OsLG3* with

known ERF TFs [23] indicated that OsLG3 belongs to group Ⅶ of the ERF subfamily and

is closely related to OsERF71 [27, 54], OsEREBP1 [25] and OsBIERF1[55], which have

been reported to be involved in stress response (**Figure 2**) .

Transactivation activity assays, in which the DNA-binding domain (GAL4-BD) of

GAL4 was fused to either the full length CDS of OsLG3 or a series of shortened fragments

created by deletions from both the N- and C-termini, indicated that OsLG3 is capable of

transcriptional activation, and that the C-terminal region (amino acids 213-334) is required

for this (**Figure 3a**). We also tested dimerization of OsLG3 protein *in vivo* using a yeast

two-hybrid system. To avoid the interference caused by self-transactivation activity, we

used the protein fragment BD-dC2 (amino acids 1-218) as bait, which lacks the

transactivation region. The constructs AD-OsLG3 and BD-dC2 were co-transformed into

yeast strain AH109 to test for a homodimer interaction. AD and BD, AD-OsLG3 and BD

were also co-transformed as negative controls for endogenous transactivation activity. Of

these combinations, only AD-OsLG3 and BD-dC2 co-transformants could grow well in the

SD/-Trp-Leu-Ade-His/X-α-gal medium (**Figure 3b**). These results indicate that OsLG3

potentially functions as a homodimer in rice. To determine the subcellular localization of

OsLG3, *Nicotiana benthamiana* leaves were infiltrated with *Agrobacterium tumefaciens*

(strain EH105) containing 35S:OsLG3-GFP and 35S::GFP, respectively. Confocal imaging

analyses showed nuclear-localized fluorescence in 35S:OsLG3-GFP transformed cells,

whilst fluorescence from free GFP was distributed throughout the whole cell (**Figure 3c**).

This indicates that OsLG3 is a nuclear-localized protein.

Expression profile of *OsLG3* under different stress treatments and in different plant

tissues

The expression profile of *OsLG3* in response to abiotic stresses and hormones in

IRAT109 was investigated using qRT-PCR. The transcript level of *OsLG3* was

significantly induced under dehydration, polyethylene glycol (PEG), hydrogen peroxide

(H_2O_2), NaCl, abscisic acid (ABA), ethylene (ETH) and gibberellin (GA) treatments but

remained unchanged under cold treatment (**Figure 4a**). To determine the spatial-temporal

expression of *OsLG3* under normal growth conditions, we isolated total RNA in eight

representative tissues (root, stem, sheath at seedling stage and root, stem, sheath, leaf,

panicle at reproductive stage) from IRAT109, and performed qRT-PCR analysis. *OsLG3*

was expressed in all of the tissues tested, and showed higher level in roots compared to

other tissues (**Figure 4b**).

Overexpression of *OsLG3* enhances drought stress tolerance in rice

To elucidate the biological functions of *OsLG3* during the stress response, 10

independent transgenic lines in which *OsLG3* was overexpressed under the control of 35S

promoter (*OsLG3*-OE) and 14 independent transgenic rice lines in which endogenous

OsLG3 expression level was suppressed by RNA interference (*OsLG3*-RNAi) were

generated. qRT-PCR was carried out to quantify *OsLG3* expression level in each transgenic

line. Two independent *OsLG3*-OE transgenic lines (OE4, OE7) with highest expression

level (**Additional file 3: Figure S3**) and two *OsLG3*-RNAi lines (RI6, RI10) with lowest

expression level (**Additional file 4: Figure S4**) of *OsLG3* were selected for further stress

analysis.

Under dehydration treatment using 20%PEG for 3 days, *OsLG3*-OE lines showed

greater resistance than WT plants (**Figure 5a**). Almost 44-81% of *OsLG3*-OE plants

survived, while only 8-15% of the WT plants survived under this treatment (**Figure 5c**). In

contrast, when four weeks old WT and *OsLG3*-RNAi plants were stressed by a slightly less

severe dehydration stress (20% PEG for 2.5 days) the relative survival of *OsLG3*-RNAi

lines (3–11%) was lower than that of WT (24-42 %) (**Figure 5b, d**).

Under severe drought soil drought stress condition (stop watering for 7 days), the

OsLG3-OE lines showed an improved survival rate compared to WT plants. Almost 48-64%

of *OsLG3*-OE survived, whereas only 17-28% of WT plants survived this treatment

(**Figure 5e, g**). In contrast, when WT and *OsLG3*-RNAi plants were treated moderate

drought (stop watering for 6 days), 36-47% of the WT plants had recovered 10 days after

watering was restored, but only 8-28% of the RNAi plants recovered (**Figure 5f, h**).

Interestingly, we found that the leaves of *OsLG3*-OE plants showed a slower rate of

water loss compared to those of WT and *OsLG3*-RNAi lines under dehydration condition

(**Additional file 5: Figure S5a**), indicating a role for *OsLG3* in reducing water loss,

especially under water deficit conditions. Expression levels of some characterized stress

response genes like *OsLEA3, OsAP37, SNAC1, RAB16C, RAB21* and *OsbZIP73* were

monitored in the WT, *OsLG3*-OE and RNAi lines under well-watered and soil drought

conditions. As shown in **Additional file 5: Figure S5b**, these genes showed significantly

higher levels under soil drought stress in *OsLG3-OE* plants when compared to WT and

RNAi lines.

We also analyzed the growth of *OsLG3*-OE and -RNAi plants under NaCl and

mannitol treatment to induce high salinity and osmotic stress, respectively. Under NaCl

treatment, OE lines showed significantly less suppression of relative shoot growth (shoot

length under stress conditions to shoot length under normal conditions) than both WT and

RNAi lines, and less suppression of relative fresh weight (fresh weight under stress

conditions to fresh weight under normal conditions) than WT and RNAi lines (**Figure 6a,**

c). Under mannitol treatment, the relative shoot growth of OE plants was significantly

higher than that of both WT and RNAi lines, and the relative fresh weight in OE lines was

higher than that of WT and RNAi lines (**Figure 6b, d**). These results show that changes in

the expression level of *OsLG3* have a significant effect on the drought tolerance in rice.

Global gene expression analysis revealed alteration in the expression of stress-related

genes and ROS-scavenging related genes.

Preliminary evidence for altered expression of stress-related genes in *OsLG3* -OE and

RNAi lines led us to investigate the pathways regulated by *OsLG3* through transcriptome

analysis. RNA samples were isolated from the leaves of ten-day-old seedlings of WT,

OsLG3-OE and RNAi plants and Digital Gene Expression (DGE) analysis was performed.

Compared to WT, 223 transcripts were found to have a greater than 2-fold change in

abundance ($P < 0.05$, FDR < 0.05) (**Figure 7a**). 159 genes were up-regulated in OE plants

and down-regulated in RNAi plants, while 64 genes were down-regulated in OE plants and

up-regulated in RNAi plants, respectively (**Figure 7b**). The expression of several of these

genes was tested independent by qRT-PCR to validate the DGE results. Of the genes

checked, six out of eight in OE plants (OE vs WT > 2) and five out of eight in RNAi plants

(RNAi vs WT < 0.5) showed expression patterns consistent with the DGE results

(**Additional file 6: Figure S6**). Gene Ontology (GO) analysis showed that the 218

differentially expressed genes affected by the *OsLG3* overexpression and suppression were

significantly enriched in three GO terms (hyper geometric test, $P < 0.01$, FDR<0.05),

including response to stress (GO: 0006950), response to stimulus (GO: 0050896), and

response to abiotic stimulus (GO: 0009628) (**Figure 7c**; **Additional file 3: Figure S3**).

These results are consistent with our proposed role for *OsLG3* in the regulation of drought

stress tolerance.

Interestingly, DGE analysis showed that ten ROS-scavenging related genes (*APX1*,

APX2, *APX4*, *APX6*, *APX8*, *CATB*, *POD1*, *POD2*, *SODcc1*, *FeSOD*) were up-regulated in

OE line and down-regulated in RNAi line. This result is consistent for a role of *OsLG3* in

the control of ROS homeostasis. To confirm this regulation, we analyzed the expression

levels of 15 genes by qRT-PCR (nine genes identified in the DGE analysis and six other

ROS-related genes) in WT, *OsLG3*-OE and -RNAi plants under well-watered and drought

conditions. 13 of the 15 tested genes were significantly up-regulated under drought stress

conditions, and the expression level of 13 genes (except *APX3* and *POD2*) were

significantly higher in OE plants than in WT and RNAi plants under drought stress

conditions (**Figure 7d**). Conversely, the expression of *APX1*, *APX2*, *POX8*, *POX22.3* and

POD1 was significantly lower in the *OsLG3*-RNAi plants compared to WT under drought

stress conditions, while the remaining nine genes did not show a significant difference in

expression between WT and RNAi line (**Figure 7d**).

$OsLG3$ participates in H_2O_2 homeostasis

The potential role of *OsLG3* in oxidative stress tolerance was examined further by using

two oxidative stress inducers, methyl viologen (MV) [56] and H_2O_2. Germinated WT and

OsLG3-OE and RNAi plants were sown on $^1/_2$ MS medium and $^1/_2$ MS medium containing

2 μM MV. Application of MV dramatically repressed seedling growth in all plants, but the

OsLG3-OE lines exhibited less growth inhibition compared to WT, while the RNAi plants

showed more severe growth inhibition than WT (**Figure 8a, b and c**). Two-week-old

seedlings were treated with 1 mM H_2O_2 or 3 μM MV for 24 hours, followed by 3,

3'-diaminobenzidine (DAB) staining to show the presence of H_2O_2 and nitroblue

tetrazolium (NBT) staining to show the presence of superoxide anion. Under control

conditions, WT or transgenic plants showed similar basal levels of H_2O_2 and superoxide,

but DAB staining and NBT staining were much stronger in the WT than in the *OsLG3*-OE

plants under H_2O_2 and MV stress treatment (**Figure 8d**). These results indicated that

overexpression of *OsLG3* in rice can enhance tolerance to oxidative stress.

Based on the above results, it is hypothesized that *OsLG3* may be involved in the

regulation of ROS homeostasis, leading to enhanced adaption to drought. To confirm this,

we stressed the WT, *OsLG3*-OE and -RNAi plants with dehydration and 20% PEG,

followed by DAB staining to detect H_2O_2 accumulation. Under well-watered conditions,

there was low H_2O_2 accumulation level in WT, *OsLG3*-OE and RNAi plants (**Figure 8e**).

However, *OsLG3*-OE leaves showed significantly less accumulation of H_2O_2 than WT

plants under drought stress conditions, whereas RNAi lines shows more H_2O_2 than WT

under drought stress. H_2O_2 accumulation under normal and drought stress conditions was

also quantified, supporting the DAB staining results. Less H_2O_2 was detected in the leaves

of *OsLG3*-OE plants than WT and *OsLG3*-RNAi plants under drought stress conditions

(**Figure 8f**). The accumulation of ROS may lead to oxidative damage, as indicated by

membrane lipid peroxidation. Monodehydroascorbate (MDA) is a biomarker for membrane

lipid peroxidation [57-59]. MDA production under normal growth conditions was similar in

WT and all transgenic plants, whereas under drought stress, MDA production was

significantly lower in the *OsLG3*-OE compared with WT and *OsLG3*-RNAi plants (**Figure**

8g). These results demonstrate that overexpression of *OsLG3* can reduce the damage to

membranes caused by drought stress.

The effect of *OsLG3*-OE on oxidative stress tolerance implies *OsLG3* may influence

ROS homeostasis. It is well known that superoxide dismutase (SOD), and peroxidase (POD)

are the major ROS-scavenging enzymes in plants under stressed conditions [58].

Four-week-old seedlings were treated with drought stress for 5 days and the activity of

SOD and POD was determined. Results suggested that under normal growth conditions,

OsLG3-OE lines have significant higher SOD than WT and RNAi plants (**Figure 8h**),

while the activity of POD did not appear to be significantly affected in *OsLG3*-OE or RNAi

plants (**Figure 8i**). Under drought stress conditions, the activities of POD and SOD were

both significantly enhanced in OE plants and significantly reduced in RNAi plants

compared to WT (**Figure 8h, i**). These results imply that the function of *OsLG3* in drought

tolerance may be associated with an enhanced antioxidant response to counteract oxidative

stress under drought.

The elite *OsLG3* allele can be used for improving *japonica* rice drought tolerance

To analyze the relationship between *OsLG3* haplotypes and drought tolerance, we

investigated the phylogenetic relationship of 1058 deep-sequenced (depth≈14.9x) rice

accessions, including 251 UR, 415 LR and 392 wild rice varieties originating from a wide

geographic range (**Additional file 10: Data Set 1**). Forty-five SNPs were identified by

minor allele frequency (MAF) > 0.05 and missing rates□≤□50%. Phylogenetic analysis

based on these variations showed that there is a clear differentiation between *japonica*-UR

and *japonica*-LR rice (**Figure 9a**). To get further insight into the phylogenetic relationship,

ten elite *japonica*_UR varieties (IRAT109, IAC150/76, IRAT266, Guangkexiangnuo,

Shanjiugu, Taitung_upland328, Jaeraeryukdo, Riku aikoku, Padi darawal, Malandi 2)

were chosen as an UR pool based on their strong drought resistance, and ten typical

japonica-LR varieties (Nipponbare, Yuefu, Guichao2, Koshihikari, Zhonghua11,

Early_chongjin, Xiushui115, IR24, Ningjing3, Xiushui114) were chosen as a LR pool.

Seven SNPs (SNP_4352414, SNP_4352886, SNP_4352960, SNP_4352792,

SNP_4352797, SNP_4353103 and SNP_4353347) that differed between the upland rice

pool and lowland rice pool were identified. These seven SNPS showed a clear

phylogenetic distinction between *japonica*-UR and *japonica*-LR (**Figure 9b**). On the basis

of these seven SNPs, we could divide the sequences of the 1058 cultivated varieties into

nine haplotypes of which there were four main variants, Hap1, Hap2, Hap3, and Hap4

(**Figure 9c, d**). *OsLG3*$^{\text{Nipponbare}}$ is representative of Hap1, which is mainly composed of

japonica-LR rice, whereas *OsLG3*$^{\text{IRAT109}}$ belongs to Hap2, which is mainly composed of

japonica-UR rice and is the second largest group. Hap3 is mainly composed of *indica* rice,

and Hap4 is mainly composed of wild rice. The results indicated that *OsLG3* has clear

differentiation in *japonica* rice, which can be divided into *japonica*-UR and *japonica*-LR

rice, while no clear division in *indica* rice. Therefore, we designated Hap1 and Hap2 as

the sensitive and tolerant alleles, respectively, of *OsLG3* in *japonica* rice.

To confirm the genetic effect of different alleles of *OsLG3* on rice drought tolerance,

two introgression lines (IL342 and IL381) were selected from a cross between IRAT109

(donor parent) and Yuefu (receptor parent). The *OsLG3* allele in these two lines shares

sequence similarity with IRAT109, with seven matching SNPs in the promoter (including

the three significant loci describe in candidate gene association analysis section), whereas

Yuefu carries the same *OsLG3* allele as Nipponbare (**Figure 9e**). Thus, the *OsLG3* allele

found in IRAT109, IL342 and IL381 was considered to be the drought-tolerant allele and

the allele in Yuefu the drought-sensitive allele, respectively. When germinated on 15%

PEG conditions, the RGR of Yuefu was 27.0% compared to germination on water, while

the RGR of   IL342, IL381 and IRAT109 was 35%, 36% and 46%, respectively (**Figure**

9f). These results support the hypothesis that the nucleotide polymorphisms in *OsLG3*

contribute to enhanced rice drought tolerance.

According to the pedigree records, Huhan3, one of green super rice accessions, is an

elite upland rice that is widely grown in Hubei province of China because of its high yield

and outstanding ability to conserve water. One of its parents is IRAT109. A re-sequencing

study showed that Huhan3 carries the allele of *OsLG3* deriving from IRAT109 (**Figure**

9g). Moreover, Zhenghan9, another elite upland rice, is derived from the cross of

IRAT109 and Yuefu. Zhenghan9 retained the favorable allele of *OsLG3* and an allele for

an unidentified grain quality-related gene derived from Yuefu (**Figure 9h**). These

observations illustrate the successful outcome of combining the *OsLG3*^{IRAT109} allele and

those alleles for other unidentified grain quality- related genes, for improving drought

tolerance and grain appearance quality, which had been employed by breeders.

Discussion

Natural variation in the promoter of *OsLG3* is associated with drought tolerance in

rice

Despite many studies investigating the transcriptional response to drought stress [5, 60-62],

how natural sequence variation is associated with phenotypic variations in drought

tolerance remains largely unknown. Because of the polygenic inheritance, low heritability,

and strong genotype-by-environment interactions of drought resistance-related traits, there

are few reports on the cloning and identification of drought resistant genes by association

analysis and positive mutant screening methods [48, 49, 63]. In this study, using candidate

gene association analysis, we detected three significant SNPs ($P < 1.0 \times 10^{-3}$)

(SNP_4352414, SNP_4352886, SNP_4352960) located within the promoter region of

OsLG3 were significantly associated with the RGR trait (**Figure 1c**). From our previous

work, transient assays showed that four out of seven SNPs are conserved and play a

significant role in regulating *OsLG3* expression [51]. In this study, transgenic experiments

further support the hypothesis that increased expression of *OsLG3* enhances rice drought

stress tolerance.

To analyze the relationship between *OsLG3* haplotypes and drought tolerance, we

investigated the phylogenetic relationship of 1058 deep-sequenced rice accessions. Based

on the seven major SNPs, we divide the sequences of the 1058 cultivated varieties into nine

haplotypes of which there were four main variants, Hap1 (mainly composed of

japonica-LR rice), Hap2 mainly composed of *japonica*-UR rice, Hap3 (mainly composed

of *indica* rice), and Hap4 (mainly composed of wild rice). The variations of *OsLG3* in these

accessions showed that there is a clear differentiation between *japonica*-UR and

japonica-LR rice, while no clear division in *indica* and wild rice (**Figure 9**). These results

implicated the Hap2 variant of *OsLG3* may improve drought tolerance of cultivated

japonica rice. Re-analysis of two introgression lines (IL342 and IL381) derived from a

cross between IRAT109 (donor parent) and Yuefu (receptor parent) indicated that selection

of Hap2 was effective in the improvement of drought tolerance (**Figure 9e**). Interestingly,

our previous work indicated that *OsLG3* acts as an important positive regulator of grain

length and could improve rice yield [51]. In fact, the pedigree records in upland rice

breeding showed that the tolerant allele of *OsLG3* had been incorporated into elite varieties

via breeding (**Figure 9g**). We propose that pyramiding the beneficial allele of *OsLG3* with

other yield- and quality-related genes could contribute to the breeding of elite rice varieties

because of its pleiotropic effects on traits.

***OsLG3* was cloned from upland rice and plays a positive role in rice drought**

tolerance.

UR has evolved enhanced drought-resistance compared to LR, derived from natural and

artificial selection over time under drought conditions. UR performs better under drought

conditions, with stronger water-retention ability, larger root volumes and higher biomass

production [64, 65]. Therefore, UR is highly suitable to be taken up as research material

regarding mechanisms underlying drought resistance. From the previous work of

expression profiles from typical LR (Nipponbare and Yuefu, drought-sensitive *japonica*

rice) and UR varieties (IRAT109 and Haogelao, drought-resistant *japonica* rice) under

water deficit stress conditions using cDNA microarray [52], we found the transcription of

OsLG3 can be induced to a greater extent in the UR varieties compared to the LR varieties

during drought stress. When IRAT109 and Nipponbare were subjected to soil drought stress,

expression of *OsLG3* in Nipponbare was not induced, whereas in IRAT109 *OsLG3* shows

significant induction under both slight drought (SLD) and moderate drought (MOD)

conditions (**Figure 1a**). Using candidate gene association analysis, we found that

nucleotide polymorphisms in the promoter region of *OsLG3* are associated with different

levels of water deficit tolerance among rice varieties at the germination stage (**Figure 1b**).

All these results indicated that *OsLG3* might play a role in the observed drought stress

response of upland rice.

To assess the effect of *OsLG3* on water deficit stress responses, we tested the growth

response of *OsLG3*-OE and -RNAi plants under simulated drought stresses. *OsLG3*-OE

lines showed higher survival under dehydration caused by 20% PEG6000 and soil drought

stress caused by stopping watering (**Figure 5**). Plants with reduced *OsLG3* expression

showed reduced survival, suggesting that this gene might be involved in response to

drought stress. A slower rate of water loss was observed in the leaves of *OsLG3*-OE

compared to WT and RNAi plants (**Additional file 5: Figure S5a**), suggesting that *OsLG3*

plays an important role in reducing water evaporation. Collectively, these results

demonstrate that *OsLG3* is a positive regulator of drought stress response in rice.

OsLG3-OE plants also showed enhanced growth compared to WT and RNAi lines under

mannitol and NaCl treatments, which suggests that *OsLG3* may be involved in osmotic and

high salinity stress tolerance through cross-talk between water deficit, osmotic and high

salinity stress response pathways.

Potential mechanisms of *OsLG3* in drought stress tolerance

We found that the expression levels of a set of stress-related genes like *OsLEA3*, *OsAP37*,

SNAC1 are increased in *OsLG3*-OE lines compared to WT, and decreased in *OsLG3*-RNAi

lines before and after drought treatment (**Additional file 5: Figure S5b**). This was

confirmed by DGE analysis (**Additional file 9: Table S3**). We investigated if the altered

response to drought was associated with global changes in the expression of stress-related

genes before stress application. Analyses indicated that many stress-related genes were

up-regulated in the OE line and down-regulated in the RNAi line (**Figure 7c**), which is

consistent with the observed phenotypes of *OsLG3*-OE and -RNAi lines. Interestingly, the

expression level of some ROS-scavenging related genes showed increased expression

levels in OE and decreased abundance in RNAi lines too. For instance, the ascorbate

peroxidase gene, *OsAPX1* plays a positive role in chilling tolerance by enhancing

H_2O_2-scavenging [66]. *DSM1* mediates drought resistance through ROS scavenging in rice

[12]. *OsCATB* prevent the excessive accumulation of H_2O_2 under water stress [67]. Thus,

these DGE analyses are consistent with our hypothesis that the function of *OsLG3* in

abiotic stress tolerance occurs through the transcriptional regulation of stress-related and

ROS scavenging-related genes.

Previous studies have demonstrated that plant response to abiotic stresses is mediated

through the regulation of ROS metabolism [9, 10, 12, 68, 69]. For example, overexpression

of a NAC gene, *SNAC3*, increased drought and heat tolerance by modulating ROS

homeostasis through regulating the expression of genes for ROS-scavenging and

production enzymes [9]. Overexpression of an ERF gene, *SERF1*, improved salinity

tolerance mainly due to the regulation of ROS-dependent signaling during the initial phase

of salt stress in rice [10]. The present data revealed overexpression of *OsLG3* led to the

up-regulation of many ROS scavenge-associated genes, while suppression of *OsLG3*

caused the down-regulation of these genes (**Figure 7d**). Furthermore, the H_2O_2 and MDA

content which accumulated in the leaves of *OsLG3*-OE plants was significantly lower than

that in the WT and *OsLG3*-RNAi plants (**Figure 8c, d**), suggesting that the improved

drought tolerance of *OsLG3*-OE plants may be due to the efficient scavenging of ROS.

Relevant to this, *OsLG3*-OE seedlings exhibited better growth under oxidative stress

caused by application of MV. In contrast, RNAi plants showed enhanced sensitivity to

oxidative stress. Therefore, it is proposed that the function of *OsLG3* in drought tolerance is

associated with the enhancing activity of ROS-scavenging.

To scavenge or detoxify excess stress-induced ROS, plants have developed a complex

antioxidant system comprising of the non-enzymatic as well as enzymatic antioxidants [7,

70]. The POD and SOD activities were found to be higher in the *OsLG3*-OE lines as

compared to WT under drought stress, and *OsLG3*-RNAi lines showed the reverse results

(**Figure 8h**). These data suggest that *OsLG3*-OE plants have enhanced activity of

ROS-scavenging enzymes which significantly contributes to the reduction of ROS

accumulation, and thereby improved drought stress tolerance.

In conclusion, here we present evidence that *OsLG3* is induced by water deficit

stresses and its induction is greater in UR IRAT109 compared to LR Nipponbare under

drought. Nucleotide polymorphisms in the promoter region of *OsLG3* are associated with

water deficit tolerance in germinating rice. Transgenic plants overexpressing *OsLG3*

showed improved growth under drought stress, probably via inducing ROS scavenging by

controlling downstream ROS-related genes. Phylogenetic analysis indicated that the

tolerant allele of *OsLG3*, identified in drought-tolerant *japonica* rice varieties, could be

introduced to rice to improve the drought tolerance. The pedigree records in upland rice

breeding showed that the tolerant allele of *OsLG3* had been incorporated into elite varieties

via breeding. Importantly, the tolerant allele of *OsLG3* is a promising genetic resource for

the development of drought-tolerant and high yield rice varieties using traditional breeding

472 approaches or genetic engineering.

473 **Methods**

474 **Plant materials and stress treatments**

475 Rice varieties IRAT109 and Nipponbare were used for quantitative real-time PCR

476 (qRT-PCR) analysis of *OsLG3* transcript levels under various stresses and hormone

477 treatments, and Nipponbare was used for all transgenic experiments. For qRT-PCR analysis

478 the expression level of *OsLG3* under simulated drought conditions, the seeds of

479 Nipponbare and IRAT109 were sown in flower pots (140 mm diameter×160 mm deep)

480 with well-mixed soil (forest soil: vermiculite in a ratio of 1:1), and grown in the greenhouse

481 under well water conditions at 28°C/ 26°C and 12 hours light/12 hours dark photoperiod.

482 Three-week old plants were subjected to drought treatment with stop watering for 0, 5, 6

483 and 7 days. From these, four stress treatment levels were categorized as no stress (NS),

484 slight drought (SLD), moderate drought (MOD) and severe drought (SED), respectively.

485 For qRT-PCR analysis the expression level of *OsLG3* under various abiotic stresses and

486 phytohormone treatments, three-week-old IRAT109 seedlings grown in Hoagland solution

487 (PPFD of 400 µmol/m^{2}s^2, 12 hours light (28°C)/12 hours dark (26°C)) were subjected to

488 different treatments with 20% (w/v) polyethylene glycol (PEG) 6000, NaCl (200 mM), cold

489 (4°C), hydrogen peroxide (H$_2$O$_2$,1 mM), abscisic acid (ABA,100 µM), ethylene (ETH,100

490 µM), GA (100 µM) and dehydration by exposure in air. Leaf tissue was harvested at 0, 1, 2,

491 4, 6, 9, 12 and 24 hours after PEG, NaCl, cold, H$_2$O$_2$, ABA, ETH, and GA treatments, and

492 at 0, 1, 2, 3, 4, 5, 6, 7 and 8 hours after dehydration treatment.

493  The seeds of WT, *OsLG3*-OE lines (OE4, OE7) and *OsLG3*-RNAi lines (RI6, RI10)

were germinated in $^1/_2$ MS medium for various stress evaluations. For dehydration

treatment, uniformly germinated seeds were transplanted into bottom removed 96-well

PCR plates, and grown hydroponically using Hoagland solution at 28°C/26°C (day/night)

with a 12 hours photoperiod. Three-week-old plants were treated with 20% (w/v) PEG6000

solution for about three days, and recovered with water for ten days. Each stress test was

repeated three times. For drought stress treatment, three-day-old WT and transgenic

seedlings were transplanted to well-mixed soil (forest soil: vermiculite in a ratio of 1:1),

and grown for four weeks under a normal watering regime. Drought stress treatment was

then applied by stopping irrigation for about seven days. After all leaves had completely

rolled, watering was resumed for 10 days. The survival ratio of each line was calculated as

the number of surviving plants over the number of treated plants in each flowerpot. To

evaluate the tolerance of rice seedlings to osmotic, high salinity and oxidative stress

treatment, three-day-old WT, *OsLG3*-OE and *OsLG3*-RNAi seedlings (10 plants per

replicate, three replicates) were transplanted to $^1/_2$ MS medium (mock treatment) or $^1/_2$ MS

medium containing 200 mM mannitol, 150 mM NaCl or 2 μM MV respectively. Plants

were grown for seven days under 12 hours light (28°C)/12 hours dark (26°C) photoperiod

conditions and then the shoot height and fresh weight of all plants was measured.

Gene expression analysis

Total RNA was extracted using RNAiso Plus (TaKaRa) according to the manufacturer's

instructions. 4 μg of the DNase-treated RNA were reverse transcribed by using M-MLV

reverse transcriptase (TaKaRa). qRT-PCR was performed in an Applied Biosystems 7500

Real Time PCR system (ABI, USA) using SYBR Premix Ex TaqTM II (TaKaRa) according

to the protocol previously described [71, 72]. The gene-specific primers used for qRT-PCR

are listed in **Supplemental Table S2**. Rice *Actin1* gene was used as the internal reference

gene for data normalization [73].

***OsLG3*-gene association analysis of rice drought tolerance among 173 rice genotypes**

173 cultivated varieties from the mini core collection of Chinese cultivated rice (Zhang et

al. 2010), including 130 *indica* and 43 *japonica* rice varieties (**Supplemental Table S1**)

were selected for the candidate gene association mapping. To perform the analysis, we

obtained phenotypic data of relative germination rates (RGR, ratio of germination rates

under stress conditions to germination rates under water conditions) from growth in water

and 15% PEG6000 treatment. Briefly, 50 seeds of each line were placed in petri dishes (90

mm diam) lined with filter paper. 10 ml water or 15% PEG6000 was added to each petri

dish as mock treatment or to induce osmotic stress, respectively. All petri dishes were

placed in a 28°C greenhouse and the germination rates were assessed after five days.

Genotype data for each line was acquired from 'The 3,000 rice genomes project' [74]. The

SNP data were filtered out. After filtering, a total of 97 SNPs remained in a 5-kb region

surrounding the *OsLG3* gene. Association analysis using a general linear model with the

population structure (Q matrix) method was conducted using TASSEL 5.2.28 software. The

Q matrix was estimated from the genomic data to control for population structure.

Plasmid construction and rice transformation

To generate the overexpression construct, the full-length cDNA of *OsLG3* was amplified

from the first-strand cDNA of IRAT109 with specific primers (**Supplemental Table S2**)

using PrimeSTAR® HS DNA Polymerase (TaKaRa), digested with *Kpn*I and *Pac*I, and

cloned into binary vector pMDC32 [75]. For construction of RNA interference (RNAi)

plasmid, a 415 bp fragment with low similarity to other rice genes located at the 5' end of

OsLG3 was amplified by PCR with specific primer, digested with *Kpn*I and *Bam*HI and

then *Spe*I and *Sac*I sites, and cloned into the pTCK303 vector as described previously [76].

Subcellular localization

For subcellular localization analysis, the full-length open reading frame (ORF) of *OsLG3*

without terminal codon was amplified, and the amplified fragments were digested with

*Kpn*I and *Pac*I, and then cloned in to pMDC83 vector, fused with the GFP reporter gene

driven by *CaMV 35S* promoter. The fusion construct (35S:OsLG3-GFP) and control

(35S:GFP) were transformed into *Agrobacterium tumefaciens* strains (EH105), and then

infiltrated to five-week-old *Nicotiana benthamiana* leaves [77]. The fluorescence signal

was examined through a confocal laser scanning microscope OLYMPUS FV1000 with

excitation at 488 nm and emission at 525 nm.

Transactivation and dimerization assay

For transactivation assay, The full length coding region and truncated fragments of *OsLG3*

generated by PCR amplification were fused in frame to the GAL4 DNA binding domain in

the vector of pGBKT7 (Invitrogen). The plasmids of FL (the full length coding region of

OsLG3, amino acids 1-334), dC1 (amino acids 1-218), dC2 (amino acids 1-167), dC3

(amino acids 1-106), dC4 (amino acids 107-334), dC5 (amino acids 168-334) and dC6

(amino acids 213-334) were constructed. These constructs were introduced into yeast strain

AH109 by LiAc-mediated yeast transformations, and screened on the selective medium

plates without Tryptophan (SD/-Trp). The PCR-verified transformants were transferred to

SD medium without Tryptophan/Histidine/Adenine (SD/-Trp/-His/-Ade) for 3 days. The

β-galactosidase activity was performed according to the *in vivo* agar plate assay (X-α-gal in

medium).

For dimerization assay, AD-OsLG3 with full length coding region of *OsLG3* fused in

frame to the GAL4 DNA activating domain in the vector of pGADT7 (Invitrogen) was

constructed. The constructs AD-OsLG3 and BD-dC2 were co-transformed into yeast strain

AH109. AD and BD empty, AD-OsLG3 and BD empty were also co-transformed as

negative controls. All transformed cells were screened on the selective medium plates

without Tryptophan and Leucine (SD/-Trp-Leu), The PCR-verified transformants were

transferred   to   SD   medium   with   X-α-gal   and   without

Tryptophan/Histidine/Adenine/Leucine (SD/-Trp-Leu-Ade-His/X-α-gal) for 3 days.

Physiological measurements

The rate of water loss under dehydration condition was measured as described previously

[72]. Ten plants of each line were used in each replicate, and three replicates were made for

each line. Histochemical assays for ROS accumulation were determined according to the

previously described method [68, 78]. Briefly, the qualitative detection of H_2O_2

accumulation was detected by DAB staining. Excised leaves were treated with DAB

staining solution (1mg/mL DAB pH 3.8) at 28°C for 12 hours in the dark. After staining,

the leaves were decolorized with Acetic acid/Ethanol (1:3) for 60 min, and rehydrated in 70%

(v/v) alcohol for 24 hours at 28°C. Each experiment was repeated on at least 10 different

plants, and representative images were shown. Superoxide anion radical accumulation was

detected by NBT staining as described previously. The leaf samples were excised and

immediately placed in 50 mM sodium phosphate buffer (pH 7.5) containing 6 mM NBT, at

28°C for 8 h in the dark. The quantitative measurement of H_2O_2 concentrations was

performed with an Amplex Red Hydrogen / Peroxidase Assay Kit (Molecular Probes)

(Invitrogen, http://www.invitrogen.com) as described by the manufacturer's instructions.

Briefly, leaf samples from both the well water and drought stressed (without water for 5

days) plants were ground in liquid nitrogen, and 100 mg of ground frozen tissue from each

sample was placed in a 2 mL Eppendorf tube and kept frozen. 1 mL precooled sodium

phosphate buffer (20 mM, pH 6.5) was immediately added into the tube and mixed. After

centrifugation (10,000 g, 4°C, 10 min) the supernatant was used for the assay.

Measurements were performed using a 96-well microplate reader (PowerWave XS2,

BioTek) at an absorbance of 560 nm. MDA content was measured as described previously

[72].

The activity of antioxidant enzymes including superoxide dismutase (SOD) and

peroxidase (POD) were measured following the protocols described previously [71]. The

units of the antioxidant enzyme activities were defined as follows: One unit of the SOD

activity was defined as the quantity of enzyme required to cause 50% inhibition of the

photochemical reduction of NBT per minute at 560 nm. One unit of POD activity was

defined as the amount enzyme required to cause a 0.01 absorbance increase per minute at

470 nm.

Digital gene expression (DGE) profiling and gene ontology (GO) analysis

WT Nipponbare and transgenic lines OE7 and RI10 were used for digital gene expression

(DGE) profiling analysis. Ten-day-old seedlings grown on $^1/_2$ MS medium were harvested

for total RNA extraction as described in qRT-PCR section. DGE was performed at the

Beijing Genomics Institute (http://www.genomics.cn) using Illumina Hiseq2000

sequencing technology. Transcripts with significant differential expression between

OsLG3-OE and WT or between *OsLG3*-RNAi and WT plants were identified as those with

$P < 0.05$ using a False Discovery Rate (FDR) of $<$ 0.05 and a fold change cutoff of >2.

Gene Ontology (GO) analysis was performed using agriGO [79]

(http://bioinfo.cau.edu.cn/agriGO/). Representative differentially expressed genes were

confirmed by qRT-PCR. The primers are listed in **Supplemental Table S2**.

Sequence analysis and phylogenetic analysis

The phylogenetic tree was analyzed by MEGA6 software based on neighbor-joining

method and bootstrap analysis (1,000 replicates). The EvolView online tool [80] was used

for visualizing the phylogenetic tree. Multiple sequence alignment was performed with

ClustalW. Gene annotation was conducted in RGAP (http://rice.plantbiology.msu.edu/). All

SNP data were obtained from the Rice functional genomics and breeding database (RFGB,

http://www.rmbreeding.cn/Index/).

Accession numbers

Sequence data from this article can be found in the Rice Genome Annotation Project

Database and Resource (RGAP) (http://rice.plantbiology.msu.edu) under following

accession numbers: *OsLG3* (LOC_Os03g08470); *Actin1* (LOC_Os10g36650); *OsLEA3*

(LOC_Os05g46480); *AP37* (LOC_Os01g58420); *SNAC1* (LOC_Os03g60080); *RAB21*

(LOC_Os11g26790); *RAB16C* (LOC_Os11g26760); *OsbZIP23* (LOC_Os02g52780); *Apx1*

(LOC_Os03g17690); *Apx2* (LOC_Os07g49400); *Apx3* (LOC_Os04g14680); *Apx5*

(LOC_Os12g07830); *Apx6* (LOC_Os12g07820); *Apx8* (LOC_Os02g34810); *OsPox8.1*

(LOC_Os07g48010); *Pox22.3* (LOC_Os07g48020); *FeSOD* (LOC_Os06g05110); *SODcc1*

(LOC_Os03g22810); *SODcc2* (LOC_Os07g46990); *POD1* (LOC_Os01g22370); *POD2*

(LOC_Os03g22010); *CATB* (LOC_Os06g51150); *DSM1* (LOC_Os02g50970); *RAB16D*

(LOC_Os11g26750); *OsNCED4* (LOC_Os07g05940); *OsNCED3* (LOC_Os03g44380);

OsDhn1 (LOC_Os02g44870); *OsSRO1C* (LOC_Os03g12820); *OsMYB48*

(LOC_Os01g74410); *OsITPK4* (LOC_Os02g26720); *OsLEA3-2* (LOC_Os03g20680).

Competing interests

The authors declare that they have no competing interests.

Author's contributions

H. X. and J. Y designed and performed most of the research and wrote the article, X. W.

and P. L. contributed to help with transgenic experiment and stress treatment, J.L., H. Z.

and Y. G. contributed to design some experiments, Y. Z. contributed to help with haplotype

analysis, Y. L and Z. Y. contributed to sample or reagents preparation, B. F., W. W., J. A.

and Z. L. help to analyze the data and revise the manuscript, and Z.L conceived the

research and assisted in writing the manuscript.

Acknowledgments

We thank Amelia Henry (International Rice Research Institute), Steven Burgess (eLIFE)

and Andrew Plackett (University of Cambridge) for critical reading and suggest revisions

to this manuscript.

Figure Legends

Figure 1. *OsLG3* is associated with drought tolerance. (a) qRT-PCR analysis of *OsLG3*

in Nipponbare and IRAT109 under different soil drought stress levels. NS, No stress; SLD,

Slight drought; MOD, Moderate drought; SED, Severe drought. **(b)** Analysis of the

association between pairwise LD of DNA polymorphisms in the *OsLG3* gene and water

deficit tolerance. Schematic of OsLG3 is shown on the X-axis and the significance of each

variation associated with seedling RGR (ratio of germination rates under 15%PEG

condition to germination rates under water conditions) is shown on the Y-axis. The SNPs

with significant variation ($P < 1.0 \times 10^{-2}$) between genotypes are connected to the pairwise

LD diagram with a solid line. Black dots in the pairwise LD diagram highlight the strong

LD of SNP_4352886 (filled red circle, $P = 2.66 \times 10^{-6}$) and two significant variations,

SNP_4352414 and SNP_4352960 (open red circles, $P < 1.0 \times 10^{-3}$). The SNP_4348903,

SNP_4352166, SNP_4352793, SNP_4353076 and SNP_4353119, which are marginally

significant ($P < 1.0 \times 10^{-2}$) are denoted by red triangles.

Figure 2. Phylogenetic tree of OsLG3 homologues. Neighbour-Joining phylogenetic

analysis of OsLG3 protein sequence in the context of other characterized AP2/ERF proteins

from rice. The phylogenetic tree was constructed using the ClustalW and MEGA programs.

Tree topology with bootstrap support is based on a percentage of 1000 replicates.

Accession   numbers   are   as   follows: ARAG1,   LOC_Os02g43970;   OsAP21,

LOC_Os01g10370; OsDREB4-1, LOC_Os02g43940; OsDREB4-2, LOC_Os04g46400;

OsDREB2A,   LOC_Os01g07120;   OsDREB2B,   LOC_Os05g27930;   OsBIERF3,

LOC_Os02g43790; OsERF1, LOC_Os04g46220; OsERF922, LOC_Os01g54890; OsWR1,

LOC_Os02g10760, OsWR4, LOC_Os06g08340; SNORKEL1, AB510478; SNORKEL2,

AB510479; OsBIERF1, LOC_Os09g26420; OsEBP-89, LOC_Os03g08460; OsEREBP1,

LOC_Os02g54160; OsLG3, LOC_Os03g08470; OsERF71, LOC_Os06g09390; Sub1A,

DQ011598; Sub1C, LOC_Os09g11480; Sub1B, LOC_Os09g11460; MFS1,

LOC_Os05g41760; OsAP2-39, LOC_Os04g52090; OsERF3/OsBIERF4/AP37,

LOC_Os01g58420; FZP, LOC_Os07g47330; OsRap2.6, LOC_Os04g32620; OsAP2-1,

LOC_Os11g03540; OsDREB1D, LOC_Os06g06970.

Figure 3. Transactivation assay and Subcellular localization of OsLG3. (a)

Transactivation assay of OsLG3 fragments. Fusion proteins of the GAL4 DNA-binding

domain and different portions of OsLG3 were expressed in yeast strain AH109 and

streaked on control plate (SD/-Trp) or selective plate (SD/-Trp-Ade-His/X-α-gal),

respectively. FL indicates the full-length CDS of OsLG3; dC1 to dC6 indicate the mutated

forms of OsLG3 (nucleotide positions are labeled in the diagrams), respectively. **(b)**

Dimerization analysis. BD with AD empty vector, AD empty with BD-dC2, AD-OsLG3

with BD empty and AD-OsLG3 with BD-dC2, were co-transformed in to yeast strain

AH109 yeast cells and streaked on control plate (SD/-Trp-Leu) or selective plate

(SD/-Trp-Leu-Ade-His/X-α-gal), respectively. The plates were incubated for 3 days. **(c)**

Subcellular localization of OsLG3 in tobacco epidermal cells. The upper panels show the

localization of OsLG3-GFP in onion cells in a transient assay, while bottom panels show

the localization of GFP as a control.

Figure 4. Expression analysis of *OsLG3*. (a) Expression level of *OsLG3* under various

abiotic stresses and hormone treatments in IRAT109. Three-week-old seedlings were

subjected to dehydration, NaCl (200 mmol), PEG 6000 (20%, w/v), cold (4°C), H_2O_2 (1

mM), ABA (100 μM), ETH (100 μM) and GA (100 μM) treatment. The relative expression

level of *OsLG3* was detected by qRT-PCR at the indicated times. Error bars indicated the

SE based on three replicates. **(b)** Detection of *OsLG3* expression in various tissues and

organs of IRAT109 using qRT-PCR. Three-week-old seedlings were used to harvest the

samples of the root, sheath and leaf at seedling stage. Plants before heading stage were used

to harvest the samples of root, stem, sheath, leaf and panicle at the reproductive growth

stage. Error bars indicate the SE based on three technical replicates.

Figure 5. *OsLG3* increases rice survival under severe drought stress. (a) Physiological

dehydration stress tolerance assay; *OsLG3*-OE plants subjected to 20% PEG for 3 days

before being allowed to recover for 10 days. **(b)** Physiological dehydration stress tolerance

assay of *OsLG3*-RNAi plants subjected to 20% PEG for 2.5 days before being allowed to

recover for 10 days. **(c)** and **(d)**, Survival rates of transgenic and WT plants testing in **(a)**

and **(b)**. Values are means ± SE (n = 3). **(e)** *OsLG3*-OE and WT plants were subjected to

severe drought stress without water for 7 days and then recovered for 10 days. **(f)**

OsLG3-RNAi and WT plants were subjected to moderate drought stress without water for 6

days and then recovered for 10 days. **(g)** and **(h)**, Survival rates of transgenic and WT

plants testing in **(c)** and **(d)**. Values are means ± SE (n = 3). * and ** indicate significant

difference at $P < 0.05$ and $P < 0.01$, respectively.

Figure 6. Growth of *OsLG3* transgenic plants under high salinity and osmotic stress

conditions. Ten day old seedlings of WT, *OsLG3*-OE and RNAi plants grown in $^1/_2$ MS

medium containing **(a)** 150 mmol/L NaCl or **(b)** 200 mmol/L mannitol. The relative shoot

length **(c)** and relative fresh weight **(d)** of the transgenic and WT plants from **(a)** and **(b)**

were compared. Values are means ± SE (n = 10). * and ** indicate significant differences at

$P < 0.05$ and $P < 0.01$, respectively.

Figure 7. DGE analysis shows that altering *OsLG3* expression in transgenic plants

affects transcription of stress response genes. (a) Scatterplots comparing the

transcriptome of *OsLG3*-OE and -RNAi with WT. The red and green dots indicate

transcripts from *OsLG3*-OE or -RNAi which have signal ratios compared to WT of greater

than 2 and less than 0.5, respectively. **(b)** Venn diagram showing the numbers of

up-regulated and down-regulated genes affected by the overexpression and suppression of

OsLG3. **(c)** Significantly enriched GO terms show representative biological processes of

up-regulated and down-regulated genes identified in *OsLG3*-OE and RNAi plants,

respectively. **(d)** Transcript levels of genes related to ROS scavenging of WT, *OsLG3*-OE

and RNAi plants under normal or drought stress conditions (withholding water for 5 days).

Values are means ± SE (n = 3).

Figure 8. *OsLG3* is involved in oxidative stress response. (a) Enhanced tolerance of

OsLG3-OE plants and enhanced sensitivity of *OsLG3*-RNAi plants to oxidative stress

caused by methyl viologen (MV). Relative shoot length **(b)** and relative fresh weight **(c)**

measurements of WT, *OsLG3*-OE and *OsLG3*-RNAi seedlings under oxidative stress

treatments. Data are the mean ±SE (n=10). * and ** indicate significant differences at $P <$

0.05 and $P < 0.01$, respectively. **(d)** DAB and NBT staining of leaves for H_2O_2 in WT,

OsLG3-OE and RNAi seedlings under oxidative stress treatments caused by H_2O_2 (100

mM) and MV (30 μM) stress treatment. **(e)** DAB staining of leaves for H_2O_2 from WT,

OsLG3-OE and RNAi seedlings under normal conditions and stress treatment

(three-week-old seedlings were treated with dehydration for 6 hours and 20% PEG6000 for

24 hours). **(f)** H_2O_2 content in leaves from WT, *OsLG3*-OE and RNAi seedlings under

normal conditions and slight drought stress treatment (withholding water for 5 days).

Relative MDA content **(g)**, SOD activity **(h)** and POD activity **(i)** in leaves from WT,

OsLG3-OE and RNAi seedlings under normal and drought conditions (withholding water

for 5 days). Data are the mean ± SE (n=3). * and ** indicate significant differences at *P* <

0.05 and *P* < 0.01, respectively.

Figure 9. The favorable allele of *OsLG3* improves drought tolerance in rice. (a)

Phylogenetic analysis of the *OsLG3* gene based on 45 SNPs in 1058 accessions. **(b)**

Phylogenetic tree of the *OsLG3* in 1058 varieties was constructed based on seven SNPs

(SNP_4352414, SNP_4352792, SNP_4352797, SNP_4352886, SNP_4352960,

SNP_4353103 and SNP_4353347). Different colors reflect the different subgroups. The

pink and green strips represent *indica* and *japonica*, respectively. Both purple strip and red

lines represent wild rice accessions. The blue and gold lines indicate upland rice and

lowland rice, respectively. **(c, d)** Haplotype analysis of *OsLG3*. Natural variation in *OsLG3*

among 1058 rice accessions of a worldwide rice collection. S1-7 denote SNP_4352414,

SNP_4352792, SNP_4352797, SNP_4352886, SNP_4352960, SNP_4353103 and

SNP_4353347, respectively. Hap., haplotype; No., number of cultivated varieties;

*Japonica*_UR, upland rice in *japonica*; *Japonica*_LR, lowland rice in *japonica*; *Indica*_UR,

upland rice in *indica*; *Indica*_LR, lowland rice in *indica*. **(e)** Haplotypes of *OsLG3* in IL342,

IL381, IRAT109 and Yuefu rice genotypes. ATG is the site of the start codon. SNP4 is

incomplete LD with other six polymorphisms in the three drought tolerant lines. These

polymorphisms are shaded in red. **(f)** Relative germination rate of IL342, IL381, IRAT109

and Yuefu rice genotypes. Data represent the mean of triplicates (*$P < 0.05$, ** $P < 0.01$).

(g, h) The pedigree of selected rice varieties. **(g)** Huhan 3 and **(h)** Zhenghan 9. The red star

refers the beneficial allele of *OsLG3*.

Additional files

Additional file 1: Figure S1. Frequency distribution of relative germination rates in the

mini core collection (MCC) population.

Additional file 2: Figure S2. Sequence analysis of OsLG3. **(a)** Schematic diagram of

OsLG3 gene structure. Representation of OsLG3 protein domain structure showing the

location of the AP2 domain and the DNA-binding sites (indicated by shaded triangles). **(b)**

Multiple sequence alignment of OsLG3 with several previously reported stress-related ERF

genes in rice with DNAMAN software. Accession numbers are as follows: OsERF71:

Oryza sativa, LOC_Os06g09390; OsBIERF1: *Oryza sativa*, LOC_Os09g26420; Sub1A:

Oryza sativa, DQ011598; OsEREBP1: *Oryza sativa*, LOC_Os02g54160; AtEBP:

Arabidopsis thaliana, AT3G16770; CaPF1: *Capsicum annuum*, AAP72289; GmEREBP1:

Glycine max, NP_001236578; JERF1: *Solanum lycopersicum*, NP_001234513. JERF3:

Solanum lycopersicum, AAQ91334. Red line, black line and red box represent

MCGGAIL/I motif, nuclear localization signal and conserved AP2 domain, respectively. **(c)**

Sequence logos of the AP2 domain were produced based on the sequences presented in **(b)**.

Height of letter (amino acid) at each position indicates degree of conservation.

Additional file 3: Figure S3. Overexpression of *OsLG3*. **(a)** *OsLG3* overexpression

construct used for rice transformation. LB, left border; HPT, *Hygromycin*

phosphotransferase; 35S, *cauliflower mosaic virus 35S* promoter; RB, right border. **(b)** The

expression level of *OsLG3* in WT and OE lines analyzed by qRT-PCR. The lines used for

further analysis were labeled in black star. **(c)** Phenotype of *OsLG3*-OE and WT plants

under normal growth conditions.

Additional file 4: Figure S4. Suppression of *OsLG3* by RNAi. **(a)** *OsLG3*-RNAi construct

used for rice transformation. **(b)** The expression level of *OsLG3* in WT and RNAi lines

analyzed by qRT-PCR. The lines used for further analysis were labeled in black star. **(c)**

Phenotype of WT and *OsLG3*-RNAi plants under normal growth conditions at seedling

stage.

Additional file 5: Figure S5. Water loss from detached leaves and expression of drought

stress-related genes in WT, *OsLG3*-OE and RNAi plants under normal and drought stress

conditions. **(a)** Water loss from detached leaves of WT, *OsLG3*-OE and -RNAi plants at

indicated time points. Data are the mean ± SE (n=3). * and ** indicate significant

differences at $P < 0.05$ and $P < 0.01$, respectively. **(b)** Expression of drought stress

responsive genes in WT, *OsLG3*-OE and RNAi under normal and drought stress conditions

(without water for 5 days). Values are means ± SE (n=3).

Additional file 6: Figure S6. qRT-PCR validation of differentially expressed genes in

OsLG3-OE **(a)** and *OsLG3*-RNAi transgenic plants **(b)** compared to WT Plants.

Additional file 7: Table S1. Information of 173 MCC varieties used for association

analysis.

Additional file 8: Table S2. Primer sequences used in this study.

Additional file 9: Table S3. GO_biological_process analysis of the differentially expressed

genes.

Additional file 10: Data Set 1. Information of 1058 *Oryza sativa* varieties and wild rice

used for haplotype analysis.

Additional file 11: Data Set 2. Differentially expressed genes in *OsLG3*-OE Plants

Compared to WT Plants.

Additional file 12: Data Set 3. Differentially expressed genes in *OsLG3*-RNAi Plants

Compared to WT Plants.

References

1.   Huke RE, Huke EH: **Rice area by type of culture: South, Southeast, and East Asia. A**

review and updated data base. *Rice Area by Type of Culture South Southeast & East Asia*

A Review & Updated Data Base 1997.

2.   Qin F, Shinozaki K, Yamaguchi-Shinozaki K: **Achievements and challenges in**

understanding plant abiotic stress responses and tolerance. *Plant & cell physiology*

2011, **52**(9):1569-1582.

3.   Mohanty S, Wassmann R, Nelson A, Moya P, Jagadish S: **Rice and climate change:**

significance for food security and vulnerability. *IRRI Discussion Paper Series No 49 Los*

Baños (Philippines): International Rice Research Institute 2013:14p.

4.   You J, Zong W, Hu H, Li X, Xiao J, Xiong L: **A STRESS-RESPONSIVE**

NAC1-regulated protein phosphatase gene rice protein phosphatase18 modulates

drought and oxidative stress tolerance through abscisic acid-independent reactive

oxygen species scavenging in rice. *Plant Physiol* 2014, **166**(4):2100-2114.

5.   Fukao T, Xiong L: **Genetic mechanisms conferring adaptation to submergence and**

drought in rice: simple or complex? *Current opinion in plant biology* 2013,

16(2):196-204.

6.   Hirayama T, Shinozaki K: **Research on plant abiotic stress responses in the post-genome**

era: past, present and future. *Plant J* 2010, **61**(6):1041-1052.

7.   Miller GAD, Suzuki N, Ciftci-Yilmaz S, Mittler RON: **Reactive oxygen species**

homeostasis and signalling during drought and salinity stresses. *Plant, Cell &*

Environment 2010, **33**(4):453-467.

8.   Hou X, Xie K, Yao J, Qi Z, Xiong L: **A homolog of human ski-interacting protein in rice**

positively regulates cell viability and stress tolerance. *Proceedings of the National*

Academy of Sciences of the United States of America 2009, **106**(15):6410-6415.

9.   Fang Y, Liao K, Du H, Xu Y, Song H, Li X, Xiong L: **A stress-responsive NAC**

transcription factor SNAC3 confers heat and drought tolerance through modulation

of reactive oxygen species in rice. *Journal of experimental botany* 2015,

66(21):6803-6817.

10.   Schmidt R, Mieulet D, Hubberten H-M, Obata T, Hoefgen R, Fernie AR, Fisahn J, San

Segundo B, Guiderdoni E, Schippers JHM *et al*: **Salt-responsive ERF1 regulates reactive**

oxygen species-dependent signaling during the initial response to salt stress in rice.

The Plant Cell 2013, **25**(6):2115-2131.

11.   Wu L, Zhang Z, Zhang H, Wang X-C, Huang R: **Transcriptional modulation of ethylene**

response factor protein JERF3 in the oxidative stress response enhances tolerance of

tobacco seedlings to salt, drought, and freezing. *Plant Physiology* 2008,

148(4):1953-1963.

12.   Ning J, Li X, Hicks LM, Xiong L: **A Raf-like MAPKKK gene *DSM1* mediates drought**

resistance through reactive oxygen species scavenging in rice. *Plant Physiology* 2010,

152(2):876-890.

13.   Yamaguchi-Shinozaki K, Shinozaki K: **Transcriptional regulatory networks in cellular**

responses and tolerance to dehydration and cold stresses. *Annu Rev Plant Biol* 2006,

57:781 - 803.

14.   Zhang H, Jin J, Tang L, Zhao Y, Gu X, Gao G, Luo J: **PlantTFDB 2.0: update and**

improvement of the comprehensive plant transcription factor database. *Nucleic Acids*

Res 2011, **39**(Database issue):D1114-1117.

15.   Ren D, Li Y, Zhao F, Sang X, Shi J, Wang N, Guo S, Ling Y, Zhang C, Yang Z *et al*:

MULTI-FLORET SPIKELET1, which encodes an AP2/ERF protein, determines

spikelet meristem fate and sterile lemma identity in rice. *Plant Physiology* 2013,

162(2):872-884.

16.   Hofmann NR: **SHAT1, a new player in seed shattering of rice**. *The Plant Cell* 2012,

24(3):839.

17.   Zhou Y, Lu D, Li C, Luo J, Zhu B-F, Zhu J, Shangguan Y, Wang Z, Sang T, Zhou B *et al*:

Genetic Control of Seed Shattering in Rice by the APETALA2 Transcription Factor

SHATTERING ABORTION1. *The Plant Cell* 2012, **24**(3):1034-1048.

18.   Komatsu M, Chujo A, Nagato Y, Shimamoto K, Kyozuka J: **FRIZZY PANICLE is**

required to prevent the formation of axillary meristems and to establish floral

meristem identity in rice spikelets. *Development* 2003, **130**(16):3841-3850.

19.   Iwase A, Mitsuda N, Koyama T, Hiratsu K, Kojima M, Arai T, Inoue Y, Seki M, Sakakibara

H, Sugimoto K *et al*: **The AP2/ERF transcription factor WIND1 controls cell**

dedifferentiation in Arabidopsis. *Current Biology* 2011, **21**(6):508-514.

20.   Yaish MW, El-kereamy A, Zhu T, Beatty PH, Good AG, Bi Y-M, Rothstein SJ: **The**

APETALA-2-like transcription factor OsAP2-39 controls key interactions between

abscisic acid and gibberellin in rice. *PLoS genetics* 2010, **6**(9):e1001098.

21.   Boutilier K, Offringa R, Sharma VK, Kieft H, Ouellet T, Zhang L, Hattori J, Liu C-M, van

Lammeren AAM, Miki BLA *et al*: **Ectopic expression of BABY BOOM triggers a**

conversion from vegetative to embryonic growth. *The Plant Cell* 2002, **14**(8):1737-1749.

22.   Zhao Y, Wei T, Yin K-Q, Chen Z, Gu H, Qu L-J, Qin G: **Arabidopsis RAP2.2 plays an**

important role in plant resistance to Botrytis cinerea and ethylene responses. *New*

Phytologist 2012, **195**(2):450-460.

23.   Nakano T, Suzuki K, Fujimura T, Shinshi H: **Genome-Wide Analysis of the ERF Gene**

Family in Arabidopsis and Rice. *Plant Physiology* 2006, **140**(2):411-432.

24.   Ambavaram MM, Basu S, Krishnan A, Ramegowda V, Batlang U, Rahman L, Baisakh N,

Pereira A: **Coordinated regulation of photosynthesis in rice increases yield and**

tolerance to environmental stress. *Nature communications* 2014, **5**:5302.

25.   Jisha V, Dampanaboina L, Vadassery J, Mithöfer A, Kappara S, Ramanan R:

Overexpression of an AP2/ERF type transcription factor OsEREBP1 confers biotic

and abiotic stress tolerance in rice. *PloS one* 2015, **10**(6):e0127831.

26.   Oh SJ, Kim YS, Kwon CW, Park HK, Jeong JS, Kim JK: **Overexpression of the**

transcription factor AP37 in rice improves grain yield under drought conditions. *Plant*

Physiol 2009, **150**(3):1368-1379.

27.   Lee DK, Jung H, Jang G, Jeong JS, Kim YS, Ha SH, Do Choi Y, Kim JK: **Overexpression**

of the OsERF71 Transcription Factor Alters Rice Root Structure and Drought Resistance**. *Plant Physiol* 2016, **172**(1):575-588.

28. Fukao T, Yeung E, Bailey-Serres J: **The Submergence Tolerance Regulator SUB1A Mediates Crosstalk between Submergence and Drought Tolerance in Rice**. *The Plant Cell* 2011, **23**(1):412-427.

29. Zhu D, Wu Z, Cao G, Li J, Wei J, Tsuge T, Gu H, Aoyama T, Qu L-J: **TRANSLUCENT GREEN, an ERF family transcription factor, controls water balance in Arabidopsis by activating the expression of aquaporin genes**. *Molecular Plant* 2014, **7**(4):601-615.

30. Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K: **Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in Arabidopsis**. *Plant Cell* 1998, **10**:1391 - 1406.

31. Haake V, Cook D, Riechmann J, Pineda O, Thomashow MF, Zhang JZ: **Transcription Factor CBF4 Is a Regulator of Drought Adaptation in Arabidopsis**. *Plant physiology* 2002, **130**(2):639-648.

32. Quan R, Hu S, Zhang Z, Zhang H, Zhang Z, Huang R: **Overexpression of an ERF transcription factor TSRF1 improves rice drought tolerance**. *Plant Biotechnology Journal* 2010, **8**(4):476-488.

33. Rong W, Qi L, Wang A, Ye X, Du L, Liang H, Xin Z, Zhang Z: **The ERF transcription factor TaERF3 promotes tolerance to salt and drought stresses in wheat**. *Plant Biotechnology Journal* 2014, **12**(4):468-479.

34.   Zhang G, Chen M, Li L, Xu Z, Chen X, Guo J, Ma Y: **Overexpression of the soybean**

GmERF3 gene, an AP2/ERF type transcription factor for increased tolerances to salt,

drought, and diseases in transgenic tobacco. *Journal of experimental botany* 2009,

60(13):3781-3796.

35.   Trujillo LE, Sotolongo M, Menéndez C, Ochogavía ME, Coll Y, Hernández I,

Borrás-Hidalgo O, Thomma BPHJ, Vera P, Hernández L: **SodERF3, a novel sugarcane**

ethylene responsive factor (ERF), enhances salt and drought tolerance when

overexpressed in tobacco plants. *Plant and Cell Physiology* 2008, **49**(4):512-525.

36.   Huang X, Wei X, Sang T, Zhao Q, Feng Q, Zhao Y: **Genome-wide association studies of**

14 agronomic traits in rice landraces. *Nature genetics* 2010, **42**(11):961-967.

37.   Huang X, Zhao Y, Wei X, Li C, Wang A, Zhao Q: **Genome-wide association study of**

flowering time and grain yield traits in a worldwide collection of rice germplasm.

Nature genetics 2012, **44**.

38.   Setter TL, Yan J, Warburton M, Ribaut JM, Xu Y, Sawkins M, Buckler ES, Zhang Z, Gore

MA: **Genetic association mapping identifies single nucleotide polymorphisms in genes**

that affect abscisic acid levels in maize floral tissues during drought. *Journal of*

experimental botany 2011, **62**(2):701-716.

39.   Yan J, Kandianis CB, Harjes CE, Bai L, Kim EH, Yang X, Skinner DJ, Fu Z, Mitchell S, Li

Q *et al*: **Rare genetic variation at Zea mays crtRB1 increases beta-carotene in maize**

grain. *Nature genetics* 2010, **42**(4):322-327.

40.   Yan J, Warburton M, Crouch J: **Association Mapping for Enhancing Maize (L.) Genetic**

Improvement. *Crop Science* 2011, **51**(2):433.

41.   Famoso AN, Zhao K, Clark RT, Tung CW, Wright MH, Bustamante C, Kochian LV,

McCouch SR: **Genetic architecture of aluminum tolerance in rice (Oryza sativa)**

determined through genome-wide association analysis and QTL mapping. *PLoS*

genetics 2011, **7**(8):e1002221.

42.   Li Q, Yang X, Bai G, Warburton ML, Mahuku G, Gore M, Dai J, Li J, Yan J: **Cloning and**

characterization of a putative GS3 ortholog involved in maize kernel development.

Theoretical and Applied Genetics 2010, **120**(4):753-763.

43.   Lu Y, Zhang S, Shah T, Xie C, Hao Z, Li X, Farkhari M, Ribaut JM, Cao M, Rong T *et al*:

Joint linkage-linkage disequilibrium mapping is a powerful approach to detecting

quantitative trait loci underlying drought tolerance in maize. *Proceedings of the*

National Academy of Sciences of the United States of America 2010, **107**(45):19585-19590.

44.   Xue Y, Warburton ML, Sawkins M, Zhang X, Setter T, Xu Y, Grudloyma P, Gethi J, Ribaut

J-M, Li W *et al*: **Genome-wide association analysis for nine agronomic traits in maize**

under well-watered and water-stressed conditions. *Theoretical and Applied Genetics*

2013, **126**(10):2587-2596.

45.   Liu S, Wang X, Wang H, Xin H, Yang X, Yan J, Li J, Tran LS, Shinozaki K,

Yamaguchi-Shinozaki K *et al*: **Genome-wide analysis of ZmDREB genes and their**

association with natural variation in drought tolerance at seedling stage of Zea mays

L. *PLoS genetics* 2013, **9**(9):e1003790.

46.   Mao H, Wang H, Liu S, Li Z, Yang X, Yan J, Li J, Tran LS, Qin F: **A transposable element**

in a NAC gene is associated with drought tolerance in maize seedlings. *Nature*

communications 2015, **6**:8326.

47. Wang X, Wang H, Liu S, Ferjani A, Li J, Yan J, Yang X, Qin F: **Genetic variation in**

ZmVPP1 contributes to drought tolerance in maize seedlings. *Nature genetics* 2016,

48(10):1233-1241.

48. Singh BP, Jayaswal PK, Singh B, Singh PK, Kumar V, Mishra S, Singh N, Panda K, Singh

NK: **Natural allelic diversity in OsDREB1F gene in the Indian wild rice germplasm**

led to ascertain its association with drought tolerance. *Plant cell reports* 2015,

34(6):993-1004.

49. Lou Q, Chen L, Mei H, Wei H, Feng F, Wang P: **Quantitative trait locus mapping of**

deep rooting by linkage and association analysis in rice. *Journal of experimental botany*

2015, **66**(15):4749-4757.

50. Hu Y, Chong K, Wang T: **OsRAF is an ethylene responsive and root abundant factor**

gene of rice. *Plant Growth Regulation* 2008, **54**(1):55-61.

51. Yu J, Xiong H, Zhu X, Zhang H, Li H, Miao J, Wang W, Tang Z, Zhang Z, Yao G *et al*:

OsLG3 contributing to rice grain length and yield was mined by Ho-LAMap. *BMC*

Biology 2017, **15**(1):28.

52. Wang H, Zhang H, Gao F, Li J, Li Z: **Comparison of gene expression between upland**

and lowland rice cultivars under water stress using cDNA microarray. *Theoretical and*

Applied Genetics 2007, **115**(8):1109.

53. Zhang H, Zhang D, Wang M, Sun J, Qi Y, Li J, Wei X, Han L, Qiu Z, Tang S: **A core**

collection and mini core collection of Oryza sativa L. in China. *Theoretical and Applied*

Genetics 2011, **122**(1):49-61.

54. Lee DK, Yoon S, Kim YS, Kim JK: **Rice OsERF71-mediated root modification affects**

shoot drought tolerance. *Plant Signal Behav* 2017, **12**(1):e1268311.

55.   Cao Y, Song F, Goodman RM, Zheng Z: **Molecular characterization of four rice genes**

encoding ethylene-responsive transcriptional factors and their expressions in response

to biotic and abiotic stress. *Journal of plant physiology* 2006, **163**(11):1167-1178.

56.   Suntres ZE: **Role of antioxidants in paraquat toxicity**. *Toxicology* 2002, **180**(1):65-77.

57.   Apel K, Hirt H: **Reactive oxygen species: metabolism, oxidative stress, and signal**

transduction. *Annual Review of Plant Biology* 2004, **55**(1):373-399.

58.   Mittler R: **Oxidative stress, antioxidants and stress tolerance**. *Trends in Plant Science*

2002, **7**(9):405-410.

59.   Yue Y, Zhang M, Zhang J, Tian X, Duan L, Li Z: **Overexpression of the AtLOS5 gene**

increased abscisic acid level and drought tolerance in transgenic cotton. *Journal of*

experimental botany 2012, **63**(10):3741-3748.

60.   Zhang ZF, Li YY, Xiao BZ: **Comparative transcriptome analysis highlights the crucial**

roles of photosynthetic system in drought stress adaptation in upland rice. *Scientific*

Reports 2016, **6**:19349.

61.   Cheah BH, Nadarajah K, Divate MD, Wickneswari R: **Identification of four functionally**

important microRNA families with contrasting differential expression profiles

between drought-tolerant and susceptible rice leaf at vegetative stage. *BMC genomics*

2015, **16**(1):692.

62.   Lyu J, Li B, He W, Zhang S, Gou Z, Jing Z, Meng L, Xin L, Tao D, Huang W: **A genomic**

perspective on the important genetic mechanisms of upland adaptation of rice. *BMC*

Plant Biology 2014, **14**(1):160.

63.   Kumar A, Dixit S, Ram T, Yadaw RB, Mishra KK, Mandal NP: **Breeding high-yielding**

drought-tolerant rice: genetic variations and conventional and molecular approaches.

Journal of experimental botany 2014, **65**(21):6265-6278.

64.   Ding X, Li X, Xiong L: **Insight into differential responses of upland and paddy rice to**

drought stress by comparative expression profiling analysis. *International Journal of*

Molecular Sciences 2013, **14**(3):5214.

65.   Lenka SK, Katiyar A, Chinnusamy V, Bansal KC: **Comparative analysis of**

drought-responsive transcriptome in Indica rice genotypes with contrasting drought

tolerance. *Plant Biotechnology Journal* 2011, **9**(3):315-327.

66.   Sato Y, Masuta Y, Saito K, Murayama S, Ozawa K: **Enhanced chilling tolerance at the**

booting stage in rice by transgenic overexpression of the ascorbate peroxidase gene,

OsAPXa. *Plant cell reports* 2011, **30**(3):399-406.

67.   Ye N, Zhu G, Liu Y, Li Y, Zhang J: **ABA controls H(2)O(2) accumulation through the**

induction of OsCATB in rice leaves under water stress. *Plant & cell physiology* 2011,

52(4):689-698.

68.   Wu A, Allu AD, Garapati P, Siddiqui H, Dortay H, Zanor M-I, Asensi-Fabado MA,

Munné-Bosch S, Antonio C, Tohge T *et al*: ***JUNGBRUNNEN1*, a reactive oxygen**

species-responsive NAC transcription factor, regulates longevity in *Arabidopsis*. *The*

Plant Cell 2012, **24**(2):482-506.

69.   Lee S, Seo PJ, Lee H-J, Park C-M: **A NAC transcription factor NTL4 promotes reactive**

oxygen species production during drought-induced leaf senescence in Arabidopsis. *The*

Plant Journal 2012, **70**(5):831-844.

70.   Noctor G, Foyer CH: **ASCORBATE AND GLUTATHIONE: Keeping Active Oxygen**

1023   **Under Control**. *Annual Review of Plant Physiology and Plant Molecular Biology* 1998,

1024   **49**(1):249-279.

1025   71.   Duan J, Zhang M, Zhang H, Xiong H, Liu P, Ali J, Li J, Li Z: **OsMIOX, a myo-inositol**

1026   **oxygenase gene, improves drought tolerance through scavenging of reactive oxygen**

1027   **species in rice (Oryza sativa L.)**. *Plant Science* 2012, **196**:143-151.

1028   72.   Xiong H, Li J, Liu P, Duan J, Zhao Y, Guo X, Li Y, Zhang H, Ali J, Li Z: **Overexpression**

1029   **of OsMYB48-1, a novel MYB-related transcription factor, enhances drought and**

1030   **salinity tolerance in rice**. *PloS one* 2014, **9**(3):e92913.

1031   73.   Livak KJ, Schmittgen TD: **Analysis of Relative Gene Expression Data Using Real-Time**

1032   **Quantitative PCR and the 2−ΔΔCT Method**. *Methods* 2001, **25**(4):402-408.

1033   74.   Li J-Y, Wang J, Zeigler RS: **The 3,000 rice genomes project: new opportunities and**

1034   **challenges for future rice research**. *GigaScience* 2014, **3**(1):8.

1035   75.   Curtis MD, Grossniklaus U: **A Gateway Cloning Vector Set for High-Throughput**

1036   **Functional Analysis of Genes in Planta**. *Plant Physiology* 2003, **133**(2):462-469.

1037   76.   Wang Z, Chen C, Xu Y, Jiang R, Han Y, Xu Z, Chong K: **A practical vector for efficient**

1038   **knockdown of gene expression in rice (*Oryza sativa* L.)**. *Plant Mol Biol Rep* 2004,

1039   **22**(4):409-417.

1040   77.   Clough SJ, Bent AF: **Floral dip: a simplified method for*Agrobacterium*-mediated**

1041   **transformation of *Arabidopsis thaliana***. *Plant J* 1998, **16**(6):735-743.

1042   78.   Li J, Pandeya D, Nath K, Zulfugarov IS, Yoo SC, Zhang H, Yoo JH, Cho SH, Koh HJ, Kim

1043   DS *et al*: **ZEBRA-NECROSIS, a thylakoid-bound protein, is critical for the**

photoprotection of developing chloroplasts during early leaf development. *Plant J*

2010, **62**(4):713-725.

79.   Du Z, Zhou X, Ling Y, Zhang Z, Su Z: **agriGO: a GO analysis toolkit for the**

agricultural community. *Nucleic Acids Research* 2010, **38**(suppl_2):W64-W70.

80.   Zhang H, Gao S, Lercher MJ, Hu S, Chen W-H: **EvolView, an online tool for visualizing,**

annotating and managing phylogenetic trees. *Nucleic Acids Research* 2012,

40(W1):W569-W572.

1051

1052

OsEREBP1
OsERF71
OsBIERF1
OsLG3
Sub1B
Sub1C
Sub1A
SNORKEL1
SNORKEL2
OsEBP-89
OsERF922
OsRap2.6
OsBIERF3
OsERF1
FZP
OsAP2-39
OsERF3/OsBIERF4/AP37
MFS1
OsAP2-1
OsDREB2A
OsDREB2B
OsWR1
OsWR4
SERF1
OsDERF1
OsAP21
ARAG1
OsDREB4-1
OsDREB4-2
OsDREB1E
OsDREB1C
OsDREB1D
OsDREB1J
OsDREB1B
OsDREB1G
OsDREB1A
OsDREB1F
HYR

VII
IX
X
IX
VIII
IV
V
II
III

0.1

(a)

FL, dC1–dC6 constructs with GAL BD domain

SD-T   SD-T-H-A+X-α-gal

(b)

	BD-dC1	BD
AD		
AD-OsLG3		

SD-T-L   SD-T-L-H-A+X-α-gal   SD-T-L   SD-T-L-H-A+X-α-gal

(c)

Pro_35S::OsLG3:GFP

Pro_35S::GFP

GFP   Merged   Bright field

(a)

(b)

(a)

WT	OE4	OE7	RI6	RI10
Normal NaCl	Normal NaCl	Normal NaCl	Normal NaCl	Normal NaCl

(b)

WT	OE4	OE7	RI6	RI10
Normal Mannitol	Normal Mannitol	Normal Mannitol	Normal Mannitol	Normal Mannitol

(c)

NaCl

Relative shoot length (%)

NaCl

Relative fresh weight (%)

(d)

Mannitol

Relative shoot length (%)

Mannitol

Relative fresh weight (%)

(a)

(b)

(c)

Hap.	S1	S2	S4	S3	S5	S6	S7	No.
Hap1	T	T	T	T	G	T	A	194
Hap2	A	A	G	A	A	G	C	153
Hap3	T	A	G	T	G	G	C	334
Hap4	T	T	T	T	G	G	C	353
Hap5	T	T	T	T	G	T	C	8
Hap6	A	T	T	A	A	G	C	4
Hap7	T	T	T	A	A	G	C	5
Hap8	T	T	T	T	A	G	C	4
Hap9	A	T	T	T	G	G	C	3

(d) Number of rice accessions

Japonica_UR
Japonica_LR
Indica_UR
Indica_LR
Wild rice

(e)

IRAT109   A A G A A G C
IL342   A A G A A G C
IL381   A A G A A G C
YF   T T T T G T A

AGT   GTA

S1 S2 S3 S4 S5 S6 S7

(f)

Relative germination rate

YF   IL342   IL381   IRAT109
*   ***   ***

(g)

IRAT109 ★
Mawannuo
P77
Mawannuo
Huhan 3 ★

(h)

IRAT109 ★
Yuefu
Zhenghan 9 ★