Natural Variation of *OsLG3* Controls Drought Stress Tolerance in Rice by Inducing ROS

Scavenging

Haiyan Xiong[1][†], Jianping Yu[1][†],Jinjie Li[1], Xin Wang[1], Pengli Liu[1], Hongliang Zhang[1], Yan Zhao[1],

Zhigang Yin[1], Yang Li[1], Yan Guo[2], Binying Fu[3], Wensheng Wang[3], Zhikang Li[3] , Jauhar Ali[4] and

Zichao Li[1]*

[1]Key Lab of Crop Heterosis and Utilization of Ministry of Education / Beijing Key Lab of Crop

Genetic Improvement, China Agricultural University, Beijing 100193, China

[2]State Key Laboratory of Plant Physiology and Biochemistry, College of Biological Sciences, China

Agricultural University, Beijing 100193, China

[3]Institute of Crop Sciences, Chinese Academy of Agricultural Sciences, Beijing, 100081, China

[4]International Rice Research Institute, DAPO Box 7777, Metro Manila 1301, Philippines

[†]These authors contributed equally to this work

*Corresponding author; E-mail: lizichao@cau.edu.cn; Tel: +86 10 62731414

ABSTRACT

Improving performance of rice under drought stress has potential to significant impact on rice
productivity. Previously we reported that *OsLG3* positively control rice grain length and yield. In this
study, we found that *OsLG3* was more strongly expressed in upland rice compared to lowland rice
under drought stress condition. Candidate gene association analysis showed that the natural variation
in *OsLG3* was associated with tolerance to water deficit stress in germinating rice seeds. Transgenic
rice with enhanced *OsLG3* expression exhibited improved tolerance to drought and that is most likely
due to enhanced ROS scavenging efficiency. Phylogenetic analysis and pedigree records indicated that
the tolerant allele of *OsLG3* has potential to improve drought tolerance of *japonica* rice. Collectively,
our work revealed that the natural variation of *OsLG3* contributes to rice drought tolerance and the
elite allele of *OsLG3* is a promising genetic resource for the development of drought-tolerant and high-
yield rice varieties.

Keywords: Upland rice; nature variation; *OsLG3*; drought tolerance; ROS

INTRODUCTION

Agriculture experience more than 50% of average yield losses worldwide due to abiotic stress, especially drought (Huke and Huke, 1997; Qin et al., 2011). Rice (*Oryza sativa* L.), one of the most important staple food for over half the world's population, requires high inputs of water during growth, which results in a number of production challenges due to water shortages and inadequate rainfall during the rice growing season (Mohanty et al., 2013). In the face of these challenges, enhanced performance of rice under drought stress has the potential to significantly improve rice productivity.

Plants have evolved complex signaling pathways that enable them to respond and adapt to unfavorable environmental conditions through morphological, physiological, and biochemical changes (Hirayama and Shinozaki, 2010; Fukao and Xiong, 2013; You et al., 2014). The generation of reactive oxygen species (ROS) is a key process in plant responsiveness to various biotic and abiotic stresses. ROS are important signal transduction molecules, but also toxic by-products of stress metabolism, depending on their overall cellular amount (Miller et al., 2010). At low to moderate concentrations, ROS are likely to function as secondary messengers in stress-signaling pathways, triggering stress defense/adaptation reactions. However, when ROS levels reach above a certain threshold, they can trigger progressive oxidative damage resulting in retarded growth and eventually cell death (Hou et al., 2009). Plants have developed flexible ROS scavenging and regulation pathways to maintain homeostasis and avoid over-accumulation of ROS in cells. There are some evidences supporting that overproduction of ROS-scavenging related genes and synthesis of various functional proteins such as superoxide dismutase (SOD), peroxidase (POD) and catalase (CAT) could increase tolerance to oxidative stress. For example, overexpression of *SNAC3* (Stress-responsive NAM, ATAF1/2, and CUC2 gene 3), increases rice drought and heat tolerance by modulating ROS homeostasis through regulating the expression of genes participated in ROS-scavenging (Fang et al., 2015). Overexpression of an Ethylene-responsive factor (ERF) family gene, *SERF1* (Salt-Responsive ERF1), improved salinity tolerance in rice, mainly due to the regulation of ROS-dependent signaling during the initial phase of salt stress (Schmidt et al., 2013). Overexpressing of *JERF3* (Jasmonate and Ethylene-Responsive Factor 3) improved the expression of genes involved in ROS-scavenging and enhanced tolerance to drought, freezing, and salt in tobacco (Wu et al., 2008). Moreover, overexpression of the

mitogen-activated protein kinase kinase kinase (MAPKKK) gene *DSM1* (Drought-hypersensitive Mutant 1) in rice, increased drought stress tolerance by regulating ROS scavenging. Conversely, deficiency in *DSM1* resulted in a decrease in ROS scavenging and increased drought hypersensitivity (Ning et al., 2010).

Transcription factors (TFs) play central roles in the regulation of gene expression in the stress signaling and adaptation networks (Yamaguchi-Shinozaki and Shinozaki, 2006; Zhang et al., 2011). One set of these is the APETALA2/Ethylene Responsive Factor (AP2/ERF) superfamily. It has been implicated that the ERF proteins play diverse roles in cellular processes involving flower development, spikelet meristem determinacy, floral meristem, plant growth, pathogens and abiotic stress tolerance (Boutilier et al., 2002; Komatsu et al., 2003; Nakano et al., 2006; Yaish et al., 2010; Iwase et al., 2011; Hofmann, 2012; Zhao et al., 2012; Zhou et al., 2012; Ren et al., 2013). There are emerging evidences to support that ERF proteins are involved in response and adaptation to drought stress. For instance, transgenic rice lines overexpression ERF TFs including SUB1A, *OsEREBP1*, *AP37*, *AP59*, *HYR* and *OsERF71*, all showed strong resistance to drought stress (Oh et al., 2009; Fukao et al., 2011; Ambavaram et al., 2014; Jisha et al., 2015; Lee et al., 2016). Other ERF genes including *HARDY*, T*RANSLUCENT GREEN* (*TG*) (Zhu et al., 2014) and DREB genes (Liu et al., 1998; Haake et al., 2002) from *Arabidopsis*, *TSRF1* (Quan et al., 2010) from tomato, *TaERF3* (Rong et al., 2014) from wheat (*Triticum aestivum*), *GmERF3* (Zhang et al., 2009) from soybean, *JERF3* (Wu et al., 2008) from tobacco and *SodERF3* (Trujillo et al., 2008) from sugarcane (*Saccharum officinarum*), have also been found to be involved in responses to water deficit stress condition. Overall, these findings suggested that ERF TFs offer the potential for engineering crops in a way that makes them more efficient under drought stress condition.

Linkage disequilibrium (LD)-based association mapping has been proven to be a powerful tool for dissecting complex agronomic traits and identifying alleles that can contribute to crop improvement (Huang et al., 2010; Setter et al., 2010; Yan et al., 2010; Huang et al., 2012). Candidate gene association analysis, an effective method to validate targets, has become easier and cheaper with the advances in next-generation sequencing (NGS) technology facilitating the discovery and detection of single nucleotide polymorphisms (SNPs) (Yan et al., 2011). This strategy has been used successfully in the

genetic dissection of allelic diversity of genes controlling fatty acid content, kernel size, ABA content,

α-tocopherol and β-carotene content and aluminum tolerance in maize and rice (Li et al., 2010; Li et

al., 2010; Yan et al., 2010; Adam N. Famoso, 2011). There have also been reports of association studies

on crop drought tolerance. For instance, Lu et al. (2010) and Xue et al. (2013) identified some QTLs

underlying drought tolerance in maize by genome-wide association analysis. Liu et al. (2013) found

that DNA polymorphisms in the promoter region of *ZmDREB2.7* were associated with maize drought

tolerance. Analysis of the association found that an 82 bp insertion in *ZmNAC111* and 366 bp insertion

in *ZmVPP1* affected drought tolerance in Maize (Mao et al., 2015; Wang et al., 2016). Recently, an

association study of 136 wild and four cultivated rice accessions identified three coding SNPs and one

haplotype in a DREB (Dehydration Responsive Element Binding) transcription factor (TF),

OsDREB1F, that are potentially associated with drought tolerance (Singh et al., 2015), and nine

candidate SNPs were identified by association mapping of the ratio of deep rooting in rice (Lou et al.,

2015). However, a role for these candidate genes or their causative variations in enhanced drought

tolerance remains to be verified experimentally.

Here, we characterize the role of an ERF family TF, *OsLG3* (LOC_Os03g08470), in rice drought

tolerance. *OsLG3* is located at the same locus as *OsERF62* (Nakano et al., 2006) and *OsRAF* (a Root

Abundant Factor gene in *Oryza sativa*) (Hu et al., 2008). In our previous work, we demonstrated that

OsLG3 plays a positive role in rice grain length without affecting the grain quality (Yu et al., 2017). In

this study, we identified the natural variation in the promoter region of *OsLG3* was associated with

tolerance to drought stress among different rice accessions. Overexpression of *OsLG3* in transgenic

lines resulted in increased drought tolerance via regulating ROS homeostasis. *OsLG3* function as a

pleiotropic gene which can contribute to rice grain length and drought stress tolerance together. These

data provide insights that the elite allele of *OsLG3* is a promising genetic resource for the genetic

improvement of rice drought tolerance and yield.

RESULTS

***OsLG3* is associated with drought stress tolerance in rice.**

In our previous cDNA microarray experiment, comparison between upland rice (UR) (IRAT109 and

Haogelao, drought-resistant *japonica* rice) and lowland rice (LR) (Nipponbare and Yuefu, drought-sensitive *japonica* rice) varieties under well-watered and water deficit conditions showed that *OsLG3* expression was induced to a significantly greater extent by water deficit stress in UR than LR (Wang et al., 2007). To confirm this we analyzed the expression level of *OsLG3* between IRAT109 and Nipponbare under increasingly severe water deficit conditions using Quantitative real-time PCR (qRT-PCR). The data confirmed that *OsLG3* is more highly expressed in IRAT109 than Nipponbare under well-watered conditions and that *OsLG3* expression is strongly induced by drought in IRAT109 but not Nipponbare (**Figure. 1A**). These results suggested that changes in *OsLG3* expression may be involved in response to drought stresses.

We conducted a candidate gene association analysis to investigate if natural variation in *OsLG3* is associated with rice drought tolerance. This approach utilized a mini core collection (MCC) panel (Zhang et al., 2011) (**Table S1**) of 173 varieties that have undergone deep sequencing to a 14.9× average depth (http://www.rmbreeding.cn/Index/). Seeds from MCC lines were germinated on water or in the presence of 15% polyethylene glycol (PEG) to simulate drought stress. The relative germination rate (RGR, ratio of germination rates under 15%PEG condition to germination rates under water condition) of each line was calculated after five days. We identified significant variation in water deficit stress tolerance between the different varieties (**Figure S1** and **Table S1**). A total of 97 SNPs within the *OsLG3* locus from these accessions were identified. To reduce the incidence of false positives, we performed general linear model (GLM) that controls population structure (Q matrix) to identify significant genotypic and phenotypic associations. The association analysis detected three significant SNPs ($P < 1.0 \times 10^{-3}$) (SNP_4352414, SNP_4352886, SNP_4352960) located within the promoter region of *OsLG3* (**Figure 1B**). SNP_4352886, located 2449 bp upstream from the start codon of *OsLG3*, showed greatest significant association with RGR ($P = 2.66 \times 10^{-6}$, **Figure 1B**), contributed 13.9% of the phenotypic variation in the MCC population. SNP_4352886 was in strong LD with two other variations (SNP_4352414, SNP_4352960) in the promoter ($r^2 \geqslant 0.8$), but not with the SNP_4348903, SNP_4352166, SNP_4352793, SNP_4353076 and SNP_4353119, which were identified as marginally significant ($P < 1.0 \times 10^{-2}$) (**Figure 1B**). SNPs identified within the coding region of *OsLG3* were not significantly associated with the RGR trait. Based on results above, we conclude that the nucleotide polymorphisms in the promoter of *OsLG3* are associated with differential

143   germination rates under water deficit.

OsLG3 is an ERF family transcription activator that functions as a homodimer

OsLG3 encodes a putative protein with 334 amino acids. Amino acids 110-159 contain a typical AP2

domain, including 11 putative DNA-binding sites, implying a strong DNA binding capacity, and one

putative nuclear localization signal (NLS) from amino acids 95-121 (**Figure S2A, B**). Phylogenetic

analysis comparing *OsLG3* with known ERF TFs (Nakano et al., 2006) indicated that OsLG3 belongs

to group Ⅶ of the ERF subfamily and is closely related to OsERF71 (Lee et al., 2016), OsEREBP1

(Jisha et al., 2015) and OsBIERF1(Cao et al., 2006), which have been reported to be involved in stress

response (**Figure 2**) .

Transactivation activity assays, in which the DNA-binding domain (GAL4-BD) of GAL4 was

fused to either the full length CDS of OsLG3 or a series of shortened fragments created by deletions

from both the N- and C-termini, indicated that OsLG3 is capable of transcriptional activation, and that

the C-terminal region (amino acids 213-334) is required for this (**Figure 3A**). We also tested

dimerization of OsLG3 protein *in vivo* using a yeast two-hybrid system. To avoid the interference

caused by self-transactivation activity, we used the protein fragment BD-dC2 (amino acids 1-218) as

bait, which lacks the transactivation region. The constructs AD-OsLG3 and BD-dC2 were co-

transformed into yeast strain AH109 to test for a homodimer interaction. AD and BD, AD-OsLG3 and

BD were also co-transformed as negative controls for endogenous transactivation activity. Of these

combinations, only AD-OsLG3 and BD-dC2 co-transformants could grow well in the SD/-Trp-Leu-

Ade-His/X-α-gal medium (**Figure 3B**). These results indicate that OsLG3 potentially functions as a

homodimer in rice. To determine the subcellular localization of OsLG3, *Nicotiana benthamiana* leaves

were infiltrated with *Agrobacterium tumefaciens* (strain EH105) containing 35S:OsLG3-GFP and

35S::GFP, respectively. Confocal imaging analyses showed nuclear-localized fluorescence in

35S:OsLG3-GFP transformed cells, whilst fluorescence from free GFP was distributed throughout the

whole cell (**Figure 3C**). This indicates that OsLG3 is a nuclear-localized protein.

Expression profile of *OsLG3* under different stress treatments and in different plant tissues

The expression profile of *OsLG3* in response to abiotic stresses and hormones in IRAT109 was investigated using qRT-PCR. The transcript level of *OsLG3* was significantly induced under dehydration, polyethylene glycol (PEG), hydrogen peroxide (H_2O_2), NaCl, abscisic acid (ABA), ethylene (ETH) and gibberellin (GA) treatments but remained unchanged under cold treatment (**Figure 4A**). To determine the spatial-temporal expression of *OsLG3* under normal growth conditions, we isolated total RNA in eight representative tissues (root, stem, sheath at seedling stage and root, stem, sheath, leaf, panicle at reproductive stage) from IRAT109, and performed qRT-PCR analysis. *OsLG3* was expressed in all of the tissues tested, and showed higher level in roots compared to other tissues (**Figure 4B**).

Overexpression of *OsLG3* enhances drought stress tolerance in rice

To elucidate the biological functions of *OsLG3* during the stress response, 10 independent transgenic lines in which *OsLG3* was overexpressed under the control of 35S promoter (*OsLG3*-OE) and 14 independent transgenic rice lines in which endogenous *OsLG3* expression level was suppressed by RNA interference (*OsLG3*-RNAi) were generated. qRT-PCR was carried out to quantify *OsLG3* expression level in each transgenic line. Two independent *OsLG3*-OE transgenic lines (OE4, OE7) with highest expression level (**Figure S3**) and two *OsLG3*-RNAi lines (RI6, RI10) with lowest expression level (**Figure S4**) of *OsLG3* were selected for further stress analysis.

Under dehydration treatment using 20%PEG for 3 days, *OsLG3*-OE lines showed greater resistance than WT plants (**Figure 5A**). Almost 44-81% of *OsLG3*-OE plants survived, while only 8-15% of the WT plants survived under this treatment (**Figure 5C**). In contrast, when four weeks old WT and *OsLG3*-RNAi plants were stressed by a slightly less severe dehydration stress (20% PEG for 2.5 days) the relative survival of *OsLG3*-RNAi lines (3–11%) was lower than that of WT (24-42 %) (**Figure 5B, D**).

Under severe drought soil drought stress condition (stop watering for 7 days), the *OsLG3*-OE

lines showed an improved survival rate compared to WT plants. Almost 48-64% of *OsLG3*-OE

survived, whereas only 17-28% of WT plants survived this treatment (**Figure 5E, G**). In contrast, when

WT and *OsLG3*-RNAi plants were treated moderate drought (stop watering for 6 days), 36-47% of the

WT plants had recovered 10 days after watering was restored, but only 8-28% of the RNAi plants

recovered (**Figure 5F, H**).

Interestingly, we found that the leaves of *OsLG3*-OE plants showed a slower rate of water loss

compared to those of WT and *OsLG3*-RNAi lines under dehydration condition (**Figure S5A**),

indicating a role for *OsLG3* in reducing water loss, especially under water deficit conditions.

Expression levels of some characterized stress response genes like *OsLEA3, OsAP37, SNAC1*,

RAB16C, *RAB21* and *OsbZIP73* were monitored in the WT, *OsLG3*-OE and RNAi lines under well-

watered and soil drought conditions. As shown in **Figure S5B**, these genes showed significantly higher

levels under soil drought stress in *OsLG3-OE* plants when compared to WT and RNAi lines.

We also analyzed the growth of *OsLG3*-OE and -RNAi plants under NaCl and mannitol treatment

to induce high salinity and osmotic stress, respectively. Under NaCl treatment, OE lines showed

significantly less suppression of relative shoot growth (shoot length under stress conditions to shoot

length under normal conditions) than both WT and RNAi lines, and less suppression of relative fresh

weight (fresh weight under stress conditions to fresh weight under normal conditions) than WT and

RNAi lines (**Figure 6A, C**). Under mannitol treatment, the relative shoot growth of OE plants was

significantly higher than that of both WT and RNAi lines, and the relative fresh weight in OE lines

was higher than that of WT and RNAi lines (**Figure 6B, D**). These results show that changes in the

expression level of *OsLG3* have a significant effect on the drought tolerance in rice.

Global gene expression analysis revealed alteration in the expression of stress-related genes and

ROS-scavenging related genes.

Preliminary evidence for altered expression of stress-related genes in *OsLG3* -OE and RNAi lines led

us to investigate the pathways regulated by *OsLG3* through transcriptome analysis. RNA samples were

isolated from the leaves of ten-day-old seedlings of WT, *OsLG3-OE* and RNAi plants and Digital Gene

Expression (DGE) analysis was performed. Compared to WT, 223 transcripts were found to have a

greater than 2-fold change in abundance ($P < 0.05$, FDR < 0.05) (**Figure 7A**). 159 genes were up-

regulated in OE plants and down-regulated in RNAi plants, while 64 genes were down-regulated in

OE plants and up-regulated in RNAi plants, respectively (**Figure 7B**). The expression of several of

these genes was tested independent by qRT-PCR to validate the DGE results. Of the genes checked,

six out of eight in OE plants (OE vs WT > 2) and five out of eight in RNAi plants (RNAi vs WT < 0.5)

showed expression patterns consistent with the DGE results (**Figure S6**). Gene Ontology (GO) analysis

showed that the 218 differentially expressed genes affected by the *OsLG3* overexpression and

suppression were significantly enriched in three GO terms (hyper geometric test, $P < 0.01$, FDR<0.05),

including response to stress (GO: 0006950), response to stimulus (GO: 0050896), and response to

abiotic stimulus (GO: 0009628) (**Figure 7C**; **Tables S3**). These results are consistent with our

proposed role for *OsLG3* in the regulation of drought stress tolerance.

Interestingly, DGE analysis showed that ten ROS-scavenging related genes (*APX1*, *APX2*, *APX4*,

APX6, *APX8*, *CATB*, *POD1*, *POD2*, *SODcc1*, *FeSOD*) were up-regulated in OE line and down-

regulated in RNAi line. This result is consistent for a role of *OsLG3* in the control of ROS homeostasis.

To confirm this regulation, we analyzed the expression levels of 15 genes by qRT-PCR (nine genes

identified in the DGE analysis and six other ROS-related genes) in WT, *OsLG3*-OE and -RNAi plants

under well-watered and drought conditions. 13 of the 15 tested genes were significantly up-regulated

under drought stress conditions, and the expression level of 13 genes (except *APX3* and *POD2*) were

significantly higher in OE plants than in WT and RNAi plants under drought stress conditions (**Figure

7D**). Conversely, the expression of *APX1*, *APX2*, *POX8*, *POX22.3* and *POD1* was significantly lower

in the *OsLG3*-RNAi plants compared to WT under drought stress conditions, while the remaining nine

genes did not show a significant difference in expression between WT and RNAi line (**Figure 7D**).

OsLG3 participates in H$_2$O$_2$ homeostasis

The potential role of *OsLG3* in oxidative stress tolerance was examined further by using two oxidative

stress inducers, methyl viologen (MV) (Suntres, 2002) and H$_2$O$_2$. Germinated WT and *OsLG3*-OE and

RNAi plants were sown on $^1/_2$ MS medium and $^1/_2$ MS medium containing 2 μM MV. Application of

MV dramatically repressed seedling growth in all plants, but the *OsLG3*-OE lines exhibited less growth

inhibition compared to WT, while the RNAi plants showed more severe growth inhibition than WT

(**Figure 8A, B and C**). Two-week-old seedlings were treated with 1 mM H_2O_2 or 3 μM MV for 24

hours, followed by 3, 3'-diaminobenzidine (DAB) staining to show the presence of H_2O_2 and nitroblue

tetrazolium (NBT) staining to show the presence of superoxide anion. Under control conditions, WT

or transgenic plants showed similar basal levels of H_2O_2 and superoxide, but DAB staining and NBT

staining were much stronger in the WT than in the *OsLG3*-OE plants under H_2O_2 and MV stress

treatment (**Figure 8D**). These results indicated that overexpression of *OsLG3* in rice can enhance

tolerance to oxidative stress.

Based on the above results, it is hypothesized that *OsLG3* may be involved in the regulation of

ROS homeostasis, leading to enhanced adaption to drought. To confirm this, we stressed the WT,

OsLG3-OE and -RNAi plants with dehydration and 20% PEG, followed by DAB staining to detect

H_2O_2 accumulation. Under well-watered conditions, there was low H_2O_2 accumulation level in WT,

OsLG3-OE and RNAi plants (**Figure 8E**). However, *OsLG3*-OE leaves showed significantly less

accumulation of H_2O_2 than WT plants under drought stress conditions, whereas RNAi lines shows

more H_2O_2 than WT under drought stress. H_2O_2 accumulation under normal and drought stress

conditions was also quantified, supporting the DAB staining results. Less H_2O_2 was detected in the

leaves of *OsLG3*-OE plants than WT and *OsLG3*-RNAi plants under drought stress conditions (**Figure**

8F). The accumulation of ROS may lead to oxidative damage, as indicated by membrane lipid

peroxidation. Monodehydroascorbate (MDA) is a biomarker for membrane lipid peroxidation (Mittler,

2002; Apel and Hirt, 2004; Yue et al., 2012). MDA production under normal growth conditions was

similar in WT and all transgenic plants, whereas under drought stress, MDA production was

significantly lower in the *OsLG3*-OE compared with WT and *OsLG3*-RNAi plants (**Figure 8G).** These

results demonstrate that overexpression of *OsLG3* can reduce the damage to membranes caused by

drought stress.

The effect of *OsLG3*-OE on oxidative stress tolerance implies *OsLG3* may influence ROS

homeostasis. It is well known that superoxide dismutase (SOD), and peroxidase (POD) are the major

ROS-scavenging enzymes in plants under stressed conditions (Mittler, 2002). Four-week-old seedlings

were treated with drought stress for 5 days and the activity of SOD and POD was determined. Results

suggested that under normal growth conditions, *OsLG3*-OE lines have significant higher SOD than

WT and RNAi plants (**Figure 8H**), while the activity of POD did not appear to be significantly affected

in *OsLG3*-OE or RNAi plants (**Figure 8I**). Under drought stress conditions, the activities of POD and

SOD were both significantly enhanced in OE plants and significantly reduced in RNAi plants

compared to WT (**Figure 8H, I**). These results imply that the function of *OsLG3* in drought tolerance

may be associated with an enhanced antioxidant response to counteract oxidative stress under drought.

The elite *OsLG3* allele can be used for improving *japonica* rice drought tolerance

To analyze the relationship between *OsLG3* haplotypes and drought tolerance, we investigated the

phylogenetic relationship of 1058 deep-sequenced (depth≈14.9x) rice accessions, including 251 UR,

415 LR and 392 wild rice varieties originating from a wide geographic range (**Supplemental Data**

Set 1). Forty-five SNPs were identified by minor allele frequency (MAF) > 0.05 and missing rates

⩽ 50%. Phylogenetic analysis based on these variations showed that there is a clear differentiation

between *japonica*-UR and *japonica*-LR rice (**Figure 9A**). To get further insight into the phylogenetic

relationship, ten elite *japonica*_UR varieties (IRAT109, IAC150/76, IRAT266, Guangkexiangnuo,

Shanjiugu, Taitung_upland328, Jaeraeryukdo, Riku aikoku, Padi darawal, Malandi 2) were chosen as

an UR pool based on their strong drought resistance, and ten typical *japonica*-LR varieties

(Nipponbare, Yuefu, Guichao2, Koshihikari, Zhonghua11, Early_chongjin, Xiushui115, IR24,

Ningjing3, Xiushui114) were chosen as a LR pool. Seven SNPs (SNP_4352414, SNP_4352886,

SNP_4352960, SNP_4352792, SNP_4352797, SNP_4353103 and SNP_4353347) that differed

between the upland rice pool and lowland rice pool were identified. These seven SNPS showed a

clear phylogenetic distinction between *japonica*-UR and *japonica*-LR (**Figure 9B**). On the basis of

these seven SNPs, we could divide the sequences of the 1058 cultivated varieties into nine haplotypes

of which there were four main variants, Hap1, Hap2, Hap3, and Hap4 (**Figure 9C, D**). *OsLG3*[Nipponbare]

is representative of Hap1, which is mainly composed of *japonica*-LR rice, whereas *OsLG3*[IRAT109]

belongs to Hap2, which is mainly composed of *japonica*-UR rice and is the second largest group.

Hap3 is mainly composed of *indica* rice, and Hap4 is mainly composed of wild rice. The results

indicated that *OsLG3* has clear differentiation in *japonica* rice, which can be divided into *japonica*-

UR and *japonica*-LR rice, while no clear division in *indica* rice. Therefore, we designated Hap1 and

Hap2 as the sensitive and tolerant alleles, respectively, of *OsLG3* in *japonica* rice.

305    To confirm the genetic effect of different alleles of *OsLG3* on rice drought tolerance, two

306  introgression lines (IL342 and IL381) were selected from a cross between IRAT109 (donor parent)

307  and Yuefu (receptor parent). The *OsLG3* allele in these two lines shares sequence similarity with

308  IRAT109, with seven matching SNPs in the promoter (including the three significant loci describe in

309  candidate gene association analysis section), whereas Yuefu carries the same *OsLG3* allele as

310  Nipponbare (**Figure 9E**). Thus, the *OsLG3* allele found in IRAT109, IL342 and IL381 was considered

311  to be the drought-tolerant allele and the allele in Yuefu the drought-sensitive allele, respectively.

312  When germinated on 15% PEG conditions, the RGR of Yuefu was 27.0% compared to germination

313  on water, while the   RGR of   IL342, IL381 and IRAT109 was 35%, 36% and 46%, respectively

314  (**Figure 9F**). These results support the hypothesis that the nucleotide polymorphisms in *OsLG3*

315  contribute to enhanced rice drought tolerance.

316    According to the pedigree records, Huhan3, one of green super rice accessions, is an elite upland

317  rice that is widely grown in Hubei province of China because of its high yield and outstanding ability

318  to conserve water. One of its parents is IRAT109. A re-sequencing study showed that Huhan3 carries

319  the allele of *OsLG3* deriving from IRAT109 (**Figure 9G**). Moreover, Zhenghan9, another elite upland

320  rice, is derived from the cross of IRAT109 and Yuefu. Zhenghan9 retained the favorable allele of

321  *OsLG3* and an allele for an unidentified grain quality-related gene derived from Yuefu (**Figure 9H**).

322  These observations illustrate the successful outcome of combining the $OsLG3^{IRAT109}$ allele and those

323  alleles for other unidentified grain quality- related genes, for improving drought tolerance and grain

324  appearance quality, which had been employed by breeders.

325  **DISCUSSION**

326  **Natural variation in the promoter of *OsLG3* is associated with drought tolerance in rice**

327  Despite many studies investigating the transcriptional response to drought stress (Fukao and Xiong,

328  2013; Lyu et al., 2014; Cheah et al., 2015; Zhang et al., 2016), how natural sequence variation is

329  associated with phenotypic variations in drought tolerance remains largely unknown. Because of the

330  polygenic inheritance, low heritability, and strong genotype-by-environment interactions of drought

331  resistance-related traits, there are few reports on the cloning and identification of drought resistant

genes by association analysis and positive mutant screening methods (Kumar et al., 2014; Lou et al.,

2015; Singh et al., 2015). In this study, using candidate gene association analysis, we detected three

significant SNPs ($P < 1.0 \times 10^{-3}$) (SNP_4352414, SNP_4352886, SNP_4352960) located within the

promoter region of *OsLG3* were significantly associated with the RGR trait (**Figure 1C**). From our

previous work, transient assays showed that four out of seven SNPs are conserved and play a

significant role in regulating *OsLG3* expression (Yu et al., 2017). In this study, transgenic experiments

further support the hypothesis that increased expression of *OsLG3* enhances rice drought stress

tolerance.

To analyze the relationship between *OsLG3* haplotypes and drought tolerance, we investigated

the phylogenetic relationship of 1058 deep-sequenced rice accessions. Based on the seven major SNPs,

we divide the sequences of the 1058 cultivated varieties into nine haplotypes of which there were four

main variants, Hap1 (mainly composed of *japonica*-LR rice), Hap2 mainly composed of *japonica*-UR

rice, Hap3 (mainly composed of *indica* rice), and Hap4 (mainly composed of wild rice). The variations

of *OsLG3* in these accessions showed that there is a clear differentiation between *japonica*-UR and

japonica-LR rice, while no clear division in *indica* and wild rice (**Figure 9**). These results implicated

the Hap2 variant of *OsLG3* may improve drought tolerance of cultivated *japonica* rice. Re-analysis of

two introgression lines (IL342 and IL381) derived from a cross between IRAT109 (donor parent) and

Yuefu (receptor parent) indicated that selection of Hap2 was effective in the improvement of drought

tolerance (**Figure 9E**). Interestingly, our previous work indicated that *OsLG3* acts as an important

positive regulator of grain length and could improve rice yield (Yu et al., 2017). In fact, the pedigree

records in upland rice breeding showed that the tolerant allele of *OsLG3* had been incorporated into

elite varieties via breeding (**Figure 9G**). We propose that pyramiding the beneficial allele of *OsLG3*

with other yield- and quality-related genes could contribute to the breeding of elite rice varieties

because of its pleiotropic effects on traits.

***OsLG3* was cloned from upland rice and plays a positive role in rice drought tolerance.**

UR has evolved enhanced drought-resistance compared to LR, derived from natural and artificial

selection over time under drought conditions. UR performs better under drought conditions, with

stronger water-retention ability, larger root volumes and higher biomass production (Lenka et al., 2011;

Ding et al., 2013). Therefore, UR is highly suitable to be taken up as research material regarding

mechanisms underlying drought resistance. From the previous work of expression profiles from typical

LR (Nipponbare and Yuefu, drought-sensitive *japonica* rice) and UR varieties (IRAT109 and Haogelao,

drought-resistant *japonica* rice) under water deficit stress conditions using cDNA microarray (Wang

et al., 2007), we found the transcription of *OsLG3* can be induced to a greater extent in the UR varieties

compared to the LR varieties during drought stress. When IRAT109 and Nipponbare were subjected

to soil drought stress, expression of *OsLG3* in Nipponbare was not induced, whereas in IRAT109

OsLG3 shows significant induction under both slight drought (SLD) and moderate drought (MOD)

conditions (**Figure 1A**). Using candidate gene association analysis, we found that nucleotide

polymorphisms in the promoter region of *OsLG3* are associated with different levels of water deficit

tolerance among rice varieties at the germination stage (**Figure 1B**). All these results indicated that

OsLG3 might play a role in the observed drought stress response of upland rice.

To assess the effect of *OsLG3* on water deficit stress responses, we tested the growth response of

OsLG3-OE and -RNAi plants under simulated drought stresses. *OsLG3*-OE lines showed higher

survival under dehydration caused by 20% PEG6000 and soil drought stress caused by stopping

watering (**Figure 5**). Plants with reduced *OsLG3* expression showed reduced survival, suggesting that

this gene might be involved in response to drought stress. A slower rate of water loss was observed in

the leaves of *OsLG3*-OE compared to WT and RNAi plants (**Figure S5A**), suggesting that *OsLG3*

plays an important role in reducing water evaporation. Collectively, these results demonstrate that

OsLG3 is a positive regulator of drought stress response in rice. *OsLG3*-OE plants also showed

enhanced growth compared to WT and RNAi lines under mannitol and NaCl treatments, which

suggests that *OsLG3* may be involved in osmotic and high salinity stress tolerance through cross-talk

between water deficit, osmotic and high salinity stress response pathways.

Potential mechanisms of *OsLG3* in drought stress tolerance

We found that the expression levels of a set of stress-related genes like *OsLEA3*, *OsAP37*, *SNAC1* are

increased in *OsLG3*-OE lines compared to WT, and decreased in *OsLG3*-RNAi lines before and after

drought treatment (**Figure S5B**). This was confirmed by DGE analysis (**Table S3**). We investigated if

the altered response to drought was associated with global changes in the expression of stress-related

genes before stress application. Analyses indicated that many stress-related genes were up-regulated in the OE line and down-regulated in the RNAi line (**Figure 7C**), which is consistent with the observed phenotypes of *OsLG3*-OE and -RNAi lines. Interestingly, the expression level of some ROS-scavenging related genes showed increased expression levels in OE and decreased abundance in RNAi lines too. For instance, the ascorbate peroxidase gene, *OsAPX1* plays a positive role in chilling tolerance by enhancing H_2O_2-scavenging (Sato et al., 2011). *DSM1* mediates drought resistance through ROS scavenging in rice (Ning et al., 2010). *OsCATB* prevent the excessive accumulation of H_2O_2 under water stress (Ye et al., 2011). Thus, these DGE analyses are consistent with our hypothesis that the function of *OsLG3* in abiotic stress tolerance occurs through the transcriptional regulation of stress-related and ROS scavenging-related genes.

Previous studies have demonstrated that plant response to abiotic stresses is mediated through the regulation of ROS metabolism (Ning et al., 2010; Lee et al., 2012; Wu et al., 2012; Schmidt et al., 2013; Fang et al., 2015). For example, overexpression of a NAC gene, *SNAC3*, increased drought and heat tolerance by modulating ROS homeostasis through regulating the expression of genes for ROS-scavenging and production enzymes (Fang et al., 2015). Overexpression of an ERF gene, *SERF1*, improved salinity tolerance mainly due to the regulation of ROS-dependent signaling during the initial phase of salt stress in rice (Schmidt et al., 2013). The present data revealed overexpression of *OsLG3* led to the up-regulation of many ROS scavenge-associated genes, while suppression of *OsLG3* caused the down-regulation of these genes (**Figure 7D**). Furthermore, the H_2O_2 and MDA content which accumulated in the leaves of *OsLG3*-OE plants was significantly lower than that in the WT and *OsLG3*-RNAi plants (**Figure 8C, D**), suggesting that the improved drought tolerance of *OsLG3*-OE plants may be due to the efficient scavenging of ROS. Relevant to this, *OsLG3*-OE seedlings exhibited better growth under oxidative stress caused by application of MV. In contrast, RNAi plants showed enhanced sensitivity to oxidative stress. Therefore, it is proposed that the function of *OsLG3* in drought tolerance is associated with the enhancing activity of ROS-scavenging.

To scavenge or detoxify excess stress-induced ROS, plants have developed a complex antioxidant system comprising of the non-enzymatic as well as enzymatic antioxidants (Noctor and Foyer, 1998; Miller et al., 2010). The POD and SOD activities were found to be higher in the *OsLG3*-OE lines as

compared to WT under drought stress, and *OsLG3*-RNAi lines showed the reverse results (**Figure 8H**).

These data suggest that *OsLG3*-OE plants have enhanced activity of ROS-scavenging enzymes which

significantly contributes to the reduction of ROS accumulation, and thereby improved drought stress

tolerance.

In conclusion, here we present evidence that *OsLG3* is induced by water deficit stresses and its

induction is greater in UR IRAT109 compared to LR Nipponbare under drought. Nucleotide

polymorphisms in the promoter region of *OsLG3* are associated with water deficit tolerance in

germinating rice. Transgenic plants overexpressing *OsLG3* showed improved growth under drought

stress, probably via inducing ROS scavenging by controlling downstream ROS-related genes.

Phylogenetic analysis indicated that the tolerant allele of *OsLG3*, identified in drought-tolerant

japonica rice varieties, could be introduced to rice to improve the drought tolerance. The pedigree

records in upland rice breeding showed that the tolerant allele of *OsLG3* had been incorporated into

elite varieties via breeding. Importantly, the tolerant allele of *OsLG3* is a promising genetic resource

for the development of drought-tolerant and high yield rice varieties using traditional breeding

approaches or genetic engineering.

MATERIALS AND METHODS

Plant materials and stress treatments

Rice varieties IRAT109 and Nipponbare were used for quantitative real-time PCR (qRT-PCR) analysis

of *OsLG3* transcript levels under various stresses and hormone treatments, and Nipponbare was used

for all transgenic experiments. For qRT-PCR analysis the expression level of *OsLG3* under simulated

drought conditions, the seeds of Nipponbare and IRAT109 were sown in flower pots (140 mm diameter

×160 mm deep) with well-mixed soil (forest soil: vermiculite in a ratio of 1:1), and grown in the

greenhouse under well water conditions at 28 ℃/ 26 ℃ and 12 hours light/12 hours dark photoperiod.

Three-week old plants were subjected to drought treatment with stop watering for 0, 5, 6 and 7 days.

From these, four stress treatment levels were categorized as no stress (NS), slight drought (SLD),

moderate drought (MOD) and severe drought (SED), respectively. For qRT-PCR analysis the

expression level of *OsLG3* under various abiotic stresses and phytohormone treatments, three-week-

old IRAT109 seedlings grown in Hoagland solution (PPFD of 400 μmol/m^{2}s^2, 12 hours light (28 ℃)/12

hours dark (26 ℃)) were subjected to different treatments with 20% (w/v) polyethylene glycol (PEG)

6000, NaCl (200 mM), cold (4 ℃), hydrogen peroxide (H$_2$O$_2$,1 mM), abscisic acid (ABA,100 μM),

ethylene (ETH,100 μM), GA (100 μM) and dehydration by exposure in air. Leaf tissue was harvested

at 0, 1, 2, 4, 6, 9, 12 and 24 hours after PEG, NaCl, cold, H$_2$O$_2$, ABA, ETH, and GA treatments, and

at 0, 1, 2, 3, 4, 5, 6, 7 and 8 hours after dehydration treatment.

The seeds of WT, *OsLG3*-OE lines (OE4, OE7) and *OsLG3*-RNAi lines (RI6, RI10) were

germinated in $^1/_2$ MS medium for various stress evaluations. For dehydration treatment, uniformly

germinated seeds were transplanted into bottom removed 96-well PCR plates, and grown

hydroponically using Hoagland solution at 28 ℃/26 ℃ (day/night) with a 12 hours photoperiod. Three-

week-old plants were treated with 20% (w/v) PEG6000 solution for about three days, and recovered

with water for ten days. Each stress test was repeated three times. For drought stress treatment, three-

day-old WT and transgenic seedlings were transplanted to well-mixed soil (forest soil: vermiculite in

a ratio of 1:1), and grown for four weeks under a normal watering regime. Drought stress treatment

was then applied by stopping irrigation for about seven days. After all leaves had completely rolled,

watering was resumed for 10 days. The survival ratio of each line was calculated as the number of

surviving plants over the number of treated plants in each flowerpot. To evaluate the tolerance of rice

seedlings to osmotic, high salinity and oxidative stress treatment, three-day-old WT, *OsLG3*-OE and

OsLG3-RNAi seedlings (10 plants per replicate, three replicates) were transplanted to $^1/_2$ MS medium

(mock treatment) or $^1/_2$ MS medium containing 200 mM mannitol, 150 mM NaCl or 2 μM MV

respectively. Plants were grown for seven days under 12 hours light (28 ℃)/12 hours dark (26 ℃)

photoperiod conditions and then the shoot height and fresh weight of all plants was measured.

Gene expression analysis

Total RNA was extracted using RNAiso Plus (TaKaRa) according to the manufacturer's instructions.

4 μg of the DNase-treated RNA were reverse transcribed by using M-MLV reverse transcriptase

(TaKaRa). qRT-PCR was performed in an Applied Biosystems 7500 Real Time PCR system (ABI,

USA) using SYBR Premix Ex TaqTM II (TaKaRa) according to the protocol previously described

(Duan et al., 2012; Xiong et al., 2014). The gene-specific primers used for qRT-PCR are listed in

Supplemental Table S2. Rice *Actin1* gene was used as the internal reference gene for data

normalization (Livak and Schmittgen, 2001).

***OsLG3*-gene association analysis of rice drought tolerance among 173 rice genotypes**

173 cultivated varieties from the mini core collection of Chinese cultivated rice (Zhang et al. 2010),

including 130 *indica* and 43 *japonica* rice varieties (**Supplemental Table S1**) were selected for the

candidate gene association mapping. To perform the analysis, we obtained phenotypic data of relative

germination rates (RGR, ratio of germination rates under stress conditions to germination rates under

water conditions) from growth in water and 15% PEG6000 treatment. Briefly, 50 seeds of each line

were placed in petri dishes (90 mm diam) lined with filter paper. 10 ml water or 15% PEG6000 was

added to each petri dish as mock treatment or to induce osmotic stress, respectively. All petri dishes

were placed in a 28 ℃ greenhouse and the germination rates were assessed after five days. Genotype

data for each line was acquired from 'The 3,000 rice genomes project' (Li et al., 2014). The SNP data

were filtered out. After filtering, a total of 97 SNPs remained in a 5-kb region surrounding the *OsLG3*

gene. Association analysis using a general linear model with the population structure (Q matrix)

method was conducted using TASSEL 5.2.28 software. The Q matrix was estimated from the genomic

data to control for population structure.

Plasmid construction and rice transformation

To generate the overexpression construct, the full-length cDNA of *OsLG3* was amplified from the first-

strand cDNA of IRAT109 with specific primers (**Supplemental Table S2**) using PrimeSTAR® HS

DNA Polymerase (TaKaRa), digested with *Kpn*I and *Pac*I, and cloned into binary vector pMDC32

(Curtis and Grossniklaus, 2003). For construction of RNA interference (RNAi) plasmid, a 415 bp

fragment with low similarity to other rice genes located at the 5' end of *OsLG3* was amplified by PCR

with specific primer, digested with *Kpn*I and *Bam*HI and then *Spe*I and *Sac*I sites, and cloned into the

pTCK303 vector as described previously (Wang et al., 2004).

Subcellular localization

For subcellular localization analysis, the full-length open reading frame (ORF) of *OsLG3* without

terminal codon was amplified, and the amplified fragments were digested with *Kpn*I and *Pac*I, and

then cloned in to pMDC83 vector, fused with the GFP reporter gene driven by *CaMV 35S* promoter.

The fusion construct (35S:OsLG3-GFP) and control (35S:GFP) were transformed into *Agrobacterium*

tumefaciens strains (EH105), and then infiltrated to five-week-old *Nicotiana benthamiana* leaves

(Clough and Bent, 1998). The fluorescence signal was examined through a confocal laser scanning

microscope OLYMPUS FV1000 with excitation at 488 nm and emission at 525 nm.

Transactivation and dimerization assay

For transactivation assay, The full length coding region and truncated fragments of *OsLG3* generated

by PCR amplification were fused in frame to the GAL4 DNA binding domain in the vector of pGBKT7

(Invitrogen). The plasmids of FL (the full length coding region of OsLG3, amino acids 1-334), dC1

(amino acids 1-218), dC2 (amino acids 1-167), dC3 (amino acids 1-106), dC4 (amino acids 107-334),

dC5 (amino acids 168-334) and dC6 (amino acids 213-334) were constructed. These constructs were

introduced into yeast strain AH109 by LiAc-mediated yeast transformations, and screened on the

selective medium plates without Tryptophan (SD/-Trp). The PCR-verified transformants were

transferred to SD medium without Tryptophan/Histidine/Adenine (SD/-Trp/-His/-Ade) for 3 days. The

β-galactosidase activity was performed according to the *in vivo* agar plate assay (X-α-gal in medium).

For dimerization assay, AD-OsLG3 with full length coding region of *OsLG3* fused in frame to

the GAL4 DNA activating domain in the vector of pGADT7 (Invitrogen) was constructed. The

constructs AD-OsLG3 and BD-dC2 were co-transformed into yeast strain AH109. AD and BD empty,

AD-OsLG3 and BD empty were also co-transformed as negative controls. All transformed cells were

screened on the selective medium plates without Tryptophan and Leucine (SD/-Trp-Leu), The PCR-

verified transformants were transferred to SD medium with X-α-gal and without

Tryptophan/Histidine/Adenine/Leucine (SD/-Trp-Leu-Ade-His/X-α-gal) for 3 days.

Physiological measurements

The rate of water loss under dehydration condition was measured as described previously (Xiong et

al., 2014). Ten plants of each line were used in each replicate, and three replicates were made for each

line. Histochemical assays for ROS accumulation were determined according to the previously

described method (Li et al., 2010; Wu et al., 2012). Briefly, the qualitative detection of H_2O_2

accumulation was detected by DAB staining. Excised leaves were treated with DAB staining solution (1mg/mL DAB pH 3.8) at 28 ℃ for 12 hours in the dark. After staining, the leaves were decolorized with Acetic acid/Ethanol (1:3) for 60 min, and rehydrated in 70% (v/v) alcohol for 24 hours at 28 ℃. Each experiment was repeated on at least 10 different plants, and representative images were shown. Superoxide anion radical accumulation was detected by NBT staining as described previously. The leaf samples were excised and immediately placed in 50 mM sodium phosphate buffer (pH 7.5) containing 6 mM NBT, at 28 ℃ for 8 h in the dark. The quantitative measurement of H_2O_2 concentrations was performed with an Amplex Red Hydrogen / Peroxidase Assay Kit (Molecular Probes) (Invitrogen, http://www.invitrogen.com) as described by the manufacturer's instructions. Briefly, leaf samples from both the well water and drought stressed (without water for 5 days) plants were ground in liquid nitrogen, and 100 mg of ground frozen tissue from each sample was placed in a 2 mL Eppendorf tube and kept frozen. 1 mL precooled sodium phosphate buffer (20 mM, pH 6.5) was immediately added into the tube and mixed. After centrifugation (10,000 g, 4 ℃, 10 min) the supernatant was used for the assay. Measurements were performed using a 96-well microplate reader (PowerWave XS2, BioTek) at an absorbance of 560 nm. MDA content was measured as described previously (Xiong et al., 2014).

The activity of antioxidant enzymes including superoxide dismutase (SOD) and peroxidase (POD) were measured following the protocols described previously (Duan et al., 2012). The units of the antioxidant enzyme activities were defined as follows: One unit of the SOD activity was defined as the quantity of enzyme required to cause 50% inhibition of the photochemical reduction of NBT per minute at 560 nm. One unit of POD activity was defined as the amount enzyme required to cause a 0.01 absorbance increase per minute at 470 nm.

Digital gene expression (DGE) profiling and gene ontology (GO) analysis

WT Nipponbare and transgenic lines OE7 and RI10 were used for digital gene expression (DGE) profiling analysis. Ten-day-old seedlings grown on $^1/_2$ MS medium were harvested for total RNA extraction as described in qRT-PCR section. DGE was performed at the Beijing Genomics Institute (http://www.genomics.cn) using Illumina Hiseq2000 sequencing technology. Transcripts with significant differential expression between *OsLG3*-OE and WT or between *OsLG3*-RNAi and WT

553 plants were identified as those with *P* < 0.05 using a False Discovery Rate (FDR) of <0.05 and a fold

554 change cutoff of >2. Gene Ontology (GO) analysis was performed using agriGO (Du et al., 2010)

555 (http://bioinfo.cau.edu.cn/agriGO/). Representative differentially expressed genes were confirmed by

556 qRT-PCR. The primers are listed in **Supplemental Table S2**.

Sequence analysis and phylogenetic analysis

558 The phylogenetic tree was analyzed by MEGA6 software based on neighbor-joining method and

559 bootstrap analysis (1,000 replicates). The EvolView online tool (Zhang et al., 2012) was used for

560 visualizing the phylogenetic tree. Multiple sequence alignment was performed with ClustalW. Gene

561 annotation was conducted in RGAP (http://rice.plantbiology.msu.edu/). All SNP data were obtained

562 from the Rice functional genomics and breeding database (RFGB, http://www.rmbreeding.cn/Index/).

Accession numbers

Sequence data from this article can be found in the Rice Genome Annotation Project Database and

Resource (RGAP) (http://rice.plantbiology.msu.edu) under following accession numbers: *OsLG3*

(LOC_Os03g08470); *Actin1* (LOC_Os10g36650); *OsLEA3* (LOC_Os05g46480); *AP37*

(LOC_Os01g58420); *SNAC1* (LOC_Os03g60080); *RAB21* (LOC_Os11g26790); *RAB16C*

(LOC_Os11g26760); *OsbZIP23* (LOC_Os02g52780); *Apx1* (LOC_Os03g17690); *Apx2*

(LOC_Os07g49400); *Apx3* (LOC_Os04g14680); *Apx5* (LOC_Os12g07830); *Apx6*

(LOC_Os12g07820); *Apx8* (LOC_Os02g34810); *OsPox8.1* (LOC_Os07g48010); *Pox22.3*

(LOC_Os07g48020); *FeSOD* (LOC_Os06g05110); *SODcc1* (LOC_Os03g22810); *SODcc2*

(LOC_Os07g46990); *POD1* (LOC_Os01g22370); *POD2* (LOC_Os03g22010); *CATB*

(LOC_Os06g51150); *DSM1* (LOC_Os02g50970); *RAB16D* (LOC_Os11g26750); *OsNCED4*

(LOC_Os07g05940); *OsNCED3* (LOC_Os03g44380); *OsDhn1* (LOC_Os02g44870); *OsSRO1C*

(LOC_Os03g12820); *OsMYB48* (LOC_Os01g74410); *OsITPK4* (LOC_Os02g26720); *OsLEA3-2*

(LOC_Os03g20680).

ACKNOWLEDGMENTS

We thank Amelia Henry (International Rice Research Institute), Steven Burgess (eLIFE) and Andrew

Plackett (University of Cambridge) for critical reading and suggest revisions to this manuscript. This

work was supported by grants from Program of S&T cooperation (2015DFG31900), National Natural

Science Foundation of China (31601278), China Postdoctoral Science Foundation funded project

(2014M560140 and 2015T80157), Ministry of Agriculture of the People's Republic of China

(2014ZX08009-003-002) and Key Program of Hainan Department of Science and Technology

(ZDYF2016217).

Figure Legends

Figure 1. *OsLG3* is associated with drought tolerance. (a) qRT-PCR analysis of *OsLG3* in

Nipponbare and IRAT109 under different soil drought stress levels. NS, No stress; SLD, Slight drought;

MOD, Moderate drought; SED, Severe drought. **(b)** Analysis of the association between pairwise LD

of DNA polymorphisms in the *OsLG3* gene and water deficit tolerance. Schematic of OsLG3 is shown

on the X-axis and the significance of each variation associated with seedling RGR (ratio of germination

rates under 15%PEG condition to germination rates under water conditions) is shown on the Y-axis.

The SNPs with significant variation ($P < 1.0 \times 10^{-2}$) between genotypes are connected to the pairwise

LD diagram with a solid line. Black dots in the pairwise LD diagram highlight the strong LD of

SNP_4352886 (filled red circle, $P = 2.66 \times 10^{-6}$) and two significant variations, SNP_4352414 and

SNP_4352960 (open red circles, $P < 1.0 \times 10^{-3}$). The SNP_4348903, SNP_4352166, SNP_4352793,

SNP_4353076 and SNP_4353119, which are marginally significant ($P < 1.0 \times 10^{-2}$) are denoted by

red triangles.

Figure 2. Phylogenetic tree of OsLG3 homologues. Neighbour-Joining phylogenetic analysis of

OsLG3 protein sequence in the context of other characterized AP2/ERF proteins from rice. The

phylogenetic tree was constructed using the ClustalW and MEGA programs. Tree topology with

bootstrap support is based on a percentage of 1000 replicates. Accession numbers are as follows:

ARAG1, LOC_Os02g43970; OsAP21, LOC_Os01g10370; OsDREB4-1, LOC_Os02g43940;

OsDREB4-2, LOC_Os04g46400; OsDREB2A, LOC_Os01g07120; OsDREB2B, LOC_Os05g27930;

OsBIERF3, LOC_Os02g43790; OsERF1, LOC_Os04g46220; OsERF922, LOC_Os01g54890;

OsWR1, LOC_Os02g10760, OsWR4, LOC_Os06g08340; SNORKEL1, AB510478; SNORKEL2,

AB510479; OsBIERF1, LOC_Os09g26420; OsEBP-89, LOC_Os03g08460; OsEREBP1,

LOC_Os02g54160; OsLG3, LOC_Os03g08470; OsERF71, LOC_Os06g09390; Sub1A, DQ011598;

Sub1C, LOC_Os09g11480; Sub1B, LOC_Os09g11460; MFS1, LOC_Os05g41760; OsAP2-39,

LOC_Os04g52090; OsERF3/OsBIERF4/AP37, LOC_Os01g58420; FZP, LOC_Os07g47330;

OsRap2.6, LOC_Os04g32620; OsAP2-1, LOC_Os11g03540; OsDREB1D, LOC_Os06g06970.

Figure 3. Transactivation assay and Subcellular localization of OsLG3. **(a)** Transactivation assay

of OsLG3 fragments. Fusion proteins of the GAL4 DNA-binding domain and different portions of

OsLG3 were expressed in yeast strain AH109 and streaked on control plate (SD/-Trp) or selective plate

(SD/-Trp-Ade-His/X-α-gal), respectively. FL indicates the full-length CDS of OsLG3; dC1 to dC6

indicate the mutated forms of OsLG3 (nucleotide positions are labeled in the diagrams), respectively.

(b) Dimerization analysis. BD with AD empty vector, AD empty with BD-dC2, AD-OsLG3 with BD

empty and AD-OsLG3 with BD-dC2, were co-transformed in to yeast strain AH109 yeast cells and

streaked on control plate (SD/-Trp-Leu) or selective plate (SD/-Trp-Leu-Ade-His/X-α-gal),

respectively. The plates were incubated for 3 days. **(c)** Subcellular localization of OsLG3 in tobacco

epidermal cells. The upper panels show the localization of OsLG3-GFP in onion cells in a transient

assay, while bottom panels show the localization of GFP as a control.

Figure 4. Expression analysis of *OsLG3*. **(a)** Expression level of *OsLG3* under various abiotic stresses

and hormone treatments in IRAT109. Three-week-old seedlings were subjected to dehydration, NaCl

(200 mmol), PEG 6000 (20%, w/v), cold (4 ℃), H_2O_2 (1 mM), ABA (100 μM), ETH (100 μM) and

GA (100 μM) treatment. The relative expression level of *OsLG3* was detected by qRT-PCR at the

indicated times. Error bars indicated the SE based on three replicates. **(b)** Detection of *OsLG3*

expression in various tissues and organs of IRAT109 using qRT-PCR. Three-week-old seedlings were

used to harvest the samples of the root, sheath and leaf at seedling stage. Plants before heading stage

were used to harvest the samples of root, stem, sheath, leaf and panicle at the reproductive growth

stage. Error bars indicate the SE based on three technical replicates.

Figure 5. *OsLG3* increases rice survival under severe drought stress. **(a)** Physiological dehydration

stress tolerance assay; *OsLG3*-OE plants subjected to 20% PEG for 3 days before being allowed to

recover for 10 days. **(b)** Physiological dehydration stress tolerance assay of *OsLG3*-RNAi plants

subjected to 20% PEG for 2.5 days before being allowed to recover for 10 days. **(c)** and **(d)**, Survival

rates of transgenic and WT plants testing in (**a**) and (**b**). Values are means ±SE (n = 3). (**e**) *OsLG3*-OE

and WT plants were subjected to severe drought stress without water for 7 days and then recovered for

10 days. (**f**) *OsLG3*-RNAi and WT plants were subjected to moderate drought stress without water for

6 days and then recovered for 10 days. (**g**) and (**h**), Survival rates of transgenic and WT plants testing

in (**c**) and (**d**). Values are means ±SE (n = 3). * and ** indicate significant difference at *P* < 0.05 and

P < 0.01, respectively.

Figure 6. Growth of *OsLG3* transgenic plants under high salinity and osmotic stress conditions.

Ten day old seedlings of WT, *OsLG3*-OE and RNAi plants grown in $^1/_2$ MS medium containing (**a**)

150 mmol/L NaCl or (**b**) 200 mmol/L mannitol. The relative shoot length (**c**) and relative fresh weight

(**d**) of the transgenic and WT plants from (**a**) and (**b**) were compared. Values are means ±SE (n = 10).

* and ** indicate significant differences at *P* < 0.05 and *P* < 0.01, respectively.

Figure 7. DGE analysis shows that altering *OsLG3* expression in transgenic plants affects

transcription of stress response genes. (**a**) Scatterplots comparing the transcriptome of *OsLG3*-OE

and -RNAi with WT. The red and green dots indicate transcripts from *OsLG3*-OE or -RNAi which

have signal ratios compared to WT of greater than 2 and less than 0.5, respectively. (**b**) Venn diagram

showing the numbers of up-regulated and down-regulated genes affected by the overexpression and

suppression of *OsLG3*. (**c**) Significantly enriched GO terms show representative biological processes

of up-regulated and down-regulated genes identified in *OsLG3*-OE and RNAi plants, respectively. (**d**)

Transcript levels of genes related to ROS scavenging of WT, *OsLG3*-OE and RNAi plants under

normal or drought stress conditions (withholding water for 5 days). Values are means ±SE (n = 3).

Figure 8. *OsLG3* is involved in oxidative stress response. (**a**) Enhanced tolerance of *OsLG3*-OE

plants and enhanced sensitivity of *OsLG3*-RNAi plants to oxidative stress caused by methyl viologen

(MV). Relative shoot length (**b**) and relative fresh weight (**c**) measurements of WT, *OsLG3*-OE and

OsLG3-RNAi seedlings under oxidative stress treatments. Data are the mean ±SE (n=10). * and **

indicate significant differences at *P* < 0.05 and *P* < 0.01, respectively. (**d**) DAB and NBT staining of

leaves for H_2O_2 in WT, *OsLG3*-OE and RNAi seedlings under oxidative stress treatments caused by

H_2O_2 (100 mM) and MV (30 μM) stress treatment. (**e**) DAB staining of leaves for H_2O_2 from WT,

OsLG3-OE and RNAi seedlings under normal conditions and stress treatment (three-week-old

seedlings were treated with dehydration for 6 hours and 20% PEG6000 for 24 hours). (**f**) H_2O_2 content in leaves from WT, *OsLG3*-OE and RNAi seedlings under normal conditions and slight drought stress treatment (withholding water for 5 days). Relative MDA content (**g**), SOD activity (**h**) and POD activity (**i**) in leaves from WT, *OsLG3*-OE and RNAi seedlings under normal and drought conditions (withholding water for 5 days). Data are the mean $\pm$ SE (n=3). * and ** indicate significant differences at $P < 0.05$ and $P < 0.01$, respectively.

Figure 9. The favorable allele of *OsLG3* improves drought tolerance in rice. (**a**) Phylogenetic analysis of the *OsLG3* gene based on 45 SNPs in 1058 accessions. (**b**) Phylogenetic tree of the *OsLG3* in 1058 varieties was constructed based on seven SNPs (SNP_4352414, SNP_4352792, SNP_4352797, SNP_4352886, SNP_4352960, SNP_4353103 and SNP_4353347). Different colors reflect the different subgroups. The pink and green strips represent *indica* and *japonica*, respectively. Both purple strip and red lines represent wild rice accessions. The blue and gold lines indicate upland rice and lowland rice, respectively. (**c, d**) Haplotype analysis of *OsLG3*. Natural variation in *OsLG3* among 1058 rice accessions of a worldwide rice collection. S1-7 denote SNP_4352414, SNP_4352792, SNP_4352797, SNP_4352886, SNP_4352960, SNP_4353103 and SNP_4353347, respectively. Hap., haplotype; No., number of cultivated varieties; *Japonica*_UR, upland rice in *japonica*; *Japonica*_LR, lowland rice in *japonica*; *Indica*_UR, upland rice in *indica*; *Indica*_LR, lowland rice in *indica*. (**e**) Haplotypes of *OsLG3* in IL342, IL381, IRAT109 and Yuefu rice genotypes. ATG is the site of the start codon. SNP4 is incomplete LD with other six polymorphisms in the three drought tolerant lines. These polymorphisms are shaded in red. (**f**) Relative germination rate of IL342, IL381, IRAT109 and Yuefu rice genotypes. Data represent the mean of triplicates (*$P < 0.05$, ** $P < 0.01$). (**g, h**) The pedigree of selected rice varieties. (**g**) Huhan 3 and (**h**) Zhenghan 9. The red star refers the beneficial allele of *OsLG3*.

Figure S1. Frequency distribution of relative germination rates in the mini core collection (MCC population).

Figure S2. Sequence analysis of OsLG3. (**A**) **Schematic** diagram of OsLG3 gene structure. Representation of OsLG3 protein domain structure showing the location of the AP2 domain and the DNA-binding sites (indicated by shaded triangles). (**B**) Multiple sequence alignment of OsLG3 with

several previously reported stress-related ERF genes in rice with DNAMAN software. Accession numbers are as follows: OsERF71: *Oryza sativa*, LOC_Os06g09390; OsBIERF1: *Oryza sativa*, LOC_Os09g26420; Sub1A: *Oryza sativa*, DQ011598; OsEREBP1: *Oryza sativa*, LOC_Os02g54160; AtEBP: *Arabidopsis thaliana*, AT3G16770; CaPF1: *Capsicum annuum*, AAP72289; GmEREBP1: *Glycine max*, NP_001236578; JERF1: *Solanum lycopersicum*, NP_001234513. JERF3: *Solanum lycopersicum*, AAQ91334. Red line, black line and red box represent MCGGAIL/I motif, nuclear localization signal and conserved AP2 domain, respectively. **(c)** Sequence logos of the AP2 domain were produced based on the sequences presented in (b). Height of letter (amino acid) at each position indicates degree of conservation.

Figure S3. Overexpression of *OsLG3*. (A) *OsLG3* overexpression construct used for rice transformation. LB, left border; HPT, *Hygromycin phosphotransferase*; 35S, *cauliflower mosaic virus 35S* promoter; RB, right border. **(B)** The expression level of *OsLG3* in WT and OE lines analyzed by qRT-PCR. The lines used for further analysis were labeled in black star. **(c)** Phenotype of *OsLG3*-OE and WT plants under normal growth conditions.

Figure S4. Suppression of *OsLG3* by RNAi. (A) *OsLG3*-RNAi construct used for rice transformation. **(B)** The expression level of *OsLG3* in WT and RNAi lines analyzed by qRT-PCR. The lines used for further analysis were labeled in black star. **(C)** Phenotype of WT and *OsLG3*-RNAi plants under normal growth conditions at seedling stage.

Figure S5. Water loss from detached leaves and expression of drought stress-related genes in WT, *OsLG3*-OE and RNAi plants under normal and drought stress conditions. (A) Water loss from detached leaves of WT, *OsLG3*-OE and -RNAi plants at indicated time points. Data are the mean $\pm$SE (n=3). * and ** indicate significant differences at $P < 0.05$ and $P < 0.01$, respectively. **(B)** Expression of drought stress responsive genes in WT, *OsLG3*-OE and RNAi under normal and drought stress conditions (without water for 5 days). Values are means $\pm$SE (n=3).

Figure S6. qRT-PCR validation of differentially expressed genes in *OsLG3*-OE (A) and *OsLG3*-RNAi transgenic plants (B) compared to WT Plants.

Table S1. Information of 173 MCC varieties used for association analysis

Table S2. Primer sequences used in this study.

Table S3. GO_biological_process analysis of the differentially expressed genes.

Data Set S1. Information of 1058 *Oryza sativa* varieties and wild rice used for haplotype analysis.

Data Set S2. Differentially expressed genes in *OsLG3*-OE Plants Compared to WT Plants.

Data Set S3. Differentially expressed genes in *OsLG3*-RNAi Plants Compared to WT Plants.

REFERENCES

Adam N. Famoso KZ, Randy T. Clark, Chih-Wei Tung, Mark H. Wright, Carlos Bustamante, Leon V.
Kochian, Susan R. McCouch (2011) Genetic Architecture of Aluminum Tolerance in Rice (Oryza
sativa) Determined through Genome-Wide Association Analysis and QTL Mapping. Plos Genetics **7:**
747-757

Ambavaram MM, Basu S, Krishnan A, Ramegowda V, Batlang U, Rahman L, Baisakh N, Pereira A (2014)
Coordinated regulation of photosynthesis in rice increases yield and tolerance to environmental stress.
Nat Commun **5:** 5302

Apel K, Hirt H (2004) Reactive oxygen species: metabolism, oxidative stress, and signal transduction. Annual
Review of Plant Biology **55:** 373-399

Boutilier K, Offringa R, Sharma VK, Kieft H, Ouellet T, Zhang L, Hattori J, Liu C-M, van Lammeren
AAM, Miki BLA, Custers JBM, van Lookeren Campagne MM (2002) Ectopic expression of BABY
BOOM triggers a conversion from vegetative to embryonic growth. The Plant Cell **14:** 1737-1749

Cao Y, Song F, Goodman RM, Zheng Z (2006) Molecular characterization of four rice genes encoding
ethylene-responsive transcriptional factors and their expressions in response to biotic and abiotic stress.
J Plant Physiol **163:** 1167-1178

Cheah BH, Nadarajah K, Divate MD, Wickneswari R (2015) Identification of four functionally important
microRNA families with contrasting differential expression profiles between drought-tolerant and
susceptible rice leaf at vegetative stage. BMC Genomics **16:** 692

Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of
Arabidopsis thaliana. The Plant Journal **16:** 735-743

Curtis MD, Grossniklaus U (2003) A Gateway Cloning Vector Set for High-Throughput Functional Analysis

of Genes in Planta. Plant Physiology **133:** 462-469

Ding X, Li X, Xiong L (2013) Insight into differential responses of upland and paddy rice to drought stress by

comparative expression profiling analysis. International Journal of Molecular Sciences **14:** 5214

Du Z, Zhou X, Ling Y, Zhang Z, Su Z (2010) agriGO: a GO analysis toolkit for the agricultural community.

Nucleic Acids Res **38:** W64-70

Duan J, Zhang M, Zhang H, Xiong H, Liu P, Ali J, Li J, Li Z (2012) OsMIOX, a myo-inositol oxygenase

gene, improves drought tolerance through scavenging of reactive oxygen species in rice (Oryza sativa

L.). Plant Science **196:** 143-151

Fang Y, Liao K, Du H, Xu Y, Song H, Li X, Xiong L (2015) A stress-responsive NAC transcription factor

SNAC3 confers heat and drought tolerance through modulation of reactive oxygen species in rice.

Journal of Experimental Botany **66:** 6803-6817

Fukao T, Xiong L (2013) Genetic mechanisms conferring adaptation to submergence and drought in rice:

simple or complex? Current Opinion in Plant Biology **16:** 196-204

Fukao T, Yeung E, Bailey-Serres J (2011) The Submergence Tolerance Regulator SUB1A Mediates Crosstalk

between Submergence and Drought Tolerance in Rice. The Plant Cell **23:** 412-427

Haake V, Cook D, Riechmann J, Pineda O, Thomashow MF, Zhang JZ (2002) Transcription Factor CBF4

Is a Regulator of Drought Adaptation in Arabidopsis. Plant Physiology **130:** 639-648

Hirayama T, Shinozaki K (2010) Research on plant abiotic stress responses in the post-genome era: past,

present and future. Plant J **61:** 1041-1052

Hofmann NR (2012) SHAT1, a new player in seed shattering of rice. The Plant Cell **24:** 839

Hou X, Xie K, Yao J, Qi Z, Xiong L (2009) A homolog of human ski-interacting protein in rice positively

regulates cell viability and stress tolerance. Proc Natl Acad Sci U S A **106:** 6410-6415

Hu Y, Chong K, Wang T (2008) OsRAF is an ethylene responsive and root abundant factor gene of rice. Plant

Growth Regulation **54:** 55-61

Huang X, Wei X, Sang T, Zhao Q, Feng Q, Zhao Y, Li C, Zhu C, Lu T, Zhang Z, Li M, Fan D, Guo Y,

Wang A, Wang L, Deng L, Li W, Lu Y, Weng Q, Liu K, Huang T, Zhou T, Jing Y, Li W, Lin Z,

Buckler ES, Qian Q, Zhang QF, Li J, Han B (2010) Genome-wide association studies of 14

agronomic traits in rice landraces. Nat Genet **42:** 961-967

Huang X, Zhao Y, Wei X, Li C, Wang A, Zhao Q, Li W, Guo Y, Deng L, Zhu C (2012) Genome-wide

774  association study of flowering time and grain yield traits in a worldwide collection of rice germplasm.

775  Nature Genetics **44:** 32-39

776 **Huke RE, Huke EH** (1997) Rice area by type of culture: South, Southeast, and East Asia. A review and updated

777  data base. Rice Area by Type of Culture South Southeast & East Asia A Review & Updated Data Base

778 **Iwase A, Mitsuda N, Koyama T, Hiratsu K, Kojima M, Arai T, Inoue Y, Seki M, Sakakibara H, Sugimoto**

779  **K, Ohme-Takagi M** (2011) The AP2/ERF transcription factor WIND1 controls cell dedifferentiation

780  in Arabidopsis. Current Biology **21:** 508-514

781 **Jisha V, Dampanaboina L, Vadassery J, Mithöfer A, Kappara S, Ramanan R** (2015) Overexpression of an

782  AP2/ERF type transcription factor OsEREBP1 confers biotic and abiotic stress tolerance in rice. PLOS

783  ONE **10:** e0127831

784 **Komatsu M, Chujo A, Nagato Y, Shimamoto K, Kyozuka J** (2003) FRIZZY PANICLE is required to prevent

785  the formation of axillary meristems and to establish floral meristem identity in rice spikelets.

786  Development **130:** 3841-3850

787 **Kumar A, Dixit S, Ram T, Yadaw RB, Mishra KK, Mandal NP** (2014) Breeding high-yielding drought-

788  tolerant rice: genetic variations and conventional and molecular approaches. Journal of Experimental

789  Botany **65:** 6265-6278

790 **Lee D-K, Jung H, Jang G, Jeong JS, Kim YS, Ha S-H, Choi YD, Kim J-K** (2016) Overexpression of the

791  OsERF71 transcription factor Alters Rice Root Structure and Drought Resistance. Plant Physiology

792 **Lee DK, Jung H, Jang G, Jeong JS, Kim YS, Ha SH, Do Choi Y, Kim JK** (2016) Overexpression of the

793  OsERF71 Transcription Factor Alters Rice Root Structure and Drought Resistance. Plant Physiol **172:**

794  575-588

795 **Lee S, Seo PJ, Lee H-J, Park C-M** (2012) A NAC transcription factor NTL4 promotes reactive oxygen species

796  production during drought-induced leaf senescence in Arabidopsis. The Plant Journal **70:** 831-844

797 **Lenka SK, Katiyar A, Chinnusamy V, Bansal KC** (2011) Comparative analysis of drought-responsive

798  transcriptome in Indica rice genotypes with contrasting drought tolerance. Plant Biotechnology Journal

799  **9:** 315-327

800 **Li J-Y, Wang J, Zeigler RS** (2014) The 3,000 rice genomes project: new opportunities and challenges for future

801  rice research. GigaScience **3:** 8

802 **Li J, Pandeya D, Nath K, Zulfugarov IS, Yoo S-C, Zhang H, Yoo J-H, Cho S-H, Koh H-J, Kim D-S, Seo**

803  **HS, Kang B-C, Lee C-H, Paek N-C** (2010) ZEBRA-NECROSIS, a thylakoid-bound protein, is critical

for the photoprotection of developing chloroplasts during early leaf development. The Plant Journal **62:**

713-725

Li Q, Yang X, Bai G, Warburton ML, Mahuku G, Gore M, Dai J, Li J, Yan J (2010) Cloning and

characterization of a putative GS3 ortholog involved in maize kernel development. Theor Appl Genet

120: 753-763

Li Q, Yang X, Bai G, Warburton ML, Mahuku G, Gore M, Dai J, Li J, Yan J (2010) Cloning and

characterization of a putative GS3 ortholog involved in maize kernel development. Theoretical and

Applied Genetics **120:** 753-763

Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K (1998) Two

transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two

cellular signal transduction pathways in drought- and low-temperature-responsive gene expression,

respectively, in Arabidopsis. Plant Cell **10:** 1391 - 1406

Liu S, Wang X, Wang H, Xin H, Yang X, Yan J, Li J, Tran LS, Shinozaki K, Yamaguchi-Shinozaki K, Qin

F (2013) Genome-wide analysis of ZmDREB genes and their association with natural variation in

drought tolerance at seedling stage of Zea mays L. PLoS Genet **9:** e1003790

Livak KJ, Schmittgen TD (2001) Analysis of Relative Gene Expression Data Using Real-Time Quantitative

PCR and the 2−ΔΔCT Method. Methods **25:** 402-408

Lou Q, Chen L, Mei H, Wei H, Feng F, Wang P (2015) Quantitative trait locus mapping of deep rooting by

linkage and association analysis in rice. J Exp Bot **66**

Lou Q, Chen L, Mei H, Wei H, Feng F, Wang P, Xia H, Li T, Luo L (2015) Quantitative trait locus mapping

of deep rooting by linkage and association analysis in rice. J Exp Bot **66:** 4749-4757

Lu Y, Zhang S, Shah T, Xie C, Hao Z, Li X, Farkhari M, Ribaut JM, Cao M, Rong T, Xu Y (2010) Joint

linkage-linkage disequilibrium mapping is a powerful approach to detecting quantitative trait loci

underlying drought tolerance in maize. Proc Natl Acad Sci U S A **107:** 19585-19590

Lyu J, Li B, He W, Zhang S, Gou Z, Jing Z, Meng L, Xin L, Tao D, Huang W (2014) A genomic perspective

on the important genetic mechanisms of upland adaptation of rice. BMC Plant Biology **14:** 160

Mao H, Wang H, Liu S, Li Z, Yang X, Yan J, Li J, Tran LS, Qin F (2015) A transposable element in a NAC

gene is associated with drought tolerance in maize seedlings. Nat Commun **6:** 8326

Miller GAD, Suzuki N, Ciftci-Yilmaz S, Mittler RON (2010) Reactive oxygen species homeostasis and

signalling during drought and salinity stresses. Plant, Cell & Environment **33:** 453-467

Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. Trends in Plant Science **7:** 405-410

Mohanty S, Wassmann R, Nelson A, Moya P, Jagadish S (2013) Rice and climate change: significance for

food security and vulnerability. IRRI Discussion Paper Series No. 49. Los Baños (Philippines):

International Rice Research Institute.**:** 14p

Nakano T, Suzuki K, Fujimura T, Shinshi H (2006) Genome-Wide Analysis of the ERF Gene Family in

Arabidopsis and Rice. Plant Physiology **140:** 411-432

Ning J, Li X, Hicks LM, Xiong L (2010) A Raf-like MAPKKK gene *DSM1* mediates drought resistance

through reactive oxygen species scavenging in rice. Plant Physiology **152:** 876-890

Noctor G, Foyer CH (1998) ASCORBATE AND GLUTATHIONE: Keeping Active Oxygen Under Control.

Annual Review of Plant Physiology and Plant Molecular Biology **49:** 249-279

Oh SJ, Kim YS, Kwon CW, Park HK, Jeong JS, Kim JK (2009) Overexpression of the transcription factor

AP37 in rice improves grain yield under drought conditions. Plant Physiol **150:** 1368-1379

Qin F, Shinozaki K, Yamaguchi-Shinozaki K (2011) Achievements and challenges in understanding plant

abiotic stress responses and tolerance. Plant Cell Physiol **52:** 1569-1582

Quan R, Hu S, Zhang Z, Zhang H, Zhang Z, Huang R (2010) Overexpression of an ERF transcription factor

TSRF1 improves rice drought tolerance. Plant Biotechnology Journal **8:** 476-488

Ren D, Li Y, Zhao F, Sang X, Shi J, Wang N, Guo S, Ling Y, Zhang C, Yang Z, He G (2013) MULTI-

FLORET SPIKELET1, which encodes an AP2/ERF protein, determines spikelet meristem fate and

sterile lemma identity in rice. Plant Physiology **162:** 872-884

Rong W, Qi L, Wang A, Ye X, Du L, Liang H, Xin Z, Zhang Z (2014) The ERF transcription factor TaERF3

promotes tolerance to salt and drought stresses in wheat. Plant Biotechnology Journal **12:** 468-479

Sato Y, Masuta Y, Saito K, Murayama S, Ozawa K (2011) Enhanced chilling tolerance at the booting stage

in rice by transgenic overexpression of the ascorbate peroxidase gene, OsAPXa. Plant Cell Rep **30:**

399-406

Schmidt R, Mieulet D, Hubberten H-M, Obata T, Hoefgen R, Fernie AR, Fisahn J, San Segundo B,

Guiderdoni E, Schippers JHM, Mueller-Roeber B (2013) Salt-responsive ERF1 regulates reactive

oxygen species-dependent signaling during the initial response to salt stress in rice. The Plant Cell **25:**

2115-2131

Setter TL, Yan J, Warburton M, Ribaut J-M, Xu Y, Sawkins M, Buckler ES, Zhang Z, Gore MA (2010)

Genetic association mapping identifies single nucleotide polymorphisms in genes that affect abscisic

acid levels in maize floral tissues during drought. Journal of experimental botany: erq308

Singh BP, Jayaswal PK, Singh B, Singh PK, Kumar V, Mishra S, Singh N, Panda K, Singh NK (2015)

Natural allelic diversity in OsDREB1F gene in the Indian wild rice germplasm led to ascertain its

association with drought tolerance. Plant Cell Rep **34:** 993-1004

Suntres ZE (2002) Role of antioxidants in paraquat toxicity. Toxicology **180:** 65-77

Trujillo LE, Sotolongo M, Menéndez C, Ochogavía ME, Coll Y, Hernández I, Borrás-Hidalgo O, Thomma

BPHJ, Vera P, Hernández L (2008) SodERF3, a novel sugarcane ethylene responsive factor (ERF),

enhances salt and drought tolerance when overexpressed in tobacco plants. Plant and Cell Physiology

49: 512-525

Wang H, Zhang H, Gao F, Li J, Li Z (2007) Comparison of gene expression between upland and lowland rice

cultivars under water stress using cDNA microarray. Theor Appl Genet **115:** 1109-1126

Wang X, Wang H, Liu S, Ferjani A, Li J, Yan J, Yang X, Qin F (2016) Genetic variation in ZmVPP1

contributes to drought tolerance in maize seedlings. Nat Genet **48:** 1233-1241

Wang Z, Chen C, Xu Y, Jiang R, Han Y, Xu Z, Chong K (2004) A practical vector for efficient knockdown

of gene expression in rice (*Oryza sativa* L.). Plant Molecular Biology Reporter **22:** 409-417

Wu A, Allu AD, Garapati P, Siddiqui H, Dortay H, Zanor M-I, Asensi-Fabado MA, Munné-Bosch S,

Antonio C, Tohge T, Fernie AR, Kaufmann K, Xue G-P, Mueller-Roeber B, Balazadeh S (2012)

JUNGBRUNNEN1, a reactive oxygen species-responsive NAC transcription factor, regulates longevity

in *Arabidopsis*. The Plant Cell **24:** 482-506

Wu A, Allu AD, Garapati P, Siddiqui H, Dortay H, Zanor M-I, Asensi-Fabado MA, Munné-Bosch S,

Antonio C, Tohge T, Fernie AR, Kaufmann K, Xue G-P, Mueller-Roeber B, Balazadeh S (2012)

JUNGBRUNNEN1, a reactive oxygen species–responsive NAC transcription factor, regulates longevity

in *Arabidopsis*. The Plant Cell Online **24:** 482-506

Wu L, Zhang Z, Zhang H, Wang X-C, Huang R (2008) Transcriptional modulation of ethylene response

factor protein JERF3 in the oxidative stress response enhances tolerance of tobacco seedlings to salt,

drought, and freezing. Plant Physiology **148:** 1953-1963

Xiong H, Li J, Liu P, Duan J, Zhao Y, Guo X, Li Y, Zhang H, Ali J, Li Z (2014) Overexpression of

OsMYB48-1, a novel MYB-related transcription factor, enhances drought and salinity tolerance in rice.

PLoS One **9:** e92913

Xue Y, Warburton ML, Sawkins M, Zhang X, Setter T, Xu Y, Grudloyma P, Gethi J, Ribaut JM, Li W,

894 **Zhang X, Zheng Y, Yan J** (2013) Genome-wide association analysis for nine agronomic traits in maize

895 under well-watered and water-stressed conditions. Theor Appl Genet **126:** 2587-2596

896 **Yaish MW, El-kereamy A, Zhu T, Beatty PH, Good AG, Bi Y-M, Rothstein SJ** (2010) The APETALA-2-

897 like transcription factor OsAP2-39 controls key interactions between abscisic acid and gibberellin in

898 rice. PLoS Genet **6:** e1001098

899 **Yamaguchi-Shinozaki K, Shinozaki K** (2006) Transcriptional regulatory networks in cellular responses and

900 tolerance to dehydration and cold stresses. Annu Rev Plant Biol **57:** 781 - 803

901 **Yan J, Kandianis CB, Harjes CE, Bai L, Kim EH, Yang X, Skinner DJ, Fu Z, Mitchell S, Li Q** (2010) Rare

902 genetic variation at Zea mays crtRB1 increases beta-carotene in maize grain. Nature Genetics **42:** 322

903 **Yan J, Warburton M, Crouch J** (2011) Association Mapping for Enhancing Maize (L.) Genetic Improvement.

904 Crop Science **51:** 433

905 **Ye N, Zhu G, Liu Y, Li Y, Zhang J** (2011) ABA controls H(2)O(2) accumulation through the induction of

906 OsCATB in rice leaves under water stress. Plant Cell Physiol **52:** 689-698

907 **You J, Zong W, Hu H, Li X, Xiao J, Xiong L** (2014) A STRESS-RESPONSIVE NAC1-regulated protein

908 phosphatase gene rice protein phosphatase18 modulates drought and oxidative stress tolerance through

909 abscisic acid-independent reactive oxygen species scavenging in rice. Plant Physiol **166:** 2100-2114

910 **Yu J, Xiong H, Zhu X, Zhang H, Li H, Miao J, Wang W, Tang Z, Zhang Z, Yao G, Zhang Q, Pan Y, Wang**

911 **X, Rashid MAR, Li J, Gao Y, Li Z, Yang W, Fu X, Li Z** (2017) OsLG3 contributing to rice grain

912 length and yield was mined by Ho-LAMap. BMC Biology **15:** 28

913 **Yue Y, Zhang M, Zhang J, Tian X, Duan L, Li Z** (2012) Overexpression of the AtLOS5 gene increased

914 abscisic acid level and drought tolerance in transgenic cotton. Journal of Experimental Botany **63:** 3741-

915 3748

916 **Zhang G, Chen M, Li L, Xu Z, Chen X, Guo J, Ma Y** (2009) Overexpression of the soybean GmERF3 gene,

917 an AP2/ERF type transcription factor for increased tolerances to salt, drought, and diseases in transgenic

918 tobacco. Journal of Experimental Botany **60:** 3781-3796

919 **Zhang H, Gao S, Lercher MJ, Hu S, Chen W-H** (2012) EvolView, an online tool for visualizing, annotating

920 and managing phylogenetic trees. Nucleic Acids Research **40:** W569-W572

921 **Zhang H, Jin J, Tang L, Zhao Y, Gu X, Gao G, Luo J** (2011) PlantTFDB 2.0: update and improvement of

922 the comprehensive plant transcription factor database. Nucleic Acids Res **39:** D1114-1117

923 **Zhang H, Zhang D, Wang M, Sun J, Qi Y, Li J, Wei X, Han L, Qiu Z, Tang S** (2011) A core collection and

mini core collection of Oryza sativa L. in China. Theoretical and Applied Genetics **122:** 49-61

Zhang ZF, Li YY, Xiao BZ (2016) Comparative transcriptome analysis highlights the crucial roles of

photosynthetic system in drought stress adaptation in upland rice. Scientific Reports **6**

Zhao Y, Wei T, Yin K-Q, Chen Z, Gu H, Qu L-J, Qin G (2012) Arabidopsis RAP2.2 plays an important role

in plant resistance to Botrytis cinerea and ethylene responses. New Phytologist **195:** 450-460

Zhou Y, Lu D, Li C, Luo J, Zhu B-F, Zhu J, Shangguan Y, Wang Z, Sang T, Zhou B, Han B (2012) Genetic

Control of Seed Shattering in Rice by the APETALA2 Transcription Factor SHATTERING

ABORTION1. The Plant Cell Online

Zhu D, Wu Z, Cao G, Li J, Wei J, Tsuge T, Gu H, Aoyama T, Qu L-J (2014) TRANSLUCENT GREEN, an

ERF family transcription factor, controls water balance in Arabidopsis by activating the expression of

aquaporin genes. Molecular Plant **7:** 601-615

Figure 1. *OsLG3* **is associated with drought tolerance.**

(A) qRT-PCR analysis of *OsLG3* in Nipponbare and IRAT109 under different soil drought stress levels. NS, No
stress; SLD, Slight drought; MOD, Moderate drought; SED, Severe drought. **(B)** Analysis of the association
between pairwise LD of DNA polymorphisms in the *OsLG3* gene and water deficit tolerance. Schematic of
OsLG3 is shown on the X-axis and the significance of each variation associated with seedling RGR (ratio of
germination rates under 15%PEG condition to germination rates under water conditions) is shown on the Y-
axis. The SNPs with significant variation ($P < 1.0 \times 10^{-2}$) between genotypes are connected to the pairwise LD
diagram with a solid line. Black dots in the pairwise LD diagram highlight the strong LD of SNP_4352886
(filled red circle, $P = 2.66 \times 10^{-6}$) and two significant variations, SNP_4352414 and SNP_4352960 (open red
circles, $P < 1.0 \times 10^{-3}$). The SNP_4348903, SNP_4352166, SNP_4352793, SNP_4353076 and SNP_4353119,
which are marginally significant ($P < 1.0 \times 10^{-2}$) are denoted by red triangles.

Figure 2.   Phylogenetic tree of OsLG3 homologues

Neighbour-Joining phylogenetic analysis of OsLG3 protein sequence in the context of other
characterized AP2/ERF proteins from rice. The phylogenetic tree was constructed using the ClustalW
and MEGA programs. Tree topology with bootstrap support is based on a percentage of 1000 replicates.
Accession numbers are as follows: ARAG1, LOC_Os02g43970; OsAP21, LOC_Os01g10370;
OsDREB4-1,   LOC_Os02g43940;   OsDREB4-2,   LOC_Os04g46400;   OsDREB2A,
LOC_Os01g07120; OsDREB2B, LOC_Os05g27930; OsBIERF3, LOC_Os02g43790; OsERF1,
LOC_Os04g46220; OsERF922, LOC_Os01g54890; OsWR1, LOC_Os02g10760, OsWR4,
LOC_Os06g08340; SNORKEL1, AB510478; SNORKEL2, AB510479; OsBIERF1,
LOC_Os09g26420; OsEBP-89, LOC_Os03g08460; OsEREBP1, LOC_Os02g54160; OsLG3,
LOC_Os03g08470; OsERF71, LOC_Os06g09390; Sub1A, DQ011598; Sub1C, LOC_Os09g11480;
Sub1B,   LOC_Os09g11460;   MFS1,   LOC_Os05g41760;   OsAP2-39,   LOC_Os04g52090;
OsERF3/OsBIERF4/AP37,   LOC_Os01g58420;   FZP,   LOC_Os07g47330;   OsRap2.6,
LOC_Os04g32620; OsAP2-1, LOC_Os11g03540; OsDREB1D, LOC_Os06g06970.

Figure 3.   Transactivation assay and Subcellular localization of OsLG3

(A) Transactivation assay of OsLG3 fragments. Fusion proteins of the GAL4 DNA-binding domain and different portions of OsLG3 were expressed in yeast strain AH109 and streaked on control plate (SD/-Trp) or selective plate (SD/-Trp-Ade-His/X-α-gal), respectively. FL indicates the full-length CDS of OsLG3; dC1 to dC6 indicate the mutated forms of OsLG3 (nucleotide positions are labeled in the diagrams), respectively. **(B)** Dimerization analysis. BD with AD empty vector, AD empty with BD-dC2, AD-OsLG3 with BD empty and AD-OsLG3 with BD-dC2, were co-transformed in to yeast strain AH109 yeast cells and streaked on control plate (SD/-Trp-Leu) or selective plate (SD/-Trp-Leu-Ade-His/X-α-gal), respectively. The plates were incubated for 3 days. **(C)** Subcellular localization of OsLG3 in tobacco epidermal cells. The upper panels show the localization of OsLG3-GFP in onion cells in a transient assay, while bottom panels show the localization of GFP as a control.

Figure 4. **Expression analysis of *OsLG3*.**

(A) Expression level of *OsLG3* under various abiotic stresses and hormone treatments in IRAT109.
Three-week-old seedlings were subjected to dehydration, NaCl (200 mmol), PEG 6000 (20%, w/v),
cold (4 ℃), H_2O_2 (1 mM), ABA (100 μM), ETH (100 μM) and GA (100 μM) treatment. The relative
expression level of *OsLG3* was detected by qRT-PCR at the indicated times. Error bars indicated the
SE based on three replicates. **(B)** Detection of *OsLG3* expression in various tissues and organs of
IRAT109 using qRT-PCR. Three-week-old seedlings were used to harvest the samples of the root,
sheath and leaf at seedling stage. Plants before heading stage were used to harvest the samples of root,
stem, sheath, leaf and panicle at the reproductive growth stage. Error bars indicate the SE based on
three technical replicates.

Figure 5. *OsLG3* **increases rice survival under severe drought stress**

(**A**) Physiological dehydration stress tolerance assay; *OsLG3*-OE plants subjected to 20% PEG for 3 days before being allowed to recover for 10 days. (**B**) Physiological dehydration stress tolerance assay of *OsLG3*-RNAi plants subjected to 20% PEG for 2.5 days before being allowed to recover for 10 days. (**C**) and (**D**), Survival rates of transgenic and WT plants testing in (**A**) and (**B**). Values are means ± SE (n = 3). (**E**) *OsLG3*-OE and WT plants were subjected to severe drought stress without water for 7 days and then recovered for 10 days. (**F**) *OsLG3*-RNAi and WT plants were subjected to moderate drought stress without water for 6 days and then recovered for 10 days. (**G**) and (**H**), Survival rates of transgenic and WT plants testing in (**C**) and (**D**). Values are means ± SE (n = 3). * and ** indicate significant difference at *P* < 0.05 and *P* < 0.01, respectively.

Figure 6. **Growth of *OsLG3* transgenic plants under high salinity and osmotic stress conditions.**

Ten day old seedlings of WT, *OsLG3*-OE and RNAi plants grown in $^1/_2$ MS medium containing **(A)** 150 mmol/L NaCl or **(B)** 200 mmol/L mannitol. The relative shoot length **(C)** and relative fresh weight **(D)** of the transgenic and WT plants from **(A)** and **(B)** were compared. Values are means $\pm$ SE (n = 10). * and ** indicate significant differences at $P < 0.05$ and $P < 0.01$, respectively.

1004

Figure 7. **DGE analysis shows that altering *OsLG3* expression in transgenic plants affects transcription of stress response genes.**

(**A**) Scatterplots comparing the transcriptome of *OsLG3*-OE and RNAi plants with WT. The red and green dots indicate transcripts from *OsLG3*-OE or -RNAi which have signal ratios compared to WT of greater than 2 and less than 0.5, respectively. (**B**) Venn diagram showing the numbers of up-regulated and down-regulated genes affected by the overexpression and suppression of *OsLG3*. (**C**) Significantly enriched GO terms show representative biological processes of up-regulated and down-regulated genes identified in *OsLG3*-OE and RNAi plants, respectively. (**D**) Transcript levels of genes related to ROS scavenging of WT, *OsLG3*-OE and RNAi plants under normal or drought stress conditions (withholding water for 5 days). Values are means ± SE (n = 3).

1015

Figure 8. *OsLG3* **is involved in oxidative stress response**

(**A**) Enhanced tolerance of *OsLG3*-OE plants and enhanced sensitivity of *OsLG3*-RNAi plants to
oxidative stress caused by methyl viologen (MV). Relative shoot length (**B**) and relative fresh weight
(**C**) measurements of WT, *OsLG3*-OE and *OsLG3*-RNAi seedlings under oxidative stress treatments.
Data are the mean ±SE (n=10). * and ** indicate significant differences at $P < 0.05$ and $P < 0.01$,
respectively. (**D**) DAB and NBT staining of leaves for H_2O_2 in WT, *OsLG3*-OE and RNAi seedlings
under oxidative stress treatments caused by H_2O_2 (100 mM) and MV (30 µM) stress treatment. (**E**)
DAB staining of leaves for H_2O_2 from WT, *OsLG3*-OE and RNAi seedlings under normal conditions
and stress treatment (three-week-old seedlings were treated with dehydration for 6 hours and 20%
PEG6000 for 24 hours). (**F**) H_2O_2 content in leaves from WT, *OsLG3*-OE and RNAi seedlings under
normal conditions and slight drought stress treatment (withholding water for 5 days). Relative MDA
content (**G**), SOD activity (**H**) and POD activity (**I**) in leaves from WT, *OsLG3*-OE and RNAi
seedlings under normal and drought conditions (withholding water for 5 days). Data are the mean ±
SE (n=3). * and ** indicate significant differences at $P < 0.05$ and $P < 0.01$, respectively.

1030

Figure 9. **The favorable allele of *OsLG3* improves drought tolerance in rice.**

(**A**) Phylogenetic analysis of the *OsLG3* gene based on 45 SNPs in 1058 accessions. (**B**) Phylogenetic tree of the *OsLG3* in 1058 varieties was constructed based on seven SNPs (SNP_4352414, SNP_4352792, SNP_4352797, SNP_4352886, SNP_4352960, SNP_4353103 and SNP_4353347). Different colors reflect the different subgroups. The pink and green strips represent *indica* and *japonica*, respectively. Both purple strip and red lines represent wild rice accessions. The blue and gold lines indicate upland rice and lowland rice, respectively. (**C, D**) Haplotype analysis of *OsLG3*. Natural variation in *OsLG3* among 1058 rice accessions of a worldwide rice collection. S1-7 denote SNP_4352414, SNP_4352792, SNP_4352797, SNP_4352886, SNP_4352960, SNP_4353103 and SNP_4353347, respectively. Hap., haplotype; No., number of cultivated varieties; *Japonica*_UR, upland rice in *japonica*; *Japonica*_LR, lowland rice in *japonica*; *Indica*_UR, upland rice in *indica*; *Indica*_LR, lowland rice in *indica*. (**E**) Haplotypes of *OsLG3* in IL342, IL381, IRAT109 and Yuefu rice genotypes. ATG is the site of the start codon. SNP4 is incomplete LD with other six polymorphisms in the three drought tolerant lines. These polymorphisms are shaded in red. (**F**)

1045 Relative germination rate of IL342, IL381, IRAT109 and Yuefu rice genotypes. Data represent the
1046 mean of triplicates (*$P < 0.05$, ** $P < 0.01$). **(G, H)** The pedigree of selected rice varieties. **(G)** Huhan
1047 3 and **(H)** Zhenghan 9. The red star refers the beneficial allele of *OsLG3*.