Time-resolved analyses of elemental distribution and concentration in living plants: An example

using manganese toxicity in cowpea leaves

F. Pax C. Blamey[1], David J. Paterson[2], Adam Walsh[2], Nader Afshar[2], Brigid A. McKenna[1], Miaomiao

Cheng[3], Caixian Tang[3], Neal W. Menzies[1], and Peter M. Kopittke[1]*

[1]The University of Queensland, School of Agriculture and Food Sciences, St. Lucia, Queensland

4072, Australia; [2]Australian Synchrotron ANSTO, Clayton, Victoria 3168, Australia; [3]La Trobe

University, Centre for AgriBioscience, Bundoora, Victoria 3086, Australia.

Author for correspondence:

Peter M. Kopittke

Tel: +61 7 3346 9149

Email: p.kopittke@uq.edu.au

Word count:

Summary: 191

Introduction: 931

Materials and Methods: 1902

Results: 1356

Discussion: 1140

Conclusion: 181

Acknowledgements: 68

Total word count (above sections):   **5769**

Number of figures: 6 (plus 3 in Supporting Information)

Figures in color: 1, 2, 3, 4, 5, (plus 3 in Supporting Information)

Number of tables: 0 (plus 1 in Supporting Information)

Summary

•   Knowledge of elemental distribution and concentration within plant tissues is crucial in the

understanding of almost every process that occurs within plants. However, analytical

limitations have hindered the microscopic determination of changes over time in the location

and concentration of nutrients and contaminants in living plant tissues.

•   We developed a novel method using synchrotron-based micro X-ray fluorescence (μ-XRF)

that allows for laterally-resolved, multi-element, kinetic analyses of plant leaf tissues *in vivo*.

To test the utility of this approach, we examined changes in the accumulation of Mn in

unifoliate leaves of 7-d-old cowpea (*Vigna unguiculata*) plants grown for 48 h at 0.2 and 30

μM Mn in solution.

•   Repeated μ-XRF scanning did not damage leaf tissues demonstrating the validity of the

method. Exposure to 30 μM Mn for 48 h increased the initial number of small spots of

localized high Mn and their concentration rose from 40 to 670 mg Mn kg^{-1} fresh mass.

Extension of the two-dimensional μ-XRF scans to a three-dimensional geometry provided

further assessment of Mn localization and concentration.

•   This method shows the value of synchrotron-based μ-XRF analyses for time-resolved *in vivo*

analysis of elemental dynamics in plant sciences.

Key words: cowpea, ionomics, manganese, micro X-ray fluorescence, synchrotron

Introduction

The concentration and distribution of nutrients and contaminants within plant tissues change over time in response to physiological stimuli, developmental stage, and changes in the broader external environment. To understand the underlying genetic and physiological processes influencing plant growth, it is necessary to determine the concomitant changes in the accumulation or decline of these elements within plant tissues. In this regard, ionomics is concerned with the examination of elements in plants, although measurements are normally conducted for bulk tissues (Salt *et al.*, 2008). Whilst such analyses provide valuable information, it is even more useful to determine laterally resolved concentrations of essential and non-essential elements within plant tissues (Conn & Gilliham, 2010). Besides its relevance in plant nutrition, ionomics involves the functional analysis of genes that directly and indirectly control plant development and physiology (Salt *et al.*, 2008; Takahashi *et al.*, 2009).

Various techniques exist for examining the distribution of elements within plant tissues but most techniques require extensive processing of sequential samples. Hence, they are not able to examine nutrient and contaminant changes over time in the same area of living plants. For example, conventional scanning electron microscopy coupled with energy-dispersive X-ray spectroscopy (SEM-EDS) needs an ultra-high vacuum that requires samples to be dehydrated; frozen samples may be used where a cryo-SEM-EDS is available (Cosio *et al.*, 2005). It is perhaps possible to examine living plants using environmental SEM (ESEM), but there are problems with sample size restrictions, electron beam damage, and a comparatively poor detection limit (Danilatos, 1981; McGregor & Donald, 2010). Though having excellent subcellular resolution, nanoscale secondary ion mass spectrometry (NanoSIMS) analysis also requires ultra-high vacuum (Moore *et al.*, 2014). Confocal microscopy with fluorophores is potentially of use for the kinetic analyses of living plants (Walczysko *et al.*, 2000; Babourina & Rengel, 2009) but there are limits imposed by the availability of fluorophores and uncertainty in their selectivity and cellular penetration. Laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) is also capable of examining living plants in ambient conditions (Salt *et al.*, 2008) but damage through ablation of the sample surface prevents kinetic analysis of the same tissue area. Autoradiography has been used for the study of plants since the 1920s (Hevesy, 1923), and macro-autoradiography can potentially be used for kinetic analyses of living plants but is limited by poor resolution, long exposure times, limited availability of suitable isotopes, and safety considerations (Solon *et al.*, 2010). Recently-developed radioisotope tracer techniques for *in vitro* analysis, such as a positron-emitting tracer imaging system (Tsukamoto *et al.*, 2006) and magnetic resonance imaging (Jahnke *et al.*, 2009), have overcome some limitations, but it is only possible to examine a single element at a time (Sugita *et al.*, 2016).

Synchrotron-based micro-X-ray fluorescence spectroscopy (μ-XRF) is of interest as there are no theoretical restrictions on sample size, analyses are conducted at ambient temperature and pressure

often at a resolution of $c. \leq 1$ µm with a detection limit of $c.$ 0.1 to 100 mg kg^{-1} fresh mass (FM). Leaf

tissues of living plants may be examined also (Scheckel *et al.*, 2004). Depending upon the element of

interest and beamline specifications, this technique can simultaneously generate maps for multiple

elements within the energy range of the beamline, often 2 to 25 keV allowing analysis from P to Ag at

the K-edge. Early analyses of plant tissues using µ-XRF analyses focused on dehydrated tissues of

hyperaccumulators (McNear *et al.*, 2005; McNear & Küpper, 2014). However, tissue dehydration

may result in experimental artifacts and the high concentrations used in studies of hyperaccumulators

are not relevant to many crop species. With progressive improvements in technology (specifically, the

development of more efficient fluorescent X-ray detectors), there has been increasing interest in the

analysis of hydrated tissues of non-hyperaccumulating species (Lombi *et al.*, 2011a; Blamey *et al.*,

2015; Kopittke *et al.*, 2015). For example, studies of Cu, Ni, Zn, and Mn rhizotoxicity in hydrated

cowpea (*Vigna unguiculata*) roots showed Cu located in the rhizodermis and outer cortex, Ni in the

inner cortex, and Zn in the stele; the meristematic zone was high in both Zn and Mn (Kopittke *et al.*,

2011; Kopittke *et al.*, 2013). However, problems may still arise with radiation damage and leaf tissue

dehydration by high-energy X-rays upon repeated scanning of the same area of leaf (Lombi & Susini,

2009) that would preclude repeated analysis of living tissues.

To examine the potential of synchrotron-based µ-XRF for the analysis of living plants, we

inspected changes in element distribution in leaves of cowpea following exposure to an adequate and

a toxic level of Mn in the rooting medium. Using µ-XRF, we previously identified high Mn in

sunflower (*Helianthus annuus*) trichomes and in vacuoles of white lupin (*Lupinus albus*) as

mechanisms of tolerance (Blamey *et al.*, 2015). Such mechanisms are not present in cowpea (an

established model species in Mn toxicity studies) and soybean (*Glycine max*), making these crop

species considerably more sensitive to high Mn^{2+} in the root environment that results from the

localized accumulation of Mn in leaf tissues (Heenan & Carter, 1976; Horst, 1983). The first visible

symptom of toxicity is the appearance of small dark spots on unifoliate leaves $c.$ 4-d after exposure of

roots to 20 µM Mn (Wissemeier & Horst, 1987). Thus, the present study tested the suitability of µ-

XRF for *in situ*, multi-element, kinetic, microscopic analyses to quantify changes in elemental

distribution in leaves of living cowpea plants. We envisage that the method developed here will be of

importance across a wide range of studies for the examination of plant responses to physiological

stimuli, developmental stage, and changes in biotic and abiotic environments.

Materials and Methods

Plant growth

Cowpea (*Vigna unguiculata* L. Walp. cv. Bunya) seeds in rolled paper towels were placed in tap

water and seedlings transplanted 4 d later into 20 L of aerated nutrient solution at pH 5.6 and ionic

strength of *c.* 3 mM (Blamey *et al.*, 2015) approximating that in soil solutions (Kopittke *et al.*, 2010).

Nominal concentrations of nutrients in the basal solution were (μM): 1000 Ca, 120 NH_4^+-N, 95 Mg,

300 K, 10 Na, 6 Fe, 0.2 Mn, 0.5 Zn, 0.2 Cu, 1250 Cl, 670 NO_3^- -N, 340 S, 5 P, 1 B, and 0.01 Mo.

After 7 d in a controlled environment room at 25 °C under fluorescent lights, plants were transferred

to the Australian Synchrotron and grown in fresh nutrient solutions at 22 °C under high-pressure

sodium lights at photosynthetically active radiation (PAR) of 1,500 μmol m^{-2} s^{-1}. Mean

concentrations of selected nutrients measured by inductively coupled optical emission spectroscopy

(ICP-OES) in solutions at the beginning and end of the initial and final periods were (μM): 1,050 Ca,

120 Mg, 360 K, 30 Na, 0.23 Mn, 10 Fe, 0.4 Zn, 0.2 Cu, 360 S, 7 P, and 4 B. Use of 15 % NH_4^+-N in

solution ensured that the initial solution pH 5.6 did not require adjustment.

A preliminary experiment tested if repeated μ-XRF scans (see below) of the same area of leaf

caused damage that would potentially modify the transport of solutes in the xylem and phloem or

cause other experimental artifacts (e.g. dehydration). This involved excision of a unifoliate leaf

followed by mounting between two layers of 4-μm Ultralene® film on a specimen holder that

provided support and limited dehydration (Fig. 1a).

The main experiment involved the transfer of four plants, each to a 50-mL polypropylene

centrifuge tube (30 × 115 mm) with roots submerged in 40 mL of basal nutrient solution. Securing the

plant stem in the neck of the tube by rolling a length of 15-mm wide geotextile covered by a slotted

cap and Parafilm M® prevented spillage onto sensitive equipment. Also secured in the neck of the

centrifuge tube were two 2-m lengths of PTFE #2 AWG thin-wall tubing (Cole-Parmer Instrument

Company, Vernon Hills IL, USA), one serving as solution input and the other as output. Tygon®

tubing with 0.7-mm internal diameter joined the PTFE tubing where necessary. The input tube

connected a 10-L reservoir of continuously aerated nutrient solution via a four-roller peristaltic pump

to the bottom of the solution in the centrifuge tube. The output tube extended from the 40-mL mark of

the centrifuge tube to the reservoir via the peristaltic pump, adjustment of which ensured no solution

overspill. Circulation of nutrient solution between the reservoir and the centrifuge tube at a measured

rate of 4 mL min^{-1} enabled 90 % renewal of the 40 mL of nutrient solution in < 30 min. Two purpose-

built sample holders, each with two centrifuge tubes, contained plants fixed to each holder. Securing a

unifoliate leaf of each plant with Ultralene® and double-sided tape protected the sensitive Be detector

window *c.* 1 mm away but also allowed transpiration via both the adaxial and abaxial leaf surfaces

(Fig. 2a). Each sample holder (i.e. replicate) with two secured plants was attached in turn to the

sample stage that allowed horizontal and vertical adjustment of the leaves and correct focusing of the

X-ray beam (Fig. 2b). With solution in each centrifuge tube connected to a 10 L-reservoir, we

imposed two treatments with nominal 0.2 (i.e. basal) and 30 μM Mn. Mean measured values at the

start and end of the 48-h experimental period were 0.33 and 30 μM Mn.

An important objective of the main experiment was ensuring optimal plant growth conditions

other than in the 30 μM Mn treatment. We had initially intended to renew the 40 mL of solution in

each centrifuge tube at appropriate intervals, but an anticipated rapid decrease in dissolved O_2

precluded this approach. Specifically, we calculated that a 7-d-old root system (2.5 g FM plant^{-1}) with

an O_2 consumption rate of 20 µmol g^{-1} FM h^{-1} (Bravo & Uribe, 1981) would deplete the 510 µmol

dissolved O_2 in 40 mL solution at 25 °C in < 30 min. Thus, root O_2 and nutrient requirements over 48

h were met by circulating 10 L of aerated solution. Use of a peristaltic pump and reservoirs placed on

the floor proved satisfactory provided the outflow rate exceeded that of the inflow to prevent overflow

of the 50-mL centrifuge tube. This ensured the safety of sensitive electronic equipment that had

previously been avoided through use of solid growth medium (Scheckel *et al.*, 2004) or a small

reservoir of solution with *Arabidopsis thaliana* plants (Fittschen *et al.*, 2017). Additionally, we had to

prevent damage to the fragile Be window of the Maia detector despite the need for unifoliate leaves

being at a distance of *c.* 1 mm for correct focus (http://www.synchrotron.org.au/aussyncbeamlines/x-

ray-fluorescence-microscopy/samples). Securing the unifoliate leaves to the sample holder with

Ultralene$^{®}$ and double-sided tape (Fig. 2a) and moving the other leaves away from the detector (Fig. 2

b) overcame this potential problem.

180   The final experiment of this study determined the bulk concentration of Mn in unifoliate

leaves of cowpea plants grown in a laboratory under high-pressure sodium lights at 25 °C using the

same procedure as above. Four plants in each of six 20-L pots were grown in basal nutrient solution

for 7 d after transplanting at which time three replicates of 0.2 and 30 µM Mn were imposed for 2 d.

Mean concentrations of selected nutrients in solution measured by ICP-OES were (µM): 820 Ca, 85

Mg, 50 K, 20 Na, 10 Fe, 0.28 Zn, 0.12 Cu, 150 S, 10 P, and 2.7 B. The mean concentration of Mn was

0.23 and 28 µM in the two treatments at the start and end of 2 d. Four unifoliate leaves were harvested

from each pot, FM and dry mass (DM) determined, and leaves digested in 5:1 HNO_3:$HClO_4$ prior to

Mn analysis using ICP-OES.

µ-XRF scans

192   Details of the XFM beamline at the Australian Synchrotron have been provided by Paterson

et al. (2011), as have details for the analysis of fresh hydrated roots by Kopittke *et al.* (2011) and

leaves by Blamey *et al.* (2015). X-rays, at an energy of 12.9 keV in the present study, were selected

using a Si(111) monochromator and focused (2 × 2 µm) by a pair of Kirkpatrick-Baez mirrors and the

X-ray fluorescence emitted by the specimen collected in a backscatter geometry using a 384-element

Maia detector system (Lombi *et al.*, 2011a; Paterson *et al.*, 2011). Elemental mapping is conducted

on-the-fly in the horizontal direction. Until recently, discrete steps in the vertical direction increased

the dwell at the edge of the scanned area (i.e. at the end of each horizontal line scan) with greater

potential for X-ray damage to hydrated tissues. A recent improvement allows significantly faster and

more accurate raster scanning via kinematically optimized fly-scans known as Rascan, a system that

wraps the motion controller into an optimal two-dimension scanner. This device renders a motion

trajectory optimized for smoothness, as well as minimal overhead times, for a given fly-scan of

known transit time (dwell) and pitch. Importantly, Rascan reduces end-of-line overheads from > 350

ms to *c*. 35 ms resulting in a *c*. 30 % reduction in scan time and less radiation damage. Using this

approach, analyses are routinely conducted with a dwell of ≤ 1 ms pixel^{-1}, comparing favorably to

many other synchrotron-based μ-XRF beamlines where a dwell of 10 to 100 ms pixel^{-1} is common

(Lombi *et al.*, 2011b). This decreased dwell using the Maia detector system and Rascan at the

Australian Synchrotron facilitated the present study with repeated scanning of the same area of living

plant leaf tissues.

The preliminary experiment determined if sequential scanning of the same area resulted in

damage from high energy X-rays. The survey scan started within 5 min of excising and mounting a

unifoliate leaf (Fig. 1a) and took 52 min to complete an area of 47 × 33 mm with a step size (i.e.

virtual pixel size) of 50 μm. From this survey scan, a 7 × 3 mm area was selected for further

examination with three detailed scans with a step size of 5 μm, each scan taking *c*. 20 min. Each of

the detailed scans covered a slightly larger area encompassing the entire previous scan to test if the X-

rays damaged the leaf that would be evident in the subsequent scans by changes in sample hydration

and elemental redistribution. A fourth, high-resolution scan with a step size of 2 μm covering an area

of 1.54 × 0.79 mm tested the possibility of damage in detail. We also used light microscopy to

examine the leaf for any visible signs of damage.

Thereafter, the main experiment involved securing the sample holder with two centrifuge

tubes, each with a living plant (Fig. 2), in the beamline and verifying basal solution (0.2 μM Mn) flow

rate. Addition of a 0.46-mL aliquot of 0.65 M $MnSO_4$ stock solution to one of the 10-L reservoirs

imposed a 30-μM Mn treatment. Two initial survey scans of 6 and 2 min identified the area for

detailed scans (Fig. 2c), the first of which started immediately thereafter (0 h). This and subsequent

detailed scans took *c*. 120 min to complete. There were six scans from 0-2, 6-8, 12-14, 18-20, 24-26,

and 48-50 h after first imposition of Mn treatments. The same procedures and scans of adjacent

unifoliate leaves followed with living plants in the second replicate. In the replicate reported here, the

detailed scan area was *c*. 120 mm^2, which included a leaf area of 55.2 and 56.4 mm^2 in the 0.2 and 30

μM Mn treatments. The detailed scans had 4 × 4 μm pixels and a velocity of 4 mm s^{-1} resulting in a

pixel transit time of 1 ms. With a total photon flux of *c*. 2 × 10^9 photon s^{-1} at an energy of 12.9 keV,

these scan parameters corresponded to *c*. 1.3 × 10^5 photon μm^{-2} for each scan (i.e. a total of 8.0 × 10^5

photon μm^{-2} for the six scans).

On completion of the scans, the CSIRO Dynamic Analysis method in GeoPIXE provided

quantitative, true-element images of X-ray fluorescence spectra (Ryan & Jamieson, 1993; Ryan,

2000) as outlined at http://www.nmp.csiro.au/dynamic.html. The survey scans required correction for

variation in leaf thickness arising from the major leaf veins by normalizing to Compton scatter.

Sections of leaves for detailed scans had only a few minor veins (Fig. 2c) that did not interfere greatly

with determination of the two-dimensional areal concentrations of elements and their distributions

(Kopittke *et al.*, 2011).

GeoPIXE provided initial quantitative analysis of Mn concentration in 2.6 × 0.08 mm

transects of detailed scans across the leaf at 30 µM Mn from 0 to 48 h that involved 650 individual

pixels in the horizontal direction × the mean of 20 pixels in the vertical direction. This was followed

by determining Mn in the high-resolution image area of 2.6 × 0.6 mm (i.e. values of individual 650 ×

150 pixels). Further analysis using GeoPIXE and ImageJ 1.48v (Schneider *et al.*, 2012) extended the

two-dimension µ-XRF scans to a three-dimensional geometry of Mn distribution and concentration.

The number and localized areas of Mn accumulation and the distributions of K, Ca, Fe, Cu, and Zn at

0 and 48 h in the two Mn treatments were determined also.

Results

Potential X-ray damage to living unifoliate leaves

The Australian Synchrotron XFM beamline Maia detector system with fast data acquisition has the

advantage of rapid pixel transit times (Paterson *et al.*, 2011) but sequential *in situ* µ-XRF analyses of

hydrated leaf tissues may still result in radiation damage and tissue dehydration as shown in micro-

tomography of a cowpea root (Lombi *et al.*, 2011a). This was evident also in an unpublished study

using X-ray absorption near edge structure (XANES) imaging of a 4.0 × 2.5 mm section of a soybean

leaf at high Mn with repeated scans of the same area at increasing energy. Compared to the rest of the

leaf, the rectangular scanned area was mildly chlorotic suggestive of radiation damage (Fig. S1a).

Furthermore, there were changes in the distribution of Ca (Fig. S1b) and dehydration damage evident

in Compton scatter (Fig. S1c) though not in Mn distribution (data not presented). Although mapping

was conducted on-the-fly in the horizontal direction with a dwell of 1 ms per virtual 0.1-mm step,

damage occurred in a previous study at ends of the horizontal scan lines because of the increased

dwell of discrete steps in the vertical direction. In contrast to the effects observed in this previous

XANES imaging study, the preliminary experiment in the present study produced no evidence of

radiation damage or dehydration to a cowpea leaf after scanning for *c*. 2 h (Fig. 1). This was attributed

to the newly-implemented Rascan system which markedly reduced overheads, and hence reduced

dwell at the end of the horizontal lines during the vertical move to the next line. We therefore

concluded that rapid scanning prevented leaf damage suggesting that experimental artifacts would not

arise with time-resolved µ-XRF scans of living leaf tissues.

Changes in elemental distribution and concentration

There were no visible symptoms of Mn toxicity on leaves of plants grown for 7 d with the basal 0.2

μM Mn in solution (Fig. 1a) but there were a few localized spots of Mn accumulation visible in the μ-

XRF scans of the detached unifoliate leaf during the preliminary experiment (Fig. 1b,c). This was

evident also at 0 h in both living cowpea unifoliate leaves (Fig. 3) but it was only after 2 d at 30 μM

Mn that dark spots indicative of Mn accumulation were visible (Fig. 2c).

Visual assessment of the detailed μ-XRF scans (Fig. 3) indicated a slight increase over time in

the number of high-Mn spots at 0.2 μM Mn, but this increase was considerably lower than that with

30 μM Mn in solution. Given that each detailed scan was *c*. 8.5 megapixels, we selected a small area

of the image of 1.56 mm^2 (2.6 × 0.6 mm) to examine changes in elemental distribution in the 30-μM

Mn treatment over the 48-h experimental period. At the start of the experiment, high Mn spots were

visible to the left of the image, the largest being *c*. 240 × 130 μm in size (Fig. 4). Elsewhere, Mn was

minimally above background. There was little change in Mn distribution after 6 h at 30 μM Mn, with

some new high Mn spots evident after 12 h and especially from 18 to 48 h. Transects across the

images from 0 to 48 h (Fig. 4) showed no visible increase in the number or concentration of high Mn

spots at 6 h, a few instances where new high Mn spots were visible at 12 h. It was only from 18 to 48

h, however, that there were clear increases in spot numbers and in their Mn concentration.

As demonstrated by Scheckel *et al.* (2004), ImageJ analysis of the high-resolution data at 30

μM Mn (Fig. 4) extended the two-dimensional μ-XRF scans to a three-dimensional geometry

providing visual images of both Mn distribution and concentration over the 48-h experimental period

(Fig. 5). As with the two-dimensional images (Figs 3, 4), there were a few spots of high Mn at the

start of the experiment with no discernable increase at 6 h. It appeared that the number and

concentration of Mn in these spots started to increase at 12 h followed by a further increase at 18 h

and marked increases at 24 h and 48 h.

Using the detailed scan areas of 55.2 and 56.4 mm^2 at 0.2 and 30 μM Mn (Fig. 3), we used

ImageJ to calculate the number of localized Mn spots at 0 h was 3.6 and 12.1 mm^{-2} in the two leaves

to be subjected to the 0.2 and 30 μM Mn treatments (Fig. 6a). The number of high-Mn spots appeared

unchanged over the first 6 h in both treatments. There was a slight increase over 48 h of 4 mm^{-2} at 0.2

μM Mn compared to the large increase of 30 mm^{-2} at 30 μM Mn. The corresponding increases in the

total area of localized high-Mn spots were < 0.001 and 0.04 mm^2 mm^{-2} (Fig. 6b). Further analyses

using ImageJ determined that most of the initial high Mn spots were < 320 μm^2 in size (Fig. S2),

largely remaining so over 48 h at 0.2 μM Mn. This contrasted to the changes in spot size in the 30

μM-Mn treatment in which the percentage of spots < 64 μm^2 decreased over 48 h. Interestingly, the

percentage of slightly larger spots of 64 to < 320 μm increased and then decreased and there was a

concomitant increase in large spots at 48 h.

With the μ-XRF scans providing quantitative, true-elemental data (Ryan & Jamieson, 1993;

Ryan, 2000), GeoPIXE analysis of the detailed μ-XRF scan of leaf sections at 48 h (Fig. 3)

determined mean values of 26 and 56 mg Mn kg^{-1} FM at 0.2 and 30 μM Mn. Analysis of the six

transects resulted in a calculated mean Mn concentration that increased from 35 to 61 mg kg^{-1} FM

over 48 h exposure to 30 µM Mn (Table S1). However, these transects indicated that the background

Mn concentration was < 100 mg kg^{-1} FM other than in the relatively few instances in which the Mn

concentration approached or exceeded 1,000 mg kg^{-1} FM. Separating the data into two classes, above

and below an estimated background value of 60 mg kg^{-1} FM, permitted further examination of leaf

Mn status. Pixel clusters with values < 60 mg Mn kg^{-1} FM had a mean concentration of 24 mg Mn kg^{-1}

FM over 48 h, markedly lower than in those with values > 60 mg kg^{-1} FM that increased from 200 to

420 mg Mn kg^{-1} over 48 h (Table S1). Data from the 2.6 × 0.6 mm area used also to determine leaf

Mn concentration increased from a mean of 69 to 170 mg kg^{-1} FM over 48 h (Table S1). These

estimates differed somewhat from the the bulk Mn concentration in entire unifoliate leaves 48 h after

imposing the high Mn treatment of 11 ± 1 and 47 ± 14 mg kg^{-1} FM at 0.2 and 30 µM Mn.

(Corresponding values were 80 ± 6 and 360 ± 110 mg kg^{-1} DM.)

Finally, we utilized the µ-XRF analyses at 0 and 48 h to generate images for multiple

elements within the energy range of the beamline. In this exercise, the incident X-rays of 12.9 keV

and the Maia detector permitted investigation of the distributions and concentrations of elements

between P and Zn. (Selection of a higher incident energy would also permit analyses of heavier

elements.) The present study permitted investigation of six elements, K, Ca, Mn, Fe, Cu, and Zn, in

unifoliate leaves that were above background concentrations (Fig. S3). At 0 h, there was relatively

even distribution of K and Ca across the leaves of plants at 0.2 µM Mn, as was that of Cu though at a

concentration close to the detection limit. In contrast, Mn and Fe accumulated in localized areas

across the leaf, but these elements were not co-located. Localized areas of high Zn were present in and

adjacent to the veins. The distributions of these elements were similar 48 h later at 0.2 µM Mn but the

distribution of Mn differed between the 0.2 and 30 µM Mn treatments as is evident in Fig. 3.

Discussion

The analysis of fresh tissues exacerbates the many challenges in µ-XRF scanning of low metal and

metalloid concentrations in biological tissues (Lombi *et al.*, 2011a). These challenges arise from

tissue hydration (often > 85 %) and the generally low concentrations of trace elements in tissues that

result in the consequent long transit times required for analysis. A high (long) dwell increases the

likelihood of radiation damage and tissue dehydration, thereby resulting in experimental artifacts that

affect element distribution (Lombi *et al.*, 2011a). The benchtop (i.e. non-synchrotron based) µ-XRF

method developed by Fittschen *et al.* (2017) ensured high sensitivity and low detection limits but

required dwells of 1 s per 40 × 40 µm pixel that resulted in substantial radiation damage to a scanned

Arabidopsis thaliana leaf. Scheckel *et al*. (2004) also noted that µ-XRF analysis using a 1-s dwell of a

6-µm pixel damaged leaves of living *Iberis intermedia* plants. As with these studies, our previous

unpublished work showed radiation damage and leaf dehydration was evident in a soybean leaf

subjected to multiple scanning (Fig. S1). Recent years, however, have seen major improvements in

overcoming these problems in synchrotron-based studies. This has been important in the study of

fresh, hydrated root (Kopittke *et al.*, 2011) and leaf tissues (Blamey *et al.*, 2015). Besides the high

spatial resolution of 0.1 to 0.2 μm at the Australian Synchrotron XFM beamline (Paterson *et al.*,

2011), there are inherent advantages of the Maia detector and Rascan in decreasing dwell. This was

evident in the preliminary experiment of the present study in which there was no visible X-ray

damage to a unifoliate cowpea leaf after two survey scans and four detailed scans with a combined

period of *c*. 2 h (Fig. 1). The recently updated raster scanning adds further advantages by decreasing

the damage to sensitive plant tissues.

Progress in μ-XRF analysis of hydrated plant tissues has been extended to leaf tissues of

living plants. For example, by growing the hyperaccumulator *I. intermedia* plants at elevated Tl in

soil, Scheckel *et al.* (2004) determined that Tl accumulated mostly in the vascular tissues. Fittschen *et*

al. (2017) constructed a benchtop μ-XRF unit to examine differences in the distributions of a number

of nutrients in the vascular system, mesophyll, and trichomes of leaves of living *A. thaliana* plants.

These studies, however, did not investigate time-resolved changes in elemental distribution and

concentration and we were not able to find instances in the literature of sequential μ-XRF analysis of

living plant tissues.

In the present study, μ-XRF detailed scans using a pixel size of 4×4 μm enabled production

of high-quality images that permitted direct visual comparisons between the effects of 0.2 and 30 μM

Mn (Fig. 3) and enabled high resolution comparisons of Mn accumulation of 30 μM Mn over 48 h

(Fig. 4). A pixel size of 1×1 μm allows even better resolution where appropriate (Blamey *et al.*,

2015). As demonstrated by Scheckel *et al.* (2004) and in the present study (Fig. 5), three-dimensional

representations of the two-dimensional scans provide further visualization of elemental distribution

and concentration.

Until recently, quantitative elemental analyses have focused on bulk tissues, from whole

plants to separate analyses of roots, shoots, stems, and leaves (Salt *et al.*, 2008) but technological

developments in synchrotron-based μ-XRF, amongst other techniques, provide greater spatial

resolution of the ionome (Punshon *et al.*, 2009; van der Ent *et al.*, 2017). However, van der Ent *et al.*

(2017) urged caution in the use of μ-XRF in foliar studies using living plants because measurements

arise from different depths, from different cell types (e.g., epidermis, palisade, mesophyll, and

vasculature), and from continued plant metabolism. The first two of these potential difficulties may be

addressed by a combination of analytical techniques as shown by Blamey *et al.* (2017) who used

NanoSIMS to confirm the suspected accumulation of Mn in the apoplast using μ-XRF by analysis.

The present study has also shown that accommodation of continued plant metabolism is possible,

albeit in only one instance of Mn accumulation. High Mn accumulation in discreet locations is of

additional benefit by overcoming the overall low concentrations of many trace elements.

It has been recognized for many years that symptoms of Mn toxicity precede a decrease in

plant growth (Foy *et al.*, 1978; Weil *et al.*, 1997), information extended by Fernando *et al*. (2016)

with Mn accumulation in leaves of wheat (*Triticum aestivum*) plants well before the appearance of

visible symptoms. This was evident in leaves of cowpea exposed to a non-toxic concentration of 0.2

μM Mn (Figs 1, 3, 4, 5). Localized spots of high Mn preceded visible symptoms associated with Mn

toxicity in the present study also with new high Mn spots first visible after 12 h growth at 30 μM Mn

(Figs 4, 5) but visible dark spots appeared only after 48 h (Fig. 2c). However, the mechanism is

unknown whereby high Mn accumulates in specific localized areas not visibly associated with leaf

anatomy (e.g., close to or distant from veins). There is similar uncertainty as to why an overall

increase in Mn in leaf tissues at 30 μM Mn (Figs 4 and 5) results in localized increases in Mn but no

general increase in background Mn concentration over time.

From an ionomics viewpoint (Salt *et al.*, 2008), the present study addressed various

microscopic measures of Mn concentration in leaf tissues based on the true-elemental data of the μ-

XRF scans. After 48 h at 0.2 μM Mn, these analyses ranged from 24 to 35 mg Mn kg^{-1} FM compared

to the unifoliate leaf bulk analysis of 11 mg Mn kg^{-1} FM. The corresponding comparison was 56 to

169 mg kg^{-1} versus a bulk leaf tissue concentration of 47 mg Mn kg^{-1} FM at 30 μM Mn. While of the

same order of magnitude, these findings suggest that further research is required in the sampling for

quantitative determination of elemental concentrations in leaf tissues. Importantly, the multi-element

analyses undertaken in the present study would provide important information on interactions

between the various elements. An example relevant to the present study might use detailed μ-XRF

studies to examine the kinetics of changes in Mn distribution (Figs 4, 5) with those of Ca (Fig. S3)

given the role of Ca in callose formation associated with Mn toxicity (Wissemeier & Horst, 1987).

Additional studies may also provide critical information as to the underlying mechanisms whereby

Mn is toxic by determining changes in Ca and Mn accumulation in the apoplast and by determining

how Mn accumulates along with Ca in vacuoles of some plant species but not in others. Finally, there

is particular promise in studies such as that by Takahashi *et al*. (2009) in combining information on

the activities of metal transporter genes with multi-element μ-XRF analysis of changes in the

distribution of Fe, Mn, Cu, and Zn in germinating rice (*Oryza sativa*) seeds.

Conclusion

Development of a synchrotron-based method has shown the feasibility of using μ-XRF analysis to

determine the distribution and concentration of Mn in living unifoliate leaves of cowpea exposed to

0.2 and 30 μM Mn in solution culture. This method may be adapted to examine the accumulation and

distribution of other elements and plant species over a range in exposure times. Besides investigations

of elemental toxicities, other examples may include determining the dynamics of nutrients upon

addition to a nutrient-deficient plant, the effects of hypoxia on nutrient status, and establishing the

relationship between transpiration and nutrient accumulation in leaf tissues.

The results and implications of the present study advance the value of ionomics, the inorganic

component of the metabolome, through quantifying changes in the microscopic distribution and

concentration of elements *in vivo* (Salt *et al.*, 2008). Though focusing the present study on the kinetics

of Mn accumulation in cowpea unifoliate leaves, μ-XRF analyses may also provide multi-element

information in plant tissues, meeting an aim of ionomics to define the functional state of an organism

driven by genetic, developmental, biotic, or abiotic influences.

Acknowledgements

F.P.C.B. acknowledges assistance from the Australian Government Research Training Program and

P.M.K. acknowledges receipt of an Australian Research Council (ARC) Future Fellowship

(FT120100277). This research was undertaken on the XFM beamline (Project AS153/XFM/11040) at

the Australian Synchrotron, part of the Australian Nuclear Science and Technology Organization

(ANSTO). We thank Dr Chris Ryan (CSIRO), Dr Martin de Jonge, and Dr Daryl Howard (Australian

Synchrotron) for assistance with synchrotron-based techniques.

Author contributions

F.P.C.B. conceived the research program that was developed in collaboration with D.J.P., A.W.,

N.W.M., and P.M.K.; A.W. and N.A. developed the requisite technology; F.P.C.B., M.C., and C.T.

conducted the plant growth experiments; F.P.C.B., D.J.P., B.A.M., M.C., and P.M.K. conducted the

μ-XRF analyses at the Australian Synchrotron; F.P.C.B and P.M.K. analyzed the data; and F.P.C.B.,

D.J.P, and P.M.K. wrote the first draft of the article to which all authors contributed.

References

Babourina O, Rengel Z. 2009. Uptake of aluminium into *Arabidopsis* root cells measured by

fluorescent lifetime imaging. *Annals of Botany* **104**: 189-195.

Blamey FPC, Hernandez-Soriano MC, Cheng M, Tang C, Paterson DJ, Lombi E, Wang WH,

Scheckel KG, Kopittke PM. 2015. Synchrotron-based techniques shed light on mechanisms

of plant sensitivity and tolerance to high manganese in the root environment. *Plant*

Physiology **169**: 2006-2020.

Blamey FPC, McKenna BA, Li C, Cheng M, Tang C, Jiang H, Howard DL, Paterson DJ,

Kappen P, Wang P, *et al.* 2017. Manganese distribution and speciation help to explain the

effects of silicate and phosphate on manganese toxicity in four crop species. *New Phytologist*.

doi: 10.1111/nph.14878.

Bravo P, Uribe EG. 1981. Temperature dependence of the concentration kinetics of absoprtion of

phosphate and potassium in corn roots. *Plant Physiology* **67**: 815-819.

Conn S, Gilliham M. 2010. Comparative physiology of elemental distributions in plants. *Annals of*

Botany **105**: 1081-1102.

Cosio C, DeSantis L, Frey B, Diallo S, Keller C. 2005. Distribution of cadmium in leaves of *Thlaspi*

caerulescens. *Journal of Experimental Botany* **56**: 765-775.

Danilatos GD. 1981. The examination of fresh or living plant material in an environmental scanning

electron microscope. *Journal of Microscopy* **121**: 235-238.

Fernando DR, Moroni SJ, Scott BJ, Conyers MK, Lynch JP, Marshall AT. 2016. Temperature

and light drive manganese accumulation and stress in crops across three major plant families.

Environmental and Experimental Botany **132**: 66-79.

Fittschen UEA, Kunz HH, Hohner R, Tyssebotn IMB, Fittschen A. 2017. A new micro X-ray

fluorescence spectrometer for *in vivo* elemental analysis in plants. *X-Ray Spectrometry* **46**:

374-381.

Foy CD, Chaney RL, White MC. 1978. The physiology of metal toxicity in plants. *Annual Review*

of Plant Physiology **29**: 511-566.

Heenan DP, Carter OG. 1976. Tolerance of soybean cultivars to manganese toxicity. *Crop Science*

16: 389-391.

Hevesy G. 1923. The absorption and translocation of lead by plants. A contribution to the application

of the method of radioactive indicators in the investigation of the change of substance in

plants. *Biochemical Journal* **17**: 439-445.

Horst WJ. 1983. Factors responsible for genotypic manganese tolerance in cowpea (*Vigna*

unguiculata). *Plant and Soil* **72**: 213-218.

Jahnke S, Menzel MI, van Dusschoten D, Roeb GW, Buhler J, Minwuyelet S, Blumler P,

Temperton VM, Hombach T, Streun M, *et al.* 2009. Combined MRI-PET dissects dynamic

changes in plant structures and functions. *Plant Journal* **59**: 634-644.

Kopittke PM, Blamey FPC, Asher CJ, Menzies NW. 2010. Trace metal phytotoxicity in solution
culture: A review. *Journal of Experimental Botany* **61**: 945-954.

Kopittke PM, Lombi E, McKenna BA, Wang P, Donner E, Webb RI, Blamey FPC, de Jonge
MD, Paterson D, Howard DL*, et al.* **2013.** Distribution and speciation of Mn in hydrated
roots of cowpea at levels inhibiting root growth. *Physiologia Plantarum* **147**: 453-464.

Kopittke PM, Menzies NW, de Jonge MD, McKenna BA, Donner E, Webb RI, Paterson DJ,
Howard DL, Ryan CG, Glover CJ*, et al.* **2011.** In situ distribution and speciation of toxic
copper, nickel, and zinc in hydrated roots of cowpea. *Plant Physiology* **156**: 663-673.

Kopittke PM, Moore KL, Lombi E, Gianoncelli A, Ferguson BJ, Blamey FPC, Menzies NW,
Nicholson TM, McKenna BA, Wang P*, et al.* **2015.** Identification of the primary lesion of
toxic aluminum in plant roots. *Plant Physiology* **167**: 1402-1411.

Lombi E, de Jonge MD, Donner E, Kopittke PM, Howard DL, Kirkham R, Ryan CG, Paterson
D. 2011a. Fast X-ray fluorescence microtomography of hydrated biological samples. *PLoS*
ONE **6**: 1-5.

Lombi E, Smith E, Hansen TH, Paterson D, de Jonge MD, Howard DL, Persson DP, Husted S,
Ryan C, Schjoerring JK. 2011b. Megapixel imaging of (micro)nutrients in mature barley
grains. *Journal of Experimental Botany* **62**: 273-282.

Lombi E, Susini J. 2009. Synchrotron-based techniques for plant and soil science: opportunities,
challenges and future perspectives. *Plant and Soil* **320**: 1-35.

McGregor JE, Donald AM 2010. The application of ESEM to biological samples. In: Baker RT ed.
Electron Microscopy and Analysis Group Conference 2009, 012021.

McNear DH, Peltier E, Everhart J, Chaney RL, Sutton S, Newville M, Rivers M, Sparks DL.
2005. Application of quantitative fluorescence and absorption-edge computed
microtomography to image metal compartmentalization in *Alyssum murale*. *Environmental*
Science & Technology **39**: 2210-2218.

McNear DHJ, Küpper JV. 2014. Mechanisms of trichome-specific Mn accumulation and toxicity in
the Ni hyperaccumulator *Alyssum murale*. *Plant and Soil* **377**: 407-422.

Moore KL, Chen Y, van de Meene AML, Hughes L, Liu WJ, Geraki T, Mosselmans F,
McGrath SP, Grovenor C, Zhao FJ. 2014. Combined NanoSIMS and synchrotron X-ray
fluorescence reveal distinct cellular and subcellular distribution patterns of trace elements in
rice tissues. *New Phytologist* **201**: 104-115.

Paterson DJ, de Jonge MD, McKinlay WLJ, Starritt A, Kusel M, Ryan CG, Kirkham R,
Moorhead G, Siddons DP. 2011. The X-ray fluorescence microscopy beamline at the
Australian synchrotron. *AIP Conference Proceedings* **1365**: 219-222.

Punshon T, Guerinot ML, Lanzirotti A. 2009. Using synchrotron X-ray fluorescence microprobes
in the study of metal homeostasis in plants. *Annals of Botany* **103**: 665-672.

Ryan CG. 2000. Quantitative trace element imaging using PIXE and the nuclear microprobe.
International Journal of Imaging Systems and Technology **11**: 219-230.
Ryan CG, Jamieson DN. 1993. Dynamic analysis: on-line quantitative PIXE microanalysis and its
use in overlap-resolved elemental mapping. *Nuclear Instruments & Methods in Physics*
Research Section B-Beam Interactions with Materials and Atoms **77**: 203-214.
Salt DE, Baxter I, Lahner B. 2008. Ionomics and the study of the plant ionome. *Annual Review of*
Plant Biology **59**: 709-733.
Scheckel KG, Lombi E, Rock SA, McLaughlin MJ. 2004. In vivo synchrotron study of thallium
speciation and compartmentation in *lberis intermedia*. *Environmental Science & Technology*
38: 5095-5100.
Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis.
Nature Methods **9**: 671-675.
Solon EG, Schweitzer A, Stoeckli M, Prideaux B. 2010. Autoradiography, MALDI-MS, and SIMS-
MS imaging in pharmaceutical discovery and development. *The AAPS Journal* **12**: 11-26.
Sugita R, Kobayashi NI, Hirose A, Saito T, Iwata R, Tanoi K, Nakanishi TM. 2016. Visualization
of uptake of mineral elements and the dynamics of photosynthates in Arabidopsis by a newly
developed real-time radioisotope imaging system (RRIS). *Plant and Cell Physiology* **57**: 743-
753.
Takahashi M, Nozoye T, Kitajima N, Fukuda N, Hokura A, Terada Y, Nakai I, Ishimaru Y,
Kobayashi T, Nakanishi H*, et al.* **2009.** In vivo analysis of metal distribution and expression
of metal transporters in rice seed during germination process by microarray and X-ray
Fluorescence Imaging of Fe, Zn, Mn, and Cu. *Plant and Soil* **325**: 39-51.
Tsukamoto T, Nakanishi H, Kiyomiya S, Watanabe S, Matsuhashi S, Nishizawa NK, Mori S.
2006. ^{52}Mn translocation in barley monitored using a positron-emitting tracer imaging system.
Soil Science and Plant Nutrition **52**: 717-725.
van der Ent A, Przybyłowicz WJ, de Jonge MD, Harris HH, Ryan CG, Tylko G, Paterson DJ,
Barnabas AD, Kopittke PM, Mesjasz-Przybyłowicz J. 2017. X-ray elemental mapping
techniques for elucidating the ecophysiology of hyperaccumulator plants. *New Phytologist*.
doi: 10.1111/nph.14810.
Walczysko P, Wagner E, Albrechtova JTP. 2000. Use of co-loaded Fluo-3 and Fura Red
fluorescent indicators for studying the cytosolic Ca^{2+} concentrations distribution in living
plant tissue. *Cell Calcium* **28**: 23-32.
Weil RR, Foy CD, Coradetti CA. 1997. Influence of soil moisture regimes on subsequent soil
manganese availability and toxicity in two cotton genotypes. *Agronomy Journal* **89**: 1-8.
Wissemeier AH, Horst WJ. 1987. Callose deposition in leaves of cowpea (*Vigna unguiculata* (L.)
Walp.) as a sensitive response to high Mn supply. *Plant and Soil* **102**: 283-286.
557

Fig. 1. Images of a detached unifoliate leaf of cowpea grown for 7 d at 0.2 μM Mn after conducting a survey scan and three detailed scans. (a) Optical image of a leaf mounted between Ultralene® films. (b) Survey μ-XRF scan of Mn distribution, the white box showing the area of subsequent detailed scans. (c,d) Detailed μ-XRF scan showing the distributions of Mn and Ca. (e) Detailed Compton scatter.

Fig. 2. Optical images of cowpea unifoliate leaf arrangement in the sample holder. (a) Leaves of two plants grown at 0.2 and 30 µM Mn between Ultralene® films in the sample holder. (b) Arrangement of living plants mounted in the X-ray beam. (c) Micrograph of unifoliate leaves after the detailed µ-XRF scan at 48 h, the red box showing the area of sequential detailed scans and the white arrow a visible dark spot of Mn accumulation at 30 µM Mn.

Fig. 3. Detailed µ-XRF scans of Mn distribution in unifoliate leaves of cowpea from 0 to 48 h after initial exposure to 30 µM Mn in solution culture. The red box at 0 h shows the area analyzed at high resolution for the distribution of Mn (Fig. 4), for three-dimensional representation of Mn distribution and concentration (Fig. 5), and for ImageJ and GeoPIXE analyses (Fig. 6).

Fig. 4. High-resolution μ-XRF scans and transects of Mn distribution and concentration (log$_{10}$ scale) from 0 to 48 h after initial exposure to 30 μM Mn in a 2.6 mm × 0.6 mm section of a cowpea unifoliate leaf (Fig. 3). The horizontal dotted line at 0 h identifies the position of transects along which the concentration of Mn was determined.

Fig. 5. Three-dimensional representation of Mn distribution and concentration in unifoliate leaf sections of cowpea from 0 to 48 h after initial exposure to 30 µM Mn. The scanned area is 2.6 mm × 0.6 mm as shown by the red box in Fig. 3 and the high-resolution µ-XRF images in Fig. 4. The Mn concentration in the large high-Mn spot (back left) exceeded 10,000 mg kg^{-1} FM from 0 to 48 h but the scale of Mn in the unifoliate leaf has been limited to 1,000 mg kg^{-1} FM to show the change in high Mn spots that developed during the 48-h experimental period.

Fig. 6. Accumulation of Mn in unifoliate leaf sections (Fig. 3) of cowpea from 0 to 48 h after initial exposure to 30 μM Mn in solution culture. (a,b) ImageJ determination of the number and area of localized spots of high Mn in detailed scans of leaf sections (Fig. 3). (c) GeoPIXE determination of the Mn concentration in identified high Mn spots (i.e. > 60 mg kg^{-1} FM) over 48 h in transects of the section of a cowpea unifoliate leaf at 30 μM Mn (Fig. 4).