1 **Structurally informed evolutionary models improve phylogenetic reconstruction for**

2 **emerging, seasonal, and pandemic influenza viruses**

4 Xueting Qiu[1,2], Justin Bahl[1,3,4]

5
[1] Center for Ecology of Infectious Diseases, Department of Infectious Diseases, College of
Veterinary Medicine, University of Georgia, Athens, GA
[2] Center for Communicable Disease Dynamics, Harvard T.H. Chan School of Public Health,
Boston, MA
[3] Department of Epidemiology and Biostatistics, College of Public Health, University of
Georgia, Athens, GA
[4] Program in Emerging Infectious Diseases, Duke-National University of Singapore Graduate
Medical School, 8 College Road, Singapore 169857, Singapore
14
15 Corresponding Authors
16 Email: xuetingqiu@hsph.harvard.edu; Justin.Bahl@uga.edu

Abstract

Precise estimation of genetic substitution patterns is critical for accurate reconstruction of pathogen phylogenies. Few studies of viral evolution account for variations of mutation rate across a single gene. This is especially true when considering evolution of segmented viruses where individual segments are short, encoding for few proteins. However, the structural and functional partitions of these proteins could provide valuable information for more accurate inference of viral evolution, due to the disparate immune selection pressure on different functional domains. Accurately reconstructed evolutionary features on specific functional domains can in turn provide biological information on viral protein and immune targets for vaccine design. In this study we developed and evaluated a structurally informed partitioning scheme that accounts for rate variation among immunogenic head and stalk domains of the surface protein hemagglutinin (HA) of influenza viruses. We evaluated the model fit and performance of four different models - HKY, SRD06 codon, HKY with a structurally informed partitioning scheme, SRD06 with a structurally informed partitioning scheme on pandemic A/H1N1pdm09, seasonal A/H1N1postpdm, A/H3N2, B-Yamagata-like and Victoria-like lineages, and two highly pathogenic avian influenza A viruses H5Nx and H7N9. Results showed that structurally informed partitioning with SRD06 performed better for all datasets with decisively statistical support. Significantly faster nucleotide substitution rates for head domain, compared to stalk domain was observed and may provide insight for stalk derived broadly-reactive vaccine design. Taken together, integrating a functionally informed partitioning scheme based on protein structures of immune targets allows for significant improvement of phylogenetic analysis and providing important biological insights.

Introduction

The importance of statistical phylogenetic methods to study the epidemiology, evolution and ecology of rapidly evolving viral pathogens has been driven by the growing availability of whole genome sequences (1). Precisely estimating the pattern of genetic variations is critical to reconstruct the accurate pathogen phylogenies (2,3). Nucleotide substitution models have been developed to describe the process of change from one nucleic state to another among viral isolates, where they often allow for rate variations between transitions and transversions (4), or incorporate nucleotide base frequencies with substitution rate parameters (5,6). Segmented viruses, such as influenza A virus (IAV), contain short gene segments that encode for few proteins (7). Consequently, the literature regarding model development of phylogeny reconstruction is dominated by the analysis on complete protein coding regions. For example, the complete viral surface glycoprotein hemagglutinin (HA) is used the most when studying influenza viral evolution and diffusion (8). However, assuming natural selection acts uniformly across the protein domains may not be justified (9). Some partitioning strategies have been developed to account for variations inside one gene segment (3,10). For example, approximate codon-models, such as the SRD06, incorporate information about the genetic code by allowing for a rate variation by defining substitution models for codon positions 1 and 2 and an independent model for codon position 3. This model accounts for the rapid accumulation of synonymous mutations in the third codon position (10).

HA glycoprotein with a spike-shaped structure binds to the receptors on the targeted host cell membrane when the virus begins the infection (11). This protein has two main domains – the globular head domain and stalk domain. Functionally different, the globular head domain contains the receptor binding sites, while the stalk domain is a main structure responsible for

membrane fusion machinery (11,12). Even though immune selection strongly drives antigenic

drift, allowing for accumulation of mutations in the head domain, the stalk domain is

functionally conserved across viral subtypes. Current seasonal influenza virus vaccines induce

humoral immune responses primarily targeting the immunodominant globular head domain

(13,14), which can provide strong protection from an infecting virus that is similar with vaccine

candidates, but have weak cross-reactivity to antigenic drift variants. Conversely, studies in the

ferret experimental model demonstrate that the stalk domain is highly conserved across IAV

subtypes and stalk specific antibodies could provide cross-reactive protection to a high diversity

of IAV subtypes (13,15). Despite the importance of understanding how substitution rates and

patterns vary across functionally conserved domains in universal vaccine design (16,17), few

studies have incorporated protein structure in phylogenetic models.

In this study we proposed a novel phylogenetic model that incorporates a protein

structure informed partitioning scheme to account for variable evolutionary rates resulting from

immune selection. We aimed to evaluate the appropriateness of the novel model to reliably to

estimate biologically informative parameters from available genetic data. We evaluated and

compared four different models: HKY, SRD06 codon, HKY with a structurally informed

partitioning scheme, SRD06 with a structurally informed partitioning scheme. We also evaluated

the model performance across multiple viral subtypes, including pandemic H1N1pdm09,

seasonal A/H1N1postpdm, A/H3N2, the two seasonal Influenza B virus lineages – Yamagata-

like and Victoria-like, and two highly pathogenic avian Influenza (HPAI) H5Nx and H7N9

viruses. We further conducted sensitivity analysis to determine whether the new model was

sensitive to sample size, data distribution and epidemic stage resulting in biased estimation.

Separate estimation of viral evolutionary rates for the head and stalk domain may be informative

for vaccine design. We therefore formally tested the hypothesis that evolutionary rates in each

domain were significantly different and further estimated the relative selection pressure for each

functional domain of HA protein.

New Approaches

This study proposed a new model to incorporate a structurally informed partitioning

scheme on a single protein into phylogenetic reconstruction. Decisive statistical support from the

model selection procedure and stably supported across multiple influenza subtypes validated the

superiority of the new model. The model can provide biological insights for viral evolution, with

reporting domain-specific evolution rates and approximate selection pressure. The tree branches

with domain-specific rate ratio $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ can inform the approximate selection pressure on each

strain, suggesting some biological explanations related to antigenic drift and emerging strains.

The structurally informed phylogenetic model may reveal novel biological insights of viral

evolution and have the potential to reveal more biological realism without over-parameterization.

Results

Model selection

To compare the four models, the log scale Bayes Factor (BF) was used, which was the

difference of log marginal likelihood estimations from two models. Marginal likelihood

estimation was conducted with two approaches – path-sampling and stepping-stone sampling,

where both generated consistent results. Based on the model selection procedure for all subtypes

(Fig. 1), the model with mixed partitioning on both codon positions and HA protein structure

(CP model in Fig. 1c) had decisive BF support for better model fitting in all datasets compared to

HKY and HKY with protein structure partitioning model (HKY model and P model in Fig. 1c).

Compared to SRD06 codon model (C model in Fig. 1c), CP model performed significantly better

for all datasets, with different levels of BF supports. HKY with protein structure model (P

model) performed significantly better than HKY model for human influenza A and B viruses, but

it was not significant in highly pathogenic avian influenza H7N9 and H5Nx datasets. This may

indicate that these emerging viruses with avian-dominated host populations have a different

immune selection pressure pattern compared to human influenza viruses.

Tree root height and substitution rates

To evaluate the accuracy and stability of model performance, tree parameters and

estimated substitution rates were used to compare different models. Results from four models

showed very similar tree root height and 95% Bayesian Credible Interval (BCI) for each

influenza subtype (Table 1). The estimated mean substitution rates (Fig. 3) estimated from all

models were similar for each dataset suggesting that the CP model generated reliable estimations

for important evolutionary parameters that were comparable to other models. Furthermore, since

both CP and P model incorporated partitioning scheme on protein structure, these two models

provided domain-specific substitution rates for STC (including signal peptide, transmembrane

domain and cytoplasmic tail), stalk and head domains, respectively. However, the domain-

specific rates reported by P model were underestimated, compared to CP model (results shown in

Supplemental Table 2). Such an observation was unsurprising since the HKY substitution model

on each protein structure partition cannot account for more realistic variations among codon

positions. One thing to note, though STC domain was not partitioned on codon positions, its

small amount of information from short nucleotide length (about 150 nucleotides) resulted in

high uncertainty of the STC substitution rates, showing long tails in the violin plots.

Comparing the overall substitution rates for each influenza subtype, we observed that the

pandemic H1N1pdm09 had the highest substitution rate ($\sim 7.5 \times 10_{-3}$ substitution/site/year).

HPAI H7N9 and H5Nx had the substitution rates around $4.0 - 4.8 \times 10_{-3}$ substitution/site/year.

Seasonal influenza viruses had the range as $2.0 - 3.5 \times 10_{-3}$ substitution/site/year, where the two

type B lineages had lower rates than the two A subtypes.

Furthermore, we compared the separate substitution rates for stalk and head domains

from the CP model. The absolute rates calculated directly from each step of the Bayesian

Markov Chain Monte Carlo (MCMC) simulation were plotted into violin plot (Fig. 3). Results

showed that the head domain had a significantly higher substitution rate than stalk domain for all

influenza subtypes, under the unit of substitutions per site per year. The Bayes Factors (BF) for

the statistical tests between the substitution rates of head and stalk domains were 1137

(H1N1pdm09), 6 (H1Npostpdm), 577 (A/H3N2), 197 (B-Victoria), 191 (B-Yamagata), 303

(H7N9), +infinity (H5Nx), respectively, with the significant criteria as BF>3.

Approximate selection pressure

With partitioning on codon positions, the CP model can estimate the ratio of substitution

rate of codon positions 1 and 2 (majority of non-synonymous changes occur in these codons)

over the rate of codon position 3 (majority changes are synonymous), that is, $d\text{C}_{1+2}/d\text{C}_3$, for stalk

domain and head domain separately. The relative rate of C_3 to C_{1+2} is a good predictor of the ω

parameter of selection pressure measure (18) and thus could reflect the domain specific selection

pressure. As shown in Table 2.2, the range of $d\text{C}_{1+2}/d\text{C}_3$ was $0.10 - 0.32$ for stalk domain and

$0.25 - 0.87$ for head domain from all subtypes of influenza viruses included in the study. The

overall $d\text{C}_{1+2}/d\text{C}_3$ for the whole HA protein calculated from the c model was lower than head

domain but higher than stalk domain for each influenza subtype. We also observed that the stalk

domain had a significantly lower $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ than head domain for each influenza subtype (Bayes

Factors shown in Table 2), which indicated that stalk domain experienced stronger purifying

selection to maintain its conservation. Further to compare pandemic H1N1pdm09 and seasonal

A/H1N1 post pandemic, we found that the seasonal H1N1 post pandemic had lower $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ in

both stalk and head domains, indicating pandemic H1N1pdm09 experienced less purifying

selection.

To observe different patterns of selection pressure on the viral populations, phylogenetic

tree branch-specific stalk $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ and head $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ were calculated and visualized via

mapping the upper and lower quantiles of the ratio values onto the phylogenetic tree. We used

seasonal H3N2 (Supplemental Fig. 1) and Egypt H5Nx (Supplemental Fig. 2) as examples. The

$d\mathrm{C}_{1+2}/d\mathrm{C}_3$ had a range of 0.63 – 0.75 for H3N2 head domain, 0.21 – 0.23 for H3N2 stalk domain,

0.12 – 1.87 for Egypt H5Nx head domain and 0.09 – 0.32 for Egypt H5Nx stalk domain. Again,

results from the branch-specific $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ demonstrated that the head domain had significantly

higher $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ compared to stalk domain, which indicated that stalk domain was under

stronger purifying selection pressure to maintain its conservation. To note, the majority of the

$d\mathrm{C}_{1+2}/d\mathrm{C}_3$ values were less than 1, which indicated purifying selection, but occasionally some

diversifying selection pressure ($d\mathrm{C}_{1+2}/d\mathrm{C}_3 >1$) was observed in Egypt H5Nx head domain in

chicken-dominated population. This high $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ in H5Nx chicken population may indicate

that the viruses circulating in chicken may experience less purifying selection pressure to

generate higher diversity with keeping more non-synonymous changes compared to that in

human population. This suggests that avian population may generate more variants with

pandemic potential.

Sensitivity analysis and model validation

With the largest dataset HPAI H5Nx in this study, sensitivity analysis was performed to

estimate the stability of model performance under different sample sizes. The same model fitting

and selection procedure was tested on randomly selected 80%, 60%, 40% and 20% of the H5Nx-

full dataset. The model selection procedure (Fig. 4) showed that all these datasets had decisive

BFs to favor the structurally informed partitioning model with SRD06 (CP model). Tree root

height estimation (Table 3) from these datasets had similar estimates and 95% BCI, where it was

only slightly overestimated in the 20% of H5Nx dataset. The substitution rates of protein

structure partitions (Fig. 5) also showed stable estimations across different sample sizes but only

had slight overestimation with the 20% H5Nx dataset. Taken together, though the new model

with partitioning on both codon positions and protein functional domains introduced more

parameters, it performed stably with different amount of data and thus was not sensitive to

sample size.

With two subsets of random selection from seasonal H3N2, model selection procedure

and evolutionary parameter estimations generated very similar results. It indicated that the new

model validated through different subsets and was not sensitive to the different distributions of

epidemiological factors, including geographic location and isolation time (shown in

Supplemental Table 3).

With dividing the H1N1pdm09 dataset into the Early stage and Later stage of an

epidemic outbreak, model selection procedure reported that the CP model was significantly

better than all other models in both datasets. The overall and domain-specific substitution rates in

the Early stage was slightly higher compared to the Later stage of the outbreak (shown in

Supplemental Table 4). And the Early stage had a higher dC_{1+2}/dC_3, indicating it experienced

weaker purifying selection pressure to be relatively diverse. This may be related to host

adaptation in the Early stage, where it generated more variants for the advantage of widely

spreading in the host population.

Discussion

In this study a novel phylogenetic model with integrating both codon positions and

protein structure was developed and tested with HA protein of influenza viruses. Results showed

that the new model was statistically supported to fit the data better and provided novel biological

information that can reveal new insights into viral evolution. The substitution rates of stalk and

head domains of HA protein were significantly different, with a higher rate in head domains for

all the tested influenza types. Approximate selection pressure constructed from codon position

partitions showed that stalk domain experienced higher purifying selection pressure to maintain

its conservation. Model validation from subsets of multiple influenza subtype reported the model

performed stably and was not sensitive to sample size, data distribution and epidemic stage.

Taken together, this new model could provide potential biological explanations related to

antigenic drift, vaccine escape and pandemic emergence to understand viral evolution and risk

assessment.

Partitioning scheme on genetic data allows for capturing similar evolutionary features

inside one partition of data and comparing different partitions to generate new insights on viral

evolution. For example, SRD06 codon model categories the first and second codon positions

together since they behave similarly to mainly generate non-synonymous changes, while the

third codon position is treated as a separate partition due to mostly generating synonymous

changes (3,10). Some models conduct partitioning scheme on multiple proteins to reconstruct the

co-evolution and estimate protein-specific rates under one model (19). These models function

well for some purposes of observing the co-evolution of multiple proteins, however, it assumes

evolutionary processes act uniformly across one protein, which may not be the biological reality

due to distinctively functional domains on one protein.

When developing the new model in this study, the primary tasks are to efficiently capture

the information regarding HA protein structural and functional domains and to properly

incorporate this information into the evolutionary models. Therefore, how to partition on the

protein structure with genetic data is critical. One solution is to compare all possible partitioning

schemes for a given genetic sequence dataset to find the best partitioning strategy (3). However,

this approach is computationally inflexible because the combination of possible partitioning

schemes is tremendous even for a very small number of data blocks (20). Consequently, many

previous studies ended up either arbitrarily choosing a single partitioning scheme or just

selecting the best scheme based on statistical model selection (3,21,22). With these limitations on

proper partitioning, the accuracy of the inferences from partitioned phylogenetic analysis

remains uncertain, even though advanced developments in phylogenetic analysis have been

achieved in recent decades (3). Due to advanced progress in viral protein structure (23),

partitioning scheme in this study is based on biological functions of different protein domains,

where HA protein has been defined to distinguish the immunogenic roles and other molecular-

level functions during viral infection. Molecular mechanism of influenza viruses, for example,

the initiation of viral fusion or antigenic-immune stimulation, is determined by the head domain

of the protein. Nucleic acid changes observed in this domain usually result from a balance

between immune selection and conserved functionality (24–26). The new partitioning model

considers the distinctive functions of head domain to mainly bind to host-cell receptors and the

stalk domain to perform membrane fusion.

Another advance in this study is that a formal model selection procedure with reliable

approaches (27) to quantitatively evaluate the model performance and accuracy of inferences for

pandemic, all seasonal and two emerging influenza viruses. Previously the harmonic mean

estimate (HME) and Akaike's information criterion through MCMC (AICM) (28) have been

commonly used. In recent years, several new approaches to perform model selection in the field

of phylogenetics, such as path- sampling (PS) (29) and stepping-stone sampling (SS) (30). Baele

et al. (31) have tested HME, AICM, PS, and SS approaches for demographic and molecular

clock model comparison, which "confirmed that HME systematically overestimates the marginal

likelihood and fails to yield reliable model classification, and PS and SS substantially outperform

HME estimators" (32). Therefore, in this study both PS and SS approaches have been used to

perform the model selection procedure to generate reliable results of model comparison.

The new model with both partitioning on codon positions and protein structure

incorporates biological functional domains inside one protein to allow for a quantitative

exportation of the evolutionary process as a function of approximate host immune selection. The

model did capture the different evolutionary process of these functional domains, which showed

that the stalk domain had a significantly lower substitution rates and lower $d\mathrm{C}_{1+2}/d\mathrm{C}_3$, indicating

that the stalk domain experienced stronger purifying selection and maintained its conserved

functionality. The estimated domain-specific $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ can reveal insights of selection pressure

on different viral isolates in different host populations. For example, diversifying selection

pressure on the head domain has been observed only in chicken population in Egypt. One

possible explanation could be that the failed efficacy of poultry influenza H5 vaccines

implemented in Egypt (33,34) may drive the genetic diversity of the viruses in the chicken

population.

Even though $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ estimated by the model is not an exact measurement of the ω

parameter, it is a good predictor of the ratio of non-synonymous changes over synonymous

changes (10). The new model can provide an informative approximation and avoid heavy

computational demands of a full codon or amino acid models (10). When assuming synonymous

changes do not affect viral fitness, the $d\mathrm{C}_{1+2}/d\mathrm{C}_3$ value can provide corresponding estimation for

selection pressure compared to mutation-selection models which assess the strength of natural

selection on specific mutations (35). Furthermore, the partitioning scheme and additional

parameter estimation did not significantly increase the analysis run time compared to other

models in this study under the same computing environment.

The structurally informed evolutionary model may provide insight for universal vaccine

design. The higher rates observed in the head domain are probably due to strong immune

selection targeted on the globular head, where some sites are mostly targeted by the neutralizing

immunity induced by seasonal influenza vaccine (13). Supported by many laboratory studies,

stalk domain could be used to induce more broad-reactive vaccines (13,15,17). This model can

more accurately reconstruct the ancestral sequence of the stalk domain with estimating domain-

specific rates and substitution patterns under a shared viral phylogeny. Therefore, if this model

was used to reconstruct common ancestral sequence for all viral types in influenza A virus Group

1 (H1, H2, H5, H6, H8, H9, H11, H12, H13, H16, and H17) and Group 2 (H3, H4, H7, H10, H14

and H15) (25), respectively, it is probably amongst the most promising approaches to design a

universal influenza vaccine.

While this model has been developed with influenza in mind, there is potential to apply

this approach to other viruses. Proteins, such as the respiratory syncytial virus F and G protein or

the coronavirus spike protein that have multiple functional domains including immune

stimulators, may be well suited to the novel approach that incorporates protein structural and

functional domains into the nucleotide substitution models. Incorporating this type of

information into comparative genomic analyses has the potential to provide important biological

information and improve both vaccine design and evaluation.

Methods

Datasets

Multiple datasets representing pandemic, seasonal and emerging influenza viruses were

used to develop and broadly test the proposed structurally informed partitioning models,

including pandemic H1N1pdm09, seasonal influenza viruses A/H1N1postpdm, A/H3N2,

B/Yamagata-like and B/Victoria-like, and two HPAI H5Nx and H7N9. Except for HPAI H7N9,

H1N1pdm09 and seasonal A/H1N1postpdm that have been directly downloaded from the Global

Initiative on Sharing All Influenza Data (GISAID, https://www.gisaid.org/) (36,37), all other

datasets were from published research, including seasonal A/H3N2 (38), HPAI H5Nx (39),

B/Yamagata-like and B/Victoria-like (40). To avoid improper weights of duplicate isolates on

the partitioning models, sequences with 100% similarity were removed for all datasets (final

datasets can be found via the GitHub link:

https://github.com/XuetingQiu/FluPartitioningModels).

Briefly, the final datasets contained 505 global isolates for H1N1pdm09 during 2009/04 –

2009/12, 635 global isolates for HPAI H7N9 during 2013-2018, and 554 global isolates for

seasonal A/H1N1postpdm during 2012-2018. HPAI H5Nx contained 1,095 isolates during 1996-

2016, after removing the vaccine candidates (39). The two influenza B virus lineages were full

genome datasets sampled during 2002-2013 in eastern Australia and New Zealand (40). For B-

Victoria, the dataset finally retained 214 isolates after randomly sampling down to 50% of the

original dataset and removing the duplicates. For B-Yamagata, 241 isolates were included after

removing the duplicates. The global dataset of seasonal A/H3N2 (38) contained 906 isolates

during 2000-2012 after the duplicates were removed and further randomly subsampled.

Definition of HA protein structural partitions

HA glycoprotein has been defined as different functional domains, including signal

peptide, stalk domain, globular head domain, transmembrane domain and cytoplasmic domain

(11,12). These protein domains labeled with sites of amino acid were translated into the linear

sites on the nucleotide sequences of HA gene (Fig. 2.1a and 2.1b). To use the full length of HA

sequences and avoid some partitions being too short and lack of information, signal peptide,

transmembrane domain, and cytoplasmic domain were combined as one partition named STC,

given the similarly evolutionary characteristics of these functional domains. Though the stalk

domain on the linear nucleotide sequence was separated by the head domain, the two gene

regions of stalk domain were merged together for analysis (supplemental table I-1 and xml files

in GitHub https://github.com/XuetingQiu/FluPartitioningModels).

Structurally informed phylogenetic models

As shown in Fig. 2.1b, four different Bayesian phylogenetic models were conducted

using the same molecular clock model and coalescent model for each dataset. Only the

partitioning strategies on the nucleotide sequences were different. The base model applied HKY

substitution model (4) with Gamma invariant distribution, where this model accounts for base

frequencies and allows for rate variations between transitions and transversions. SRD06 codon

model (C model in Fig. 2.1c) (10) partitioned all codons of the HA gene reading frame into

codon positions 1+2 and codon position 3, where these two partitions also used HKY

substitution model. Protein structure partitioning model (P model in Fig. 2.1c) contained STC,

stalk and head functional domains with HKY substitution model for each domain. The last model

(CP model in Fig. 2.1c) combined the partitioning strategy in both C and P models, that is, codon

positions in each protein functional domain were further partitioned into codon positions $1 + 2$

and codon position 3.

Tree likelihood construction in the partitioning models

After partitioning with codon positions or/and protein structural domains, the overall tree

likelihood calculations need to be explained. The general Bayesian inferences are based on the

posterior probability of a hypothesis, for example, a given tree τ. Then the posterior probability

of the tree based on the data can be obtained using Bayes theorem,

349
$$Pr(\tau|x) = \frac{Pr(x|\tau)Pr(\tau)}{\sum Pr(x|\tau)Pr(\tau)} \quad [1]$$

350 In which, *Pr(τ)* is the prior probability of the tree hypothesis τ based on data x; *Pr(x/τ)* is

351 the probability of observing the data at a specific site.

352 Then assuming independence of substitution across sites, the general likelihood based on

353 a specific phylogenetic model, i.e. the probability of observing the aligned matrix X of all j

354 sequences is,

355
$$f(X|\tau_j, v_j, \theta, \phi) = \prod_{i=1}^{c} f(x_i|\tau_j, v_j, \theta, \phi) \quad [2]$$

In which, c is the number of total sites along the sequences; τ represents the tree hypothesis, v

represents the branch lengths, θ is the substitution model parameters, and ϕ represents all other

model parameters.

The proposed structurally informed partitioning model based on HA protein functional

domains extends the formula [2] of the tree likelihood estimation. Each of the defined partition

has its tree likelihood estimated independently, therefore, the overall tree likelihood is the

products of each partition-specific likelihood, that is,

$$f(X|\tau_j, v_j, \theta, \phi)$$

$$= \prod_{i=1}^{c_1} f(x_i|\tau_j, v_j, \theta, \phi)$$

$$* \prod_{i=1}^{c_2} f(x_i|\tau_j, v_j, \theta, \phi) * \dots * \prod_{i=1}^{c_n} f(x_i|\tau_j, v_j, \theta, \phi)$$

In which, the sum of c_1, c_2, …, c_n is equal to c, the total number of nucleotide sites along

the sequences; and n is the number of partitions in the model.

Phylogenetic model simulations

All models were simulated in the package of Bayesian Evolutionary Analysis Sampling

Trees (BEAST) v1.8.4 (41). To allow for rate variations across lineages, the uncorrelated

lognormal relaxed molecular clock was used with an initial mean of 0.0033 with a uniform prior

ranging from 0.0 to 1.0. Based on prior knowledge, a smooth and time-aware Gaussian Markov

random field (GMRF) process prior on the population sizes was applied in the Skyride calescent

model (42). Each model was performed at least four independent runs of 100 million Markov

Chain Monte Carlo (MCMC) generations. To report the substitution rates and phylogenetic trees,

four runs for each model were combined after removal of burn-in to achieve an Effective Sample

Size (ESS) of >200 as diagnosed in Tracer v 1.5. Visualized violin plots for substitution rates

were generated via personal R scripts (GitHub

https://github.com/XuetingQiu/FluPartitioningModels).

Model selection procedure and criteria

The goal of model selection procedure is to compare model superiority and identify a

model that is sufficiently complex to capture the biological realism and evolutionary processes

that have occurred but to avoid overparameterized models with more parameters than these can

be reliably estimated from the available data (3,31). In this study, both path-sampling (PS) (29)

and stepping-stone sampling (SS) (30) were used to compute the marginal likelihood estimation

to perform the model selection procedure. PS and SS approaches account for both the number of

parameters and the appropriateness of prior distributions for these parameters.

The marginal likelihood is the probability of the data (that is, likelihood) given the model

type, not assuming any particular model parameters (31). The marginal likelihood estimation for

both PS and SS with Beta path step distributions is set to run 100 path steps and length of chains

as 1 million. Bayes Factors (BF), the ratio of marginal likelihood estimations from two models,

are used to evaluate the significance of the model comparison. When converted to a log scale, BF

becomes the difference between two log marginal likelihood estimations. The mathematical

conversion is below (43):

For a model with parameters Θ, the marginal likelihood for the model M with data X is,

396
$$p\ (X\ /\ M) = \int p\ (X|\Theta, M)\, p(\Theta|M) d\Theta \quad [3]$$

397   Then the Bayes factor can be calculated for model M1 against model M2,

398
$$BF_{12} = \frac{p(X|M_1)}{p(X|M_2)} = \frac{\int p\ (X|\Theta_1, M_1)\, p(\Theta_1|M_1) d\Theta_1}{\int p\ (X|\Theta_2, M_2)\, p(\Theta_2|M_2) d\Theta_2} \quad [4]$$

399   The log scale of BF is derived as below,

400
$$logBF_{12} = log\ p(X|M_1) - log\ p(X|M_2) = dB \quad [5]$$

With log scale BF as dB, the criteria for model selection are: dB <0 means the selection

supports M2; $0 \leq$ dB <5 means no evidence for supporting M1; $5 \leq$ dB < 10 means substantial

strength of evidence for supporting M1; $10 \leq$ dB <15 means strong evidence of supporting M1;

$15 \leq$ dB < 20 means very strong evidence of supporting M1; dB ≥ 20 means decisive evidence of

supporting M1.

Model performance, validation and sensitivity analysis

To compare model performance among four models for each dataset, two basic

evolutionary parameters were estimated and reported: the root heights and overall substitution

rates. If the parameters for each dataset were similar across these four models, then it indicated

that the model can provide accurate and stable estimation no matter how many partitions it has

included.

To evaluate whether the structurally informed partitioning model was sensitive to sample

size, sensitivity analysis was performed on HPAI H5Nx from 1996-2017, the largest final dataset

with 1,095 isolates in this study. With retaining the oldest isolate of H5Nx, the rest was

randomly subsampled to keep 80%, 60%, 40% and 20% of the dataset, respectively. Four runs of

100 million MCMC chain length of the four models for the full dataset and these subsampled

datasets were conducted with marginal likelihood estimations for model selection. Tree height

parameters, domain-specific substitution rates, and model selection BFs were reported to

evaluate the data sufficiency for model stability.

To further conduct model validation, two more approaches were applied. Firstly, to

evaluate whether the structurally informed partitioning model was sensitive to data distribution,

the full seasonal A/H3N2 dataset was randomly and repeated sampled 40% of the full dataset

into two subsets. These two subsets had different geographical and temporal distributions of

A/H3N2 isolates. Secondly, to assess whether the new model was sensitive to the epidemic

stage. With the H1N1pdm09 dataset, the Early stage of the pandemic was defined as April to

middle of July, 2009, and the Later stage of the pandemic was July-December, 2009. The same

model variations and model selection procedure were conducted to all these subsets.

Domain-specific $d\text{C}_{1+2}/d\text{C}_3$ recorded to tree branches

The new model can report domain-specific rates and $d\text{C}_{1+2}/d\text{C}_3$ values. To visualize these

specific rates or ratios on the tree branches, the BEAST xml files (GitHub link:

https://github.com/XuetingQiu/FluPartitioningModels) were coded to compute and load the

specific branch rates for stalk domain codon positions 1+2, stalk domain codon position 3, head

domain codon positions 1+2, and head domain codon position 3. Python script (GitHub link:

https://github.com/XuetingQiu/FluPartitioningModels) was constructed to compute and write the

$d\text{C}_{1+2}/d\text{C}_3$ for stalk and head domain from the median rate of each partition to the final

summarized tree branches. Seasonal A/H3N2 and all Egypt H5Nx data were used as examples to

show the domain-specific $d\text{C}_{1+2}/d\text{C}_3$ changes in A/H3N2 with different vaccine isolates

introduced and the changes in different hosts (human v.s. avian) in Egypt H5Nx.

Acknowledgements

This study was partially funded by NIAID Centers of Excellence for Influenza Virus Research

and Surveillance (CEIRS, HHSN272201400006C). These funders had no role in study design,

data collection and analysis, decision to publish, or preparation of the manuscript. We gratefully

acknowledge the authors, originating and submitting laboratories of the sequences from

GISAID's EpiFluTM Database, for making tremendous data publicly available and largely

benefiting the science community.

References

1.   Campbell F, Strang C, Ferguson N, Cori A, Jombart T. When are pathogen genome

sequences informative of transmission events? Parrish C, editor. PLOS Pathog [Internet].

2018 Feb 8 [cited 2019 Jun 29];14(2):e1006885. Available from:

https://dx.plos.org/10.1371/journal.ppat.1006885

2.   Biek R, Pybus OG, Lloyd-Smith JO, Didelot X. Measurably evolving pathogens in the

genomic era. Trends Ecol Evol [Internet]. 2015 Jun [cited 2019 Jan 9];30(6):306–13.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/25887947

3.   Lanfear R, Calcott B, Ho SYW, Guindon S. PartitionFinder: Combined Selection of

Partitioning Schemes and Substitution Models for Phylogenetic Analyses. Mol Biol Evol

[Internet]. 2012 Jun 1 [cited 2019 Feb 20];29(6):1695–701. Available from:

https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/mss020

4.   Hasegawa M, Kishino H, Yano T. Dating of the human-ape splitting by a molecular clock

of mitochondrial DNA. J Mol Evol [Internet]. 1985 [cited 2019 Feb 20];22(2):160–74.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/3934395

5.   Lanave C, Preparata G, Sacone C, Serio G. A new method for calculating evolutionary

substitution rates. J Mol Evol [Internet]. 1984;20(1):86–93. Available from:

https://doi.org/10.1007/BF02101990

6.   Tavaré S. Some Probabilistic and Statistical Problems in the Analysis of DNA Sequences.

Am Math Soc Lect Math Life Sci [Internet]. 1986;17:57–86. Available from: citeulike-

article-id:6732633

7.   Webster RG, Bean WJ, Gorman OT, Chambers TM, Kawaoka Y. Evolution and ecology

of influenza A viruses. Microbiol Rev [Internet]. 1992 Mar 1 [cited 2019 Feb

12];56(1):152–79. Available from: http://www.ncbi.nlm.nih.gov/pubmed/1579108

8. Baele G, Suchard MA, Rambaut A, Lemey P. Emerging Concepts of Data Integration in Pathogen Phylodynamics. Syst Biol [Internet]. 2017 Jun 1 [cited 2018 Dec 2];66(1):e47–65. Available from: https://academic.oup.com/sysbio/article/doi/10.1093/sysbio/syw054/2670001/Emerging-concepts-of-data-integration-in-pathogen

9. Koonin E, Galperin M. Evolutionary Concept in Genetics and Genomics. In: Sequence - Evolution - Function: Computational Approaches in Comparative Genomics. Boston: Kluwer Academic; 2003.

10. Shapiro B, Rambaut A, Drummond AJ. Choosing Appropriate Substitution Models for the Phylogenetic Analysis of Protein-Coding Sequences. Mol Biol Evol [Internet]. 2006 Jan 1;23(1):7–9. Available from: http://dx.doi.org/10.1093/molbev/msj021

11. Riwilaijaroen BNS, Uzuki YS. Review Molecular basis of the structure and function of H1 hemagglutinin of in fl uenza virus. 2012;88:226–49.

12. Wiley DC, Skehel JJ. The Structure and Function of the Hemagglutinin Membrane Glycoprotein of Influenza Virus. Annu Rev Biochem [Internet]. 1987 Jun 1 [cited 2019 Feb 13];56(1):365–94. Available from: http://www.ncbi.nlm.nih.gov/pubmed/3304138

13. Nachbagauer R, Miller MS, Hai R, Ryder AB, Rose JK, Palese P, et al. Hemagglutinin Stalk Immunity Reduces Influenza Virus Replication and Transmission in Ferrets. 2016;90(6):3268–73.

14. Wang TT, Palese P. Catching a Moving Target. Science (80-) [Internet]. 2011 Aug 12;333(6044):834 LP – 835. Available from: http://science.sciencemag.org/content/333/6044/834.abstract

15. Krammer F, Hai R, Yondola M, Tan GS, Leyva-Grado VH, Ryder AB, et al. Assessment

of influenza virus hemagglutinin stalk-based immunity in ferrets. J Virol [Internet]. 2014

Mar 15 [cited 2017 Nov 16];88(6):3432–42. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/24403585

16. Krammer F, Palese P. Influenza virus hemagglutinin stalk-based antibodies and vaccines.

Curr Opin Virol [Internet]. 2013;3(5):521–30. Available from:

http://www.sciencedirect.com/science/article/pii/S187962571300134X

17. Subbarao K, Matsuoka Y. The prospects and challenges of universal vaccines for

influenza. Trends Microbiol [Internet]. 2013 Nov 15;21(7):350–8. Available from:

http://dx.doi.org/10.1016/j.tim.2013.04.003

18. Shapiro B, Rambaut A, Drummond AJ. Choosing Appropriate Substitution Models for the

Phylogenetic Analysis of Protein-Coding Sequences. Mol Biol Evol [Internet]. 2006 Jan 1

[cited 2019 Feb 20];23(1):7–9. Available from:

http://academic.oup.com/mbe/article/23/1/7/1193608/Choosing-Appropriate-Substitution-

Models-for-the

19. Mintaev RR, Alexeevski A V, Kordyukova L V. Co-evolution analysis to predict protein–

protein interactions within influenza virus envelope. J Bioinform Comput Biol [Internet].

2014 Mar 6;12(02):1441008. Available from:

https://doi.org/10.1142/S021972001441008X

20. Li C, Lu G, Ortí G. Optimal Data Partitioning and a Test Case for Ray-Finned Fishes

(Actinopterygii) Based on Ten Nuclear Loci. Buckley T, editor. Syst Biol [Internet]. 2008

Aug 1 [cited 2019 Apr 18];57(4):519–39. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/18622808

21.   Brandley MC, Schmitz A, Reeder TW. Partitioned Bayesian Analyses, Partition Choice,

and the Phylogenetic Relationships of Scincid Lizards. Anderson F, editor. Syst Biol

[Internet]. 2005 Jun 1 [cited 2019 Apr 18];54(3):373–90. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/16012105

22.   McGuire JA, Witt CC, Altshuler DL, Remsen J V. Phylogenetic Systematics and

Biogeography of Hummingbirds: Bayesian and Maximum Likelihood Analyses of

Partitioned Data and Selection of an Appropriate Partitioning Strategy. Zamudio K,

Sullivan J, editors. Syst Biol [Internet]. 2007 Oct 1 [cited 2019 Apr 18];56(5):837–56.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/17934998

23.   Swanson KA, Settembre EC, Shaw CA, Dey AK, Rappuoli R, Mandl CW, et al. Structural

basis for immunization with postfusion respiratory syncytial virus fusion F glycoprotein

(RSV F) to elicit high neutralizing antibody titers. Proc Natl Acad Sci U S A [Internet].

2011 Jun 7 [cited 2019 Mar 12];108(23):9619–24. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/21586636

24.   Kordyukova L. Structural and functional specificity of Influenza virus haemagglutinin and

paramyxovirus fusion protein anchoring peptides. Virus Res [Internet].

2017;227(Supplement C):183–99. Available from:

http://www.sciencedirect.com/science/article/pii/S0168170216304063

25.   Lu X, Shi Y, Gao F, Xiao H, Wang M, Qi J, et al. Insights into Avian Influenza Virus

Pathogenicity: the Hemagglutinin Precursor HA0 of Subtype H16 Has an Alpha-Helix

Structure in Its Cleavage Site with Inefficient HA1/HA2 Cleavage. J Virol [Internet]. 2012

Dec 25;86(23):12861–70. Available from:

http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3497694/

26.   Skehel JJ, Wiley DC. Receptor Binding and Membrane Fusion in Virus Entry: The

Influenza Hemagglutinin. Annu Rev Biochem [Internet]. 2000 Jun 1;69(1):531–69.

Available from: https://doi.org/10.1146/annurev.biochem.69.1.531

27.   Baele G, Lemey P, Bedford T, Rambaut A, Suchard MA, Alekseyenko A V. Improving

the Accuracy of Demographic and Molecular Clock Model Comparison While

Accommodating Phylogenetic Uncertainty Research article. 2012;29(9):2157–67.

28.   Raftery A, Newton M, Satagopan J, Krivitsky P. Estimating the Integrated Likelihood via

Posterior Simulation Using the Harmonic Mean Identity. Meml Sloan-Kettering Cancer

Center, Dept Epidemiol Biostat Work Pap Ser [Internet]. 2006 Apr 14 [cited 2017 Nov

16]; Available from: http://biostats.bepress.com/mskccbiostat/paper6

29.   Lartillot N, Philippe H, Lewis P. Computing Bayes Factors Using Thermodynamic

Integration. Syst Biol [Internet]. 2006 Apr 1;55(2):195–207. Available from:

http://dx.doi.org/10.1080/10635150500433722

30.   Xie W, Lewis PO, Fan Y, Kuo L, Chen M-H. Improving Marginal Likelihood Estimation

for Bayesian Phylogenetic Model Selection. Syst Biol [Internet]. 2011 Mar 1 [cited 2019

May 5];60(2):150–60. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21187451

31.   Baele G, Lemey P, Bedford T, Rambaut A, Suchard MA, Alekseyenko A V. Improving

the accuracy of demographic and molecular clock model comparison while accomodating

phylogenetic uncertainty. Mol Biol Evol [Internet]. 2012 Jan; Available from:

http://www.ncbi.nlm.nih.gov/pubmed/17065594

32.   Baele G, Lok W, Li S, Drummond AJ, Suchard MA, Lemey P. Accurate Model Selection

of Relaxed Molecular Clocks in Bayesian Phylogenetics Letter Fast Track.

2012;30(2):239–43.

33.   Kandeil A, Sabir JSM, Abdelaal A, Mattar EH, El-Taweel AN, Sabir MJ, et al. Efficacy of

commercial vaccines against newly emerging avian influenza H5N8 virus in Egypt. Sci

Rep. 2018 Dec 1;8(1).

34.   Hamid Samaha MP. Avian influenza vaccination in Egypt: Limitations of the current

strategy. J Mol Genet Med. 2009;03(02).

35.   Spielman SJ, Wilke CO. The relationship between dN/dS and scaled selection

coefficients. Mol Biol Evol [Internet]. 2015 Apr [cited 2019 Jul 19];32(4):1097–108.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/25576365

36.   Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative

contribution to global health. Glob Challenges [Internet]. 2017 Jan 1 [cited 2019 Apr

29];1(1):33–46. Available from: http://doi.wiley.com/10.1002/gch2.1018

37.   Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data – from vision

to reality. Eurosurveillance [Internet]. 2017 Mar 30 [cited 2019 Apr 28];22(13):30494.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/28382917

38.   Bedford T, Riley S, Barr IG, Broor S, Chadha M, Cox NJ, et al. Global circulation

patterns of seasonal influenza viruses vary with antigenic drift. Nature [Internet]. 2015 Jun

8 [cited 2019 Feb 7];523(7559):217. Available from:

http://www.nature.com/articles/nature14460

39.   Qiu X, Duvvuri VR, Gubbay JB, Webby RJ, Kayali G, Bahl J. specific epitope profiles for

HPAI H5 pre- - pandemic vaccine selection and evaluation. 2017;(July):445–56.

40.   Vijaykrishna D, Holmes EC, Joseph U, Fourment M, Su YCF, Halpin R, et al. The

contrasting phylodynamics of human influenza B viruses. Elife. 2015;4:1–23.

41.   Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian Phylogenetics with BEAUti

and the BEAST 1.7. Mol Biol Evol [Internet]. 2012 Aug [cited 2019 Feb 5];29(8):1969–

73. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22367748

42.   Minin VN, Bloomquist EW, Suchard MA. Smooth Skyride through a Rough Skyline:

Bayesian Coalescent-Based Inference of Population Dynamics. Mol Biol Evol [Internet].

2008 Apr 3 [cited 2019 Feb 20];25(7):1459–71. Available from:

https://academic.oup.com/mbe/article-lookup/doi/10.1093/molbev/msn090

43.   Kass RE, Raftery AE. Bayes Factors. J Am Stat Assoc [Internet]. 1995 Jun

1;90(430):773–95. Available from:

http://www.tandfonline.com/doi/abs/10.1080/01621459.1995.10476572

Tables and Figures

Table 1. Tree root height estimation and its 95% Bayesian credible interval (BCI) from each model for each influenza subtype.
Results from four models show very similar tree root height and 95% BCI for each influenza subtype. The unit of the root height is in
years.
602
603 **Models: HKY model** is a substitution model that considers different base frequencies and assigns different rates for transitions v.s.
604 transversions. The **c model** represents SRD06 codon position model. The partitioning strategy is to analyze codon positions 1 + 2 and
605 codon position 3 separately. The **p model** takes the protein structure partitions into account based on the amino acid positions for each
606 domain on the linear diagram. The **cp model** combines both c and p models, which estimates codon positions in protein structural
607 partitions. All models use HKY substitution model.
608

Models	Pandemic			Seasonal													Emerging					
	H1N1pdm09			H1N1postpdm			A/H3N2			B-Victoria			B-Yamagata			H7N9			H5Nx-full		
	Height	95% BCI		Height	95% BCI		Height	95% BCI		Height	95% BCI		Height	95% BCI		Height	95% BCI		Height	95% BCI	
HKY	1.12	0.99	1.29	8.01	7.70	8.34	16.06	14.46	18.86	11.44	11.10	11.84	13.81	12.88	14.86	6.31	5.34	7.89	21.25	21.08	21.48
c	1.12	0.99	1.29	8.01	7.68	8.34	16.06	14.45	18.67	11.46	11.13	11.87	13.79	12.87	14.86	6.30	5.37	7.90	21.28	21.08	21.48
p	1.12	0.98	1.30	8.00	7.71	8.35	16.07	14.38	18.80	11.47	11.13	11.88	13.79	12.87	14.83	6.29	5.32	7.86	21.27	21.08	21.49
cp	1.12	0.99	1.29	8.01	7.69	8.31	16.06	14.44	18.80	11.46	11.12	11.87	13.80	12.84	14.81	6.29	5.34	7.88	21.27	21.08	21.49

609
610

Table 2. Substitution rate and $d\text{C}_{1+2}/d\text{C}_3$ of stalk and head domain for each influenza subtype. Compared to the head domain, the stalk domain with lower value of $d\text{C}_{1+2}/d\text{C}_3$ experiences stronger purifying selection to maintain its conserved functionality.

Datasets	Substitution rates (subs/site/year)				$d\text{C}_{1+2}/d\text{C}_3$*			
	Stalk 1+2*	Stalk 3**	Head 1+2	Head 3	Stalk	Head	BF++	Overall^
H1N1pdm09	3.80E-03	1.18E-02	6.78E-03	1.42E-02	0.32	0.48	53	0.42
H1N1postpdm	1.26E-03	6.37E-03	2.16E-03	6.07E-03	0.20	0.36	+Inf	0.28
A/H3N2	1.39E-03	6.28E-03	3.66E-03	5.42E-03	0.22	0.68	+Inf	0.43
B-Victoria	5.44E-04	4.04E-03	1.51E-03	4.63E-03	0.13	0.33	8804	0.21
B-Yamagata	5.12E-04	4.93E-03	1.39E-03	5.56E-03	0.10	0.25	8875	0.18
H7N9	1.48E-03	7.48E-03	3.05E-03	7.80E-03	0.20	0.39	+Inf	0.30
H5Nx-full	1.74E-03	7.23E-03	6.11E-03	7.03E-03	0.24	0.87	+Inf	0.43

+: $d\text{C}_{1+2}/d\text{C}_3$ means the ratio of substitution rate of codon positions 1 and 2 over the rate of codon position 3.

++: BF is Bayes Factor. BF>3 represents statistical significance.

*: 1+2 represent the mean substitution rates of the codon positions 1 and 2.

**: 3 means the codon position 3.

^: overall $d\text{C}_{1+2}/d\text{C}_3$ is calculated from the SRD06 codon model for the whole HA gene.

Table 3. Tree root height estimation and its 95% Bayesian credible interval (BCI) from each model for all H5Nx datasets. The statistical significance of the new model holds across different subsampling datasets. Only 20% dataset has the trend to slightly overestimate the root height and would not report the precise evolution history of H5Nx. In sum, the new model is not sensitive to sample size. The unit of root height is in years.

Models: HKY model is a substitution model that considers different base frequencies and assigns different rates for transitions v.s. transversions. The **c model** represents SRD06 codon position model. The partitioning strategy is to analyze codon positions 1 + 2 and codon position 3 separately. The **p model** takes the protein structure partitions into account based on the amino acid positions for each domain on the linear diagram. The **cp model** combines both c and p models, which estimates codon positions in protein structural partitions. All models use HKY substitution model.

Models	H5Nx-full			H5Nx-80%			H5Nx-60%			H5Nx-40%			H5Nx-20%		
	Height	95% BCI		Height	95% BCI		Height	95% BCI		Height	95% BCI		Height	95% BCI	
HKY	21.27	21.08	21.48	21.31	21.08	21.57	21.34	21.07	21.62	21.42	21.07	21.79	21.64	21.07	22.19
c	21.29	21.08	21.48	21.30	21.08	21.57	21.32	21.07	21.61	21.43	21.07	21.81	21.65	21.08	22.20
p	21.28	21.08	21.49	21.32	21.08	21.57	21.32	21.07	21.59	21.41	21.07	21.79	21.65	21.08	22.20
cp	21.28	21.08	21.49	21.33	21.08	21.58	21.32	21.07	21.59	21.42	21.07	21.80	21.66	21.09	22.23

Figure 1. Conceptual diagram for influenza virus Hemagglutinins (HA) domains and four
models tested

1a. HA protein 3D structure: Stalk and Head domains. Head domain is the immunodominant
with high plasticity, which contains the receptor binding sites. Stalk domain is immune-
subdominant and more conserved. *1b. Linear diagram for functional domains on HA*
nucleotide sequence. The stalk domain has two parts separated by head domain on the linear
sequence. To compensate for short length, signal peptide, transmembrane domain, and
cytoplasmic tail are combined into one partition, referred to as "STC" in the analysis. *1c. Model*
conceptual design. Four models are applied and compared for each dataset. **HKY model** is a
substitution model that considers different base frequencies and assigns different rates for
transitions v.s. transversions. **C model** represents SRD06 codon position model. The partitioning
strategy is to analyze codon positions 1 + 2 and codon position 3 separately. **P model** takes the
protein structure partitions into account based on the amino acid positions for each domain on
the linear diagram. **CP model** combines both C and P models, which estimates codon positions
in protein structural partitions. All models use HKY substitution model.

Figure 2. Log scale Bayes Factors for model selection procedure for each influenza subtype.
All these Bayes Factors (BF) are calculated to compare with HKY via path-sampling (upper
panel) and stepping-stone sampling (lower panel) approaches to compute the marginal likelihood
estimates for each model. The asterisk * means a significant supportive BF for the structurally
informed codon model (cp), compared to SRD06 codon only model (c model). Both cp and c
models perform better than HKY model with decisive BF supports in all datasets. The HKY plus
structurally informed model (p model) perform better in all datasets except for H7N9.

658
659 Log scale BF criteria: BF< 5 represents no significance; $5 \leq$ BF< 10 means substantial support;
660 $10 \leq$ BF< 15 means strong support; $15 \leq$ BF < 20 means very strong support; BF $\geq$ 20 means
661 decisive support.

662
Models: HKY model is a substitution model that considers different base frequencies and
assigns different rates for transitions v.s. transversions. The **c model** represents SRD06 codon
position model. The partitioning strategy is to analyze codon positions $1 + 2$ and codon position
3 separately. The **p model** takes the protein structure partitions into account based on the amino
acid positions for each domain on the linear diagram. The **cp model** combines both c and p
models, which estimates codon positions in protein structural partitions. All models use HKY
substitution model.

Figure 3. Overall mean substitution rates estimated from four models and domain-specific
substitution rates from structurally informed partitioning model. Violin plots show the
substitution rates from each step of Bayesian simulation. The results show overall mean
substitution rates from four models for each dataset are similar, but the head domain has a
significantly faster rate than that of stalk domain based on the new model (cp model). Generally,
STC domain has more uncertain estimation with long tails of the violin plot, which is probably
due to short length of nucleotides. Bayes Factors for the statistical tests on the differences of
head and stalk domain substitution rates are 1,137 for H1N1pdm09, 6 for H1N1postpdm, 577 for
A/H3N2, 197 for B-Victoria-like, 191 for B-Yamagata-like, 303 for H7N9 and +Infinity for
H5Nx.
682
Models: HKY model is a substitution model that considers different base frequencies and
assigns different rates for transitions v.s. transversions. The **c model** represents SRD06 codon
position model. The partitioning strategy is to analyze codon positions $1 + 2$ and codon position
3 separately. The **p model** takes the protein structure partitions into account based on the amino
acid positions for each domain on the linear diagram. The **cp model** combines both c and p
models, which estimates codon positions in protein structural partitions. All models use HKY
substitution model.

Figure 4. Log scale Bayes Factors for sensitivity analysis on full and subsampled HPAI

H5Nx datasets. All these Bayes Factors (BF) are calculated to compare with HKY via path-

sampling (upper panel) and stepping-stone sampling (lower panel) approaches to compute the

marginal likelihood estimates for each model. The asterisk * represents a significant supportive

BF for the structurally informed codon model (cp), compared to SRD06 codon only model (c

model). Both cp and c models perform better than HKY model with decisive BF supports. The

HKY plus structurally informed model (p model) performs better with supportive BFs in all

datasets.

Log scale BF criteria: BF< 5 represents no significance; $5 \leq$ BF< 10 means substantial support;

$10 \leq$ BF< 15 means strong support; $15 \leq$ BF < 20 means very strong support; BF ≥ 20 means

decisive support.

Models: HKY model is a substitution model that considers different base frequencies and

assigns different rates for transitions v.s. transversions. The **c model** represents SRD06 codon

position model. The partitioning strategy is to analyze codon positions $1 + 2$ and codon position

3 separately. The **p model** takes the protein structure partitions into account based on the amino

acid positions for each domain on the linear diagram. The **cp model** combines both c and p

models, which estimates codon positions in protein structural partitions. All models use HKY

substitution model.

Figure 5. Sensitivity analysis: overall mean substitution rates and domain-specific
substitution rates from structurally informed partitioning model for HPAI H5Nx
subsampled datasets. Violin plots show the substitution rates from each step of the Bayesian
simulations. Only 20% dataset has the trend to slightly exaggerate the rates and would not report
the precise evolution history of H5Nx. The model superiority generally holds in the sensitivity
analysis and the model is not sensitive to sample size.
719
720 **Models:** The **cp model** combines both c and p models, which estimates codon positions in
721 protein structural partitions. It uses HKY substitution model.