Title: Identifying Important Interaction Modifications in Ecological Systems.

Authors:

J. Christopher D. Terry (Corresponding Author: james.terry@zoo.ox.ac.uk)[1]

Michael B. Bonsall[1,2]

Rebecca J. Morris[1,3]

[1] Department of Zoology, University of Oxford, OX1 3PS, UK;

[2] St. Peter's College, New Inn Hall Street, Oxford, OX1 2DL UK;

[3] Biological Sciences, Faculty of Natural and Environmental Sciences, Life Sciences Building 85,

University of Southampton, Highfield Campus, Southampton, SO17 1BJ, UK.

Abstract

Trophic interaction modifications, where a consumer-resource link is affected by additional

species, are widespread and significant causes of indirect effects in ecological networks. The

sheer number of potential interaction modifications in ecological systems poses a considerable

challenge, making prioritisation for empirical study essential. Here, we introduce measures to

quantify the topological relationship of individual interaction modifications relative to the

underlying network. We use these, together with measures for the strength of trophic

interaction modifications to identify modifications that are most likely to exert significant

effects on the dynamics of whole systems. Using a set of simulated food webs and randomly

distributed interaction modifications, we test whether a subset of interaction modifications

important for the local stability and directional determinacy of complex networks can be

identified. We show that trophic interaction modifications from low-density species, those

affecting high-flux interactions, those that connect species otherwise distantly linked, and those

that act from high trophic level modifiers to interactions lower in the web have particular

importance for dynamics. In contrast, the centrality of modifications provided little information.

This work demonstrates that analysis of interaction modifications can be tractable at the

network scale and highlights the importance of understanding the relationship between trophic

and non-trophic effect distributions.

Key words: interaction modification; food web; non-trophic effect; predictability; trait-

mediated indirect effect; trophic interaction; ecological networks;

Introduction

The study of the network dynamics of whole ecological communities is dominated by trophic

interactions (Dunne and Pascual 2006). However, it is increasingly recognised that non-trophic

effects must be considered to improve our understanding of the dynamics of ecosystems (Kéfi

et al. 2012). Interaction modifications (Wootton 1993), where third-party species influence the

strength of interactions, are a significant cause of non-trophic effects and are known to be

widespread in ecological communities (Werner and Peacor 2003, Kéfi et al. 2015). By grouping

the diverse processes that can be represented as modifications of functional responses, the

trophic interaction modification (TIM) approach (Golubski and Abrams 2011, Terry et al. 2017)

provides a route to investigate a significant portion of non-trophic effects.

Studies in small empirical model systems have demonstrated that the impact of TIMs can be

considerable, comparable with trophic interactions (Schmitz et al. 2004), and capable of driving

community dynamics (van Veen et al. 2005). Interaction modifications have often been

identified as the cause of unexpected ecosystem responses to perturbation (Doak et al. 2008,

Peckarsky et al. 2008, Tack et al. 2011). However, despite growing interest (Ohgushi et al. 2012,

Levine et al. 2017), applications to complex natural systems have been slower to develop (Kéfi

et al. 2012). The number of possible interaction modifications rises rapidly with increasing

community size. Identification and prioritisation of the most important processes (Aufderheide

et al. 2013) is therefore necessary to accelerate the improvement of our understanding of

ecological networks.

Interaction modifications do not fit directly within conventional network theory (Newman

2010) as they cannot be depicted as a set of nodes linked by edges. Instead, the nodes directly

affect edges. Approaches to deal with this discontinuity within conventional network theory

have generally taken two approaches. Most commonly, studies focus on resultant pairwise

effects, drawing conventional 'non-trophic interaction' links between the modifier and both

interactors (Kéfi et al. 2012) which can be analysed as multi-layer networks (Pilosof et al. 2017)

while keeping non-trophic and trophic networks distinct. Alternatively, hyper-graph approaches

can be used where edges can link any number of nodes (Golubski et al. 2016). Both these

approaches have significant drawbacks in terms of maintaining the distinct features of

interaction modifications (Terry et al. 2017).

Here we introduce analogues of conventional network metrics that can describe how individual

interaction modifications are situated within the wider trophic network. Using simulated food

webs with random distributions of trophic interaction modifications, we show that it is possible

to detect a valuable subset of interaction modifications, the identification of which can offer

significant gains in directional determinacy and the estimation of community stability.

Methods

Generating Communities

We generated a set of *in silico* communities, using empirically derived allometries, to which we added a random set of TIMs and conducted our dynamic analyses. We describe these systems at an equilibrium by modelling the interactions between each population with a community matrix (Novak et al. 2016). All analyses were carried out using R v. 3.4.3 (R Core Team 2017) and all code is available in Supplementary Information.

Initial trophic network topologies were generated using the niche model algorithm (Williams and Martinez 2000) with species number and trophic connectance set at 25 and 0.1, respectively. This simple algorithm has been generally validated as a useful approach to generate networks that have similar structural properties to natural systems (Williams and Martinez 2008). Cannibalistic interactions were removed and trophic levels were assigned from the network structure following the unweighted pathway method of Levine (1980). Species body-masses, m_i, were assigned based on trophic level following empirical distributions of consumer-resource ratios (Brose et al. 2006, detailed in SI 1).

The biomass density of each species, B_i, was drawn from a distribution based on the individual body mass as $B_i = 10^{-2.36+g+\epsilon} m_i^g$ following the approach of Tang et al. (2014), where $\epsilon \sim \mathcal{N}(0, \sigma = 0.2)$. The scaling factor between body mass and biomass-density, g, can vary considerably between communities (Reuman et al. 2009). We therefore repeat our analyses with three values of g within the range of empirical estimates (0.25, 0.1 and -0.1). To

ease comparison between different scaling terms, the mean density across each community

was standardised to 1.

Trophic Interactions

To define the effect of trophic interactions between populations, we used body-mass

allometries (Yodzis and Innes 1992, Woodward et al. 2005) to parameterise a Type I functional

response model:

91
$$\frac{dB_i}{dt} = \sum_{i \subset \text{prey of } i}^{i} e_{ij} a_{ij} B_i B_i - \sum_{j \subset \text{predators of } i}^{j} a_{ij} B_i B_j \qquad \text{(Eq. 1)}$$

This allows us to specify a community matrix **A**, where each element $\mathbf{A}_{ij} = \frac{\partial(dB_i/dt)}{\partial B_j}$, detailing

the change in the growth rate of the population of i in response to a small change in the

population of j.

Each element of the community matrix representing the trophic effect of the population of

predator j on a population of prey i is specified as $\mathbf{A}_{ij} = -a_{ij} B_i$ while the reverse effect,

$\mathbf{A}_{ji} = e_{ij} a_{ij} B_j$. Conversion efficiencies, e_{ij}, were drawn from a uniform distribution $\mathcal{U}(0.1, 0.2)$.

Biomass-density specific attack rates, a_{ij}, were defined based on the documented relationship

between consumer-resource size ratios and per-capita attack rates (Pawar et al. (2012) ,

detailed in SI 1) and incorporating a penalty for predator generality.

Self-regulation terms

The diagonal elements of the community matrix A_{ii} represent the self-regulation of each

species. We tested two approaches to specify these terms. First, each species was assigned a

negative self-regulation term equal to twice the mean of the absolute size of interactions

exerted upon it. In a second approach, producers were assumed to be more strongly self-

regulating than consumers. Each producer was assigned terms equal to twice the strongest

trophic interaction in the network while each consumer was assigned a term 100 times smaller.

TIMs

We model interaction modifications as a multiplicative term, where the resultant attack rate is

dependent on the relative divergence of the modifier species density, B_k, from their

equilibrium:

$$a'_{ij} = a_{ij} \left(1 + \sum_{k \in modifiers}^{k} c_{ijk} \log \left(\frac{B_k}{B_k^*} \right) \right)$$

. (Eq. 2)

Species are assumed to not modify their own trophic interactions ($k \neq i, j$), or rather any effect

is considered subsumed into the self-regulatory terms. A positive slope parameter, c_{ijk}, causes

an increased level of the modifier to strengthen the underlying trophic interaction. A negative

value would weaken the interaction. Each TIM results in two additional non-trophic effect

terms to be added to the community matrix:

$$\frac{\partial B_i}{\partial B_k} = \ldots - a_{ij} B_i B_j c_{ijk} \frac{1}{B_k^*} \; , \quad \frac{\partial B_j}{\partial B_k} = \ldots + e_{ij} a_{ij} B_i B_j c_{ijk} \frac{1}{B_k^*}$$

. (Eq. 3)

We assume that the trophic interaction specified by the allometries incorporates the influence

of the modifier at the equilibrium density, i.e. it is already at 'post-modification' strength. This

allows the effect of interaction modifications to be clearly distinguished from disruption of the

trophic network, something that has not been possible in previous dynamic analyses of the

effect of interaction modifications on system dynamics (e.g. Arditi et al. 2005, Goudard and

Loreau 2008, Lin and Sutherland 2013) where the introduction of interaction modification shifts

the underlying trophic interaction distribution.

Each potential TIM (combination of consumer, resource, and third-party species) had an equal

and independent probability (0.05) of being present. This resulted in an expectation of 69 TIMs

per community of 25 species and trophic connectance 0.1. Slope parameters (c_{ijk}) for each

extant TIM were drawn from a uniform distribution $\mathcal{U}(-0.5, 0.5)$. The resultant log-normal

distribution of non-trophic effect components largely overlapped that of the trophic

interactions but tended to be weaker (SI. 2).

Quantifying TIMs

Topology

We quantify the location of a TIM within a trophic network in seven aspects of topology,

illustrated in Fig. 1. The simplest method records the *trophic height* of the modifier species and

of the interaction (the mean trophic level of the two interactor species). Here, we calculate

trophic level using the unweighted trophic network and the pathway method of Levine (1980).

These can then be compared to measure what we term the '*trophic direction*' of the

modification. We define this as the trophic height of the interaction minus the trophic level of

the modifier. Therefore, positive values imply modifications to interactions higher in the web

than the modifier and negative values imply the opposite. The '*trophic span*' of a modification

constitutes the mean number of trophic links from the modifying species to each interactor by

the shortest route and measures the extent to which a TIM links otherwise disconnected

species.

The *centrality* of the species involved in a TIM can be used to determine the connectivity of the

TIM to the wider trophic network. The centrality of a node within a network can be measured

by a diversity of algorithms (Newman 2010). Here we use *'betweenness'*, the number of

shortest paths between all nodes that pass through a given node. We only consider unweighted

trophic links (excluding TIM connectance) when calculating centrality as this information is

more widely available. We measure the *'interactor betweenness'* as the mean of the centrality

of the two interactors. Lastly, the centrality of the interaction being modified was measured as

the *'interaction betweenness'*, the number of shortest paths that pass through a given

interaction.

Metric	Modification:	
	·······•	– – – •
Trophic Height of Modifier	3	2
Trophic Height of Interaction	2.5	3.5
Trophic Span	1	2
Trophic Direction	-0.5	+1.5
Modifier Betweenness	0	24.5
Interactor Betweenness	19.5	3.75
Interaction Betweenness	17.5	7.5

Figure 1. Example values of topological metrics for interaction modifications (dashed lines) within a simple trophic

network (solid lines).

Strength of TIMs

Interaction strengths can be simultaneously considered with multiple and useful approaches

(Laska and Wootton 1998, Berlow et al. 2004). This extends to TIMs which can be meaningfully

considered from a range of perspectives (Terry et al. 2017). We measure the strength of a

modification in our linearised equilibrium systems in four distinct ways (Table 1). Here we solely

measure strength with respect to the prey species since in our simple model the impact on the

predator is very closely related to the impact of prey. In addition, we test the information

contained in the density of the modifier species and the size of the underlying trophic

interaction as a proxy for interaction modification strength, as measured by the attack rate,

and the biomass flux rate,   .

Table 1: Description of the TIM strength metrics used to rank TIMs in the model systems. In each case *i,j* and *k*

indices denote the prey, predator and modifier species respectively.

Metric	Description	General Specification	Calculation
TIM Function Slope	The strength of the relationship between the modifier species density and the interaction rate.	$$\frac{\delta}{\delta B_k} a_{ij}$$	
Direct Non-Trophic Effect of TIM	The contribution of an individual TIM to the effect from the modifier to an interactor. Also described as	$$\left| \mathbf{A}_{ik}^{Full} - \mathbf{A}_{ik}^{c_{ijk}=0} \right|$$	$$a_{ij} B_i B_j |c_{ijk}| \frac{1}{B_k^*}$$

	the trait-mediated indirect effect (Abrams 2008, Okuyama and Bolker 2012).		
Impact on Interaction	The change in the direct interaction strength caused by the TIM. This is a measure of instantaneous effect of a TIM.	$\left\| \dfrac{\delta}{\delta B_k} \mathbf{A}_{ij} \right\|$	$a_{ij} B_i \|c_{ijk}\| \dfrac{1}{B_k^*}$
Impact on Net-Effects between Interactors.	The change in the 'net-effect' between species as the modifier species density changes. Contextualises the role the trophic interaction plays in the overall interaction between species.	$\left\| \dfrac{\delta}{\delta B_k^{mod}} [\mathbf{A}^{-1}]_{ij} \right\|$	Calculated numerically changing only the density of the modifier term in the TIM function to isolate the TIM effect from those mediated by other interaction chains through the modifier species.

Assessment of Metrics

The ability of metrics to identify influential TIMs was assessed by testing how well a system

containing only a subset of interaction modifications that had been selected using a metric

under examination was able to capture the dynamics of the complete system (Figure 2). We

assess dynamics in two ways: 1) accuracy in estimating local stability and 2) the ability to

determine the direction of response to sustained perturbations.

Figure 2. Schematic showing overall approach to test methods for identifying interaction modifications that are

influential in the dynamics of model ecological communities. The trophic interaction modifications in the full

community model are quantified and 20% retained using a particular metric. The capacity for the subset model to

represent the dynamics of the dull model system are then assessed stability estimation and directional

determinacy.

Stability

Local asymptotic stability, whether a system will eventually return to its previous equilibrium

after a small perturbation, is a widely used dynamic system property and can be determined

from the sign of the real part of the largest eigenvalue of a system's community matrix,   .

Where this is negative the system can be considered locally stable. With sufficient self-

regulation any system can be stabilised. To focus our analysis on the impact of the interaction

modifications, rather than the distribution of self-regulation terms (van Altena et al. 2016,

Barabás et al. 2017), we set each $\mathbf{A}_{ii}$ to 0. In this case, $\Re\left(\lambda_1^{\mathbf{A}}\right)$ is always positive but can be

interpreted as a degree of the self-regulation that would be necessary to stabilise the

community. For simplicity we will refer to this quantity as 'stability'.

We generated 2000 communities with interaction modifications for each value of biomass

density scaling factor g. For each web, the TIM metrics described in the previous section were

calculated for each TIM and the 'top' 20% of TIMs were identified. Where there was no obvious

expectation about which extreme would be more important, for example, centrality, both the

highest and lowest values were tested independently. A random subset of 20% of TIMs was also

generated for each web to establish a baseline to test metrics against. For each web and each

subset of TIMs, the community matrix was recalculated including the trophic interactions and

the selected TIMs. We then calculated the absolute difference in $\log(\Re\left(\lambda_1^{\mathbf{A}}\right))$ between the full

model and the subset model. The statistical significance of a metric's ability to identify a subset

of TIMs that estimate stability differently to a random subset of TIMs was tested with a

Wilcoxon signed rank test, paired by the underlying web topology.

Direction of response to perturbation

The negative inverse of a community matrix $(-\mathbf{A}^{-1})$ encapsulates the responses (via all

pathways, interaction chains and interaction modifications) of each species to a sustained,

small, perturbation to every other species (Bender et al. 1984) to generate a 'net-effects

matrix'. This relies on the assumption that the perturbation is sufficiently small that any non-

linearities in the response are well represented by the linear approximation of the underlying

community matrix. While estimates of the exact values of responses would be empirically

unfeasible, the *sign* of the net-effects between species, i.e. whether a change in a particular

population will lead to an increase or a decrease in another population is a reasonable objective

(Yodzis 1988, Novak et al. 2011, Aufderheide et al. 2013).

We tested the proportion of pairwise net-effect relationships between species where the sign

of the net-effects matrix of the full model matches that of the model containing only a subset

of the TIMs. This directional determinacy, also termed 'predictability' (Novak et al. 2011, Iles

and Novak 2016) therefore ranges from a minimum of approximately 0.5, where the subset

model is no better than chance at correctly identifying species responses, to 1 where the subset

model is able to perfectly predict the 'true' response. Analytic expressions of the impact on

directional determinacy due to the misspecification of **A** have recently been derived (Koslicki

and Novak 2018). However, since for multiple error terms the expressions become very

complex, we directly calculate the effect numerically.

Because local asymptotic stability is a prerequisite for calculating directional determinacy,

additional communities were generated and unstable communities excluded to create 2000

locally stable communities for each of the three values of g and the two approaches to

including self-regulation terms. We then followed the same procedure as for local stability to

test each metric's ability to identify valuable TIMs. For each web we compared the directional

determinacy of a model including the selected subset of interaction modifications to a model

including just the trophic interactions.

Results

Figure 3. Comparison of the capacity of different metrics to identify TIMs that contribute to the dynamics of whole

systems. Arrows indicate whether TIMs were selected for inclusion based on high or low values of the given

property. Median error in the estimation of the logarithm of the system stability is plotted on the x-axis, median

improvement in the percentage of direction of inter-specific net effects correctly estimated on the y-axes. Grey

dotted lines show a baseline when TIMs are chosen at random. Those metrics in the top-right quadrant are

therefore better than average at identify TIMs valuable for both types of dynamics, while those in the lower-left

quadrant are worse for both. Points marked with symbols were not significantly different to a random draw for the

stability (+), the directional determinacy (x) or both (*) Wilcoxon test (n=2000, $\alpha = 0.05$), paired by underlying

network.

It was possible to identify the trophic interaction modifications that influenced the system

dynamics. In general, those TIMs affecting directional determinacy also affected the local

stability of the system. Results for the case of g=0.1 and stronger self-regulation terms on

producers are shown in Figure 3. Results for other values of body mass-biomass density scaling

and different self-regulation terms (SI 3) were broadly similar except in specific cases discussed

below. The variability in changes to directional determinacy was related to the median

improvement – where a metric tended to give a larger improvement, there was also a greater

chance that including those TIMs would result in a reduced model that was considerably worse

than not including any TIMs at all (SI 3). This effect was not noticeable for stability estimation.

The strength properties of TIMs were correlated with each other but topological features

considerably less so (Figure 4, SI 6). Interaction flux was still negatively correlated with

interaction trophic level when g=0.1. The positive body-mass - density relationship was

counteracted by reduced attack rates from higher consumer resource body-mass ratios at

higher trophic levels.

Figure 4. Correlation plot showing relationships between the properties of the TIMs in a model food web

community (g = 0.1, producers higher self-regulation than consumers). Ellipses depict Spearman's rank

correlations, blue shapes indicating positive correlations and red negative. Corresponding plots for the other

model structures are shown in SI 6.

In general, those metrics that addressed aspects of the strength of the interaction modification

were the most valuable (shown in green in Fig. 3) and the very best allowed a near perfect

estimation of stability. Metrics based on the centrality of the interaction modifications were the

least effective at identifying TIMs, with many not being significantly different to a random draw.

Modifications caused by species with a lower biomass density were more valuable than those

with high density.

Interaction modifications with large slope parameters of either sign were valuable. However,

including just the large strengthening (+ c_{ijk}) interaction modifications gave a better estimate

of the local stability than the strongly negative interaction modifications. This difference was

particularly notable for the g=-0.75 case (SI 3). Conversely, selecting interaction modifications

for directional determinacy but not for stability.

The influence of trophic height depended on the relationship between trophic level and

biomass density (controlled by the g parameter) with modifications of high-trophic level

interactions becoming comparatively more important when the webs were more top-heavy.

The impact of trophic direction was highly dependent on the distribution of self-regulation

terms. Where producers are more strongly self-regulated than consumers, 'upwards' TIMs are

less influential for directional determinacy.

<u>Discussion</u>

We have presented methods to quantify the structure of interaction modifications within a

trophic network and shown that it is possible to identify properties of trophic interaction

modifications that are particularly influential in the dynamics of model ecosystems. This is

informative both on a theoretical level in helping to discern how TIMs affect dynamics, but also

is useful from a practical perspective in highlighting which TIMs might be most valuable to

study empirically. Our work also highlights areas where empirical information is lacking, in

particular the need to understand empirical relationships between trophic and non-trophic

interaction strengths.

TIMs introduce a high level of complexity to ecosystem dynamics. It is therefore reassuring to

note that relatively simple and easily observable measures, for example interaction flux, are

able to identify the TIMs that drive the system dynamics. Similar sets of TIMs were identified as

being important in two distinct types of dynamics – response to press perturbations (directional

determinacy) and response to pulse perturbations (local stability). By identifying comparatively

high-impact TIMs we can make progress despite the present dearth of empirical distributional

data. Below we focus our discussion on specific results and their implications, before looking

forward to developing this approach in future work.

This study required assumptions regarding the appropriate strength and distribution of TIM

values to use. In particular, our results regarding interaction strength are necessarily dependent

on the range of the distribution from which the various properties the model are chosen from.

If for instance the range of slope parameters was larger compared to the range of biomass

densities, the c_{ijk} parameter in our model would have been a more valuable in comparison to

the other TIM strength parameters. Empirical distributions of interaction modification strength

in a network do not (to our knowledge) yet exist. Meta-analyses of non-trophic effect strength

(Bolnick and Preisser 2005, Preisser et al. 2005) suggest an approximate correspondence with

trophic interactions. However, the occasional very strong non-trophic effect at the upper tail of

the distribution we used (SI 2) is not necessarily unrealistic as the impact of behavioural effects

can in certain circumstances be much faster and stronger than consumptive interactions

(Abrams 2001). At the other end of the spectrum, it is necessary to set a threshold where a

species' influence on an interaction is considered too weak to include. This decision trades-off

the mean strength and the overall connectance of the modification distribution. An additional

complication is how interaction modifications combine to influence an interaction (Golubski

and Abrams 2011). For example, interaction modifications may act antagonistically to each

other, with combined modifications exerting less effect than would be expected from the

individual effects. Although strengths in our study this effect is mitigated by our assumption

that the trophic interactions are at the post-modification strength this could be significant in

more dynamic, non-equilibrium, systems.

The increased impact of low-density modifiers via interaction modifications runs contrary to

usual expectations for interaction strength distributions. In our study this results from our

assumption of the dependence on *relative* change in modifier populations (Eq. 2). Hence, a

change in the density of a low-density modifier will result in a greater modification than the

same absolute change in a more abundant modifier. Whether this is a reasonable assumption is

to return to the question of how the strength of TIMs is distributed in natural communities. This

effect is opposed by trophic interactions being stronger between high-density populations,

leading to opposing effects in mixed-process interaction chains. Consider a case where A

consumes B which then modifies an interaction involving C. In our model, if B had a lower

density, the A-B trophic interaction would be weaker, but this would be more than outweighed

by an increase in the B~C non-trophic effect. Further work will be needed to discern how these

processes act in nature, but this result highlights that TIMs can greatly increase the importance

of species that might otherwise be neglected or dismissed and may disrupt assumptions of the

structural patterns of interaction strengths in webs based on trophic interactions only.

Interpretation of the value of topological measures is more challenging as a significant part of

their effect can be attributed to correlations with strength-based metrics. In this study we

considered topology based on trophic interactions only, as these are currently more widely

available and better understood. However, interaction modifications themselves are part of the

dynamic topology of TIMs and ideally should be considered in relation to each other.

The most valuable topological metric was the 'direction' of the TIM. Those TIMs acting 'up' the

network were generally more valuable, although the reduced signal with more negative body

mass – density relationships suggests this is due to the relationship between trophic height and

density rather than pure topological effects. Nonetheless, strongly self-regulated producers

may be more resistant to change and hence modifications deriving from them may well be less

influential. This highlights that the variability or sensitivity of a species is of equal importance as

the magnitude of the modification when considering likely impact.

Trophic net-effects tend to rapidly decline with the number of trophic links between the

interactors (Neutel et al. 2002). Where the trophic net-effect is weak, there is a greater

potential for a TIM to 'short-circuit' the system and dominate the pairwise net-effect between

those species. This can be seen in the value of the TIM span metric for directional determinacy,

however the comparatively poor potential suggests that other significant features of TIMs may

be able to override this expectation. Although TIMs have the potential to reduce the dynamic

breadth (the number of interactions between the most distant species) of food webs, given that

it is estimated that the majority of species are no more than two trophic links apart (Williams et

al. 2002) the extent of this effect may be limited in practice.

The poor indicative potential of TIM span for local stability, shows that the increased

connectance that long-range TIMs has little impact on local stability, as simple models may

predict (May 1973). The powerful strengthening TIMs (that benefit the consumer to the

detriment of the resource) influence the stability more than weakening TIMs. This is a distinct

effect to that caused by mutualism or competitive direct interactions (e.g. Mougi and Kondoh

2012, Coyte et al. 2015) since every TIM exerts both a negative and a positive influence. This

effect could be driven by at least two mechanisms. Firstly, in our model the assumption of top-

down dominated interaction strength imbalance caused by the assimilation term e causes non-

trophic effects on prey to be larger than on predators (Eq. 3). This has the result that

strengthening TIMs ($+c_{ijk}$) exert larger negative effects (on prey) than positive effects (on

predators). Secondly, there will be a topological effect – strengthening TIMs will lead to

consistently positive non-trophic effects on high trophic level species and consistently negative

impacts on basal species, which will interact with the underlying tropic interaction distribution.

Outlook

For this study we made a number of assumptions about the form, topology and strength of

TIMs to construct a plausible distribution. The lack of empirical data at a community level is a

particular challenge, although identifying non-trophic effects at the network level is possible

(Pocock et al. 2012, Kéfi et al. 2015) and shows that networks of pairwise non-trophic

interactions have considerable structure (Kéfi et al. 2016). Our study emphasises the tight

relationship between the underlying trophic network and the consequences of interaction

modifications. Isolating an interaction modification from its context, for example in a lab study,

will result in the loss of valuable information. The properties of TIMs identified here (Fig. 5) are

a first step in building a profile that could be used to identify processes that are more likely to

have a considerable influence. Without empirical estimates for the distribution of TIMs it is not

yet possible to quantify the potential impact in the manner of Novak et al. (2011).

Figure 5. Properties of interaction modifications of relevance to determining their impact.

The instances where directional determinacy actually reduced on the inclusion of TIMs (SI 3)

shows the challenges in attempting to improve ecological understanding. Our results suggest

that there are no easy shortcuts, but that the problem is not intractable. One avenue that will

need particular work is the degree of accuracy to which interaction modifications must be

known in order to contribute meaningfully to improving ecological forecasts (Petchey et al.

2015). Our assumption that all pairwise interaction strengths are known, and that included

TIMs are perfectly quantified, constitutes rather wishful thinking for any system of realistic size.

Although for small systems, qualitative analyses (Dambacher et al. 2003, Dambacher and

Ramos-Jiliberto 2007) can be useful for mapping out the potential responses of systems, for

larger systems they can rapidly become indeterminate. Exploratory analyses (SI 7) testing the

importance in correctly identifying the slope of the TIM function show that as long as the sign

of the modification is correct, choosing either random slopes or a fixed value for the slope can

give improvements of mean directional determinacy of 10.5% and 15%, respectively. Future

work should more robustly test alternative approximation approaches, such as 'binning' effects

into broad categories, that have been shown to give good results for estimating trophic

interaction strengths (Barabás and Allesina 2015).

Equilibrium studies such as ours allow a precise analysis of particular processes in isolation from

other sources of variation. However, this approach does not incorporate the full spectrum of

impacts TIMs can have in dynamic systems and future work should seek to address this gap.

Furthermore, trophic interaction strengths are not purely determined by biotic factors – there

is also extensive dependence on the abiotic environment (Wootton and Emmerson 2005, Poisot

et al. 2015). The balance of the impact of exogenous and endogenous sources of interaction

strength variation is unknown. The two processes can interact, with TIMs varying depending on

abiotic conditions (Sentis et al. 2017) and interaction modifications allowing species to resist

abiotic variability (Bruno et al. 2003). The extent to which interaction strength variation will

frustrate ecological prediction is unclear. However, at least some of this variation can be

accounted for by dependence on other species in the community which, in principle, is

quantifiable (Terry et al. 2017).

Conclusion

To improve our understanding of ecosystems it will be necessary to incorporate important

dynamic processes whether they are consumer-resource links, direct non-trophic effects or

interaction modifications (Fontaine et al. 2011). This potential complexity poses a daunting

challenge but not all interaction modifications are equal. We have shown that considering

interaction modifications as distinct entities opens up a range of analyses and that although the

underlying mechanisms may be complex, relatively simple heuristics may be useful in

identifying significant processes in food webs. Where interaction modifications are suspected,

but unquantified, our work provides guidance to determine where further research could bring

the greatest improvement in understanding. Fully incorporating interaction modifications into

the fold of ecosystem ecology will be a long, but rewarding, endeavour. We hope that our work

will encourage further studies, both experimental and theoretical, to understand these critical

components of ecosystem dynamics.

Data Statement

All code and raw simulation outputs are available on the Open Science Framework:

https://osf.io/49jgr/

Acknowledgements:

JCDT is funded by the Natural Environmental Research Council through the Oxford

Environmental Research Doctoral Training Program (NE/L002612/1). RJM was funded by a

Royal Society University Research Fellowship.

References

Abrams, P. A. 2001. Describing and Quantifying Interspecific Interactions: A Commentary on

Recent Approaches. - Oikos 94: 209–218.

Abrams, P. A. 2008. Measuring the impact of dynamic antipredator traits on predator-prey-

resource interactions. - Ecology 89: 1640–1649.

Arditi, R. et al. 2005. Rheagogies: Modelling non-trophic effects in food webs. - Ecol. Complex.

2: 249–258.

Aufderheide, H. et al. 2013. How to predict community responses to perturbations in the face

of imperfect knowledge and network complexity. - Proc. R. Soc. London B Biol. Sci. 280:

20132355.

Barabás, G. and Allesina, S. 2015. Predicting global community properties from uncertain

estimates of interaction strengths. - J. R. Soc. Interface 12: 20150218.

Barabás, G. et al. 2017. Self-regulation and the stability of large ecological networks. - Nat. Ecol.

Evol. 1: 1870–1875.

Bender, E. A. et al. 1984. Perturbation Experiments in Community Ecology: Theory and Practice.

- Ecology 65: 1–13.

Berlow, E. L. et al. 2004. Interaction Strengths in Food Webs: Issues and Opportunities. - J.

Anim. Ecol. 73: 585–598.

Bolnick, D. I. and Preisser, E. L. 2005. Resource competition modifies the strength of trait

mediated predator–prey interactions: a meta-analysis. - Ecology 86: 2771–2779.

Brose, U. et al. 2006. Consumer-resource body-size relationships in natural food webs. - Ecology

87: 2411–2417.

Bruno, J. F. et al. 2003. Inclusion of facilitation into ecological theory. - Trends Ecol. Evol. 18:

119–125.

Coyte, K. Z. et al. 2015. The ecology of the microbiome: Networks, competition, and stability. -

Science 350: 663–666.

Dambacher, J. M. and Ramos-Jiliberto, R. 2007. Understanding and Predicting Effects of

Modified Interactions Through a Qualitative Analysis of Community Structure. - Q. Rev.

Biol. 82: 227–250.

Dambacher, J. M. et al. 2003. Qualitative predictions in model ecosystems. - Ecol. Modell. 161:

79–93.

Doak, D. F. et al. 2008. Understanding and predicting ecological dynamics: are major surprise

inevitable? - Ecology 89: 952–961.

Dunne, J. A. and Pascual, M. 2006. Ecological Networks: Linking Structure to Dynamics in Food

Webs. - Oxford University Press.

Fontaine, C. et al. 2011. The ecological and evolutionary implications of merging different types

of networks. - Ecol. Lett. 14: 1170–1181.

Golubski, A. J. and Abrams, P. A. 2011. Modifying modifiers: what happens when interspecific

interactions interact? - J. Anim. Ecol. 80: 1097–1108.

Golubski, A. J. et al. 2016. Ecological Networks over the Edge: Hypergraph Trait-Mediated

Indirect Interaction (TMII) Structure. - Trends Ecol. Evol. 31: 344–354.

Goudard, A. and Loreau, M. 2008. Nontrophic Interactions, Biodiversity, and Ecosystem

Functioning: An Interaction Web Model. - Am. Nat. 171: 91–106.

Iles, A. C. and Novak, M. 2016. Complexity Increases Predictability in Allometrically Constrained

Food Webs. - Am. Nat. 188: 87–98.

Kéfi, S. et al. 2012. More than a meal… integrating non-feeding interactions into food webs. -

Ecol. Lett. 15: 291–300.

Kéfi, S. et al. 2015. Network structure beyond food webs: mapping non-trophic and trophic

interactions on Chilean rocky shores. - Ecology 96: 291–303.

Kéfi, S. et al. 2016. How Structured Is the Entangled Bank? The Surprisingly Simple Organization

of Multiplex Ecological Networks Leads to Increased Persistence and Resilience. - PLOS

Biol. 14: e1002527.

Koslicki, D. and Novak, M. 2018. Exact probabilities for the indeterminacy of complex networks

as perceived through press perturbations. - J. Math. Biol. 76: 877–909.

Laska, M. S. and Wootton, J. T. 1998. Theoretical Concepts and Empirical Approaches to

Measuring Interaction Strength. - Ecology 79: 461–476.

Levine, S. 1980. Several measures of trophic structure applicable to complex food webs. - J.

Theor. Biol. 83: 195–207.

Levine, J. M. et al. 2017. Beyond pairwise mechanisms of species coexistence in complex

communities. - Nature 546: 56–64.

Lin, Y. and Sutherland, W. J. 2013. Color and degree of interspecific synchrony of environmental

noise affect the variability of complex ecological networks. - Ecol. Modell. 263: 162–173.

May, R. M. 1973. Stability and Complexity in Model Ecosystems. - Princeton University Press.

Mougi, A. and Kondoh, M. 2012. Diversity of Interaction Types and Ecological Community

Stability. - Science 337: 349–351.

Neutel, A.-M. et al. 2002. Stability in Real Food Webs: Weak Links in Long Loops. - Science 296:

1120–1123.

Newman, M. 2010. Networks: An Introduction. - Oxford University Press.

Novak, M. et al. 2011. Predicting community responses to perturbations in the face of

imperfect knowledge and network complexity. - Ecology 92: 836–846.

Novak, M. et al. 2016. Characterizing Species Interactions to Understand Press Perturbations:

What Is the Community Matrix? - Annu. Rev. Ecol. Evol. Syst. 47: 409–432.

Ohgushi, T. et al. 2012. Trait-Mediated Indirect Interactions: Ecological and Evolutionary

Perspectives. - Cambridge University Press.

Okuyama, T. and Bolker, B. M. 2012. Model-based, response-surface approaches to quantifying

indirect interactions. - In: Ohgushi, T. et al. (eds), Trait-Mediated Indirect Interactions:

Ecological and Evolutionary Perspectives. Cambridge University Press, pp. 186–204.

Pawar, S. et al. 2012. Dimensionality of consumer search space drives trophic interaction

strengths. - Nature 486: 485–489.

Peckarsky, B. L. et al. 2008. Revisiting the Classics?: Considering Nonconsumptive Effects in

Textbook Examples of Predator – Prey Interactions. - Ecology 89: 2416–2425.

Petchey, O. L. et al. 2015. The ecological forecast horizon, and examples of its uses and

determinants (D Vasseur, Ed.). - Ecol. Lett. 18: 597–611.

Pilosof, S. et al. 2017. The multilayer nature of ecological networks. - Nat. Ecol. Evol. 1: 101.

Pocock, M. J. O. et al. 2012. The robustness and restoration of a network of ecological

networks. - Science 335: 973–7.

Poisot, T. et al. 2015. Beyond species: why ecological interaction networks vary through space

and time. - Oikos 124: 243–251.

Preisser, E. L. et al. 2005. Scared to death? The effects of intimidation and consumption in

predator-prey interactions. - Ecology 86: 501–509.

R Core Team 2017. R: A Language and Environment for Statistical Computing. in press.

Reuman, D. C. et al. 2009. Allometry of Body Size and Abundance in 166 Food Webs. - Adv. Ecol.

Res. 41: 1–44.

Schmitz, O. J. et al. 2004. Trophic cascades: the primacy of trait-mediated indirect interactions. -

Ecol. Lett. 7: 153–163.

Sentis, A. et al. 2017. Predator diversity and environmental change modify the strengths of

trophic and nontrophic interactions. - Glob. Chang. Biol. 23: 2629–2640.

Tack, A. J. M. et al. 2011. Can we predict indirect interactions from quantitative food webs? - an

experimental approach. - J. Anim. Ecol. 80: 108–118.

Tang, S. et al. 2014. Correlations between interation strengths drives stability in large ecological

networks - Supp Info. - Ecol. Lett. 17: 1094–1100.

Terry, J. C. D. et al. 2017. Trophic interaction modifications: an empirical and theoretical

framework. - Ecol. Lett. 20: 1219–1230.

van Altena, C. et al. 2016. Patterns in intraspecific interaction strengths and the stability of food

webs. - Theor. Ecol. 9: 95–106.

van Veen, F. J. F. et al. 2005. Stable Coexistence in Insect Communities Due to Density- and

Trait-Mediated Indirect Effects. - Ecology 86: 3182–3189.

Werner, E. E. and Peacor, S. D. 2003. A Review of Trait-Mediated Indirect Interactions in

Ecological Communities. - Ecology 84: 1083–1100.

Williams, R. J. and Martinez, N. D. 2000. Simple rules yield complex food webs. - Nature 404:

180–183.

Williams, R. J. and Martinez, N. D. 2008. Success and its limits among structural models of

complex food webs. - J. Anim. Ecol. 77: 512–519.

Williams, R. J. et al. 2002. Two degrees of separation in complex food webs. - Proc. Natl. Acad.

Sci. 99: 12913–12916.

Woodward, G. et al. 2005. Body size in ecological networks. - Trends Ecol. Evol. 20: 402–409.

Wootton, J. T. 1993. Indirect Effects and Habitat Use in an Intertidal Community: Interaction

Chains and Interaction Modifications. - Am. Nat. 141: 71–89.

Wootton, J. T. and Emmerson, M. 2005. Measurement of Interaction Strength in Nature. -

Annu. Rev. Ecol. Evol. Syst. 36: 419–444.

Yodzis, P. 1988. The Indeterminacy of Ecological Interactions as Perceived Through Perturbation

Experiments. - Ecology 69: 508–515.

Yodzis, P. and Innes, S. 1992. Body Size and Consumer-Resource Dynamics. - Am. Nat. 139:

1151–1175.