Running Head: Key Interaction Modifications

Title: Identifying Important Interaction Modifications in Ecological Systems.

Type: Article

Authors:

J. Christopher D. Terry (Corresponding Author: james.terry@zoo.ox.ac.uk)[1]

Michael B. Bonsall[1,2]

Rebecca J. Morris[1,3]

[1] Department of Zoology, University of Oxford, OX1 3PS, UK;

[2] St. Peter's College, New Inn Hall Street, Oxford, OX1 2DL UK;

[3] Biological Sciences, Faculty of Natural and Environmental Sciences, Life Sciences Building 85,

University of Southampton, Highfield Campus, Southampton, SO17 1BJ, UK.

Abstract

Trophic interaction modifications, where a consumer-resource link is affected by additional

species, are widespread and significant causes of indirect effects in ecological networks. The

sheer number of potential interaction modifications in ecological systems poses a significant

challenge, making prioritisation for empirical study essential. We use the trophic interaction

modification framework to identify modifications that are most likely to exert significant effects

on the dynamics of whole systems, introducing tools to quantify the topological relationship of

interaction modifications relative to the underlying network. Using a set of simulated food

webs and randomly distributed interaction modifications we test whether a subset of

interaction modifications important for determining the local stability and directional

determinability of complex networks can be identified. We show that trophic interaction

modifications from rare species, those affecting high-flux interactions, those that connect

species otherwise distantly linked, and those that act from high trophic level modifiers to

interactions lower in the web have particular importance for dynamics. This work demonstrates

that analysis of interaction modifications can be tractable at the network scale and highlights

the importance of the inter-relationship between the structure of trophic networks and

interaction modifications.

Key words: interaction modification; food web; non-trophic effect; predictability; trait-

mediated indirect effect; trophic interaction; ecological networks;

Introduction

The study of ecological network dynamics has been dominated by trophic interactions (Dunne

and Pascual 2006), but it is increasingly recognised that to improve our understanding of

ecosystems non-trophic effects must be considered (Kéfi et al. 2012). Interaction modifications

(Wootton 1993), where non-interactor species alter the strengths of links, are a significant

cause of non-trophic effects and are known to be widespread in ecological communities

(Werner and Peacor 2003, Kéfi et al. 2015). By grouping the diverse effects that can be

represented as modifications of functional responses, the trophic interaction modification (TIM)

approach (Golubski and Abrams 2011, Terry et al. 2017) provides a route to investigate a

significant portion of non-trophic effects.

Studies in small model systems have shown that the effects of TIMs can be considerable,

comparable with trophic interactions (Schmitz et al. 2004, Bolnick and Preisser 2005), and

capable of driving community dynamics (van Veen et al. 2005). Interaction modifications have

often been identified as the cause of unexpected ecosystem responses to perturbation (Doak et

al. 2008, Tack et al. 2011). However, despite growing interest (Ohgushi et al. 2012, Levine et al.

2017) applications to complex natural systems have been slower to develop (Kéfi et al. 2012).

The number of possible interaction modifications rises rapidly with increasing community size.

Identification and quantification of the most important processes (Novak et al. 2011) is

therefore necessary to improve our understanding of ecological networks.

Here we introduce methods to examine the topological structure of interaction modifications in

food webs. Using simulated food webs with random distributions of trophic interaction

modifications we show that it is possible to detect a valuable subset of interaction

modifications, the identification of which can offer significant gains in predictability and

estimation of stability.

Methods

Food web model specification

We used artificial communities at a steady-state equilibrium as a baseline for our analysis. To

generate these, initial community trophic network topologies were generated using the niche

model algorithm (Williams and Martinez 2000) with species number and trophic connectance

fixed at 25 and 0.15. This simple algorithm has been generally validated as a useful approach to

generate networks that have similar structural properties to natural systems (Williams and

Martinez 2008). Trophic levels were assigned from the network structure following the pathway

method of Levine (1980). The system dynamics were modelled and parameterised following the

Allometric Trophic Network (ATN) model approach (Williams and Martinez 2004, Brose et al.

2006b, Delmas et al. 2016), which uses metabolic body-size scaling relationships to parametrise

bio-energetic differential equation models (Eq. 1). This model was chosen because of its wide

use, limited number of parameters and successful application to experimental data (Boit et al.

2012). The model is of the form:

1)
$$\frac{dB_i}{dt} = r_i B_i G_i - x_i B_i + \sum_{j\, \in\, prey} x_i y_{ij} F_{ij} B_i - \sum_{j\, \in\, predators} x_j \frac{y_{ji}}{e_{ji}} F_{ji} B_j$$

70   2)
$$F_{ij} = \frac{\omega_i B_j^2}{B_0^2 + B_j^2}$$

71   3)
$$G_i = 1 - \frac{B_i}{K}$$

where all parameters and definitions are given in Table 1. Non-trophic dynamic links in the

standard ATN model were excluded as they could obscure the TIM effects under study. Hence,

the alternative prey interference terms in the functional response (Koen-Alonso 2007) were not

incorporated, since prey switching is a cause of TIMs (Abrams 2010). Instead a direct type III

functional response was used (Eq. 2). A Type III response was chosen to increase the likelihood

of reaching a stable equilibrium rather than a limit cycle or chaotic attractor. Predators with

multiple prey had a reduced attack rate on each determined by ω, a predation choice term,

which was set at 1/n, where n is the number of prey species the predator consumes. Producer

growth was modelled without interspecific producer competition (Eq. 3), with each producer

having a separate, identical, carrying capacity K. Species body-masses M_i were assigned based

on trophic level following empirical consumer-resource ratios (Brose et al. 2006a, detailed in SI

1). Model parameters were assigned values following previous studies wherever possible (Table

1).

Community population densities were determined by initiating each species at a biomass of 0.1

and running each system to a steady state before the inclusion of interaction modifications (see

SI 1). Of 1000 initial webs, 797 successfully reached a steady-state equilibrium density with a

complete network of at least 10 extant species (density greater than 10^{-12}). These were used to

form the set of baseline systems. Summary statistics of the structure of the model communities

are given in SI 1.

Adding TIMs

TIMs were represented with a function that links the density of the modifier species with the

functional response (Eq. 4). Here we used an extension of the sigmoidal function suggested by

Kéfi et al. (2012), parameterised such that at the equilibrium density of the modifier the

resultant modification term is 1 (i.e. there is no change to the interaction strength at the

equilibrium point) and the point of inflexion of the function is at the equilibrium point:

4) $$F'_{ij} = F_{ij} \prod_{k \, \in \, modifiers}^{k} \left(M_0 + \frac{M_{max} - M_0}{(1 + e^{-c_{ijk}(log_{10}(B_k) - log_{10}(B_k^*))})^{1/\nu_k}} \right) \quad , \quad \nu_k = \frac{log(2)}{log(\frac{M_{max}}{1 - M_0})} \, .$$

Through the use of this parameterisation and by conducting analyses at a fixed set of species

densities, the distribution of trophic interaction strengths is maintained. This allows the effect

of interaction modifications to be clearly distinguished from disruption of the trophic network,

something that has not been possible in previous dynamic analyses of the effect of interaction

modifications on system dynamics (e.g. Arditi et al. 2005, Goudard and Loreau 2008, Lin and

Sutherland 2013). Since the Jacobian-based analyses we conduct are linearisations, an

equivalent linear TIM function would have the same result, however this sigmoidal function

allows consistency with additional analyses using extended dynamics. The TIM slope parameter

(c_{ijk}) controls the direction and magnitude of the modification. A positive c_{ijk} causes an

increased level of the modifier to increase the underlying trophic interaction while a negative

value results in a reduction in the interaction as the modifier density increases. The sign of the

realized functional response rate is always unaffected and so energy flow between species is

never reversed. Although reversals are possible (Barkai and McQuaid 1988), they are unlikely to

be widespread. The dependence on the logarithm of the modifier biomass ensures that

modifications are scaled to the relative change in density of the modifier.

When multiple TIMs affect a single interaction, they are combined multiplicatively. The most

prevalent combination of interaction modifications in nature is unknown, although there have

been suggestions that interaction modifications are likely to act antagonistically to each other

(Golubski and Abrams 2011), meaning that combined modifications have less effect than would

be expected from the individual effects. For this analysis, since the starting condition is

assumed to be the post-modification interaction strengths, the complications such

considerations might introduce is reduced.

TIMs were added so that each potential TIM (combination of an existing trophic interaction and

a third-party species) was present with probability of 0.1 (non-trophic connectance *sensu*

Goudard and Loreau (2008)). This choice of TIM connectance was arbitrary and would not

affect the results since the aim of the study is to determine relative importance of TIMs. TIMs

were assigned a slope parameter (c_{ijk}) drawn from a uniform distribution -0.3:0.3. This

distribution of slope parameters resulted in a distribution of Jacobian elements components

due to TIMs in approximately the range of trophic interactions (SI 2).

Empirical distributions of interaction modification strength in a network do not (to our

knowledge) yet exist although an approximate correspondence with trophic interaction

strengths is a reasonable estimate from meta-analyses (Preisser et al. 2005). In our simulations,

average direct pairwise TIM effects were smaller than trophic effects, but there was a notable

upper tail of interactions (SI 2). 3.17% of non-zero TIM-attributable Jacobian element

components had a magnitude larger than 1, the upper bound of trophic effects. This occasional

very strong direct effect caused by an interaction modification is not necessarily unrealistic as

the impact of behavioural effects can in certain circumstances be much faster and stronger

than consumptive interactions (Abrams 2001). Nonetheless, we repeated the analysis capping

such outlier interactions in line with the maximum trophic interaction element and found

similar results (SI 2).

Ranking by TIM metrics and Generation of Reduced Models

The TIM metrics listed in Table 2 and discussed below were calculated for each TIM.

Correlations between the TIM metrics across all the webs were assessed with Spearman's rank

correlation tests. For each web the TIMs were ranked according to each metric, using the

ranking methods detailed in Table 2. For each web, for each metric, the 'top' 30% of TIMs were

identified and a 'Reduced' system was generated including just those TIMs. A 'random' draw of

30% of TIMs was also generated for each web to establish a baseline. 30% was chosen as giving

a good spread between the metrics – a subset with too few TIMs for inclusion led to very little

change in predictability, while if too many were included the difference between the metrics

was less clear (SI 3). The Jacobian matrices (**J**) for each of the 'Full', 'Reduced' and 'No TIM'

models were calculated numerically at the equilibrium density with the *numDeriv* R package.

Strength of TIMs

It is well established that no single aspect of interaction strength is sufficient to categorise

trophic interactions completely (Laska and Wootton 1998, Berlow et al. 2004). Similarly the

strength of a TIM can be considered from a range of perspectives and with different valid

methods (Terry et al. (2017). Metrics applicable to the equilibrium systems studied here are:

TIM Slope parameter (c_{ijk}): The parameter in the model that relates the modifier species

density to the functional response relating the interactors.

Flux of Modified Interaction: The flow rate of biomass between the interactor species.

Direct Effect of modifier on interactors attributable to TIM: The derivative of the change in each

of the interactor growth rates with respect to the modifier density when using the full model,

minus the same value calculated without the presence of the TIM.

Derivatives of Jacobian elements representing the interaction with respect to the modifier:

Corresponds to the modifier's effect on the direct effect of the consumer on the resource at

equilibrium and vice versa.

Derivatives of the net-effects matrix elements representing the consumer-resource interaction

with respect to the modifier: Corresponds to the modifier's effect on the net- effect of the

consumer on the resource at equilibrium and vice versa.

Topology of TIMs

Interaction modifications do not fit directly within conventional network theory (Newman

2010) as they cannot be depicted as a set of nodes linked by edges. Instead, they link nodes to

edges. Approaches to deal with this discontinuity within conventional network theory have

generally taken two approaches. Most commonly, studies focus on resultant pairwise effects,

drawing conventional 'non-trophic interaction' links between the modifier and both interactors

(Kéfi et al. 2012). Multi-layer networks (Pilosof et al. 2017) may also be used to analyse these

while keeping non-trophic and trophic networks distinct. Alternatively, hyper-graph approaches

can be used where edges can link any number of nodes (Golubski et al. 2016). Both these

approaches have significant drawbacks in terms of maintaining the dynamic differences

between interaction modifications and pairwise interactions (Terry et al. 2017). Here we

introduce analogues of conventional network metrics that can describe how individual

interaction modifications sit within the wider trophic network illustrated in Fig. 1. Aggregate

metrics and approaches to consider the relationships of TIMs with each other are discussed in

SI 4.

Range of modification: The mean number of trophic links from the modifying species to each

interactor by the shortest route.

Trophic direction of modification: How the trophic level of the modifier compares to the trophic

level interactors. The direction of a TIM is calculated as the mean trophic level of the

interactors minus the trophic level of the modifier. Therefore, positive values imply

modifications to interactions higher in the web than the modifier, negative values imply that

species higher in the food web are modifying the interactions of those lower down.

Centrality of modifier and interactor species: The centrality of a species within a network can be

measured by a diversity of algorithms (Newman 2010) and be applied to TIMs by calculating the

centrality of the modifier species or the mean centrality of the interactors in terms of the their

trophic linkages. Two common measures for how central a node is within a web are

'betweenness', which assess how many shortest paths between nodes go through a given node

and PageRank, the 'Google algorithm' which assesses how much each node is linked to,

weighted by the importance of each linked node.

Betweenness of modified interaction: The centrality of the modified interaction can be assessed

by the edge-betweenness of the interaction – how often the shortest path between nodes goes

via a given edge.

Assessment of predictability

Predictability (Novak et al. 2011, Iles and Novak 2016) is defined as the directional consistency

in how species population changes in response to press perturbations (Yodzis 1988).

Predictability was calculated as the fraction of matching signs of the net effects matrix ($-\mathbf{J}^{-1}$)

between the 'Full' model and the 'Reduced' model. Ten systems were found to be uninvertible

– these were removed from further consideration. To standardise across webs we calculated

the predictability when using the 'No TIM' model and report the difference in predictability

between the 'Reduced' model and the 'No TIM' model. The statistical significance of a metric's

ability to identify important TIMs differently to a random draw was tested with a Wilcoxon

signed rank test paired by underlying web.

Assessment of TIM metrics for stability

Local stability, whether the system will return to its previous equilibrium after a small

perturbation, can be determined from the sign of the real part of the largest eigenvalue of a

system's Jacobian matrix. Where this is negative the system can be considered locally stable.

Stability was assessed for each 'Full' and 'Reduced' Model. If the sign of the dominant

eigenvalue of the 'Reduced' model matched the 'Full' model it was deemed to successfully

categorise local stability. The ability of each metric to identify valuable sets of TIMs was

assessed with a binomial test with an expected value 0.82, the fraction correctly identified by

random draws of included TIMs

Alternative Network structures

To isolate TIM topology from other attributes, we repeated the predictability analysis using

underlying networks more restricted structure. Species body masses, parameters and

equilibrium densities were set to fixed values to equalise trophic direct effects (SI 5). Web

topologies (S=20, n=500) were created using either the niche model (C=0.1) or a simple 'grid'

model. The grid model consisted of four trophic levels each with 5 species, where each had 'in'

and 'out' degree of 2 as appropriate to its trophic level. Parameters were chosen to either

maintain the asymmetry between resource effect on consumer and consumer effect on

resources, or with all direct interactions fixed to a constant value regardless of direction.

Results

Predictability

Non-topological metrics

The capacity of all of the metrics to identify TIMs that are important for predictability was

significantly different from random draws (Fig. 2, results of significance tests given in SI 6)

although the divergence varied greatly. The variability in changes to predictability was related

to the median improvement – where a metric tended to give a larger improvement in

predictability, there was also a greater chance that including those TIMs would result in a

reduced model that was considerably worse than the No TIM model. Combinations of metrics

were not particularly valuable - trying to optimise over multiple measures simultaneously did

not bring additional information. There were strong correlations between a number of the

metrics (see SI 7), clustering into topological and non-topological groups, although the two

groups are linked by a correlation between interactor trophic level and trophic interaction

strength (Spearman's ρ = -0.58).

Interaction flux was the best method for identifying the size of the underlying interaction,

better than the direct effects between the interactor species. Measures of the dependence on

the modifier species density of the interaction itself or the interactor species growth rate gave

amongst the greatest improvements in predictability. Dependence on the net-effects between

the interactors on the modifier density was less successful. The density of the modifier was

more valuable than the magnitude of the slope parameter c_{ijk} at providing useful information

about the TIM function specification. There was no significant difference between selecting

TIMs by the most positive or most negative c_{ijk} values (paired Wilcoxon signed-rank test

p=0.0651, n=787).

Topological Metrics

The most valuable topological metric in the main set was the 'direction' of the TIM (Fig. 2).

Those TIMs 'down' the network, where the modifier had a higher trophic level than the

interactors, were particularly important. As lower interactions were likely to have a larger flux

and higher trophic level species were likely to be rarer, the measure of direction is correlated

with both the modifier biomass and interaction flux (Spearman's ρ: 0.31 and -0.31 respectively).

Measures of centrality generally had poor comparative performance. The two measures of

centrality (node-betweenness and PageRank) were relatively uncorrelated with each other

(Spearman's ρ for modifiers: 0.33, for interactors -0.08). Approaches that identified TIMs

involving high centrality modifiers or low-centrality interactions were generally better than

random. However, effect size is weak in all sets of webs and effects are attributable to other

correlated aspects of TIMs that have stronger impacts. TIMs where the modifier and interactor

were widely trophically separated were more valuable than near TIMs, but difference in the

median predictability change was smaller than other metrics.

Results for the sets with fixed direct interaction strengths and species density are detailed in

Fig. 3. The four sets of networks each show different metrics as being able to identify distinct

and important groups of TIMs. In general, as the networks become more rigidly structured,

fewer topological metrics were able to identify subsets of TIMs consistently better than

random. Downwards acting TIMs from high trophic level modifier species were more impactful

in set B whereas upwards TIMs were more valuable in sets A and C. Distributional changes of

net effects with trophic level are shown in SI 1. TIM range was able to identify valuable TIMs in

the niche model topology case but not in the grid topology case.

Local Stability

Metrics varied considerably in their ability to identify a subset of TIMs that accurately

determined local stability by the sign of the dominant eigenvalue (Fig. 4, statistical significance

tests in SI 6). Metrics that were better at identifying stability gained the most by increasing the

proportion of unstable full systems that were correctly categorised as unstable with the

reduced model. Metrics that performed well at identifying important TIMs for predictability

were also valuable for identifying those that were important for categorising local stability

(Spearman's ρ = 0.96, SI 6).

Nonetheless there were some notable differences. Selecting TIMs for inclusion that reached

further across the web was significantly worse than selecting TIMs at random. There was also a

sharp difference between selecting TIMs of differing signs: including the 'most negative' TIMs

was better than random draws, successfully identifying more of the unstable systems, while

including the 'most positive' was less successful than random.

<u>Discussion</u>

We have presented methods to quantify the structure of interaction modifications within a

trophic network and shown that it is possible to identify properties of trophic interaction

modifications that are particularly influential in the dynamics of model ecosystems. This is

informative both on a theoretical level in helping to discern how TIMs affect dynamics, but also

from a practical perspective in highlighting which TIMs are most valuable to study empirically.

TIMs introduce a high level of complexity to ecosystem dynamics. It is therefore reassuring to

note that relatively simple measures are able to identify TIMs that are important for dynamics.

Similar sets of TIMs were identified as being important in two distinct types of dynamics –

response to press perturbations (predictability) and response to pulse perturbations (local

stability). By identifying relatively high-impact TIMs we can make progress despite the present

dearth of empirical distributional data. Below we focus our discussion on specific results and

their implications, before looking forward to developing this approach in future work.

Selecting TIMs by interaction flux was a better method than assessing the direct effects

between the interactors. This is a helpful result for experimentally identifying TIMs since

interaction flux is easier to estimate than dependence of the interactors on each other as it can

be directly observed rather than requiring experimental manipulation (Wootton and Emmerson

2005). The need to contextualise TIMs in the wider community is reiterated by some of the

greatest improvements in predictability deriving from measures of the dependence on the

modifier species density of the interaction itself or the interactor species growth rate. Both are

measures of instantaneous effect, the first from a TIM perspective (dependence of the

interaction on the modifier), the second taking a pairwise non-trophic approach (dependence

of the interactors on the modifier through trait-mediated indirect interactions (Abrams 2008,

Okuyama and Bolker 2012)).

The importance of the rarity of the modifier in our study can be traced back to dependence of

the modification function on the logarithm of modifier density. This has the effect that a change

in the density of a rare modifier will result in a greater modification than the same absolute

change in a more common modifier. Nonetheless, this dependence on relative changes seems a

reasonable assumption and this result shows that TIMs can greatly increase the importance of

species that might otherwise be neglected or dismissed. Furthermore, even in this linearised

system it can be seen that a simple description of the relationship between the modifier species

density and the interaction depends on both the starting density and a slope parameter.

Interpretation of the value of topological measures in the main set of webs is more challenging

as a significant part of their effect can be attributed to correlations with strength-based

metrics. The most valuable topological metric in the main set was the 'direction' of the TIM.

Those TIMs acting 'down' the network were particularly important, although trophic height is

correlated with both interaction flux and equilibrium density. However, a 'direction' effect can

also be observed in web sets A and B (where interaction flux and equilibrium is fixed)

demonstrating a distinct mechanism by which the direction of TIM can identify its importance.

The low improvement when selecting TIMs by centrality and the opposing effect direction in

sets A and B (with identical trophic topology), suggests the mechanism is not due to pure

network topological effects.

An explanation can be found from differences in the trophic net-effects affecting species at

different trophic heights. Although direct effects were held constant, the self-limitation terms

(Jacobian matrix diagonals) still vary between species. These are composed of contributions

from intrinsic rates of change (which differ between producers and consumers) as well as terms

from density-dependent predation rates leading to structural patterns in net-effects (SI 1.3).

Where mean net-effects increase with trophic height (Set B and the 'realistic' main set),

'downwards' TIMs can be seen to have a greater impact, driven by trophic height of the

modifier. Greater sensitivity of higher trophic level species to extinction provides some support

that this is a common natural relationship. Where the modifier species is more sensitive to

other species, the TIM can act as a stronger conduit for other species to spread their impact

indirectly through the modification. When mean net-effects decrease with trophic height (set

A) the reverse is true –TIMs from low trophic level modifiers have more of an impact.

Trophic net-effects tend to decline with the number of trophic links between the interactors

(Neutel 2002). Where the trophic net effect is weak, there is a greater potential for a TIM to

dominate the pairwise net-effect between those species and therefore be important for

predictability. This can be seen in the effect of range in the fixed webs. The comparatively poor

indicative potential of TIM range in the main set suggests that other significant features of TIMs

(discussed above) may be able to override this expectation, as there is a strong net-effects –

distance relationship. Although TIMs have the potential to greatly reduce the dynamic breadth

of food webs, given that it is estimated that the majority of species are no more than two

trophic links apart (Williams et al. 2002) the extent of this effect may be limited in practice.

In two particular cases the results from the stability estimation test diverged from those of

predictability. Range of TIM was notably poor at the estimation of stability – including just the

long-range TIMs lead to a preponderance of stable web estimates. This is surprising given that

long-range TIMs would be expected to contribute more to increasing dynamic connectivity and

consequent instability (May 1973). Selecting negative TIMs (that benefit the resource to the

detriment of the consumer) result in more stable estimates of systems, while selecting positive

TIMs lead to more estimates that the resultant system would be unstable. This is a distinct

effect to that caused by mutualism or competitive direct interactions e.g. (Mougi and Kondoh

2012, Coyte et al. 2015) since every TIM exerts both a negative and a positive influence and

should be investigated further.

Outlook

For this study we had to make a number of assumptions about the form, distribution and

strength of TIMs. The lack of empirical data at a community level is a particular challenge,

although identifying non-trophic effects at the network level is possible (Pocock et al. 2012, Kéfi

et al. 2015) and shows that networks of pairwise non-trophic interactions have considerable

structure (Kéfi et al. 2016). Our study emphasises the tight relationship between the underlying

trophic network and the consequences of interaction modifications. Isolating an interaction

modification from its context will result in the loss of valuable information. The five properties

of TIMs identified here (Fig. 5) build up a profile that could be used to identify processes that

are more likely to have a considerable effect on dynamics in response to perturbations.

One avenue that will need particular work is the degree of accuracy to which interaction

modifications must be known in order to contribute to improving ecological forecasts (Petchey

et al. 2015). Our assumption that all pairwise interaction strengths are known, and that

included TIMs are perfectly quantified, constitutes rather wishful thinking for any system of

realistic size. For small systems, qualitative analyses (Dambacher et al. 2003, Dambacher and

Ramos-Jiliberto 2007) can be useful for mapping out the potential responses of systems

although for larger systems they can rapidly become indeterminate. As a testament to the

importance of quantifying TIMs we conducted a randomisation study (SI 8) which showed

randomising slope parameter values while maintaining the sign yielded an average directional

predictability of 0.57, barely better than chance. Without empirical estimates for the

distribution of TIMs it is difficult to assess the potential impact on overall predictability in the

manner of Novak et al. (2011) but it is clear that there is the potential for a significant impact.

Future work should test alternative approximation approaches, such as 'binning' effects into

broad categories, that have been shown to give good results for estimating trophic interaction

strengths (Barabás and Allesina 2015).

Equilibrium studies such as ours allow a precise analysis of particular processes in isolation from

other sources of variation. However, this approach does not incorporate the full spectrum of

impacts TIMs can have in dynamic systems and future work should seek to address this gap.

Furthermore, trophic interaction strengths are not purely determined by biotic factors – there

is also extensive dependence on the abiotic environment. The balance of the impact of

exogenous impacts on a system compared to endogenous sources of interaction strength

variation is unknown. The two processes can interact, with TIMs varying depending on abiotic

conditions (Sentis et al. 2017) and interaction modifications allowing species to resist abiotic

variability (Bruno et al. 2003). The extent to which interaction strength variation will frustrate

ecological prediction is unclear. However, at least some of this variation can be accounted for

by dependence on other species in the community which, in principle, is quantifiable (Terry et

al. 2017).

Conclusion

To improve our understanding of ecosystems it will be necessary to incorporate important

dynamic processes whether they are consumer-resource links, direct non-trophic effects or

interaction modifications (Fontaine et al. 2011). This potential complexity poses a daunting

challenge but not all interaction modifications are equal. We have shown that considering

interaction modifications as distinct entities opens up a range of analyses and that although the

underlying mechanisms may be complex, relatively simple heuristics may be useful in

identifying significant processes in food webs. Where interaction modifications are suspected,

but unquantified, our work provides guidance to determine where further research could bring

the greatest improvement in understanding. Fully incorporating interaction modifications into

the fold of ecosystem ecology will be a long, but rewarding, endeavour. We hope that our work

will encourage further studies, both experimental and theoretical, to understand these critical

components of ecosystem dynamics.

Acknowledgements:

JCDT is funded by the Natural Environmental Research Council through the Oxford

Environmental Research Doctoral Training Program (NE/L002612/1). RJM was funded by a

Royal Society University Research Fellowship.

Literature Cited:

Abrams, P. A. 2001. Describing and Quantifying Interspecific Interactions: A Commentary on

Recent Approaches. Oikos 94:209–218.

Abrams, P. A. 2008. Measuring the impact of dynamic antipredator traits on predator-prey-

resource interactions. Ecology 89:1640–1649.

Abrams, P. A. 2010. Implications of flexible foraging for interspecific interactions: lessons from

simple models. Functional Ecology 24:7–17.

Arditi, R., J. Michalski, and A. H. Hirzel. 2005. Rheagogies: Modelling non-trophic effects in food

webs. Ecological Complexity 2:249–258.

Barabás, G., and S. Allesina. 2015. Predicting global community properties from uncertain

estimates of interaction strengths. Journal of The Royal Society Interface 12:20150218.

Barkai, A., and C. McQuaid. 1988. Predator-prey role reversal in a marine benthic ecosystem.

Science 242:62–4.

Berlow, E. L., A.-M. Neutel, J. E. Cohen, P. C. de Ruiter, B. Ebenman, M. Emmerson, J. W. Fox, V.

A. A. Jansen, J. I. Jones, G. D. Kokkoris, D. O. Logofet, A. J. McKane, J. M. Montoya, and O.

Petchey. 2004. Interaction Strengths in Food Webs: Issues and Opportunities. Journal of

Animal Ecology 73:585–598.

Boit, A., N. D. Martinez, R. J. Williams, and U. Gaedke. 2012. Mechanistic theory and modelling

of complex food-web dynamics in Lake Constance. Ecology Letters 15:594–602.

Bolnick, D. I., and E. L. Preisser. 2005. Resource competition modifies the strength of trait

mediated predator–prey interactions: a meta-analysis. Ecology 86:2771–2779.

Brose, U., T. Jonsson, E. L. Berlow, P. Warren, C. Banasek-Richter, L.-F. Bersier, J. L. Blanchard, T.

Brey, S. R. Carpenter, M.-F. C. Blandenier, L. Cushing, H. A. Dawah, T. Dell, F. Edwards, S.

Harper-Smith, U. Jacob, M. E. Ledger, N. D. Martinez, J. Memmott, K. Mintenbeck, J. K.

Pinnegar, B. C. Rall, T. S. Rayner, D. C. Reuman, L. Ruess, W. Ulrich, R. J. Williams, G.

Woodward, and J. E. Cohen. 2006a. Consumer-resource body-size relationships in natural

food webs. Ecology 87:2411–2417.

Brose, U., R. J. Williams, and N. D. Martinez. 2006b. Allometric scaling enhances stability in

complex food webs. Ecology Letters 9:1228–1236.

Brown, J. H., J. F. Gillooly, A. P. Allen, V. M. Savage, and G. B. West. 2004. Toward a Metabolic

Theory of Ecology. Ecology 85:1771–1789.

Bruno, J. F., J. J. Stachowicz, and M. D. Bertness. 2003. Inclusion of facilitation into ecological

theory. Trends in Ecology and Evolution 18:119–125.

Coyte, K. Z., J. Schluter, and K. R. Foster. 2015. The ecology of the microbiome: Networks,

competition, and stability. Science 350:663–666.

Dambacher, J. M., H. W. Li, and P. A. Rossignol. 2003. Qualitative predictions in model

ecosystems. Ecological Modelling 161:79–93.

Dambacher, J. M., and R. Ramos-Jiliberto. 2007. Understanding and Predicting Effects of

Modified Interactions Through a Qualitative Analysis of Community Structure. The

Quarterly Review of Biology 82:227–250.

Delmas, E., U. Brose, D. Gravel, D. B. Stouffer, and T. Poisot. 2016. Simulations of biomass

dynamics in community food webs. Methods in Ecology and Evolution.

Doak, D. F., J. A. Estes, B. S. Halpern, U. Jacob, D. R. Lindberg, J. Lovvorn, D. H. Monson, M. T.

Tinker, T. M. Williams, J. T. Wootton, I. Carroll, M. Emmerson, F. Micheli, and M. Novak.

2008. Understanding and predicting ecological dynamics: are major surprise inevitable?

Ecology 89:952–961.

Dunne, J. A., and M. Pascual. 2006. Ecological Networks: Linking Structure to Dynamics in Food

Webs. Oxford University Press, Oxford.

Fontaine, C., P. R. Guimarães, S. Kéfi, N. Loeuille, J. Memmott, W. H. van der Putten, F. J. F. van

Veen, and E. Thébault. 2011. The ecological and evolutionary implications of merging

different types of networks. Ecology Letters 14:1170–1181.

Golubski, A. J., and P. A. Abrams. 2011. Modifying modifiers: what happens when interspecific

interactions interact? Journal of Animal Ecology 80:1097–1108.

Golubski, A. J., E. E. Westlund, J. Vandermeer, and M. Pascual. 2016. Ecological Networks over

the Edge: Hypergraph Trait-Mediated Indirect Interaction (TMII) Structure. Trends in

Ecology & Evolution 31:344–354.

Goudard, A., and M. Loreau. 2008. Nontrophic Interactions, Biodiversity, and Ecosystem

Functioning: An Interaction Web Model. The American Naturalist 171:91–106.

Iles, A. C., and M. Novak. 2016. Complexity Increases Predictability in Allometrically Constrained

Food Webs. The American Naturalist 188:87–98.

Kéfi, S., E. L. Berlow, E. A. Wieters, L. N. Joppa, S. A. Wood, U. Brose, and S. A. Navarrete. 2015.

Network structure beyond food webs : mapping non-trophic and trophic interactions on

Chilean rocky shores. Ecology 96:291–303.

Kéfi, S., E. L. Berlow, E. A. Wieters, S. A. Navarrete, O. L. Petchey, S. A. Wood, A. Boit, L. N.

Joppa, K. D. Lafferty, R. J. Williams, N. D. Martinez, B. A. Menge, C. A. Blanchette, A. C. Iles,

and U. Brose. 2012. More than a meal… integrating non-feeding interactions into food

webs. Ecology Letters 15:291–300.

Kéfi, S., V. Miele, E. A. Wieters, S. A. Navarrete, and E. L. Berlow. 2016. How Structured Is the

Entangled Bank? The Surprisingly Simple Organization of Multiplex Ecological Networks

Leads to Increased Persistence and Resilience. PLOS Biology 14:e1002527.

Koen-Alonso, M. 2007. A Process-Oriented Approach to the Multispecies Functional Response.

Pages 1–36 *in* N. Rooney, K. S. McCann, and D. L. G. Noakes, editors.From Energetics to

Ecosystems: The Dynamics and Structure of Ecological Systems. Springer Netherlands.

Laska, M. S., and J. T. Wootton. 1998. Theoretical Concepts and Empirical Approaches to

Measuring Interaction Strength. Ecology 79:461–476.

Levine, J. M., J. Bascompte, P. B. Adler, and S. Allesina. 2017. Beyond pairwise mechanisms of

species coexistence in complex communities. Nature 546:56–64.

Levine, S. 1980. Several measures of trophic structure applicable to complex food webs. Journal

of Theoretical Biology 83:195–207.

Lin, Y., and W. J. Sutherland. 2013. Color and degree of interspecific synchrony of

environmental noise affect the variability of complex ecological networks. Ecological

Modelling 263:162–173.

May, R. M. 1973. Stability and Complexity in Model Ecosystems. Princeton University Press,

Princeton.

Mougi, A., and M. Kondoh. 2012. Diversity of Interaction Types and Ecological Community

Stability. Science 337:349–351.

Neutel, A.-M. 2002. Stability in Real Food Webs: Weak Links in Long Loops. Science 296:1120–

1123.

Newman, M. 2010. Networks: An Introduction. Oxford University Press, Oxford.

Novak, M., J. T. Wootton, D. F. Doak, M. Emmerson, J. A. Estes, and M. T. Tinker. 2011.

Predicting community responses to perturbations in the face of imperfect knowledge and

network complexity. Ecology 92:836–846.

Ohgushi, T., O. Schmitz, and R. D. Holt. 2012. Trait-Mediated Indirect Interactions: Ecological

and Evolutionary Perspectives. Cambridge University Press, Cambridge.

Okuyama, T., and B. M. Bolker. 2012. Model-based, response-surface approaches to quantifying

indirect interactions. Pages 186–204*in* T. Ohgushi, O. Schmitz, and R. D. Holt, editors.Trait-

Mediated Indirect Interactions: Ecological and Evolutionary Perspectives. Cambridge

University Press, Cambridge.

Petchey, O. L., M. Pontarp, T. M. Massie, S. Kéfi, A. Ozgul, M. Weilenmann, G. M. Palamara, F.

Altermatt, B. Matthews, J. M. Levine, D. Z. Childs, B. J. McGill, M. E. Schaepman, B. Schmid,

P. Spaak, A. P. Beckerman, F. Pennekamp, and I. S. Pearse. 2015. The ecological forecast

horizon, and examples of its uses and determinants. Ecology Letters 18:597–611.

Pilosof, S., M. A. Porter, M. Pascual, and S. Kéfi. 2017. The multilayer nature of ecological

networks. Nature Ecology & Evolution 1:101.

Pocock, M. J. O., D. M. Evans, and J. Memmott. 2012. The robustness and restoration of a

network of ecological networks. Science 335:973–7.

Preisser, E. L., D. I. Bolnick, and M. F. Benard. 2005. Scared to death? The effects of intimidation

and consumption in predator-prey interactions. Ecology 86:501–509.

Schmitz, O. J., V. Krivan, and O. Ovadia. 2004. Trophic cascades: the primacy of trait-mediated

indirect interactions. Ecology Letters 7:153–163.

Sentis, A., C. Gémard, B. Jaugeon, and D. S. Boukal. 2017. Predator diversity and environmental

change modify the strengths of trophic and nontrophic interactions. Global Change Biology

23:2629–2640.

Tack, A. J. M., S. Gripenberg, and T. Roslin. 2011. Can we predict indirect interactions from

quantitative food webs? - an experimental approach. Journal of Animal Ecology 80:108–

118.

Terry, J. C. D., R. J. Morris, and M. B. Bonsall. 2017. Trophic interaction modifications: an

empirical and theoretical framework. Ecology Letters 20:1219–1230.

van Veen, F. J. F., P. D. van Holland, and H. C. J. Godfray. 2005. Stable Coexistence in Insect

Communities Due to Density- and Trait-Mediated Indirect Effects. Ecology 86:3182–3189.

Werner, E. E., and S. D. Peacor. 2003. A Review of Trait-Mediated Indirect Interactions in

Ecological Communities. Ecology 84:1083–1100.

Williams, R. J., E. L. Berlow, J. A. Dunne, A.-L. Barabási, and N. D. Martinez. 2002. Two degrees

of separation in complex food webs. Proceedings of the National Academy of Sciences

99:12913–12916.

Williams, R. J., and N. D. Martinez. 2000. Simple rules yield complex food webs. Nature

404:180–183.

Williams, R. J., and N. D. Martinez. 2004. Stabilization of chaotic and non-permanent food-web

dynamics. The European Physical Journal B - Condensed Matter 38:297–303.

Williams, R. J., and N. D. Martinez. 2008. Success and its limits among structural models of

complex food webs. Journal of Animal Ecology 77:512–519.

Wootton, J. T. 1993. Indirect Effects and Habitat Use in an Intertidal Community: Interaction

Chains and Interaction Modifications. The American Naturalist 141:71–89.

Wootton, J. T., and M. Emmerson. 2005. Measurement of Interaction Strength in Nature.

Annual Review of Ecology, Evolution, and Systematics 36:419–444.

Yodzis, P. 1988. The Indeterminacy of Ecological Interactions as Perceived Through Perturbation

Experiments. Ecology 69:508–515.

Table 1: Parameters used in the main set of simulations.

Parameter	Value	Meaning and source
B_i	(from integration)	Species Density
K	5	Producer carrying capacity
e	0.45 (herbivory) 0.85 (carnivory)	Assimilation efficiencies (Brose et al. 2006b)
y	10	Maximum ingestion rate (Boit et al. 2012)
B_0	1	Half-saturation density
x_i	$a_i M_i^{-0.25}$	Species-specific metabolic rate following ¼ power-law (Brown et al. 2004)
a_i	Producers: 1 Trophic Level 2: 0.314 Trophic Level 3+: 0.88	Allometric constant (Brose et al. 2006b)
r	1	Producer growth rate
M_{max}	10	Greatest increasing modification
M_0	0	Greatest reducing modification
c_{ijk}	-0.3 : 0.3	TIM slope parameter. Impact of modifier k on the consumption of i by j.
B_k^*	(from integration)	Modifier equilibrium density

552 Table 2: Metrics used to rank TIMs to determine inclusion in reduced model.

	Metric to Rank by	**Ranking Method**
Strength	TIM parameter c_{ijk}	Magnitude, most +, most -
	Interaction flux	Largest
	Direct effect of modifier on interactors, due to TIM	Largest magnitude , effect on prey & effect on predator
	Derivative of direct effect between interactors w.r.t modifier	Largest magnitude. Effect of prey on predator and vice-versa
	Derivative of net- effect between interactors w.r.t modifier	Largest magnitude. Effect of prey on predator and vice-versa.
	Equilibrium density of modifier	Highest, Lowest
Topological	Distance	Largest (furthest), smallest (nearest)
	Direction of TIMs	Most upwards, most downwards
	Mean PageRank of interactors	Highest, Lowest
	Mean betweenness of interactors	Highest, Lowest
	PageRank centrality of modifier	Highest, Lowest
	Betweenness centrality of modifier	Highest, Lowest
	Betweenness of interaction	Highest, Lowest
	Mean trophic level of interactors	Highest, Lowest
	Trophic level of modifier	Highest, Lowest
Combinations	Largest flux & modifier rarity	Average rank across metrics
	Largest flux, modifier rarity & downwardsness	Average rank across metrics

Figure 1. Example of metrics to describe the topology of trophic interaction modifications

relative to the underlying trophic network.

Figure 2. Change in directional predictability when 30% of trophic interaction modifications

(TIMs) selected by different methods are included in a reduced model compared to a No-TIM

model. Red line demarks median predictability increase when drawing TIMs at random. All

metrics gave predictability results significantly different to random draws (paired Wilcoxon test,

$p<0.05$, n= 787. See SI 6 for full results).

Figure 3. Results from webs with fixed direct interaction strengths and equilibrium densities.

Change in directional predictability when 30% of trophic interaction modifications (TIMs)

selected by different methods are included in a reduced model compared to a no-TIM model.

Each panel is standardised to the median change in predictability when drawing TIMs at

random form comparison. Colour indicates significance of difference in predictability compared

to the random draws (paired Wilcoxon test, green = $p<0.05$, blue = $p>0.05$, n= 500)

Figure 4. Match between estimate of local stability from a reduced model including TIMs

selected by different metrics and the stability of the full model. Black line shows which metrics

did not correctly estimate local stability (shown in blue) significantly differently to the random

draw (Binomial test, expectation = 0.82, n=787, $\alpha = 0.05$. Full results of significance tests are in

SI 6.)

Figure 5. Properties of trophic interaction modifications that tend to increase their impact on

system dynamics.

Figure 1.

	Modification:	
Metric:	Blue	Brown
Range	1	2
Direction	-0.5	+1.5
Centrality of Modifier	0	24.5
Centrality of Interactors	19.5	3.75
Modified Interaction Betweenness	17.5	7.5

577   Figure 2.

580   Figure 3.

 Figure 4.

585   Figure 5