Antiretroviral treatment, prevention of transmission, and modeling the HIV epidemic: why the ART efficacy and effectiveness parameter matters

Reuben Granich[1], Somya Gupta[2], Matt Wollmers[3], Brian Williams[4]

1) Independent Public Health Consultant, San Francisco, California, United States
2) Independent Public Health Consultant, Delhi, India
3) Independent Public Health Consultant, Washington D.C., United States
4) South African Centre for Epidemiological Modelling and Analysis (SACEMA), Stellenbosch, South Africa

Address correspondence and reprint requests to:

Reuben Granich, MD, MPH
San Francisco, California
Phone: (202) 615-6913
Email: rgranich@gmail.com

Introduction

Despite progress, HIV is still a major public health threat with over 75 million deaths, 2 million infections a year and over 1 million HIV-associated TB cases a year[1, 2]. The impact of the HIV epidemic prompted an unprecedented response and we now know more about HIV than any other pathogen in history. The discovery of effective antiretroviral treatment (ART) in 1996 and subsequent evidence regarding the prevention of illness, death and transmission transformed the epidemic from an unending, unmitigated disaster into something that could be prevented and even ended someday[3-9,10]. The potential for expanded access to ART to curb the epidemic and the concept of treatment as prevention was introduced in 2006[5] and later formalized by WHO[11]. Over the past decade, the global and local HIV strategy has shifted from "test and wait" to "test and treat" focusing on achieving the 90-90-90 target (73% of people on ART and virally suppressed) and ending AIDS (defined as universal treatment, one AIDS case and one AIDS related death per 1000 population) [12 13, 14].

Policy discussions around treatment-as-prevention focused on how to increase access to testing and how early to provide diagnosis and treatment. Global HIV leadership and stakeholders turned to modeling to answer key questions about the risks and benefits of expanding access to treatment. Before and after the prevention impact of ART was understood, researchers used models to explore the impact of possible ART expansion scenarios[15-19]. While a few models explored expansion of ART access beyond existing WHO guidelines, most models limited testing and treatment to those who were severely immunocompromised[19]. Similarly, traditional costing efforts could be classified as "doomsday costing" insofar as they took a health sector perspective and only looked at the costs of providing earlier treatment while ignoring the potential prevention benefits and cost savings of earlier treatment[20, 21]. More modern "second generation" approaches to economic modeling took into account the prevention impact of scaling up treatment along with other interventions[18,22-24]. These models explored treatment as prevention of illness, death, and transmission. In some cases, the models were combined with a costing framework to examine the costs, cost benefits, and costs savings of various scale-up scenarios[22-25]. The dominant model used by UNAIDS, the Global Fund and the United States government and many countries now includes the prevention impact of ART and is used to determine the health and transmission impact, costs and cost-savings for various HIV response scenarios[23]. The UNAIDS model

incorporates data from available surveys and other surveillance information and makes forward projections of incidence, prevalence and resources needs according to their financial framework categories[23]. In the 2016 UNAIDS HIV Update and in the UNAIDS resource needs projections, the incidence and prevalence were reported as stable for the 5 years from 2010 to 2015 in all regions of the world except for Eastern Europe where the rates are increasing[1, 23]. During this time the world spent an estimated US$200 Bn on attempting to control HIV and the conclusion that can be drawn from the report is that the significant investment has had little or no impact on incidence or mortality. If true, this has major implications for future resource needs as well as for global HIV control strategy since the impact of treatment appears to be far less than expected. However, there are reasons to question the flat-line UNAIDS estimates of incidence and prevalence as these results contrast with other models and the scientific evidence regarding the potential impact of ART and other prevention interventions. Recent population-based studies from a number of countries suggest a marked decline in incidence, prevalence and deaths, mostly likely due to treatment expansion, in many countries in East and Southern Africa[26].

Methods

The marked contrast in model outcomes, one predicting flat-line incidence and prevalence in four of five regions[1, 23] while others projecting a steady decline to elimination, prompted us to explore the importance of ART efficacy, effectiveness and coverage parameters. Specifically, for 2015 and 2016 we reviewed global and national mathematical modeling studies that looked at impact of ART (with or without other HIV prevention interventions) OR 90-90-90 on either new HIV infections or investment or both. We reviewed these HIV epidemiologic and costing models for their structure and parameterization around ART.

Results

Table 1 describes the 9 models including the available parameters used to derive ART effectiveness expressed in terms of percentage reduction in HIV transmission. The modeling parameters for ART effectiveness by 2020 ranged from 20% to 95% for ART effectiveness. This disparity in ART Effectiveness is further highlighted in Figure 1 that shows the comparison between the GOALS and SACEMA models for Mozambique (other model comparisons not shown). The GOALS model gives a more pessimistic prediction of the trend in incidence and

mortality than does the SACEMA mode in the left panel of the figure. However, the GOALS model limits eligibility for ART initiation and assumes that at full coverage 80% of those infected with HIV and with a CD4+ cell count below 350 cells/μL. With these assumptions about half of all those infected with HIV will be on ART. The SACEMA model, on the other hand, assumes that we reach 90-90-90 by 2020 and then continue to roll-out ART at the same rate while ART reduces transmission by 96%; this gives a much more optimistic forecast. If we apply the SACEMA model but assume that under full coverage 65% of those infected with HIV are on ART and ART reduces transmission by 65% the SACEMA model gives results that are very close to the GOALS model. Clearly, these two parameters are the critical determinants of the impact of treatment on incidence and mortality. Comparisons with other models show similar results with the ART effectiveness parameter driving outcomes.

Discussion

Modeling plays an important role in exploring potential outcomes while also illustrating what data are missing for more accurate predictions. While parametrization relies on studies and other surrogate information, it is critical to be clear about the judgements involved in selecting critical values for interventions such as ART coverage and effectiveness. Studies now show that over 90% of people will agree to an HIV test if it is offered, that people started on ART adhere to a very high level, that ART suppresses the virus and reduces transmission by close to 100%, retention is probably much higher than predicted, costs have fallen significantly, other interventions have some value but are not cost-drivers, and high levels of ART and viral suppression can be achieved. Being clear about what parameters to use and how they are applied to the epidemiology and costing is critical when using models to guide the HIV response. Our brief review suggests that many of the current models have likely underestimated the impact of ART. This overly conservative assessment could account for past decisions to under invest in expanding access to treatment while using remaining resources for other budget categories[27]. Using more realistic parameters around ART suggests that as we expand access and support the achievement of sustainable viral suppression it will be possible to significantly reduce transmission and eliminate HIV in many settings.

Table 1: Description of Models with Calculated ART Effectiveness

STUDY	SETTING	PARAMETERS	ART EFFECTIVENESS
Williams (SACEMA model), Current HIV AIDS Research 2015 Epidemiological Trends for HIV in Southern Africa: Implications for Reaching the Elimination Targets	Southern Africa	• Proportion on ART virally suppressed: 90% • Reduction in transmission on ART * virally suppressed 96% * not virally suppressed 88% • Full coverage (2020) People at risk tested on average twice a year and started on treatment immediately.	**Effectiveness: 95%** **Calculation:** 1-(0.9*0.04+0.1*0.12) =95%
Smith (Imperial model), Lancet HIV 2016 Maximizing HIV prevention by balancing the opportunities of today with the promises of tomorrow: a modeling study	South Africa	• Efficacy (protection afforded by perfect use of a product) of early ART: **85%** (source is Cohen NEJM 2011) • Effective coverage (proportion of people who fully adhere to a product such that they benefit from its protection): Constant: 0%; medium: 40%; maximum: **60%**	**Effectiveness: 51%** **Calculation:** 1-(0.6x0.15+0.4) = 51%
McGillen (Imperial model), Lancet HIV 2016 Optimum resource allocation to reduce HIV incidence across SSA: a mathematical modeling study	18 countries from sub-Saharan Africa (80% of adult HIV burden in the region)	• Effectiveness of early ART as prevention (reduction in risk of onward transmission): **70%** (reports that figure is used in UNAIDS Fast track report 2014) Note: Early ART refers to a prevention method comprising outreach testing programmes and the offer of treatment to all PLHIV. This method is distinct from the assumption of provision of treatment for PLHIV who present for care (typically with low CD4 cell counts) • Achievable coverage: 33% among heterosexual men and low-risk women and 66% among MSM and FSW (this is the	**Effectiveness: 29%** **Calculation:** 1-(0.63x0.7+0.27) = 29%

STUDY	SETTING	PARAMETERS	ART EFFECTIVENESS
McGillen (Imperial model) -continued		coverage of early ART for PLHIV who have not already presented for care i.e. their CD4 is above 200 initially and above 350 later when ART eligibility criteria shifted) • Author's reply: When early ART is selected, then the proportion of people living with HIV who are virally suppressed reaches **63%**	
Stover (GOALS model), PLoS ONE 2016 What Is Required to End the AIDS Epidemic as a Public Health Threat by 2030? The Cost and Impact of the Fast-Track Approach	45 countries (86% of new infections globally)	• 95% reduction in infectiousness among those virally suppressed (source: Cohen NEJM 2011) • Adult ART 2020 coverage: 81% (90% started, 90% retained); 90% of them are virally suppressed • Adult ART 2030 coverage: 90% (95% started, 95% retained); 95% of them are virally suppressed • Eligibility for treatment expands to all PLHIV by 2018	**Effectiveness: 81%** **Calculation:** 81% (in 2020)
Kripke (DMPPT 2.1 model), PLoS ONE 2016 Impact and Cost of Scaling Up Voluntary Medical Male Circumcision for HIV Prevention in the Context of the New 90-90-90 HIV Treatment Targets	Lesotho, Malawi, South Africa, Uganda	• "ART effect" parameter (ratio of infectiousness with ART to without ART) was used to model the level of viral suppression: base value was 0.25 (till 2015) and reduced to 0.1 (by 2020) and 0.05 (by 2030) under 90-90-90 scenario • Adult ART 2020 coverage: 81% • Adult ART 2030 coverage: 90% • Eligibility for treatment expands to all PLHIV by 2017	**Effectiveness: 73%** **Calculation:** 1-(0.81x0.1+0.19) = 73% (in 2020)
Korenromp (GOALS model), PLoS ONE 2015	Mozambique	ART reduces infectivity of HIV patients by **80%**, as an average effectiveness between	**Effectiveness: 34%**

STUDY	SETTING	PARAMETERS	ART EFFECTIVENESS
Impact and Cost of the HIV/AIDS National Strategic Plan for Mozambique, 2015-2019—Projections with the Spectrum/Goals Model		relevant recent studies including a 96% reduced infectivity found in a clinical trial across multiple–mainly developed, western–countries with very high adherence [Cohen NEJM 2011; Attia AIDS 2009], a 38% reduction in a high-coverage ART program in rural South Africa [Tanser Science 2013], and 85% viral suppression observed in Swaziland [SHIMS 2010-12; Justman CROI 2013] • Scenario 'current targets': ART is scaled-up from 56% to 76% of adults with CD4 <350 in North region, from 65% to 81% in Center, and from 57% to 85% in South; additionally eligibility includes TB/HIV-co-infected adults and pregnant women (from 2012 and 2014, respectively), in all scenarios irrespective of CD4 count. • Scenario 'Accelerated scale-up': ART is further scaled-up to 85% of eligible PLWH with CD4 <350 and all FSW irrespective of CD4 count • Retention on ART, at 3 years after enrolment: 52% in current targets scenario and 70% in Accelerated scale-up scenario	**Calculation:** 1-(0.85*0.5*0.2+(1-0.85*0.5)) = 34%
Walensky (CEPAC model), Ann Intern Med 2016	South Africa	• UNAIDS Target strategy: 73% suppression in 5 years from 24%	**Effectiveness: 20%**

STUDY	SETTING	PARAMETERS	ART EFFECTIVENESS
The Anticipated Clinical and Economic Impact of 90-90-90 in South Africa		• HIV transmission rates by disease stage and viral load: 0.16–9.03/100PY (Attia AIDS 2009; Wawer JID 2005) • Mean ART efficacy, % virologic suppression at 48 weeks: 72% (Kranzer JAIDS 2013; Barth LID 2010)	**Calculation:** 1-(0.73x0.72+0.27) = 20%
Olney (Imperial model), Lancet HIV 2016 Evaluating strategies to improve HIV care outcomes in Kenya: a modelling study	Kenya	• Table S5: Infectiousness weights by health state HIV-positive, on ART and virally suppressed: 0.1 (estimate) • Proportion of individuals initiating ART who adhere to ART and achieve viral suppression: 86% (Boender CID 2015)	**Effectiveness: 51%** **Calculation:** 1-(0.6x(0.86x0.1+0.14x0.5)+0.4)=0.51 Assuming a coverage of 60% and for those not virally suppressed transmission fall by 0.5.
Hontelez (STDSIM model), AIDS 2016 Changing HIV treatment eligibility under health system constraints in sub-Saharan Africa: investment needs, population health gains, and cost-effectiveness	10 sub-Saharan African countries (80% regional burden)	• ART reduces infectiousness of HIV by 90% (Attia AIDS 2009)	54% **Calculation:** 1-(0.6x0.1+0.4)=0.54

Figure 1. Annual HIV incidence (blue) and mortality (red) for Mozambique.

Dots: GOALS model[1]; Lines: SACEMA model.[2]

Left: GOALS projections compared to SACEMA projections.

Right: Adjustment of SACEMA model to assume 65% of those infected with HIV are on ART and ART reduces transmission by 65% the SACEMA model gives results that are very close to the GOALS model.

1. Korenromp, E., et al., Impact and Cost of the HIV/AIDS National Strategic Plan for Mozambique, 2015-2019—Projections with the Spectrum/Goals Model. PLOS One, 2016. 10 (11): e0142908. doi:10.1371.
2 Williams, B.G., et al., Epidemiological Trends for HIV in Southern Africa: Implications for Reaching the Elimination Targets. Current HIV/AIDS reports, 2015. 12: p. 1-11.

References

1.   UNAIDS Global Update (Geneva 2016)
http://www.unaids.org/en/resources/documents/2016/Global-AIDS-update-2016.
2.   WHO TB Fact Sheet (WHO 2016)
http://www.who.int/tb/challenges/mdr/mdr_tb_factsheet.pdf.
3.   Das M, Chu PL, Santos GM, et al. Decreases in community viral load are accompanied by reductions in new HIV infections in San Francisco. *PloS one* 2010; **5**(6): e11068.
4.   Group ISS, Lundgren JD, Babiker AG, et al. Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection. *The New England journal of medicine* 2015; **373**(9): 795-807.
5.   Montaner JS, Hogg R, Wood E, et al. The case for expanding access to highly active antiretroviral therapy to curb the growth of the HIV epidemic. *Lancet* 2006; **368**(9534): 531-6.
6.   Group TAS, Danel C, Moh R, et al. A Trial of Early Antiretrovirals and Isoniazid Preventive Therapy in Africa. *The New England journal of medicine* 2015; **373**(9): 808-22.
7.   Attia S, Egger M, Muller M, Zwahlen M, Low N. Sexual transmission of HIV according to viral load and antiretroviral therapy: systematic review and meta-analysis. *AIDS* 2009; **23**(11): 1397-404.
8.   Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *The New England journal of medicine* 2011; **365**(6): 493-505.
9.   Tanser F, Barnighausen T, Grapsa E, Zaidi J, Newell ML. High coverage of ART associated with decline in risk of HIV acquisition in rural KwaZulu-Natal, South Africa. *Science* 2013; **339**(6122): 966-71.
10.   Rodger AJ, Cambiano V, Bruun T, et al. Sexual Activity Without Condoms and Risk of HIV Transmission in Serodifferent Couples When the HIV-Positive Partner Is Using Suppressive Antiretroviral Therapy. *Jama* 2016; **316**(2): 171-81.
11.   Antiretroviral treatment as prevention (TASP) of HIV and TB: Programmatic update (2012). WHO Geneva.
http://apps.who.int/iris/bitstream/10665/70904/1/WHO_HIV_2012.12_eng.pdf.
12.   UNAIDS 2014 An ambitious treatment target to help end the AIDS epidemic
http://www.unaids.org/sites/default/files/media_asset/90-90-90_en_0.pdf.
13.   Gupta S GR. When will sub-Saharan Africa adopt HIV treatment for all? S Afr J HIV Med. 2016;17(1), a459. http://dx.doi.org/10.4102/sajhivmed.v17i1.459.
14.   Guidelines on When to Start Antiretroviral Therapy and on Pre-exposure Prophylaxis for HIV. September WHO 2015
http://apps.who.int/iris/bitstream/10665/186275/1/9789241509565_eng.pdf.
15.   Garnett GP, Anderson RM. Antiviral therapy and the transmission dynamics of HIV-1. *The Journal of antimicrobial chemotherapy* 1996; **37 Suppl B**: 135-50.
16.   Velasco-Hernandez JX, Gershengorn HB, Blower SM. Could widespread use of combination antiretroviral therapy eradicate HIV epidemics? *The Lancet Infectious diseases* 2002; **2**(8): 487-93.
17.   Auvert B, Males S, Puren A, Taljaard D, Carael M, Williams B. Can highly active antiretroviral therapy reduce the spread of HIV?: A study in a township of South Africa. *Journal of acquired immune deficiency syndromes* 2004; **36**(1): 613-21.

18.   Granich RM, Gilks CF, Dye C, De Cock KM, Williams BG. Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: a mathematical model. *Lancet* 2009; **373**(9657): 48-57.

19.   Schwartlander B, Stover J, Hallett T, et al. Towards an improved investment approach for an effective response to HIV/AIDS. *Lancet* 2011; **377**(9782): 2031-41.

20.   Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection: recommendations for a public health approach (2013). WHO, Geneva. http://apps.who.int/iris/bitstream/10665/85321/1/9789241505727_eng.pdf.

21.   Hamilton A, Garcia-Calleja JM, Vitoria M, et al. Changes in antiretroviral therapy guidelines: implications for public health policy and public purses. *Sexually transmitted infections* 2010; **86**(5): 388-90.

22.   Granich R, Kahn JG, Bennett R, et al. Expanding ART for treatment and prevention of HIV in South Africa: estimated cost and cost-effectiveness 2011-2050. *PloS one* 2012; **7**(2): e30216.

23.   Stover J, Bollinger L, Izazola JA, et al. What Is Required to End the AIDS Epidemic as a Public Health Threat by 2030? The Cost and Impact of the Fast-Track Approach. *PloS one* 2016; **11**(5): e0154893.

24.   Walensky RP, Wolf LL, Wood R, et al. When to start antiretroviral therapy in resource-limited settings. *Annals of internal medicine* 2009; **151**(3): 157-66.

25.   Barnighausen T, Bloom DE, Humair S. Economics of antiretroviral treatment vs. circumcision for HIV prevention. *Proceedings of the National Academy of Sciences of the United States of America* 2012; **109**(52): 21271-6.

26.   Centers for Disease Control, USA New PHIA Survey Data Show Critical Progress Towards Global HIV Targets https://www.cdc.gov/globalhivtb/who-we-are/events/world-aids-day/phia-surveys.html Accessed 21 May 2017.

27.   Granich R, Gupta S, Montaner J, Williams B, Zuniga JM. Pattern, Determinants, and Impact of HIV Spending on Care and Treatment in 38 High-Burden Low- and Middle-Income Countries. *Journal of the International Association of Providers of AIDS Care* 2016; **15**(2): 91-100.