Reconciling age-related changes in behavioural and neural indices of human perceptual decision making

David P. McGovern[1][*][§], Aoife Hayes[1][§], Simon P. Kelly[2] & Redmond O'Connell[1]

[1] Trinity College Institute of Neuroscience and School of Psychology, Trinity College Dublin, College Green, Dublin 2, Ireland
[2] School of Electrical & Electronic Engineering, University College Dublin, Dublin 4, Ireland

[*]Email: mcgoved1@tcd.ie

[§] These authors contributed equally to this work

Ageing impacts on decision making behaviour across a wide range of cognitive tasks and scenarios. Computational modeling has proven highly valuable in providing mechanistic interpretations of these age-related differences; however, the extent to which model parameter differences accurately reflect changes to the underlying neural computations has yet to be tested. Here, we measured neural signatures of decision formation as younger and older participants performed motion discrimination and contrast-change detection tasks, and compared the dynamics of these signals to key parameter estimates from fits of a prominent accumulation-to-bound model (drift diffusion) to behavioural data. Our results indicate marked discrepancies between the age-related effects observed in the model output and the neural data. Most notably, while the model predicted a higher decision boundary in older age for both tasks, the neural data indicated no such differences. To reconcile the model and neural findings, we used our neurophysiological observations as a guide to constrain and adapt the model parameters. In addition to providing better fits to behaviour on both tasks, the resultant neurally-informed models furnished novel predictions regarding other features of the neural data which were empirically validated. These included a slower mean rate of evidence accumulation amongst older adults during motion discrimination and a beneficial reduction in between-trial variability in accumulation rates on the contrast-change detection task, which was linked to more consistent attentional engagement. Our findings serve to highlight how combining human brain signal measurements with computational modelling can yield unique insights into group differences in neural mechanisms for decision making.

Ageing is associated with a progressive decline in many aspects of cognitive function including episodic memory[1], working memory[2], speed of processing[3] and task-switching[4]. In an effort to better understand these wide-ranging changes, researchers have sought to pinpoint core mental operations that are impacted by ageing (e.g. refs. [3,5,6]). One such operation is perceptual decision making, the process whereby sensory information is translated into goal-directed actions.

Much of what we know about how ageing affects perceptual decision making comes from behavioural studies employing sequential sampling models[7–9]. The core principle underlying these models is that noisy sensory evidence is accumulated until a predetermined quantity has accrued in favour of one of the decision alternatives. The two most consistent findings to emerge from this work have been an age-dependent widening of decision boundaries[10–16], consistent with older adults adopting more cautious decision policies[17] and an age-related slowing of non-decision processing indicating a delay in sensory encoding and/or motor execution[10–12,14,18–21]. Perhaps surprisingly, ageing effects on the parameter that dictates the mean rate of evidence accumulation ('drift rate') vary substantially across studies and appear to be highly task-dependent (e.g. refs. [10,14,21,22]). For instance, whereas older and younger adults had similar drift rates on a signal detection task[10], older adults had lower drift rates on a letter discrimination task[21] and higher drift rates during motion discrimination[16].

While these mathematical models have provided valuable insights into the effects of ageing on decision making, there are limits on what can be gleaned from the modelling of behavioural data alone (see [23] for a recent review). For instance, it is difficult to ascertain whether any age-related differences in drift rate emanate from differences in the quality of sensory encoding, the integrity of the evidence accumulation process itself or engagement of other systems that play a supporting role in perceptual decision making, such as neuromodulatory and attention systems which are known to be affected by ageing (e.g. ref. [24]). More generally, abstract decision models that quantitatively capture behaviour do not necessarily reflect the neural computations underlying decision formation, as evidenced in recent non-human primate research[25–27]. Some animal neurophysiology studies have thus begun to use neural signatures of decision formation directly to construct novel model variants that are more representative of the neural implementation of the decision process as well as its behavioural output[25,28]. The recent identification of analogous decision signals in non-invasive recordings presents an opportunity to similarly develop neurally-informed models of human decision making. These human brain signals bear all of the same decision-predictive characteristics as have been reported for build-to-threshold decision signals observed in single-unit recordings[29–31], including a gradual buildup whose rate predicts reaction time and is proportional to evidence strength, and a fixed amplitude immediately prior to decision reports

consistent with a boundary crossing effect[32–34]. Two functionally-distinct categories of human decision signals have been characterised: effector-selective signals that represent the translation of sensory evidence into a specific motor plan, such as lateralised oscillatory activity in the beta-band[33–36], and a domain-general signal found in the event-related potential, termed the centroparietal positivity (CPP), that exhibits the same evidence accumulation properties irrespective of whether a response is immediate, delayed[37] or not required at all[33]. The buildup of the CPP reliably precedes evidence-selective motor preparation signals[32], suggesting that it reflects a processing level that intermediates between sensory encoding and motor preparation.

To investigate the impact of ageing on the neural decision process and to examine correspondences with the predictions derived from behavioural modelling, we asked a group of younger and older participants to perform a continuous version of a random dot motion task[32] and a gradual contrast-change detection task[33], while we recorded 64-channel electroencephalography (EEG). This approach allowed us to isolate both effector-selective and domain-general indices of decision formation whose dynamics we compared to key parameter values derived from fitting the most popular sequential sampling variant, the drift diffusion model[9], to the behavioural data. For the continuous random dot motion task, the diffusion model accounted for age-related performance deficits (longer reaction times and more misses) in terms of a widening of decision boundaries in older adults. However, there were no differences in the amplitude of either the effector-selective or domain-general decision signals at the time of the decision report. Instead, older adults showed slower decision signal buildup of neural evidence accumulation signals, which is equally consistent with the performance deficits in this task. Meanwhile, in the contrast-change detection task, older subjects performed better (less variable reaction times and fewer misses), and while the model fits explained this by an increase in both drift rate and decision bound, no age-related differences in the corresponding neural signal measurements were observed. We go on to show that if the diffusion model is constrained to take account of these neurophysiological observations, the model provides a better fit to the motion discrimination data and a comparable fit to behaviour on the contrast-change detection task despite having fewer free parameters, as well as producing estimates of the non-constrained parameters that accord with the corresponding neural measures. Specifically, when decision bounds were constrained to be equal and starting-point variability was added as a free parameter to take account of premature evidence accumulation observed in older adults on the motion task, the model was then able to capture the drift rate difference evident in the neural data. In the contrast-change task, reduced across-trial variability in drift rate in older adults that became apparent in the neurally-constrained model was reflected in reduced variability both in the buildup rate of the decision signals and in pretarget alpha-band activity, suggesting more consistent attentional engagement. Together,

our data suggest that human neurophysiological signals can play an important role in constraining models of perceptual decision making and revealing key mechanistic differences that may go undetected using behavioural modelling alone.

Results

Motion Discrimination

Behaviour

In a first experiment, 35 younger (17 males, age range: 18-38, mean age: 22 years old) and 31 older participants (13 males, age range: 66-83, mean age: 74 years old) performed a continuous version of a random dot motion task in which they were required to discriminate the direction of intermittent periods of coherent motion (randomly set to 30 or 60%) that occurred within a continuous stream of incoherent motion (Figure 1a; see also ref. [32]). For both young and old participants, higher sensory evidence strength (motion coherence) resulted in fewer misses (p<0.001 for both age groups, Wilcoxon signed-rank test) and faster reaction times (RTs; p<0.001 for both age groups, Wilcoxon signed-rank test). In keeping with previous studies examining the effects of ageing on motion discrimination[16,38,39], older adults were significantly less accurate than their younger counterparts for both levels of motion coherence (Figure 1b; 30%: U=250, p<0.001, 60%: U=167, p<0.001, Mann-Whitney test) and also displayed significantly slower RTs (Figure 1b; 30%: U=263, p<0.001, 60%: U=114, p<0.001, Mann-Whitney test). Further analysis revealed that older adults made more erroneous responses (30% coherence: 2.6% ± 5.4 vs. 0.5 ± 1.2%, U=357, p=0.005; 60%: 5.4% ± 6.6 vs. 0.8% ± 1.1, U=199, p<0.001) and more misses (30%: 6.6% ± 10.3 vs. 1.3 % ± 2.1, U=344, p=0.007; 60%: 0.4% ± 1.1 vs. 0% of targets, U=473, p=0.04) than their younger counterparts, while there were no differences in the false alarm rates between the two age groups (Younger (mean±SD): 0.39± 0.5 per block, Older: 0.54 ± 0.7 per block, U=497, p=0.55, Mann-Whitney test).

To gain a better understanding of what aspects of the decision process led to these differences in behaviour, we fit a drift diffusion model to the accuracy and RT data for both age groups (Figure 2a). We first identified the key parameters required to fit the model to data pooled across participants from each group (decision bound, drift rate, non-decision time, across-trial drift rate variability; see Supplementary Fig. 1 and *Methods* for further information on our approach), before fitting the model to each individual's data from the two groups (Figure 2c). An analysis of incorrect responses concluded that they were highly unlikely to have arisen from incorrect decision boundary crossings through evidence accumulation (see *Methods* for further details) and therefore incorrect responses were omitted from the fitting procedure. The model produced good fits to the pooled group data for both the younger ($G^2=$ 51.51) and older ($G^2=$ 70.81) groups. Consistent with previous behavioural modelling studies of motion discrimination and other two-alternative perceptual decisions[10,16,21], both the group-level and individual fits suggested that older adults had significantly larger boundary separations than younger adults (Figure 2c; U=91, p<0.001, Mann-

Figure 2: Comparison of diffusion model fits to motion discrimination data with decision boundary free to vary and constrained. a) Schematic of the two-choice drift diffusion model that was fit to the data. Evidence accumulation begins at starting point, *z,* and a response is initiated when the cumulative evidence reaches the correct, *a*, or incorrect, *0*, response boundary. The mean rate of accumulation is determined by the drift rate, *v*, which can vary from trial-to-trial. This variability is assumed to be normally distributed with a standard deviation of *η*. In the neurally-informed model, the starting point of the accumulation process could also vary on a trial-to-trial basis, *sz*, and this variability was assumed to be uniformly distributed. On some trials, the cumulative evidence does not reach either decision boundary before the response deadline and in these cases the trial is classified as a miss. b) Neurally-informed model fits (solid lines) to observed reaction time distributions (histograms) pooled across participants. Inset of each panel: Observed and model estimates of target misses. c) Mean parameter estimates derived from model fits to individual data where decision boundary was free to vary. d) Mean parameter estimates derived from model fits to individual data where decision boundary was constrained to be the same across age groups and where starting point variability was added as a free parameter.

Whitney test; see also Supplementary Fig. 1b), with no significant differences in drift rate (Figure 2c; *vL*: t(64)=0.23,p=0.82; *vH*: t(64)=0.58, p=0.56). Older adults also had significantly shorter non-decision times (t(64)=2.1, p=0.04).

Neurophysiology

To measure the impact of ageing on neural evidence accumulation processes, we recorded EEG while participants performed the random dot motion task. In line with our earlier findings[32,40], the onset of coherent motion elicited a rising positivity in the

event-related potential over centroparietal scalp regions in both younger and older
participants (Figure 3a). Our previous work has demonstrated that this centroparietal
positivity (CPP) exhibits the key properties of bounded accumulation that are central
to sequential sampling models of perceptual decision making [32,33], while other work
has identified similar properties in lateralised oscillatory activity in Mu/Beta band
(e.g. [33–35]). Specifically, these signals exhibit a gradual buildup whose rate scales with
evidence strength and reaches a stereotyped amplitude at the time of response.
Importantly, both of these signals have also been shown to be sensitive to
experimental manipulations previously shown to modulate participants' decision
boundaries, such as predictive cueing paradigms and experiments that manipulate the
speed/accuracy emphasis of a task (e.g. [34,36,41]). Here, we took advantage of these
evidence accumulation properties to examine the effect of ageing on the buildup and
amplitude of response-locked CPP and oscillatory Mu/Beta activity and compare the
findings to our modelling results.
220
Initial inspection of the data revealed that the buildup of the CPP of the older group
actually commenced before the onset of coherent motion suggesting a tendency
towards premature evidence accumulation (see Supplementary Fig. 2). To account for
this, we baseline-corrected the ERP waveforms of both age groups relative to a 100
ms interval preceding this initial build-up (600 to 500 ms prior to coherent motion;
see *Discussion* for further consideration of this analysis step).
227
Across both groups, the CPP shared two of the key characteristics of the theoretical
decision variable central to sequential sampling models, namely a gradual buildup
whose rate increased in proportion to evidence strength (i.e. motion coherence,
$F(1,64)=25.13$, $p<0.001$, $BF_{01}<0.001$), and a stereotyped amplitude immediately prior
to response execution that did not differ as a function of coherence ($F(1, 64)= 0.8078$,
$p=0.37$, $BF_{01}=3.53$). The rate of CPP buildup differed between the age groups, with a
slower rate of accumulation in the older group compared to the young group ($F(1, 64)$
$= 10.73$, $p=0.002$, $BF_{01}=0.005$), while there was no significant difference between age
groups in CPP amplitude at response ($F(1, 64) = 1.72$, $p=0.19$). Although our Bayes
factor analysis indicated that the evidence was inconclusive as to whether there was a
significant difference in the CPP amplitude at response ($BF_{01}=1.42$), it should be
noted that the numerical trend favoured a *lower* CPP amplitude in the older group,
and is thus at odds with the prevailing view to emerge from the modelling literature
that older adults implement higher decision bounds (e.g. [10-16]). A qualitatively similar
pattern of results was observed at the level of effector-selective motor preparation,
with contralateral Mu/Beta activity (Figure 3b) reaching a fixed amplitude before
response execution irrespective of evidence strength ($F(1, 64)= 0.61$, $p=0.44$,
$BF_{01}=3.9$), although it was again inconclusive as to whether the amplitude at response
varied as function of age group with anecdotal evidence in favour of there being no
difference ($F(1,64)=2.76$, $p=0.1$, $BF_{01}=1.84$). Also in keeping with the CPP results,

the buildup rate of contralateral Mu/Beta was faster for higher motion coherence
$(F_{(1,64)}=10.85, p=0.002, BF_{01}=0.007)$ and in younger adults $(F_{(1,64)}=5.9, p=0.018,$
$BF_{01}=0.34)$.
251
252
253

Figure 3: During motion discrimination older adults exhibit a slower rate of evidence accumulation relative to younger adults, with both age groups displaying similar amplitudes at response. a) Stimulus-locked (left) and response-locked (right) CPP waveforms for young and older participants performing the continuous random dot motion task. The rate of CPP buildup differed between the age groups, with a slower rate of accumulation in the older group compared to the young group, while there was no difference in CPP amplitude between the different age groups at the time of decision report (black solid line on right panel). b) A similar pattern of results was observed for contralateral Mu/Beta activity. Shaded areas indicate ±1 SEM of data points.

 Neurally-informed modelling

Our neurophysiological data indicate that age-related decrements in motion
discrimination performance arise due to slower accumulation of motion information
in older adults, with little or no difference in the decision boundary positions of
younger and older participants. However, these observations are at odds with our
behavioural modelling results and those of previous studies (e.g. ref. [16]). In an effort
to address this discrepancy, we fit an additional model to the behavioral data in
which, based on our neurophysiological observations, the decision bound was fixed
to the mean of the young and older group parameter values from the original group-
level model fits (see *Methods*). To account for the early buildup of the CPP observed
in older adults (see Figure 3a), we also included starting point variability as a free
parameter in the model on the grounds that it would capture any variance in
performance and decision-signal buildup due to premature evidence accumulation
(e.g. ref. [42]) irrespective of the particular accumulation strategy that was adopted (e.g.
leaky versus non-leaky). The model fits from this neurally-informed model were
better than the original model for both the younger ($G^2 = 46.9$) and older ($G^2 = 54.3$)
groups despite both models having the same number of free parameters. Moreover,
the age group differences in the parameter values from the neurally-informed model
were more in keeping with our neurophysiological data, with significantly lower drift
rates in older subjects for both low and high coherence stimuli (Figure 2d; *vL*:
U=264, p<0.001, *vH*: U=242, p<0.001, Mann-Whitney tests; see also Supplementary
Fig. 1c). Older adults also displayed significantly higher levels of starting point
variability compared to their younger counterparts (t(64)=4.49, p<0.001), which
accords with the observation of larger pretarget CPP build-up in this group; since the
CPP manifests as a positive deflection irrespective of which decision alternative the
evidence favours, its average amplitude at target onset should be larger when there is
greater starting point variability. Between-trial drift rate variability was also higher in
younger participants than older participants, although this effect was not statistically
significant (t(64)=1.94, p=0.056; see *Contrast-change detection task* for further
analysis and discussion). We also note that even when starting point variability was
not free to vary between groups, a model with constrained bounds produced more
parsimonious fits (BICs=93.29 and 108.37 for younger and older groups,
respectively) than a model with unconstrained bounds (BICs=95.17 and 113.69) and
again highlighted significantly higher drift rates in older participants (*vL*: U=220,
p<0.001, *vH*: U=182, p=0.001, Mann-Whitney tests). Thus, adapting the diffusion
model according to our neurophysiological observations produced better fits to
behaviour while also providing a better account of the age-related differences
observed in our neural measures of evidence accumulation.

Contrast-change Detection

Behaviour

To assess whether the age-related effects we observed on the random dot motion task generalised to other perceptual decision making tasks, the same participants performed a contrast-change detection task[33] within the same testing session, with the order of the tasks counterbalanced across participants. Participants were required to continuously monitor a checkerboard stimulus for intermittent reductions in contrast, which were reported via a right-hand button click (Figure 4a). Older adults surprisingly outperformed their younger counterparts on this task, detecting more of the contrast reductions (Figure 4b; U=456, p=0.005, Mann-Whitney test) despite maintaining similar average response times (Figure 4b; t(74)=0.179, p=0.86) and false alarm rates (Younger: 1.6 per block ± 1.4, Older: 1.6 per block ± 1.9, U=636, p=0.37, Mann-Whitney Test). The younger group did, however, display increased RT variability compared to the older group (Figure 4c; Mean coefficient of variation across participants: Younger= 0.234, Older=0.209, t(74)=2.5, p=0.015). To establish what aspects of the decision process gave rise to the older adults outperforming their younger counterparts on this task, we fit the data with a one-choice drift diffusion model (Figure 5a; ref. [43]), with the same free parameters as in the analysis of the motion discrimination (decision bound, drift rate, non-decision time, between-trial drift variability). A key difference in this model, however, was that we assumed that the drift rate rose linearly over the duration of the target stimulus to reflect the ramping evidence signal (i.e. decreasing contrast; see *Methods*). This model provided a very good fit to the pooled group data (Supplementary Fig. 3a) for both the younger (G^2=84.24) and older (G^2=87.33) groups. The parameter estimates from the fits of this model to the individual data indicated that the behavioural advantage in the older group was linked to the combined effect of two distinct parameter differences: higher drift rates (Figure 5c; t(74)=3.26, p=0.002; see also Supplementary Fig. 3b), as well as wider decision boundary separations than young adults (t(74)=2.4, p=0.02).

Figure 4: Gradual contrast-change detection task and associated behaviour: a) Schematic of contrast-change detection task. Participants monitored a checkerboard annulus for intermittent decreases in stimulus contrast. b) Older adults detected more of the contrast targets than younger adults, while maintaining similar reaction times. c) Reaction time distributions for young and older age groups.

Neurophysiology

To examine whether comparable effects were observed in the neural indices of decision formation, we again measured the CPP and left hemisphere beta (LHB) as assays of sensory evidence accumulation and motor preparation, respectively. In addition to these measures, the gradual contrast detection task also allowed us to track the basic sensory encoding of contrast over time by measuring the amplitude of the occipital steady-state visual evoked potential (SSVEP; refs. [33,44]) generated by the on-off flicker of the checkerboard stimulus. Although the amplitude of the SSVEP was greatly reduced in the older group compared to the young group, when the signals were normalised to take account of these baseline differences (see *Methods* for further details), the SSVEPs for both age groups reliably tracked the decrease in stimulus contrast displaying remarkably similar stimulus- and response-aligned waveforms (Figure 6a). Furthermore, both the CPP (Figure 6b) and LHB (Figure 6c) were very similar across age groups, with no significant differences in the buildup

361 rates (CPP: t(74)=1.18, p=0.24, BF_{01}=3.02; LHB: t(74)=0.99, p=0.32, BF_{01}=3.64) or
362 peak amplitudes (CPP: t(74)=0.68, p=0.5, BF_{01}=4.62; LHB: t(74)=0.16, p=0.88,
363 BF_{01}=5.66) of these signals in the response-aligned waveforms.

370
Figure 5: Comparison of diffusion model fits to contrast-change detection data with decision
boundary and drift rate free to vary and constrained. a) Schematic of the one-choice drift
diffusion model that was fit to the data. Evidence accumulation begins at starting point, *z,* and a
response is initiated when the cumulative evidence reaches the decision boundary, *a*. The rate of
accumulation is called the drift rate, *v*, which can vary from trial-to-trial. This variability is assumed
to be normally distributed with a standard deviation of *η*. On some trials the cumulative evidence
does not reach the decision boundary before the response deadline and in these cases the trial is
classified as a miss. b) Neurally-informed model fits (solid lines) to observed reaction time
distributions (histograms) pooled across participants. Inset of each panel: Observed and model
estimates of target misses. c) Mean parameter estimates derived from model fits to individual data
where decision boundary and drift rate were free to vary. d) Mean parameter estimates derived from
model fits to individual data where decision boundary and drift rate were constrained.

384
385 **Figure 6: Sensory encoding, decision formation and motor preparation signals exhibit similar**

dynamics in young and older participants on the contrast-change detection task. a) Stimulus-locked (left) and response-locked (right) SSVEP displays similar buildup rates and amplitudes at response for both age groups. A similar pattern was observed for both the CPP (b) and for left hemisphere beta (LHB) band activity (c). Dashed lines on stimulus-locked CPP and LHB indicate mean reaction times of both age groups.

Neurally-informed modelling

Our initial neural signals analysis did not uncover any differences at the sensory encoding, decision formation or motor preparation processing levels that could account for the enhanced contrast detection performance of the older participants. As in the random dot motion data, these electrophysiological data appear to be at odds with our findings from a fit of a standard model, which suggested that older adults displayed higher rates of evidence accumulation and elevated decision boundaries. In an attempt to reconcile these findings, we fixed the boundary criterion and drift rate parameters to intermediate values between the younger and older values derived from the initial model fit (4 free parameters) and fit the resulting model (2 free parameters) to the behavioural data (Figure 5b). While the fits of this reduced model to the pooled data were marginally worse compared to the unconstrained model fits for both younger (G^2=97.31) and older (G^2=90.72) groups, the BIC values indicated that the constrained model (Younger: 113.79, Older: 107.2) provided a more parsimonious fit to the data compared to the original model (Younger: 117.19, Older: 120.3) with 4 free parameters. Fitting this neurally-constrained model to the individual data suggested that the key factor underlying the age-related behavioural advantage was a higher degree of between-trial drift rate variability in younger adults compared to older adults (Figure 5d; U=359, p<0.001, Mann-Whitney test; see Supplementary Fig. 3c for parameter values from fits to pooled data). In the following section we sought evidence in our neurophysiological data to support these model findings.

Linking neurally-informed modelling to neurophysiology

The results from the neurally-constrained diffusion model suggest that the main factor driving the older adults' behavioural advantage on the contrast-change detection task is reduced between-trial drift rate variability. To establish whether there was a corresponding age-related reduction in the variability of the neural signatures of decision formation, we calculated single-trial estimates of the CPP slope for the two age groups. This analysis revealed that the buildup rate of the CPP was significantly more variable in the young group compared to the older group (Mean SD across participants: Young=0.31, Old=0.21, t(74)=5.7, p<0.001) in keeping with the modelling results. Furthermore, a similar age-related reduction in the variability of the buildup rate of the CPP was also observed in the data from the random dot

427 motion task (Median SD across participants: Young=0.16, Old=0.13, U=385, p=0.04,
428 Mann-Whitney Test) suggesting that this effect generalises across different
429 perceptual tasks.
430
431
One potential reason for the age-dependent decrease in between-trial variability of
evidence accumulation is that older adults maintained a greater level of engagement
with the task (e.g. [16]), while younger participants may have been more susceptible to
fluctuations of attention over the course of the experiment. Although the current
experiment did not directly manipulate the effect of attention on task performance,
the amplitude of pretarget alpha-band activity over posterior scalp sites is a well-
known electrophysiological signature of endogenous attention[32,45,46]. In line with the
idea that younger adults may be more susceptible to fluctuations of attention, across-
trial variability of alpha band activity (coefficient of variation, CV) was higher for
younger participants compared to their older counterparts (Young: 0.231, Old: 0.185,
t(74)=3.14, p=0.002).
443
444
445 # Discussion
446
Our results highlight ways in which human brain recordings and abstract
computational models can be used to reciprocally inform one another to gain deeper
insights into the factors that shape perceptual decision making. For both motion
discrimination and contrast-change detection decisions, the initial behavioural
modelling and neural signal measurements pointed to very different explanations for
age-related changes in decision making performance. However, adapting the models
to take account of our neurophysiological observations by fixing certain model
parameters (i.e. bound for motion discrimination, bound and drift rate for contrast-
change detection) and adding in others (i.e. starting point variability for motion
discrimination) produced estimates of the remaining unconstrained parameters that
could account for the observed ageing effects on the neural decision signals as well as
behaviour. Furthermore, the neurally-constrained model fits uncovered additional,
novel effects which were in turn confirmed in follow-up neural signal analysis.
460
Like many other studies to examine the effects of ageing using sequential sampling
models[10–16,19,20], our initial model fits to the behavioural data suggested that the main
age-related adjustment in decision making was a widening of the decision
boundaries. However, these modelling results ran contrary to our neural signals
analysis, which showed no significant age-related differences in the amplitude of
either the domain-general or effector-selective signatures of evidence accumulation at
the time of decision report. While a Bayes factor analysis suggested that there was

strong evidence to indicate a lack of age group differences in the amplitude of either
the CPP or Mu/Beta activity at the time of response for the contrast-change detection
task, the results for the motion discrimination task were less conclusive. However, it
should be noted that the numerical group trends went in opposite directions for these
two decision-related signals, with older adults showing smaller CPPs and greater
Mu/Beta activity at response compared to their younger counterparts. Thus, across
the two experimental tasks we observed little neurophysiological evidence to support
the hypothesis that older adults set higher decision boundaries. Importantly, both of
these decision-related signals have previously been shown to be modified in a manner
consistent with a change in decision threshold via experimental manipulations
commonly used to influence decision policies, such as altering the speed/accuracy
emphasis of a task and the presentation of predictive cues[34,36,41], providing confidence
that these signals are sensitive enough to detect group differences in decision
boundary. Thus, these neurophysiological observations suggest that, in the current
experiments, younger and older adults applied similar decision policies. This is not to
say that older adults do not exhibit enhanced response caution in some situations. The
requirement to continuously monitor stimuli for temporally unpredictable feature
changes renders our tasks qualitatively different to those used in previous work in
which trials are discretely presented and the stimulus onset provides a cue for the
initiation of evidence accumulation. Here, the unpredictable evidence onsets and
instructions to respond only when sure likely encouraged the implementation of
conservative decision policies in both groups. It may be that older adults do exhibit
relatively higher decision boundaries when placed under greater time pressure but our
results highlight that any such differences do not generalise to all scenarios.
492
A further advantage of this modelling approach is that it generated novel predictions
regarding the neural data which would otherwise have been overlooked. Our initial
analysis of the sensory encoding and decision signals elicited during the contrast-
change detection task provided no clear explanation as to why the older adults
performed better than younger adults. While the initial model fits to the behavioural
data indicated significant age group differences in both drift rate and boundary
separation, we did not observe any differences in the corresponding neural
measurements (decision signal build-up rates and amplitudes at response,
respectively). When the model was constrained to match these neural observations, it
accounted for the poorer performance of younger adults via increased trial-to-trial
variability in drift rate. Thus, the neurally-constrained model yielded a new,
empirically testable prediction that young participants should exhibit greater trial-to-
trial variability in the build-up rate of the CPP and, indeed, we found this to be the
case during both the contrast and motion discrimination tasks. The increased
variability of drift rates and more variable decision signal build-up of the young
group were also accompanied by increased variability in posterior alpha-band
activity, an established oscillatory marker of attentional engagement[45–47]. These

results suggest that older participants were able to maintain attentional engagement
more consistently across the duration of the tasks. It cannot be determined here
whether this effect reflects a compensatory strategy on the part of the older group or
increased motivation to perform well on the task. Alternatively, given the established
links between heightened arousal and greater variability in evidence accumulation[48],
it may be that younger adults were more aroused as they completed the experiment.
This heightened arousal state could also help to explain the surprising higher miss
rate of the younger group; increased drift variability in the absence of other parameter
differences should result in elevated miss rates. Regardless of the precise mechanisms
underlying the increased drift rate variability of younger adults, these results serve to
highlight how combining neurophysiological data with behavioural modelling make
it possible to disentangle positive and negative ageing effects.
522
Our work builds on an increasing trend to use decision-related signals in the human
brain to constrain computational models of perceptual decision making (e.g. [16,49–53]).
For instance, Turner and colleagues[49] have pioneered a powerful neurally-informed
computational approach in which model parameters are estimated by simultaneously
fitting a model to behavioural data and brain-wide BOLD activations, leading to
better model fits and more accurate predictions about withheld behavioural data. This
data-driven approach has proven to be particularly beneficial in overcoming the
limited temporal resolution of fMRI to isolate decision-related brain structures. Our
approach is distinct from that of Turner et al. to the extent that it focuses on the
dynamics of a specific set of neural signals that have been empirically verified to
trace decision formation in a model-independent fashion[32,33]. Rather than
simultaneously fitting the model to the neural and behavioural data, we first fit a
standard diffusion model to our behavioural data and the recovered parameter
estimates were used to generate predictions about the neural data. We then examined
these decision-related neural signals and, in cases where discrepancies arose between
the model results and the signal analyses, the neural data was used to constrain the
model. In both experiments the constrained model was found to provide a marginally
better fit to the behavioural data than the standard model. More importantly, however,
the constrained model provided a much better account of the key group differences
observed in our neural signatures of decision formation including the slower rate of
evidence accumulation and greater noise accumulation at target onset in older adults
during motion discrimination, and greater across-trial variability in evidence
accumulation rates in younger subjects on the contrast-change detection task.
546
The continuous monitoring tasks employed in the current study placed a strong
emphasis on the detection of stimulus feature changes. In the version of the random
dot motion task implemented here, participants frequently missed targets but
incorrect discriminations were rare. Consequently, the behavioural data provided
fewer constraints for modelling purposes than the discrete trial forced-choice tasks

that are more typically employed in the decision making literature. Nevertheless,
there is good evidence to suggest that the diffusion model behaves appropriately with
these data. First, both the initial and neurally-constrained models provided excellent
fits to the behavioural data from both the motion discrimination and contrast-change
detection tasks (see Figures 2 & 5, as well as Supplementary Figures 1 & 3). Second,
the recovered parameter estimates from the unconstrained models produced age-
related effects that are consistent with those reported in previous studies that
employed standard two-alternative forced-choice tasks. Most notably, in keeping with
the previous literature[10–16], our modelling results indicated an age-dependent
widening of decision boundaries for both the motion discrimination and contrast-
change detection tasks. Thirdly, our neurally-informed models also captured all of the
key group differences in evidence accumulation dynamics observed in our
neurophysiological data. In this sense, the current study highlights how neural data
can provide additional constraints for modelling data from behavioural scenarios that
are in other ways not ideal for model fitting.
567
Whereas older adults displayed smaller drift rate parameter estimates coupled with
slower mean buildup rates of the abstract and effector-selective decision signals on
the random dot motion task, no such effects were observed on the contrast-change
task. Thus, age-related drift rate reductions are likely due to task-specific deficits in
sensory encoding affecting the quality of evidence entering the decision process.
These findings are in keeping with the literature on the effects of age on sensory
processing; while there is ample psychophysical and neurophysiological[54–56] evidence
to suggest that motion processing declines with increasing age, sensitivity to stimuli
of relatively low spatial frequency and high contrast, such as the checkerboard stimuli
used in the current study, appears to be preserved in older adults[57–59]. Thus, taken
together, our neurophysiological data suggest that ageing does not lead to a
fundamental decline in perceptual decision formation processes per se, but rather
impacts on certain sensory inputs. An implication of these observations is that
remediation strategies aimed at enhancing perceptual decisions in older adults should
target those specific aspects of sensory processing that are subject to age-related
decline.
584
One of the unexpected findings to emerge from the neural data was the
early build-up of the CPP in older adults on the random dot motion task. We tested
the hypothesis that this early build-up reflected greater premature (i.e. pre-target)
accumulation in the older group by including starting point variability parameter as a
free parameter in our model on the grounds that it would capture any variance in
performance due to evidence accumulation during the inter-trial interval[42,60]. The
addition of this parameter improved the fit to behaviour and also indicated higher
levels of starting point variability in the older group, consistent with our CPP
observations; because the CPP manifests as a positive deflection irrespective of the

participant's choice [61,62], its amplitude at onset of coherent motion should be larger
with higher levels of starting point variability. The reasons for this early buildup are
unclear. One possibility is that older participants implemented a different evidence
accumulation strategy in comparison to the younger group. Previous work from our
lab has shown that in cases of temporal uncertainty of target onset, younger
participants rely on early target selection signals to initiate neural evidence
accumulation and these findings motivated our initial decision to commence the
accumulation process in our models at the time of evidence onset. However, the
premature buildup of the CPP exhibited by the older adults in the motion
discrimination tasks suggests that it is unlikely that the decision process is started at
evidence onset in this cohort. Rather, the current CPP data suggest that older adults
accumulate evidence over the course of the inter-trial interval, perhaps owing to a
reduced sensitivity to coherent motion onset. In this case, it is possible that both
young and older participants implement a continuous "leaky" accumulation-to-bound
process in which older samples of evidence are discounted as time passes[63]. The
increased pre-target build-up of the CPP in the older adults could arise from reduced
leakage, possibly as a strategic adaptation to avoid missed targets. While the
introduction of starting point variability to our model allowed us to account for
variations in behaviour and CPP arising from pretarget accumulation, further research
will be required to establish the precise accumulation strategies that are implemented
in continuous monitoring contexts.
615
Leveraging human brain signatures of decision formation to constrain computational
signals has potential benefits that extend beyond research on ageing. Recent studies
have suggested that the full drift diffusion model may be more complex than required
in certain circumstances and this complexity can lead to more variable parameter
estimates. The key message from these studies is that the sensitivity of diffusion
models to between-group effects can be enhanced by reducing or constraining some
of its parameters. However, determining which parameters should be constrained in a
given experiment is not straightforward. For example, the EZ diffusion model[64] does
not include between-trial variability in any of its parameters and has been shown to
be a more powerful tool at detecting simulated between-group effects as a result of
this simplification[65]. Yet, our neurophysiological data suggest that its exclusion in the
present context would overlook an important aspect of how ageing impacts the
decision-making process. Here, we show that inspection of neurophysiological
signals reflecting decision formation processes can provide a principled way of
constraining the parameters of abstract decision models (see refs. [25,28,36] for similar
approaches). Given the success of this approach in reconciling the differences in our
behavioural and neural indices of decision making, this study paves the way for using
similar techniques to address questions in clinically-relevant groups with impaired
decision making, such as those with ADHD (e.g. ref. [66]) and those suffering from
addiction (e.g. ref. [67]).

Methods

Participants

Thirty-nine younger and 42 older adults volunteered to take part in the study. The younger participants were recruited from the student population of Trinity College Dublin and compensated with research credits for their time, while older participants were recruited from the Trinity College Institute of Neuroscience volunteer participant panel. Criteria for inclusion in the study were right-handedness; normal or corrected-to-normal vision; no personal history of neurological or psychiatric illness, brain injury, abuse of substances or use of psychotropic drugs; no personal history or family history of epilepsy, unexplained fainting or sensitivity to flickering light; and a minimum score of 26 on the Mini Mental State Examination (MMSE). The age groups were matched for gender and years of education (Younger: 15.1, Older: 15.9), however, the younger group had a significantly higher average score on the MMSE than the older group (Young: 28.8, Older: 27.7, $p<0.001$). All participants were naive to the purposes of the study and provided written consent to participate. Participants were excluded if more than two-thirds of their trials were rejected due to EEG artifacts. This led to the rejection of two young participants and four older participants on the random dot motion task and one young participant and four older participants on the contrast-change detection task. A further two young and seven older participants were excluded from the random dot motion task due to technical issues during data acquisition. Following participant rejection for each experiment, there was a final sample of 35 younger adults (17 males, age range: 18-38, mean age: 22 years old) and 31 older adults (13 males, age range: 66-83, mean age: 74 years old) on the random dot motion task and a sample of 38 younger adults (15 males, age range: 18-38, mean age: 22 years old) and 38 older adults (16 males, age range: 66-85 years old, mean age: 74 years old) on the contrast-change detection task. All recruitment and experimental procedures were approved by the School of Psychology Research Ethics Committee, Trinity College Dublin in accordance with the principles of the Declaration of Helsinki.

Procedure

Participants performed a continuous version of the random dot motion discrimination task[32] and a contrast-change detection task[33]. Both tasks were completed in the same session in a darkened and sound-attenuated room, with the order of the tasks pseudo-randomised across participants. In the same testing session, participants also participated in a visual oddball task, the results of which will be reported in a subsequent publication. Stimuli were presented on a 51 cm CRT monitor operating at 85 Hz and a resolution of 1024x768. Participants were seated at a distance of 55 cm

677   from the display and were instructed to fixate a centrally-presented fixation point at
678   all times during task performance. Prior to each task, participants carried out a
679   practice block to familiarise themselves with the task and stimuli. During practice
680   sessions, participants were given feedback on hits, misses and false alarms.
681
682   *Random dot motion task*
683
Participants continuously monitored a patch of incoherently moving dots for
intermittent targets defined by a period of coherent motion in the leftward or
rightward direction (Figure 1a). Motion direction and coherence level (30% or 60%)
were varied independently and randomly on a target-by-target basis. To facilitate the
measurement of motor preparation signals, participants were asked to indicate
leftward motion with a left-hand button press and rightward motion with a right-hand
button press. Participants were instructed to avoid guessing and to respond as soon as
they were certain they perceived coherent motion. The inter-target interval, during
which the incoherent motion was continuously displayed, lasted 3, 5 or 7 seconds and
was randomly chosen on a target-by-target basis.
694
Motion stimuli were presented within an 8-degree aperture centred on the fixation
point and were displayed against a black background. During incoherent motion, an
average of 150 white dots (each 6 x 6 pixels) were placed randomly and
independently within the circular aperture on each of a sequence of 58.8 ms frames
played at 17 frames per second. During coherent motion, a proportion of the dots
were randomly selected on each frame to be displaced in either a leftward or
rightward direction on the following frame at a speed of 6 degrees per second. In
order to accommodate very slow responses from older participants, coherent motion
targets were displayed for a maximum of 10 seconds or until 500 ms after the
participant responded. Participants completed 6 blocks of the task, each lasting
approximately 5 minutes and comprising 30 targets.
706
707
708   *Gradual contrast-change detection task*
709
Participants continuously monitored a flickering (21.25Hz) annular checkerboard
pattern for intermittent targets defined by a gradual decrease in contrast. The
checkerboard stimulus (inner radius=3 degrees, outer radius=8 degrees) consisted of
alternating light and dark radial segments presented against a dark grey background
and was located in the centre of the display. Targets consisted of a linear decrease in
contrast from 70 to 40% over a period of 1.6 seconds followed by a return to 70%
contrast over a further 0.8 seconds (Figure 4s). The inter-target interval, during which
the stimulus flickered at 70% contrast, lasted 3, 5 or 8 seconds and was randomly
chosen on a target-by-target basis. Participants were instructed to avoid guessing and

to make a mouse button press with their right index finger as soon as they were
certain that the annulus was decreasing in contrast. Participants completed 4 blocks of
the task, each lasting approximately 4 minutes and comprising 24 targets.
722
EEG acquisition and preprocessing

Continuous EEG data were recorded using an ActiveTwo system (Biosemi, The
Netherlands) from 64 scalp electrodes and digitised at 512 Hz. Electrodes were
arranged using the standard 10/20 setup. Vertical and horizontal eye movements were
recorded from four electro-occulogram (EOG) electrodes located above and below
the left eye and at the left and right outer canthus, respectively. Data were analysed
using custom-made scripts in MATLAB (MathWorks) drawing on EEGLAB routines
for reading data files and for spherical interpolation of noisy channels[68]. EEG data
were re-referenced offline to the average reference and low-pass filtered to 40Hz
using a two-way least-squares FIR filter.

EEG data were segmented into epochs of -750 to 2500 ms and -750 to 2100 ms for
the random dot motion and gradual contrast-change detection tasks, respectively. For
the gradual contrast-change detection task, epochs were baseline-corrected relative to
the average signal in the 250 ms interval preceding target onset. Inspection of the data
from the random dot motion task revealed that, on average, the onset of the CPP of
the older group occurred before the onset of coherent motion (see Supplementary Fig.
2), suggesting a tendency towards premature evidence accumulation onset due to
temporal uncertainty of the stimulus onset. Epochs were therefore baseline-corrected
to the average signal from -600 to -500 ms with respect to coherent motion onset.
Trials were rejected if the bipolar vertical EOG signal (upper minus lower) exceeded
an absolute value of 200 μV or if any scalp electrode signal exceeded 100 μV within
a window of 750 ms pre-stimulus to 150 ms post-response. EEG data were converted
to current source density[69] to increase spatial selectivity and to reduce the spatial
blurring effect of volume conduction.

Signal analysis

SSVEP: In the contrast-change detection task the contrast-dependent steady-state
visual evoked potential (SSVEP) provided a cortical representation of the sensory
evidence. The SSVEP was measured at a frequency of 21.25 Hz using the discrete
Fourier transform over a window of exactly 10 cycles (470 ms) of the stimulus flicker
frequency in order to attenuate spectral leakage. We first identified the region of
maximum amplitude of the SSVEP on the grand-average scalp topography over 10
cycles of the SSVEP immediately prior to target onset. For each participant, the
pretarget SSVEP amplitude was averaged across trials and normalised by dividing by
the average amplitude at 21.25 Hz across all electrodes and trials. This normalisation
step was required to account for inter-individual differences in the amplitude of the

SSVEP[70,71]. A further baseline correction step was then applied by dividing the
normalised SSVEP at each electrode by the average amplitude measured across all
frequencies. On the basis of the resulting topographies, the SSVEP was averaged
over 6 electrodes centred on standard sites Oz and POz for both age groups.

The temporal evolution of the SSVEP was measured for each participant by
calculating the average SSVEP amplitude across trials over a 10-cycle window at the
start of the epoch and progressively moving the window forward by one sample (step
size= 50 ms) until the SSVEP was calculated across the entire stimulus- and
response-locked epochs. The SSVEP was then normalised relative to the 470 ms
pretarget window for each participant.

CPP: The timecourse of evidence accumulation was indexed by a centro-parietal
positivity (CPP) in the event-related potential[32,33]. Stimulus-locked CPP waveforms
were generated for each participant by averaging the single-trial epochs defined in
EEG acquisition and preprocessing. Response-locked CPPs were derived by
extracting epochs from -900 to 300 ms relative to the time of the response, retaining
the same prestimulus baseline intervals as the stimulus-locked waveforms. For both
tasks and age groups, CPP amplitude and latency were measured at a single electrode
centred on the region of maximum component amplitude identified in the grand-
average response-locked scalp topography (standard site Pz). The peak magnitude of
the response-locked CPP was calculated as the maximum voltage within the -100 ms
to 100 ms window centred on the individual response time. The buildup rate of the
response-locked CPP was measured as the slope of a straight line fitted to the
unfiltered ERP waveform over a time window of -250 to -100 ms.

Mu/Beta: On the random dot motion task, motor preparation signals were indexed as
a decrease in lateralised oscillatory activity in the Mu and Beta bands (8-30Hz)
excluding the single frequency bin at precisely 17 Hz (stimulus flicker frequency).
Oscillatory power in the Mu/Beta band was measured in the hemisphere contralateral
to the hand used to indicate the direction of dot motion. Mu/Beta power was
measured over electrodes C3 and CP3 for right responses and C4 and CP4 for left
responses with the electrode sites selected on the basis of the grand-average response-
locked scalp topographies. Contralateral signals for right and left responses were
averaged to produce a single contralateral waveform for each motion coherence
condition. The temporal evolution of the Mu/Beta power was measured via a sliding
boxcar window of 412 ms with a 58 ms step size. For each participant, Mu/Beta
amplitude was normalised relative to the average signal from -600 to -500 ms with
respect to coherent motion onset.

On the contrast-change detection task, lateralised Mu/Beta power was measured as
the oscillatory power in the 8-30 Hz range excluding the 21.25 Hz frequency bin
(stimulus flicker frequency) over the standard left hemisphere motor site C3. The

temporal evolution of the Mu/Beta power was measured via a sliding boxcar window
of 470 ms with a 50 ms step size. For each participant, Mu/Beta amplitude was
normalised relative to the 470 ms pretarget window. The slope of the response-locked
Mu/Beta power was measured over a time window of -250 to 50 ms in the response-
locked waveform, while the trough of the Mu/Beta power was measured within the -
100 ms to 100 ms window centred on the individual response time.

Alpha: The degree of attentional engagement during task performance was indexed
by variability in pretarget alpha-band activity. For each participant, the average alpha
amplitude (8-14 Hz) was calculated over the 707 ms (exactly 15 cycles of the
SSVEP) interval prior to target onset across all 17 posterior electrodes. To assess the
variability of alpha power between the two age groups, the coefficient of variation
was calculated by dividing the standard deviation of pretarget alpha amplitude by the
mean activity. The coefficient of variation is closely related to the standard deviation
of a sample; however, it is not dependent on the sample mean[72] and is therefore an
appropriate measure for situations in which there is potentially a difference in sample
means.

Bayes factor analysis: To determine the relative strength of evidence behind our
approach of fixing parameters in the drift diffusion model based on our neural signals
analysis, we conducted a Bayes factor analysis of the slope and amplitude measures
of the CPP and Mu/Beta power for each task using JASP[73]. The Bayes factor
overcomes some of the issues associated with null hypothesis significance testing by
quantifying the relative likelihood of the data under the null versus the alternative
hypothesis. Specifically, we calculated the Jeffrey, Zellner and Siow (JZS; see ref. [74])
Bayes factor with an effect size of 1 to determine the strength of evidence in
favour/against a group-level difference in the slope and peak amplitude of the neural
signals of interest in each task. A JZS Bayes factor can be interpreted such that a
value of three favours the null hypothesis three times more than the alternative
hypothesis, while a value of one third favours the alternative three times more than
the null.
826
Drift diffusion modelling

Behavioural data from the gradual contrast-change detection and random dot motion
tasks were fit with one-choice[43] and two-choice[75] drift diffusion models, respectively.
Drift diffusion models assume that decisions are made through a noisy accumulation
process in which sensory evidence is accumulated over time from a starting point, z,
and a response is initiated when the cumulative evidence reaches the correct, a, or
incorrect, 0, response boundary (see Figure 2a). The rate of accumulation is called the
drift rate, v, and is assumed to reflect the quality of information driving the decision
process. The mean drift rate can vary across trials and this variability is assumed to

be normally distributed with a standard deviation of η. There is also within-trial
variability, or noise, in the evidence accumulation process. This allows that processes
with the same mean drift rate terminate at different times, leading to a distribution of
response times (RTs), and occasionally at the wrong boundary, leading to incorrect
responses. The noise within a trial is also assumed to be normally distributed with a
standard deviation of s and is fixed at 0.1 to scale the other parameters[76]. All non-
decision related processing is accounted for by a single non-decision parameter, t_{er},
that incorporates additive delays associated with sensory encoding and motor
execution. For the contrast-change detection task, we assumed that the drift rate rose
linearly over the duration of the target stimulus to reflect the ramping evidence signal
(i.e. decreasing contrast) in this task. Guided by our neurophysiological observations
from the motion discrimination task, we also introduced across-trial starting point
variability (sz) into the model for that experiment and this was assumed to be
uniformly distributed. Given the continuous nature of the experimental tasks, we
made an additional assumption that if the evidence accumulation process had not
terminated at one of the response boundaries by a time deadline, the trial was
classified as a miss (see also ref. [77]). The response deadlines were set to 10,000ms
and 1750ms for the random dot motion and contrast-change detection tasks,
respectively. We also made the additional assumption that the evidence accumulation
process commenced following target onset. This decision was motivated by previous
work from our lab showing that in scenarios involving temporal uncertainty of target
onset, early target selection signals appear to play a role in initiating the neural
evidence accumulation process[40].

Unlike discrete versions of the random dot motion task, erroneous discriminations on
the continuous version of the task were rare (~1% and 4% of targets in young and
older participants, respectively). Several features of the data suggest that the few
errors that did occur were likely not the result of evidence accumulation towards the
incorrect decision boundary, but rather arose from erroneous action selection. First,
older adults made significantly more errors when coherence was high than when it
was low (5.4% vs. 2.6%, U=260, p<0.001, Mann-Whitney test), a pattern inconsistent
with the predictions of the diffusion model (for example, see ref. [75]). Second, error
rates were greater than false alarms rates for both younger and older adults (Younger:
0.004 vs. 0.002 per second, p=0.01, Older: 0.015 vs. 0.004 per second, p<0.001,
Wilcoxon signed rank test). Given that false alarms occur in conditions with no
sensory evidence (i.e. drift rate=0) and errors occur with a positive drift rate diffusing
towards the correct decision boundary, if errors reflected crossings of the incorrect
decision boundary we would expect to see fewer errors than false alarms. Together
these observations suggest that most of the errors we observe on the motion
discrimination task do not arise from the evidence accumulation process itself. As a
result, errors were excluded from the diffusion modelling analysis of the motion
discrimination data.

We fit a number of diffusion models to the behavioural data from each perceptual
task with varying parameter constraints and estimated parameter values by
minimising the G^2 statistic with a SIMPLEX minimisation routine. In order to fit the
model to the data, five RT quantiles (0.1, 0.3, 0.5, 0.7, 0.9) were calculated from the
RT distribution on correct trials and the proportion of trials lying between those
quantiles were multiplied by the total number of trials to yield observed values. All
RTs lying between 0 ms and the deadline in each task were included in these fits.
These quantiles were then fed into the drift diffusion model to calculate the simulated
proportion of trials that lay between these RT quantiles and were multiplied by the
total number of trials to yield the model-derived expected values. The goodness-of-fit
between the observed values and expected values was calculated via a G^2 test and this
statistic was minimised to provide estimates of the key model parameters. In models
where parameters were constrained to reflect a lack of an age group difference in our
neurophysiological observations, parameters were fixed to the mean of the young and
older estimates from the unconstrained model fit to the pooled data. Model
comparisons were performed using Bayes Information Criterion (BIC). The BIC
provides a trade-off between model complexity and goodness-of-fit, favouring a
model with less parameters if the differences in the degrees of freedom outweigh the
gains associated with a better model fit. The preferred model for each task was
chosen based on which produced the smallest BIC value.

For each experiment, we fit the model to the data in two ways. First, we pooled data
across participants and fit the model separately to the younger and older group data.
Here our approach was to find the most parsimonious version of the model (fewest
number of parameters) to adequately fit the data in order to avoid overfitting (see ref.
[65] for discussion of this topic). To this end, we first attempted to fit the pooled data
with just the core components of the drift diffusion model free to vary (drift rate,
decision boundary and non-decision time). However, this initial model did not
provide a good fit to the data and through model simulations we identified that we
also needed to include between-trial drift rate variability as a free parameter to
capture the shape of the RT distributions. The addition of between-trial drift rate
variability greatly reduced G^2 values in both the younger and older data. Therefore,
we fit this model to each participant's data individually and averaged the resulting
parameter values across participants. These recovered parameter values were then
subject to inferential statistical analyses.

Code availability. Custom code used for conducting EEG analysis and drift diffusion
modelling available upon request to the corresponding author.

Data availability. The data that support the findings of the study are available upon
request to the corresponding author.

Acknowledgements

This work was supported by a European Research Council (ERC) Starting Grant (to ROC) under the European Union's Horizon 2020 research and innovation program (Grant Agreement No. 638289), by a Science Foundation Ireland ERC Support Grant (to ROC) and an Irish Research Council Postgraduate Scholarship (to AH). SPK is supported by a Career Development Award from Science Foundation Ireland (15/CDA/3591).

Author contributions

ROC, SK and AH conceived and designed the experiments. AH collected the data. DPM analysed the data and fit the models. DPM, ROC and SK wrote the manuscript and all authors approved the final version.

References

1. Levine, B., Svoboda, E., Hay, J. F., Winocur, G. & Moscovitch, M. Aging and autobiographical memory: dissociating episodic from semantic retrieval. *Psychol. Aging* **17,** 677–689 (2002).
2. Gazzaley, A., Cooney, J. W., Rissman, J. & D'Esposito, M. Top-down suppression deficit underlies working memory impairment in normal aging. *Nat. Neurosci.* **8,** 1298–1300 (2005).
3. Salthouse, T. A. Constraints on theories of cognitive aging. *Psychon. Bull. Rev.* **3,** 287–299 (1996).
4. Wasylyshyn, C., Verhaeghen, P. & Sliwinski, M. J. Aging and task switching: a meta-analysis. *Psychol. Aging* **26,** 15–20 (2011).
5. Stern, Y. What is cognitive reserve? Theory and research application of the reserve concept. *J. Int. Neuropsychol. Soc.* **8,** 448–460 (2002).
6. Park, D. C. & Reuter-Lorenz, P. The adaptive brain: aging and neurocognitive scaffolding. *Annu. Rev. Psychol.* **60,** 173–196 (2009).
7. Laming, D. *Information theory of choice-reaction times*. (London: Academic Press, 1968).
8. Link, S. W. & Heath, R. A. A sequential theory of psychological discrimination. *Psychometrika* **40,** 77–105 (1975).
9. Ratcliff, R. A Theory of Memory Retrieval. *Psychol. Rev.* **85,** 59–108 (1978).
10. Ratcliff, R., Thapar, A. & McKoon, G. The effects of aging on reaction time in a signal detection task. *Psychol. Aging* **16,** 323–341 (2001).

11. Ratcliff, R., Thapar, A. & McKoon, G. A diffusion model analysis of the
effects of aging on brightness discrimination. *Percept. Psychophys.* **65,** 523–535
(2003).
12. Ratcliff, R., Thapar, A. & McKoon, G. Aging, practice, and perceptual tasks: a
diffusion model analysis. *Psychol. Aging* **21,** 353–371 (2006).
13. Starns, J. J. & Ratcliff, R. The effects of aging on the speed-accuracy
compromise: Boundary optimality in the diffusion model. *Psychol. Aging* **25,** 377–
390 (2010).
14. Ratcliff, R., Thapar, A. & McKoon, G. Individual differences, aging, and IQ in
two-choice tasks. *Cognit. Psychol.* **60,** 127–157 (2010).
15. Spaniol, J., Voss, A. & Grady, C. L. Aging and emotional memory: cognitive
mechanisms underlying the positivity effect. *Psychol. Aging* **23,** 859–872 (2008).
16. Forstmann, B. U. *et al.* The speed-accuracy tradeoff in the elderly brain: a
structural model-based approach. *J. Neurosci.* **31,** 17242–17249 (2011).
17. Rabbitt, P. How Old and Young Subjects Monitor and Control Responses for
Accuracy and Speed. *Br. J. Psychol.* **70,** 305–11 (1979).
18. Ratcliff, R., Thapar, A., Gomez, P. & McKoon, G. A diffusion model analysis
of the effects of aging in the lexical-decision task. *Psychol. Aging* **19,** 278–289
(2004).
19. Ratcliff, R., Thapar, A. & McKoon, G. Aging and individual differences in
rapid two-choice decisions. *Psychon. Bull. Rev.* **13,** 626–635 (2006).
20. Ratcliff, R., Thapar, A. & McKoon, G. Application of the diffusion model to
two-choice tasks for adults 75-90 years old. *Psychol. Aging* **22,** 56–66 (2007).
21. Thapar, A., Ratcliff, R. & McKoon, G. A diffusion model analysis of the
effects of aging on letter discrimination. *Psychol. Aging* **18,** 415–429 (2003).
22. Ratcliff, R., Thapar, A. & McKoon, G. Effects of aging and IQ on item and
associative memory. *J. Exp. Psychol. Gen.* **140,** 464–487 (2011).
23. Dully, J., McGovern, D. P. & O'Connell, R. G. The impact of natural aging on
computational and neural indices of perceptual decision making: A review. *Behav.*
Brain Res. (2018). doi:10.1016/j.bbr.2018.02.001
24. Robertson, I. H. A noradrenergic theory of cognitive reserve: implications for
Alzheimer's disease. *Neurobiol. Aging* **34,** 298–308 (2013).
25. Hanks, T., Kiani, R. & Shadlen, M. N. A neural mechanism of speed-accuracy
tradeoff in macaque area LIP. *eLife* **3,** (2014).
26. Hanks, T. D., Mazurek, M. E., Kiani, R., Hopp, E. & Shadlen, M. N. Elapsed
decision time affects the weighting of prior probability in a perceptual decision
task. *J. Neurosci.* **31,** 6339–6352 (2011).
27. Heitz, R. P. & Schall, J. D. Neural chronometry and coherency across speed-
accuracy demands reveal lack of homomorphism between computational and
neural mechanisms of evidence accumulation. *Philos. Trans. R. Soc. Lond. B. Biol.*
Sci. **368,** 20130071 (2013).

28. Purcell, B. A. & Kiani, R. Neural Mechanisms of Post-error Adjustments of
Decision Policy in Parietal Cortex. *Neuron* **89,** 658–671 (2016).
29. Hanes, D. P. & Schall, J. D. Neural control of voluntary movement initiation.
Science **274,** 427–430 (1996).
30. Roitman, J. D. & Shadlen, M. N. Response of neurons in the lateral
intraparietal area during a combined visual discrimination reaction time task. *J.*
Neurosci. **22,** 9475–9489 (2002).
31. Ratcliff, R., Hasegawa, Y. T., Hasegawa, R. P., Smith, P. L. & Segraves, M. A.
Dual diffusion model for single-cell recording data from the superior colliculus in a
brightness-discrimination task. *J. Neurophysiol.* **97,** 1756–1774 (2007).
32. Kelly, S. P. & O'Connell, R. G. Internal and External Influences on the Rate of
Sensory Evidence Accumulation in the Human Brain. *J. Neurosci.* **33,** 19434–
19441 (2013).
33. O'Connell, R. G., Dockree, P. M. & Kelly, S. P. A supramodal accumulation-
to-bound signal that determines perceptual decisions in humans. *Nat. Neurosci.* **15,**
1729–1735 (2012).
34. de Lange, F. P., Rahnev, D. A., Donner, T. H. & Lau, H. Prestimulus
oscillatory activity over motor cortex reflects perceptual expectations. *J. Neurosci.*
33, 1400–1410 (2013).
35. Donner, T. H., Siegel, M., Fries, P. & Engel, A. K. Buildup of choice-
predictive activity in human motor cortex during perceptual decision making. *Curr.*
Biol. **19,** 1581–1585 (2009).
36. Murphy, P. R., Boonstra, E. & Nieuwenhuis, S. Global gain modulation
generates time-dependent urgency during perceptual choice in humans. *Nat.*
Commun. **7,** 13526 (2016).
37. Twomey, D. M., Kelly, S. P. & O'Connell, R. G. Abstract and Effector-
Selective Decision Signals Exhibit Qualitatively Distinct Dynamics before Delayed
Perceptual Reports. *J. Neurosci.* **36,** 7346–7352 (2016).
38. Ball, K. & Sekuler, R. Improving visual perception in older observers. *J.*
Gerontol. **41,** 176–182 (1986).
39. Billino, J., Bremmer, F. & Gegenfurtner, K. R. Differential aging of motion
processing mechanisms: evidence against general perceptual decline. *Vision Res.*
48, 1254–1261 (2008).
40. Loughnane, G. M. *et al.* Target Selection Signals Influence Perceptual
Decisions by Modulating the Onset and Rate of Evidence Accumulation. *Curr.*
Biol. **26,** 496–502 (2016).
41. Steinemann, N. A., O'Connell, R. G. & Kelly, S. P. Decisions are expedited
through multiple neural adjustments spanning the sensorimotor hierarchy. *Nat.*
Commun. (In press).
42. Jepma, M., Wagenmakers, E.-J. & Nieuwenhuis, S. Temporal expectation and
information processing: a model-based analysis. *Cognition* **122,** 426–441 (2012).

43.   Ratcliff, R. & Van Dongen, H. P. A. Diffusion model for one-choice reaction-
time tasks and the cognitive effects of sleep deprivation. *Proc. Natl. Acad. Sci. U.*
S. A. **108,** 11285–11290 (2011).
44.   Di Russo, F. *et al.* Spatiotemporal analysis of the cortical sources of the steady-
state visual evoked potential. *Hum. Brain Mapp.* **28,** 323–334 (2007).
45.   Hanslmayr, S. *et al.* Prestimulus oscillations predict visual perception
performance between and within subjects. *NeuroImage* **37,** 1465–1473 (2007).
46.   O'Connell, R. G. *et al.* Uncovering the neural signature of lapsing attention:
electrophysiological signals predict errors up to 20 s before they occur. *J. Neurosci.*
29, 8604–8611 (2009).
47.   Dockree, P. M. *et al.* The Effects of Methylphenidate on the Neural Signatures
of Sustained Attention. *Biol. Psychiatry* **82,** 687–694 (2017).
48.   Murphy, P. R., Vandekerckhove, J. & Nieuwenhuis, S. Pupil-Linked Arousal
Determines Variability in Perceptual Decision Making. *PLOS Comput. Biol.* **10,**
e1003854 (2014).
49.   Turner, B. M., van Maanen, L. & Forstmann, B. U. Informing cognitive
abstractions through neuroimaging: The neural drift diffusion model. *Psychol. Rev.*
122, 312–336 (2015).
50.   Turner, B. M., Forstmann, B. U., Love, B. C., Palmeri, T. J. & Van Maanen, L.
Approaches to analysis in model-based cognitive neuroscience. *J. Math. Psychol.*
76, 65–79 (2017).
51.   Turner, B. M. *et al.* A Bayesian framework for simultaneously modeling neural
and behavioral data. *NeuroImage* **72,** 193–206 (2013).
52.   Turner, B. M., Rodriguez, C. A., Norcia, T. M., McClure, S. M. & Steyvers,
M. Why more is better: Simultaneous modeling of EEG, fMRI, and behavioral
data. *NeuroImage* **128,** 96–115 (2016).
53.   Frank, M. J. *et al.* fMRI and EEG Predictors of Dynamic Decision Parameters
during Human Reinforcement Learning. *J. Neurosci.* **35,** 485 (2015).
54.   Yang, Y. *et al.* Aging affects contrast response functions and adaptation of
middle temporal visual area neurons in rhesus monkeys. *Neuroscience* **156,** 748–
757 (2008).
55.   Yang, Y. *et al.* Aging affects the neural representation of speed in Macaque
area MT. *Cereb. Cortex* **19,** 1957–1967 (2009).
56.   Liang, Z. *et al.* Aging affects the direction selectivity of MT cells in rhesus
monkeys. *Neurobiol. Aging* **31,** 863–873 (2010).
57.   Owsley, C., Sekuler, R. & Siemsen, D. Contrast sensitivity throughout
adulthood. *Vision Res.* **23,** 689–699 (1983).
58.   Elliott, D., Whitaker, D. & MacVeigh, D. Neural contribution to
spatiotemporal contrast sensitivity decline in healthy ageing eyes. *Vision Res.* **30,**
541–547 (1990).
59.   Habak, C. & Faubert, J. Larger effect of aging on the perception of higher-
order stimuli. *Vision Res.* **40,** 943–950 (2000).

60. Laming, D. Choice reaction performance following an error. *Acta Psychol. (Amst.)* **43,** 199–224 (1979).

61. Kelly, S. P. & O'Connell, R. G. The neural processes underlying perceptual decision making in humans: Recent progress and future directions. *J. Physiol.-Paris* **109,** 27–37 (2015).

62. Afacan-Seref, K., Steinemann, N. A., Blangero, A. & Kelly, S. P. Dynamic Interplay of Value and Sensory Information in High-Speed Decision Making. *Curr. Biol.* **28,** 795-802.e6 (2018).

63. Usher, M. & McClelland, J. L. The time course of perceptual choice: the leaky, competing accumulator model. *Psychol. Rev.* **108,** 550–592 (2001).

64. Wagenmakers, E.-J., van der Maas, H. L. J. & Grasman, R. P. P. P. An EZ-diffusion model for response time and accuracy. *Psychon. Bull. Rev.* **14,** 3–22 (2007).

65. van Ravenzwaaij, D., Donkin, C. & Vandekerckhove, J. The EZ diffusion model provides a powerful test of simple empirical effects. *Psychon. Bull. Rev.* **24,** 547–556 (2017).

66. Hauser, T. U., Fiore, V. G., Moutoussis, M. & Dolan, R. J. Computational Psychiatry of ADHD: Neural Gain Impairments across Marrian Levels of Analysis. *Trends Neurosci.* **39,** 63–73 (2016).

67. Bechara, A. Decision making, impulse control and loss of willpower to resist drugs: a neurocognitive perspective. *Nat. Neurosci.* **8,** 1458–1463 (2005).

68. Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J. Neurosci. Methods* **134,** 9–21 (2004).

69. Kayser, J. & Tenke, C. E. Principal components analysis of Laplacian waveforms as a generic method for identifying ERP generator patterns: I. Evaluation with auditory oddball tasks. *Clin. Neurophysiol.* **117,** 348–368 (2006).

70. Silberstein, R. B. *et al.* Steady-state visually evoked potential topography associated with a visual vigilance task. *Brain Topogr.* **3,** 337–347 (1990).

71. Silberstein, R. B., Nunez, P. L., Pipingas, A., Harris, P. & Danieli, F. Steady state visually evoked potential (SSVEP) topography in a graded working memory task. *Int. J. Psychophysiol.* **42,** 219–232 (2001).

72. Barlow, J. S. *The Electroencephalogram: Its Patterns and Origins*. (MIT Press, 1993).

73. JASP team. *JASP (Version 0.8.3) [Computer software]*. (2017).

74. Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D. & Iverson, G. Bayesian t tests for accepting and rejecting the null hypothesis. *Psychon. Bull. Rev.* **16,** 225–237 (2009).

75. Ratcliff, R. & McKoon, G. The diffusion decision model: theory and data for two-choice decision tasks. *Neural Comput.* **20,** 873–922 (2008).

76.   Ratcliff, R. & Tuerlinckx, F. Estimating parameters of the diffusion model:
approaches to dealing with contaminant reaction times and parameter variability.
Psychon. Bull. Rev. **9,** 438–481 (2002).
77.   Murphy, P. R., Robertson, I. H., Harty, S. & O'Connell, R. G. Neural evidence
accumulation persists after choice to inform metacognitive judgments. *eLife* **4,**
(2015).
1122