A simple mung bean infection model for studying the virulence of *Pseudomonas aeruginosa*

Sneha Garge[1], Sheyda Azimi[2*] & Stephen P. Diggle[2]

[1]Department of Microbiology and Biotechnology center, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara, India; [2] School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA 30032, USA.

*Correspondence: sheyda.azimi@biosci.gatech.edu

Keywords: Mung bean; *Pseudomonas aeruginosa*; virulence model; quorum sensing

Abstract

Much of our knowledge about bacterial virulence and pathogenesis has been derived from studying cells growing in test tubes, under conditions which are often not representative of those found during infection. In addition, many current *in vivo* models have biotic and abiotic limitations, which include failing to fully mimic *in vivo* environments, financial implications, small sample sizes, and ethical constraints. Therefore, new models which fully or partially address these issues are desirable. Here we highlight the development of a simple and high throughput mung bean model to study virulence in the opportunistic pathogen *Pseudomonas aeruginosa*. The model is easy to setup and cultivate under controlled conditions in the laboratory, and infection and virulence can be monitored for up to 10 days. In a first test of the model, we found that mung bean seedlings infected with PAO1 showed poor development of roots and high mortality rates compared to un-infected controls. We also found that a quorum sensing (QS) mutant strain was significantly less virulent when compared with the PAO1 wild type. Our work introduces a new tool for studying virulence in *P. aeruginosa*, that will allow for high throughput virulence studies of mutants, and for testing the *in vivo* efficacy of new therapies at a time when new antimicrobial drugs are desperately needed.

Main text

Pseudomonas aeruginosa is an opportunistic pathogen that can be isolated from diverse habitats including water, soil, animals and plants (1-3). In humans, it is a leading cause of infection, morbidity and mortality in cystic fibrosis (CF) lungs, and is a problematic pathogen in burn wounds, chronic diabetic wounds and immunocompromised individuals (1, 3, 4). Because of the problems that *P. aeruginosa* causes in human hosts, researchers have developed a number of different *in vivo* infection models for assessing *P. aeruginosa* virulence, pathogenesis and disease(5-7). These include animal models such as waxmoths (8) (9), fruit flies (10) , nematodes, mice (11-13) (14-16), pigs (17) and *ex vivo* pig lungs (18, 19). Because *P. aeruginosa* uses a number of the same virulence factors to infect and cause disease in both animals and plants, plant infection models such as the *Arabidopsis* model have also been widely used (20, 21). A number of factors are important to consider when designing and using a virulence model. These include cost effectiveness, ethical considerations, ease of growth in a laboratory, measures of virulence, and the ability to handle large sample sizes. In this study we describe the development of a mung bean

infection model for *P. aeruginosa*, and describe how this easy to establish model can be used
to assess virulence.

Fig. 1. Mung bean infection method and comparison of growth defect in infected plants. (a) The
procedure to infect mung bean seeds with *P. aeruginosa,* and the comparison of PAO1 and PAO-JG1
treated mung bean plants; (b) root length; (c) shoot length and (d) number of root branches. Values
represent the mean of six trials. Bars indicate standard deviation of the mean. Each trial has five
replicates. Statistical analysis was done using one-way ANOVA Bonferroni's multiple comparison
test (* = $p<0.05$, ** = $p< 0.01$ and *** = $p< 0.0001$).

We purchased mung bean seeds from a general supermarket store (Tesco U.K.). To avoid any
fungal or other bacterial contamination, we first surface sterilized the seeds by soaking in
70% ethanol in a sterile flask for 30 seconds and then we soaked them in 0.01% Mercury
Chloride ($HgCl_2$) for 30 seconds three times. We then washed the seeds with sterile water.
We transferred surface sterilised seeds onto soft agar plates (0.8% w/v agar), and added 2 ml
of sterile water to the surface of the agar. We then incubated the plates for 24 hours at 37℃
under humidified conditions to allow for the germination of seeds. We used 8 germinated
seeds per group for the infection studies. As a first test of our model, we wanted to determine
whether quorum sensing (QS) is important for virulence in mung beans, because QS has
previously been shown to be an important regulator of virulence in a number of different
hosts including mice, *Arabidopsis*, lettuce, nematodes, and insects (22) (11). We used a

PAO1 *lasI⁻ rhlI⁻* double mutant (PAO-JG1) (23) to determine the role of QS on pathogenesis, colonisation and disease development in mung beans.

To infect the seeds, we inoculated the PAO1 wildtype strain and PAO-JG1 into 100 ml Lysogeny Broth (LB) from overnight start-up cultures, and incubated at 30°C/200 rpm to reach a cell density of ~10^9 CFU/ml. We centrifuged stationary phase cultures at 7000 rpm for 10 mins and we re-suspended pellets into 4 ml of sterile PBS (pH 7). We added 8 geminated mung bean seeds to this bacterial suspension and then incubated at 30°C for 24 hours under static conditions to achieve a good colonization of bacterial cells on the germinated seeds. We incubated equal numbers of seeds in 4 ml of PBS as uninfected controls. After the incubation, we took 3 seeds from each treatment group, and re-suspended them in 1 ml of sterile PBS and vortexed for 30 secs. We used this suspension to perform viable counts (CFU/ml), to determine the number of adhered bacteria that coated each seed. The cell counts recorded are shown in Table 1.

	Initial CFU/ml	CFU/ml adhered to seedlings	CFU/ml of roots 10 days post inoculation	CFU/mg of roots 10 days post inoculation
PAO1	$1.16\pm0.65\times10^9$	$1.92\pm1.64\times10^8$	$5.4\pm5.0\times10^6$	$1.6\pm0.6\times10^6$
PAO-JG1	$1.36\pm0.68\times10^9$	$1.80\pm1.65\times10^6$	$5.1\pm6.4\times10^6$	$3.5\pm1.4\times10^5$

Table 1. Viable counts of colonized bacteria in seedlings and roots by PAO1 or PAO-JG1. Values are represented as mean± SD for six independent trials with five technical replicates.

For plant growth, we prepared Murashige-skoog agar media [Murashige-skoog 4.4 g/L, (Sigma aldrich) and 0.8% w/v agar (Oxoid)]. We poured 30 ml aliquots of media into large glass tubes (30×200 mm, Duran groups) and sterilized by autoclaving for 20 mins. We aseptically transferred both infected and PBS control mung bean seedlings (5 seeds per group) into the tubes. We allowed the seedlings to grow in a natural daylight cycle for 10 days in gnotobiotic conditions. To determine viable bacterial counts (CFU/mg of root weight) of both the strains colonizing the roots of the mung bean seeds during the incubation of 10 days, we cut the root of each plant with sterile scissors, and re-suspended these in 1 ml of sterile PBS and vortexed for 30 secs. We used this suspension to perform a viable count of bacteria to determine the number of bacteria colonizing the root of each plant. We normalized the CFUs on each root by the respective root weight (mg) of each plant root. We show that both PAO1 and PAO-JG1 strains were viable on the roots throughout the infection period (Table 1). We determined the health of the plants by measuring parameters such as mortality rate of seedlings, shoot length, root length and the number of root branches due to the infection. We defined mortality rate as the percentage of seedlings that died from the total number of seedlings treated with bacterial culture suspension in each trial. We found that the mortality rate of the seedlings treated with PAO1 was higher compared to PAO-JG1 treated seedlings (Table 2). We observed all the seedlings infected with PAO-JG1 developed into healthy plants (Fig. 1, a). We found that seedlings which survived infection with PAO1, demonstrated severely attenuated growth (Fig. 1, a). These plants displayed disease symptoms such as water soaked lesions on their roots, and blackening of the root and shoot.

Our infection model examined three parameters to determine and score plant health: root length, shoot length and the number of root branches. In seedlings that survived PAO1 treatment, we observed a significant decrease in the root and shoot length measures when

compared to seedlings treated with PAO-JG1 and healthy PBS control plants (Fig. 1, b, c and d). We found that the root development of PAO1 treated plants was also impaired as the number of root branches were reduced when compared to PAO-JG1 treated plants and healthy PBS control plants (Fig. 1, d).

	Mortality Rate (%)					
	Trial 1	Trial 2	Trial 3	Trial 4	Trial 5	Trial 6
PAO1	60%	20%	40%	40%	100%	No mortality
PAO-JG1	No mortality	No mortality	No mortality	No mortality	No mortality	No mortality

Table 2. Mortality rates in mung bean treated seedlings by PAO1 and PAO-JG1. There were five culture suspension treated seedlings in each trial.

In 6 independent trials, each with 5 technical replicates, we found that PAO1 either caused the death of the seedlings or the seedlings survived with impaired growth. In contrast, we found that PAO-JG1 did not cause mortality of seedlings in any of the trials and furthermore, the seedlings developed into significantly healthy plants after 10 days (Fig. 1 and Table 2). This data confirms previous observations that the *las* and *rhl* QS systems regulate the virulence of *P. aeruginosa* in plant tissues (24-26). Although we found that QS is important for virulence and disease in mung beans, infections were not completely prevented, suggesting the role of other factors responsible for pathogenesis in mung bean plants which are QS-independent.

Our data demonstrates that we have developed a simple plant model to study mechanisms of pathogenesis and virulence during *P. aeruginosa* infection. Our model has advantages compared to the popular *Arabidopsis* model, which takes 4-6 weeks to grow the plants, and then 5 days post inoculation for observing disease symptoms (20, 24). The mung bean model takes a maximum of 12 days, which includes 1 day to obtain germinated mung bean seedlings, 1 day for infecting the germinated seeds of mung bean with bacterial cells and 7-10 days to grow the plants and allow for the development of disease symptoms. The ease of the methods involved in this model also offers a number of conveniences. It is easy and cheap to acquire mung bean seeds, germinate them and maintain the plants in most lab facilities and without any ethical considerations. Coating germinated seedlings with culture suspension to achieve infection is a simple microbiological technique, and coating allows researchers to evaluate infection in terms of colonization and mortality of seedlings. This can be used as an important parameter for qualitative studies of the colonization of a host, virulence and phenotypic traits involved in the pathogenesis of *P. aeruginosa*. You can also score the pathogenesis of *P. aeruginosa* in plants by measuring parameters such as root length, shoot length and the number of root branches. These are useful parameters where levels of infection by different mutants or strain can reveal important information about virulence factors and virulence in general.

Large sample sizes is not a limitation for the model, so it can be used as a high-throughput virulence screen. Due to the clear effects that we found of QS on virulence; screening libraries of chemical compounds such as *N*-acyl homoserine lactone (AHL) mimics, analogues, inhibitors, antagonists, or quorum quenching bacteria for controlling the infection, could be easily tested. Functional studies of the genes and regulatory proteins which are

involved in phenotypes such as motility, adhesion, colonisation, pathogen survival, and their
impact on virulence can be conducted using the model. Finally, if seedlings are coated with
sub-lethal doses of *P. aeruginosa,* which do not cause the death of plants, this could allow for
sociomicrobiology studies relating to social behaviours, to determine how spatial and
temporal interactions within and between bacterial communities progress in a natural
environment. In summary, we have developed a cheap, reproducible and simple to cultivate
plant host model which can be used to study *P. aeruginosa* virulence, and it also may also
have relevance for other bacterial species.

Acknowledgments

158 S.G. was sponsored by an EMBO Short Term Fellowship (ASTF number: 27-2015). This
159 work was supported by a Human Frontier Science Program Young Investigators grant to
160 S.P.D. (RGY0081/2012). We thank Freya Harrison for helpful comments on the work.

References

1.   Lin CK, Kazmierczak BI. Inflammation: A Double-Edged Sword in the Response to
Pseudomonas aeruginosa Infection. J Innate Immun. 2017;9(3):250-61.
2.   Barr HL, Halliday N, Camara M, Barrett DA, Williams P, Forrester DL, et al. Pseudomonas
aeruginosa quorum sensing molecules correlate with clinical status in cystic fibrosis. Eur Respir J.
2015;46(4):1046-54.
3.   Gellatly SL, Hancock RE. Pseudomonas aeruginosa: new insights into pathogenesis and host
defenses. Pathog Dis. 2013;67(3):159-73.
4.   Whiteley M, Diggle SP, Greenberg EP. Progress in and promise of bacterial quorum sensing
research. Nature. 2017;551(7680):313-20.
5.   Folkesson A, Jelsbak L, Yang L, Johansen HK, Ciofu O, Hoiby N, et al. Adaptation of
Pseudomonas aeruginosa to the cystic fibrosis airway: an evolutionary perspective. Nat Rev
Microbiol. 2012;10(12):841-51.
6.   Williams BJ, Dehnbostel J, Blackwell TS. Pseudomonas aeruginosa: host defence in lung
diseases. Respirology. 2010;15(7):1037-56.
7.   Roberts AE, Kragh KN, Bjarnsholt T, Diggle SP. The Limitations of In Vitro Experimentation in
Understanding Biofilms and Chronic Infection. J Mol Biol. 2015;427(23):3646-61.
8.   Harrison F, Browning LE, Vos M, Buckling A. Cooperation and virulence in acute
Pseudomonas aeruginosa infections. BMC Biol. 2006;4:21.
9.   Racey D, Inglis RF, Harrison F, Oliver A, Buckling A. The effect of elevated mutation rates on
the evolution of cooperation and virulence of Pseudomonas aeruginosa. Evolution. 2010;64(2):515-
21.
10.   Sibley CD, Duan K, Fischer C, Parkins MD, Storey DG, Rabin HR, et al. Discerning the
complexity of community interactions using a Drosophila model of polymicrobial infections. PLoS
Pathog. 2008;4(10):e1000184.
11.   Mahajan-Miklos S, Tan MW, Rahme LG, Ausubel FM. Molecular mechanisms of bacterial
virulence elucidated using a Pseudomonas aeruginosa-Caenorhabditis elegans pathogenesis model.
Cell. 1999;96(1):47-56.
12.   Papaioannou E, Wahjudi M, Nadal-Jimenez P, Koch G, Setroikromo R, Quax WJ. Quorum-
quenching acylase reduces the virulence of Pseudomonas aeruginosa in a Caenorhabditis elegans
infection model. Antimicrob Agents Chemother. 2009;53(11):4891-7.
13.   Brackman G, Cos P, Maes L, Nelis HJ, Coenye T. Quorum sensing inhibitors increase the
susceptibility of bacterial biofilms to antibiotics in vitro and in vivo. Antimicrob Agents Chemother.
2011;55(6):2655-61.

14.   Stieritz DD, Holder IA. Experimental studies of the pathogenesis of infections due to
Pseudomonas aeruginosa: description of a burned mouse model. J Infect Dis. 1975;131(6):688-91.

15.   Rumbaugh KP, Diggle SP, Watters CM, Ross-Gillespie A, Griffin AS, West SA. Quorum sensing
and the social evolution of bacterial virulence. Curr Biol. 2009;19(4):341-5.

16.   Moser C, Van Gennip M, Bjarnsholt T, Jensen PO, Lee B, Hougen HP, et al. Novel
experimental Pseudomonas aeruginosa lung infection model mimicking long-term host-pathogen
interactions in cystic fibrosis. APMIS. 2009;117(2):95-107.

17.   Stoltz DA, Meyerholz DK, Pezzulo AA, Ramachandran S, Rogan MP, Davis GJ, et al. Cystic
fibrosis pigs develop lung disease and exhibit defective bacterial eradication at birth. Sci Transl Med.
2010;2(29):29ra31.

18.   Harrison F, Muruli A, Higgins S, Diggle SP. Development of an ex vivo porcine lung model for
studying growth, virulence, and signaling of Pseudomonas aeruginosa. Infect Immun.
2014;82(8):3312-23.

19.   Harrison F, Diggle SP. An ex vivo lung model to study bronchioles infected with
Pseudomonas aeruginosa biofilms. Microbiology. 2016;162(10):1755-60.

20.   Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. Common virulence
factors for bacterial pathogenicity in plants and animals. Science. 1995;268(5219):1899-902.

21.   He P, Chintamanani S, Chen Z, Zhu L, Kunkel BN, Alfano JR, et al. Activation of a COI1-
dependent pathway in Arabidopsis by Pseudomonas syringae type III effectors and coronatine. Plant
J. 2004;37(4):589-602.

22.   Rahme LG, Ausubel FM, Cao H, Drenkard E, Goumnerov BC, Lau GW, et al. Plants and
animals share functionally common bacterial virulence factors. Proc Natl Acad Sci U S A.
2000;97(16):8815-21.

23.   Cornforth DM, Popat R, McNally L, Gurney J, Scott-Phillips TC, Ivens A, et al. Combinatorial
quorum sensing allows bacteria to resolve their social and physical environment. Proc Natl Acad Sci
U S A. 2014;111(11):4280-4.

24.   Walker TS, Bais HP, Deziel E, Schweizer HP, Rahme LG, Fall R, et al. Pseudomonas
aeruginosa-plant root interactions. Pathogenicity, biofilm formation, and root exudation. Plant
Physiol. 2004;134(1):320-31.

25.   Quinones B, Dulla G, Lindow SE. Quorum sensing regulates exopolysaccharide production,
motility, and virulence in Pseudomonas syringae. Mol Plant Microbe Interact. 2005;18(7):682-93.

26.   Ortiz-Castro R, Martinez-Trujillo M, Lopez-Bucio J. N-acyl-L-homoserine lactones: a class of
bacterial quorum-sensing signals alter post-embryonic root development in Arabidopsis thaliana.
Plant Cell Environ. 2008;31(10):1497-509.

229