Running Title: *Drosophila* descending neuron function

Title

Optogenetic dissection of descending behavioral control in *Drosophila*.

Authors

Jessica Cande[1*], Gordon J. Berman[2,3*], Shigehiro Namiki[1,4], Jirui Qiu[3], Wyatt Korff[1],

Gwyneth Card[1], Joshua W. Shaevitz[5] and David L. Stern[1].

Author Affilitations

[1]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147.

[2]Department of Biology, Emory University, Atlanta, Georgia 30322.

[3]Department of Physics, Emory University, Atlanta, Georgia 30322.

[4]Research Center for Advanced Science and Technology, University of Tokyo, Tokyo

153-8904, Japan.

[5]Department of Physics and the Lewis-Sigler Institute for Integrative Genomics,

Princeton University, Princeton, New Jersey 08544.

[*]These authors contributed equally to the manuscript.

Correspondence (for submission purposes): David L. Stern

Email: sternd@janelia.hhmi.org

Telephone: 571-209-4237

19700 Helix Dr., Ashburn, VA 20147

Abstract

In most animals, the brain makes behavioral decisions that are transmitted by descending

neurons to the nerve cord circuitry that produces behaviors.   In insects, only a few

descending neurons have been associated with specific behaviors. To explore how these

neurons control an insect's movements, we developed a novel method to systematically

assay the behavioral effects of activating individual neurons on freely behaving terrestrial

D. melanogaster. We calculated a two-dimensional representation of the entire behavior

space explored by these flies and associated descending neurons with specific behaviors

by identifying regions of this space that were visited with increased frequency during

optogenetic activation. Applying this approach across a population of descending

neurons, we found, that (1) activation of most of the descending neurons drove

stereotyped behaviors, (2) in many cases multiple descending neurons activated similar

behaviors, and (3) optogenetically-activated behaviors were often dependent on the

behavioral state prior to activation.

Introduction

As animals navigate a dynamic environment, their survival depends on their ability to execute specific motor programs and to adjust motor output in response to external stimuli. While the brain performs computations essential for behavior, the motor circuits that directly control behavior are located close to the muscles that they control in the vertebrate spinal cord and insect ventral nerve cord. Information to drive motor patterns must therefore be transmitted from the brain to the nerve cord to direct behavior. Since there are many fewer descending neurons than neurons in the central brain, descending neurons generate an information processing bottleneck, which may generate a fundamental problem in information coding.

In flies, descending commands from the brain to the ventral nerve cord are transmitted through an estimated 250-550 pairs of descending neurons that arborize in 20 highly-conserved clusters in the brain involved in sensory processing and motor behavior (Gronenberg & Strausfeld, 1990; Hsu & Bhandawat, 2016). Each descending neuron extends a single axon through the neck connective to the ventral nerve cord, where they synapse onto interneurons associated with leg, neck, and wing motor circuitry (Namiki et al, 2017).

Little is known about how so few neurons—approximately 0.5% of all neurons in the fly (Alivisatos et al., 2012)—encode signals from the brain to control the full range of movements performed by a freely moving fly. Several potential models have been suggested. One possibility is that, as with vertebrates, many stereotypical insect behaviors, such as walking, flying, or "singing" can be decomposed into individual motor modules controlled by central pattern generators located in the ventral nerve cord.

Several recent findings in *Drosophila melanogaster*, together with earlier

electrophysiological studies in larger insects, support this idea. Activation of some

individually-identifiable descending neurons triggers specific motor outputs, such as

courtship song (von Philipsborn et al., 2011a), backwards walking (Bidaye, Machacek,

Wu, & Dickson, 2014), or escape behavior (King & Wyman, 1980). However, some

descending neurons modify motor programs, rather than trigger them. For example,

cricket walking initiation, speed, and turning appear to depend on separately encoded

descending commands (Böhm & Schildberger, 1992; Gras & Kohstall, 1998).

Alternatively, motor activity may result from the summed activity of multiple descending

neurons (Heinrich, 2002). For example, a cluster of descending neurons linking fly visual

centers in the brain to the flight apparatus in the ventral nerve cord (Strausfeld &

Gronenberg, 1990; Namiki et al, 2017) supports the idea that at least some descending

neurons may function this way.

Descending neuron function may also be gated by behavioral state. For example,

a single descending neuron may cause a different or modified behavior if the animal is

walking versus flying. For example, DN sensory responses have been shown to be

modified by locomotor state (Staudacher & Schildberger, 1998). This kind of gating has

been observed in other contexts, such as the effect of the neuromodulator pyrokinin on

the oscillatory mechanisms underlying the crustaecean gastric mill central pattern

generator (Marder & Goaillard, 2006). However, it has not previously been possible to

undertake a systematic analysis of the context dependency across the DN population.

Systematic dissection of descending motor control is challenging for two reasons.

First, it has been difficult to precisely manipulate a large number of descending neurons

individually in freely behaving animals. Second, we have not had a high-throughput,

unbiased behavioral phenotyping pipeline capable of objectively categorizing all of an

individual's movements. Historically, insect descending neuron anatomy, connectivity

and function have been described by backfilling neurons with dye and recording from

individual neurons in locusts, grasshoppers, and cockroaches (for a review see

(Strausfeld, Bassemir, & Singh, 1984)), with more recent studies performing similar

experiments in flies (Hsu & Bhandaway, 2016). While this approach has allowed

researchers to describe the anatomy and electrophysiological responses of individual

neurons, it is inherently low throughput and biased towards larger or otherwise more

accessible neurons.   Additionally, because experiments are typically carried out on

immobile preparations, only in rare cases have investigators been able to link individual

neurons to behavior (e.g. (E. Staudacher & Schildberger, 1998)). While recent technical

and genetic advances in the model fly *Drosophila melanogaster* have improved our

ability to access and manipulate individual descending neurons, to date only a handful of

Drosophila descending neurons have been linked to specific motor outputs (e.g. (Bidaye

et al., 2014; von Philipsborn et al., 2011b; von Reyn et al., 2014).

To assess how descending neurons control motor behaviors on a systems scale, it

will be necessary to move beyond isolated examples and to describe the behavioral

functions of large numbers of descending neurons. Our goal was to identify all of the

behavioral phenotypes observable in one particular setting, freely behaving flies moving

within a two-dimensional arena, for many descending neurons, without any *a priori*

expectation about the neurons' effects on behavior. Namiki et al (Namiki et al, 2017)

created a collection of transgenic *Drosophila* strains that target descending neurons using

the split-GAL4 intersectional system (Pfeiffer et al., 2010) in a cell-type specific manner.

We screened 130 of the sparsest lines in this collection, targeting approximately 160

neurons that are divisible into 58 distinct anatomical cell-types. 40 of these cell types

consist of a single pair of bilaterally symmetric descending neurons, while the remaining

18 categories target populations of 3 to 15 descending neurons with similar

neuroanatomy. We used this split-GAL4 collection to drive the expression of the red-

shifted channelrhodopsin *CsChrimson* (Klapoetke et al., 2014a) in specified subsets of

descending neurons, allowing us to photo-activate these neurons in a temporally precise

fashion. We combined these genetic reagents with a recently described method for

objective, quantitative analysis of behavior (Berman, Choi, Bialek, & Shaevitz, 2014) to

comprehensively identify the behaviors associated with the activation of specific neurons

in an unbiased fashion. Unlike supervised machine learning approaches for classifying

behavior, this approach does not rely on a human-trained classifier to decide which

behaviors are of interest. Instead it captures a wide range of movements by converting

high-dimensional postural dynamics into a two-dimensional map using dimensionality

reduction techniques (Berman et al., 2014). Using this method, we associated 80% of the

descending neurons in our collection with specific behaviors.

We have generated a behavioral dataset from freely walking animals that

comprehensively describes the activation phenotypes of roughly one third to one half of

the total number of fly descending neurons. The size of this dataset has allowed us to

move beyond individual examples to extract general features of descending neuron

function, and therefore to consider how these neurons might encode information to

modulate behaviors. We find that, with a few exceptions, descending neuron control of

behavior appears to be largely modular. In addition, we find many cases in which

descending neuron function is context dependent, even within the confines of a single fly

confined to a two-dimensional substrate.

Results

Establishing a framework for large scale analysis of descending neuron activation

phenotypes

Mapping fly behavior using postural dynamics requires high temporal and spatial

resolution video data from a large number of animals. Accordingly, we built a red light

activation apparatus that consisted of an array of 12 USB cameras that allowed us to film

12 flies in separate chambers simultaneously at high resolution (Figure 1A). We crossed

each split-GAL4 line to a UAS-CsChrimson line, and we filmed six experimental

progeny that had been fed retinal, a co-factor necessary for neuronal activation via

channelrhodopsin, and six genetically identical control flies whose food had not been

supplemented with retinal. The flies were backlit using custom light tables, each

consisting of an array of infrared and red LEDs covered by a diffuser. Each chamber was

a 3 cm "fly bubble" (Klibaite (2017)), which had sloping sides coated with silicone. This

encouraged the flies to remain on the flat floor of the chamber, where they could roam

freely and would remain in the focal plane of the camera (Berman et al., 2014). For each

split-GAL4 line, we recorded 30 trials consisting of a 15-second pulse of red light

followed by a 45 second recovery interval (Figure 1A).

If the descending neuron(s) labeled by a particular split-GAL4 line are involved in

triggering, maintaining, or modulating a particular behavior, then activating these

neurons with *CsChrimson* may be sufficient to activate that behavior. To identify

behavioral phenotypes in an unbiased manner, we utilized the behavior mapping methods

described in Berman et al. (2014). First, we generated a comprehensive "behavior space"

of stereotyped actions that single flies could produce in our assay. We collected a dataset

of approximately 700 million images, which included behaviors recorded from activation

of descending interneuron split-GAL4 lines, previously characterized sparse GAL4

drivers (*fruitless-GAL4* and *pIP10*) that trigger courtship-related behaviors (Stockinger,

Kvitsiani, Rotkopf, Tirián, & Dickson, 2005; von Philipsborn et al., 2011b), and

interneuron drivers targeting the flight neuropil. The additional lines that are not part of

the descending neuron screen were included to sample fly behaviors as widely as

possible, allowing for higher resolution mapping within the space of behaviors. We

computed the behavior space by (1) aligning video images (Figure 1A), (2) decomposing

the pixel value dynamics (which correspond to the fly's posture changes) into a low-

dimensional basis set using principal component analysis (Figure S1), (3) projecting the

original pixel values onto this basis set and transforming those values using a spectral

wavelet function to produce a time series that was (4) embedded into a two-dimensional

"behavior space" (Figure 1B) using *t*-distributed Stochastic Neighbor Embedding (t-

SNE) (van der Maaten & Hinton, 2008).

Each position in the behavior space corresponds to a unique set of postural

dynamics. Nearby points represent similar motions, i.e. those involving related body

parts executing similar temporal patterns.   By observing the video data underlying sub-

regions of the behavior space (Figure S2 and movies S1-S5), we generated a human-

curated version of the behavior space to aid interpretation (Figure 1C). In this behavior

space, anterior directed movements such as eye/antennal grooming and proboscis extension are located at the top (supplemental movie S1). Anterior-directed foreleg movements are on the upper left side (movie S2). Extremely slow or still postures are on the upper right side (movie S3), and complex wing and abdomen movements such as body and abdomen grooming, abdomen bending and wing extension are in the center (movie S4). Locomotion, ranging from slow (left) to fast (right) is at the bottom (movie S5).

Red peaks, or density maxima, represent the fly behaviors observed most frequently in our data set. These tend to be repetitive, stereotyped behaviors, such as walking or grooming, that our analysis methodology is most sensitive at detecting. By definition, we could not detect behaviors occuring over time-scales faster than 50 Hz, the Nyquist frequency of our system. Approximately 93% of all video image data points could be embedded in this space, irrespective of whether the red light was on or off (Figure S3), indicating that the majority of red light activated behaviors are well represented in the behavior space. Un-embeddable imaging errors, such as the fly wandering partially out of frame, are randomly distributed within the dataset.

Entropy of behavior space density provides a quantitative and sensitive measure of optogenetic activation phenotypes

Having established a behavior space representing the full repertoire of fly behaviors that could be captured with our apparatus, we next examined which parts of this space were occupied when individual or subsets of descending neurons were optogenetically activated by CsChrimson. We focused on 130 split-GAL4 lines that

targeted descending neurons with little, or no, extraneous expression in other neurons.

We first considered the timing and duration of red light triggered behaviors. If

descending neuron activation triggered a particular behavior represented in the behavior

space, then we expect that the density of that line in the behavior space should shift into

the region that represents that behavior during periods of red light activation. For

example, upon red light activation, retinal-fed flies expressing *CsChrimson* in a

descending neuron line targeting DNg07 and DNg08 (SS02635) groomed their heads

(supplemental movie S6). We identified regions in the behavior space that experienced a

statistically significant shift in density for experimental flies during the first three seconds

of red light compared to a window at the end of the recovery period when the red light

was off (Figure 2B, C, Figure S4). This same region in the behavior space did not

undergo a significant shift in the control flies (Wilcoxon rank-sum test $p < 0.05$ using the

Dunn–Šidák correction for multiple hypotheses (Šidák, 1967)). Likewise, when

considering densities over the whole behavior space in three second sliding windows, the

experimental, but not the control, flies shift into the head grooming region (arrowheads,

Figure 2C).

The shift in behavior in the experimental animals, and the timing of this shift

relative to red light activation, can be detected as a reduction in the entropy of the

behavior space density during this epoch (Figure 2A). Entropy measures the degree of

disorder inherent in the distribution of the flies in the behavior space. When the red light

was off, flies exhibited a range of different behaviors, and the probability that they

performaned any one behavior was low. This results in a low probability density

distributed throughout the behavior space and correspondingly high entropy (Figure 2A).

Upon red light activation, the experimental fly line engaged in red light triggered

behaviors at the expense of other natural behaviors. This increased the probability that

they occupied a small region within the behavior space, generating a drop in entropy

whose timing and duration mirrored that of the red light triggered behaviors (Figure 2A

movie S7). We can therefore use entropy as a proxy for the duration and onset of red light

triggered movements in the behavior space without needing to know, *a priori*, which

behaviors are activated (i.e. which part of the behavior space to examine).

The region density and entropy are quantitative measurements sensitive to small

changes in behavior map distribution. We therefore used these values to identify subtle

phenotypes that could not be easily identified by manual inspection of the movies. For

example, the rapid activation of descending neuron DNg25 induced a transient rapid

running phenotype (Figure 2E & 2F) that was identified by the transient drop in entropy

in the behavior space (Figure 2D) and the transient increase in density in the fast

locomotion region of the space (Figure 2E & 2F, supplemental movie S8).

Comprehensive characterization of descending neuron split-Gal4 line activation

phenotypes

To characterize the time-course and likelihood of optogenetically-induced

phenotypes across the entire collection of descending neuron lines, we examined the

entropy time course of each line (Figure 3). We found that most lines displayed the

largest entropy drop immediately after red light activation (Figure 3A). For roughly a

third of the lines, this entropy drop persisted throughout the entire red light activation

window (Figure 3B). For most of the rest of the lines, however, the entropy drop was

transient and diminished after several seconds (Figure 3B). For a minority of lines, the

entropy minimum occurred near the middle or end of the activation window (Figure 3B).

We reviewed the raw video data for these lines and found that most of these flies

performed some action upon red light activation, followed by a freeze. This explained

why the entropy was lower in the later part of the activation window, because consistent

stillness is a low entropy state (see Figure S5 for a line by line description of

phenotypes). We therefore performed our system-wide analysis using the first 3 seconds

of the red light activation period, because this time period captured the majority of

CsChrimson activated behaviors.

In our initial analysis, we looked for behaviors produced when our descending

neuron lines were activated using a comparatively low level of red light, (5 mW/cm^2).

Under these conditions, 91 of the 130 lines (69%) displayed a statistically significant

increase in density of some area of the behavior space. We then re-tested most of the 41

lines that did not produce a significant density increase by driving CsChrimson at higher

levels by growing the flies on food containing an increased retinal concentration and

exposting flies to higher intensity red light (9 mW/cm^2). Under these conditions, 80% of

the lines that had previously displayed no phenotype produced a statistically significant

increase in density in the behavior space.

Pooling the data from the low and high activation protocols, we detected

statistically significant increases in the behavior space in 119 of the 130 (90%)

descending neuron lines (Figure 4A). In 86 cases, we observed an increased density in

only a single statistically significant region in the behavior space. However some lines

generated density increases in multiple non-contiguous regions of the behavior space

(Figure 4A, examples shown in Figure 4B-D, supplemental movies S9-S11, supplemental

figure S5). In many cases, these multi-region lines reflect multiple behaviors performed

approximately simultaneously by the flies. For example, a line targeting DNp10 induced

anterior reaching movements and wing flicking with similar timing (Figure 4D, red and

blue regions respectively). However, in other cases, multiple activated regions reflect a

stereotyped sequence of behaviors. For example, the DNp09 line shown in Figure 4B

repeatedly ran and then paused throughout the entire 15 second activation period. The

increased density in the run region of the behavior space (Figure 4B, red) appeared before

the increased density in the paused region (Figure 4B, blue), reflecting the sequential

timing of the two behaviors. However, the flies rapidly became asynchronous as they

repeated this series of behaviors, so this behavioral series is detected as density increases

in the running and still regions throughout the red light activation window. A line

targeting descending neuron DNb01 displayed a simple behavior series; flies produced an

anteriorly directed twitch of the front legs when the red light was turned on (Figure 4C,

red region), then froze for the majority of the red light activation period (Figure 4C, black

region), and then twitched when the light was turned off (Figure 4C, blue region). Thus,

examining the timing of density shifts illuminates the more complicated behavior series

produced by red light activation. This level of analysis is provided for all lines in

supplemental Figure S5.

Behavioral result of descending neuron activation is often context dependent

Why does activation of some descending neurons result in multiple, distinct

behavioral outputs? One possibility is that the behavioral output of some descending

neurons depends on the behavioral context of the fly when the descending neuron is

activated. To address this possibility, we calculated the mutual information between the

density distribution of the experimental flies in the behavior space at 1.5 to 0.5 seconds

before the red light was turned on versus the first second after red light activation. Mutual

information is a non-linear measure of the degree of dependence between two variables

and is typically measured in units of bits (Cover & Thomas, 2005). The higher the mutual

information, the more the first variable, here the behavior of flies immediately prior to

red light activation as measured by their distribution in the behavior space at t = -1.5 to -

0.5 seconds, informs the value of the second variable, the region of the behavior space

occupied in the first second of red light activation.

We found that, in all cases, experimental animals displayed non-zero mutual

information between the pre- and post-stimulation behaviors (Figure 5A). In addition, for

most lines, more information was available in the experimental flies than in the controls

(Figure 5B). This means that even in those cases where red light activation produced only

one significant region in the behavior space, the fly's activity prior to red light activation

influenced whether or not it performed the behavior. However, lines with multiple red

light activated regions in the behavior space were also those with a relatively high level

of mutual information (Figure 5A-B). Thus, a given fly's behavior before red light

activation was highly informative of which behavior that fly would perform after red

light activation, as indicated by the different significantly activated regions in the

behavior space. Figure 5C displays this phenomenon for one of the lines with the highest

mutual information, SS02542 (asterisk in figure 5A-B, also shown in Figure 4C). Here,

if the flies were performing an action in the wind/abdomen movement regions of the

behavior space prior to the stimulation, they were likely to perform an anterior movement

(region 1) immediately following stimulation.   Similarly, flies performing anterior

grooming were likely to transition to the small anterior twitch region (region 2), and flies

that were initially still tended to remain still post-stimulation (region 3).

Individual descending neurons produce mainly stereotyped, modular behaviors

So far, we have analyzed split-GAL4 lines as if they were a proxy for individual

descending neurons or anatomical classes of descending neurons. However, these lines

vary in both their strength of expression and in the number and identity of additional cells

labeled. To estimate phenotypes for individual descending neurons, we therefore

averaged the behavior space densities of multiple lines for those cases where we had

multiple lines targeting the same descending neuron (Figure 6). Using this method, and

combining it with those descending neurons for which we had only a single

representative split-GAL4 line, we estimated phenotypes for 47 of the 58 descending

neuron cell types. We have also included six lines and line averages that target two

different types of descending neurons cleanly, but for which we have no lines that target

each type individually. Twenty-six descending neurons drove locomotion phenotypes and

ten drove anterior directed foreleg movements. We also identified six new descending

neurons that triggered wing and abdomen movements (plus the previously published

pIP10 (von Philipsborn et al., 2011b)), two that drove anterior grooming, one that drove

abdomen stroking, and four that drove still or slow behaviors.

In general, we found that activation of each type of descending neuron drove

behaviors that mapped to a relatively small region of the behavior space. For example,

some descending neurons drove slow locomotion, whereas others drove fast locomotion.

Only a few, such as DNa01, DNa02 and DNp26, seemed to produce a global increase in

locomotor activity. Likewise, we found descending neurons that produced different types

of grooming, such as head grooming (DNg07 & DNg08, and DNg12) or abdomen

grooming (DNp29), different types of anterior reaching movements (DNg10 versus

DNg13) and different types of slow movements (e.g. DNd02 versus DNp02).

Discussion

Using optogenetic activation and automated behavioral quantification, we

assigned behavioral phenotypes to 80% of the descending neurons cell types in our

collection of lines, or one third to one half of the estimated total number of descending

neurons present in the fly. Using a dataset of this scope, it is possible for the first time to

move beyond isolated examples to consider systems-level trends in how descending

neurons control behaviors.

For several reasons, this is unlikely to be a comprehensive categorization of the

activation effects of these descending neurons. First, behaviors performed more quickly

than the Nyquist frequency of 50Hz for our movies could not be detected. Second, we

assayed only behaviors that can be activated when flies are standing and walking.

Descending neurons controlling behaviors gated by flight, for example, would not be

detected. Third, we assayed only males, so any female-specific behaviors may not be

identified. Finally, we assayed solitary flies, so any behaviors gated by social

interactions, for example courtship, may not have been detected.

There are several, non-mutually exclusive ways a limited number of seemingly

highly modular descending neurons could encode the wide range of behaviors undertaken

by freely moving animals. First, descending neurons could be more important for

triggering and maintaining behaviors than for controlling individual details of a given

motor program (Heinrich, 2002). Many motor programs, particularly those controlling

repetitive, rhythmic actions such as walking or stridulation, can function in the absence of

descending control ((Bentley, 1977; Kien, 1983), for a review on walking circuits see

(Ritzmann & Bü Schges, 2007)). For example, Hedwig (1994) identified two pairs of

descending neurons that control stridulation in grasshoppers. In this system, tonic

activation of the descending neurons was sufficient to induce and modulate the activity of

the stridulation central pattern generator in the thorax, indicating that the descending

neurons play only a limited role in patterning stridulation. Several of our lines, including

the DNg07 & DNg08 head grooming line (Figure 2A & 2B), appear to reflect a similar

phenomenon, driving a repeated stereotyped behavior during the entire CsChrimson

activation window.

Second, behaviors might be controlled not by single descending neurons acting as

command neurons, but by combinations of descending neurons acting in concert

(Heinrich, 2002). Neuroanatomy suggests this possibility; roughly a third of described

descending neurons appear to have unique projection patterns in *Drosophila* and

Calliphora, while the rest share common input and/or output regions in the brain and

ventral nerve cord with other descending neurons (Gronenberg & Strausfeld, 1990; Milde

& Strausfeld, 1990; Namiki et al, 2017). In addition, several examples illustrate that

multiple descending neurons control the same specific behaviors (Griss & Rowell, 1986;

Gronenberg & Strausfeld, 1990; Hensler, 1992; Kanzaki, Ikeda, & Shibuya, 1994; Kien,

1983, 1990; Milde & Strausfeld, 1990; Rowell & Reichert, 1986; E. M. Staudacher,

2001). In this case, Larimer et al (1988) suggest that sufficiently strong stimulation of one

neuron in a command cohort or module is sufficient to recruit the activity of the other

descending neurons, triggering the behavior. Our data do not allow us to definitively

address this question. However, the large number of descending neurons that drive

similar patterns of fast locomotion, slow locomotion and anterior reaching suggest that,

for these motor circuits at least, this is a possibility. Alternatively, it is possible that many

of these descending neurons modulate distinct aspects of these motor programs.

Third, another way to generate behavioral complexity is through coding different

behaviors via combinations of descending neurons. We examined a few lines that target

multiple descending neurons. We compared behaviors produced by these "multi-hit"

split-GAL4 lines with lines that targeted the individual neurons and found only weak

evidence for the emergence of new behaviors when descending neurons were triggered in

combination. For example, both DNa05 and DNd02 produce slightly different

phenotypes when activated in combination with DNa07 and DNd03, respectively, as

compared to when lines targeting these neurons are activated alone (see Figure S5).

However, our collection contains, by design, few lines driving expression in

combinations of descending neuron types. Therefore, further exploration of this idea will

require the generation and characterization of additional lines.

Finally, descending neurons could be re-used in multiple behavioral contexts.

While there are to date no published examples of a single descending neuron triggering

different context-dependent behaviors, there are multiple cases in which descending

neurons exhibit different physiological responses depending on the state of the animal

(e.g. walking, flying, courting, etc.) (Böhm & Schildberger, 1992; B Hedwig, 2000; R.

M. Olberg, 1983; R. Olberg & Willis, 1990; E. M. Staudacher, 2001; E. Staudacher &

Schildberger, 1998; Strausfeld & Bassemir, 1985; Zorović & Hedwig, 2011). Our results

strongly support a role for context dependency for two reasons. First, the high level of

mutual information between behaviors immediately before and after red light activation

seen in lines that have multiple red light activated regions indicates that even within the

relatively confined system of our assay, the behavior of the fly immediately before

descending neuron activation biases the behavioral output in many cases. Second, our

observation that substrate specific behaviors, such as foreleg tapping, reaching, and

locomotion are strongly represented in our dataset, while flight and courtship behaviors

are less prevalent suggests that descending neuron outputs may be context dependent. By

forcing the flies to remain on a two dimensional substrate in isolation, we may have

observed predominantly indirect results of behaviors that would normally take place in a

different context. For example, when we activated a line expressing in DNp01, the giant

fiber, a neuron known to elicit a rapid escape response initiated by a jump when

optogenetically activated (Lima & Miesenbock, 2005) we detected the flies running after

returning back to the ground because the jump was too fast (~30 ms) to be detected in our

assay. It is also possible that some of these descending neurons are never naturally

activated in the two-dimensional context of walking and that proprioceptive feedback

may have generated abnormal behaviors in our assay.

Our objective, quantitative assessment of a descending neuron activation screen

provides a foundation for understanding descending neuron functions more broadly.

Using similar analytical approaches to study the results of descending neuron activation

and inactivation in other behavioral settings in the future will broaden our understanding

of how descending neurons direct motor patterns in specific behavioral contexts and

reveal how the fly's rich behavioral repertoire can be encoded with only a few hundred

neurons.

Materials and Methods

Fly stocks and fly handling

The descending neuron split-GAL4 driver collection is described in Namiki et al

(Namiki et al 2017). Male flies were crossed to virgin females carrying *20xUAS-*

CsChrimson-mVenus (Klapoetke et al., 2014b) integrated into the *attP18* landing site

(Markstein, Pitsouli, Villalta, Celniker, & Perrimon, 2008) and transferred to Dickson lab

power food (1L water, 10g agar, 80g Brewer's yeast, 20g yeast extract, 20g peptone, 30g

sucrose, 60g dextrose, 0.5g $MgSO_4*6H20$, 0.5g $CaCl_2*2H20$, 6mL propionic acid and

7mL 15% Nipagin). For the initial screen, experimental animals were raised on power

food supplemented with 0.2 mM retinal. This concentration was increased to 0.4 mM for

animals that were re-assayed at a higher light intensity. All flies (except parental stocks)

were handled under 453 nm blue LEDs and reared in dark blue acrylic boxes (acrylic

available from McMaster-Carr, # 8505K84) at 22°C on a 12 hour lights on:12 hour lights

off day:night cycle. Individual male flies were collected upon eclosion and housed singly

in 2mL wells in a 96 well "condo," with power food (with or without retinal) deposited in

the bottom of each well, which was sealed at the top with an airpore sheet (Qiagen

#195761). Flies were imaged at age 7-12 days, within 4 hours of lights on.

 Data Collection

Single flies were loaded into individual trays made from 4.5 mm clear acrylic

topped with a fly "bubble" 3 cm in diameter and 4 mm at its tallest point, which was

vacuum molded from clear 0.020" PETG thermoform plastic (WidgetWorks, available on

Amazon) (Klibaite (2017)). PETG was placed in a frame, heated in a Oster Convection

Bake pizza oven set at 350°F until the plastic started to deform (about 20 seconds), then

placed on a vacuum manifold. To further encourage the flies to remain on the 2

dimensional acrylic surface, the bubbles were coated with Sigmacote siliconizing reagent

one day prior to imaging and lightly wiped with ethanol to remove the excess silicone.

For each descending neuron split-GAL4 line six retinal-fed experimental animals

and six non-retinal-fed control animals were imaged simultaneously. For imaging, flies

were placed in individual fly bubbles atop custom light tables (3 identical light tables,

each imaging 4 flies). These tables consisted of a custom light board topped with a 0.75"

3D printed white plastic standoff that was lined with infrared reflective tape, and which

was capped with a diffuser made from 0.125" white plexiglass acrylic (available from

eplastics.com, # ACRY24470.125PM24X48), which had 50% light transmittance. The

light board itself consisted of an array of 256 IR LEDs (Osram Opto SFH 4050-Z, 850

nm wavelength) arrayed in a 16 x16 pattern, spaced 7.14 mm apart, and 64 red LEDs

(Philips Lumileds, LXM2-PD01-0050, 627 nm) arranged in an 8x8 pattern, spaced 14.28

mm apart. IR and red LED intensity was controlled separately by 0-2.5V control

voltages, yielding 0-100mA for the IR LEDS and 0-400 mA for the red LEDs. We set the

IR LEDs to 1V, which provided even illumination without overheating the flies. We used

0.2V (4.5 mW/cM^2) red light in the initial screen and 1.0V (9 mW/cm^2) red light when

re-screening a subset of lines. All three light tables were connected to a 68-Pin

unshielded I/O connector block (National Instruments, CB-68LP), then to an M Series

multifunction DAQ board (National Instruments, NI USB-6281), so that all tables could

be run simultaneously from a single computer. Each light table was topped with a 10"

square frame constructed from off the shelf parts from Thorlabs, which supported four

1.3 MP grayscale USB cameras (Point Grey FL3-U3-13Y3M-C, 1 camera per fly) on

optical rails, whose X/Y/Z coordinates could be adjusted relative to the fly bubble. Each

camera was fitted with an HR F2/35mm lens from Thorlabs and an 800 nm longpass

filter (Thorlabs, FEL0800). Each set of four cameras was connected to a separate Dell

Precision T3600 Tower Workstation. Each was fitted with two 100 GB internal solid

state drives, so that 2 cameras wrote to each SSD.

486   Cameras were programmed using NI-MAX and custom software written in

Labview (National Instruments). Data acquisition and the LED light tables were

controlled by custom software written in Labview. In brief, a master computer ran a

single program that (1) turned the red and infrared (Berman et al., 2014) LEDs on for all

three light tables, the former running a program of 15 seconds, then off for 45 seconds,

for 30 cycles; (2) started all 12 cameras recording; (3) recorded the position of the fly's

centroid for each frame for each camera; and (4) grabbed the frame number for each

camera over the network every 2-3 frames, and wrote the frame number and red LED

status from the light tables to a single text file. All movies were recorded as

uncompressed avi files at 100 frames per second. Each camera was set to 1024 x 1024

pixel resolution that encompassed the entire 3 cM arena. However, flies were tracked

using a blob detector and only a 150x150 pixel box centered on each fly was saved and

used for analysis.

Behavior Space Generation

Our approach for generating a behavior space largely follows the methodology

originally described in Berman, 2014 (Berman et al., 2014), which describes much of the

procedure in additional detail. We first segmented flies using Canny's method for edge

detection (Canny, 1986) and morphological dilation to find the outline of the fly. All

pixels within the corresponding closed curve were considered part of the fly. We assumed

that all flies had identical morphology but variable sizes. We calculated a rescaling factor

for each fly by segmenting 100 randomly-selected images from a single fly and finding

the pixels belonging to that fly's body (head, thorax, and abdomen) in each of them,

ignoring pixels associated with the wings and legs. Body pixels were assigned via a two-

component Gaussian mixture model, and the average value of the number of pixels was

chosen as the body area. All frames from a single movie were then uniformly re-scaled to

make the number of body pixels in the average image equal to that in a reference image

of a fly. We then rotationally aligned segmented, recalled images by finding the maximal

angular cross-correlation of the magnitudes of the two-dimensional polar Fourier

transforms between the image and a reference image. This reference image was common

to all aligned images. Translational registration was then performed by maximizing the

spatial cross-correlation.

Postural decomposition was performed as described in Berman (2014). Images

were Radon-transformed using a 2 degree spacing, and the 9,781 Radon-space pixels that

contained the most variance were kept for further analysis (>95% of the total variance).

We then performed principal components analysis (PCA) on these data, keeping the 50

modes capturing the most variance (>90% of the total variance). We projected the

segmented and aligned images onto the found eigenvectors to create a set of time series

that were representative of the postural movements of the fly. To obtain dynamic

information about these time series, we applied a Morlet continuous wavelet transform to

these time series. We transformed each mode separately, using 25 frequency channels

that were dyadically spaced between 1 Hz and 50 Hz, retaining only the amplitudes of the

resulting complex numbers.

Low dimensional embedding of these wavelet time series using t-Distributed

Stochastic Neighbor Embedding (t-SNE) (van der Maaten & Hinton, 2008) largely

followed the approach in (Berman et al., 2014) as well. A distance metric between points

in time was calculated via the the Kullback-Leibler divergence (Cover & Thomas, 2005)

between their associated normalized mode-frequency spectra. Because this data set

contains several orders of magnitude more data than can be calculated through brute-

force minimization of the t-SNE cost function, we used the sub-sampling technique

described in (Berman et al., 2014) to identify 600 representative data points from each of

the recording sessions. From here, points were randomly assigned subsequent groupings

such that each of these groups contained 36,000 data points. The same sub-sampling

process was performed amongst these data points, but now keeping twice as many data

points as in the previous iteration. This process was repeated until a data set of 36,000

points was obtained. We minimized t-SNE for this data set to create a low-dimensional

embedding. We used the re-embedding procedure described in (Berman et al., 2014) to

include data from outside the 36,000-point training set into the embedding, resulting in

the overall density seen in Figure 1.

Statistical Analysis

Our main goal for the statistical analysis of the behavior space data was to isolate

regions of the map that were significantly affected by optogenetic stimulation. Here, we

assessed significance by (1) comparing the flies' behavior when the LED was on versus

when the LED was off, and (2) requiring that the effect of the LED stimulation be larger

in the experimental flies than in the control flies. Specifically, we compared the flies'

behavior during the first three seconds of stimulation (t=0s to t=3s, where the LED turns

on at t=0) to their behavior between stimulation (t=30s to t=45s). To statistically assess

whether a particular region of the behavior space was significantly affected by the

stimulus, we first defined $\rho_{i,n}^{on}(x,y)$ to be the average behavior space density for fly i

during the nth cycle at location (x,y) during the first 3 seconds of excitation and

$\rho_{i,n}^{off}(x,y)$ to be the same, but during the 15s window furthest from the stimulation. We

then tested whether $\rho_{i,n}^{on}(x,y)$ was significantly different from $\rho_{i,n}^{off}(x,y)$ through a

Wilcoxon rank sum test with Šidák corrections (p < .05 after corrections) (Šidák, 1967).

To calculate the number of corrections, we conservatively assumed that the number of

measurements was equal to 2^H, where H was the entropy of the mean density of the

behavior space. This is likely an over-estimate of the number of comparisons, but it

provides an upper-bound for the number of distinctions that could be made.

To compare the effect of the optogenetic stimulus on the experimental flies to that

of the effect on the control flies, we computed the quantity $\chi_{i,n}(x,y) = \rho_{i,n}^{on}(x,y) -$

$\frac{1}{2}(\rho^{off}_{i,n-1}(x,y) + \rho^{off}_{i,n}(x,y))$, which was the behavior space density during light

stimulation compared to the average of the two preceding time periods with no light

stimulation. We thus assessed statistical significance by using a Wilcoxon rank sum test

with Šidák corrections (p < .05 after corrections) to compare

$\{\chi_{i,n}(x,y)\}_{i \in experimental\ flies}$ with $\{\chi_{i,n}(x,y)\}_{i \in control\ flies}$. For a point, (x,y), in the

behavior space to be considered significantly affected by the stimulus, we required that

both of these tests—within experimental flies test and experimental versus control flies

test—yielded a significant result.

Behavior activation maps for individual descending interneurons (Figure 6) were

calculated by averaging together the maps of significantly significant activations

($E[\chi_{i,n}(x,y)]_{i,n} > 0$) from each of the lines exciting that neuron.

Stimulation-response entropy curves (Figure 2A & D) were generated by first

aligning each time point to its associated phase within the 60 second LED on-off cycle.

For each phase within the cycle, we found all embedding points from all relavent trials

that were detected within ± 200 ms (using periodic boundary conditions). We then

generated a histogram of these points, normalized and convolved the resulting values

with a symmetric two-dimensional Gaussian of width $\sigma = 2$, to generate a probability

density function, $p_t(x,y)$. From this, the entropy curve value at phase t was given by

$H(t) = \int dx\, dy\, p_t(x,y) \log p_t(x,y)$. We then pooled data from all individuals of a

specific type together (i.e. all control flies from a given line or all experimental flies from

a given line) to calculate these curves.

Mutual information between pre-stimulus behavior space densities and post-

stimulus regions was computed by numerically integrating the integral:

$$MI\left(\rho_{pre}; R_{post}\right) = \sum_{k=0}^{m} \int d\vec{x} \, p_{pre}(\vec{x}|R_k) \log_2 \frac{p_{pre}(\vec{x}|R_k)}{\sum_{\ell=0}^{m} p_{pre}(\vec{x}|R_k)p(R_k)}, \text{ where } p_{pre}(\vec{x}|R_k) \text{ is}$$

the conditional probability of observing the fly's behavior to be at location $\vec{x}$ between 1.5

and 0.5 seconds before the stimulus onset and $p(R_k)$ is the probability that the fly

transitions to region R_k following the stimulus onset. Finite data-size corrections were

performed by drawing subsets of the data with replacement and extrapolating to an

infinite number of trials, and error bars were generated by extrapolating the calculated

variance in a similar manner (Bialek, 2012). The region of transition for each trial was

assigned by finding the mode of the behavior space distribution during the first second

subsequent to the onset of the stimulus. If the location of the mode of the distribution for

that trial was within or closer than 5-pixels to the edge of a region, it was assigned to that

region, unless another region was closer. Trials not assigned to any of the regions were

given a "zero" label, as reflected in the previous equation.

To provide a sense of scale, if there are N significantly activated regions, the

maximum possible mutual information one could potentially measure between the prior

distribution and the activated region would be $\log_2(N)$ bits. Note, however, that we

assigned an additional state corresponding to the fly performing a behavior outside of the

significantly activated regions subsequent to the light turning on, thus making the

maximal possible mutual information $\log_2(N + 1)$. This additional "zero" state is

necessary to account for the possibility that the significant regions might be exhibited

only in a context-dependent manner, leading to no significant phenotype when the fly is

performing some behaviors at the onset of red light stimulation and leading to a

phenotype if other actions are being exhibited.

Acknowledgements

We thank Vivek Jayaraman for reagents and feedback, Jan Ache and Ugne Klibaite for discussions, Todd Laverty and members of the Janelia Fly Core for their support, and Steven Sawtelle, Igor Negroshov, Ben Arthur and Roger Rogers for help with the rig, fly bubble design and fabrication. The driver lines were developed as part of the Descending Interneuron Project Team at the Janelia Research Campus. This project was supported by the Janelia Research Campus Visitor Program and NIH GM098090.

References

Alivisatos, A. P., Chun, M., Church, G. M., Greenspan, R. J., Roukes, M. L., & Yuste, R. (2012). The brain activity map project and the challenge of functional connectomics. *Neuron*, *74*(6), 970–4. http://doi.org/10.1016/j.neuron.2012.06.006

Bentley, D. (1977). Control of cricket song patterns by descending interneurons. *Journal of Comparative Physiology*, *116*(1), 19–38. http://doi.org/10.1007/BF00605514

Berman, G. J., Choi, D. M., Bialek, W., & Shaevitz, J. W. (2014). Mapping the stereotyped behaviour of freely moving fruit flies. *Journal of the Royal Society, Interface*, *11*(99), 20140672. http://doi.org/10.1098/rsif.2014.0672

Bialek, W. (2012). *Biophysics: searching for principles*. Princeton: Princeton University Press.

Bidaye, S. S., Machacek, C., Wu, Y., & Dickson, B. J. (2014). Neuronal control of Drosophila walking direction. *Science*, *344*(6179), 97–101. http://doi.org/10.1126/science.1249964

Böhm, H., & Schildberger, K. (1992). Brain Neurones Involved in the Control of

Walking in the Cricket Gryllus Bimaculatus. *Journal of Experimental Biology*.

Canny, J. (1986). A Computational Approach to Edge Detection. *IEEE Transactions on*

Pattern   Analysis   and   Machine   Intelligence,   *PAMI-8*(6),   679–698.

http://doi.org/10.1109/TPAMI.1986.4767851

Cover, T. M., & Thomas, J. A. (2005). *Elements of Information Theory*. *Elements of*

Information Theory. http://doi.org/10.1002/047174882X

Gras, H., & Kohstall, D. (1998). Current injection into interneurones of the terminal

ganglion modifies turning behaviour of walking crickets. *Journal of Comparative*

Physiology. A, Neuroethology, Sensory, Neural, and Behavioral Physiology, *182*(3),

351–361.

Griss, C., & Rowell, C. H. F. (1986). Three descending interneurons reporting deviation

from course in the locust. *Journal of Comparative Physiology. A, Neuroethology,*

Sensory, Neural, and Behavioral Physiology, *158*(6), 765–774.

Gronenberg, W., & Strausfeld, N. J. (1990). Descending neurons supplying the neck and

flight motor of Diptera: physiological and anatomical characteristics. *The Journal of*

Comparative Neurology, *302*(4), 973–991.

Hedwig, B. (1994). A cephalothoracic command system controls stridulation in the

acridid grasshopper Omocestus viridulus L. *J Neurophysiol*, *72*(4), 2015–2025.

Hedwig, B. (2000). Control of Cricket Stridulation by a Command Neuron: Efficacy

Depends on the Behavioral State. *J Neurophysiol*, *83*(2), 712–22.

Heinrich, R. (2002). Impact of descending brain neurons on the control of stridulation,

walking, and flight in orthoptera. *Microscopy Research and Technique*, *56*(4), 292–

301.

Hensler, K. (1992). Neuronal co-processing of course deviation and head movements in

locusts. *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural,*

and Behavioral Physiology, *171*(2), 257–271.

Hsu, C. T., & Bhandawat, V. (2016). Organization of descending neurons in Drosophila

melanogaster. *Scientific Reports*, *6*, 20259. http://doi.org/10.1038/srep20259

Kanzaki, R., Ikeda, A., & Shibuya, T. (1994). Morphological and physiological

properties of pheromone-triggered flipflopping descending interneurons of the male

silkworm moth, Bombyx mori. *Journal of Comparative Physiology. A,*

Neuroethology, Sensory, Neural, and Behavioral Physiology, *175*(1), 1–14.

Kien, J. (1983). The Initiation and Maintenance of Walking in the Locust: An Alternative

to the Command Concept. *Proceedings of the Royal Society B: Biological Sciences*,

219(1215), 137–174. http://doi.org/10.1098/rspb.1983.0065

Kien, J. (1990). Neuronal activity during spontaneous walking--I. Starting and stopping.

Comparative Biochemistry and Physiology. A, Comparative Physiology, *95*(4), 607–

621.

King, D. G., & Wyman, R. J. (1980). Anatomy of the giant fibre pathway in Drosophila.

I. Three thoracic components of the pathway. *Journal of Neurocytology*, *9*(6), 753–

770. http://doi.org/10.1007/BF01205017

Klapoetke, N. C., Murata, Y., Kim, S. S., Pulver, S. R., Birdsey-Benson, A., Cho, Y. K.,

… Boyden, E. S. (2014b). Independent optical excitation of distinct neural

populations. *Nature Methods*, *11*(3), 338–46. http://doi.org/10.1038/nmeth.2836

Klibaite, U., Berman, G. J., Cande, J., Stern, D. L., & Shaevitz, J. W. (2017). An

unsupervised method for quantifying the behavior of paired animals. *Physical*

Biology, *14*(1), 015006. http://doi.org/10.1088/1478-3975/aa5c50

Larimer, J. L. (1988). The command hypothesis: a new view using an old example.

Trends in Neurosciences, *11*(11), 506–510. http://doi.org/10.1016/0166-

2236(88)90013-6

Lima, S. Q. & Miesenböck, G. (2005). Remote control of behavior through genetically

targeted photostimulation of neurons. *Cell*, *121*(1), 141-152.

http://doi.org/10.1016/j.cell.2005.02.004

Marder, E. & Goaillard, J.-M. (2006). Variability, compensation and homeostasis in

neuron and network function. *Nature Review Neuroscience*, *7*, 563-574.

http://doi.org/10.1038/nrn1949

Markstein, M., Pitsouli, C., Villalta, C., Celniker, S. E., & Perrimon, N. (2008).

Exploiting position effects and the gypsy retrovirus insulator to engineer precisely

expressed transgenes. *Nature Genetics*, *40*(4), 476–83. http://doi.org/10.1038/ng.101

Milde, J. lgen J., & Strausfeld, N. J. (1990). Cluster organization and response

characteristics of the giant fiber pathway of the blowfly Calliphora erythrocephala.

The Journal of Comparative Neurology, *294*(1), 59–75.

http://doi.wiley.com/10.1002/cne.902940106

Namiki, S., Dickinson, M.H., Wong, A.M., Korff, W., and Card, G.M. (2017) The

functional organization of descending sensory-motor pathways in *Drosophila.*

bioRxiv.

Okada, R., Sakura, M., & Mizunami, M. (2003). Distribution of dendrites of descending

neurons and its implications for the basic organization of the cockroach brain. *The*

Journal of Comparative Neurology, *458*(2), 158–174.

Olberg, R. M. (1983). Pheromone-triggered flip-flopping interneurons in the ventral

nerve cord of the silkworm moth,Bombyx mori. *Journal of Comparative Physiology*

A, *152*(3), 297–307. http://doi.org/10.1007/BF00606236

Olberg, R., & Willis, M. (1990). Pheromone-modulated optomotor response in male

gypsy moths, Lymantria dispar L.: Directionally selective visual interneurons in the

ventral nerve cord. *Journal of Comparative Physiology A*, *167*(5).

http://doi.org/10.1007/BF00192665

Pfeiffer, B. D., Ngo, T.-T. B., Hibbard, K. L., Murphy, C., Jenett, A., Truman, J. W., &

Rubin, G. M. (2010). Refinement of tools for targeted gene expression in

Drosophila. *Genetics*, *186*(2), 735–755.

Ritzmann, R. E., & Bü Schges, A. (2007). Adaptive motor behavior in insects. *Current*

Opinion in Neurobiology, *17*, 629–636. http://doi.org/10.1016/j.conb.2008.01.001

Rowell, C. H., & Reichert, H. (1986). Three descending interneurons reporting deviation

from course in the locust. II. Physiology. *Journal of Comparative Physiology. A,*

Sensory, Neural, and Behavioral Physiology, *158*(6), 775–94.

Šidák, Z. (1967). Rectangular Confidence Regions for the Means of Multivariate Normal

Distributions. *Journal of the American Statistical Association*, *62*(318), 626–633.

http://doi.org/10.1080/01621459.1967.10482935

Simon, J. C., & Dickinson, M. H. (2010). A new chamber for studying the behavior of

Drosophila. *PLoS ONE*, *5*(1), e8793.

Staudacher, E. (1998). Distribution and morphology of descending brain neurons in the

cricket gryllus bimaculatus. *Cell and Tissue Research*, *294*(1), 187–202.

Staudacher, E. M. (2001). Sensory responses of descending brain neurons in the walking

cricket, Gryllus bimaculatus. *Journal of Comparative Physiology. A*, *187*(1), 1–17.

Staudacher, E. & Schildberger, K. (1998). Gating of sensory responses of descending

brain neurones during walking in crickets. *Journal of Experimental Biology*.

Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirián, L., & Dickson, B. J. (2005). Neural

circuitry that governs Drosophila male courtship behavior. *Cell*, *121*(5), 795–807.

Strausfeld, N. J., & Bassemir, U. K. (1985). Lobula plate and ocellar interneurons

converge onto a cluster of descending neurons leading to neck and leg motor

neuropil in Calliphora erythrocephala. *Cell and Tissue Research*, *240*(3), 617–640.

http://doi.org/10.1007/BF00216351

Strausfeld, N. J., Bassemir, U., Singh, R. N., & Bacon, J. P. (1984). Organizational

principles of outputs from Dipteran brains. *Journal of Insect Physiology*, *30*(1), 73-

96. https://doi.org/10.1016/0022-1910(84)90109-4

Strausfeld, N. J., & Gronenberg, W. (1990). Descending neurons supplying the neck and

flight motor of Diptera: organization and neuroanatomical relationships with visual

pathways. *The Journal of Comparative Neurology*, *302*(4), 954–972.

van der Maaten, L., & Hinton, G. (2008). Visualizing Data using t-SNE. *Journal of

Machine Learning Research*, *9*, 2579–2605.

von Philipsborn, A. C., Liu, T., Yu, J. Y., Masser, C., Bidaye, S. S., & Dickson, B. J.

(2011a). Neuronal control of Drosophila courtship song. *Neuron*, *69*(3), 509–522.

von Philipsborn, A. C., Liu, T., Yu, J. Y., Masser, C., Bidaye, S. S., & Dickson, B. J.

(2011b). Neuronal control of Drosophila courtship song. *Neuron*, *69*(3), 509–22.

http://doi.org/10.1016/j.neuron.2011.01.011

von Reyn, C. R., Breads, P., Peek, M. Y., Zheng, G. Z., Williamson, W. R., Yee, A. L.,

… Card, G. M. (2014). A spike-timing mechanism for action selection. *Nature*

Neuroscience, *17*(7), 962–970.

Zorović, M., & Hedwig, B. (2011). Processing of species-specific auditory patterns in the

cricket brain by ascending, local, and descending neurons during standing and

walking. *Journal of Neurophysiology*, *105*(5), 2181–94.

http://doi.org/10.1152/jn.00416.2010

Figures and Figure Legends

Figure 1.

Figure 1. Descending neuron phenotyping pipeline and behavior space. (A) The red light

activation rig. Six no retinal control flies and six retinal fed experimental flies were

mounted in parallel in individual 3 cm diameter plexiglass bubbles on top of three custom

light boards with constant 850 nm infrared light and variable 617 nm red light. The red

LEDs were repeatedly turned off and on for 45 sec and 15 sec, respectively. Each fly was

filmen at 150x150 pixel, 100 fps resolution by a single camera. Video data was then

aligned and processed, and the line was characterized for its occupancy in the descending

neuron behavior space with respect to red light activation and controls. (B) A 2D

representation of behaviors in the descending neuron video dataset was generated by

applying a probability density function to all the embedded data points (scale bar), which

was then convolved with a Gaussian (σ = 1.5) (C) Localization of various behaviors

within the descending neuron behavior space seen in (B), based on human curation of

watershedded regions in the space (Supplemental Figure 2, Supplemental movies S1-S6).

Figure 2.

Figure 2. Analysis of the head grooming DNg07 & DNg08 line (SS02635) and the

transient fast-running DNg25 line (SS01602). (A, D) Digital entropy of the distribution of

the retinal fed experimental flies (red) and non-retinal fed control flies (blue) in the

behavior space relative to the timing of red light stimulus onset (red light turned on at t =

0 sec, the time of red light activation is indicated by dashed lines). Experimental flies

experience a decrease in entropy when they perform a specific set of behaviors, because

they shift from the full range of normal fly behaviors to a subset of red light-activated

behaviors. (B, E) Average density +/- the standard deviation in the head grooming map

region indicated in red (inset, upper right) in experimental flies (red) and controls (blue)

relative to red light activation. The head grooming region was calculated as the region in

the map that experienced a statistically significant shift in density in experimental flies

but not controls when comparing the first 3 seconds (green bar) of the activation period to

the last 15 seconds of the recovery period (Wilcoxon rank sum test, p < 0.05, using the

Dunn–Šidák correction for multiple hypotheses). (C, F) Average density in the map over

a series of 3 second windows (calculated from 6 animals, 30 trials each). Red and blue

indicate regions of high and low density, respectively.

Figure 3.

Figure 3. Entropy for all of of descending neuron split-GAL4 lines. Trials were aligned

such that the onset of red light activation occured at t = 0 sec (A, B). The red light was

turned off at t = 15 sec. (A) Baseline subtracted entropy (Y-axis) versus time (X-axis) for

all experimental animals and all trials. Colors indicate a probability distribution

describing the entropy of experimental animals at a given time in the trial. (B) Entropy

levels of experimental animals over the course of the aligned trials (X-axis) shown by

line (Y-axis). Warm and blue colors indicate high and low entropy, respectively.

Figure 4

Figure 4. The timing of density shifts in descending neuron lines that occupy multiple

behavior space regions. (A) Most lines (86) increase density in only one region of the

behavior space upon red light activation. However, some lines occupy multiple regions in

the behavior space (B-D) Examples of lines that occupy multiple discontinuous regions

upon red light activation. Time is indicated on the X-axis, with the red light turned on at

T = 0 sec and off at 15 sec. Change in density in the color coded regions in the

experimental animals above the controls is indicated on the Y-axis. (B) Line SS01540

targeting descending neuron DNp09; (C) SS02542 targeting descending neuron DNb01;

(D) SS01049 targeting descending neuron DNp10.

Figure 5. Mutual information between behaviors performed before and during red-light

activation. (A) Mutual information for experimental flies, calculated using the density of

the flies in the behavior space at t = -1.5 to -0.5 seconds prior to red light activation

versus density at t = 0 to1.0 seconds after red light activation. Y-axis indicates the mutual

information, X-axis is all the lines in order of most to least mutual information, error bars

indicate the standard deviation. Lines are color coded by the number of significant

regions produced by red light activation, the key is indicated in the upper right, line

SS02542 is indicated by an asterisk. (B) Mutual information of experimental flies (X-

axis) plotted against that of non-retinal fed control flies (Y-axis), with the same color

coding as (A). (C) Partial mutual information for SS02542, showing the density at t = -

1.5 to 0.5 that has mutual information with the different regions of the behavior space

after red light activation indicated in the key in the upper left portion of the figure.

Figure 6.

Figure 6. Averages of representative lines for individual descending neurons. Descending

neurons are organized according to the region of the behavior space that they activate

(Figure 1). (A) Example illustrating the averaging of two lines to produce an estimated

phenotype. Colors indicate the degree to which particular regions are represented in the

average. Red regions are highly represented, blue less so, and white not at all. (B)

Phenotypes for 53 of the 58 descending neurons in the collection, plus pIP10 (von

Philipsborn et al., 2011b). Some descending neurons are represented by a single clean

line, others by averaging multiple clean lines. For averages, the number of lines are

indicated under the descending neuron name.

Supplementary Information

Figure S1.

Figure S1. Postural eigenmodes used to build the descending neuron behavior space

(Figure 1B).

Figure S2.

Figure S2. Watershedded regions in the descending neuron behavior space. (A) A

watershedding algorithm was used on the behavior space in Figure 1B to identify local

density maxima, which represent stereotyped behaviors. (B) Numbered watershedded

regions. These correspond to videos underlying all the watershedded regions (Movies

S1…S6) which were examined to create the human curated version of the behavior space

in Figure 1C.

Figure   S3.

Figure S3. Fraction of video data points for each movie embedded in the behavior space,

during the red light stimulus window (Y axis) and during the recovery period (X axis).

For most movies, between $90 - 95\%$ of frames embedded, with very little difference

between when the red light is on and when the red light is off, indicating that

CsChrimson and red light induced behaviors are well represented in the behavior space.

Failure to embed is usually indicative of imaging flaws, for example when part of the fly

is out of the field of view of the camera.

Figure S4.

Figure S4. Density in experimental (red) and control (blue) animals in the regions defined

in Figure 2A and 2D (A and B, this figure, respectively). Densities shown in Fig. 2B and

2E were averaged for six animals over 30 minutes. Gray bars indicate periods of red light

activation.

901
902   Figure S5. Analysis of all the descending neuron split-GAL4 lines. The line by line

903   analysis shown here follows the model of the selected examples described in Figure 2 of

904   the main text.

906   *Available at: http://www.biology.emory.edu/Berman/files/FigureS5_Cande_et_al.pdf*

907   (Large file size)

Figure S6. Choosing the red light intensity levels. (A) While most lines were screened at 5 mW/cm[2], 49 lines that had no or only a weak phenotype at this light level were repeated at 9 mW/[cm]2 and a higher retinal concentration. 43 of these lines acquired a phenotype or a strengthened statistical significance of the previously observed phenotype under these conditions. No lines lost their phenotype at the higher intensity. (B) 10 transient locomotion lines, overlaid in the same behavior space in different colors, were repeated at the lower and higher light intensities and retinal concentrations. The boundaries of the significant regions in the behavior space shift slightly between treatments, but overall the phenotypes remain largely unchanged.