Mutation accumulation in selfing populations under fluctuating selection

Authors: Eddie K.H. Ho[*1] and Aneil F. Agrawal[1]

*Corresponding author (contact: eddie.ho@mail.utoronto.ca)

[1]Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks Street, M5S 3B2, Toronto, Ontario

Key words: mutation accumulation, selfing, fluctuating selection, stochastic simulation

Running head: Mut. accum. under fluctuating selection

Author contributions: EKHH performed the simulations and analyses. EKHH and AFA wrote the paper.

Acknowledgements: This work was supported by NSERC Discovery grant 2015-05387 to Aneil Agrawal. The authors have no conflict of interests to declare.

Abstract

Selfing species are prone to extinction, possibly because highly selfing populations can suffer

from a continuous accumulation of deleterious mutations, a process analogous to Muller's

ratchet in asexual populations. However, current theory provides little insight into which types of

genes are most likely to accumulate deleterious alleles and what environmental circumstances

may accelerate genomic degradation. Here we investigate temporal changes in the environment

that cause fluctuations in the strength of purifying selection. We simulate selfing populations

with genomes containing a mixture of loci experiencing constant selection and loci experiencing

selection that fluctuates in strength (but not direction). Even when both types of loci experience

the same average strength of selection, loci under fluctuating selection contribute

disproportionately more to deleterious mutation accumulation. Moreover, the presence of loci

experiencing fluctuating selection in the genome increases the deleterious fixation rate at loci

under constant selection; under most realistic scenarios this effect of linked selection can be

attributed to a reduction in N_e. Fluctuating selection is particularly injurious when selective

environments are strongly autocorrelated over time and when selection is concentrated into rare

bouts of strong selection. These results imply that loci under fluctuating selection are likely

important drivers of extinction in selfing species.

Introduction

Self-fertilization has evolved many times independently in plants and animals (Goodwillie *et al*. 2005, Jarne and Auld 2006). Despite frequent transitions from outcrossing to self-fertilization, highly selfing populations are relatively rare in nature and relegated at the tips of phylogenies (Stebbins 1957, Igic and Kohn 2006). Stebbins (1957) proposed that highly selfing species suffer from low rates of adaptation and are more prone to extinction; classically called the 'dead end' hypothesis. Population genetics theory provides insights into why selfing is a dead end. Selfing reduces the effective population size, N_e, which diminishes the efficacy of selection. Under neutrality, N_e is reduced two-fold in obligately selfing populations because gametes are not sampled independently (Pollak 1987). In addition, but perhaps more importantly, high homozygosity in selfing populations reduces the effectiveness of recombination and can further reduce N_e due to linked selection effects (Nordborg 1997, Roze 2016). Explicit contrasts of selfing and outcrossing species confirm lower rates of adaptation and higher rates of deleterious mutation accumulation in both theory (Glemin and Ronfort 2013; Kamran-Disfani and Agrawal 2014) and reality (Slotte *et al*. 2010, Arunkumar *et al*. 2014, Burgarella *et al*. 2015). Further, phylogenetic analyses infer higher extinction rates in self-compatible *Solanacaea* (Goldberg *et al*. 2009) relative to their self-compatible counterparts. The disadvantage resulting from very strong linked selection can maintain low levels of outcrossing in species considered strong selfers (Kamran-Disfani and Agrawal 2014).

While both reduced rates of adaptation and higher rates of deleterious mutation accumulation may contribute to extinction, our focus here is on better understanding the latter. Though existing theory makes it clear that deleterious mutations accumulate more readily in highly selfing populations, this literature does not aim to provide insight into the types of genes

most likely to degrade or the types of environmental circumstances that might hasten genomic

degradation. Here we investigate how loci that experience fluctuations in intensity (but not

direction) affect deleterious mutation accumulation in highly selfing populations. As described

below, fluctuations in selection intensity affect fixation probabilities in both outcrossing and

selfing species but we focus on the latter because deleterious mutation accumulation is, in

general, expected to be much more prevalent in selfers.

There is abundant evidence that selection is environmentally sensitive in natural and in

laboratory experiments (Leimu and Fischer 2008, Bergland *et al*. 2014, Latta *et al*. 2015, Rutter

et al. 2017). Conceptually, selection can vary in direction and magnitude between environments

(Martin and Lenormand 2006), but empirically, selection varies mostly in magnitude. For

example, 'stressful' environments may alleviate or intensify the strength of selection, but rarely

reverses the direction of selection (Kishony and Leibler 2003, Agrawal and Whitlock 2010,

Kraemer *et al*. 2015). Hillenmeyer *et al*. (2008) found that 97% of all viable singe gene

knockouts in yeast reduce fitness in at least one environment while having little effect in other

environments after performing over 1000 chemical assays. This suggests that selection against

mutations can fluctuate in magnitude from neutrality to deleterious between environments (i.e.,

conditionally neutral mutations may be common). Other types of studies also indicate that

temporal changes in the environment create fluctuations in the strength of selection (Grant and

Grant 2002, Bell 2010, Siepielski *et al*. 2017). Despite this evidence, most mutation

accumulation models assume selection is constant. This likely underestimates deleterious

fixation rates if selection fluctuates temporally (Wardlaw and Agrawal 2012, Cvijovic *et al*.

2015).

Though classic single locus models for the fixation probability assume selection is

constant (Haldane 1927, Kimura 1962), fixation probabilities are known to be highly sensitive to

selection changing over time (Kimura and Ohta 1970, Gillespie 1978, Peischl and Kirkpatrick

2012). Often neglected are changes in selection that are stochastic and temporally autocorrelated,

especially for mutations that are on average deleterious. Environments can fluctuate over many

different time scales under various degrees of temporal autocorrelation (Halley 1996, Vasseur

and Yodviz 2004, Furguson *et al*. 2016) and evolutionary processes can be strongly dependent

on both the temporal autocorrelation in selection and the average strength of selection

(Charlesworth 1993, Pieschl and Kirkpatrick 2012, Wardlaw and Agrawal 2012). In general, if

changes in selection occur rarely (very high temporal autocorrelation), then the fixation

probability for the mutation is largely dependent on the selective environment in which it arose

and can be estimated from models assuming constant selection. If fluctuations are very frequent

(very low temporal autocorrelation), fixation probabilities can also be approximated by constant

selection models but using a time-averaged strength in selection. However, in the intermediate

range where selection and environmental change operate on similar time scales, classic equations

become inaccurate. Within this range, Cvijovic *et al*. (2015) have shown in single locus models

that selection fluctuating in direction increases the probability of fixation for all mutations.

We examine selection that is always negative (or zero) but fluctuates in intensity (but not

direction) in a system with strong linked selection due to selfing. Interference between

deleterious mutations leads to mutation accumulation in asexual and highly selfing populations

(Muller 1932, Haigh 1978, Charlesworth and Charlesworth 1993, 1997, Kamran-Disfani and

Agrawal 2014). However, little is known about selective interference when some loci experience

fluctuations in selection intensity. In a related study, Wardlaw and Agrawal (2012) found that

temporal autocorrelation in selection intensity is an important determinant of Muller's ratchet in

asexuals because the ratchet can turn quickly during periods of relaxed selection. The current

work extends their findings to selfing populations with some recombination, explores more

realistic environmental scenarios, separately quantifies fixation rates for loci experiencing

fluctuating versus constant selection within the same genome, examines the extent to which the

linked selection effects from fluctuations can be accounted for through N_e, and considers

selective interference involving beneficial mutations.

Our goal here is to investigate mutation accumulation in selfers under different models of

selection (rather than to contrast selfers and outcrossers). We find that genomes containing a

larger proportion of loci under fluctuating selection experience higher fixation rates genome

wide, both because of higher fixation rates at the fluctuating loci themselves but also through a

linked selection effect on other loci, including those under constant selection. These results

imply that loci under fluctuating selection contribute disproportionately to mutation

accumulation and fitness decline.

Methods

In most respects, the simulation follows standard assumptions so we only describe the important

features here but provide a detailed description in Supplementary File Part 2 (simulation code

available from https://github.com/EddieKHHo/SelfingMutAccum). The population consists of N

diploid hermaphrodites. Each genome consists of 5000 bi-allelic loci that affect fitness as well as

three neutral quantitative loci that are used to estimate effective population size, N_e. Loci are

arranged on a single linear chromosome that experiences, on average, $M = 1$ recombination event

per gamete. Fitness acts multiplicatively across loci and we assume deleterious alleles are

partially recessive, $h = \frac{1}{4}$. Reproduction occurs following standard assumptions assuming a

sporophitic selfing of $S = 0.98$. Each offspring receives, on average, U new deleterious

mutations. We confirmed the program produces the expected population genetic results under

constant selection (detailed in Supplementary File Part 3).

Base model

Our "base model" assumes individuals carry two types of loci that affect fitness. "C-loci"

experience a constant selection coefficient of s_c in all generations. "F-loci" experience a strength

of selection that changes over time, $s_f(t)$, taking the value 0 or s_{max}, i.e., mutations at F-loci are

conditionally neutral. A proportion, p_f, of the fitness affecting loci in the genome are F-loci and

the remainder are C-loci; location assignment of locus types within the genome was randomized

every simulation run. All F-loci experience the same selection in a given generation (0 or s_{max}) as

if they all are affected by the same environmental factor. The parameter f controls the temporal

autocorrelation in selection. With probability f selection remains the same as the previous

generation. With probability, $1-f$, $s_f(t)$ is randomly chosen to be 0 or s_{max} with probability $1-\phi$

and ϕ, respectively; ϕ is the expected proportion of generations that $s_f(t) = s_{max}$. The geometric

average strength of selection experienced by loci under fluctuating selection is $\bar{s} = 1-(1-s_{max})^{\phi}$.

For all simulations we chose s_{max} such that $\bar{s} = s_c$, so that F-loci are expected to experience the

same geometric average selection strength as C-loci to provide the fairest comparison between

these types of loci.

After a burn-in period, we calculated the per locus fixation rates for C- and F-loci

separately as the number of fixations at each type of loci divided by the number of generations

over which the data was collected and divided by the number of loci for each type. Throughout,

we calculated the *relative fixation rate* as the fixation rate in populations with $p_f > 0$ divided by

the fixation rate in populations with $p_f = 0$ (holding all other parameters equal). The relative

fixation rate for C-loci quantifies the extent to which fixation rates are affected by residing in

genomes that experience fluctuating selection rather than residing in genomes that only

experience constant selection (i.e., the classic model). The relative fixation rate for F-loci

quantifies the direct effect of fluctuating selection as well as effect of residing in a genome

experiencing fluctuating selection.

Simulations with exponentially distributed selection

We modified the base model by allowing selection strength to vary over time following an

exponential distribution (as opposed to the discrete "on/off" version of conditional neutrality

assumed in the base model). Each generation, consecutive environments remain the same with

probability f. With probability, 1-f, the selection strength is chosen from an exponential

distribution with mean $1/\bar{s}$. Similar to the base model, C- and F-loci experienced the $\bar{s}$. We

focused on simulations with $\bar{s} = 0.025$ where the geometric mean and the variance in selection

strength under the exponential model was almost identical to that of the conditionally neutral

mutations with $\bar{s} = 0.025$ and $\phi = 0.5$ in the base model.

Simulations with two fluctuating ecological variables

An unrealistic assumption of the base model is that all F-loci experience fluctuating selection in

exactly the same way. As the simplest way to relax this assumption, we allowed for two sets (F1,

F2) of loci under fluctuating selection and we imagine that each set responds to a different

ecological variable. These sets comprise p_{f1} and p_{f2} of the genome and the remaining proportion,

1-p_{f1}-p_{f2}, are C-loci. We assumed equally sized sets of fluctuating loci, p_{f1} = p_{f2} (and p_{f1} + p_{f2} =

p_f).

If mutations at F-loci are assumed to be conditionally neutral (as in the base model), both

F-loci sets experience fluctuations in selection between 0 and s_{max} with the same $\bar{s}$. Each

generation, the environment remains in the same state with probability f. With probability, 1-f,

the environment is randomly selected among four possible states: $\{s_{f1}(t) = s_{max}, s_{f2}(t) = s_{max}\}$,

$\{s_{f1}(t) = s_{max}, s_{f2}(t) = 0\}$, $\{s_{f1}(t) = 0, s_{f2}(t) = s_{max}\}$, $\{s_{f1}(t) = 0, s_{f2}(t) = 0\}$ with probabilities ϕ^2+d,

$\phi(1$-$\phi)$-d, $(1$- $\phi)\phi$-d, $(1$- $\phi)(1$-$\phi)$+d, respectively. The value of d determines correlation ρ in

selective states between sets F1 and F2. We also examined simulations with sets F1 and F2 in

which selection fluctuates over time following an exponential distribution; selection between sets

F1 and F2 was made to be either completely correlated, independent or negatively correlated (ρ =

1, 0, -0.645, respectively; detailed in Supplementary file Part 2).

Simulations with beneficial mutations

Lastly, we extended the base model by allowing beneficial mutations to occur at C-loci with a

genome wide rate of U_b, assuming U_b << U. We estimated the fixation probability for a new

beneficial mutation as the number of beneficial fixations divided by the expected number of

beneficials that were introduced (i.e., N * U_b * number of generations).

<u>Results</u>

Heuristic model

We first outline predictions for the fixation probability of a new deleterious mutation occurring

at a locus experiencing temporal fluctuations in the intensity of negative selection, denoted as P_f.

These predictions are useful for interpreting the results of our multi-locus simulations. If

selection fluctuates temporally between two selection coefficients $s = \{0, s_{max}\}$ with equal

frequency ($\phi = 0.5$), we can make simple predictions for P_f in the limiting cases where selection

fluctuates very often or very rarely. In the first case, selection fluctuates so often that a new

mutation will experience many periods of neutrality and strong selection before fixation and thus

experiences its expected (geometric) average fitness. We predict that P_f would be similar to the

fixation probability under time constant selection with $s = \bar{s} = 1-(1-s_{max})^{0.5}$; we denote this

prediction as $P_{\bar{s}}$. In the second case, selection fluctuates so rarely that a new mutation is almost

certain to be lost or fixed before selection changes. The expected fixation probability is then $\bar{P} =$

$(0.5) \, 1/(2N) + (0.5) \, P_{s_{max}}$, where $1/(2N)$ is the fixation probability for a neutral mutation ($s = 0$)

and $P_{s_{max}}$ is the fixation probability when $s = s_{max}$.

Given these two limiting cases, we make two predictions regarding P_f. Because the

fixation probability for a new deleterious mutation is a convex function of s (Fig. S1a; Kimura

1962), it follows that $\bar{P} > P_{\bar{s}}$ (via Jensen's inequality) and we predict P_f to lie within the

boundaries represented by the limiting cases of $\bar{P}$ and $P_{\bar{s}}$ (i.e., $\bar{P} \geq P_f \geq P_{\bar{s}}$; Fig. S1b, c). Second,

as $\bar{s}$ increases, $P_{\bar{s}}$ decays quickly and asymptotes at 0 while $\bar{P}$ decays more slowly and

asymptotes at $1/(4N)$ (Fig. S1b) so there is a larger proportional change in $P_{\bar{s}}$ than $\bar{P}$ with respect

to changes in $\bar{s}$ (Fig. S1e). This means that $P_{\bar{s}}$ is more sensitive to changes in s_{max} than $\bar{P}$; we

predict P_f to be intermediate in its sensitivity to changes in $\bar{s}$ (Fig. S1b, e). We see a similar

pattern when changing the population size, N. If we hold $\bar{s}$ constant, we observe that $P_{\bar{s}}$ is more

sensitive to changes in N compared to P_f and $\bar{P}$ (Fig. S1c, d, f).

The heuristic arguments above are based on single locus theory. Below we investigate

mutation accumulation in multi-locus models of a highly selfing species where selective

interference plays an important role in elevating fixation rates; nonetheless, we show that the

main ideas of the heuristic model apply. Within the context of our multi-locus simulations,

comparing the deleterious fixation rates of loci experiencing constant selection (C-loci) and loci

experiencing fluctuation selection (F-loci) is similar to comparing $P_{\bar{s}}$ and P_f, respectively

(assuming both loci experience the same $\bar{s}$). Thus, we predict fixation rates at F-loci to be higher

than that of C-loci (because $P_f > P_{\bar{s}}$). We also predict fixation rates at F-loci to be less sensitive

to changes in $\bar{s}$ and N_e than fixation rates at C-loci.

Temporal autocorrelaition in selection

Our heuristic model implies that the fixation probability of mutations should depend on

the temporal autocorrelation of selection, f, (i.e., $P_f \rightarrow \bar{P}$ as $f \rightarrow 1$ and $P_f \rightarrow P_{\bar{s}}$ as $f \rightarrow 0$). We

confirmed this in our multi-locus model of highly selfing populations (Fig. S3, S4); Wardlaw and

Agrawal (2012) found similar results for the deleterious fixation rate in obligately asexual

populations. Rather than further exploring the effects of f on fixation rates, we assume that

selection undergoes moderate degrees of temporal autocorrelation ($f = 0.95$, $\phi = 0.5$) such that

the correlation in selection between time points separated by 1, 10, and 100 generations is 0.95,

0.60, and 0.006, and the average run length between switches in the environment is ~40

generations. This regime elevates fixation rates at loci under fluctuating loci, but is moderate in

its effects; fixation rates can be much higher with higher values of f and lower values of ϕ (Fig.

S3, S4, S15). In the remaining sections, we focus on exploring how loci under fluctuating

selection in the genome alter fixation rates at linked sites and genome wide.

Varying the composition of the genome, p_f

We examine fixation rates in genomes where a fraction p_f of loci are under fluctuating selection

(F-loci) and the remaining loci are under constant selection (C-loci). Unless otherwise noted, we

focus on simulations of highly selfing populations ($S = 0.98$) with $N = 10^4$, $\bar{s} = 0.025$, and $h =$

0.25 with $f = 0.95$ and $\phi = 0.5$; results for obligately selfing populations ($S = 1$) can be found in

Supplementary File Part 1. Per locus fixation rates within genomes experiencing fluctuating

selection ($p_f > 0$) are typically higher than fixation rates in the classic model ($p_f = 0$); this is true

for both C- and F-loci, but the effect is stronger for the latter (Fig. 1) The fact that per locus

fixation rates vary with p_f indicates that changing the density of F-loci in the genome alters the

effects of linked selection on both F- and C-loci. When less than half the genome consists of F-

loci ($p_f < 0.5$), increasing p_f generally elevates fixation rates at both C- and F-loci and, thereby,

the genome-wide average fixation rate. At high mutation rates ($U = 1$) further increasing p_f

reduces fixation rates for genomes that contain more than 50% F-loci (Fig. 1a). As it seems

unlikely that a high fraction of loci experience strong fluctuations in selection, the results for $p_f <$

0.5 are the most biologically relevant.

When mutation rates are lower (so absolute fixation rates are lower), the effect of

fluctuating selection on the *relative* fixation rate is more dramatic for both C- and F-loci. For the

parameter sets examined in Fig. 1, the inclusion of F-loci maximally increases genome-wide

fixation rate by a factor of 1.38 vs. 2.42 at the high and low mutation rate, respectively. This

effect of mutation rate holds in simulations using other values of autocorrelation f and frequency

of selection ϕ and in obligately selfing populations (Fig. S4, S5). Although the relative impact of

fluctuating selection is larger at lower mutation rates, in the remaining sections we focus on

results from high mutation rates ($U = 1$) because simulations at lower mutation rates require

much longer runs to obtain accurate estimates of fixation rates.

Figure 1. Relative fixation rates (mean ± SE) for C- (blue) and F-loci (red) in highly selfing
populations with varying genomic composition, p_f. Fixation rates under high mutation rates in (a)
$U = 1$ and low mutation rates (b) $U = 0.5$. Fixation rates are given relative to the fixation rate for
C-loci in the classic model ($p_f = 0$), which is represented by the horizontal dashed line at 1.0 (the
grey shading denotes SE). The absolute fixation rates in the classic model ($p_f = 0$) for panels a,b
are 7.47×10^{-6} and 8.74×10^{-7} per locus per generation respectively. Other parameters: $N = 10^4$,
$S = 0.98$, $f = 0.95$, $\phi = 0.5$, $h = 0.25$, $\bar{s} = 0.025$.

The density of F-loci alters the effective population size, N_e (Fig. 2a shows N_e for the

simulations represented by Fig. 1; Fig. S6 for obligately selfing populations). This might be the

reason for the variation in fixation rates in relation to p_f (Fig. 1a). If so, fixation rates should be

equal for simulations with different values of p_f if they experienced the same N_e. To test this, we

ran additional simulations changing the census population size, N, of the simulations to raise or

lower N_e for a given value of p_f. (We observe that N_e is not linearly related to N (Fig. 2b) as has

been observed previously in simulations of asexual populations accumulating deleterious

mutations due to Muller's ratchet (Gordo *et al.* 2002).) Consistent with the idea that the

relationship between p_f and fixation rate results from the effect of p_f on N_e, fixation rates are

similar across different values of p_f when N_e is similar, but only if p_f is not too high ($0 < p_f \leq 0.5$;

Figs. 2c, d). However, when comparing simulations with similar estimates of N_e, very high

values of p_f (i.e., $p_f = 0.8, 0.95$) result in higher rates fixation rates than low or moderate values

of p_f (i.e., $p_f \leq 0.5$). This indicates that when p_f is large (and the mutation rate is high), selective

interference between C- and F-loci affects fixation rates in ways that cannot be captured by a

genome-wide estimate of N_e.

Figure 2. (a) Estimates of effective population size (mean ± SE) for genomes containing
different densities of F-loci, p_f, in a population of $N = 10^4$. Shown are N_e values for highly selfing
($S = 0.98$) populations under high ($U = 1$) and lower ($U = 0.5$) mutation rates; these results are
from the same simulations represented in Figs. 1a,b. The horizontal dashed line indicates the N_e
(grey shading is SE) for the corresponding classic model ($p_f = 0$). (b) Estimates of N_e plotted
against N ($U = 1$) for genomes with different values of p_f as depicted in the legend. The open
symbol indicates the N_e when $N = 10^4$ (i.e., corresponding to simulations in Fig. 1a). Fixation
rates plotted against N_e (c) loci under constant selection (d) loci under fluctuating selection ($U =$
1).

Fixation bias at F-loci

As predicted from the heuristic model, the fixation rate of deleterious mutations at F-loci is

always greater than that for C-loci within all mixed genomes (Figs. 1, 2c, d). This bias is clearly

stronger when selection fluctuates in ways that directly increase fixation rates at F-loci (i.e.,

larger f and smaller ϕ; Figs. 3a, S4). Moreover, this bias is affected by the 'traditional'

parameters affecting mutation accumulation such as the strength of selection ($\bar{s}$), the mutation

rate (U) and the population size (N). To examine how this bias depends on aspects of selection,

we performed simulations where half the genome consist of F-loci ($p_f = 0.5$) with an average

selection strength of $s_c = \bar{s} = \{0.005, 0.025, 0.05\}$. The bias in fixations at F-loci increases with $\bar{s}$

as stronger selection reduces fixation rates at C-loci relatively more than at F-loci (Fig. 3b).

These results can be interpreted in light of the heuristic model that predicts F-loci will be less

sensitive than C-loci to increases in $\bar{s}$: mutations at F-loci can accumulate during episodes where

selection is weak regardless of how large s_{max} becomes in episodes when selection is strong.

Overall, conditions that slow down mutation accumulation, such as stronger $\bar{s}$ (Fig. 3), lower U

(Figs. 4, S4, S14b), larger N (Fig. S9), and less selfing (Fig. S10) reduce fixation rates at C-loci

more so than at F-loci, increasing the relative contribution of F-loci to total mutation

accumulation; see Fig. S8 for simulations of obligately selfing populations.

Figure 3. (a) Ratio of fixation rates at F-loci to fixation rates at C-loci when half of the genome

consists of F-loci (p_f = 0.5); error bars are SE (often too small to be visible). When ϕ = 0.5 (solid

black line), selection is non-neutral (s_{max}) half the time but when ϕ = 0.25 (double-dashed grey

line), selection is non-neutral only a quarter of the time. To maintain $\bar{s}$ = {0.005, 0.025, 0.05}

when ϕ is reduced by half we approximately double the value of s_{max} (Methods), therefore

making selection concentrated into shorter but more intense events. (b) Fixation rates at F-loci

(red) and C-loci (blue) when ϕ = 0.5 (solid line) and when ϕ = 0.25 (double-dashed line). Other

parameters: S = 0.98, N = 10^4, U = 1, h = 0.25, f = 0.95.

Figure 4. Ratio of fixation rates (mean ± SE) at F-loci to fixation rates at C-loci in populations

with different proportions of F-loci under high (U = 1, circles) and low (U = 0.5, triangles)

mutation rates. Other parameters: S = 0.98, N = 10^4, h = 0.25, f = 0.95, ϕ = 0.5.

Exponentially distributed selection over time

In the base model we assumed that mutations at F-loci are conditionally neutral with selection

fluctuating discretely between 0 and s_{max}. As a test of whether the patterns above are robust to

that assumption, we consider simulations where the selection strength at F-loci is randomly

sampled from an exponential distribution when the environment changes. Both models have a

similar mean and standard deviation in selection strength ($\bar{s} \cong 0.025$, standard deviation in $s_f(t)$ =

0.025). Using f = 0.95 in the exponential model (as in the base model), the correlation in

selection between time points separated by 1, 10, and 100 generations is 0.95, 0.60, and ~0,

respectively. The fixation rates for both C- and F-loci are slightly lower under the exponential

model than the discrete conditional neutrality model (Figs. 5, S12). Mutations fix

disproportionately at F-loci under the exponential selection model, but this bias is not quite as

strong as observed in the conditional neutrality model. As in the base model, reducing U (from U

= 1 to $U = 0.5$) under the exponential model increases relative fixation rates and the proportion

of mutations that fix at F-loci (Figs. 5, S14). Though the base model and exponential model

represent qualitatively different forms of temporal heterogeneity in selection, the effects of these

models on fixation rates are reasonably similar. We speculate that any form of fluctuations in

selection intensity are likely to elevate fixation rates and fitness decline in highly selfing

populations as long as the variance in selection over time is sufficiently large and there is some

degree of temporal autocorrelation.

Figure 5. Relative fixation rates (mean ± SE) at C- (blue) and F-loci (red) in populations with

different genomic compositions, p_f, when selection strength is modelled as discrete conditional

neutrality (solid line) or following an exponential distribution (double-dashed line) over time

under (a) high mutation rates ($U = 1$) and (b) low mutation rates ($U = 0.5$). Relative fixation rates

under the conditional neutrality model are re-plotted from Figs. 1a,b. The geometric time

average and the variance in selection strength and for the conditional neutrality and exponential

selection model are approximately the same ($\bar{s} = 0.025$, standard deviation in $s_f(t) = 0.025$).

Fixation rates are given relative to the fixation rate for C-loci in the classic model ($p_f = 0$), which

is represented by the horizontal dashed line at 1.0 (the grey shading denotes SE). Other

parameters: $S = 0.98$, $f = 0.95$, $N = 10^4$, $h = 0.25$.

Two components of fluctuating selection

In the previous sections, we assumed that all F-loci simultaneously respond to the same

environmental factor that fluctuates in time. As the simplest way to relax this assumption, we

allow for two equally sized sets of F-loci (F1, F2) that respond to two separate environmental

factors. The two environmental factors may fluctuate in perfect synchrony, causing a complete

correlation in selective states between loci within F1 and F2 ($\rho = 1$), which is equivalent to the

base model. Alternatively, the two environment factors fluctuate in anti-synchrony, causing a

negative correlation in selective states between F1 and F2 ($\rho = -1$). We perform simulations for a

range of different values of ρ, the correlation in selective states between set F1 and F2.

The environmental correlation ρ has little effect on the fixation rates of C- and F-loci

when the genome consists of mostly C-loci ($p_f \leq 0.5$); ρ is negatively related to fixation rate

when the genome consists of mostly F-loci, making our base model ($\rho = 1$) conservative in this

respect (Fig. 6a); obligately selfing populations are less insensitive to changes in ρ (Fig. S12).

We observe qualitatively similar effects of ρ when selection between the two groups is

exponentially distributed in time (Figs. 6b, S12, S13). In sum, under biologically realistic ranges

of p_f (i.e., $p_f < 0.5$), fixation rates do not appear to be strongly dependent on how fluctuations in

selection are correlated between two equally sized sets of F-loci.

Figure 6. Relative fixation rates (mean ± SE) for C-loci (blue) and F-loci (red) as a function of the correlation in environmental variables responsible for fluctuating selection ρ. Loci under fluctuating selection are equally distributed between two sets of F-loci (F1, F2). The correlation in selective states between F1 and F2, ρ, can be positive or negative or zero. The F-loci (summing both F1 and F2) comprise a fraction p_f of the genome. (a) Selection under the model of discrete conditional neutrality; ρ = {1, 0.5, 0, -0.5, 1}. (b) Selection is exponentially distributed over time; ρ = {1, 0, -0.645}. Fixation rates are shown for genomes with different frequencies of fluctuating loci, p_f. Fixation rates are given relative to the fixation rate for C-loci in the classic model (p_f = 0), which is represented by the horizontal dashed line at 1.0 (the grey shading denotes SE); the actual fixation rates is 7.47 x 10^{-6}. Other parameters: S = 0.98, f = 0.95, ϕ = 0.5, $\bar{s}$ = 0.025, N = 10^4, U = 1, h = 0.25.

Beneficial mutations

We performed simulations in which beneficial mutations occurred at C-loci at much lower rates than deleterious mutations ($U_b \leq 0.001$) and had smaller, larger or equal fitness effects compared to deleterious mutations (beneficials: s_b = {0.01, 0.025, 0.05, 0.08}, h_b = 0.5; deleterious: $\bar{s}$ = 0.025, h = 0.025). F-loci only received deleterious mutations and experienced selection fluctuating discretely over time between 0 and s_{max} (f = 0.95, ϕ = 0.5).

The fixation probability of beneficial mutations was lower in simulations with F-loci ($p_f =$

0.2) than those without F-loci ($p_f = 0$) for all values of s_b (Table 1) indicating that deleterious

mutations at F-loci caused more interference with beneficials than those at C-loci. Beneficials

with smaller or similar effect size to deleterious mutations (i.e. $s_b = 0.01, 0.025$) had little effect

on the deleterious fixation rate. However, beneficials with larger fitness effects ($s_b = 0.05, 0.08$)

increased deleterious fixation rates at both C- and F-loci (Table 1). The deleterious fixation rate

of C-loci changed proportionately more with increases in the strength of beneficials (s_b)

compared to that of F-loci (Table 1), consistent with our heuristic model. Recall that comparing

fixation rates at F-loci and C-loci is similar to comparing P_f and $P_{\bar{s}}$, respectively, and $P_{\bar{s}}$ is

proportionally more sensitive to changes in N and s compared to P_f (Fig. S1e, f). Because an

increase in s_b causes a decrease in N_e (Table 1), we expect C-loci ($P_{\bar{s}}$) to be more sensitive to this

change than F-loci (P_f).

Because the F-locus rate without beneficials is larger than the C-locus rate, the fact that

the absolute increase in the deleterious fixation rate caused by beneficials is larger for F-loci than

C-loci is obscured by only looking at their "proportional increase". This is clearer by taking a

second perspective in which we calculate the average number of additional fixed deleterious

mutations per fixed beneficial (Table S3). When the beneficial effect size is larger than

deleterious ($s_b = 0.05, 0.08$), each beneficial that fixes is responsible for, on average, less than

one additional deleterious fixation but these are not evenly distributed between F- and C-loci. F-

loci are only 25% as abundant as C-loci (i.e., $p_f = 0.2$), but the number of additional fixed

deleterious F-alleles per fixed beneficial is more than 25% the number of additional fixed

deleterious C-alleles. From this perspective, F-loci experience stronger interference from

beneficials, in an absolute sense, than do C-loci even though C-loci experience a greater

proportional interference effect than F-loci (Table 1).

Table 1. Summary of simulations with beneficial mutations

U	U_b	s_b	p_f	N_e (SE)	Ben prob fix x 10^{-3} (SE)	Del fix rate relative to NO ben (SE) C-loci	F-loci
0.5	0.0001	0.01	0	99.7	0.172	0.992	
				(0.35)	(0.004)	(0.006)	
			0.2	96.5	0.164	0.999	1.032
				(0.36)	(0.005)	(0.005)	(0.013)
		0.025	0	99.7	0.752	1.021	
				(0.37)	(0.011)	(0.009)	
			0.2	96.9	0.703	1.019	1.072
				(0.48)	(0.012)	(0.007)	(0.018)
		0.05	0	96.6	3.798	1.312	
				(0.25)	(0.015)	(0.008)	
			0.2	93.5	3.499	1.240	1.159
				(0.33)	(0.025)	(0.008)	(0.024)
		0.08	0	87.5	11.429	2.373	
				(0.34)	(0.040)	(0.011)	
			0.2	85.9	10.631	2.092	1.664
				(0.34)	(0.044)	(0.012)	(0.022)
1	0.001	0.01	0	73.8	0.121	0.994	
				(0.33)	(0.002)	(0.003)	
			0.2	71.2	0.119	0.980	1.039
				(0.24)	(0.002)	(0.003)	(0.018)
		0.025	0	72.8	0.384	1.016	
				(0.34)	(0.003)	(0.003)	
			0.2	71.0	0.356	1.005	1.008
				(0.33)	(0.004)	(0.003)	(0.011)
		0.05	0	71.0	1.531	1.199	
				(0.31)	(0.007)	(0.004)	
			0.2	67.7	1.405	1.152	1.129
				(0.31)	(0.007)	(0.004)	(0.018)
		0.08	0	62.5	3.916	1.774	
				(0.29)	(0.010)	(0.003)	
			0.2	60.9	3.672	1.650	1.478
				(0.28)	(0.008)	(0.004)	(0.012)

U, U_b are the genomic wide mutation rates of deleterious and beneficial mutations, respectively. s_b is the fitness advantage of a beneficial mutation. 'Ben prob fix' is the fraction of beneficial mutations that fixed per generation in the simulation assuming that the number of beneficial mutations introduced per generation was $U_b * N$. 'Del fix rate relative to NO ben' is the fixation rate of C-loci and F-loci in simulations with beneficial mutations relative to the fixation rate of C-loci and F-loci in simulations without beneficial mutations (holding all other parameters constant). Other parameters: $S = 0.98$, $N = 10000$, $h = 0.25$, $\bar{s} = 0.025$, $h_b = 0.5$, $f = 0.95$, $\phi = 0.5$, $\rho = 1$. See Table S2, S3 for further information on these simulations.

Discussion

Because of the reduced efficacy of selection associated with high selfing rates, such populations

can accumulate and permanently fix deleterious mutations at high rates leading to rapid fitness

decline and possibly to extinction (Heller and Maynard Smith 1979, Charlesworth and

Charlesworth 1993). Environmental changes occur on many different times scales (Halley 1996,

Vasseur and Yodviz 2004, Furguson *et al*. 2016) and can induce fluctuations in the intensity and

direction of selection (Siepielski *et al*. 2017, Bell 2010) but the consequences for mutation

accumulation in selfers is unstudied. Loci experiencing selection that fluctuates in intensity, F-

loci, fix deleterious mutations at a higher rate than loci under constant selection, C-loci,

experiencing the same time averaged selection strength ($s_c = \bar{s}$). The extent to which F-loci

contribute disproportionately to total mutation accumulation depends on aspects of

environmental heterogeneity. The bias is stronger with increasing temporal autocorrelation in

selection, *f*, and when selection is concentrated into rarer but more intense episodes (low values

of ϕ), holding $\bar{s}$ constant (Fig. S3, S4, S14). This bias also becomes stronger with changes to

"traditional" parameters that lower the overall rate of mutation accumulation, i.e., higher values

of $\bar{s}$ or *N* as well as lower values of *U* (Fig. 3, 4, S8, S9). In sum, fluctuating selection can

increase the rate of mutation accumulation in highly selfing populations by several fold and the

loci under fluctuating selection contribute disproportionately to mutational decline. These

conclusions appear qualitatively robust to the exact nature of fluctuations (Fig. 5, 6, S12, S13).

In our model, deleterious mutations fix as a consequence of selective interference

between linked sites. Supporting this, simulations with the same parameter values but at higher

rates of outcrossing ($S \leq 0.9$) fix few to no mutations (Table S1). Obligately asexual organisms

(Wardlaw and Agrawal 2012) and obligate selfers ($S = 1$) have high rates of mutation

accumulation that are further increased by fluctuating selection. These effects remain important

even with low levels of outcrossing ($S = 0.98$) that permit some genetic mixing. Interference

between constantly selected loci can lead to the accumulation of deleterious mutations in asexual

and highly selfing populations, a process classically called Muller's ratchet (Haigh 1978, Heller

and Maynard Smith 1979). However, little attention has been given to understanding of selective

interference between loci under fluctuating selection and loci under constant selection.

In genomes containing a mix of F- and C-loci, F-loci have inherently higher rates of

fixation but also tend to increase fixation rates at C-loci and other F-loci as illustrated in Fig. 1

where increasing the fraction of F-loci in the genome increases the per-locus fixation rates at

both C- and F-loci (for $0 < p_f < 0.5$). This indicates that F-loci cause more selective interference

(with both C- and F-loci) than C-loci. Similarly, we found that deleterious mutations at F-loci

reduce fixation rates of beneficials more than deleterious mutations at C-loci do (Table 1, S2).

Previous work in asexual populations where selection is constant over time but varies among loci

shows that interference from strongly deleterious alleles increase fixation of more weakly

deleterious alleles (Gordo and Charlesworth 2001, Soderberg and Berg 2007). From this we can

infer that strong interference caused by F-loci occurs during periods when selection is strong,

$s_f(t) = s_{max} > s_c$. However, during periods where $s_f(t) = 0$, F-loci do not cause interference. Given

the observed net increase in fixation rate at C-loci, we infer that the increased interference when

selection at F-loci is strong ($s_f(t) = s_{max}$) outweighs the reduced interference when selection is

weak ($s_f(t) = 0$). On the other hand, we observed that strongly selected beneficials interfered

more strongly with deleterious alleles at C-loci than at F-loci in terms of the proportional change

in their fixation rates (Table 1). Viewing interference from the beneficials as a reduction in N_e,

the result above is consistent with the heuristic model prediction that fixation probabilities for F-

loci are less sensitive to N_e than for C-loci.

Indeed the effect of selective interference on fixation rates can be interpreted as resulting

from a reduction in the effective population size in some cases (Hill and Robertson 1966,

McVean and Charlesworth 2000, Gordo and Charlesworth 2002). We observed that per locus

fixation rates varied as a function of p_f for both C- and F-loci (Fig. 1). In the range where p_f was

not too large ($0 < p_f \leq 0.5$), fixation rates for each locus type (F and C) were nearly equal for

different values of p_f when we adjusted N to obtain similar N_e values (Fig. 2). Because neutral

diversity (used to infer N_e) and fixation rates are similarly affected by selective interference

within this range of p_f values, we conclude that selective interference in this regime can be

thought of as causing a reduction in N_e. Good *et al.* (2014) showed that in asexual populations

experiencing selective interference the variance in fitness explains neutral diversity much better

than the N_e expected under deterministic mutation-selection balance. However, the time-

averaged variance in fitness did not explain the variation in fixation rates in relation to p_f better

than our diversity-based estimates of N_e (Fig. S7).

Neither N_e nor the variance in fitness can account for the variation in fixation rates when

most loci undergo fluctuating selection ($p_f > 0.5$). Others have noted that the effects of selective

interference on fixation rates cannot always be captured through N_e. A sweep of a strongly

beneficial mutation differentially affects neutral diversity and fixation probability at a linked

weakly beneficial mutation (Stephan *et al.* 1992, Barton 1994). Fluctuating selection may have

some sweep-like properties because genotypes favoured during the selective period ($s_f(t) = s_{max}$)

can be different from those favoured during the neutral period ($s_f(t) = 0$) due to the shift in the

relative importance of F- and C-loci in determining fitness. Recurrent sweeps of highly fit

genotypes caused by changing selective environments may represent another dynamical process

that differentially affects fixation probability at selected loci and genetic drift at linked neutral

loci (Comeron *et al*. 2008).

There are several questions in assessing the relevance of our model to natural systems. As

highlighted in the Introduction, numerous lab studies show selection on mutations varies in

strength across the environmental conditions. However, we do not know the extent to which this

variation occurs in nature. Further, we have no estimate of the proportion of loci in the genome

that experiences fluctuating selection, p_f. Excluding mutations causing intrinsic lethality, we

speculate that almost every locus experiences temporal change in selection strength as it seems

implausible selection strength is exactly the same in all conditions. However, for many loci,

fluctuations in selection may be minor. Perhaps only a minority of mutations undergo

fluctuations in selection strength on the order of magnitude that we modeled and with at least a

moderate level of temporal autocorrelation.

Environmental changes that cause fluctuations in selection may also cause fluctuations in

population size (e.g., Bergland *et al*. 2014). Our main goal was to isolate the effects of

fluctuating selection on its own but population size changes may be important on their own

and/or interact with fluctuating selection. As described in Supplementary File Part 5, we briefly

explored the interaction between fluctuating N and s. As expected, simulations with only

fluctuating N fixed deleterious mutations at a rate similar to one with constant $N = N_h$, where N_h

represents the harmonic time average population size (Otto and Whitlock 1997). Fluctuations in s

alone increased deleterious fixation rates compared with one with constant $s = \bar{s}$. Keeping N_h

constant, the addition of fluctuating N into models of fluctuating s did not greatly change the

deleterious fixation rate of the latter regardless of the (positive and negative) correlations in the

fluctuations of N and s. In summary, fluctuations in N and s have different effects on deleterious

fixation rates; we cannot simply use fluctuations in the product Ns as a substitute for fluctuations

in each component alone.

Our study was motivated by the 'dead end' hypothesis for highly selfing species

(Stebbins 1957, Igic and Busch 2013) but we do not explicitly model extinction. Similar to

previous examinations of mutation accumulation in genomes with low rates of recombination

(Haigh 1978, Heller and Maynard Smith 1979, Charlesworth and Charlesworth 1997), we

hypothesize that deleterious fixations should lead to fitness loss and eventual extinction. Models

using realistic parameters to predict the expected time to extinction under Muller's ratchet

assume deleterious mutations directly reduce population growth rates (Gabriel *et al*. 1993,

Loewe 2006, Loewe and Lamatsch 2008). However, it is not obvious how the deleterious fitness

effects of mutations directly affect ecological success and species persistence (Agrawal and

Whitlock 2012). That said, our model predicts that mutations accumulate disproportionately at

loci experiencing fluctuating selection. We might expect that populations are driven to extinction

when deleterious mutations fix at loci required for surviving rare but severe climatic events.

Prior to extinction, the central prediction is that F-loci contribute disproportionately to mutation

accumulation. In reality, this would be challenging to test because it would be difficult to

categorize the genome into F- and C-loci. Even if one did so (perhaps based on plasticity in

expression), it would be extremely difficult to identify subsets of loci from each category that

have comparable average selection strengths.

We focused on the mutation accumulation in highly selfing populations but the dynamics

of deleterious mutations can affect the evolution of the selfing rate itself. In finite populations,

the build up of negative linkage disequilibrium (LD) at sites under negative selection can

increase deleterious fixation rates and favour the evolution of increased recombination and

outcrossing (Keightley and Otto 2006, Gordos and Campos 2008, Kamran-Disfani and Agrawal

2014). Kamran-Disfani and Agrawal (2014) found that selfing rates evolve to just below where

Muller's ratchet begins turning at high rates. Because the presence of F-loci in the genome

increases the deleterious fixation rate, it is natural to ask whether it also affects the evolution of

selfing. In Supplementary File Part 4, we briefly explored this by allowing selfing rates to evolve

in our model. We found that the equilibrium level of selfing was similar if F-loci are present ($p_f =$

0.25) or absent ($p_f = 0$), even though deleterious fixation rates are higher if F-loci are present.

554   Highly selfing species appear to be prone to extinction though we lack a detailed

understanding of how this occurs. Our work indicates fluctuations in the strength of purifying

selection can be an important contributor to this process. The results presented may

underestimate the relative importance given that we observe stronger effects under lower

mutation rates (Fig. 1, S13, S14). Even if rare in the genome, loci experiencing fluctuating

selection make a disproportionately larger contribution to mutation accumulation and fitness

decline than loci under constant selection. Moreover, mutation accumulation at such loci may

make selfing species particularly susceptible to extinction to rare environmental conditions

where the functions of such loci are particularly important. Although detecting and measuring

the fitness impact of loci under fluctuating selection will be empirically challenging, our results

demonstrate a plausible parameter space where fluctuating selection can strongly impact the

mutation accumulation in highly selfing species.

<u>References</u>

Agrawal, A.F. and Whitlock, M.C. 2010. Environmental duress and epistasis: how does stress

affect the strength of selection on new mutations? Trends Ecol. Evol. 25:450-458.

Agrawal, A.F., and Whitlock, M.C. 2012. Mutation load: the fitness of individuals in populations

where deleterious alleles are abundant. Annu. Rev. Ecol. Evol. Syst. 43: 115-135.

Arunkumar, R., Ness, R. W., Wright, S. I. and Barrett, S. C. H. 2015. The evolution of selfing is

accompanied by reduced efficacy of selection and purging of deleterious mutations. Genetics

199: 817-829.

Barton, N. H. 1994. The reduction in fixation probability caused by substitutions at linked loci.

Genet. Res. Camb. 64: 199-208.

Bell, G. 2010. Fluctuating selection: the perpetual renewal of adaptation in variable

environments. Phil. Trans. R. Soc. B. 365: 87-97.

Bergland, A.O., Behrman, E.L., O'Brien, K.R., Schmidt, P.S. and Petrov, D.A. 2014. Genomic

evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. PLoS

Genet. 10: e1004775.

Burgarella, C., Gayral, P., Ballenghien, M. *et al*. 2015. Molecular evolution of freshwater snails

with contrasting mating systems. Mol. Biol. Evol. 32: 2403-2416.

Charlesworth, D. Morgan, M.T. and Charlesworth, B. 1993. Mutation accumulation in finite

outbreeding and inbreeding populations. Genet. Res. Camb. 61: 39-56.

Charlesworth, B. 1993. Directional selection and evolution of sex and recombination. Genet.

Res. Camb. 61: 205-224.

Charlesworth, B. and Charlesworth, D. 1997. Rapid fixation of deleterious alleles can be caused

by Muller's ratchet. Genet. Res. Camb. 70: 63-73.

Comeron, J.M., Williford, A. and Kliman, R.M. 2008. The Hill-Robertson effect: evolutionary

consequences of weak selection and linkage in finite populations. Heredity 100: 19-31.

Cvijovic, I., Good, B.H., Jerison, E.R. and Desai, M.M. 2015. Fate of a mutation in a fluctuating

environment. P. Nat. Acad. Sci. USA. 112: E5021-E5028.

Ferguson, J.M., Carvalho, F., Murillo-Garcia, O., Taper, M.L. and Ponciano, J.M. 2016. An

updated perspective on the role of environmental autocorrelation in animal populations. Theor.

Ecol. 9: 129-148.

Gabriel, W., Lynch, M. and Burger, R. 1993. Muller's ratchet and mutational meltdowns.

Evolution 74: 1744-1757.

Gillespie, J.H. 1978. A general model to account for enzyme variation in natural populations. V.

the SAS-CFF model. Thoer. Pop. Biol. 14: 1-45.

Glemin, S., and Ronfort, J. 2013. Adaptation and maladaptation in selfing and outcrossing

species: new mutations versus standing variation. Evolution 67: 225-240.

Goldberg, E.E., Kohn, J.R., Lande, R., Robertson, K.A., Smith, S.A. and Igic, B. 2010. Species

selection maintains self-incompatibility. Science 330: 493-495.

Good, B.H., Walczak, A.M., Neher, R.A. and Desai, M.M. 2014. Genetic diversity in the

interference selection limit. PLoS Genet. 10: e1004222.

Goodwillie, C., Kalisz, S. and Eckert, C. G. 2005. The evolutionary enigma of mixed mating

systems in plants: occurrence, theoretical explanations, and empirical evidence. Annu. Rev. Ecol.

Evol. Syst. 36: 47-79.

Gordo, I. and Campos, P.R.A. 2008. Sex and deleterious mutations. Genetics 179: 621-626.

Gordo, I. and Charlesworth, B. 2001. The speed of Muller's ratchet with background selection

and the degeneration of Y chromosomes. Genet. Res. Camb. 78: 149-161.

Gordo, I., Navarro, A. and Charlesworth, B. 2002. Muller's ratchet and the pattern of variation at

a neutral locus. Genetics 161: 835-848.

Grant, P.R. and Grant, B.R. 2002. Unpredictable evolution in a 30-year study of Darwin's

finches. Science 296: 707-711.

Haigh, J. 1978. The accumulation of deleterious genes in a population-Muller's ratchet. Theor.

Popul. Biol. 14: 251-267.

Haldane, J.B.S. 1927. A mathematical theory of natural and artificial selection Part V: selection

and mutation. Proc. Camb. Philos. Soc. 23: 838-844.

Halley, J.M. 1996. Ecology, evolution and the 1/f-noise. Trends Ecol. Evol. 11: 33-37.

Heller, R. and Maynard Smith, J. 1979. Does Muller's ratchet work with selfing? Genet. Res.

Camb. 32: 289-293.

Hill, W. G. and Robertson, A. 1966. The effects of linkage on limits to artificial selection. Genet.

Res. Camb. 8: 269-294.

Hillenmeyer, M.E., Fung, E., Wildenhain, J., Pierce, S.E., Hoon, S., Lee, W., Proctor, M.,

St.Onge, R.P., Tyers, M., Koller, D. *et al*. 2008. The chemical genomic portrait of yeast:

uncovering a phenotype for all genes. Science 320: 362-365.

Igic, B. and Busch, J.W. 2013. Is self-fertilization an evolutionary dead end? New Phytol. 198:

386-397.

Igic, B. and Kohn, J.R. 2006. The distribution of plant mating systems: study bias against

obligately outcrossing species. Evolution 60: 1098-1103.

Jarne, P. and Auld, J.R. 2006. Animals mix it up too: the distribution of self-fertilization among

hermaphroditic animals. Evolution 60: 1816-8124.

Kamran-Disfani, A. and Agrawal, A.F. 2014. Selfing, adaptation and background selection in

finite populations. J. Evol. Biol. 27:1360-1371.

Keightley, P.D. and Otto, S. P. 2006. Interference among deleterious mutations favour sex and

recombination in finite populations. Nature Letters 443: 89-92.

Kimura, M. 1962. On the probability of fixation of mutant genes in a population. Genetics 47:

713-719.

Kimura, M., and Ohta, T. 1969. Probability of fixation of a mutant gene in a finite population

when selective advantage decreases with time. Genetics 65: 525-534.

Kishony, R. and Leibler, S. 2003. Environmental stress can alleviate the average deleterious

effect of mutations. J. Biol. 2: 14.

Kraemer, S.A., Morgan, A.D., Ness, R.W., Keightley, P.D. and Colegrave, N. 2015. Fitness

effects of new mutations in *Chlamydomonas reinhardtii* across two stress gradients. J. Evol.

Biol. 29: 583-593.

Latta IV, L.C., Peacock, M., Civitello, D.J., Dudycha, J.L., Meik, J.M. and Schaack, S. 2015.

The phenotypic effects of spontaneous mutations in different environments. Am. Nat 185: 243-

252.

Leimu, R. and Fischer, M. 2008. A meta-analysis of local adaptation in plants. PLoS One 3:

e4010.

Loewe, L. 2006. Quantifying the genomic decay paradox due to Muller's ratchet in human

mitochondrial DNA. Genet. Res. Camb. 87: 133-159.

Loewe, L. and Lamatsch, D.K. 2008. Quantifying the threat of extinction from Muller's ratchet

in the diploid Amazon molly (*Poecilia formosa*). BMC Evol. Biol. 8: 88.

Martin, L. and Lenormand, T. 2006. The fitness effect of mutations across environments: a

survey in light of fitness landscape models. Evolution 60: 2413-2427.

McVean, G.A.T. and Charlesworth, B. 2000. The effects of Hill-Robertson interference between

weakly selected mutations on patterns of molecular evolution and variation. Genetics 155: 929-

944.

Muller, H.J. 1932. Some genetic aspects of sex. Am. Nat. 66: 118-138.

Nordborg, M. 1997. Structured coalescent processes on different time scales. Genetics 146:

1501-1514.

Peischl, S. and Kirkpatrick, M. 2012. Establishment of new mutations in changing environments.

Genetics 191: 895-906.

Pollak, E. 1987. On the theory of partially inbreeding finite populations. I. partial selfing.

Genetics 117: 353-360.

Roze, D. 2016. Background selection in partially selfing populations. Genetics 203: 937-957.

Rutter, M.T., Wieckowski, Y.M., Murren, J. and Strand, A.E. 2017. Fitness effects of mutations:

testing genetic redundancy in *Arabidopsis thaliana*. J. Evol. Biol. 30: 1124-1135.

Siepielski, A.M. Morrissey, M.B., Buoro, M., Carlson, S.M., Caruso, C.M., Clegg, S.M.,

Coulson, T., Dibattista, J., Gotanda, K.M., Francus, C.D. *et al*. 2017. Precipitation drives global

variation in natural selection. Science 355: 959-962.

Slotte, T., Foxe, J. P., Hazzouri, K. M. and Wright, S. I. 2010. Genome-wide evidence for

efficient positive and purifying selectionin *Capsella grandiflora*, a plant species with a large

effective population size. Mol. Biol. Evol. 27: 1813-1821.

Soderberg, R.J. and Berg. O.G. 2007. Mutational interference and the progression of Muller's

ratchet when mutations have a broad range of deleterious effects. Genetics 177: 971-986.

Stebbins, G. L. 1957. Self fertilization and population variability in the higher plants. Am. Nat.

91: 337-354.

Stephan W., Wiehe, T.H.E. and Lenz, M.W. 1992. The effect of strongly selected substitutions

on neutral polymorphisms: Analytical results based on diffusion theory. Theor. Pop. Biol. 41:

237-254.

Vasseur, D.A. and Yodvis, P. 2004. The color of environmental noise. Ecology 85: 1146-1152.

Wardlaw, A.M., and Agrawal, A.F. 2012. Temporal variation in selection accelerates mutational

decay by Muller's ratchet. Genetics 191: 907-916.