11 Jan 2018

Warning SINEs: *Alu* elements, evolution of the human brain, and the spectrum of

neurological disease

Peter A. Larsen[*1,2], Kelsie E. Hunnicutt[1], Roxanne J. Larsen[3], Anne D. Yoder[1,2], and Ann M.

Saunders[4]

[1]Department of Biology, Duke University, Durham, NC 27712

[2]Duke Lemur Center, Duke University, Durham, NC 27712

[3]Duke University School of Medicine, Duke University, Durham, NC 27710

[4]Zinfandel Pharmaceuticals, Inc., Research Triangle Park, NC 27709

Running title: *Alu* elements and the spectrum of neurological disease

*Corresponding Author:

Peter A. Larsen

130 Science Drive, Box 90338

Department of Biology

Duke University

Durham, NC 27708

Office phone: 1-919-613-8727

peter.larsen@duke.edu

Abstract

Alu elements are a highly successful family of primate-specific retrotransposons that have
fundamentally shaped primate evolution, including the evolution of our own species. *Alu*s
play critical roles in the formation of neurological networks and the epigenetic regulation of
biochemical processes throughout the central nervous system (CNS), and thus are
hypothesized to have contributed to the origin of human cognition. Despite the benefits that
*Alu*s provide, deleterious *Alu* activity is associated with a number of neurological and
neurodegenerative disorders. In particular, neurological networks are potentially vulnerable
to the epigenetic dysregulation of *Alu* elements operating across the suite of nuclear-encoded
mitochondrial genes that are critical for both mitochondrial and CNS function. Here, we
highlight the beneficial neurological aspects of *Alu* elements as well as their potential to
cause disease by disrupting key cellular processes across the CNS. We identify at least 37
neurological and neurodegenerative disorders wherein deleterious *Alu* activity has been
implicated as a contributing factor for the manifestation of disease and, for many of these
disorders, this activity is operating on genes that are essential for proper mitochondrial
function. We conclude that the epigenetic dysregulation of *Alu* elements can ultimately
disrupt mitochondrial homeostasis within the CNS. This mechanism is a plausible source for
the incipient neuronal stress that is consistently observed across a spectrum of sporadic
neurological and neurodegenerative disorders.

Keywords: A-to-I editing, Alzheimer's disease, brain connectome, epigenetics,
mitochondria, mosaic brain

 List of Abbreviations:

A-to-I: adenosine-to-inosine

AD: Alzheimer's Disease

ADAR: adenosine deaminase acting on RNA

ALS: Amyotrophic Lateral Sclerosis

AMPA: α-amino-3-hydroxy-5methyl-4-isoxazole propionate

APP: amyloid precursor protein

circRNAs: circular RNAs

CNS: central nervous system

FLAM: free left *Alu* monomer

LINE: long interspersed element

L1: long interspersed element-1

LTR: long-terminal repeat

mRNA: messenger RNA

PD: Parkinson's Disease

pre-mRNA: precursor messenger RNA

SEDs: super-enhancer domains

SINE: short-interspersed element

TADs: topologically associating domains

TOMM: translocase of outer mitochondrial membrane

Introduction

Retrotransposons are mobile genetic elements that utilize an RNA intermediate to

copy and paste themselves throughout the genome. There are two primary groups of

retrotransposons, those having long-terminal repeats (LTRs) and those without (non-LTR)

(Cordaux and Batzer 2009). In the human genome, non-LTR retrotransposons consist of long

interspersed elements (LINEs) and short interspersed elements (SINEs) and these collectively

account for a remarkable ~33% of total genome sequence (Cordaux and Batzer 2009). *Alu*

elements are primate-specific SINEs that are approximately 300 nucleotides in length and are

abundant in the human genome, with over 1.3 million elements accounting for at least 11% of

overall DNA sequence (Deininger et al. 2003; Hancks and Kazazian 2016). Although once

considered to be useless 'junk DNA', the prevalence, diversity, and non-random distribution

of *Alu* elements across primate genomes is suggestive of a functional advantage. Indeed, a

large body of evidence documents that *Alu* elements have directly influenced primate

evolution by facilitating genome innovation through: novel gene formation, elevated

transcriptional diversity, long non-coding RNA and microRNA evolution (including circular

RNAs), transcriptional regulation, and creation of novel response elements (Vansant and

Reynolds 1995; Norris et al. 1995; Britten 1997; Lev-Maor et al. 2003; Polak and Domany

2006; Laperriere et al. 2007; Lin et al. 2008, 2016; Lehnert et al. 2009; Cordaux and Batzer

2009; Shen et al. 2011; Jeck et al. 2013; Töhönen et al. 2015; Luco 2016; Chen and Yang

2017). Moreover, *Alus* fundamentally alter the three-dimensional architecture and spatial

organization of primate genomes by defining the boundaries of chromatin interaction

domains (i.e., topologically associating domains (TADs); Dixon et al. 2012). A growing body

of evidence indicates that genome architecture has a direct influence on biological function,

and the observation that *Alus* are enriched within both TADs and super-enhancer domains

(SEDs) supports the hypothesis that *Alus* directly influence a wide range of critically

important processes within primates across multiple levels, from overall genome stability to

tissue-specific gene regulation (Huda et al. 2009; Dixon et al. 2012; Soibam 2017; Glinsky

2018). In light of the functional benefits that *Alus* provide primates, it is interesting to note

that *Alu* retrotransposition events occurred at an estimated 15-fold higher rate in the human,

chimpanzee, and bonobo lineage (as compared to other great apes) and a 2.2-fold higher rate

in humans when compared to chimpanzee and bonobo (Hedges et al. 2004; Prüfer et al. 2012;

Hormozdiari et al. 2013). These evolutionary patterns indicate that positive selection is acting

to maintain *Alu* elements in primate genomes, especially within humans (Mattick and Mehler

2008; Tsirigos and Rigoutsos 2009).

One of the most fascinating and biologically important aspects of *Alu* elements is that

they serve an important role in the formation and function of the brain connectome (Oliver

and Greene 2011; Li and Church 2013; Smalheiser 2014; Sakurai et al. 2014; Prendergast et

al. 2014; Linker et al. 2017; Bitar and Barry 2018). Many lines of evidence connect *Alu*

elements with neurogenesis and critical neuronal biochemical processes, including: somatic

retrotransposition in developing neurons (in parallel to L1 retrotransposition; Baillie et al.

2011; Kurnosov et al. 2015), formation of regulatory circRNAs that are enriched in the

central nervous system (CNS) and concentrated at synapses (Jeck et al. 2013; Rybak-Wolf et

al. 2015; Chen and Schuman 2016; Floris et al. 2017), regulation of genes that are essential

for proper neuron function (e.g. *ACE*, *SMN1*, *SMN2*, *SLC6A4*; Wu et al. 2013; Ottesen et al.

2017; Schneider et al. 2017), and elevated adenosine-to-inosine (A-to-I) RNA editing in the

brain (Mehler and Mattick 2007; Kurnosov et al. 2015; Behm and Öhman 2016). In

particular, epigenetic A-to-I editing plays a significant role in mediating neuronal gene

expression pathways (Tariq and Jantsch 2012) with *Alus* serving as the primary target for

RNA editing in primates (Picardi et al. 2015; Behm and Öhman 2016). Beyond RNA editing

mechanisms, human neuronal gene pathways are regulated by noncoding RNAs originating

from *Alu* elements (e.g., BC200 and NDM29) and specific *Alu* subfamilies contain retinoic

acid response elements which help to regulate neural patterning, differentiation, and axon

outgrowth (Vansant and Reynolds 1995; Laperriere et al. 2007; Maden 2007; Castelnuovo et

al. 2010; Smalheiser 2014). Moreover, recent discoveries indicate *Alu* elements underlie the

formation of a vast number of human-specific circRNAs that are hypothesized to play

important roles in neurological gene expression pathways (Jeck et al. 2013; Rybak-Wolf et al.

2015; Chen and Schuman 2016; Dong et al. 2017). There is a deep connection between *Alus*

and the formation and function of primate neurological networks, and this has led to the

hypothesis that *Alu* elements were essential for development of the transcriptional diversity

and regulation required for the genesis of human cognitive function (Mattick and Mehler

2008; Oliver and Greene 2011; Li and Church 2013; Sakurai et al. 2014).

Despite the functional benefits that *Alus* have provided primate genomes, *Alu*

elements can disrupt gene expression and function through many pathways (Figure 1;

Deininger and Batzer 1999; Deininger 2011; Tarallo et al. 2012; Ade et al. 2013; Elbarbary et

al. 2016; Varizhuk et al. 2016). For this reason, the genome tightly regulates *Alus* using both

DNA methylation and histone (H3K9 methylation) modification in order to control their

expression and *de novo* retrotransposition (Varshney et al. 2015; Elbarbary et al. 2016; Mita

and Boeke 2016) and there is mounting evidence indicating that the loss of these epigenetic

control mechanisms (due to aging, cellular senescence, environmental factors and stress)

contributes to many forms of cancer, diabetes, osteoporosis, and several mental and

neurodegenerative disorders (Szpakowski et al. 2009; Belancio et al. 2010; Muotri et al.

2010; Jintaridth et al. 2013; Dannlowski et al. 2014; Erwin et al. 2014; Bundo et al. 2014;

Sun et al. 2014; Goodier 2016; Neven et al. 2016; Bedrosian et al. 2016; Shpyleva et al.

2017; Thongsroy et al. 2017). With respect to deleterious *Alu* pathways and neurological

disease, there are at least 37 mental and neurodegenerative disorders wherein *Alu* elements

are hypothesized to disrupt key cellular processes, thereby resulting in or contributing to the

diseased state (Table 1).

Given the tight connection between *Alu* elements and the formation and function of

the nervous system, it is likely that the dysregulation of *Alu* elements contributes to many

sporadic or idiopathic neurological disorders observed across the global human population

(Larsen et al. 2017). Here, we highlight both the beneficial neurological aspects of *Alu*

elements as well as their potential to cause neurological disease. We focus on a novel

hypothesis that identifies a potential epigenetic vulnerability to neurological networks that

has likely escaped purifying selection. The *Alu* neurodegeneration hypothesis (sensu Larsen

et al. 2017) posits that the epigenetic dysregulation of *Alu* elements ultimately serves to

disrupt mitochondrial homeostasis in neurological networks, thereby setting the stage for

increased neuronal stress and neurodegeneration. Given this hypothesis, it is noteworthy that

many of the *Alu*-disrupted genes associated with neurological disorders are related to

mitochondrial function and trafficking, including nuclear-encoded mitochondrial genes (i.e.,

mitonuclear) which help to regulate oxidative stress and metabolic processes in the CNS

(Table 1). Mitochondrial dysfunction is implicated across the spectrum of neurological and

neurodegenerative disorders that are observed in humans and this pattern is suggestive of a

genetic vulnerability that has evolved in humans. Considering this, we begin by reviewing the

integral role that *Alu* elements have played in human evolution through brain-specific

epigenetic A-to-I RNA editing pathways and neurological network formation. Although these

Alu-related processes are hypothesized to have contributed to the origin of human cognition,

they are likely accompanied by age or stress-related epigenetic vulnerabilities to the CNS,

with mitochondrial pathways being especially sensitive.

Alu elements, A-to-I editing, and evolution of the human brain

172   *Alu* elements are non-randomly distributed throughout the genome. They occur most

frequently within introns and are enriched within genes involved in metabolic, mitochondrial,

cellular transport, and binding pathways (Grover et al. 2003; de Andrade et al. 2011; Larsen

et al. 2017). *Alu* nucleotide sequences and lengths (~300 bp) are generally conserved (Batzer

and Deininger 2002) and it is this seemingly simple aspect of *Alu* biology that is of

monumental biological importance. When inserted within a gene at opposite orientations and

at close proximity, *Alus* bind upon themselves post-transcriptionally, resulting in the

formation of a duplex stem-loop structure that is stabilized by the *Alu* nucleotide sequence

and length (Figure 2; Athanasiadis et al. 2004). These *Alu*-based secondary structures

fundamentally alter the shape of pre-mRNA molecules and serve as the primary binding site

for ADAR proteins, which bind to the double-stranded pre-mRNA duplex and edit adenosine

(A) residues to inosine (I) thereby recoding pre-mRNAs (Figure 2). When operating in

coding regions (either directly or indirectly), the translation machinery interprets the resulting

I residues as guanosine (G) and this mechanism accounts, in part, for the incredible diversity

observed in the human proteome that is not encoded within the original DNA sequence

(Nishikura 2016). However, the vast majority of A-to-I editing operating on *Alu* elements

occurs within pre-mRNA introns and 3' UTRs and this can directly influence gene regulation

and function in a surprising number of ways, including: the creation of novel splice donor

and acceptor sites that result in *Alu* exonization and alternative gene splicing (Nishikura

2016); recoding of exons immediately adjacent to *Alus* (Daniel et al. 2014); disruption of

RNAi pathways (Chen and Carmichael 2008); production of novel micro-RNA regulatory

sites (Borchert et al. 2009); and increased nuclear retention of promiscuously edited mRNAs

(Chen and Carmichael 2008).

Although A-to-I editing plays an essential role in generating transcriptional diversity

across eukaryotes, *Alu* elements provide primate-specific RNA editing opportunities. An

example of this is found when comparing the rodent-specific SINE B1 family to *Alu*. Both

B1 and *Alu* SINE families originated from 7SL RNA (Ullu and Tschudi 1984; Vassetzky et

al. 2003) yet rodent-specific B1 elements are approximately half the length (~140 bp) of

primate *Alus* (~300 bp) and have greater levels of intra-nucleotide variation. When

considering the functional mechanics of A-to-I editing (Figure 2), the shorter lengths and

more variable rodent B1 elements result in the formation of shorter and less-stable double-

stranded stem-loop structures in rodent pre-mRNAs. Thus, the molecular properties that

separate rodent B1 from primate *Alu* translate to key functional genomic differences that have

influenced evolutionary processes within each lineage (Eisenberg et al. 2005; Neeman et al.

2006; Picardi et al. 2015; Tan et al. 2017).

A-to-I editing associated with *Alu* elements is perhaps one of the most functionally

important yet underappreciated aspect of *Alu* biology. Approximately 90% of A-to-I editing

within primate gene networks centers on *Alu* elements and this has fundamentally shaped

primate evolution, including the evolution of our own species where A-to-I editing is

estimated to occur at over 100 million sites in the human transcriptome (Bazak et al. 2014).

Moreover, recent data supports a connection between A-to-I editing and neurological

network formation, with elevated editing levels occurring throughout neurogenesis (Behm

and Öhman 2016). Genes encoding for key neurological proteins involved in

neurotransmission, neurogenesis, gliogenesis, and synaptogenesis are subject to enhanced A-

to-I editing and thus a number of studies have hypothesized a strong link between A-to-I

RNA editing pathways and brain development and function (Schmauss and Howe 2002;

Mehler and Mattick 2007; Tan et al. 2009; Sakurai et al. 2014; Liscovitch et al. 2014; Behm

and Öhman 2016; Hwang et al. 2016; Picardi et al. 2017b). A recent analysis of A-to-I

editing in over 8,500 human samples identified tissue specific editing patterns with elevated

editing levels in the brain, including unique patterns in the cerebellum (Tan et al. 2017). The

vast majority of these neurologic A-to-I editing events are operating on *Alu* elements and,

when combined with human-specific *Alu* evolution (Hedges et al. 2004; Cordaux and Batzer

2009; Prüfer et al. 2012; Hormozdiari et al. 2013), this observation serves as the foundation

for the hypothesis that *Alu* elements and *Alu*-related pathways contributed to the evolution of

enhanced human cognitive abilities (Mattick and Mehler 2008; Li and Church 2013).

Given the relationship between *Alu*-centric A-to-I editing and the formation and

function of the CNS, it is important to expand upon the neuro-specific functions of ADAR

proteins. Three ADAR proteins are identified (ADAR1, ADAR2, and ADAR3) and these

proteins have distinct tissue-specific expression patterns (Picardi et al. 2015; Tan et al. 2017).

ADAR1 and ADAR2 co-opt to regulate neuronal activity by editing key neurotransmitter

receptors and ion channels in the CNS (Hood and Emeson 2011). Interestingly, both ADAR2

and ADAR3 have unique brain-specific expression patterns with ADAR2 being highly

expressed in the brain and ADAR3 exclusively expressed in the brain (Mehler and Mattick

2007). Until recently the functional role of the brain-specific ADAR3 protein was largely

unknown, however, Oakes et al. (2017) discovered that ADAR3 competes with ADAR2 to

regulate glutamate receptor subunit B (*GRIA2*) A-to-I editing. The *GRIA2* protein forms a

critical subunit of α-amino-3-hydroxy-5methyl-4-isoxazole propionate (AMPA) receptors,

which regulate synaptic calcium and are involved with synaptic plasticity, memory, and

learning (Wright and Vissel 2012). Remarkably, A-to-I editing of a specific adenosine

nucleotide within *GRIA2* results in an amino acid change that alters the GluR-2 protein

conformation, thus disrupting calcium permeability of the AMPA receptor and potentially

contributing to epilepsy, amyotrophic lateral sclerosis (ALS), and schizophrenia (see Oakes

et al. 2017). In light of Oakes et al. (2017), the brain-specific expression pattern of ADAR3

indicates that this protein helps to offset A-to-I editing by ADAR2, perhaps serving to

mediate enhanced RNA editing processes throughout the CNS.

Considering the essential role that A-to-I editing processes play in the CNS, the

dysregulation of these processes can have a profound impact on the stability of neurological

networks (Mehler and Mattick 2007; Rice et al. 2012; Hwang et al. 2016). With respect to

ADAR proteins, mutations within ADAR1 have been linked to Aicardi-Goutières syndrome

(characterized by severe brain dysfunction; Rice et al. 2012) and *Alu*-related alternative

splicing events of ADAR2 are linked to glioma (Li et al. 2015). Disruption of ADAR1

editing increases production of unedited RNAs which interact with MAV proteins in the

outer mitochondrial membrane, ultimately serving to activate inflammatory response

pathways (Bajad et al. 2017; Gallo et al. 2017) and perhaps providing a mechanism for

inflammatory diseases of the CNS (Hofer and Campbell 2016). ADAR2 knockout mice

display epileptic seizures and neuronal death caused by an influx of calcium owing to the

disruption of *GRIA2* editing (see above). The interference of A-to-I editing processes

associated with the *KCNA1* gene (encoding a protein essential for potassium regulation and

neuron excitability) is hypothesized to underlie Episodic Ataxia Type-1 disorder, a disease of

the CNS characterized by seizures, stress-induced ataxia, and myokymia (Ferrick-Kiddie et

al. 2017). Moreover, a reduction of A-to-I editing has been observed within hippocampal

tissues of Alzheimer's brains versus healthy controls (Khermesh et al. 2016). From a broader

perspective, the disruption of A-to-I editing processes across the CNS has been linked to a

wide variety of mental and neurodegenerative disorders including major depression and

suicide, epilepsy, schizophrenia, Alzheimer's disease (AD), and ALS (Gurevich et al. 2002;

Kawahara et al. 2004; Kwak and Kawahara 2005; Maas et al. 2006; Kubota-Sakashita et al.

2014; Khermesh et al. 2016; Weissmann et al. 2016; Gal-Mark et al. 2017).

***Alu* elements, neurogenesis, and the human brain connectome**

There is a strong connection between *Alu* A-to-I editing and the development and

function of the brain, therefore it is impossible to disentangle *Alus* from the formation and

function of neurologic networks (Mehler and Mattick 2007; Tan et al. 2009; Behm and

Öhman 2016). It is estimated that the human brain is comprised of over 100 billion neurons

that are organized into functional hubs or parcels collectively forming the brain connectome

(Van Den Heuvel and Sporns 2013). Beyond major structures of the brain (e.g., cerebellum,

frontal cortex, hippocampus, etc.), the existence of connectome parcels shared across

unrelated individuals is indicative of an evolutionary conserved process underlying

neurological network formation and operating throughout neurogenesis. For example, a

recent study mapped the cerebral cortex using multi-modal magnetic resonance imaging and

identified 180 connectome parcels that were largely shared across 210 healthy adults (Glasser

et al. 2016). Understanding the molecular processes that contribute to the formation of the

human brain connectome is essential for understanding the origin of human-specific

neurological disorders and diseases observed across the global distribution of our species.

This is especially true for neurodegenerative conditions that are hypothesized to originate in

functional network hubs and progress along neuronal network connections (e.g., AD; Seeley

2017; Cope et al. 2018).

A growing body of evidence indicates that retrotransposons (including both LINEs

and SINEs) are active throughout neurogenesis and contribute to mosaic neuron genomes that

ultimately form the human brain connectome (Muotri et al. 2005; Erwin et al. 2014;

Kurnosov et al. 2015; Evrony 2016; Paquola et al. 2016; Linker et al. 2017). Although

somatic L1 retrotransposition events within developing neurons have received much

attention, it is noteworthy that *Alu* retrotransposition occurs in parallel with L1 throughout

neurogenesis (Baillie et al. 2011; Kurnosov et al. 2015), thus providing primate-specific

aspects of neurologic network formation. Furthermore, there is evidence that unites *Alu*

elements with retinoic acid regulation (Vansant and Reynolds 1995; Laperriere et al. 2007),

which is essential for neuronal patterning and differentiation throughout neurogenesis and is

a potential regulator of neuron regeneration (Maden 2007). Retinoic acid is vital for the

establishment, maintenance, and repair of neuronal networks and, given the presence of

retinoic acid response elements in *Alu* elements, it is possible that *Alu* activity during

neurogenesis is connected to retinoic acid signaling processes.

Considering the *Alu* regulatory pathways discussed above, it is of great interest to

note that retrotransposition of *Alu* elements is hypothesized to occur at elevated levels within

the dentate gyrus of the hippocampus, the putative site of adult neurogenesis (Kurnosov et al.

2015). Moreover, A-to-I editing levels steadily increase as neural progenitor cells develop

into adult neurons (Behm and Öhman 2016). These data indicate that at least two

retrotransposon-centric processes (somatic retrotransposition of both LINES and SINES and

enhanced A-to-I editing operating primarily on *Alu* elements) are major contributors to

neurogenesis, perhaps serving to establish the neuronal and biochemical diversity that

underlies the ~100 billion neuron brain connectome. Remarkably, emerging data suggests

that a third *Alu*-centric process is associated with the formation and function of neurological

networks, this being the production of circRNAs that are enriched in the brain and

concentrated at synaptic junctions (Jeck et al. 2013; Rybak-Wolf et al. 2015; Chen and

Schuman 2016). Identifying vulnerabilities to each of these retrotransposon-centric processes

will likely contribute to the identification of novel mechanisms underlying mental disorders

and neurologic disease and could lead to novel therapeutic interventions.

Pathways to incipient neuronal stress and neurological disease

The disruption of *Alu*-centric epigenetic RNA editing processes is implicated across

the entire spectrum of neurologic disorders (see above). In light of this observation, it is

interesting to note that another, seemingly unrelated, feature of many neurological disorders

is mitochondrial dysfunction (Lin and Beal 2006; Rugarli and Langer 2012; Gottschalk et al.

2014; Petschner et al. 2017). However, we have previously shown that mitonuclear genes are

enriched with *Alu* elements when compared to random (Larsen et al. 2017), which is

consistent with earlier observations regarding the non-random insertion of *Alu* elements into

genes associated with transcriptionally active regions of the genome (Grover et al. 2003; de

Andrade et al. 2011). Thus, it is likely that *Alu*-mediated gene regulatory processes are

actively influencing mitonuclear gene expression, regulation, and protein function through

the pathways discussed above and reviewed in Chen and Yang (2017). Knowing this, the

dysregulation of epigenetic *Alu* regulatory pathways is a plausible source for mitochondrial

stress and dysfunction, with the CNS being particularly vulnerable (Larsen et al. 2017). Such

a mechanism could contribute to the initial activation of complex mitochondrial stress

pathways and incipient neuronal stress associated with sporadic neurologic disorders (e.g.,

inflammation, immune response, mitophagy, etc.). Importantly, these processes would

precede macroscopic pathologies such as protein aggregation and neuronal atrophy observed

in neurodegenerative diseases (Swerdlow et al. 2010; Larsen et al. 2017; Swerdlow 2017).

The *Alu* neurodegeneration hypothesis (*sensu* Larsen et al. 2017) proposes a 'double-

edged sword', whereby the beneficial *Alu*-related processes that underlie neuron diversity and

function also have the potential to disrupt mitochondrial homeostasis across neurological

networks through deleterious cascade events that are facilitated by eroding tissue-specific *Alu*

epigenetic control mechanisms. The stability of the brain's connectome and the entire CNS

depends on healthy mitochondrial populations within neurons, astrocytes, microglia and

supporting cells (Cai et al. 2011; Viader et al. 2011; Schwarz 2013; Jackson and Robinson

2017). Mitochondria play critical roles for a wide range of essential neuronal processes

including glucose and lipid metabolism, metal ion biosynthesis, cellular trafficking along

axons, neurotransmitter relay across synapses, and synaptic calcium homoeostasis (Schwarz

2013; Harbauer et al. 2014). Therefore, molecular mechanisms that are known to disrupt gene

expression and protein folding of genes that are essential for mitochondrial function can

ultimately disrupt neurological function.

Interference of mitochondrial dynamics across the CNS is consistently hypothesized

to occur during the earliest stages of mental, neurological, and neurodegenerative disorders

ranging from depression, epilepsy, and schizophrenia to ALS, AD, and Parkinson's disease

(PD; Lu 2009; Rezin et al. 2009; Kim et al. 2010; Coskun et al. 2012; Martin 2012;

Gottschalk et al. 2014; Larsen et al. 2017; Flippo and Strack 2017; Petschner et al. 2017).

Collectively, these disorders are estimated to impact approximately 250 million people

globally, accounting for at least 10.2% of the global disease burden (GBD 2015 Neurologic

Disorders Collaborator Group 2017). The occurrence of sporadic forms of human-specific

neurologic disorders (e.g. non-familial schizophrenia, ALS, late-onset AD, PD, etc.) across

the entire distribution of our species is suggestive of a common yet complex genetic

mechanism that evolved in primates and is amplified in humans (Larsen et al. 2017).

Considering this, we expand on the mitocentric view of idiopathic neurologic disease

manifestation by reviewing the evidence that unites primate-specific *Alu* activity with

incipient neurologic mitochondrial dysfunction.

Eukaryotic mitochondria are hypothesized to have originated from an endosymbiotic

alphaproteobacterium which, over expansive evolutionary time, evolved in parallel with host

genomes into the mitochondrial organelles that we observe today (Roger et al. 2017). The

human mitochondrial genome encodes only 13 proteins yet it is estimated that human

mitochondria depend on approximately ~2,000 genes encoded within the nuclear genome for

their functionality (Calvo et al. 2015; Johnson et al. 2017). These mitonuclear genes are thus

subject to deleterious *Alu* activity and *Alu*-related deleterious events have been linked to

many neurologic and neurodegenerative disorders, including epilepsy, Wilson's disease,

Leigh syndrome, PD, ALS, and AD (Table 1 and references therein; Figure 3). When

considering the incipient mitochondrial dysfunction observed across the spectrum of

neurological neurodegenerative disorders, it is possible that tissue-specific epigenetic

dysregulation of *Alu* elements within the CNS can ultimately manifest into distinct disease

phenotypes (Larsen et al. 2017).

Several interesting patterns emerge when examining the key neurologic processes that

are disrupted through deleterious *Alu* activity (Table 1). For example, mitochondria play an

essential role in maintaining intra-cellular metal ion homeostasis (e.g., iron, copper, and

zinc), the disruption of which can result in the increased production of free radicals that

damage mitochondria and contribute to the increased production of reduced oxygen species

(ROS; Rossi et al. 2004; Madsen and Gitlin 2007). The brain is especially sensitive to ROS

production, and iron, copper, and zinc-related oxidative stress has been linked to many

neurodegenerative disorders including AD, PD, and Wilson's disease (Rossi et al. 2004;

Madsen and Gitlin 2007). It is notable that deleterious *Alu* activity has been identified in

several genes that are essential to maintaining proper iron and copper homeostasis, including

FXN, *ATP7A*, *ATP7B*, *HMBS*, *NDUFS2*, *SLC30A6*, and *PARK7* (*DJ-1*) (Table 1; Gu et al.

2007; Kaler 2011; Girotto et al. 2014). Knowing this, it is possible that either global or tissue-

specific dysregulation of *Alu* elements within mitonuclear genes can alter mitochondrial

metal ion processing pathways thereby contributing to increased ROS production leading to

neurologic stress.

A second interesting pattern with respect to deleterious neurologic *Alu* activity

concerns metabolic pathways. The efficient processing of glucose and lipids across the CNS

is critical for the stability and function of neurons, and the disruption of mitochondrial-

mediated metabolic pathways has been linked to many neurologic disorders including AD

and peripheral neuropathies (Viader et al. 2013; De La Monte and Tong 2014). Deleterious

Alu activity occurs in genes that are critical for glucose and lipid metabolism, including

ABCD1, *ACAT1*, *ALMS1*, *APOB*, *GK*, *GLA*, *HPRT*, *LPL*, *PDHA1*, *PMM2*, *PSEN1*, *SOD2* and

SPAST (Table 1). Several of these genes encode for mitochondrial-related proteins that have

been implicated in metabolic diseases that directly, or indirectly, contribute to neurological

dysfunction. The connection between *Alu* elements and metabolic pathways is consistent with

the observation that *Alu* elements preferentially insert into metabolic genes, and this has led

to the hypothesis that *Alus* regulate the expression of genes related to Type 1 Diabetes

(Grover et al. 2003; Mirza et al. 2014; Kaur and Pociot 2015). Moreover, *Alu* RNAs act to

suppress the expression of both endothelial nitric oxide synthase (eNOS) and superoxide

dismutase 2 (SOD2) during hyperglycemic conditions (Wang et al. 2016), suggesting a

regulatory role of *Alu* elements during oxidative stress and strengthening the link between

Alu element activity and diabetes.

There is growing evidence linking sporadic AD with dysfunctional metabolic

pathways, leading some to consider AD as a 'Type 3 Diabetes' wherein glycolysis and lipid

homeostasis is altered (Steen et al. 2005; De La Monte et al. 2006; De La Monte and Wands

2008; De La Monte and Tong 2014; De Felice and Lourenco 2015; Mittal et al. 2016). The

most well-documented risk factor for AD is a variant within *APOE* (*APOE* ε4), a gene which

encodes for a glycoprotein that that mediates cholesterol and lipid transport (Saunders et al.

1993; Strittmatter et al. 1993; Mahley and Rall 2000). The *APOE* ε4 allele is strongly

associated with earlier onset of AD, and it is hypothesized that this is a result of the

disruption of cholesterol processing and subsequent accumulation of amyloid precursor

proteins (APP; i.e., the Amyloid cascade hypothesis). Although the 'Amyloid cascade

hypothesis' has dominated Alzheimer's research for decades (Hardy and Higgins 1992;

Selkoe 2000; McKhann et al. 2011), the failure of multiple drug trials targeting amyloid

pathways has led many in the Alzheimer's research community to search for alternative

hypotheses that can help explain the origin of neurodegenerative disease as well as novel

molecular pathways with therapeutic potential (Herrup 2015).

It is of particular interest then to note that a second genetic risk factor for AD,

TOMM40, is located immediately adjacent to *APOE* on human chromosome 19, and the two

genes are in tight linkage disequilibrium (Lyall et al. 2014; Roses et al. 2016a). *TOMM40*

encodes for a beta-barrel protein that ultimately forms a central pore in the outer

mitochondrial membrane (Shiota et al. 2015) and, much like *APOE*, genetic variants of

TOMM40 are linked to cognitive impairment and neurodegenerative disease (Roses 2010;

Gottschalk et al. 2014; Greenbaum et al. 2014; Roses et al. 2016b; Arpawong et al. 2017).

The most-well known of these *TOMM40* variants is the rs10524523 (rs523) homopolymer

repeat, a variable stretch of deoxythymidine (T) located within *TOMM40* intron 6 (Roses

2010). The rs523 poly-T varies in length from approximately 12 to 46 nucleotides, and the

longer variants are statistically associated with thinning of the hippocampus (independent of

the *APOE* ε4 allele; Burggren et al. 2017) and earlier onset of AD (Lutz et al. 2010; Roses et

al. 2010). Interestingly, rs523 is embedded within tandemly repeated *Alu* elements and

originated from an *Alu* insertion event (Larsen et al. 2017). At least 149 *Alu* A-to-I editing

events are identified within *TOMM40*, the majority of which are associated with *Alu* elements

surrounding the rs523 repeat and intron 9 (Picardi et al. 2017a).

There is a potentially important link that unties *APOE* APP processing with the

functional mechanics of pre-protein transport through the TOMM pore. It is possible that

conformational changes of the Tom40 protein, potentially originating from *Alu*-mediated

events (see above, reviewed in Elbarbary et al. 2016; Chen and Yang 2017; Larsen et al.

2017), can ultimately serve to restrict the passage of lipids across the outer-mitochondrial

membrane (Larsen et al. 2017). When combined with altered APP processing, this process

could account for the initial site of intra-cellular protein accumulation that is hypothesized to

precede extra-cellular plaque formation during very early stages of AD (Skovronsky et al.

1998; D'Andrea et al. 2001; Takahashi et al. 2002). Consistent with this hypothesis is the

direct observation of APP accumulation at the TOMM pore (Devi et al. 2006) as well as

functional tolerance of Tom40 conformational changes by mitochondria (Mager et al. 2011;

Kuszak et al. 2015). Importantly, this mechanism could help to explain the common patterns

of protein accumulation (e.g., amyloid plaques and alpha-synuclein Lewy bodies) observed

across the spectrum of neurodegenerative disease, including ALS, AD, and PD (Ross and

Poirier 2004; Gottschalk et al. 2014; Larsen et al. 2017). An age or stress-related component

to Tom40 conformational changes comes with the epigenetic dysregulation of *Alu* elements

associated with the aging process or traumatic stress (see Larsen et al. 2017). Whether or not

these processes are directly mediated by deleterious *Alu* events remains to be tested, however,

it is notable that *Alu* exons and *Alu* somatic retrotransposition events have been identified in

several TOM genes that are required for the stability of the translocase of the outer

mitochondrial membrane and pre-protein import, including *TOMM5*, *TOMM7*, *TOMM22*,

TOMM40, and *TOMM40L* (Baillie et al. 2011; de Andrade et al. 2011; Lin et al. 2016).

With respect to *Alu* elements, mitochondrial dysfunction, and the broader pathological

scope of AD and other neurodegenerative diseases, there is evidence suggesting that *Alu*-

derived peptides interact with tau proteins, perhaps serving a regulatory role for tau

phosphorylation (Hoenicka et al. 2002). Tau is a microtubule associated protein that

functions to stabilize axonal microtubules and to transport mitochondria along axons, and

taupathies (including tau hyperphosphorylation) are a characteristic feature of several

neurodegenerative diseases including AD, progressive supranuclear palsy, corticobasal

degeneration, and Pick's disease (Ittner and Götz 2011; Khanna et al. 2016). The *MAPT* gene

encodes for tau and alternative splicing events of *MAPT* result in multiple tau isoforms

(Reddy 2011). Approximately 86 *Alu* elements (including FLAMs) are distributed throughout

MAPT introns and A-to-I editing is occurring at 315 *Alu* related sites with elevated levels at

the 3' end of *MAPT* (REDIportal database; Picardi et al. 2017a). When considering the

potential for *Alu* structural variants of *MAPT* (including DNA and pre-mRNA secondary

structures) and evidence of *Alu* RNAs interacting with tau proteins (Hoenicka et al. 2002), we

recommend additional studies aimed at elucidating the regulatory impacts that *Alu* elements

might have on *MAPT* gene expression and tau phosphorylation.

A-to-I editing and the potential for mitochondrial stress

Although several neurological disorders are hypothesized to be the result of disruptive

A-to-I editing processes across the CNS (see above), it is presently unknown whether or not

these processes are actively influencing mitochondrial function. What evidence is there

indicating that post-transcriptional modification of mitonuclear genes can alter gene

expression or function? Are there particular neurological or neurodegenerative disorders that

are associated with mitonuclear genes that have elevated levels of A-to-I editing? To provide

insights into these questions, we searched the REDIportal A-to-I editing database (Picardi et

al. 2017a) for mitonuclear genes where 1) A-to-I editing has been identified within *Alu*

elements in coding regions and 2) A-to-I editing has contributed to non-synonymous amino

acid changes. We identified 57 mitonuclear genes with A-to-I editing occurring within

putative *Alu* exons and in 52 of these genes the post-transcriptional modification resulted in

nonsynonymous amino acid changes (Supplementary Table 1). Many of these genes are

involved with essential neuronal processes including calcium binding and transport, zinc

transport, apoptosis regulation, voltage-gated ion channels, and mitochondrial elongation

with notable examples including *ADSL*, *BAX*, *CASP2*, *COQ2*, *DFFB*, *FBXO18*, *LYRM4*,

PACRG, and *SLC30A6* (Supplementary Table 1).

From a broader perspective, we identified enhanced A-to-I editing across 134

mitonuclear genes that are associated with a spectrum of neurologic and neurodegenerative

disorders ranging from depression, tobacco use disorder, and bipolar disorder to ALS, Leigh

syndrome, PD, and AD (Supplementary Table 2). In light of these patterns, we hypothesize

that system-wide or tissue-specific epigenetic dysregulation of *Alu* A-to-I editing within the

CNS can serve to disrupt key mitochondrial biochemical processes, thus potentially

contributing to incipient mitochondrial and neuronal stress (Figure 3).

Conclusions

Enhanced somatic retrotransposon throughout neurogenesis contributes to the mosaic

brain, however, such activity likely contributes to mosaic pathways leading to disease (Erwin

et al. 2014). Elucidating these pathways might ultimately provide insight into the sporadic

nature of idiopathic diseases that are impacting the global human population. The disruption

of *Alu*-mediated pathways that underlie gene regulation is a plausible mechanism for the

origin of complex human-specific neurologic and neurodegenerative disorders. Although

many of these disorders have similar phenotypes (e.g., mitochondrial dysfunction), it is

possible that these phenotypes arise from deleterious activity operating across tissue-specific

gene networks. If correlated with eroding or fluctuating epigenetic control mechanisms of

retrotransposons that are associated with aging, cellular senescence, and/or cellular stress

(Belancio et al. 2010; Pal and Tyler 2016; Schneider et al. 2017), then such mechanisms

might largely escape purifying selection and would be difficult to detect using traditional

methods (e.g., genome-wide association studies). It is important to note that the *Alu*-centric

mechanisms discussed herein collectively provide a unified framework for multiple

hypotheses that have been put forth regarding the origin of neurodegenerative disease

including inflammation, oxidative-stress, metabolic dysfunction, and accumulation of protein

bodies (see above).

Alu elements have played a pivotal role in the evolution of the human epigenome

(Prendergast et al. 2014), and both hyper- and hypomethylation of *Alu* elements have been

correlated with a number of age-related disorders including Alzheimer's disease, multiple

sclerosis, osteoporosis, and many forms of cancer (Bollati et al. 2009; Jintaridth and

Mutirangura 2010; Belancio et al. 2010; Jintaridth et al. 2013; Neven et al. 2016). In light of

these patterns, as well as the newly discovered regulatory roles of *Alu* elements (Polak and

Domany 2006; Chen and Carmichael 2008; Chen and Yang 2017), we recommend additional

research that focuses on the epigenetic interplay between *Alu* elements and mitochondrial

gene networks in the central nervous system.

Acknowledgments

We thank A.D. Brown, W.K. Gottschalk, and M. Mihovilovic for helpful discussion. C.

Hohoff, B. Ciapa, and an anonymous reviewer kindly reviewed the manuscript and provided

helpful comments. This is Duke Lemur Center publication number 1389.

References

Ade C, Roy-Engel AM, Deininger PL (2013) Alu elements: An intrinsic source of human
genome instability. Curr. Opin. Virol. 3:639–645.
Antonell A, Gelpi E, Sánchez-Valle R, et al (2012) Breakpoint sequence analysis of an AβPP
locus duplication associated with autosomal dominant Alzheimer's disease and severe
cerebral amyloid angiopathy. J Alzheimers Dis 28:303–3088
Arnoldi A, Tonelli A, Crippa F, et al (2008) A clinical, genetic, and biochemical
characterization of SPG7 mutations in a large cohort of patients with hereditary spastic
paraplegia. Hum Mutat 29:522–531.
Arpawong TE, Pendleton N, Mekli K, et al (2017) Genetic variants specific to aging-related
verbal memory: Insights from GWASs in a population-based cohort. PLoS One
12:e0182448.
Athanasiadis A, Rich A, Maas S (2004) Widespread A-to-I RNA editing of Alu-containing
mRNAs in the human transcriptome. PLoS Biol 2:e391.
Baillie JK, Barnett MW, Upton KR, et al (2011) Somatic retrotransposition alters the genetic
landscape of the human brain. Nature 479:534–537.
Bajad P, Jantsch MF, Keegan L, O'Connell M (2017) A to I editing in disease is not fake
news. RNA Biol 14:1223–1231.
Batzer MA, Deininger PL (2002) Alu repeats and human genomic diversity. Nat Rev Genet

3:370–379.
Bazak L, Haviv A, Barak M, et al (2014) A-to-I RNA editing occurs at over a hundred
million genomic sites, located in a majority of human genes. Genome Res 24:365–376.
Bedrosian TA, Linker S, Gage FH (2016) Environment-driven somatic mosaicism in brain
disorders. Genome Med 8:58.
Behm M, Öhman M (2016) RNA Editing: A Contributor to Neuronal Dynamics in the
Mammalian Brain. Trends Genet 32:165–175.
Belancio VP, Roy-Engel AM, Deininger PL (2010) All y'all need to know 'bout
retroelements in cancer. Semin Cancer Biol 20:200–210.
Bhattacharjee A, Yang H, Duffy M, et al (2016) The Activity of Menkes Disease Protein
ATP7A Is Essential for Redox Balance in Mitochondria. J Biol Chem 291:16644–
16658.
Bitar M and Barry G (2017) Multiple innovations in genetic and epigenetic mechanisms
cooperate to underpin human brain evolution. Mol Biol Evol, msx303, doi:
10.1093/molbev/msx303
Bollati V, Schwartz J, Wright R, et al (2009) Decline in genomic DNA methylation through
aging in a cohort of elderly subjects. Mech Ageing Dev 130:234–239.
Bondurand N, Fouquet V, Baral V, et al (2012) Alu-mediated deletion of SOX10 regulatory
elements in Waardenburg syndrome type 4. Eur J Hum Genet 20:990–994. doi:
10.1038/ejhg.2012.29
Bondy-Chorney E, Crawford Parks TE, Ravel-Chapuis A, et al (2016) Staufen1 Regulates
Multiple Alternative Splicing Events either Positively or Negatively in DM1 Indicating
Its Role as a Disease Modifier. PLoS Genet 12:e1005827. doi:
10.1371/journal.pgen.1005827
Bonifati V, Rizzu P, van Baren MJ, et al (2003) Mutations in the DJ-1 gene associated with
autosomal recessive early-onset parkinsonism. Science 299:256–9. doi:
10.1126/science.1077209
Boone PM, Yuan B, Campbell IM, et al (2014) The Alu-rich genomic architecture of SPAST
predisposes to diverse and functionally distinct disease-associated CNV alleles. Am J
Hum Genet 95:143–161.
Borchert GM, Gilmore BL, Spengler RM, et al (2009) Adenosine deamination in human
transcripts generates novel microRNA binding sites. Hum Mol Genet 18:4801–7. doi:
10.1093/hmg/ddp443
Bouchet C, Vuillaumier-Barrot S, Gonzales M, et al (2007) Detection of an Alu insertion in
the POMT1 gene from three French Walker Warburg syndrome families. Mol Genet
Metab 90:93–96. doi: 10.1016/j.ymgme.2006.09.005
Britten RJ (1997) Mobile elements inserted in the distant past have taken on important
functions. Gene 205:177–82.
Brooks EM, Branda RF, Nicklas JA, O'Neill JP (2001) Molecular description of three macro-
deletions and an Alu-Alu recombination-mediated duplication in the HPRT gene in four
patients with Lesch-Nyhan disease. Mutat Res 476:43–54.
Bundo M, Toyoshima M, Okada Y, et al (2014) Increased l1 retrotransposition in the
neuronal genome in schizophrenia. Neuron 81:306–13. doi:
10.1016/j.neuron.2013.10.053
Burggren AC, Mahmood Z, Harrison TM, et al (2017) Hippocampal thinning linked to longer
TOMM40 poly-T variant lengths in the absence of the APOE ε4 variant. Alzheimer's
Dement 13:739–748. doi: 10.1016/j.jalz.2016.12.009
Cai Q, Davis ML, Sheng Z-H (2011) Regulation of axonal mitochondrial transport and its
impact on synaptic transmission. Neurosci Res 70:9–15. doi:
10.1016/j.neures.2011.02.005

Calvo SE, Clauser KR, Mootha VK (2015) MitoCarta2.0: An updated inventory of
mammalian mitochondrial proteins. Nucleic Acids Res 44:D1251–D1257. doi:
10.1093/nar/gkv1003
Castelnuovo M, Massone S, Tasso R, et al (2010) An Alu-like RNA promotes cell
differentiation and reduces malignancy of human neuroblastoma cells. FASEB J
24:4033–4046. doi: 10.1096/fj.10-157032
Checler F, Goiran T, da Costa AC (2017) Presenilins at the crossroad of a functional
interplay between PARK/PARKIN and PINK1 to control mitophagy: implications for
neurodegenerative diseases. Autophagy 15:1–2. doi: 10.1080/15548627.2017.1363950
Chen L-L, Carmichael GG (2008) Gene regulation by SINES and inosines: biological
consequences of A-to-I editing of Alu element inverted repeats. Cell Cycle 7:3294–
3301. doi: 10.4161/cc.7.21.6927
Chen L-L, Yang L (2017) ALUternative Regulation for Gene Expression. Trends Cell Biol
27:480–490. doi: 10.1016/j.tcb.2017.01.002
Chen W, Schuman E (2016) Circular RNAs in Brain and Other Tissues: A Functional
Enigma. Trends Neurosci 39:597–604. doi: 10.1016/j.tins.2016.06.006
Choi B-O, Kim NK, Park SW, et al (2011) Inheritance of Charcot–Marie–Tooth disease 1A
with rare nonrecurrent genomic rearrangement. Neurogenetics 12:51–58. doi:
10.1007/s10048-010-0272-3
Conceição Pereira M, Loureiro JL, Pinto-Basto J, et al (2012) Alu elements mediate large
SPG11 gene rearrangements: further spatacsin mutations. Genet Med 14:143–151. doi:
10.1038/gim.2011.7
Chong-Chong X, Denton KR, Wang Z, et al (2016) Abnormal mitochondrial transport and
morphology as early pathological changes in human models of spinal muscular atrophy.
Dis Model Mech 9:39–49.
Cope TE, Rittman T, Borchert RJ, et al. (2018) Tau burden and the functional connectome in
Alzheimer's disease and progressive supranuclear palsy. Brain, awx347,
doi.org:10.1093/brain/awx347
Cordaux R, Batzer MA (2009) The impact of retrotransposons on human genome evolution.
Nat Rev Genet 10:691–703. doi: 10.1038/nrg2640
Coskun P, Wyrembak J, Schriner SE, et al (2012) A mitochondrial etiology of Alzheimer and
Parkinson disease. Biochim Biophys Acta (BBA)-General Subj 1820:553–564.
D'Andrea MR, Nagele RG, Wang HY, et al (2001) Evidence that neurones accumulating
amyloid can undergo lysis to form amyloid plaques in Alzheimer's disease.
Histopathology 38:120–134. doi: 10.1046/j.1365-2559.2001.01082.x
Daniel C, Silberberg G, Behm M, Öhman M (2014) Alu elements shape the primate
transcriptome by cis-regulation of RNA editing. Genome Biol 15:R28. doi: 10.1186/gb-
2014-15-2-r28
Dannlowski U, Kugel H, Redlich R, et al (2014) Serotonin transporter gene methylation is
associated with hippocampal gray matter volume. Hum Brain Mapp 35:5356–5367. doi:
10.1002/hbm.22555
Dawson G, Kilkus J, Siakotos AN, et al (1995) Mitochondrial abnormalities in CLN2 and
CLN3 forms batten disease. Mol Chem Neuro 29:227–235.
De Andrade A, Wang M, Bonaldo MF, et al (2011) Genetic and epigenetic variations
contributed by Alu retrotransposition. BMC Genomics 12:617.
De Felice FG, Lourenco M V (2015) Brain metabolic stress and neuroinflammation at the
basis of cognitive impairment in Alzheimer's disease. Front Aging Neurosci 7:94. doi:
10.3389/fnagi.2015.00094
De La Monte SM, Tong M (2014) Brain metabolic dysfunction at the core of Alzheimer's
disease. Biochem. Pharmacol. 88:548–559.

De La Monte SM, Tong M, Lester-Coll N, et al (2006) Therapeutic rescue of neurodegeneration in experimental type 3 diabetes: relevance to Alzheimer's disease. J Alzheimers Dis 10:89–109. doi: 10.2165/11597760-000000000-00000

De La Monte SM, Wands JR (2008) Alzheimer's Disease is Type 3 Diabetes—Evidence Reviewed. J Diabetes Sci Technol 2:1101–1113. doi: 10.1177/193229680800200619

Deininger P (2011) Alu elements: know the SINEs. Genome Biol 12:1.

Deininger PL, Batzer MA (1999) Alu repeats and human disease. Mol Genet Metab 67:183–193.

Deininger PL, Moran J V, Batzer MA, Kazazian HH (2003) Mobile elements and mammalian genome evolution. Curr Opin Genet Dev 13:651–658.

Devi L, Prabhu BM, Galati DF, et al (2006) Accumulation of amyloid precursor protein in the mitochondrial import channels of human Alzheimer's disease brain is associated with mitochondrial dysfunction. J Neurosci 26:9057–9068.

Dixon JR, Selvaraj S, Yue F, et al. (2012) Topological domains in mammalian genomes identified by analysis of chromatin interactions. Nature 485:376–380.

Dobrovolny R, Nazarenko I, Kim J, et al (2011) Detection of large gene rearrangements in X-linked genes by dosage analysis: identification of novel α-galactosidase A (GLA) deletions causing Fabry disease. Hum Mutat 32:688–695. doi: 10.1002/humu.21474

Dong R, Ma XK, Chen LL, Yang L (2017) Increased complexity of circRNA expression during species evolution. RNA Biol 14:1064–1074.

Eisenberg E, Nemzer S, Kinar Y, et al (2005) Is abundant A-to-I RNA editing primate-specific? Trends Genet 21:77–81. doi: 10.1016/j.tig.2004.12.005

Elbarbary RA, Lucas BA, Maquat LE (2016) Retrotransposons as regulators of gene expression. Science 351:aac7247.

Erwin JA, Marchetto MC, Gage FH (2014) Mobile DNA elements in the generation of diversity and complexity in the brain. Nat Rev Neurosci 15:497–506.

Evrony GD (2016) One brain, many genomes. Science 354:557–558.

Ferrari L, Scuvera G, Tucci A, et al (2017) Identification of an atypical microdeletion generating the RNF135-SUZ12 chimeric gene and causing a position effect in an NF1 patient with overgrowth. Hum Genet 136:1329–1339. doi: 10.1007/s00439-017-1832-5

Ferrick-Kiddie EA, Rosenthal JJC, Ayers GD, Emeson RB (2017) Mutations underlying Episodic Ataxia type-1 antagonize Kv1.1 RNA editing. Sci Rep 7:41095. doi: 10.1038/srep41095

Flippo KH, Strack S (2017) An emerging role for mitochondrial dynamics in schizophrenia. Schizophr. Res. 187:26–32.

Floris G, Zhang L, Follesa P, Sun T (2017) Regulatory Role of Circular RNAs and Neurological Disorders. Mol Neurobiol 54:156–5165.

Gal-Mark N, Shallev L, Sweetat S, et al (2017) Abnormalities in A-to-I RNA editing patterns in CNS injuries correlate with dynamic changes in cell type composition. Sci Rep 7:43421. doi: 10.1038/srep43421

Gallo A, Vukic D, Michalík D, et al (2017) ADAR RNA editing in human disease; more to it than meets the I. Hum Genet 136:1265–1278. doi: 10.1007/s00439-017-1837-0

Gallus GN, Cardaioli E, Rufa A, et al (2010) Alu-element insertion in an OPA1 intron sequence associated with autosomal dominant optic atrophy. Mol Vis 10:178–183.

GBD 2015 Neurological Disorders Collaborator Group 2015 NDC (2017) Global, regional, and national burden of neurological disorders during 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. Lancet Neurol 16:877–897. doi: 10.1016/S1474-4422(17)30299-5

Girotto S, Cendron L, Bisaglia M, et al (2014) DJ-1 is a copper chaperone acting on SOD1 activation. J Biol Chem 289:10887–99. doi: 10.1074/jbc.M113.535112

Glasser MF, Coalson TS, Robinson EC, et al (2016) A multi-modal parcellation of human
cerebral cortex. Nature 536:171–178. doi: 10.1038/nature18933
Glinsky GV (2018) Contribution of transposable elements and distal enhancers to evolution
of human-specific features of interphase chromatin architecture in embryonic stem cells.
Chromosome Res *in press*
Goodier JL (2016) Restricting retrotransposons: a review. Mob DNA 7:16. doi:
10.1186/s13100-016-0070-z
Gottschalk WK, Lutz MW, He YT, et al (2014) The broad impact of TOM40 on
neurodegenerative diseases in aging. J Parkinsons Dis Alzheimers Dis 1.
Greenbaum L, Springer RR, Lutz MW, et al (2014) The TOMM40 poly-T rs10524523
variant is associated with cognitive performance among non-demented elderly with type
2 diabetes. Eur Neuropsychopharmacol 24:1492–1499.
Grover D, Majumder PP, Rao CB, et al (2003) Nonrandom Distribution of Alu Elements in
Genes of Various Functional Categories: Insight from Analysis of Human
Chromosomes 21 and 22. Mol Biol Evol 20:1420–1424. doi: 10.1093/molbev/msg153
Gu S, Yuan B, Campbell IM, et al (2015) Alu-mediated diverse and complex pathogenic
copy-number variants within human chromosome 17 at p13.3. Hum Mol Genet
24:4061–4077. doi: 10.1093/hmg/ddv146
Gu Y, Kodama H, Watanabe S, et al (2007) The first reported case of Menkes disease caused
by an Alu insertion mutation. Brain Dev 29:105–108. doi:
10.1016/j.braindev.2006.05.012
Gurevich I, Tamir H, Arango V, et al (2002) Altered editing of serotonin 2C receptor pre-
mRNA in the prefrontal cortex of depressed suicide victims. Neuron 34:349–356. doi:
10.1016/S0896-6273(02)00660-8
Hancks DC, Kazazian HH (2016) Roles for retrotransposon insertions in human disease. Mob
DNA 7:1.
Harbauer AB, Zahedi RP, Sickmann A, et al (2014) The protein import machinery of
mitochondria—a regulatory hub in metabolism, stress, and disease. Cell Metab 19:357–
372.
Hardy J, Higgins G (1992) Alzheimer's disease: the amyloid cascade hypothesis. Science
(80-) 256:184–185. doi: 10.1126/science.1566067
Hedges DJ, Callinan PA, Cordaux R, et al (2004) Differential Alu mobilization and
polymorphism among the human and chimpanzee lineages. Genome Res 14:1068–1075.
Herrup K (2015) The case for rejecting the amyloid cascade hypothesis. Nat Neurosci
18:794–799. doi: 10.1038/nn.4017
Hoenicka J, Arrasate M, de Yebenes JG, Avila J (2002) A two-hybrid screening of human
Tau protein: interactions with Alu-derived domain. Neuroreport 13:343–9.
Hofer MJ, Campbell IL (2016) Immunoinflammatory diseases of the central nervous system -
the tale of two cytokines. Br J Pharmacol 173:716–728. doi: 10.1111/bph.13175
Hood JL, Emeson RB (2011) Editing of Neurotransmitter Receptor and Ion Channel RNAs in
the Nervous System. Springer, Berlin, Heidelberg, pp 61–90
Hormozdiari F, Konkel MK, Prado-Martinez J, et al (2013) Rates and patterns of great ape
retrotransposition. Proc Natl Acad Sci USA 110:13457–62. doi:
10.1073/pnas.1310914110
Huang LS, Ripps ME, Korman SH, et al (1989) Hypobetalipoproteinemia due to an
apolipoprotein B gene exon 21 deletion derived by Alu-Alu recombination. J Biol Chem
264:11394–400.
Hunter R, Gagnidze K, McEwen B, Pfaff D. (2015) Stress and the dynamic genome: steroids,
epigenetics, and the transposome. Proc Natl Acad Sci USA 112:6828–6833.
Huda A, Mariño-Ramírez L, Landsman D, Jordan IK (2009) Repetitive DNA elements,

nucleosome binding and human gene expression. Gene 436:12–22.

Hwang T, Park C-K, Leung AKL, et al (2016) Dynamic regulation of RNA editing in human
brain development and disease. Nat Neurosci 19:1093–9. doi: 10.1038/nn.4337

Ittner LM, Götz J (2011) Amyloid-β and tau — a toxic pas de deux in Alzheimer's disease.
Nat Rev Neurosci 12:65–72. doi: 10.1038/nrn2967

Jackson JG, Robinson MB (2017) Regulation of mitochondrial dynamics in astrocytes:
Mechanisms, consequences, and unknowns. Glia. doi: 10.1002/glia.23252

Jeck WR, Sorrentino JA, Wang K, et al (2013) Circular RNAs are abundant, conserved, and
associated with ALU repeats. RNA 19:141–157. doi: 10.1261/rna.035667.112

Jiang J, Wu X, Shen D, et al (2017) Analysis of RP2 and RPGR Mutations in Five X-Linked
Chinese Families with Retinitis Pigmentosa. Sci Rep 7:44465. doi: 10.1038/srep44465

Jintaridth P, Mutirangura A (2010) Distinctive patterns of age-dependent hypomethylation in
interspersed repetitive sequences. Physiol Genomics 41:194–200.

Jintaridth P, Tungtrongchitr R, Preutthipan S, Mutirangura A (2013) Hypomethylation of Alu
elements in post-menopausal women with osteoporosis. PLoS One 8:e70386. doi:
10.1371/journal.pone.0070386

Jobic F, Morin G, Vincent-Delorme C, et al (2017) New intragenic rearrangements in non-
Finnish mulibrey nanism. Am J Med Genet Part A 173:2782–2788. doi:
10.1002/ajmg.a.38381

Johnson SC, Gonzalez B, Zhang Q, et al (2017) Network analysis of mitonuclear GWAS
reveals functional networks and tissue expression profiles of disease-associated genes.
Hum Genet 136:55–65. doi: 10.1007/s00439-016-1736-9

Kaler SG (2011) ATP7A-related copper transport diseases—emerging concepts and future
trends. Nat Rev Neurol 7:15–29. doi: 10.1038/nrneurol.2010.180

Kaneko H, Dridi S, Tarallo V, et al (2011) DICER1 deficit induces Alu RNA toxicity in age-
related macular degeneration. Nature 471:325–30. doi: 10.1038/nature09830

Kaur S, Pociot F (2015) Alu Elements as Novel Regulators of Gene Expression in Type 1
Diabetes Susceptibility Genes? Genes (Basel) 6:577–591. doi: 10.3390/genes6030577

Kawahara Y, Ito K, Sun H, et al (2004) Glutamate receptors: RNA editing and death of motor
neurons. Nature 427:801. doi: 10.1038/427801a

Khanna MR, Kovalevich J, Lee VMY, et al (2016) Therapeutic strategies for the treatment of
tauopathies: Hopes and challenges. Alzheimer's Dement. 12:1051–1065.

Khermesh K, D'Erchia AM, Barak M, et al (2016) Reduced levels of protein recoding by A-
to-I RNA editing in Alzheimer's disease. RNA 22:290–302. doi:
10.1261/rna.054627.115

Kim J, Moody JP, Edgerly CK, et al (2010) Mitochondrial loss, dysfunction and altered
dynamics in Huntington's disease. Hum Mol Genet 19:3919–3935.

Kim Y, Tarallo V, Kerur N, et al (2014) DICER1/Alu RNA dysmetabolism induces Caspase-
8-mediated cell death in age-related macular degeneration. Proc Natl Acad Sci U S A
111:16082–7. doi: 10.1073/pnas.1403814111

Knight Johnson A, Schaefer GB, Lee J, et al (2017) Alu-mediated deletion of PIGL in a
Patient with CHIME syndrome. Am J Med Genet Part A 173:1378–1382. doi:
10.1002/ajmg.a.38181

Kubota-Sakashita M, Iwamoto K, Bundo M, Kato T (2014) A role of ADAR2 and RNA
editing of glutamate receptors in mood disorders and schizophrenia. Mol Brain 7:5. doi:
10.1186/1756-6606-7-5

Kurnosov AA, Ustyugova S V, Nazarov VI, et al (2015) The evidence for increased L1
activity in the site of human adult brain neurogenesis. PLoS One 10:e0117854. doi:
10.1371/journal.pone.0117854

Kuszak AJ, Jacobs D, Gurnev PA, et al (2015) Evidence of Distinct Channel Conformations

and Substrate Binding Affinities for the Mitochondrial Outer Membrane Protein
Translocase Pore Tom40. J Biol Chem 290:26204–26217. doi:
10.1074/jbc.M115.642173

Kutsche K, Ressler B, Katzera H, et al (2002) Characterization of breakpoint sequences of
five rearrangements in L1CAM and ABCD1 (ALD) genes. Hum Mutat 19:526–535.

Kwak S, Kawahara Y (2005) Deficient RNA editing of GluR2 and neuronal death in
amyotropic lateral sclerosis. J Mol Med (Berl) 83:110–20. doi: 10.1007/s00109-004-
0599-z

Lapp HE, Hunter RG (2016) The dynamic genome: transposons and environmental
adaptation in the nervous system. Epigenomics 8:237–249.

Laperriere D, Wang T-T, White JH, Mader S (2007) Widespread Alu repeat-driven
expansion of consensus DR2 retinoic acid response elements during primate evolution.
BMC Genomics 8:23. doi: 10.1186/1471-2164-8-23

Larsen PA, Lutz MW, Hunnicutt KE, et al (2017) The Alu neurodegeneration hypothesis: A
primate-specific mechanism for neuronal transcription noise, mitochondrial dysfunction,
and manifestation of neurodegenerative disease. Alzheimer's Dement. 13:828–838.

Le Guennec K, Veugelen S, Quenez O, et al (2017) Deletion of exons 9 and 10 of the
Presenilin 1 gene in a patient with Early-onset Alzheimer Disease generates longer
amyloid seeds. Neurobiol Dis 104:97–103. doi: 10.1016/j.nbd.2017.04.020

Lehnert S, Loo P Van, Thilakarathne PJ, et al (2009) Evidence for Co-Evolution between
Human MicroRNAs and Alu-Repeats. doi: 10.1371/journal.pone.0004456

Lerner T, Boustany RMN, Anderson JW, et al (1995) Isolation of a novel gene underlying
Batten disease, CLN3. Cell 82:949–57.

Lev-Maor G, Sorek R, Shomron N, Ast G (2003) The birth of an alternatively spliced exon:
3'splice-site selection in Alu exons. Science 300:1288–1291.

Li JB, Church GM (2013) Deciphering the functions and regulation of brain-enriched A-to-I
RNA editing. Nat Neurosci 16:1518–1522. doi: 10.1038/nn.3539

Li, TH and Schmid CW (2001) Differential stress induction of individual Alu loci:
implications for transcription and retrotransposition. Gene 276:135–141.

Li, W, Jin Y, Prazak L, et al (2012) Transposable elements in TDP-43-mediated
neurodegenerative disorders. PLOS One e44099:1–10.

Li Z, Tian Y, Tian N, et al (2015) Aberrant alternative splicing pattern of ADAR2
downregulates adenosine-to-inosine editing in glioma. Oncol Rep 33:2845–52. doi:
10.3892/or.2015.3907

Lin L, Jiang P, Park JW, et al (2016) The contribution of Alu exons to the human proteome.
Genome Biol 17:15.

Lin L, Shen S, Tye A, et al (2008) Diverse Splicing Patterns of Exonized Alu Elements in
Human Tissues. PLoS Genet 4:e1000225. doi: 10.1371/journal.pgen.1000225

Lin MT, Beal MF (2006) Mitochondrial dysfunction and oxidative stress in
neurodegenerative diseases. Nature 443:787–795.

Linker SB, Marchetto MC, Narvaiza I, et al (2017) Examining non-LTR retrotransposons in
the context of the evolving primate brain. BMC Biol 15:68. doi: 10.1186/s12915-017-
0409-z

Liscovitch N, Bazak L, Levanon EY, Chechik G (2014) Positive correlation between ADAR
expression and its targets suggests a complex regulation mediated by RNA editing in the
human brain. RNA Biol 11:1447–1456. doi: 10.4161/15476286.2014.992286

López E, Casasnovas C, Giménez J, et al (2015) Characterization of Alu and recombination-
associated motifs mediating a large homozygous SPG7 gene rearrangement causing
hereditary spastic paraplegia. Neurogenetics 16:97–105. doi: 10.1007/s10048-014-0429-
6

Lu B (2009) Mitochondrial Dynamics and Neurodegeneration. Curr Neurol Neurosci Rep 9:212.

Luco RF (2016) Retrotransposons jump into alternative-splicing regulation via a long noncoding RNA. Nat Struct Mol Biol 23:952–954. doi: 10.1038/nsmb.3318

Lutz MW, Crenshaw DG, Saunders AM, Roses AD (2010) Genetic variation at a single locus and age of onset for Alzheimer's disease. Alzheimer's Dement 6:125–131.

Lyall DM, Harris SE, Bastin ME, et al (2014) Alzheimer's disease susceptibility genes APOE and TOMM40, and brain white matter integrity in the Lothian Birth Cohort 1936. Neurobiol Aging. doi: 10.1016/j.neurobiolaging.2014.01.006

Maas S, Kawahara Y, Tamburro KM, Nishikura K (2006) A-to-I RNA editing and human disease. RNA Biol 3:1–9.

Maden M (2007) Retinoic acid in the development, regeneration and maintenance of the nervous system. Nat Rev Neurosci 8:755–765. doi: 10.1038/nrn2212

Madsen E, Gitlin JD (2007) Copper and Iron Disorders of the Brain. Annu Rev Neurosci 30:317–337. doi: 10.1146/annurev.neuro.30.051606.094232

Mager F, Gessmann D, Nussberger S, Zeth K (2011) Functional refolding and characterization of two Tom40 isoforms from human mitochondria. J Membr Biol 242:11–21.

Mahley RW, Rall SC (2000) Apolipoprotein E: far more than a lipid transport protein. Annu Rev Genomics Hum Genet 1:507–537. doi: 10.1146/annurev.genom.1.1.507

Mameli E, Lepori MB, Chiappe F, et al (2015) Wilson's disease caused by alternative splicing and Alu exonization due to a homozygous 3039-bp deletion spanning from intron 1 to exon 2 of the ATP7B gene. 569:276–279.

Mariner PD, Walters RD, Espinoza CA, et al (2008) Human Alu RNA is a modular transacting repressor of mRNA transcription during heat shock. Mol Cell 29:499–509.

Martin LJ (2012) Biology of mitochondria in neurodegenerative diseases. Prog Mol Biol Transl Sci 107:355–415. doi: 10.1016/B978-0-12-385883-2.00005-9

Mattick JS, Mehler MF (2008) RNA editing, DNA recoding and the evolution of human cognition. Trends Neurosci 31:227–33. doi: 10.1016/j.tins.2008.02.003

McKhann GM, Knopman DS, Chertkow H, et al (2011) The diagnosis of dementia due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. Alzheimer's Dement. 7:263–269.

Mefford HC, Zemel M, Geraghty E, et al (2015) Intragenic deletions of ALDH7A1 in pyridoxine-dependent epilepsy caused by Alu-Alu recombination. Neurology 85:756–62. doi: 10.1212/WNL.0000000000001883

Mehler MF, Mattick JS (2007) Noncoding RNAs and RNA editing in brain development, functional diversification, and neurological disease. Physiol Rev 87:799–823.

Mirza AH, Kaur S, Brorsson CA, Pociot F (2014) Effects of GWAS-associated genetic variants on lncRNAs within IBD and T1D candidate loci. PLoS One. doi: 10.1371/journal.pone.0105723

Mita P, Boeke JD (2016) How retrotransposons shape genome regulation. Curr Opin Genet Dev 37:90–100. doi: 10.1016/j.gde.2016.01.001

Mittal K, Mani RJ, Katare DP (2016) Type 3 Diabetes: Cross Talk between Differentially Regulated Proteins of Type 2 Diabetes Mellitus and Alzheimer's Disease. Sci Rep 6:25589. doi: 10.1038/srep25589

Morais S, Bastos-Ferreira R, Sequeiros J, Alonso I (2016) Genomic mechanisms underlying PARK2 large deletions identified in a cohort of patients with PD. Neurol Genet 2:e73. doi: 10.1212/NXG.0000000000000073

Muotri AR, Chu VT, Marchetto MCN, et al (2005) Somatic mosaicism in neuronal precursor

cells mediated by L1 retrotransposition. Nature 435:903–910.

Muotri AR, Marchetto MCN, Coufal NG, et al (2010) L1 retrotransposition in neurons is
modulated by MeCP2. Nature 468:443–446. doi: 10.1038/nature09544

Mustajoki S, Ahola H, Mustajoki P, Kauppinen R (1999) Insertion of Alu element
responsible for acute intermittent porphyria. Hum Mutat 13:431–8.

Neeman Y, Levanon EY, Jantsch MF, Eisenberg E (2006) RNA editing level in the mouse is
determined by the genomic repeat repertoire. RNA 12:1802–9. doi: 10.1261/rna.165106

Neven KY, Piola M, Angelici L, et al (2016) Repetitive element hypermethylation in
multiple sclerosis patients. BMC Genet 17:84. doi: 10.1186/s12863-016-0395-0

Nishikura K (2016) A-to-I editing of coding and non-coding RNAs by ADARs. Nat Rev Mol
Cell Biol 17:83–96. doi: 10.1038/nrm.2015.4

Norris J, Fan D, Aleman C, et al (1995) Identification of a new subclass of Alu DNA repeats
which can function as estrogen receptor-dependent transcriptional enhancers. J Biol
Chem 270:22777–82. doi: 10.1074/JBC.270.39.22777

Nozu K, Iijima K, Ohtsuka Y, et al (2014) Alport syndrome caused by a COL4A5 deletion
and exonization of an adjacent AluY. Mol Genet genomic Med 2:451–453.

Okubo M, Horinishi A, Saito M, et al (2007) A novel complex deletion–insertion mutation
mediated by Alu repetitive elements leads to lipoprotein lipase deficiency. Mol Genet
Metab 92:229–233. doi: 10.1016/J.YMGME.2007.06.018

Oliver KR, Greene WK (2011) Mobile DNA and the TE-Thrust hypothesis: supporting
evidence from the primates. Mob DNA 2:8. doi: 10.1186/1759-8753-2-8

Ottesen EW, Seo J, Singh NN, Singh RN (2017) A Multilayered Control of the Human
Survival Motor Neuron Gene Expression by Alu Elements. Front Microbiol 8:2252. doi:
10.3389/fmicb.2017.02252

Pal S, Tyler JK (2016) Epigenetics and aging. Sci Adv 2:e1600584–e1600584. doi:
10.1126/sciadv.1600584

Pandolfo M (2006) Friedreich Ataxia: Detection of GAA Repeat Expansions and Frataxin
Point Mutations. In: Congenital Heart Disease. Humana Press, New Jersey, pp 197–216

Paquola ACM, Erwin JA, Gage FH (2016) Insights into the role of somatic mosaicism in the
brain. Current Opinion in Systems Biology 1:90-94.

Payer LM, Steranka JP, Yang WR, et al (2017) Structural variants caused by Alu insertions
are associated with risks for many human diseases. Proc Natl Acad Sci USA
114:E3984–E3992. doi: 10.1073/pnas.1704117114

Pehlivan D, Beck CR, Okamoto Y, et al (2016) The role of combined SNV and CNV burden
in patients with distal symmetric polyneuropathy. Genet Med 18:443–451. doi:
10.1038/gim.2015.124

Petschner P, Gonda X, Baksa D, et al (2017) Genes linking mitochondrial function, cognitive
impairment and depression are associated with endophenotypes serving precision
medicine. Neuroscience. doi: 10.1016/J.NEUROSCIENCE.2017.09.049

Picardi E, D'Erchia AM, Lo Giudice C, Pesole G (2017a) REDIportal: a comprehensive
database of A-to-I RNA editing events in humans. Nucleic Acids Res 45:D750–D757.
doi: 10.1093/nar/gkw767

Picardi E, Horner DS, Pesole G (2017b) Single-cell transcriptomics reveals specific RNA
editing signatures in the human brain. RNA 23:860–865. doi: 10.1261/rna.058271.116

Picardi E, Manzari C, Mastropasqua F, et al (2015) Profiling RNA editing in human tissues:
towards the inosinome Atlas. Sci Rep 5:14941. doi: 10.1038/srep14941

Polak P, Domany E (2006) Alu elements contain many binding sites for transcription factors
and may play a role in regulation of developmental processes. BMC Genomics 7:133.
doi: 10.1186/1471-2164-7-133

Prendergast JGD, Chambers E V, Semple CAM (2014) Sequence-level mechanisms of

human epigenome evolution. Genome Biol Evol 6:1758–71. doi: 10.1093/gbe/evu142

Prudencio M, Gonzales PK, Cook CN, et al (2017) Repetitive element transcripts are
elevated in the brain of C9orf72 ALS/FTLD patients. Hum Mol Genet 26:3421–3431.
doi: 10.1093/hmg/ddx233

Prüfer K, Munch K, Hellmann I, et al (2012) The bonobo genome compared with the
chimpanzee and human genomes. Nature 486:nature11128. doi: 10.1038/nature11128

Reddy PH (2011) Abnormal tau, mitochondrial dysfunction, impaired axonal transport of
mitochondria, and synaptic deprivation in Alzheimer's disease. Brain Res. 1415:136–
148.

Reynolds WF, Rhees J, Maciejewski D, et al (1999) Myeloperoxidase Polymorphism Is
Associated with Gender Specific Risk for Alzheimer's Disease. Exp Neurol 155:31–41.
doi: 10.1006/exnr.1998.6977

Rezin GT, Amboni G, Zugno AI, et al (2009) Mitochondrial Dysfunction and Psychiatric
Disorders. Neurochem Res 34:1021–1029. doi: 10.1007/s11064-008-9865-8

Rice GI, Kasher PR, Forte GMA, et al (2012) Mutations in ADAR1 cause Aicardi-Goutières
syndrome associated with a type I interferon signature. Nat Genet 44:1243–8. doi:
10.1038/ng.2414

Roger AJ, Muñoz-Gómez SA, Kamikawa R (2017) The Origin and Diversification of
Mitochondria. Curr Biol 27:R1177–R1192. doi: 10.1016/j.cub.2017.09.015

Roses AD (2010) An inherited variable poly-T repeat genotype in TOMM40 in Alzheimer
disease. Arch Neurol 67:536–541.

Roses AD, Sundseth S, Saunders A, et al (2016a) Understanding the genetics of APOE and
TOMM40 and role of mitochondrial structure and function in clinical pharmacology of
Alzheimer's disease. Alzheimers Dement 12:687–694. doi: 10.1016/j.jalz.2016.03.015

Roses AD, Akkari PA, Chiba-Falek O, et al (2016b) Structural variants can be more
informative for disease diagnostics, prognostics and translation than current SNP
mapping and exon sequencing. Expert Opin Drug Metab Toxicol 12:135–147. doi:
10.1517/17425255.2016.1133586

Roses AD, Lutz MW, Amrine-Madsen H, et al (2010) A TOMM40 variable-length
polymorphism predicts the age of late-onset Alzheimer's disease. Pharmacogenomics J
10:375–384.

Ross CA, Poirier MA (2004) Protein aggregation and neurodegenerative disease. Nat Med
S10-17.

Rossi L, Lombardo MF, Ciriolo MR, Rotilio G (2004) Mitochondrial Dysfunction in
Neurodegenerative Diseases Associated with Copper Imbalance. Neurochem. Res.
29:493–504.

Rugarli EI, Langer T (2012) Mitochondrial quality control: a matter of life and death for
neurons. EMBO J 31:1336–1349.

Rybak-Wolf A, Stottmeister C, Glazar P, et al (2015) Circular RNAs in the Mammalian
Brain Are Highly Abundant, Conserved, and Dynamically Expressed. Mol Cell 58: 870–
885.

Sakurai M, Ueda H, Yano T, et al (2014) A biochemical landscape of A-to-I RNA editing in
the human brain transcriptome. Genome Res 24:522–34. doi: 10.1101/gr.162537.113

Saunders AM, Schmader K, Breitner JC, et al (1993) Apolipoprotein E epsilon 4 allele
distributions in late-onset Alzheimer's disease and in other amyloid-forming diseases.
Lancet (London, England) 342:710–1. doi: 10.1016/0140-6736(93)91709-U

Schanze D, Neubauer D, Cormier-Daire V, et al (2014) Deletions in the 3′ Part of the *NFIX*
Gene Including a Recurrent Alu-Mediated Deletion of Exon 6 and 7 Account for
Previously Unexplained Cases of Marshall-Smith Syndrome. Hum Mutat 35:1092–
1100. doi: 10.1002/humu.22603

Schmauss C, Howe JR (2002) RNA editing of neurotransmitter receptors in the mammalian brain. Sci STKE 2002:pe26. doi: 10.1126/stke.2002.133.pe26

Schneider I, Kugel H, Redlich R, et al. (2017) Association of Serotonin Transporter Gene AluJb Methylation with Major Depression, Amygdala Responsiveness, 5-HTTLPR/rs25531 Polymorphism, and Stress. Neuropsychopharmacology *online early* 1–9. doi:10.1038/npp.2017.273

Schollen E, Keldermans L, Foulquier F, et al (2007) Characterization of two unusual truncating PMM2 mutations in two CDG-Ia patients. Mol Genet Metab 90:408–413. doi: 10.1016/j.ymgme.2007.01.003

Schwahn U, Lenzner S, Dong J, et al (1998) Positional cloning of the gene for X-linked retinitis pigmentosa 2. Nat Genet 19:327–32. doi: 10.1038/1214

Schwarz TL (2013) Mitochondrial trafficking in neurons. Cold Spring Harb Perspect Biol 5:a011304.

Seeley WW (2017) Mapping Neurodegenerative Disease Onset and Progression. Cold Spring Harb Perspect Biol 9:a023622. doi: 10.1101/cshperspect.a023622

Selkoe DJ (2000) Toward a comprehensive theory for Alzheimer's disease. Hypothesis: Alzheimer's disease is caused by the cerebral accumulation and cytotoxicity of amyloid beta-protein. Ann N Y Acad Sci 924:17–25. doi: 10.1111/j.1749-6632.2000.tb05554.x

Shen S, Lin L, Cai JJ, et al (2011) Widespread establishment and regulatory impact of Alu exons in human genes. Proc Natl Acad Sci 108:2837–2842.

Shiota T, Imai K, Qiu J, et al (2015) Molecular architecture of the active mitochondrial protein gate. Science 349:1544–1548.

Shpyleva S, Melnyk S, Pavliv O, et al (2017) Overexpression of LINE-1 Retrotransposons in Autism Brain. Mol Neurobiol 1–10. doi: 10.1007/s12035-017-0421-x

Skovronsky DM, Doms RW, Lee VM (1998) Detection of a novel intraneuronal pool of insoluble amyloid beta protein that accumulates with time in culture. J Cell Biol 141:1031–9.

Smalheiser NR (2014) The RNA-centred view of the synapse: non-coding RNAs and synaptic plasticity. Philos Trans R Soc B Biol Sci 369:20130504–20130504. doi: 10.1098/rstb.2013.0504

Soibam B (2017) Super-lncRNAs: identification of lncRNAs that target super-enhancers via RNA:DNA:DNA triplex formation. RNA 23:1729–1742.

Steen E, Terry BM, Rivera EJ, et al (2005) Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease--is this type 3 diabetes? J Alzheimers Dis 7:63–80.

Strittmatter WJ, Saunders AM, Schmechel D, et al (1993) Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. Proc Natl Acad Sci 90:1977–1981. doi: 10.1073/pnas.90.5.1977

Sun Z, Wu Y, Ordog T, et al (2014) Aberrant signature methylome by DNMT1 hot spot mutation in hereditary sensory and autonomic neuropathy 1E. Epigenetics 9:1184–1193. doi: 10.4161/epi.29676

Swerdlow RH (2017) Mitochondria and Mitochondrial Cascades in Alzheimer's Disease. J Alzheimer's Dis Preprint:1–14. doi: 10.3233/JAD-170585

Swerdlow RH, Burns JM, Khan SM (2010) The Alzheimer's disease mitochondrial cascade hypothesis. J Alzheimer's Dis JAD 20:265.

Szpakowski S, Sun X, Lage JM, et al (2009) Loss of epigenetic silencing in tumors preferentially affects primate-specific retroelements. Gene 448:151–67. doi: 10.1016/j.gene.2009.08.006

Takahashi RH, Milner TA, Li F, et al (2002) Intraneuronal Alzheimer abeta42 accumulates in multivesicular bodies and is associated with synaptic pathology. Am J Pathol 161:1869–

79.

Tan B, Huang H, Lam R, Soong T (2009) Dynamic regulation of RNA editing of ion
channels and receptors in the mammalian nervous system. Mol Brain 2:13. doi:
10.1186/1756-6606-2-13

Tan MH, Li Q, Shanmugam R, et al (2017) Dynamic landscape and regulation of RNA
editing in mammals. Nature 550:249–254. doi: 10.1038/nature24041

Tarallo V, Hirano Y, Gelfand BD, et al (2012) DICER1 loss and Alu RNA induce age-related
macular degeneration via the NLRP3 inflammasome and MyD88. Cell 149:847–59. doi:
10.1016/j.cell.2012.03.036

Tariq A, Jantsch MF (2012) Transcript diversification in the nervous system: A to I RNA
editing in CNS function and disease development. Front Neurosci 1–13. doi:
10.3389/fnins.2012.00099

Taşkesen M, Collin GB, Evsikov A V, et al (2012) Novel Alu retrotransposon insertion
leading to Alström syndrome. Hum Genet 131:407–13. doi: 10.1007/s00439-011-1083-9

Thongsroy J, Patchsung M, Mutirangura A (2017) The association between Alu
hypomethylation and severity of type 2 diabetes mellitus. Clin Epigenetics 9:93. doi:
10.1186/s13148-017-0395-6

Töhönen V, Katayama S, Vesterlund L, et al (2015) Novel PRD-like homeodomain
transcription factors and retrotransposon elements in early human development. Nat
Commun 6:8207. doi: 10.1038/ncomms9207

Tsirigos A, Rigoutsos I (2009) Alu and B1 Repeats Have Been Selectively Retained in the
Upstream and Intronic Regions of Genes of Specific Functional Classes. PLOS Comput
Biol 5:e1000610. doi: 10.1371/JOURNAL.PCBI.1000610

Udaka T, Okamoto N, Aramaki M, et al (2007) An Alu retrotransposition-mediated deletion
of CHD7 in a patient with CHARGE syndrome. Am J Med Genet Part A 143:721–726.
doi: 10.1002/ajmg.a.31441

Ullu E, Tschudi C (1984) Alu sequences are processed 7SL RNA genes. Nature 312:171–
172. doi: 10.1038/312171a0

Van Den Heuvel MP, Sporns O (2013) Network hubs in the human brain. Trends Cogn Sci
17:683–696. doi: 10.1016/j.tics.2013.09.012

Vandewalle J, Bauters M, Van Esch H, et al (2013) The mitochondrial solute carrier
SLC25A5 at Xq24 is a novel candidate gene for non-syndromic intellectual disabilty.
Hum Genet 132:1177-1185.

Vansant G, Reynolds WF (1995) The consensus sequence of a major Alu subfamily contains
a functional retinoic acid response element. Proc Natl Acad Sci U S A 92:8229–33.

Varizhuk AM, Sekridova AV, Tankevich MV, et al (2016) Conformational polymorphism of
G-rich fragments of DNA Alu-repeats. II. the putative role of G-quadruplex structures in
genomic rearrangements. Biomeditsinskaya Khimiya 62:630–637. doi:
10.18097/pbmc20166206630

Varshney D, Vavrova-Anderson J, Oler AJ, et al (2015) SINE transcription by RNA
polymerase III is suppressed by histone methylation but not by DNA methylation.

Vassetzky NS, Ten OA, Kramerov DA (2003) B1 and related SINEs in mammalian genomes.
Gene 319:149–60.

Viader A, Golden JP, Baloh RH, et al (2011) Schwann cell mitochondrial metabolism
supports long-term axonal survival and peripheral nerve function. J Neurosci 31:10128–
40. doi: 10.1523/JNEUROSCI.0884-11.2011

Viader A, Sasaki Y, Kim S, et al (2013) Aberrant Schwann cell lipid metabolism linked to
mitochondrial deficits leads to axon degeneration and neuropathy. Neuron 77:886–98.
doi: 10.1016/j.neuron.2013.01.012

Wang W, Li L, Lin W, et al (2013) The ALS disease-associated mutant TDP-43 impairs

mitochondrial dynamics and function in motor neurons. Hum Mol Genet 22:4706–4719.

Wang W, Wang WH, Azadzoi KM, et al. (2016) Alu RNA accumulation in hyperglycemia

augments oxidative stress and impairs eNOS and SOD2 expression in endothelial cells.

Mol Cell Endrocrinol 426:91–100.

Weissmann D, van der Laan S, Underwood MD, et al (2016) Region-specific alterations of

A-to-I RNA editing of serotonin 2c receptor in the cortex of suicides with major

depression. Transl Psychiatry 6:e878. doi: 10.1038/tp.2016.121

Wimmer K, Callens T, Wernstedt A, Messiaen L (2011) The NF1 gene contains hotspots for

L1 endonuclease-dependent De Novo insertion. PLoS Genet. doi:

10.1371/journal.pgen.1002371

Wright A, Vissel B (2012) The essential role of AMPA receptor GluR2 subunit RNA editing

in the normal and diseased brain. Front Mol Neurosci 5:34. doi:

10.3389/fnmol.2012.00034

Wu S-J, Hsieh T-J, Kuo M-C, et al (2013) Functional regulation of Alu element of human

angiotensin-converting enzyme gene in neuron cells. Neurobiol Aging 34:1921.e1-

1921.e7. doi: 10.1016/j.neurobiolaging.2013.01.003

Zempel H, Mandelkow E (2015) Tau missorting and spastin-induced microtubule disruption

in neurodegeneration: Alzheimer disease and hereditary spastic paraplegia. Mol

Neurodegener 10:68. doi: 10.1186/s13024-015-0064-1

Zhang G, Fukao T, Sakurai S, et al (2006) Identification of Alu-mediated, large deletion-

spanning exons 2-4 in a patient with mitochondrial acetoacetyl-CoA thiolase deficiency.

Mol Genet Metab 89:222–6. doi: 10.1016/j.ymgme.2006.06.010

Zhang Y-H, Dipple KM, Vilain E, et al (2000) AluY insertion (IVS4-52ins316alu) in the

glycerol kinase gene from an individual with benign glycerol kinase deficiency. Hum

Mutat 15:316–323. doi: 10.1002/(SICI)1098-1004(200004)15:4<316::AID-

HUMU3>3.0.CO;2-9

1134

1135

1136

1137   **Figure Legends**

1138

Figure 1. Select mechanisms whereby *Alu* elements can alter gene expression and function

(also see Elbarbary et al. 2016). **A:** Sequence homology and orientation of *Alu* elements

contributes to the formation of distinct secondary structures in both DNA and RNA. DNA

Alu G-quadruplex structures can alter transcription kinetics (Varizhuk et al. 2016) and pre-

mRNA *Alu* binding forms stem-loop structures that are the primary site for A-to-I editing

(see Figure 2). **B:** Recombination of intra-gene *Alu* elements resulting in exon deletion. **C:**

Exonification of intronic *Alus* contributing to the production of alternative mRNAs. **D:**

Environmental or traumatic stress cascades resulting in increased expression of *Alu* RNAs

that contribute to inflammation (Li and Schmid 2001; Tarallo et al. 2012; Hunter et al. 2015;

Lapp et al. 2016), the disruption of global gene transcription through Pol II binding (Mariner

et al. 2008), and an increase of H3K9 histone methylation that alters *Alu* epigenetic pathways

(Varshney et al. 2015; Lapp and Hunter 2016; Larsen et al. 2017).

1151

Figure 2. Intronic *Alu* elements located in close proximity (**A**) can bind to each other within

pre-mRNAs (**B**) thereby producing a stable stem-loop secondary structure that is the primary

substrate for A-to-I editing in primates. ADAR proteins bind to pre-mRNA *Alu* structures

(**C**) and convert adenosine residues to inosine. If occurring in coding regions, the translation

machinery then interprets the inosine residues as guanosine and this can contribute to amino

acid changes and alternative protein conformations (**D**).

1158

Figure 3. Deleterious *Alu* activity operating on mitonuclear genes can disrupt mitochondrial

function in the CNS and contribute to a number of diseased phenotypes (see Table 1). The

type and severity of associated neurological and neurodegenerative disorders depends on the

deleterious *Alu* mechanism of action, the mitonuclear gene pathways involved, the time or

developmental stage of induction, level or severity of traumatic stress, and tissue specificity

(see Larsen et al. 2017). If operating across the suite of mitonuclear genes through epigenetic

pathways, the mechanism helps to explain the origin of incipient mitochondrial stress and

CNS connectome destabilization that is observed across the spectrum of neurological and

neurodegenerative disorders.

1168

1169

1170

A.

B.

C.

D.

Figure 1.

A. DNA

Alu   *Alu*

B. pre-mRNA

C. mRNA

Alu   *Alu*

GUGAGACUCCIUCUCNNNUCUCUUAUUUUUCUG

D. Protein

| V | R | L | R | L | X | S | L | I | F | L |

 Figure 2.

Deleterious *Alu* Mechanisms

Exonization
Alternative splicing
Stress-induced activation
Dysregulation of A-to-I editing
Non-homologous recombination
Dysregulation of DNA methylation and histone modification

↓

Mitonuclear Genes (n = ~2,000)

Variable impact on mitochondrial function

Normal   Abnormal   Dysfunctional

↓

Variable Disruption of Key Mitochondrial Processes

Apoptosis
Mitophagy
ROS production
Calcium buffering
Fatty acid processing
Inflammatory response
Pre-protein importation
Iron, zinc, copper processing
Electron transport chain function

↓

Spectrum of Sporadic Disease

Variable impact on CNS function and connectome stability

↓

Epilepsy   ALS
Depression   Alzheimer's Disease
Bipolar Disorder   Parkinson's Disease
Schizophrenia   Pick's Disease
PTSD
CTE

1186

1187

1188

1189

1190   **Figure 3.**

Table 1. Genes associated with neurological and neurodegenerative disorders wherein deleterious *Alu* activity has been documented experimentally or is hypothesized to disrupt gene function. Gene names in bold identify genes essential for mitochondrial function and transport and/or are associated with mitochondrial abnormalities (sensu Dawson et al. 1995, Calvo et al. 2015, Zempel and Mandelkow 2015, Bhattachargee et al. 2016, Chong-Chong et al. 2016, Checler et al. 2017, Johnson et al. 2017, Wang et al. 2013). For additional *Alu* associated diseases see Hancks and Kazazian (2016) and Payer et al. (2017). Asterisks identify genes where mutations result in dysregulation of *Alu* elements.

Gene Name	Disorder	*Alu* Mechanism of Disruption	Reference
ABCD1	Adrenoleukodystrophy	Deletion events	Kutsche et al. 2002
ACAT1 (T2)	Mitochondrial acetoacetyl-CoA thiolase deficiency	Deletion event	Zhang et al. 2006
ACE	Alzheimer's disease	Insertion events	Wu et al. 2013
ADAR2	Glioma	Exonization	Li et al. 2015
ALDH7A1	Pyridoxine-dependent epilepsy	Recombination	Mefford et al. 2015
ALMS1	Alström syndrome	Insertion event	Taşkesen et al. 2012
APOB	Hypobetalipoproteinemia	Recombination	Huang et al. 1989
ATP5J	Alzheimer's disease	Duplication	Antonell et al. 2012
ATP7A	Menkes disease	Insertion event	Gu et al. 2007; Bhattacharjee et al. 2016
ATP7B	Wilson's disease	Alternative splicing	Mameli et al. 2015
C9orf72	ALS, FTLD	Loss of epigenetic control, elevated *Alu* transcripts	Prudencio et al. 2017
CHD7	CHARGE syndrome	Deletion	Udaka et al. 2007
CLN3	Batten disease	Deletion	Lerner et al. 1995
COL4A5	Alport syndrome	Deletion and exonization	Nozu et al. 2014
DICER1*	Age-related macular degeneration	*Alu* RNA build-up with reduced DICER1 activity	Kaneko et al. 2011; Kim et al. 2014
FXN	Friedreich's Ataxia	*Alu* repeat expansion, alternative splicing events	Pandolfo 2006
GK	Glycerol kinase deficiency	Insertion event	Zhang et al. 2000
GLA	Fabry disease	Deletion event	Dobrovolny et al. 2011
HPRT	Lesch-Nyhan disease	Recombination	Brooks et al. 2001
HMBS	Acute intermittent prophyria	Insertion event	Mustajoki et al. 1999
LPL	Lipoprotein lipase deficiency	Complex deletion-insertion	Okubo et al. 2007
MFN2 (CMT2a)	Charcot-Marie-Tooth type 2A	Copy number variants	Pehlivan et al. 2016
MPO	Alzheimer's disease	*Alu* hormone response variant; estrogen dysregulation	Reynolds et al. 1999

NDUFS2	Leigh Syndrome	Exonization	Larsen et al. 2017
NF1	Neurofibromatosis type I	Deletion and chimeric gene fusion	Wimmer et al. 2011; Ferrari et al. 2017
NFIX	Marshall-Smith syndrome	Deletions	Schanze et al. 2014
OPA1	Autosomal Dominant Optic Atrophy	Alternative splicing events	Gallus et al. 2010
PARK2	Parkinson's disease	Recombination	Morais et al. 2016
PARK7 (DJ-1)	Parkinson's disease	Deletion	Bonifati et al. 2002
PDHA1	Pyruvate Dehydrogenase Deficiency	Exonization	Larsen et al. 2017
PIGL	CHIME syndrome	Deletion	Knight Johnson et al. 2017
PMM2	Congenital disorders of glycosylation type Ia	Complex deletion	Schollen et al. 2007
POMT1	Walker Warburg syndrome	Insertion	Bouchet et al. 2007
PSEN1	Alzheimer's disease	Deletion	Le Guennec et al. 2017
PXMP2 (PMP22)	Charcot-Marie-Tooth type 2A	*Alu-Alu*-mediated rearrangement	Choi et al. 2011; Gu et al. 2015
RP2 (NUDT19)	X-linked retinitis pigmentosa	*Alu*-L1 recombination	Schwahn et al. 1998, Jiang et al. 2017
SLC6A4	Depression, reduced hippocampal volume	Altered promoter methylation	Dannlowski et al. 2014
SLC25AC	Intellectual disability	Deletion	Vandewalle et al. 2013
SLC30A6	Alzheimer's disease, Dementia, ALS	Gene fusion event	Boone et al. 2014
SMN1	Spinal muscular atrophy	Exonization, deletion events, circularization	Ottesen et al. 2017
SOD2	Hyperglycemia	Repressed expression	Wang et al. 2016
SOX10	Waardenburg syndrome type 4	Deletion	Bondurand et al. 2012
SPAST	Autosomal-dominant spastic paraplegia 4	Deletions, CNVs, gene fusion events	Boone et al. 2014
SPG7	Hereditary spastic paraplegia	Deletion, recombination	Arnoldi et al. 2008; López et al. 2015
SPG11	Hereditary spastic paraplegia	Deletion	Conceição et al. 2012
STAU1	Myotonic Dystrophy Type 1	Alternative splicing regulation	Bondy-Chorney et al. 2016
TDP-43	ALS, frontotemporal lobar degeneration	Transposable element dysregulation	Li et al. 2012
TOMM40	Late-Onset Alzheimer's Disease	*Alu* repeat expansion, putative alternative splicing events	Larsen et al. 2017
TRIM37	Mulibrey nanism	Deletion events	Jobic et al. 2017