Seasonal temperature variation influences climate suitability for dengue,

chikungunya, and Zika transmission

John H. Huber[1, 2], Marissa L. Childs[3], Jamie M. Caldwell[2], Erin A. Mordecai[2*]

[1]Department of Applied and Computational Mathematics and Statistics, University

of Notre Dame, Notre Dame, Indiana, USA

[2]Department of Biology, Stanford University, Stanford, California, USA

[3]Emmett Interdisciplinary Program in Environment and Resources, Stanford

University, Stanford, California, USA

*Corresponding Author: emordeca@stanford.edu

Short title: Seasonal climate affects vector transmission

ABSTRACT

Dengue, chikungunya, and Zika virus epidemics transmitted by *Aedes aegypti*

mosquitoes have recently (re)emerged and spread throughout the Americas,

Southeast Asia and the Pacific Islands, and elsewhere. Understanding how

environmental conditions affect epidemic magnitude and duration is critical for

predicting and responding to the geographic and seasonal spread of disease. Here,

we develop the first dynamic disease transmission model for dengue virus and

Aedes aegypti mosquitoes that integrates mechanistic, empirically parameterized,

and independently validated mosquito and virus trait thermal responses under

seasonally varying temperatures. With the model, we examine the influence of

variation in seasonal temperature regime on epidemic dynamics. We find that at

both constant and seasonally varying temperatures, warmer temperatures at the

start of epidemics promote more rapid epidemics due to faster burnout of the

susceptible population. By contrast, intermediate temperatures (~24°C) at epidemic

onset produced the largest epidemics in a seasonally varying temperature regime.

When seasonal temperature variation was low, 25-35°C annual average

temperatures produced the largest epidemics, but this range shifts to cooler

temperatures as seasonal temperature variation increases. Tropical and sub-

tropical cities such as Rio de Janeiro, Fortaleza, and Salvador, Brazil; Cali, Cartagena,

and Baranquilla, Colombia; Delhi, India; Guangzhou, China; and Manila, Philippines

have mean annual temperatures and seasonal temperature ranges that produce the

largest epidemics. However, more temperate cities like Shanghai, China had high

epidemic suitability because large seasonal variation offset moderate annual

average temperatures. By accounting for seasonal variation in temperature, our

model provides a baseline for mechanistically understanding environmental

suitability for virus transmission by *Aedes aegypti*. Overlaying the impact of human

activities and socioeconomic factors onto this mechanistic temperature-dependent

framework is critical for understanding likelihood and magnitude of outbreaks.

NON-TECHNICAL SUMMARY (150-200 WORDS)

Mosquito-borne viruses like dengue, Zika, and chikungunya have recently caused

large epidemics that are partly driven by temperature. Using a mathematical model

built from laboratory experimental data for *Aedes aegypti* mosquitoes and dengue

virus, we examine the impact of variation in seasonal temperature regimes on

epidemic size and duration. At constant temperatures, both low and high

temperatures (20°C and 35°C) produce small epidemics, while intermediate

temperatures like 25°C and 30°C produce much larger epidemics. In seasonally

varying temperature environments, epidemics peak more rapidly at higher starting

temperatures. Intermediate starting temperatures produce the largest epidemics.

Seasonal mean temperatures of 25-35°C are most suitable for large epidemics when

seasonality is low, but in more variable seasonal environments epidemic suitability

peaks at lower annual average temperatures. Tropical and sub-tropical cities have

the highest temperature suitability for epidemics, but more temperate cities with

high seasonal variation also have the potential for very large epidemics.

INTRODUCTION

Over the last 30-40 years, arboviral outbreaks have dominated the public health

landscape globally (1). These viruses, most notably chikungunya (CHIKV), dengue

(DENV), and Zika (ZIKV), can cause symptoms ranging from rash, arthralgia, and

fever to hemorrhagic fever (DENV), long-term arthritis (CHIKV), Guillain-Barré

syndrome and microcephaly (ZIKV) (2–4). DENV, which historically spread

worldwide along shipping routes (5), places 3.97 billion individuals at risk

worldwide (6) and causes an estimated 390 million cases annually, including 96

million symptomatic cases (7). CHIKV was introduced into the Americas in

December 2013 after an outbreak in St. Martin Island (8). Since then, autochthonous

transmission has been reported in 45 countries (9), and 1.3 billion people

worldwide are at risk of contracting CHIKV (10). More recently, the ZIKV epidemic

in the Americas captured global attention after the World Health Organization

(WHO) designated it a Public Health Emergency of International Concern in

February 2016 in response to its association with neurological disorders. Following

the first reported case in Brazil in May 2015, ZIKV has spread to 48 countries and

territories where it is transmitted autochthonously (11). Because DENV, CHIKV, and

ZIKV are mostly transmitted by *Aedes aegypti* mosquitoes, they may have similar

geographic distributions and risk factors.

Informed public health decisions to limit the spread and magnitude of these

arboviral epidemics depend on a robust understanding of transmission dynamics.

One mechanistic modeling framework, the Susceptible – Infected – Recovered (SIR)

model, has been implemented successfully to model the dynamics of outbreaks of

influenza, measles, and vector-borne diseases such as CHIKV and ZIKV (12–14). This

approach tracks virus population dynamics by compartmentalizing individuals by

their state in an epidemic (i.e., Susceptible (S), Infected (I), Recovered (R)). This

framework can be extended to include additional compartments, such as a latency

stage, or to incorporate the dynamics of the mosquito population for vector

transmission.

Arbovirus dynamics are strikingly seasonal and geographically restricted to

relatively warm climates (15). This arises because several life history traits of the

mosquitoes that transmit DENV, CHIKV, and ZIKV are strongly influenced by

temperature and seasonality (16–23). For simplicity, many existing models assume

static life history traits (14), and those that address seasonal forcing tend to

incorporate sinusoidal variation as a single transmission parameter, β (24).

However, decades of experimental work have demonstrated strongly nonlinear

(often unimodal) relationships between mosquito and pathogen traits and

temperature that are not well captured in a single sinusoidal forcing function (25).

Efforts by Yang et al. (26,27) addressed the need to include seasonal variation by

adopting an SEI-SEIR compartmental framework with time-varying entomological

parameters and fitting the model to DENV incidence data in Campinas, Brazil.

Here, we expand upon an SEI-SEIR model (26,27) to incorporate nonlinear,

temperature-dependent vector parameters (25). We use the model to evaluate the

effect of temperature on epidemic length and size, producing outputs that are more

directly comparable to observed epidemics than static measures like R_0 (28).

Importantly, we move beyond mean temperature to examine the impact of seasonal

temperature variation on epidemics, by allowing entomological parameters to vary

seasonally with temperature. Specifically, we use the model to ask: (1) How does

final epidemic size vary across constant temperatures? (2) Under seasonally varying

temperatures, how does the temperature at the start of the epidemic affect the final

epidemic size and duration? (3) How do temperature mean and seasonal range

interact to determine epidemic size? (4) Based on the model, which geographic

locations have high epidemic suitability based on climate?

METHODS

Model

Model Framework

We adopted an SEI-SEIR compartmental modeling framework to simulate arboviral

transmission by the *Aedes aegypti* vector (Fig. 1). We introduced temperature-

dependence into the model by using fitted thermal response curves for the

mosquito life history traits provided by Mordecai et al. (25). The full model is:

$$\frac{dS_V}{dt} = EFD(T) * pEA(T) * MDR(T) * N_V * \left(1 - \frac{N_V}{K(T)}\right) - \left(a(T) * pMI(T) * \frac{I_H}{N_H} + \mu(T)\right) * S_V \qquad (1)$$

$$\frac{dE_V}{dt} = a(T) * pMI(T) * \frac{I_H}{N_H} * S_V - \left(PDR(T) + \mu(T)\right) * E_V \qquad (2)$$

$$\frac{dI_V}{dt} = PDR(T) * E_V - \mu(T) * I_V \qquad (3)$$

$$\frac{dS_H}{dt} = -a(T) * b(T) * \frac{I_V}{N_H} * S_H \qquad (4)$$

$$\frac{dE_H}{dt} = a(T) * b(T) * \frac{I_V}{N_H} * S_H - \delta * E_H \qquad (5)$$

$$\frac{dI_H}{dt} = \delta * E_H - \eta * I_H \qquad (6)$$

$$\frac{dR_H}{dt} = \eta * I_H \tag{7}$$

Fig 1 Caption. Compartmental model of transmission. S_H, E_H, I_H, and R_H represent

the susceptible, exposed (or latent), infectious, and recovered segments of the

human population, respectively. Likewise, S_V, E_V, and I_V represent the susceptible,

exposed (or latent), and infectious segments of the mosquito population. Solid

arrows signify the directionality of transition from one compartment to the next,

and dashed arrows indicate the directionality of transmission.

The SEI portion of the model describes the vector population, where S_V

represents the number of susceptible mosquitoes, E_V is the number of mosquitoes in

the latency stage, and I_V is the number of infectious mosquitoes. We assumed that

Aedes aegypti mosquitoes remain infectious until they die. In equations 1-3, *EFD(T)*

is the number of eggs laid per female per day, *pEA(T)* is the probability of mosquito

egg-to-adult survival, *MDR(T)* is the mosquito egg-to-adult development rate, N_V is

the total mosquito population at time t (i.e., $S_v + E_v + I_v$), *K(T)* is the carrying capacity

for the mosquito population, *a(T)* is the per mosquito biting rate, *pMI(T)* is the

probability of mosquito infection per bite on an infectious host, *μ(T)* is the adult

mosquito mortality rate, and *PDR(T)* is the parasite development rate. Each life

history trait of the *Aedes aegypti* mosquito is a temperature-dependent function fit

from experimental laboratory data (Table 1).

Table 1. Fitted thermal responses for *Aedes aegypti* life history traits. Traits were fit

to a Brière $[cT(T - T_0)(T_m - T)^{\frac{1}{2}}]$ or a quadratic $[c(T - T_m)(T - T_0)]$ function where

T represents temperature. T_0 and T_m are the critical thermal minimum and

maximum, respectively, and c is the rate constant. Thermal responses were fit by

(29).

Trait	Definition	Function	Fitted Parameters		
a	Biting rate (day^{-1})	Brière	c = 2.71e-04	T_{min} = 14.67	T_{max} = 41.00
EFD	Eggs laid per female per day	Brière	c = 2.08e-02	T_{min} = 14.06	T_{max} = 32.03
pEA	Probability of mosquito egg-to-adult survival	Quadratic	c = -3.36e-03	T_{min} = 7.68	T_{max} = 38.31
MDR	Mosquito egg-to-adult development rate (day^{-1})	Brière	c = 1.49e-04	T_{min} = 15.12	T_{max} = 37.67
lf	Adult mosquito lifespan (days)	Quadratic	c = -1.24e+00	T_{min} = 16.63	T_{max} = 31.85
b	Probability of mosquito infectiousness	Brière	c = 9.86e-04	T_{min} = 12.05	T_{max} = 32.79
pMI	Probability of mosquito infection	Brière	c = 5.23e-04	T_{min} = 1.51	T_{max} = 34.74
PDR	Virus extrinsic incubation rate (day^{-1})	Brière	c = 1.04e-04	T_{min} = 11.50	T_{max} = 38.97

The SEIR portion of the model describes the human population, where S_H

represents the number of susceptible individuals, E_H the number of latent (or

exposed) individuals, I_H the number of infectious individuals, and R_H the number of

recovered individuals. We assumed a static population size, N_H, that was neither

subject to births nor deaths because the human lifespan far exceeds the duration of

an epidemic. Further, we binned asymptomatic and symptomatic individuals into a

single infectious class since asymptomatic infections have been shown to transmit

DENV (30) and exhibit similar viremic profiles as symptomatic patients in CHIKV

(31). Based on previous arboviral outbreaks (32,33), we assumed that an infection

conferred long-term immunity to an individual. Thus, a previously infectious

individual entering the recovered class is protected from subsequent re-infection

for the remainder of the epidemic. In the case of dengue, where there are four

unique serotypes, we consider single-season epidemics of a single serotype. In

equations 4-7, *b(T)* is the probability of human infection per bite by an infectious

mosquito, δ^{-1} is the intrinsic incubation period, and η^{-1} is the human infectivity

period. Since human components of the transmission cycle are not seasonal, we

used constants of 5.9 days for the intrinsic incubation period, $1/\delta$, and 5.0 days for

the infectious period, $1/\eta$ (34). All temperature-independent parameter values are

given in Table 2.

Table 2. Values of temperature-invariant parameters used in the model, and their

sources.

Parameter	Definition	Value	Source
δ^{-1}	Intrinsic incubation period (days)	5.9	(34)
η^{-1}	Human infectivity period (days)	5.0	(34)
I_0^H / N	Proportion of initially infectious humans	0.0025	(34)
I_0^V / M	Proportion of initially infectious mosquitoes	0.015	(34)

M/N Ratio of mosquitoes-to-humans at 29°C   2.0   (35)

174   Further, because the lifespan of an adult mosquito is short relative to the

175   timespan of an epidemic, we allowed mosquito birth and death rates to drive

176   population dynamics. Additionally, the birth rate of susceptible mosquitoes was

177   regulated by a temperature-dependent carrying capacity, K (equation 8), which we

178   modeled as a modified Arrhenius equation (36):

$$K(T) = \frac{EFD(T_0) * pEA(T_0) * MDR(T_0) - \mu(T_0)}{EFD(T_0) * pEA(T_0) * MDR(T_0)} * N_m * e^{\frac{-E_A*(T-T_0)^2}{\kappa_B*(T+273)*(T_0+273)}} \qquad (8)$$

Here, T_0 is defined as the reference temperature (i.e., the temperature at

which the carrying capacity is greatest) in Celsius, N_m is the maximum carrying

capacity, and κ_B is Boltzmann constant (8.617 x 10^{-5} eV/K). EFD is the number of

eggs laid per female per day, pEA is the probability of egg-to-adult mosquito

survival, MDR is the mosquito egg-to-adult development rate, and μ is the adult

mosquito mortality rate. We calculated these values for the reference temperature.

E_A is the activation energy, which we set to 0.5 and represents the temperature

dependence of the carrying capacity, a conservative estimate as we lacked sufficient

data on estimates of the carrying capacity of *Aedes aegypti* and its underlying

temperature dependence. To convert from Celsius to Kelvin, we incremented the

temperature T and the reference temperature T_0 by 273. Equation (8) was adopted

from (36) and modified to allow the distribution to be unimodal. We set the

reference temperature, T_0, to 29°C, the optimal transmission temperature for *Aedes*

aegypti transmission found by Mordecai et al. (25).

Seasonal Forcing

To address seasonality in the model, we allowed temperature to vary over

time. We modeled temperature as a sinusoidal curve with a period of 365 days of

the form:

$$T(t) = \frac{T_{max} - T_{min}}{2} * \sin\left(\frac{2\pi}{365}t\right) + T_{mean} \tag{9}$$

Here, T_{max}, T_{mean}, and T_{min} represent the average monthly maximum, mean, and

minimum temperatures across a calendar year, respectively, and t is measured in

days. By modeling temperature as a function of time, we allowed the life history

traits of the *Aedes aegypti* vector to vary across time for the duration of the

epidemic.

Data

Life History Traits

To incorporate seasonal forcing into the compartmental modeling framework, we

used fitted mechanistic thermal response curves (25). Mordecai et al. (25) examined

published data on thermal responses for life history traits of the *Aedes aegypti*

vector and adopted a Bayesian approach for fitting quadratic ($Q(T)$; Eq. 10) or

Brière ($B(T)$; Eq. 11) curves.

$$Q(T) = c * (T - T_{min}) * (T - T_{max}) \tag{10}$$

$$B(T) = c * T * (T - T_{min}) * \sqrt{T_{max} - T} \tag{11}$$

Here, c is a rate constant, T_{min} is the critical temperature minimum, and T_{max} is the

critical temperature maximum. Fitted parameters for each of the traits used in the

model are provided in Table 1. Consistent with Mordecai et al. (25) we assumed

values above and below the critical thermal maxima and minima, respectively, were

equal to zero. By doing so, we constrained the distribution of each life history trait

to biologically relevant and empirically driven measures.

Mordecai et al. (25) provided a fitted thermal response for adult mosquito

lifespan (Table 1), which is equal to the inverse of the adult mosquito mortality rate

(μ), in days^{-1}, used in our model. At temperatures greater than or equal to the

critical thermal maximum and less than or equal to the critical thermal minimum,

adult mosquito mortality was set to 24 days^{-1} (i.e., mosquitoes survive for one hour

at temperatures outside of the T_{min} to T_{max} range).

Historical Weather Data

To identify areas of epidemic suitability across the globe, we extracted

monthly mean temperatures for 2016 from Weather Underground

(wunderground.com) for twenty different cities (Table 3). For each city, we

calculated the mean, minimum, and maximum from the average monthly mean

temperatures, to estimate temperature seasonality. This provided a range of the

average monthly temperature over the span of a calendar year. We chose this time

period because it provided the most recent full calendar year to demonstrate

seasonal fluctuations in temperature.

Table 3. Estimates of epidemic suitability for major cities. Epidemic suitability was

calculated as the proportion of the population that became infected in simulations

run under default conditions (see Appendix). Each city was simulated with its

respective temperature regime from the 2016 calendar year.

City	Annual Mean Temperature (°C)	Annual Temperature Amplitude (°C)	Epidemic Suitability
Buenos Aires, Argentina	16.5	8.0	0.0313
Sao Paulo, Brazil	20.6	5.0	0.509
Rio de Janeiro, Brazil	24.3	4.0	0.999
Salvador, Brazil	26.3	2.0	0.998
Fortaleza, Brazil	27.8	0.50	0.999
Belo Horizonte, Brazil	21.9	3.0	0.499
Recife, Brazil	27.2	1.5	0.999
Shanghai, China	17.6	12.5	0.993
Beijing, China	12.8	16	0.405
Guangzhou, China	22.9	8.0	0.999
Bogotá, Colombia	14.7	1.0	0.0025
Medellin, Colombia	17.9	1.0	0.00484
Cali, Colombia	25.1	1.5	0.986
Barranquilla, Colombia	28.8	1.0	1.00
Cartagena, Colombia	28.6	1.0	1.00
Delhi, India	26.3	9.5	0.926
Tokyo, Japan	17.0	10.5	0.621

Kobe, Japan	17.4	11	0.899
Manila, Philippines	29.0	1.5	1.00
New York, USA	13.8	12	0.0328

Variability in Epidemic Dynamics with Constant Temperature

We first examined how epidemic dynamics varied across different constant

temperatures. Here, we did not introduce seasonal forcing into the model but rather

assumed static life history traits for *Aedes aegypti* for the simulation period. We

simulated the model under default starting conditions (see Appendix) at different

constant temperatures ranging from 0.0°C to 40.0°C in increments of 0.1°C.

Variability in Epidemic Dynamics with Starting Temperature

Using the model that included seasonal variation in temperature, we examined how

the dynamics of an epidemic varied due to the temperature at which the epidemic

began under two temperature regimes. To address this variation, we set T_{max} =

40.0°C, T_{mean} = 25.0°C, and T_{min} = 10.0°C in the time-varying seasonal temperature

model under default parameters provided in the Appendix. We then varied the

temperature at the start of the epidemic from 10.0°C to 40.0°C in increments of

0.1°C and examined the response of final epidemic size, epidemic length, and

maximum number of infected individuals at any given point in time. We then

considered the temperature regime with T_{max} = 30.0°C, T_{mean} = 25.0°C, and T_{min} =

20.0°C. The same metrics were computed using the time-varying seasonal

temperature module with the same parameterization while varying the start of the

epidemic from 20.0°C to 30.0°C in increments of 0.1°C

Seasonal Variability of Final Epidemic Size

Using the compartmental modeling framework with the default starting conditions,

we examined the variation in final epidemic size as a result of seasonal forcing. To

do so, we simulated over a wide range of temperature mean and seasonal variance

regimes. The mean annual temperature varied from 0.0°C to 40.0°C in increments of

0.1°C, while the seasonal fluctuation about the mean (i.e., $\frac{T_{max} - T_{min}}{2}$) ranged from

0.0°C to 20.0°C in increments of 0.1°C. Many of these temperature regimes are

unlikely to be observed empirically. However, the simulated temperature regimes

spanned the full range of feasible temperature conditions. We recorded the final

epidemic size, measured as the number of individuals in the recovered

compartment at the end of the simulation, for each unique combination of mean

annual temperature and seasonal fluctuation.

We then compared simulated climate regimes with actual climates in major

cities, to measure relative epidemic suitability of the following cities: São Paulo,

Brazil; Rio de Janeiro, Brazil; Salvador, Brazil; Fortaleza, Brazil; Belo Horizonte,

Brazil; Recife, Brazil; Bogotá, Colombia; Medellín, Colombia; Cali, Colombia;

Barranquilla, Colombia; Cartagena, Colombia; Tokyo, Japan; Delhi, India; Manila,

Philippines; Shanghai, China; Beijing, China; New York City, USA; Guangzhou, China;

Kobe, Japan; and Buenos Aires, Argentina. These cities were chosen because they

represent some of the most populous urban areas across South America and

throughout the world.

Model Sensitivity and Uncertainty Analysis

To characterize uncertainty in the model, we simulated epidemic suitability at a

constant temperature, ranging from 0.0°C to 40.0°C in increments of 0.5°C, while

sampling 1,000 estimates for c, T_{min}, and T_{max} for each life history trait from the

marginal posterior distributions provided by Mordecai et al. (25). We calculated the

95% credible interval for the epidemic suitability at each temperature.

RESULTS

Variability in Epidemic Dynamics with Constant Temperature

Holding temperature constant, we examined variability in epidemic dynamics

across four temperatures: 20°C, 25°C, 30°C, and 35°C. As temperature increased

from 20°C to 30°C, the number of susceptible individuals depleted more rapidly

(Fig. 2, S_H). At 20°C and 35°C, the epidemics were small and burned out rapidly.

Although simulations run at 25°C and 30°C produced similar final epidemic sizes of

92.96% vs. 99.95% of the population infected, respectively (Fig. 2, R_H), the epidemic

peaked much faster at 30°C. Finally, smaller epidemics of 1.51% and 6.17% of the

population infected occurred at 20°C and 35°C.

Fig 2 Caption. Variation in epidemic dynamics by temperature. The model was

simulated under default parameters at four constant temperatures: 20°C, 25°C,

30°C, and 35°C. The number of humans in each compartment is represented on a log

scale.

Variability in Epidemic Dynamics with Starting Temperature

Next, we examined variability in epidemic dynamics due to the temperature at

which the epidemic began, given two fixed temperature regimes (25°C mean and

amplitude of either 5°C or 15°C). Consequently, starting temperature was allowed to

vary from 10°C to 40°C or 20°C to 30°C depending on the regime.

312   Given that an epidemic occurred, epidemic length monotonically decreased

as a function of starting temperature for both temperature regimes (Fig. 3): warmer

temperatures at the start of the epidemic produced shorter epidemics, and vice

versa. When temperature varied from 10°C to 40°C, the longest epidemic simulated

was 134.97 days and occurred at starting temperatures of 11.3°C and 11.4°C, and

the shortest epidemic lasted 16.09 days and occurred when the temperature at the

epidemic start was less than 10.3°C. When the temperature was 35.9°C or higher,

no epidemic was observed. Similarly, when temperature was constrained between

20°C and 30°C, the longest epidemic simulated was 112.9 days at a starting

temperature of 20°C, and the shortest epidemic lasted 71.3 days at a starting

temperature of 30°C.

Fig 3 Caption. Epidemiological indices as a function of starting temperature,

within a given seasonal temperature regime. The red curve represents the

maximum number of humans in the infected class (I_H) at any given point during the

simulation. The blue curve represents the final (or cumulative) epidemic size (R_H at

the final time step). The green curve represents the length of the epidemic (i.e., the

point at which the number of infected individuals was below one). Here, simulations

were run with the temperature conditions: T_{min} = 10°C, T_{mean} = 25°C, and T_{max} = 40°C

(A) and T_{min} = 20°C, T_{mean} = 25°C, and T_{max} = 30°C (B).

In contrast to epidemic length, both final epidemic size and maximum

infected responded unimodally to the temperature at the epidemic onset. The total

number of people infected peaked at a starting temperature of 24.1°C for both

temperature regimes (Fig. 3). The maximum instantaneous number of infected

individuals peaked at 24.4°C in both temperature regimes, with 8.42% of the

population infected. Together, these results show that epidemics introduced at

different times within identical seasonal temperature regimes produce dramatically

different final sizes and maximum infection rates.

Seasonal Variability of Final Epidemic Size

To address how mean temperature and seasonal variance combined to influence the

final epidemic size, we simulated over a wide range of temperature regimes that

accounted for variation in the mean and temperature range over a calendar year.

We calculated relative epidemic suitability, defined as the final epidemic size as a

proportion of the human population, for twenty major cities worldwide (Table 3).

In a constant thermal environment, a band of mean temperatures between

approximately 25°C and 35°C supports the highest epidemic suitability (Fig. 4). As

the seasonal temperature range increases, lower mean temperatures are capable of

supporting large epidemics. However, outside this narrow band of temperature

regimes, epidemic suitability rapidly diminishes and most temperature regimes did

not produce epidemics.

Fig 4 Caption. Variation in epidemic suitability across different seasonal

temperature regimes. The heat map shows the epidemic suitability (represented

as the proportion of the total human population infected during an epidemic) as a

function of mean annual temperature and temperature range. Here, temperature

range is defined as the seasonal fluctuation about the annual mean temperature.

Twenty large, globally important cities are plotted to illustrate their epidemic

suitability.

Of the focal 20 major cities, those with high mean temperature and small

average temperature fluctuations exhibited the highest epidemic suitability. For

instance, Manila, Philippines, which has a monthly mean temperature of 29°C and

average seasonal amplitude in mean temperature of 1.50°C, had an epidemic

suitability of 1.00. Cartagena and Barranquilla, Colombia also had epidemic

suitability of 1.00. On the other hand, areas with low average temperature and

greater temperature fluctuations, such as Beijing and New York, exhibited lower—

but still non-zero—epidemic suitabilities of 0.405 and 0.0328 respectively. Notably,

Guangzhou and Shanghai, China have high epidemic suitability (0.999 and 0.993,

respectively) despite high seasonal variation in temperature due to more moderate

mean temperatures (22.9 and 17.6°C, respectively), while high seasonal variation

reduced suitability to 0.926 in Delhi, India, which has a high mean temperature of

26.3°C (Fig. 4).

Model Sensitivity and Uncertainty Analysis

Within the 95% credible interval previously estimated for each trait thermal

response (25), epidemic suitability as a function of temperature is not highly

sensitive to the values of c, T_{min}, and T_{min} (Fig. 5). The 95% credible interval on

epidemic suitability across a range of constant temperatures is narrow, suggesting

low uncertainty in the predicted final epidemic sizes arising from trait thermal

responses.

Fig 5 Caption. Sensitivity of epidemic suitability to the parameterization of life

history traits. Epidemic suitability (represented as the proportion of the total

human population infected during an epidemic) as a function of a constant

temperature. Temperature was held constant, and 1,000 samples of c, T_{min}, and T_{max}

were taken from the marginal posterior distribution of each trait thermal response

from Mordecai et al. (25). The 95% credible interval is represented by the shaded

blue region, and the median is plotted in blue.

DISCUSSION

Recent outbreaks of DENV, CHIKV, and ZIKV in Latin America and across the

globe have captured the attention of the public health community and underscore

the importance of preparation for future outbreaks. As temperatures rise, the global

landscape suitable for such outbreaks will expand and shift geographically,

potentially placing a larger proportion of the world's population at risk.

Understanding how local temperature regimes govern epidemic dynamics becomes

increasingly important in resource allocation and control interventions (37).

Epidemic duration and size (i.e., proportion of individuals who eventually

become infected) are two key metrics of epidemic dynamics. We explored how

epidemic size and duration varied across temperature regimes by extending a

mechanistic modeling framework developed by (26) to incorporate nonlinear

thermal responses of vector traits and seasonal forcing (25). We found that the

temperature mean, seasonal variation, and seasonal timing of epidemic onset

differentially affected epidemic size versus duration.

The most notable variation in epidemic dynamics at constant temperature

was in the rate at which susceptible individuals were depleted. Epidemics simulated

at 25°C and 30°C produced similar final epidemic sizes despite the epidemic at 25°C

proceeding at a much slower rate. This "slow burn" phenomenon occurs because

slower depletion of susceptible individuals can produce epidemics of similar size to

epidemics that infect people very rapidly.

Within a given seasonal temperature regime, epidemic size peaked at

intermediate starting temperatures (24.1°C for the regimes simulated here). More

generally, the unimodal relationship suggests that even in highly suitable seasonal

environments, epidemics may vary dramatically depending on when they begin.

Compared to epidemic size, the maximum number of infected individuals at a time

peaked at slightly higher intermediate starting temperatures. We attribute this

difference between the optimal temperatures for final epidemic size and the

maximum number of infected individuals to the "slow burn" phenomenon: larger

epidemics occurred at slightly lower starting temperatures than the instantaneous

maximum number of people infected because the former epidemics last longer.

Epidemic suitability was sensitive to the interaction between annual mean

temperatures and seasonal temperature variation. At low seasonal amplitude, we

found a narrow band of annual mean temperatures (approximately 25-35°C) with

the highest epidemic suitability. Outside this band of temperature regimes,

suitability diminishes rapidly. Larger seasonal variation in temperature lowers the

range of optimal annual mean temperatures (i.e., suitability is high in cooler places

with larger seasonal variation in temperature). It is important to note that these

epidemic suitabilities should be treated as an upper bound on the potential for large

epidemics. That is, within highly suitable climate regimes, epidemics can vary in

magnitude due to human population size and movement dynamics (38), effective

vector control, and other mitigating factors. Likewise, our estimates are conditioned

on an *Aedes aegypti* population being present and virus introduction to support an

outbreak.

Previous modeling efforts have addressed seasonal forcing by allowing

transmission parameters to vary sinusoidally (24). This assumption of sinusoidal

variation does not fully capture the nonlinear relationship between vector traits and

temperature. Our mechanistic modeling framework provides the most

comprehensive treatment of nonlinear, seasonally varying temperature-dependence

for *Ae. aegypti* virus transmission to date. To do so, it incorporates experimentally

measured thermal responses for all transmission-relevant vector traits into a

dynamic transmission model, and these traits are forced with realistic seasonal

temperature variation. To aid future studies in adopting this framework, code is

available online to reproduce our results and to expand on this seasonal arboviral

transmission model. Accurately describing epidemic dynamics of emerging and

established vector-borne pathogens will ultimately require integrating realistic

models of environmental suitability, as presented here, with demographic, social,

and economic factors that promote or limit disease transmission (39) .

ACKNOWLEDGEMENTS

JHH and EAM were funded by the National Science Foundation (DEB-1518681).

EAM was also funded by NSF (DEB-1640780) and the Stanford University Center for

Innovation in Global Health. EAM and JMC were funded by the Stanford University

Woods Institute for the Environment, Environmental Ventures Program. MLC

acknowledges funding from the Lindsay Family E-IPER Fellowship.

REFERENCES

1. Gubler DJ. The Global Emergence / Resurgence of Arboviral Diseases As Public Health
Problems. Arch Med Search. 2002;33:330–42.
2. Faria NR, Azevedo R do S da S, Kraemer MUG, Souza R, Cunha MS, Hill SC, et al. Zika virus in
the Americas: Early epidemiological and genetic findings. Science [Internet].
2016;5036(March):aaf5036. Available from:
http://science.sciencemag.org/content/early/2016/03/23/science.aaf5036.abstract
3. Staples JE, Breiman RF, Powers AM. Chikungunya Fever: An Epidemiological Review of a Re-
Emerging Infectious Disease. Clin Infect Dis [Internet]. 2009;49(6):942–8. Available from:
http://cid.oxfordjournals.org/lookup/doi/10.1086/605496
4. World Health Organization (WHO) Regional Office for South-East Asia. Dengue: guidelines for
diagnosis, treatment, prevention, and control. Special Programme for Research and Training
in Tropical Diseases. 2009.
5. Gubler DJ. Dengue/dengue haemorrhagic fever: history and current status. Novartis Found
Symp [Internet]. 2006;277:3-16-22, 71–3, 251–3. Available from:
http://europepmc.org/abstract/MED/17319151
6. Brady OJ, Gething PW, Bhatt S, Messina JP, Brownstein JS, Hoen AG, et al. Refining the Global
Spatial Limits of Dengue Virus Transmission by Evidence-Based Consensus. PLoS Negl Trop
Dis. 2012;6(8).
7. Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution
and burden of dengue. Nature [Internet]. 2013;496(7446):504–7. Available from:
http://dx.doi.org/10.1038/nature12060
8. Khan K, Bogoch I, Brownstein JS, Miniota J, Nicolucci A, Hu W, et al. Assessing the origin of and
potential for international spread of chikungunya virus from the Caribbean. PLoS Curr
[Internet]. 2014;6:1–11. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4055609&tool=pmcentrez&ren
dertype=abstract
9. Perkins TA, Metcalf CJE, Grenfell BT, Tatem AJ. Estimating drivers of autochthonous
transmission of Chikungunya virus in its invasion of the Americas. PLoS Curr.
2015;7(OUTBREAKS):1–26.
10. Nsoesie EO, Kraemer MUG, Golding N, Pigott DM, Brady OJ, Moyes CL, et al. Global distribution
and environmental suitability for chikungunya virus, 1952 to 2015. Eurosurveillance.
2016;21(20).
11. PAHO. Regional Zika Epidemiological Update (Americas) August 25, 2017 [Internet]. Pan
American Health Organization. 2017 [cited 2017 Dec 4]. Available from:
http://www.paho.org/hq/index.php?option=com_content&view=article&id=11599:regional-
zika-epidemiological-update-americas&Itemid=41691&lang=en
12. Bjornstad ON, Finkenstadt BF, Grenfell BT. Dynamics of measles epidemics: Estimating
scaling of transmission rates using a Time series SIR model. Ecol Monogr. 2002;72(2):169–
84.
13. Coburn BJ, Wagner BG, Blower S. Modeling influenza epidemics and pandemics: insights into
the future of swine flu (H1N1). BMC Med [Internet]. 2009;7(1):30. Available from:
http://bmcmedicine.biomedcentral.com/articles/10.1186/1741-7015-7-30
14. Kucharski AJ, Funk S, Eggo RM, Mallet H, Edmunds WJ, Nilles EJ. Transmission dynamics of
Zika virus in island populations : a modelling analysis of the 2013 – 14 French Polynesia
outbreak. bioRxiv. 2016;1–15.
15. Ryan SJ, McNally A, Johnson LR, Mordecai EA, Ben-Horin T, Paaijmans K, et al. Mapping
Physiological Suitability Limits for Malaria in Africa Under Climate Change. Vector-Borne
Zoonotic Dis [Internet]. 2015;15(12):718–25. Available from:
http://online.liebertpub.com/doi/10.1089/vbz.2015.1822
16. Eisen L, Monaghan AJ, Lozano-Fuentes S, Steinhoff DF, Hayden MH, Bieringer PE. The impact
of temperature on the bionomics of Aedes (Stegomyia) aegypti, with special reference to the
cool geographic range margins. J Med Entomol [Internet]. 2014;51(3):496–516. Available

513     from: http://www.ncbi.nlm.nih.gov/pubmed/24897844

514 17. Rueda LM, Patel KJ, Axtell RC, Stinner RE. Temperature-Dependent Development and Survival
515     Rates of Culex quinquefasciatus and Aedes aegypti (Diptera: Culicidae). J Med Entomol
516     [Internet]. 1990;27(5):892–8. Available from: https://academic.oup.com/jme/article-
517     lookup/doi/10.1093/jmedent/27.5.892

518 18. Tun-Lin W, Burkot TR, Kay BH. Effects of temperature and larval diet on development rates
519     and survival of the dengue vector Aedes aegypti in north Queensland, Australia. Med Vet
520     Entomol. 2000;14(1):31–7.

521 19. Kakimura K, Matsuse IT, Takahashi H, Fukuda T, Suzuki K, Aratani M, et al. Effect of
522     temperature on the development of Aedes aegypti and Aedes albopictus. Med Entomol Zool.
523     2002;53(1):53–8.

524 20. Beserra EB, Fernandes CRM, Silva SA de O, Silva L a. Da, Santos JW Dos. Efeitos da
525     temperatura no ciclo de vida, exigências térmicas e estimativas do número de gerações anuais
526     de Aedes aegypti (Diptera, Culicidae). Iheringia Série Zool [Internet]. 2009;99(2):142–8.
527     Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0073-
528     47212009000200004&lng=pt&nrm=iso&tlng=pt

529 21. Couret J, Dotson E, Benedict MQ. Temperature, larval diet, and density effects on development
530     rate and survival of Aedes aegypti (Diptera: Culicidae). PLoS One. 2014;9(2).

531 22. Watts DM, Burke DS, Harrison BA, Whitmire RE, Nisalak A. Effect of temperature on the
532     vector efficiency of Aedes aegypti for dengue 2 virus. Am J Trop Med Hyg. 1987;36(1):143–
533     52.

534 23. McLean DM, Clarke AM, Coleman JC, Montalbetti CA, Skidmore AG, Walters TE, et al. Vector
535     capability of *Aedes aegypti* mosquitoes for California encephalitis and dengue viruses at
536     various temperatures. Can J Microbiol [Internet]. 1974;20(2):255–62. Available from:
537     http://www.nrcresearchpress.com/doi/abs/10.1139/m74-040

538 24. Altizer S, Dobson A, Hosseini P, Hudson P, Pascual M, Rohani P. Seasonality and the dynamics
539     of infectious diseases: Seasonality and infectious diseases. Ecol Lett. 2006 Mar;9(4):467–84.

540 25. Mordecai EA, Cohen JM, Evans M V., Gudapati P, Johnson LR, Lippi CA, et al. Detecting the
541     impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic
542     models. PLoS Negl Trop Dis. 2017;11(4).

543 26. Yang HM, Macoris MLG, Galvani KC, Andrighetti MTM, Wanderley DM V. Assessing the effects
544     of temperature on dengue transmission. Epidemiol Infect. 2009;137(8):1179–87.

545 27. Yang HM, Boldrini JL, Fassoni AC, Freitas LFS, Gomez MC, Lima KKB de, et al. Fitting the
546     Incidence Data from the City of Campinas, Brazil, Based on Dengue Transmission Modellings
547     Considering Time-Dependent Entomological Parameters. PLoS One [Internet].
548     2016;11(3):e0152186. Available from: http://dx.plos.org/10.1371/journal.pone.0152186

549 28. Heffernan JM, Smith RJ, Wahl LM. Perspectives on the basic reproductive ratio. J R Soc
550     Interface. 2005;2(4):281–93.

551 29. Mordecai E, Cohen J, Evans M V, Gudapati P, Johnson LR, Miazgowicz K, et al. Temperature
552     determines Zika, dengue and chikungunya transmission potential in the Americas. 2016 Jul.
553     Report No.: biorxiv;063735v1.

554 30. Duong V, Lambrechts L, Paul RE, Ly S, Lay RS, Long KC, et al. Asymptomatic humans transmit
555     dengue virus to mosquitoes. Proc Natl Acad Sci [Internet]. 2015;112(47):14688–93. Available
556     from: http://www.pnas.org/lookup/doi/10.1073/pnas.1508114112

557 31. Appassakij H, Khuntikij P, Kemapunmanus M, Wutthanarungsan R, Silpapojakul K. Viremic
558     profiles in asymptomatic and symptomatic chikungunya fever: A blood transfusion threat?
559     Transfusion. 2013;53(10 PART 2):2567–74.

560 32. Murphy BR, Whitehead SS. Immune Response to Dengue Virus and Prospects for a Vaccine.
561     Annu Rev Immunol [Internet]. 2011;29(1):587–619. Available from:
562     http://www.annualreviews.org/doi/10.1146/annurev-immunol-031210-101315

563 33. Weaver SC, Osorio JE, Livengood JA, Chen R, Stinchcomb DT. Chikungunya virus and
564     prospects for a vaccine. Expert Rev Vaccines [Internet]. 2012;11(9):1087–101. Available
565     from: http://www.tandfonline.com/doi/full/10.1586/erv.12.84

566 34. Kucharski AJ, Funk S, Eggo RM, Mallet H-P, Edmunds WJ, Nilles EJ. Transmission Dynamics of
567     Zika Virus in Island Populations: A Modelling Analysis of the 2013–14 French Polynesia

Outbreak. Barker CM, editor. PLoS Negl Trop Dis. 2016 May;10(5):e0004726.
35. Le Menach A, McKenzie FE, Flahault A, Smith DL. The unexpected importance of mosquito
oviposition behaviour for malaria: non-productive larval habitats can be sources for malaria
transmission. Malar J [Internet]. 2005;4:23. Available from:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1164431&tool=pmcentrez&ren
dertype=abstract
36. Palamara GM, Childs DZ, Clements CF, Petchey OL, Plebani M, Smith MJ. Inferring the
temperature dependence of population parameters: The effects of experimental design and
inference algorithm. Ecol Evol. 2014;4(24):4736–50.
37. Perkins TA. Retracing Zika's footsteps across the Americas with computational modeling.
Proc Natl Acad Sci [Internet]. 2017;114(22):5558–60. Available from:
http://www.pnas.org/lookup/doi/10.1073/pnas.1705969114
38. Wesolowski A, Qureshi T, Boni MF, Sundsøy PR, Johansson MA, Rasheed SB, et al. Impact of
human mobility on the emergence of dengue epidemics in Pakistan. Proc Natl Acad Sci
[Internet]. 2015;112(38):11887–92. Available from:
http://www.pnas.org/lookup/doi/10.1073/pnas.1504964112
39. Zhang Q, Sun K, Chinazzi M, Pastore y Piontti A, Dean NE, Rojas DP, et al. Spread of Zika virus
in the Americas. Proc Natl Acad Sci [Internet]. 2017;114(22):E4334–43. Available from:
http://www.pnas.org/lookup/doi/10.1073/pnas.1620161114
40. R Development Core Team. R: A Language and Environment for Statistical Computing
[Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2014. Available:
http://www.R-project.org
590
591

APPENDIX

Starting Conditions

The model requires estimates of the mosquito-to-human ratio and the

number of initially infectious humans and vectors to initiate each simulation. For

these initial conditions, we used measures provided in previous models. In a model

of the ZIKV outbreak in French Polynesia from 2013-2014, Kucharski et al. provided

marginal posterior estimates for the proportion of initially infectious mosquitoes

and the number of initially infectious humans on each island (34). We digitized the

data for Tahiti and calculated the weighted average to arrive at estimates of 0.015

for the proportion of initially infectious vectors. Kucharski et al. estimated the mean

number of initially infectious humans to be 450 (34). Assuming the 2012 population

size of 183,645 for Tahiti, we arrived at an estimate of 0.0025 for the proportion of

initially infectious humans.

The literature on the ratio of mosquitoes to humans is limited. One model

(35) that measured mosquito oviposition behavior adopted a patch modeling

framework in which the overall ratio of mosquitoes to humans was equal to two,

which we adopted here.

With estimates for the mosquito-to-human ratio and the number of initially

infectious individuals and vectors, we were able to calibrate our model. The total

human population size, N_H, was set to a constant 10,000. Based on the proportion

derived from (34), 25 individuals were initially infectious and the remaining

population (9,975 individuals) was completely susceptible. E_H and R_H were initially

set to zero.

Since the carrying capacity and adult mosquito mortality rate were both

dependent on temperature, the mosquito population size was subject to

temperature effects. We assumed that the carrying capacity was greatest at 29°C

(based on studies by Mordecai et al., 2016 and Le Menach et al., 2005) and totaled

20,000. To equilibrate the model, we set the initial mosquito population size (M_0) to

the carrying capacity calculated from eq. 8 at the starting temperature. The quantity

of initially infectious mosquitoes was calculated as 1.5% of the starting population

size (34). The remaining mosquitoes (98.5%) were susceptible. The initial

conditions were thus:

$$(S_V, E_V, I_V, S_H, E_H, I_H, R_H)_0 = (0.985 M_0, 0, 0.015 M_0, 9975, 0, 25, 0)$$

626   *Simulations*

627   Simulations were performed using a temperature-sensitive mechanistic modeling

628   framework developed in C++. Code and supporting documentation for this model

629   can be accessed at https://github.com/jhuber3/temperature-sensitive-sir.

630   Simulation outputs were processed in the R Programming Language 4.3.1 (40).