The Combined activities of FtsW and MurJ are required for lipid II translocation during cell division in *Escherichia coli*

Xiaolong Liu[1], Nils Y. Meiresonne[1], Ahmed Bouhss[2,3] and Tanneke den Blaauwen[1]*

[1]*Bacterial Cell Biology, Swammerdam Institute for Life Sciences, Faculty of Science, University of Amsterdam, Amsterdam, The Netherlands*

[2]*Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ Paris-Sud, Université Paris-Saclay, 91198, Gif-sur-Yvette, France.*

[3]*Laboratoire Structure-Activité des Biomolécules Normales et Pathologiques (SABNP), Univ Evry, INSERM U1204, Université Paris-Saclay, 91025 Evry, France*

*Corresponding author

Abstract

Peptidoglycan (PG) is the unique cell shape-determining component of the bacterial envelope, and is a key target for antibiotics. PG synthesis requires the transmembrane movement of the precursor lipid II by its flippase. Evidence has been shown that the inner membrane protein MurJ is able to provide this activity in *E. coli*. However, how MurJ functions has thus far not been reported. Here, by using functional fluorescent MurJ fusion proteins, we show *in vivo* that MurJ localizes both in the lateral membrane and at midcell, and is recruited to midcell simultaneously with late-localizing divisome proteins and the proteins MraY and MurG, which are required for Lipid II synthesis. We show that this septal localization of MurJ is dependent on the divisome activation, lipid II synthesis and FtsW activity. Investigations on localization of MurJ cysteine variants whose function can be inactivated by the cysteine-binding molecule MTSES indicate that the inward-facing conformation is the default state for MurJ. The loss-of-function MurJ mutant proteins are in this conformation and can still be recruited, while MTSES traps functional mutants A29C and S263C in the outward-facing conformation that cannot be recruited, and causes loss of localization and function of MurJ. Our results also indicate that the flipping of lipid II coincides with the MurJ conformational change from inward-facing into outward-facing conformation. Combining the available data, a model is deduced in which binding and presenting of lipid II by FtsW recruits MurJ and allows lipid II flipping by MurJ.

Key wards: peptidoglycan synthesis, lipid II, flippase, divisome, conformational changes, *E. coli*

Significance Statement

Biosynthesis of bacterial peptidoglycan requires the flipping of lipid-linked disaccharide-pentapeptide precursor lipid II from the inside to the outside of the cytoplasm membrane by the so-called flippase. To date, the conflicting evidences on the flippase candidates RodA, FtsW and MurJ have prevented the unequivocal identification of the flippase. Here we provide evidence that supports MurJ as the

Lipid-II flippase and FtsW as the provider of lipid II to MurJ reconciling the thus far

conflicting data.

Introduction

Rod-shaped bacteria such as the Gram-negative bacterium *Escherichia coli* grow by

elongation and binary fission. One of the requirements for proliferation is the

enlargement of their cell envelope, which contains an inner membrane (IM) and an

outer membrane (OM). The shape of the bacterium is determined by the shape of its

peptidoglycan (PG) layer, which is attached to the outer membrane in the periplasmic

space located between both membranes of the envelope (1). The PG layer is a mesh-

like heteropolymeric macromolecule of alternating *N*-acetylmuramyl-peptides

(MurNAc-pentapeptides) and *N*-acetylglucosamine (GlcNAc) disaccharides that are

connected by a β-(1,4) bond forming glycan chains that are crosslinked between *meso*

diaminopimelic acid (DAP) and D-Ala at the third and fourth position of the acceptor

and donor stem peptide, respectively (2). The biosynthesis of PG is a complicated

process that can be divided into 3 stages. In the cytoplasmic stage, the nucleotide

precursors UDP-GlcNAc and UDP-MurNAc-pentapeptide are synthesized from

fructose-6-phosphate by the GlmSMU and MurABCDEF proteins, respectively (3). In

the IM stage, the undecaprenyl-pyrophosphate- (C_{55}) linked intermediates lipid I (C_{55}-

MurNAc-pentapeptide) and lipid II (C_{55}-MurNAc(-pentapeptide)-GlcNAc) are

formed by the MraY and MurG proteins, respectively (2). During this stage, lipid II,

containing the disaccharide-peptide moiety, the building unit of peptidoglycan, is

translocated by a "flippase" across the IM (3–7). In the periplasmic stage, the

MurNAc-pentapeptide-GlcNAc component of lipid II is inserted into the PG layer by

the glycosyltransferase and transpeptidase activities of various PBPs (8, 9), and the

lipoyl moiety C_{55}-PP is cleaved off and recycled (10).

In *E. coli,* new PG-building units are inserted into the existing PG layer during length

growth by a protein complex termed the elongasome, whereas during division new

poles are synthesized from completely new material (11) by proteins that collectively

have been termed the divisome (12). The elongasome is organized by the actin

homolog MreB, which localizes underneath the IM in patches in a helix-like

distribution. MreB is bound to the membrane by its amphipathic helix, and interacts

with the bitopic membrane protein RodZ, the integral membrane proteins MreD and

MreC, and the PG synthesis essential proteins RodA and PBP2 (12–16). How the

elongasome is regulated is not yet very clear whereas in contrast, cell division has

been well investigated. During cell division, at least 20 proteins are recruited to

midcell to form the divisome. The tubulin homologue FtsZ is attached to the IM at

midcell by the anchor proteins FtsA and ZipA, and polymerizes to form the Z-ring (17,

18). Simultaneously, other proteins, such as the Zap proteins and FtsEX, are recruited

to the ring to form the early divisome or protoring. The protoring then recruits the

later divisome proteins FtsK, FtsQ, FtsL, FtsB, FtsW, FtsI (PBP3), PBP1B, FtsN (19),

and a number of hydrolases and regulatory proteins (20) to form the mature divisome

(21). The arrival of FtsN is believed to initiate PG synthesis by elevating the

repression of the FtsQLB complex (22–24). The assembled divisome functions as a

scaffold to initiate septal PG synthesis, OM constriction, and the cleavage of the PG

layer to separate daughter cells (20). Septal PG synthesis is mainly orchestrated by

PBP3 and PBP1B (9, 25), but PBP1A can substitute for PBP1B (26, 27). OM

constriction is organized by the Tol/Pal system, which forms a complex connecting

IM and OM (28, 29). Tightly controlled amidase activity is involved in the separation

of the daughter cells (30).

Although the stages of PG synthesis are well understood, conflicting information on

the flipping of lipid II across the IM has been published over the last few years. In *E.*

coli, RodA and FtsW that belong to the SEDS (shape, elongation, division,

sporulation) family, and MurJ that belongs to the MOP (multidrug/oligo-saccharidyl-

lipid/polysaccharide) export super family (31), are reported as candidates as lipid II

flippase, either functioning in elongation or division only or in both (6, 32, 33). An

alternative activity as glycosyltransferase was recently reported for RodA in *Bacillus*

subtilis (34), while the debate which protein is responsible for lipid II flipping is still

going on.

FtsW is an essential integral membrane cell division protein with 10 trans-membrane

helixes (TMHs) and is conserved in most bacterial species that have a PG cell wall

(35). FtsW and PBP3 form a complex that is recruited to midcell by the FtsQLB

complex (36–38). FtsW was shown *in vitro* to have lipid II binding and flipping

activity, for which two charged residues in TMH 4, arginine 145 and lysine 153,

appeared to be essential, while no lipid II interaction or flipping activity was detected

for MurJ *in vitro* (38–40).

MurJ is an essential IM protein that contains 14 TMHs with both C- and N-termini in

the cytoplasm (41, 42). Earlier functional and structural studies of MurJ reveal an

outward-facing central cavity that is formed by TMHs 1, 2, 7 and 8, which contain

several charged residues that are essential for MurJ function (41, 42). The observation

of an open inward-facing conformation in the crystal structure of MurJ from

Thermosipho africanus (MurJ$_{TA}$) suggests alternative conformational changes of

MurJ (43). *In vivo* evidence favors MurJ over FtsW as the lipid II flippase, and

depletion of MurJ causes the accumulation of lipid II (31, 32, 44, 45). Interestingly,

the function of MurJ in *E. coli* can be replaced by the flippases WzK, Amj and YtgP

from other species that lack sequence identify and similarity between them (46–48).

Although published data presents evidence that MurJ is the lipid II flippase, it is still

not clear where and how MurJ functions in bacterial cells. In this study, by visualizing

the cellular localization of MurJ with functional fluorescent protein fusions, we

provided new evidence to support MurJ as the flippase and show how MurJ performs

its function during PG synthesis.

Results

Fluorescent-protein fusions to the N-terminus of MurJ are functional. We

generated fluorescent protein (FP) fusions to MurJ in *E. coli* and investigated their

function and localization in cells. Initially, two low-copy-number plasmids, pTHV037

and pSAV057, were used for expression of fused FP-MurJ under control of a down-

regulated P$_{trcdown}$ promoter to avoid high overexpression (49). Based on the physical

properties of fluorescent proteins (50, 51), green fluorescent protein mNeonGreen

(mNG) and red fluorescent protein mCherry (mCh) were chosen for MurJ fusions.

Fusions were tested for their ability to complement a MurJ-depletion strain (Material

and Methods).

FP fusions to either the N-terminus or the C-terminus of MurJ with different linkers

of Gly-Gly-Ser (GGS) repeats (52) were constructed. The N-terminal MurJ fusions

were fully functional in the absence of MurJ and localized to the IM, while C-terminal

fusions were less functional or not functional at all (Table 1). Among these, the N-

terminal mNG fusion with a (GGS)$_2$ linker showed the most stable and strongest

fluorescence signals, while for mCh, the fusion without linker was better, and thus

they were chosen for further studies. The crystal structure of MurJ from *T. africanus*

reveals that TMHs 13 and 14 form a hydrophobic groove, which penetrates into the

central transport cavity and is hypothesized to be important for lipid II binding (43).

The loss of function of MurJ C-terminal fusions might be caused by defects in folding

and groove formation.

MurJ localizes in the lateral wall and also at midcell. MurJ is reported to be

present at 674 and 78 molecules per cell when grown in rich and minimal medium,

respectively (53). In order to characterize MurJ at its endogenous expression level, the

MurJ fusion was introduced into the *E. coli* chromosome at its native locus. The

existence of the native promoter p_{murJ} was confirmed by inserting a divergently

transcribed chloramphenicol cassette between the stop codon of gene *yceM* upstream

of *murJ* and the putative *murJ* promoter (Fig. S1A). However, attempts to introduce

FP-fused MurJ under the native *murJ* promoter failed, apart from a mCh fusion,

which yielded a strain with defects in cell morphology and growth (Fig. S1B). Since

the plasmid expressed fusions were fully functional, a likely explanation for the

morphological defect could be that the decreased expression of the fusion under the

native *murJ* promoter is not enough to support growth. Therefore, we replaced the

chromosomal *murJ* gene with *fp-murJ* constructs transcribed under the control of

$p_{trcdown}$ promoter to generate strains XL08 and XL09 (Material and Methods). No

defects on growth or morphology were observed (Fig. S1 C and D). Strain XL08 was

grown to steady state in minimal glucose medium (Gb1) at 28°C to be able to

correlate its length to its cell division cycle age (18, 54), and the localization of MurJ

in living *E. coli* cells was analyzed using ImageJ and the Coli-Inspector of ObjectJ

(54). The mass double time (MD) of this strain is 83 minutes, which is close to 80

minutes of wild-type *E. coli* (54). MurJ localized in the cylindrical part of the cells as

well as specifically at midcell in dividing cells (Fig. 1). After normalizing and plotting

the fluorescence profiles into 10 age groups, a clear midcell localization was observed

starting from 50%-60% of the cell cycle. To show that this accumulation was not an

effect of double membrane formation during cell invagination, the localization of

mNG-(GGS)$_2$-GlpT was introduced as control. GlpT (glycerol-3-phosphate
transporter) is the major transporter for the uptake of *sn*-glycerol 3-phosphate in *E.*
coli. It was reported to have 12 TMHs and to function as monomer (55), which
diffuses homogenously in the IM (56). It was therefore expected to show homogenous
membrane localization. Like for MurJ, the localization of mNG-(GGS)$_2$-GlpT was
imaged and analyzed. As expected, GlpT localized homogenously in the membrane
without midcell accumulation (Fig. 1B). After plotting into 10 age groups, GlpT
showed a clear difference with MurJ, and no enhanced fluorescence was observed at
midcell during constriction (Fig. 1A). The differences between the localization
patterns of the two proteins are also obvious in images of the cells (Fig. 1C). The
specific localization pattern of MurJ suggests that MurJ is likely involved in both
length growth and cell division.

MurJ is recruited to midcell simultaneously with the PG synthesis proteins.
Since potential changes in MurJ cylindrical localization is not easily visualized, we
focused on the MurJ septal localization. The midcell localization of MurJ promoted us
to investigate at which cell division cycle age it was recruited. Because XL08 is not a
wild-type strain, we could not directly compare the timing of localization of MurJ
with data from the literature (70). Therefore, the timing of localization of MurJ, the
early and late localizing proteins FtsZ and FtsN, respectively, and the proteins MraY
and MurG, which synthesize Lipid I and lipid II, respectively (21, 23, 57–60) was
determined in steady-state grown strain XL08. Cells from the same culture and optical
density were either imaged live for MurJ localization or were fixed with FAGA (2.8%
formaldehyde and 0.04% glutaraldehyde) and subsequently immunolabeled with
antibodies specific for the above mentioned proteins (61). As shown in Fig 2, the
early divisome protein FtsZ was recruited at midcell at about 25% of cell cycle, and
the core divisome assembly completed by the midcell arrival of FtsN at 40% of cell
cycle. Interestingly, MraY, MurG and MurJ accumulated at midcell simultaneously
between 40%-50% of cell cycle at the later stage of divisome assembly. These results
indicate that MurJ is very likely functioning together with the PG synthesis complex,
which would be in agreement with its function as a lipid II flippase.

MurJ requires a mature divisome for midcell localization. The timing of MurJ
midcell localization raised the question whether its localization is dependent on

particular divisome proteins. To answer this, the mNG-(GGS)$_2$-MurJ chromosomal

fusion was introduced into a series of temperature-sensitive cell division mutant

strains. First was established that MurJ still localized at midcell in a wild-type

background at the restrictive temperature (Fig. 3A). FtsA is an essential early

divisome protein that anchors the Z-ring to the IM (62, 63). The absence of FtsA

causes filamentous growth of *E. coli*, as the downstream divisome proteins are not

recruited to midcell (64). In the FtsA(*Ts*) background, MurJ showed normal midcell

localization when grown at permissive temperature. However, when grown at 42 °C

MurJ midcell localization was lost, and MurJ only localized in the filamentous

cylindrical membrane (Fig. 3B). This indicates that MurJ midcell localization is

dependent on the presence of the early divisome. Like in the FtsA(*Ts*) cells, MurJ

midcell localization was only observed when grown at permissive conditions, but not

at non-permissive conditions in temperature-sensitive strains of the later localizing

divisome proteins, FtsQ, FtsW, FtsI (65, 66) or in an FtsN depleted strain (Fig. 3 C-

F). These results suggest that MurJ requires a mature divisome for midcell

localization. To determine to what extent MurJ is dependent on cell division, we also

examined MurJ localization in strains that were deleted for the genes encoding

PBP1A, PBP1B, Pal, TolA, or the amidases AmiABC. The observation of a clear

MurJ midcell localization in the absence of these proteins, even when cells grow as

chains, indicates that MurJ midcell localization is independent of these processes (Fig.

3 G-K). All these results support the conclusion that MurJ is specifically recruited to

midcell during septal PG synthesis.

MurJ midcell localization requires PBP3 activity. As the putative flippase of lipid

II, the next obvious question would be whether the localization of MurJ is dependent

on septal PG synthesis. Septal PG synthesis requires the activity of PBP3 (FtsI) and

PBP1B, although the latter can be replaced by PBP1A (26). The specific inhibition of

PBP3 activity by aztreonam blocks septal PG synthesis without initially inhibiting the

localization of the divisome, resulting in filamentous growth of cells with divisomes

localized at regular unit cell distances (59, 67). When grown at 28 °C to steady state

in Gb1 medium, MurJ localization was determined in the presence of 1 mg.L^{-1}

aztreonam. Already after 30 min of division inhibition, MurJ midcell localization was

lost (Fig. S2), and after 1 MD and 2 MDs, filamentous morphology of cells was

observed, and MurJ midcell localization was completely abolished (Fig. S2 and Fig.
4). After 2 MDs treatment, cells were fixed and immunolabeled with antibodies
against FtsZ and FtsN to show the localization of divisomes, and against MraY and
MurG to show the localization of the lipid II-synthesizing complex. As expected, we
found that FtsZ, FtsN, MurG and MraY localized at potential division sites in the
filaments (Fig. 4). This strongly suggests that MurJ localization is dependent on the
activity of PBP3. The midcell localization is specifically dependent on the septal PG
synthesis, since MurJ localization was not affected in the presence of elongasome
inhibitors A22 and mecillinam for 2 MDs, which inactivate MreB and PBP2,
respectively (67–70), although cells started to grow spherical (Fig. S3). Together
these results indicate that the lipid II synthesizing proteins MraY and MurG are
dependent on the physical presence of the divisome, while the flippase MurJ is not
only dependent on the presence of the divisome, but also critically on the activity of
septal PG synthesis.

MurJ midcell localization requires lipid II synthesis. As the putative lipid II
flippase, we wondered whether MurJ localization would be lipid II-dependent. The
biosynthesis of lipid II requires the activity of two essential enzymes in the
cytoplasmic stage, the pyridoxal 5'-phosphate (PLP) dependent alanine racemase,
Alr, which converts L-alanine into D-alanine (3, 71), and the ATP-dependent D-
ala:D-ala ligase, Ddl, which catalyzes the formation of D-alanyl-D-alanine from two
molecules D-alanine (3, 72). D-cycloserine (DCS) is a structural analogue of D-
alanine that inactivates Alr and Ddl, and causes rapid cell lysis (73–76). At
concentration of 1 μg.ml^{-1} DCS is sufficient to deplete the lipid I precursor UDP-
MurNac-pentapeptide pool in 30 minutes when *E. coli* is grown in Gb1 (77, 78).
Therefore, steady-state grown XL08 cells were treated with different concentrations
of DCS, and MurJ localization was determined at 30 min, 1 MD and 2 MDs after
adding DCS. At concentrations of 0.5 μg.mL^{-1} DCS or higher, growth inhibition and
rapid cell lysis was observed after 30 min, while lower concentrations of DCS caused
lysis after 1 MD or longer (Fig. 5A). Interestingly, DCS also showed a dose-
dependent effect on MurJ localization. In 30 min, the increasing DCS concentrations
eventually blocked the midcell localization of MurJ, and when the DCS concentration
was higher than 0.5 μg.mL^{-1}, MurJ was mostly absent from midcell (Fig. 5B). In
addition, low concentrations of DCS showed a concentration-dependent inhibition on

MurJ midcell localization (Fig. S4). Together, these results indicate that MurJ midcell

localization is dependent on the synthesis of lipid II.

MurJ midcell localization requires FtsW activity. Our results so far showed that

MurJ midcell localization is dependent on the assembly of the divisome, the activity

of PBP3 and also the synthesis of lipid II. But because FtsW has been reported to bind

lipid II *in vitro* whereas MurJ did not, it could be possible that FtsW was somehow

involved in the flipping activity of MurJ. Investigation of divisome assembly and

regulation showed that the recruitment of the last divisome protein FtsN to midcell

depresses the inactivity of the FtsQLB, which subsequently activates the FtsW-PBP3

complex that initiates the synthesis of septal PG (23, 79). Very recently, *in vitro* data

showed that FtsW, PBP3 and PBP1B form a ternary complex, and the high lipid II

binding affinity of FtsW inhibits the polymerization of lipid II by PBP1B in the

absence of PBP3, while the presence of PBP3 stimulates the release of lipid II from

FtsW, and activates its polymerization by PBP1B (38). These observations raised the

possibility that FtsW is needed to present lipid II to MurJ, which implied that MurJ

would not be able to localize at midcell in the presence of inactive FtsW. To validate

this hypothesis, plasmids that expressed two non-functional FtsW mutants, FtsW

R145A and FtsW K153N were introduced into strain XL08 to investigate the

localization of MurJ. These two mutants were shown to localize at midcell but failed

to complement the FtsW(*Ts*) strain at non-permissive temperature, and their

expression in LMC500 wild-type cells showed a dominant-negative filamentous

morphology. In addition, *in vitro* data showed that these mutants lost the lipid II

"flipping" activity of FtsW (39). In our study, to eliminate the role of the wild-type

FtsW, 20 μM IPTG was used to overexpress these two mutants, and thus diluted the

functional copy of FtsW from the XL08 genome. When cells were grown to steady

state in minimal medium, cell length increased due to the defect on cell division,

especially in cells expressing mutant K153N (Fig. 6B). MurJ midcell localization was

lost in the presence of non-functional FtsW mutants for 2 MDs, despite the

overexpression of MurJ as it is also expressed under the IPTG inducible $P_{trcdown}$

promoter. Whereas the divisome, which was monitored by immunolabeling of FtsN,

was still localized at midcell (Fig. 6). Moreover, when grown in LB medium, strains

expressing these mutants showed more defects on morphology and MurJ localization.

At early exponential phase ($OD_{600} = 0.2$), filamentous morphology was observed for

all cultures, even for cells grown with glucose that suppressed the expression of the
FtsW mutants (Fig. S5A). Interestingly, in cultures at late log phase expressing the
FtsW K153N mutant or in cultures at stationary phase expressing the FtsW R145A
mutant, some cells started to restore the MurJ localization, especially in the cells that
were induced with IPTG (Fig. S5A). Since these mutants are toxic, we wondered
whether the XL08 strain lost these FtsW expressing plasmids at high frequency,
restoring MurJ localization. A spot assay was carried out to test our hypothesis: cells
from each culture were diluted to the same OD_{600} value and spotted on LB and LB-
ampicillin agar plates (these two mutants plasmids are ampicillin resistant). For both
strains, no obvious growth difference was observed on both plates when grown with
glucose, indicating that most cells still contained the mutant plasmids. However, for
cells that had been induced with IPTG, a large population of cells had lost the FtsW
mutant plasmids, as much less growth was observed on the ampicillin plate compared
with LB plate (Fig. S5B). These results strongly suggest that MurJ midcell
localization requires FtsW activity, rather than its physical presence.

MurJ flipping activity is not required for its localization. Earlier studies revealed
several conserved charged residues, R18, R24, R52 and R270 in the central cavity of
MurJ that are essential for its function (41–43). To verify whether MurJ needed to be
functional to allow midcell localization, an mNG fused MurJ R18A mutant was
expressed from plasmid in LMC500. The cells were grown in Gb1 and expression
was induced with 20 μM IPTG. As shown in Fig. 7A, the non-functional mutant
R18A localized both in the lateral wall and at midcell as was observed for the wild
type-MurJ fusion. Although MurJ is likely functional as monomer based on its
structure, the wild-type copy of MurJ from LMC500 genome might still contribute to
the localization of this non-functional mutant. To eliminate this potential contribution,
localization of R18A mutant was also determined in strain XL30, a MurJ-depletion
strain in which native MurJ production is under an arabinose-inducible promoter (31).
In the presence of arabinose, no growth defect was observed, and the R18A mutant
showed the same localization as in the LMC500 strain (Fig. 7B, left). When grown in
the presence of glucose, native MurJ was eventually depleted as morphology defects
were observed after 3 hours, however, the mutant R18A was still localized in lateral
wall and at midcell (Fig. 7B, right). The localization of the other loss-of-function-
mutants showed the same results as R18A (Fig 9 and Fig. S7). These results suggest

that the MurJ function is not required for its localization. Since we showed that MurJ

localization is driven by its lipid II substrate, it is likely that these residues are not

responsible for the recognition of lipid II, but for flipping lipid II from the central

cavity of MurJ.

Function and localization of MurJ single-cysteine A29C and S263C variants are

sensitive to MTSES. The function of a number of MurJ single-cysteine mutants,

A29C, N49C, S263C and E273C, was reported to be sensitive to MTSES (41).

MTSES covalently modifies reduced cysteine residues that are exposed to the

periplasm. In the absence of MTSES, these mutant proteins are functional, but not

when MTSES is bound to their cysteine residues (44, 45, 80). To investigate how

MTSES affected the functionality of these mutants, the localization of their mNG

fusions was determined in the presence of MTSES. The optimal MTSES

concentration was determined to be 0.5 mM for the Gb1 grown strain XL03 that

expressed these functional mutants (Material and Methods and Fig. S6). Expression

was induced with 20 μM IPTG and 40 μM IPTG to show their functional and

localization sensitivities to MTSES, respectively. Surprisingly, MTSES was found to

act on these mutants through two different mechanisms, although the function of all

these mutants were affected by MTSES as cell lysis was observed after MTSES

addition (Fig. 8 A-E). Midcell localization of the mutants A29C and S263C was

blocked in the presence of MTSES (Fig. 8 A, B and Fig. S8 B, C). The latter was

more sensitive to MTSES as rapid cell lysis was observed in 20 minutes and MurJ

localization could only be determined at 10 min and 20 min after MTSES addition. In

contrast, the localization of mutants N49C and E273C was not at all affected by

MTSES (Fig. 8 C, D and Fig. S8 D, E). Moreover, E273C was reported to be partially

functional (42), and indeed morphology defects were observed in many cells, while

the MurJ localization was not influenced (Fig. S8E). This supports the notion that

MurJ functionality is not essential for its localization. Additionally, the functional

cysteine-free mutant (MurJ Cys⁻) was not affected by MTSES, neither with respect to

growth nor on localization (Fig. 8E). These results indicate that, in the presence of

MTSES, the loss of function of mutants A29C and S263C is caused by the loss of

their recruitment while for N49C and E273C, only the flipping activity is blocked but

not their localization. Recently, a crystal structural study of MurJ$_{TA}$ predicted the

existence of the inward- and outward-facing states, and these 4 residues are accessible

to MTSES when MurJ$_{TA}$ is in the outward-facing state. Moreover, it was predicted

that mutants A29C and S248C (like S263C in *E. coli*) will be trapped in the out-

facing state in the presence of MTSES (43). Together, these results strongly suggest

that A29 and S263 are involved in the conformational change from outward- to

inward-facing state, and when trapped in the outward-facing state by MTSES, MurJ

cannot be recruited to midcell anymore. While mutants N49C and E273C are still able

to change back into the inward-facing state and localize. Consequently, the inward-

facing state is the default state of MurJ, and only in this state, MurJ can be recruited

and localized.

MTSES does not influence the localization of non-functional MurJ mutants.

Previous research showed that the non-functional MurJ mutants R18C, R24C, R52C,

and R270C are also accessible to MTSES (41). And we showed that MurJ

functionality is not needed for its localization. To verify whether MTSES influences

the localization of these mutants, similar MTSES experiments were performed on

these mNG-fused mutants in the LMC500 background. To eliminate the potential

influence of MTSES on wild type MurJ from the LMC500 genome, the MurJ deletion

strain expressing mNG fused wild-type MurJ was firstly checked. In agreement with

the evidence that the two native cysteine residues in wild-type MurJ are not labeled

by MTSES, no defect on cell growth, morphology or MurJ localization was observed

when MTSES was present (Fig. 8F and Fig. S8F), (41). Similarly, no growth or

morphology defect was observed after addition of MTSES to the non-functional

single cysteine mutants, and MurJ was still localized at midcell (Fig. 8G, Fig. S7 and

Fig. S8 G-K). We should point out that, when induced with 20 μM or 40 μM IPTG,

mutant R24C showed strongly reduced fluorescence and a cytoplasmic-like

localization, compared to other mutants. However, when induced with 100 μM IPTG,

R24C showed a normal MurJ localization and comparable fluorescence, indicating

that mutation R24C does not affect MurJ localization (Fig. S7D). Surprisingly, a

spontaneous suppressor mutation R447H that is situated at the cytoplasmic end of

TMH13 restored fluorescence and showed a typical MurJ localization when induced

with 40 μM IPTG (Fig. S7E-G). Since the R24C mutant was reported not to affect

protein level (42), R24C might slightly change the structure of MurJ, and affect mNG

folding and fluorescence, while mutation R447H could suppress this defect. These

results suggest that the non-functional mutants stay in the inward-facing

conformation, since we showed that only in this state, MurJ could be recruited to
midcell. Furthermore, these results indicate that these mutations cause the loss of
MurJ flipping activity, rather than lipid II binding activity, since MurJ midcell
localization is substrate dependent.
440
Localization of non-functional double mutant A29C-R52C is not sensitive to
MTSES. Our results so far indicate that the inward-facing state is essential for MurJ
midcell localization. To verify that the still-localized MTSES treated non-functional
mutants (R18C, R24C, R52C, and R270C, and N49C and E273C) are fixed in the
inward open conformation, an mNG fused MurJ A29C-R52C double mutant was
constructed. The mutant R52C is nonfunctional but still localizes when bound to
MTSES, whereas the localization and function of mutant A29C are disrupted by
MTSES. The double mutant is expected to be nonfunctional, and if it would be in the
inward-facing conformation not able to flip. MTSES is expected not to affect its
localization because residue A29C is not accessible in this state. Indeed, the double
mutant failed to complement MurJ depletion indicating the loss of MurJ function.
Interestingly, in contrast with single mutant A29C but like the non-functional single
mutant R52C, this double mutant showed clear midcell localization in the presence of
MTSES (Fig. 8H). Apparently R52C prevented the accessibility of A29C to MTSES.
This strongly suggests that the localized-loss-of-functional mutants are not able to
convert to the outward-facing conformation, and are fixed in the default inward-
facing state, which could still be recruited to midcell. More importantly, these results
suggest that the flipping of lipid II coincides with its conformational change from
inward-facing state into outward-facing state. The subsequent release of lipid II into
the periplasm would allow MurJ to switch back to the inward-facing state.
461

Discussion
Recruitment of MurJ to midcell and the coordination of cell division and septal
PG synthesis. The synthesis of PG has been studied for decades, and FtsW and RodA
have been always considered as the lipid II flippases in elongasome and divisome,
respectively (12, 33, 39, 40, 81–83). Only recently, MurJ was identified as the lipid II
flippase (31, 44, 45, 47, 48). FtsW has been shown to bind and flip lipid II and other
lipids *in vitro* but not *in vivo* (39, 40), whereas evidence has been provided that MurJ
is flipping *in vivo* but not *in vitro* (38, 44, 45). Our study indicates that these

observations can be very well all valid, as both FtsW and MurJ are required for lipid

II flipping during cell division.

Our results show for the first time the specific MurJ cellular localization in *E.*

coli. Its midcell localization is critically dependent on maturation of the divisome, on

PBP3 and FtsW activity, and lipid II synthesis. MurJ can only recognize the substrate

and localize to midcell when it is in the inward-facing conformation. In *E. coli*, MurJ

can be replaced by non-homologous flippases from other organisms (43, 46–48, 84).

Probing the interaction of MurJ with cell division proteins such as PBP3, FtsW, MurG

using our cytoplasmic FRET assay (85) did not support interaction between these

proteins and MurJ. Although negative FRET results are not at all a guarantee for the

absence of interactions, together with the complementation of MurJ by other flippases,

it suggests that protein-protein interactions are not involved in MurJ localization,

which implies that its localization at midcell is driven by its substrate lipid II. *In vivo*,

inactivation of MurJ results in accumulation of lipid II (31, 45), indicating that lipid II

synthesis is independent of the flipping process and MurJ's presence is needed for

lipid II translocation but not synthesis. Because MurJ midcell localization is driven by

the presence of its substrate, the failure of MurJ to localize at an incomplete divisome

or upon inactivation of PBP3 (by aztreonam) or FtsW (by mutation) signifies that

lipid II is inaccessible to MurJ under these conditions, despite the proper localization

of MraY (lipid I production) and MurG (lipid II production) (Fig 3, 4 and 6). But why

would lipid II not be accessible to MurJ? FtsW is clearly able to bind lipid II *in vitro*

(38), and the absence of PBP3 causes FtsW to hold lipid II and prevents its

polymerization by PBP1B, whereas presence of PBP3 stimulates the release of lipid II

from FtsW and its polymerization by PBP1B (38). Depletion of RodA and FtsW

results in the decrease in levels of PG lipid intermediates (44). Adding that a complete

divisome (FtsN arrival) is required to activate the FtsW-PBP3 complex by elevating

the repression by FtsQLB (23, 79), a logical explanation would be that Lipid II

synthesis at the divisome is stalled in the absence of a functional FtsW. If FtsW would

be able to bind and hold Lipid II, it would explain both its lipid II binding *in vitro* and

its ability to stall further lipid II synthesis when the divisome is inactivated. Therefore,

we propose that FtsW presents lipid II to MurJ, which will ensure that MurJ is only

flipping lipid II at the precise position needed.

Why would it be necessary to present lipid II at midcell to MurJ and does this imply

that RodA as an FtsW homolog is also presenting lipid II to MurJ during lateral

growth? The midcell position is very precisely determined by the cell (86, 87). If lipid

II would be synthesized randomly and flipped randomly, it could lead to loss of

precision of the binary fission into identical daughter cells. The observation that

inhibition of the elongasome did not affect division or MurJ midcell localization, and

the inhibition of the divisome did not influence cell length growth (Fig. 4 and Fig. S3),

suggest that lipid II flipping is precisely organized at the position where new PG

synthesis should occur. This can be achieved by allowing it only to flip as closely as

possible where the substrate is needed, and by only allowing it to bind lipid II next to

the proteins that are activated to synthesize the new cell wall.

Role of important residues on MurJ function. MurJ is predicted to cycle between

inward-facing and outward-facing conformations (43), and some charged residues

were reported to be essential for MurJ function (42), whereas a number of functional

cysteine mutants were inactivated by MTSES. Interestingly, most of these residues

locate in the central cavity of MurJ (44). But how do these residues contribute to the

function and conformation of MurJ? The functional mutants A29C and S263C are not

recruited to midcell in the presence of MTSES (Fig. 8 A and B), indicating that they

cannot recognize the substrate lipid II anymore. Since these residues are predicted to

be accessible in the open-outward conformation (43), binding to MTSES traps MurJ

in this conformation, and causes the loss of localization and function of MurJ.

Consequently, the inward-facing state is the default structure for MurJ and only in this

conformation MurJ can be recruited. This is consistent with the fact that crystalized

MurJ showed this conformation (88). However, the other two functional mutants

N49C and E273C that are also sensitive to MTSES (Fig. 9 D and E) and the total-

loss-of-function mutants R18C, R24C, R52C and R270C are still able to localize at

midcell, suggesting that these mutants are in the inward-facing conformation, and

therefore are able to recognize substrate but have lost flippase activity. The

observation that the MurJ double mutant A29C-R52C is still able to localize at

midcell in the presence of MTSES strongly suggests that these loss-of-function

mutants stay only in the inward-facing conformation, since A29C is no longer

accessible to MTSES, and further indicates that the flipping of lipid II coincides with

the MurJ conformational change from inward-facing state into outward-facing state.

Narrowing the pore channel by binding MTSES at the non-essential residues N49C

and E273C interrupts the passthrough of lipid II during flipping, and traps MurJ in the inward-facing conformation only, which results in the observations that they are originally functional and sensitive to MTSES, but behave like the total-loss-of-function mutants. The conclusion that these identified important residues are responsible for lipid II flipping rather than recognition fits the fact that they are not conserved in the flippases from other bacteria (Fig. S9), which could complement MurJ depletion in *E. coli*. This indicates that Gram negative and Gram positive MurJ might have different flipping mechanisms.

Model of MurJ function in septal PG synthesis. Based on our evidence and that of others (23, 38, 39, 41, 43, 44, 90, 91), we arrived at a model for the mechanism of MurJ recruitment and lipid II flipping during septal PG synthesis (Fig. 9). For MurJ recruitment, the arrival of the last divisome protein FtsN activates the divisome by depressing the inhibition of the FtsQLB complex, and subsequently activating the FtsW-PBP3-PBP1B complex. This promotes the presenting and release of lipid II by FtsW, which recruits MurJ to midcell (Fig. 9 A). This ensures the precise positioning of septal PG synthesis. Interruption of lipid II synthesis by D-cycloserine, or inactivation of PBP3 by aztreonam, or inactivation of FtsW will block lipid II accessibility and MurJ fails to be recruited to midcell. Once recruited, MurJ flips lipid II by changing from the inward-facing state to an outward-facing conformation and presents lipid II for new PG synthesis in the periplasm (Fig. 9 B and C). Subsequently, the unloaded outward facing MurJ switches back to the default inward-facing state, and can be recruited for new cycles (Fig. 9D). Charged residues R18, R24, R52, R270 are essential for the flipping process, and a mutation at these positions, or the binding of MTSES to residues N49C and E273C will trap MurJ in the inward-facing state that is still able to localize (Fig. 9B). While the binding of MTSES on residues A29C and S263C will trap MurJ in the outward-facing state, and prevents it from running new cycles (Fig. 9D). Here, we showed how MurJ flips lipid II during cell division, it should be noted that although we mainly focused on the septal MurJ, some evidence, like the D-cycloserine experiments, MurJ depletion experiment, and MTSES experiments, showed a global effect on *E. coli*, indicating that MurJ very likely works similarly in elongasome, probably with FtsW homologous protein RodA.

Materials and Methods

Strains, plasmids and primers used in this study can be found in *SI Appendix, Tables S1* and *S2*. Information on strains and plasmids constructing, functionality test, immunolocalization assay, MTSES assay, microscopy and data analysis are described in *SI Materials and Methods*.

ACKNOWLEDGEMENTS. We would like to thank Dr. N. Ruiz for critically reading the manuscript and for the gift of plasmids pFLAGMurJΔCys and pRC7MurJ for cloning and the functionality testing.

X.L. was supported by the Chinese Scholarship Council (File No.201406220123), and N.Y.M. by NWO ALW open program (822.02.019).

X.L. and T.B. designed most experiments. X.L performed most experiments and N.Y.M designed and performed some of the experiments. X.L and TdB analyzed the data and wrote the manuscript. A.B. isolated MraY and made antibodies against the protein. All authors critically read the manuscript and gave feedbacks.

References

1.   Vollmer W, Blanot D, De Pedro MA (2008) Peptidoglycan structure and
architecture. *FEMS Microbiol Rev* 32(2):149–167.
2.   Bouhss A, Trunkfield AE, Bugg TDH, Mengin-Lecreulx D (2008) The
biosynthesis of peptidoglycan lipid-linked intermediates. *FEMS Microbiol Rev*
32(2):208–233.
3.   Barreteau H, et al. (2008) Cytoplasmic steps of peptidoglycan biosynthesis.
FEMS Microbiol Rev 32(2):168–207.
4.   Pomorski T, Menon AK (2006) Lipid flippases and their biological functions.
Cell Mol Life Sci 63(24):2908–2921.
5.   Ruiz N (2016) Filling holes in peptidoglycan biogenesis of *Escherichia coli*.
Curr Opin Microbiol 34:1–6.
6.   Ruiz N (2015) Lipid Flippases for Bacterial Peptidoglycan Biosynthesis. *Lipid*
Insights 8(Suppl 1):21–31.
7.   Scheffers DJ, Tol MB (2015) LipidII: Just Another Brick in the Wall? *PLoS*
Pathog 11(12):1–12.
8.   Macheboeuf P, Contreras-Martel C, Job V, Dideberg O, Dessen A (2006)
Penicillin binding proteins: Key players in bacterial cell cycle and drug
resistance processes. *FEMS Microbiol Rev* 30(5):673–691.
9.   Sauvage E, Kerff F, Terrak M, Ayala JA, Charlier P (2008) The penicillin-
binding proteins: Structure and role in peptidoglycan biosynthesis. *FEMS*
Microbiol Rev 32(2):234–258.
10.   Manat G, et al. (2014) Deciphering the Metabolism of Undecaprenyl-
Phosphate: The Bacterial Cell-Wall Unit Carrier at the Membrane Frontier.
Microb Drug Resist 20(3):199–214.
11.   Pedro MA De, Quintela JC, Höltje J V, Schwarz H (1997) Murein segregation
in *Escherichia coli*. *Microbiology* 179(9):2823–2834.
12.   Den Blaauwen T, De Pedro MA, Nguyen-Distèche M, Ayala JA (2008)
Morphogenesis of rod-shaped sacculi. *FEMS Microbiol Rev* 32(2):321–344.
13.   van Teeffelen S, et al. (2011) The bacterial actin MreB rotates, and rotation
depends on cell-wall assembly. *Proc Natl Acad Sci* 108(38):15822–15827.
14.   Morgenstein RM, et al. (2015) RodZ links MreB to cell wall synthesis to
mediate MreB rotation and robust morphogenesis. *Proc Natl Acad Sci U S A*
112(40):12510–12515.
15.   Fenton AK, Mortaji L El, Lau DTC, Rudner DZ, Bernhardt TG (2016) CozE is
a member of the MreCD complex that directs cell elongation in Streptococcus
pneumoniae. *Nat Microbiol* 2(December 2016):16237.
16.   Contreras-Martel C, et al. (2017) Molecular architecture of the PBP2–MreC
core bacterial cell wall synthesis complex. *Nat Commun* 8(1):776.
17.   Sun Q, Margolin W (1998) FtsZ Dynamics during the Division Cycle of Live
Escherichia coli Cells. 180(8):2050–2056.
18.   Blaauwen T Den, Buddelmeijer N, Aarsman MEG, Hameete CM, Nanninga N
(1999) Timing of FtsZ assembly in *Escherichia coli*. *J Bacteriol* 181(17):5167–
5175.
19.   den Blaauwen T, Hamoen LW, Levin PA (2017) The divisome at 25: the road
ahead. *Curr Opin Microbiol* 36:85–94.
20.   Gray AN, et al. (2015) Coordination of peptidoglycan synthesis and outer
membrane constriction during *Escherichia coli* cell division. *Elife* 4(MAY):1–
29.
21.   Aarsman MEG, et al. (2005) Maturation of the *Escherichia coli* divisome

occurs in two steps. *Mol Microbiol* 55(6):1631–1645.

22.   Lutkenhaus J (2009) FtsN - Trigger for septation. *J Bacteriol* 191(24):7381–
7382.

23.   Weiss DS (2015) Last but not least: New insights into how FtsN triggers
constriction during *Escherichia coli* cell division. *Mol Microbiol* 95(6):903–
909.

24.   Gerding MA, et al. (2009) Self-enhanced accumulation of FtsN at division sites
and roles for other proteins with a SPOR domain (DamX, DedD, and RlpA) in
Escherichia coli cell constriction. *J Bacteriol* 191(24):7383–7401.

25.   Bertsche U, et al. (2006) Interaction between two murein (peptidoglycan)
synthases, PBP3 and PBP1B, in *Escherichia coli*. *Mol Microbiol* 61(3):675–
690.

26.   Yousif SY, Broome-Smith JK SB (1985) Lysis of *Escherichia coli* by beta-
lactam antibiotics: deletion analysis of the role of penicillin-binding proteins
1A and 1B. *J Gen Microbiol* 131(10):2839–45.

27.   Suzuki H, Nishimura Y, Hirota Y (1978) On the process of cellular division in
Escherichia coli: a series of mutants of *E. coli* altered in the penicillin-binding
proteins. *Proc Natl Acad Sci U S A* 75(2):664–668.

28.   Derouichei R, Bénédetti H, Lazzaroni JC, Lazdunski C, Lloubès R (1995)
Protein complex within *Escherichia coli* inner membrane: TolA N-terminal
domain interacts with TolQ and TolR proteins. *J Biol Chem* 270(19):11078–
11084.

29.   Dubuisson JF, Vianney A, Lazzaroni JC (2002) Mutational Analysis of the
TolA C-Terminal Domain of *Escherichia coli* and Genetic Evidence for an
Interaction between TolA and TolB. *J Bacteriol* 184(16):4620–4625.

30.   Priyadarshini R, De Pedro MA, Young KD (2007) Role of peptidoglycan
amidases in the development and morphology of the division septum in
Escherichia coli. *J Bacteriol* 189(14):5334–5347.

31.   Ruiz N (2008) Bioinformatics identification of MurJ (MviN) as the
peptidoglycan lipid II flippase in *Escherichia coli*. *Proc Natl Acad Sci U S A*
105(40):15553–7.

32.   Young KD (2014) A flipping cell wall ferry. *Microbiology* 345(6193):139–
140.

33.   Ikeda M, et al. (1989) Structural similarity among *Escherichia coli* FtsW and
RodA proteins and *Bacillus subtilis* SpoVE protein, which function in cell
division, cell elongation, and spore formation, respectively. *J Bacteriol*
171(11):6375–6378.

34.   Meeske AJ, et al. (2016) SEDS proteins are a widespread family of bacterial
cell wall polymerases. *Nature* 537(7622):1–15.

35.   Khattar MM, Begg KJ, Donachie WD (1994) Identification of FtsW and
characterization of a new *ftsW* division mutant of *Escherichia coli*. *J Bacteriol*
176(23):7140–7147.

36.   Goehring NW, Gueiros-filho F, Beckwith J (2005) Premature targeting of a cell
division protein to midcell allows dissection of divisome assembly in
Escherichia coli. *Genes Dev* 19(1):127–137.

37.   Goehring NW, Gonzalez MD, Beckwith J (2006) Premature targeting of cell
division proteins to midcell reveals hierarchies of protein interactions involved
in divisome assembly. *Mol Microbiol* 61(1):33–45.

38.   Leclercq S, et al. (2017) Interplay between Penicillin-binding proteins and
SEDS proteins promotes bacterial cell wall synthesis. *Sci Rep* 7:43306.

39.   Mohammadi T, et al. (2014) Specificity of the transport of lipid II by FtsW in
Escherichia coli. *J Biol Chem* 289(21):14707–14718.

40.   Mohammadi T, et al. (2011) Identification of FtsW as a transporter of lipid-
linked cell wall precursors across the membrane. *EMBO J* 30(8):1425–1432.

41.   Butler EK, Davis RM, Bari V, Nicholson PA, Ruiz N (2013) Structure-function
analysis of MurJ reveals a solvent-exposed cavity containing residues essential
for peptidoglycan biogenesis in *Escherichia coli*. *J Bacteriol* 195(20):4639–
4649.

42.   Butler EK, Tan WB, Joseph H, Ruiz N (2014) Charge requirements of lipid II
flippase activity in *Escherichia coli*. *J Bacteriol* 196(23):4111–4119.

43.   Kuk ACY, Mashalidis EH, Lee S-Y (2016) Crystal structure of the MOP
flippase MurJ in an inward-facing conformation. *Nat Struct Mol Biol* 24(2):1–
8.

44.   Sham L-T, et al. (2014) MurJ is the flippase of lipid-linked precursors for
peptidoglycan biogenesis. *Science (80-)* 345(6193):220–222.

45.   Qiao Y, et al. (2017) Lipid II overproduction allows direct assay of
transpeptidase inhibition by [beta]-lactams. *Nat Chem Biol* 13(7):793–798.

46.   Ruiz N (2009) *Streptococcus pyogenes* YtgP (Spy-0390) complements
Escherichia coli strains depleted of the putative peptidoglycan flippase MurJ.
Antimicrob Agents Chemother 53(8):3604–3605.

47.   Meeske AJ, et al. (2015) MurJ and a novel lipid II flippase are required for cell
wall biogenesis in Bacillus subtilis. *Proc Natl Acad Sci U S A* 112(20):6437–
42.

48.   Elhenawy W, et al. (2016) The o-antigen flippase wzk can substitute for MurJ
in peptidoglycan synthesis in helicobacter pylori and *Escherichia coli*. *PLoS*
One 11(8):1–16.

49.   van der Ploeg R, Goudelis ST, den Blaauwen T (2015) Validation of FRET
Assay for the Screening of Growth Inhibitors of *Escherichia coli* Reveals
Elongasome Assembly Dynamics. *Int J Mol Sci* 16(8):17637–54.

50.   Cranfill PJ, et al. (2016) Quantitative assessment of fluorescent proteins. *Nat*
Methods 13(7):557–562.

51.   Meiresonne NY, Ploeg R van der, Hink MA, Tanneke den Blaauwen (2017)
Activity-Related Conformational Changes in D,D-Carboxypeptidases Revealed
by In Vivo Periplasmic Förster Resonance Energy Transfer Assay in
Escherichia coli. *MBio* 8(5):e01089-17.

52.   Chen X, Zaro JL, Shen WC (2013) Fusion protein linkers: Property, design and
functionality. *Adv Drug Deliv Rev* 65(10):1357–1369.

53.   Li GW, Burkhardt D, Gross C, Weissman JS (2014) Quantifying absolute
protein synthesis rates reveals principles underlying allocation of cellular
resources. *Cell* 157(3):624–635.

54.   Vischer NOE, et al. (2015) Cell age dependent concentration of *Escherichia*
coli divisome proteins analyzed with ImageJ and ObjectJ. *Front Microbiol*
6(JUN):1–18.

55.   Huang Y (2003) Structure and Mechanism of the Glycerol-3-Phosphate
Transporter from *Escherichia coli*. *Science (80-)* 301(5633):616–620.

56.   Oswald F, Varadarajan A, Lill H, Peterman EJG, Bollen YJM (2016) MreB-
Dependent Organization of the E. coli Cytoplasmic Membrane Controls
Membrane Protein Diffusion. *Biophys J* 110(5):1139–1149.

57.   Mengin-Lecreulx D, Texier L, Rousseau M, Van Heijenoort J (1991) The
murG gene of *Escherichia coli* codes for the UDP-N-acetylglucosamine:N-

acetylmuramyl-(pentapeptide) pyrophosphoryl-undecaprenol N-
acetylglucosamine transferase involved in the membrane steps of
peptidoglycan synthesis. *J Bacteriol* 173(15):4625–4636.

58. Bouhss A, Mengin-Lecreulx D, Le Beller D, Van Heijenoort J (1999)
Topological analysis of the MraY protein catalysing the first membrane step of
peptidoglycan synthesis. *Mol Microbiol* 34(3):576–585.

59. Mohammadi T, et al. (2007) The essential peptidoglycan glycosyltransferase
MurG forms a complex with proteins involved in lateral envelope growth as
well as with proteins involved in cell division in *Escherichia coli*. *Mol
Microbiol* 65(4):1106–1121.

60. Bisson Filho AW, et al. (2016) Treadmilling by FtsZ filaments drives
peptidoglycan synthesis and bacterial cell division. *bioRxiv*.
doi:10.1101/077560.

61. Buddelmeijer N, Aarsman M, Blaauwen T den (2013) Immunolabeling of
Proteins in situ in *Escherichia coli* K12 Strains. *Bio-Protocol* 3(15):1–5.

62. Pichoff S, Lutkenhaus J (2005) Tethering the Z ring to the membrane through a
conserved membrane targeting sequence in FtsA. *Mol Microbiol* 55(6):1722–
1734.

63. Rico AI, García-Ovalle M, Mingorance J, Vicente M (2004) Role of two
essential domains of *Escherichia coli* FtsA in localization and progression of
the division ring. *Mol Microbiol* 53(5):1359–1371.

64. Pichoff S, Lutkenhaus J (2002) Unique and overlapping roles for ZipA and
FtsA in septal ring assembly in *Escherichia coli*. *EMBO J* 21(4):685–693.

65. Taschner PE, Huls PG, Pas E, Woldringh CL (1988) Division behavior and
shape changes in isogenic *ftsZ*, *ftsQ*, *ftsA*, *pbpB*, and *ftsE* cell division mutants
of *Escherichia coli* during temperature shift experiments. *J Bacteriol*
170(4):1533–1540.

66. Ishino F, et al. (1989) New mutations *fts-36*, *lts-33*, and *ftsW* clustered in the
mra region of the *Escherichia coli* chromosome induce thermosensitive cell
growth and division. *J Bacteriol* 171(10):5523–5530.

67. Den Blaauwen T, Aarsman MEG, Vischer NOE, Nanninga N (2003)
Penicillin-binding protein PBP2 of *Escherichia coli* localizes preferentially in
the lateral wall and at mid-cell in comparison with the old cell pole. *Mol
Microbiol* 47(2):539–547.

68. White CL, Gober JW (2012) MreB: Pilot or passenger of cell wall synthesis?
Trends Microbiol 20(2):74–79.

69. Vollmer W, et al. (2005) Structural and mechanistic basis of penicillin-binding
protein inhibition by lactivicins. *FEMS Microbiol Rev* 30(2):565–569.

70. Shih YL, Kawagishi I, Rothfield L (2005) The MreB and Min cytoskeletal-like
systems play independent roles in prokaryotic polar differentiation. *Mol
Microbiol* 58(4):917–928.

71. Wood BWA, Gunsalusa IC (1959) D-Alanine formation; a racemase in
Streptococcus faecalis. *J Biol Chem* 190:403–416.

72. Neuhaus FC (1962) The enzymatic synthesis of D-alanyl-D-alanine. I.
Purification and properties of D-alanyl-D-alanine synthetase. *J Biol Chem*
237(3).

73. Wang E, Walsh C, Walsh C (1978) Suicide substrates for the alanine racemase
of *Escherichia coli*. *Biochemistry* 17(7):1313–1321.

74. Neuhaus FC, Lynch JL (1972) Studies on the inhibition of D-alanyl-D-alanine
synthetase by the antibiotic D-cycloserine. *Biochem Biophys Res Commun*

8(5):377–382.

75.   Azam MA, Jayaram U (2015) Inhibitors of alanine racemase enzyme: a review. *J Enzyme Inhib Med Chem* 6366(August):1–10.

76.   Prosser G a, de Carvalho LPS (2013) Kinetic mechanism and inhibition of *Mycobacterium tuberculosis* D-alanine:D-alanine ligase by the antibiotic D-cycloserine. *FEBS J* 280(4):1150–66.

77.   de Roubin MR, Mengin-Lecreulx D, van Heijenoort J (1992) Peptidoglycan biosynthesis in *Escherichia coli*: variations in the metabolism of alanine and D-alanyl-D-alanine. *J Gen Microbiol* 138 Pt 8(1 992):1751–7.

78.   Mengin-lecreulx D, et al. (1983) Pool Levels of UDP N-Acetylglucosamine and UDP N- Acetylglucosamine-Enolpyruvate in *Escherichia coli* and Correlation with Peptidoglycan Synthesis. 154(3):1284–1290.

79.   Liu B, Persons L, Lee L, de Boer PAJ (2015) Roles for both FtsA and the FtsBLQ subcomplex in FtsN-stimulated cell constriction in *Escherichia coli*. *Mol Microbiol* 95(6):945–970.

80.   Chamakura KR, et al. (2017) A viral protein antibiotic inhibits lipid II flippase activity. *Nat Microbiol*. doi:10.1038/s41564-017-0023-4.

81.   Matsuzawa H, Hayakawa K, Sato T, Imahori K (1973) Characterization and genetic analysis of a mutant of *Escherichia coli* K-12 with rounded morphology. *J Bacteriol* 115(1):436–442.

82.   Sieger B, Schubert K, Donovan C, Bramkamp M (2013) The lipid II flippase RodA determines morphology and growth in *Corynebacterium glutamicum*. *Mol Microbiol* 90(5):966–982.

83.   Dai K, Lutkenhaus J (1991) *ftsZ* is an essential cell division gene in *Escherichia coli*. *J Bacteriol* 173(11):3500–3506.

84.   Fay A, Dworkin J (2009) *Bacillus subtilis* homologs of MviN (MurJ), the putative *Escherichia coli* lipid II flippase, are not essential for growth. *J Bacteriol* 191(19):6020–6028.

85.   Alexeeva S, Gadella TWJ, Verheul J, Verhoeven GS, Den Blaauwen T (2010) Direct interactions of early and late assembling division proteins in *Escherichia coli* cells resolved by FRET. *Mol Microbiol* 77(2):384–398.

86.   Rowlett VW, Margolin W (2015) The Min system and other nucleoid-independent regulators of Z ring positioning. *Front Microbiol* 6(MAY):1–10.

87.   Ortiz C, Natale P, Cueto L, Vicente M (2015) The keepers of the ring: Regulators of FtsZ assembly. *FEMS Microbiol Rev* 40(1):57–67.

88.   Pastoret S, et al. (2004) Functional Analysis of the Cell Division Protein FtsW of *Escherichia coli*. *J Bacteriol* 186(24):8370–8379.

89.   Tsang MJ, Bernhardt TG (2015) Guiding divisome assembly and controlling its activity. *Curr Opin Microbiol* 24:60–65.

90.   Du S, Pichoff S, Lutkenhaus J (2016) FtsEX acts on FtsA to regulate divisome assembly and activity. *Proc Natl Acad Sci*:201606656.

1
Fig. 1. MurJ localizes both in the lateral wall and at midcell. Strain XL08 containing the
chromosomal mNG-MurJ fusion shows a cell-cycle-dependent localization in contrast to the
homogenously localizing integral membrane protein mNG-GlpT. Cells were grown in Gb1
minimal medium to steady state at 28°C. Left: MurJ profiles; Right: membrane control protein
GlpT profiles. (A) For each protein, the diameter (black lines) and fluorescence (green lines)
profiles along normalized cell length are shown in 10 % age class bins. (B) Maps of diameter
profiles (magenta) and fluorescence profiles (green). Cells are plotted by length, ascending
from left to right. (C) Phase contrast image (left) and the corresponding fluorescence image
(right) for each protein. Scale bar equals 5 μm. At least 1200 cells were included for both
analyses.
12

13
14
Fig. 2. Timing of MurJ midcell localization. mNG-(GGS)$_2$-MurJ cellular localization was
compared with the localization of divisomal and PG precursor synthesizing proteins. XL08
strain was grown to steady state in Gb1 medium. MurJ localization was determined in living
cells. After fixation with FAGA, the rest of cells were immunolabeled with antibodies against
FtsZ, FtsN, MraY and MurG. Diameter (black lines) and Fluorescence (colored lines) profiles
were plotted into 10 % age class bins along the normalized cell length in %. At least 1200
cells were included.

Fig. 3. MurJ requires the core divisome for its midcell localization. Chromosomal mNG-

(GGS)$_2$-MurJ fusion was introduced into strains that harbor temperature sensitive, depletable,

or deleted division proteins (except that the mNG-(GGS)$_2$-MurJ plasmid was expressed in

ΔtolA and Δpal strains, and the mCherry-MurJ plasmid was expressed in *ΔamiABC* strain).

(A-E) MurJ localization in XL08 (WT), *ftsA 1882* (Ts), *ftsQ1* (Ts), *ftsW* (Ts) and *ftsI2185* (Ts)

backgrounds. Left: Localization at permissive temperature of 28 °C. Right: Localization after

growth at the non-permissive temperature of 42 °C for 2 mass doublings (MD)s. (F) MurJ

localization in FtsN depletion background at 30 °C. Left: localization in the presence of 0.2%

w/v arabinose. Right: Localization of MurJ in the absence of arabinose for 2 MDs. (G-H) mCh-

MurJ localization in *ΔmrcA* (PBP1A), *ΔmrcB* (PBP1B), *Δpal, ΔtolA* and *ΔamiABC* background

at 30 °C, respectively. Scale bar equals 5 µm.

Fig. 4. Localization of cell division proteins in the presence of the PBP3 inhibitor aztreonam for 2 MDs. Cell division is inhibited but the localization and assemblage of the divisome continues, whereas the localization of MurJ is lost. In Gb1 at 28 °C, steady state growing XL08 was treated with 1 mg.L^{-1} aztreonam, MurJ localization was determined in living cells, and FtsZ, FtsN, MraY and MurG were immunolabeled with antibodies after fixation. Diameter (black lines) and fluorescence (colored lines) profiles were plotted into 10 % age class bins along the normalized cell length. At least 1200 cells were included.

Fig. 5 Inhibition of lipid II biogenesis blocks MurJ midcell localization. In Gb1 at 28 °C, steady state growing XL08 was treated with increasing concentrations of D-cycloserine (DCS), and mNG-MurJ localization was determined at 30 minutes after addition of DCS. (A) Growth curves were plotted as time against the natural logarithm of the optical density of the cells. The black arrow indicates the addition of DCS. (B) Fluorescence maps of MurJ localization. Cells are plotted according to increasing cell length from top to bottom. From left to right, the maps correspondent to 0, 0.05, 0.1, 0.5, 1 and 2 µg.L^{-1} DCS. At least 1200 cells were included.

Fig. 6. Non-functional FtsW mutants R145A (A) and K153N (B) abolish MurJ midcell localization. XL08 expressing mutant FtsW R145A or FtsW K153N was grown in Gb1 minimal medium to steady state at 28 °C. IPTG (20 μM) was used to induce expression of the FtsW mutants from plasmid. MurJ localization was determined in living cells and the presence of the divisome was confirmed by immunolabeling of FtsN, after fixation. Top, average localization profiles plotted along normalized cell length (n ≥1200 cells). Black lines indicate cell diameter. Green lines and red lines indicate MurJ and FtsN localization, respectively. Middle, map of profiles of cell diameter and MurJ or FtsN localization sorted by ascending cell length. Bottom, phase contrast and fluorescence microscopy images. Scale bar equals 5 μm.

74
75
76
Fig. 7. MurJ non-functional mutant R18A does not influence its localization. (A) The plasmid
expressing the mNG-MurJ R18A fusion was introduced into LMC500. This strain was grown
in Gb1 minimal medium to steady state at 28 °C, and expression was induced with 20 µM
IPTG for 2 MDs. Localization was determined and presented as outlined in Fig 1. At least
1200 cells were included. Scale bar equals 5 µm. (B) The plasmid expressing the mNG-MurJ
R18A fusion was introduced into the MurJ depletion strain XL20. Mutant localization was
determined in the presence of arabinose (wild type MurJ expression) or glucose (wild type
MurJ depletion). Scale bar equals 1 µm.

Fig. 8. MurJ mutants behave differently in the presence of MTSES. Strains expressing MurJ single-cysteine-mutants from plasmid were grown to steady state in Gb1 medium at 28 °C, and expression was induced with IPTG for 2 MDs. MurJ localization was determined at 10 minutes and 40 minutes after the addition of MTSES (for mutant S263C, localization was determined at 10 minutes and 20 minutes since it is more sensitive to MTSES), Average of MurJ localization profiles were plotted along normalized cell axis. Black lines indicate the MurJ localization in the absence of MTSES, green and red lines indicate the MurJ localization at 10 and 40 min in the presence of MTSES, respectively. (A and B) Growth and localization of functional mutants A29C and S263C are affected by MTSES. (C and D) Only growth but not localization of functional mutants N49C and E273C are affected by MTSES. (E) Growth and localization of cysteine free mutants are not altered by MTSES. (F) Growth and localization of wild type MurJ are not changed by MTSES. (G) Localization of non-functional MurJ mutant R52C is not affected by MTSES. (H) Localization of non-functional MurJ double mutant A29C-R52C is not influenced by MTSES.

120
121
Figure 9. Model of MurJ recruitment and function in septal peptidoglycan synthesis. (A)
Recruitment of MurJ to midcell when FtsW is activated. (B) Presenting of lipid II from FtsW to
MurJ. (C) MurJ flips lipid II across the IM by changing its conformation structure from the
inward-facing state into outward-facing state. (D) Unloaded MurJ changes back into default
inward-facing state to participate in the new cycle. (i) Functional MurJ mutants that are
inactivated by MTSES still localize but are not able to change from inward-facing
conformation into outward-facing conformation. (ii) Loss-of-function MurJ mutants that
localize but are not able to change from inward-facing conformation into outward-facing
conformation. (iii) Functional MurJ mutants that are fixed in the outward open conformation
upon binding of MTSES.
132
133
134

Table 1. Functionality and localization of MurJ fluorescence proteins fusions with different linkers at N- or C- terminus

Fusions	Plasmids	Complementation[a]	Localization[b]	Function[c]
No fusion	pTHV037-MurJ	≈16h	ND	+
	pSAV057-MurJ	≈16h	ND	+
N-terminus	pTHV037-mCh-(GGS)$_n$-MurJ	≈16h	+	+
	pSAV057-mNG-(GGS)$_n$-MurJ	≈16h	+	+
	pSAV057-mKO-MurJ	≈16h	+	+
C-terminus	pTHV037-MurJ-(GGS)$_n$-mCh	≥3 days	−	−
	pSAV057-MurJ-(GGS)$_n$-mNG	≥3 days	−	−

n: Number of GGS repeats (n = 0, 1, 2, 3)

[a] after transformation, the time period needed for visible white colony appearance.

[b] +, FP fused MurJ is fluorescent and localizes in the membrane; −, FP fused MurJ is fluorescent and localizes in the cytosol.

[c] +, functional fusion; −, loss-of-function fusion.