1 **FtsW activity and lipid II synthesis are required for recruitment of**
2 **MurJ to midcell during cell division in *Escherichia coli***
3

Xiaolong Liu[1], Nils Y. Meiresonne[1], Ahmed Bouhss[2,3] and Tanneke den

Blaauwen[1*]

[1]Bacterial Cell Biology, Swammerdam Institute for Life Sciences, Faculty of Science,

University of Amsterdam, Amsterdam, The Netherlands.

[2]Institute for Integrative Biology of the Cell (I2BC), CEA, CNRS, Univ Paris-Sud,

Université Paris-Saclay, 91198, Gif-sur-Yvette, France.

[3]Laboratoire Structure-Activité des Biomolécules Normales et Pathologiques

(SABNP), Univ Evry, INSERM U1204, Université Paris-Saclay, 91025 Evry, France

*Corresponding author

Abstract

Peptidoglycan (PG) is the unique cell shape-determining component of the bacterial envelope, and is a key target for antibiotics. PG synthesis requires the transmembrane movement of the precursor lipid II, and MurJ has been shown to provide this activity in *E. coli*. However, how MurJ functions *in vivo* has not been reported. Here we show that MurJ localizes both in the lateral membrane and at midcell, and is recruited to midcell simultaneously with late-localizing divisome proteins and proteins MraY and MurG. MurJ septal localization is dependent on the presence of a complete and active divisome, lipid II synthesis and PBP3/FtsW activities. Inactivation of MurJ, either directly by mutation or through binding with MTSES, did not affect the midcell localization of MurJ. Our study visualizes MurJ localization *in vivo* and reveals a possible mechanism of how MurJ functions during cell division, which gives possibilities for future investigations and further antibiotics developments.

Key words: peptidoglycan synthesis, lipid II, flippase, divisome, *E. coli*

Introduction

Rod-shaped bacteria such as the Gram-negative bacterium *Escherichia coli* grow by elongation and binary fission. One of the requirements for proliferation is the enlargement of their cell envelope, which contains an inner membrane (IM) and an outer membrane (OM). The shape of the bacterium is determined by the shape of its peptidoglycan (PG) layer, which is attached to the outer membrane in the periplasmic space located between both membranes of the envelope (Vollmer, Blanot, & De Pedro, 2008). The PG layer is a mesh-like heteropolymeric macromolecule of alternating *N*-acetylmuramyl-peptides (MurNAc-pentapeptides) and *N*-acetylglucosamine (GlcNAc) disaccharides that are connected by a β-(1,4) bond forming glycan chains that are crosslinked between *meso* diaminopimelic acid (DAP) and D-Ala at the third and fourth position of the acceptor and donor stem peptide, respectively (Bouhss, Trunkfield, Bugg, & Mengin-Lecreulx, 2008). The biosynthesis of PG is a complicated process that can be divided into 3 stages. In the cytoplasmic stage, the nucleotide precursors UDP-GlcNAc and UDP-MurNAc-pentapeptide are synthesized from fructose-6-phosphate by the GlmSMU and MurABCDEF proteins, respectively (Barreteau et al., 2008). In the IM stage, the undecaprenyl-pyrophosphate- (C_{55}) linked intermediates lipid I (C_{55}-MurNAc-pentapeptide) and lipid II (C_{55}-MurNAc(-pentapeptide)-GlcNAc) are formed by the MraY and MurG proteins, respectively (Bouhss et al., 2008). During this stage, lipid II, the building unit of peptidoglycan, is translocated by a "flippase" across the IM (Barreteau et al., 2008; Pomorski & Menon, 2006; Ruiz, 2015, 2016; Scheffers & Tol, 2015). In the periplasmic stage, the MurNAc-pentapeptide-GlcNAc component of lipid II is inserted into the PG layer by the glycosyltransferase and transpeptidase activities of various PBPs (Macheboeuf, Contreras-Martel, Job, Dideberg, & Dessen, 2006; Sauvage, Kerff, Terrak, Ayala, & Charlier, 2008), and the lipoyl moiety C_{55}-PP is cleaved off and recycled (Manat et al., 2014).

In *E. coli,* new PG-building units are inserted into the existing PG layer during length growth by a protein complex termed the elongasome, whereas during division new poles are synthesized from completely new material (Pedro, Quintela, Höltje, & Schwarz, 1997) by proteins that collectively have been termed the divisome (Den Blaauwen, De Pedro, Nguyen-Distèche, & Ayala, 2008). The elongasome is organized by the actin homolog MreB, which localizes underneath the IM in patches

in a helix-like distribution. MreB is bound to the membrane by its amphipathic helix,

and interacts with the bitopic membrane protein RodZ, the integral membrane

proteins MreD and MreC, and the PG synthesis essential proteins RodA and PBP2

(Contreras-Martel et al., 2017; Den Blaauwen et al., 2008; Fenton, Mortaji, Lau,

Rudner, & Bernhardt, 2016; Morgenstein et al., 2015; van Teeffelen et al., 2011).

How the elongasome is regulated is not yet very clear whereas in contrast, cell

division has been well investigated. During cell division, at least 20 proteins are

recruited to midcell to form the divisome. The tubulin homologue FtsZ is attached to

the IM at midcell by the anchor proteins FtsA and ZipA, and polymerizes to form the

Z-ring (Blaauwen, Buddelmeijer, Aarsman, Hameete, & Nanninga, 1999; Sun &

Margolin, 1998). Simultaneously, other proteins, such as the Zap proteins and FtsEX,

are recruited to the ring to form the early divisome or protoring. The protoring then

recruits the later divisome proteins FtsK, FtsQ, FtsL, FtsB, FtsW, FtsI (PBP3), PBP1b,

FtsN (den Blaauwen, Hamoen, & Levin, 2017), and a number of hydrolases and

regulatory proteins (Gray et al., 2015) to form the mature divisome (Aarsman et al.,

2005). The arrival of FtsN completes the assembly of the core divisome, and signals

to initiate septal PG synthesis (Gerding et al., 2009; Lutkenhaus, 2009; Weiss, 2015).

The assembled divisome functions as a scaffold to initiate septal PG synthesis, OM

constriction, and the cleavage of the PG layer to separate daughter cells (Gray et al.,

2015). Septal PG synthesis is mainly orchestrated by PBP3 and PBP1b (Bertsche et

al., 2006; Sauvage et al., 2008), but PBP1a can substitute for PBP1b (Suzuki,

Nishimura, & Hirota, 1978; Yousif SY, Broome-Smith JK, 1985). OM constriction is

organized by the Tol/Pal system, which forms a complex connecting IM and OM

(Derouichei, Bénédetti, Lazzaroni, Lazdunski, & Lloubès, 1995; Dubuisson, Vianney,

& Lazzaroni, 2002). Tightly controlled amidase activity is involved in the separation

of the daughter cells (Priyadarshini, De Pedro, & Young, 2007).

Although the stages of PG synthesis are well understood, conflicting

information on the flipping of lipid II across the IM has been published over the last

few years. In *E. coli*, RodA and FtsW that belong to the SEDS (shape, elongation,

division, sporulation) family, and MurJ that belongs to the MOP (multidrug/oligo-

saccharidyl-lipid/polysaccharide) export super family (Ruiz, 2008), are reported as

candidates of lipid II flippase, either functioning in elongation or division only or in

both (Ikeda et al., 1989; Ruiz, 2015; Young, 2014). An alternative activity as

glycosyltransferase was recently reported for RodA in *Bacillus subtilis* (Meeske et al.,

2016), while the debate which protein is responsible for lipid II flipping is still going

on. FtsW is an essential integral membrane cell division protein with 10 trans-

membrane helixes (TMHs) and is conserved in most bacterial species that have a PG

cell wall (Khattar, Begg, & Donachie, 1994). FtsW, PBP3 and PBP1b form a complex

that is recruited to midcell at the later stage of divisome assembly (Goehring,

Gonzalez, & Beckwith, 2006; Goehring, Gueiros-filho, & Beckwith, 2005; Leclercq

et al., 2017). FtsW was shown *in vitro* to have lipid II binding and flipping activity,

for which two charged residues in TMH 4, arginine 145 and lysine 153, appeared to

be essential (Leclercq et al., 2017; Mohammadi et al., 2011, 2014).

MurJ is an essential IM protein that contains 14 TMHs with both C- and N-

termini in the cytoplasm (Butler, Davis, Bari, Nicholson, & Ruiz, 2013; Butler, Tan,

Joseph, & Ruiz, 2014). Depletion of MurJ will cause irregularly shaped cell, and

finally result into cell lysis (Ruiz, 2008). Earlier functional and structural studies of

MurJ reveal an outward-facing central cavity that is formed by TMHs 1, 2, 7 and 8,

which contain several charged residues that are essential for MurJ function (Butler et

al., 2013, 2014), while the observation of an open inward-facing conformation in the

crystal structure of MurJ from *Thermosipho africanus* (MurJ$_{TA}$) suggests alternative

conformational changes of MurJ (Kuk, Mashalidis, & Lee, 2017). *In vivo* evidence

favors MurJ over FtsW as the lipid II flippase, and depletion or inhibition of MurJ

caused the accumulation of lipid II in cells (Qiao et al., 2017; Ruiz, 2008; Sham et al.,

2014; Young, 2014), while *in vitro* flipping activity has thus far only be established

for FtsW (Mohammadi et al., 2011, 2014). Interestingly, the function of MurJ in *E.

coli* can be replaced by other flippases that lack sequence similarity, such as the O-

antigen flippase Wzk from *E. coli*, and Amj and YtgP from other species (Elhenawy

et al., 2016; Hong, Liu, & Reeves, 2018; Meeske et al., 2015; Ruiz, 2009). Two *in

vitro* studies on lipid II binding of MurJ showed conflicting results, as one reported a

higher lipid II binding affinity of MurJ compared with FtsW (Bolla et al., 2018),

while the other one showed lipid II binding only by FtsW but not by MurJ (Leclercq

et al., 2017). Although these data support MurJ to be involved in flipping lipid II, it is

still not clear where and how MurJ functions in bacterial cells. In this study, by

visualizing the cellular localization of MurJ with functional fluorescent protein

fusions, we provide new evidence on how MurJ performs its function during PG

synthesis.

Results

Construct a functional fluorescent protein fusion to the N-terminus of MurJ. To investigate the MurJ function and localization *in vivo*, fluorescent protein (FP) fusions to either the N-terminus or the C-terminus of MurJ with different repeats of Gly-Gly-Ser (GGS) linkers were constructed (X. Chen, Zaro, & Shen, 2013). Two low-copy-number plasmids, pTHV037 and pSAV057, were used to express the fused proteins under the control of a down-regulated $P_{trcdown}$ promoter (van der Ploeg, Goudelis, & den Blaauwen, 2015). Based on the physical properties of fluorescent proteins (Cranfill et al., 2016; Meiresonne, van der Ploeg, Hink, & den Blaauwen, 2017), green fluorescent protein mNeonGreen (mNG) and red fluorescent protein mCherry (mCh) were chosen for MurJ fusions. The functionality of the MurJ fusions was tested by a complementation assay (Material and Methods).

As shown in Table 1, the N-terminal fusions of MurJ were fully functional, while C-terminal fusions were less functional or not functional at all. Among these, the N-terminal mNG fusion with a $(GGS)_2$ linker showed the most stable and strongest fluorescence signals. For mCh, the fusion without linker was the best. Thus they were chosen for further studies. The crystal structure of MurJ from *T. africanus* reveals that TMHs 13 and 14 form a hydrophobic groove, which penetrates into the central transport cavity and is hypothesized to be important for lipid II binding (Kuk et al., 2017). The loss function of MurJ C-terminal fusions might due to the defects in folding and groove formation.

MurJ localizes in the lateral wall and also at midcell. MurJ is reported to be present at 674 and 78 molecules per cell when grown in rich and minimal medium, respectively (Li, Burkhardt, Gross, & Weissman, 2014). In order to characterize MurJ at its endogenous expression level, the MurJ fusion was introduced into the *E. coli* chromosome at its native locus. The existence of the native promoter p_{murJ} was confirmed by inserting a divergently transcribed chloramphenicol cassette between the stop codon of gene *yceM* upstream of *murJ* and the putative *murJ* promoter (Fig. S1A). However, attempts to introduce FP-fused MurJ under the native *murJ* promoter failed, apart from a mCh fusion, which yielded a strain with defects in cell morphology and growth (Fig. S1B). Since the plasmid expressed fusions were fully functional, a likely explanation for the morphological defect could be that the

translation efficiency of the fused proteins was less than that of MurJ only.
Consequently, the expression of the FP fused MurJ under the native *murJ* promoter
was not enough to support growth. Therefore, we replaced the chromosomal *murJ*
gene with *fp-murJ* constructs transcribed under the control of $p_{trcdown}$ promoter to
generate strains XL08 and XL09 (Material and Methods). No defect on growth or cell
morphology was observed even when grown without IPTG induction (Fig. S1 C and
D).

Strain XL08 was grown to steady state in minimal glucose medium (Gb1) at
28°C to be able to correlate its length to its cell division cycle age (Blaauwen et al.,
1999; Vischer et al., 2015), and the localization of MurJ in living *E. coli* cells was
analyzed using ImageJ and the Coli-Inspector of ObjectJ (Vischer et al., 2015). To
avoid overexpression, no IPTG induction was applied. The mass double time (MD) of
this strain is 83 minutes, which is close to 80 minutes of wild-type *E. coli* (Vischer et
al., 2015). MurJ localized in the cylindrical part of the cells as well as specifically at
midcell in dividing cells (Fig. 1). After normalizing and plotting the fluorescence
profiles into 10 age groups, a clear midcell localization was observed starting from
50% of the cell cycle. The septal proportion of MurJ, which is indicated as the "Ring
fraction", was analyzed as function of the division cell cycle, and a maximal
proportion of 8.58% was observed at 82.5% of cell cycle (Fig. S2). To show that this
accumulation was not an effect of double membrane formation during cell
invagination, the localization of mNG-(GGS)$_2$-GlpT was introduced as control. GlpT
(glycerol-3-phosphate transporter) is the major transporter for the uptake of *sn*-
glycerol 3-phosphate in *E. coli*. It was reported to have 12 TMHs and to function as
monomer (Huang, 2003), which diffuses homogenously in the IM (Oswald,
Varadarajan, Lill, Peterman, & Bollen, 2016). It was therefore expected to show
homogenous membrane localization. Like for MurJ, the localization of mNG-(GGS)$_2$-
GlpT was imaged and analyzed. As expected, GlpT localized homogenously in the
membrane without midcell accumulation. After plotting into 10 age groups, GlpT
showed a clear difference with MurJ, and no enhanced fluorescence was observed at
midcell during constriction (Fig. 1A and B). The differences between the localization
patterns of the two proteins are also obvious in images of the cells (Fig. 1C). The
specific localization pattern of MurJ suggests that MurJ is likely involved in both
length growth and cell division.

MurJ is recruited to midcell simultaneously with the PG synthesis proteins.

The observation of MurJ midcell localization prompted us to the next question: at

which point of the cell division cycle MurJ is recruited. To reveal this, localization of

MurJ, the early and late localizing proteins FtsZ and FtsN, respectively, and the

proteins MraY and MurG, which synthesize Lipid I and lipid II, respectively

(Aarsman et al., 2005; Bisson Filho et al., 2017; Bouhss, Mengin-Lecreulx, Le Beller,

& Van Heijenoort, 1999; Mengin-Lecreulx, Texier, Rousseau, & Van Heijenoort,

1991; Mohammadi et al., 2007; Weiss, 2015) was determined in steady-state grown

strain XL08. Cells from the same culture and optical density were either imaged live

for MurJ localization or were fixed with FAGA (2.8% formaldehyde and 0.04%

glutaraldehyde) and subsequently immunolabeled with antibodies specific for the

above mentioned proteins (Buddelmeijer, Aarsman, & Blaauwen, 2013). The early

divisome protein FtsZ was recruited at midcell at about 25% of cell cycle, and the

core divisome assembly was completed by the midcell arrival of FtsN at 40%-50% of

the cell division cycle (Fig.2). The measured extra fluorescence at mid cell (FCplus)

of these proteins plotted as function of the cell division cycle age (Fig. S3) is in

agreement with a previous study (Van der Ploeg et al., 2013). Interestingly, MraY,

MurG and MurJ accumulated at midcell simultaneously at approximately 50% of cell

cycle at the later stage of divisome assembly. These results indicate that MurJ is very

likely functioning together with the PG synthesis complex, which would be in

agreement with its function as a lipid II flippase.

MurJ requires a mature divisome for midcell localization. The timing of MurJ

midcell localization raised the question whether its localization is dependent on

particular divisome proteins. To answer this, the mNG-(GGS)$_2$-MurJ chromosomal

fusion was introduced into a series of temperature-sensitive cell division mutant

strains. FtsA is an essential early divisome protein that anchors the Z-ring to the IM

(Pichoff & Lutkenhaus, 2005; Rico, García-Ovalle, Mingorance, & Vicente, 2004).

The absence of FtsA causes filamentous growth of *E. coli*, as the downstream

divisome proteins are not recruited to midcell (Pichoff & Lutkenhaus, 2002). In the

FtsA(*Ts*) background, MurJ showed normal midcell localization when grown at

permissive temperature. However, when grown at 42 °C, MurJ midcell localization

was lost, and MurJ only localized in the cylindrical membrane of the filamentous cells

(Fig. 3B and Fig. S4). This indicates that MurJ midcell localization is dependent on

the presence of the early divisome. Like in the FtsA(*Ts*) cells, depletion of the later

divisome proteins, FtsQ, FtsW, FtsI or FtsN (Ishino et al., 1989; Taschner, Huls, Pas,

& Woldringh, 1988), abolished MurJ midcell localization in the filamentous cells

(Fig. 3 C-F, and Fig. S4). To point out, some localization bands of MurJ were

observed in FtsQ(*Ts*) filaments at 42 °C, which was likely caused by the

relocalization of FtsQ proteins during the imaging at room temperature. As control,

MurJ still localized at midcell in a wild-type background at 42 °C (Fig. 3A). These

results suggest that MurJ requires a mature divisome for midcell localization. To

determine to what extent MurJ is dependent on cell division, we also examined MurJ

localization in strains that were deleted for PBP1a, PBP1b, Pal, TolA, or the amidases

AmiABC. The observation of a clear MurJ midcell localization in the absence of

these proteins, even when cells grow as chains, indicates that MurJ midcell

localization is independent of these proteins (Fig. 3 G-K). All these results are

consistent with the function of MurJ as the lipid II flippase during PG synthesis.

MurJ midcell localization requires PBP3 activity. After observation of MurJ

midcell localization, the next obvious question would be whether this localization is

dependent on septal PG synthesis. Septal PG synthesis requires the activity of PBP3

(FtsI) and PBP1b, although the latter can be replaced by PBP1a (Yousif SY, Broome-

Smith JK, 1985). The specific inhibition of PBP3 activity by aztreonam blocks septal

PG synthesis without initially inhibiting the localization of the divisome, resulting in

filamentous cells with divisomes localized at regular cell division distances (Den

Blaauwen, Aarsman, Vischer, & Nanninga, 2003; Mohammadi et al., 2007). After

growing the chromosomal mNG-MurJ containing strain XL08 at 28 °C to steady state

in Gb1 medium, MurJ localization was determined in the presence of 1 mg.L^{-1}

aztreonam. A block of MurJ midcell localization was already observed after 30 min,

without much change in cell morphology (Fig. S5). After 1 MD and 2 MDs, MurJ

midcell localization was completely abolished in the typical filamentous cells (Fig. S5

and Fig. 4). Subsequently, cells were fixed and immunolabeled with antibodies

against FtsZ and FtsN to show the localization of divisomes, and against MraY and

MurG to show the localization of the lipid II-synthesizing complex. As expected, we

found that FtsZ, FtsN, MurG and MraY localized at potential division sites in the

filaments (Fig. 4). Likewise, MurJ localization was determined in the presence of elongasome inhibitors A22 and mecillinam for 2 MDs, which inactivate MreB and PBP2, respectively (Den Blaauwen et al., 2003; Shih, Kawagishi, & Rothfield, 2005; Vollmer et al., 2005; White & Gober, 2012). No influence on MurJ midcell localization was observed although cells started to grow spherically (Fig. S6). Together, these results suggest that MurJ localization is specifically dependent on the activity of PBP3 and septal PG synthesis.

MurJ midcell localization requires lipid II synthesis. Next, we asked whether MurJ localization would be lipid II-dependent. The biosynthesis of lipid II requires the activity of two essential enzymes in the cytoplasmic stage, the pyridoxal 5'-phosphate (PLP) dependent alanine racemase, Alr, which converts L-alanine into D-alanine (Barreteau et al., 2008; Wood & Gunsalusa, 1959), and the ATP-dependent D-ala:D-ala ligase, Ddl, which catalyzes the formation of D-alanyl-D-alanine from two molecules of D-alanine (Barreteau et al., 2008; Neuhaus, 1962). D-cycloserine (DCS) is a structural analogue of D-alanine that inactivates Alr and Ddl, and causes rapid cell lysis (Azam & Jayaram, 2015; Neuhaus & Lynch, 1972; Prosser & de Carvalho, 2013; Wang, Walsh, & Walsh, 1978). A concentration of 1 μg.ml^{-1} DCS is sufficient to deplete the UDP-MurNac-pentapeptide pool in 30 minutes when *E. coli* is grown in minimal medium (de Roubin, Mengin-Lecreulx, & van Heijenoort, 1992; Mengin-lecreulx et al., 1983). Therefore, steady-state grown XL08 cells were treated with different concentrations of DCS, and MurJ localization was determined at 30 min after adding DCS. At concentrations of 0.5 μg.mL^{-1} DCS or higher, growth inhibition and rapid cell lysis was observed after 30 min, while lower concentrations of DCS caused lysis after 1 MD or even longer incubation periods (Fig. 5A). Interestingly, DCS also showed a concentration-dependent effect on MurJ localization. Increasing DCS concentrations eventually blocked the MurJ midcell localization, and when the DCS concentration was higher than 0.5 μg.mL^{-1}, MurJ was mostly absent from midcell (Fig. 5B and Fig. S7A). In addition, localization of MraY and MurG was somewhat reduces at midcell in the presence of DCS (Fig. S7, B and C). On the contrary, the core divisome monitored by the presence of FtsN remained intact (Fig. S7D). Together, these results indicate that MurJ midcell localization is dependent on the synthesis of lipid II.

MurJ midcell localization requires FtsW activity. Our results so far showed that MurJ midcell localization is dependent on the assembly of the divisome, the activity of PBP3 and also the synthesis of lipid II. Previous investigation of divisome assembly and regulation suggests that FtsQLB normally keeps FtsW/PBP3 inactive, and once the divisome is completely assembled, the FtsW/PBP3 complex is somehow activated (Liu, Persons, Lee, & de Boer, 2015; Weiss, 2015). Very recently, *in vitro* data showed that FtsW, PBP3 and PBP1b form a ternary complex, and the lipid II binding activity of FtsW inhibits the polymerization of lipid II by PBP1b in the absence of PBP3, while the presence of PBP3 stimulates the release of lipid II from FtsW, and activates its polymerization by PBP1b (Leclercq et al., 2017). These observations raised the possibility that FtsW is needed to ensure the lipid II accessibility to MurJ, which implied that MurJ would not be able to localize at midcell in the presence of inactive FtsW. To validate this hypothesis, plasmids that expressed two non-functional FtsW mutants, FtsW R145A and FtsW K153N were introduced into strain XL08 to investigate the localization of MurJ. These two mutants were shown to localize at midcell but failed to complement the FtsW(*Ts*) strain at non-permissive temperature, and their expression in LMC500 wild-type cells showed a dominant-negative filamentous morphology (Mohammadi et al., 2014). In addition, *in vitro* data showed that these mutants lost the lipid II "flipping" activity of FtsW (Mohammadi et al., 2014). In our study, to eliminate the role of the wild-type FtsW, 20 μM IPTG was used to overexpress these two mutants, and thus diluted the functional wild type FtsW that expressed from the XL08 genome. When cells were grown to steady state in minimal medium, cell length increased due to the defect on cell division, especially in cells expressing mutant K153N (Fig. 6B). MurJ midcell localization was lost in the presence of non-functional FtsW mutants for 2 MDs, despite the overexpression of MurJ as it is also expressed under the IPTG inducible $P_{trcdown}$ promoter. In contrast, the divisome, which was monitored by immunolabeling of FtsN, was still localized at midcell (Fig. 6). Midcell localization of MurJ was not affected when wild type FtsW was overexpressed (Fig. S8), indicating that MurJ needs a functional FtsW for its localization at mid cell. Moreover, when grown in LB medium, strains expressing these mutants showed strong defects on morphology and MurJ localization. At early exponential phase (OD$_{600}$ = 0.2), filamentous morphology was observed for all cultures, even for cells grown with glucose that suppressed the expression of the FtsW mutants (Fig. S9A). Interestingly, in cultures at late log phase

expressing the FtsW K153N mutant or in cultures at stationary phase expressing the
FtsW R145A mutant, some cells started to restore the MurJ localization, especially in
the cells that were induced with IPTG (Fig. S9A). Since these mutants are toxic, we
wondered whether the XL08 strain lost these FtsW expressing plasmids at high
frequency, restoring MurJ localization. A spot assay was carried out to test our
hypothesis: cells from each culture were diluted to the same OD_{600} value and spotted
on LB and LB-ampicillin agar plates (these two plasmids expressing the FtsW
mutants are ampicillin resistant). For both strains, no obvious growth difference was
observed on both plates when grown with glucose, indicating that most cells still
contained the mutant plasmids. However, for cells that had been induced with IPTG, a
large population of cells had lost the FtsW mutant plasmids, as much less growth was
observed on the ampicillin plate compared with LB plate (Fig. S9B). These results
strongly suggest that MurJ midcell localization requires FtsW activity, rather than its
physical presence.
381
MurJ activity is not required for its localization. Earlier studies revealed several
conserved charged residues, R18, R24, R52 and R270 in the central cavity of MurJ
that are essential for its function (41–43). To verify whether MurJ needed to be
functional to allow midcell localization, an mNG fused MurJ R18A mutant was
expressed from plasmid in LMC500. The cells were grown in Gb1 and expression
was induced with 20 μM IPTG. As shown in Fig. 7A, the non-functional mutant
R18A localized both in the lateral wall and at midcell as was observed for the wild
type-MurJ fusion. Although MurJ is likely functional as a monomer based on its
structure, the wild-type copy of MurJ from LMC500 genome might still contribute to
the localization of this non-functional mutant. To eliminate this potential contribution,
localization of the R18A mutant was also determined in strain XL30, a MurJ-
depletion strain in which native MurJ production is under an arabinose-inducible
promoter (Ruiz, 2008). In the presence of arabinose, no growth defect was observed,
and the R18A mutant showed the same localization as in the LMC500 strain (Fig. 7B,
left). When grown in the presence of glucose, native MurJ was eventually depleted as
morphology defects were observed after 3 hours, however, the mutant R18A was still
localized in lateral wall and at midcell (Fig. 7B, right). The localization of the other
loss-of-function-mutants showed the same results as R18A (see below). These results
suggest that the MurJ function is not required for its localization.

MTSES does not influence the localization of MurJ single-cysteine variants. The function of a number of MurJ single-cysteine mutants, A29C, N49C, S263C and E273C, was reported to be sensitive (only partially in the case of E273C) to MTSES (Butler et al., 2013). MTSES covalently modifies reduced cysteine residues that are exposed to the periplasm. In the absence of MTSES, these mutant proteins are functional, but not when MTSES is bound to their cysteine residues (Chamakura et al., 2017; Qiao et al., 2017; Sham et al., 2014). To investigate whether MTSES affects the functionality of these mutants through the disruption of their recruitment, the localization of their mNG fusions was determined in the presence of MTSES. Having optimized the induction conditions, MurJ cysteine mutants were induced with 40 μM IPTG, and their localization was determined at 10 min and 40 min (20 min for S263C variant due to the higher sensitivity) after addition of 0.5 mM MTSES (Material and Methods and Fig. S10). We found that MTSES influences only the functionality but not the localization of these mutants, as MurJ midcell localization was still observed, in spite of cell lysis after MTSES addition (Fig. 8 A-D, and Fig. S11). As a control, neither the functionality nor localization of cysteine free variant (MurJ Cys⁻) was influenced by MTSES (Fig. 8 E).

Next, we also investigated whether MTSES influences the localization of the total-loss-of-function mutants, R18C, R24C, R52C, and R270C. Similar MTSES experiments were performed on these mNG-fused mutants in the LMC500 background. To eliminate the potential influence of MTSES on wild type MurJ from the LMC500 genome, the MurJ deletion strain expressing mNG fused wild-type MurJ was firstly checked. In agreement with the evidence that the two native cysteine residues in wild-type MurJ are not labeled by MTSES (Butler et al., 2013), no defect on cell growth, morphology or MurJ localization was observed when MTSES was added (Fig. 8F and Fig. S11F). Similarly, no growth or morphology defect was observed after addition of MTSES to the non-functional single cysteine mutants, and MurJ still localized at midcell (Fig. 8G-H, Fig. S11, G-K and Fig. S12). To point out, when induced with 40 μM IPTG, mutant R24C showed strongly reduced fluorescence and a cytoplasmic-like localization, compared to other mutants. However, when induced with 100 μM IPTG, R24C showed a normal MurJ localization and comparable fluorescence, indicating that mutation R24C does not affect MurJ localization (Fig. S12C). Surprisingly, a spontaneous suppressor mutation R447H that

is situated at the cytoplasmic end of TMH13 restored fluorescence and showed a
typical MurJ localization when induced with 40 μM IPTG (Fig. S7D-F). Since the
R24C mutant was reported not to affect protein level (Butler et al., 2014), R24C
might slightly change the structure of MurJ, and affect mNG folding and
fluorescence, while mutation R447H could somehow suppress this defect.

Together, the investigation of MTSES on MurJ cysteine variants supports the
notion that MurJ activity is not essential for its localization.

442
443

Discussion
Recruitment of MurJ to midcell and the coordination of cell division and septal
PG synthesis. The synthesis of PG has been studied for decades, and FtsW and RodA
have been always considered as the lipid II flippases in elongasome and divisome,
respectively (Dai & Lutkenhaus, 1991; Den Blaauwen et al., 2008; Ikeda et al., 1989;
Matsuzawa, Hayakawa, Sato, & Imahori, 1973; Mohammadi et al., 2011, 2014;
Sieger, Schubert, Donovan, & Bramkamp, 2013). Only recently, MurJ was identified
as the lipid II flippase (Elhenawy et al., 2016; Meeske et al., 2015; Qiao et al., 2017;
Ruiz, 2008; Sham et al., 2014). FtsW has been shown to bind and flip lipid II and
other lipids *in vitro* but not *in vivo* (Mohammadi et al., 2011, 2014), whereas evidence
has been provided that MurJ has the lipid II flipping activity *in vivo*, but only lipid II
binding activity so far *in vitro* (Bolla et al., 2018; Leclercq et al., 2017; Qiao et al.,
2017; Sham et al., 2014). Our study indicates that these observations can be very well
all valid, as both FtsW and MurJ are required for lipid II flipping during cell division.

Our results show for the first time the specific MurJ cellular localization in *E.*
coli. Its midcell localization is critically dependent on maturation of the divisome, on
PBP3 and FtsW activity, and lipid II synthesis. Interestingly, in *E. coli*, MurJ can be
replaced by non-homologous flippases from other organisms (Elhenawy et al., 2016;
Fay & Dworkin, 2009; Kuk et al., 2017; Meeske et al., 2015; Ruiz, 2009). Probing the
interaction of MurJ with cell division proteins such as PBP3, FtsW, MurG using our
cytoplasmic FRET assay (Alexeeva, Gadella, Verheul, Verhoeven, & Den Blaauwen,
2010) did not support interaction between these proteins and MurJ, which is also
supported by the loss of MurJ localization in aztreonam treated cells that still have an
intact divisome at mid cell (Fig. 4). Although negative FRET results are not at all a
guarantee for the absence of protein interactions, together with the smoothly

replacement of MurJ by other non-homologous flippases, it suggests that protein-
protein interactions are likely not involved in MurJ recruitment, which implies that its
localization at midcell is driven by its substrate lipid II. This is also in agreement with
the delocalization of MurJ in the presence of D-cycloserine that inhibits the
production of Lipid-II (Fig. 5). Inactivation of MurJ *In vivo* results in accumulation of
lipid II (Qiao et al., 2017; Ruiz, 2008), indicating that lipid II synthesis is independent
of the flipping process and MurJ's presence is needed for lipid II translocation but not
synthesis. The failure of MurJ to localize at an incomplete divisome or upon
inactivation of PBP3 (by aztreonam) or FtsW (by mutation) signifies that lipid II is
inaccessible to MurJ under these conditions (Fig 3, 4 and 6).

But why would lipid II not be accessible to MurJ under these conditions? One
possible explanation could concern the regulation between PBP3 and FtsW to bind
lipid II. We showed that MurJ requires FtsW activity for midcell localization. FtsW is
clearly able to bind lipid II *in vitro* (Leclercq et al., 2017), and the absence of PBP3
causes FtsW to hold lipid II and prevents its polymerization by PBP1b, whereas
presence of PBP3 stimulates the release of lipid II from FtsW and allows its
polymerization by PBP1b (Leclercq et al., 2017). Adding that an incomplete divisome
will keep the FtsW-PBP3 complex activity in check (probably) by the FtsQLB
subcomplex (Liu et al., 2015; Weiss, 2015), a logical explanation would be that the
inactivated FtsW, either caused by the incomplete divisome assembly, or inactivation
of PBP3, or mutations (FtsW R145A and FtsW K513N), is unable to release lipid II to
MurJ, and thus blocks MurJ midcell recruitment. An alternative explanation could be
that FtsW, stimulated by PBP3, is required to take over the flipped lipid II from MurJ
to use it or present it for PG synthesis. The inactivated PBP3 and/or FtsW will keep
MurJ from resuming its flipping cycle. In this scenario MurJ would diffuse away from
the division site and would provide any available transglycosylase with lipid II.
Taking over lipid II from MurJ would be in agreement with the suggested
transglycosylase function of FtsW (Meeske et al., 2016). The requirement of
FtsW/PBP3 activities would also explain why MurJ is not flipping lipid II *in vitro*. A
third possibility, which we cannot exclude presently, is that FtsW and MurJ are both
flipping lipid to insert multiple glycan stands simultaneously in the septum as was
suggested in the three for one or four for two models of PG insertion during cell
division (Egan & Vollmer, 2015).

Why would it be necessary to present lipid II at midcell to MurJ (or take it

over from MurJ) and does this imply that RodA as an FtsW homolog is also

presenting (or taking over) lipid II to/from MurJ during lateral growth? The midcell

position is very precisely determined by the cell (Ortiz, Natale, Cueto, & Vicente,

2015; Rowlett & Margolin, 2015). If lipid II would be synthesized randomly and

flipped randomly, it could lead to loss of precision of the binary fission into identical

daughter cells. The observation that inhibition of the elongasome did not affect

division or MurJ midcell localization, and the inhibition of the divisome did not

influence cell length growth (Fig. 4 and Fig. S6), suggest that lipid II flipping is

precisely organized at the position where new PG synthesis should occur. This can be

achieved by allowing it only to flip as closely as possible where the substrate is

needed, and by only allowing it to bind lipid II next to the proteins that are activated

to synthesize the new cell wall.

Other studies suggested that MurJ alters between inward-facing and outward-

facing conformations during substrate transport (Kuk et al., 2017), and observed that

cysteine mutations at the charged residues, R18, R24, R52 and R270, in the cavity

center cause the loss function of MurJ, while some other cysteine mutants at position

A29, N49, S263 and E273 are functional but not in the presence of cysteine-reactive

agent MTSES (Butler et al., 2014; Sham et al., 2014). But how do these residues

contribute to the function of MurJ is not clear. Our results show that the functionality

of MurJ is not required for its midcell recruitment, despite the inactivation caused

directly by mutation or by MTSES treatment (Fig. 7 and 8). As we mentioned above,

MurJ midcell recruitment is likely driven by its substrate lipid II, our results on the

MurJ mutants suggest that these mutants are able to recognize at least part of the

substrate lipid II, but cannot flip. Also, these important residues are not conserved in

other flippases that are able to replace MurJ and therefore more likely involved in the

flipping mechanism than in initial substrate recognition (Fig. S13).

However, residues A29 and S263 are situated at the interface between the two

lobes of MurJ, and thus are predicted to be important for MurJ conformational

changes from the outward-facing to the inward-facing state, and binding of MTSES at

these two positions will trap the protein at the outward-facing state (Kuk et al., 2017;

Sham et al., 2014). Also, recent *in vitro* evidence suggests that binding of MTSES at

A29C reduced the lipid II binding affinity of MurJ protein (Bolla et al., 2018). This

would predict delocalization for the MTSES bound A29C MurJ mutant, whereas we

observe the typical midcell localization *in vivo*. If this MTSES bound mutant does not

recognize lipid II, its localization would require protein-protein interaction, which is
unlikely in view of MurJ's smoothly replacement by completely non-homologous
flippases. It would also imply that the dependence of MurJ recruitment on divisome
assembly, lipid II synthesis, FtsW activity and PBP3 activity would then be indirect,
and that an important unknown factor that determines MurJ localization is missing. *In*
vitro data are not always completely comparable to the *in vivo* situation, and given the
conflicting lipid II binding evidences *in vitro* (Bolla et al., 2018; Leclercq et al., 2017),
we assume that the MurJ A29C MTSES mutant still has a partial ability to recognize
lipid II. According to the present published models (Chamakura et al., 2017; Kuk et
al., 2017; Qiao et al., 2017), this would require changing to the inward open
conformation, which is the only conformation observed thus far in MurJ crystals (Kuk
et al., 2017).

Model of MurJ function in septal PG synthesis. Based on our evidence and that of
others (23, 38, 39, 41, 43, 44, 90, 91), we arrived at a model for the mechanism of
MurJ recruitment and lipid II flipping during septal PG synthesis (Fig. 9). After
maturation of the core divisome complex, MurJ is recruited to midcell through the
recognition of its substrate lipid II (Fig. 9 A), and ensures the precise positioning of
septal PG synthesis. Interruption of lipid II synthesis by D-cycloserine, or inactivation
of PBP3 by aztreonam, or inactivation of FtsW will likely make lipid II inaccessible
for MurJ, and block the recruitment of MurJ to midcell. Once recruited, MurJ flips
lipid II by changing from the inward-facing state to an outward-facing conformation,
and the flipped lipid II can be used for new PG synthesis (Fig. 9 B and C).
Subsequently, the unloaded outward facing MurJ switches back to the inward-facing
state, and can be recruited for new cycles (Fig. 9D). Inactivation of MurJ, either by
direct mutation at the charged residues R18, R24, R52, R270, or by binding of
MTSES to residues A29C, N49C, S263C and E273C, will disrupt the translocation of
lipid II across the membrane (Fig. 9B). Although we cannot totally rule out the
possibility of involvement of protein-protein interaction recruitment of MurJ during
septal PG synthesis, our data show a possible mechanism of how MurJ functions
during cell division *in vivo*, and visualize of MurJ localization will give possibilities
for future investigations and further antibiotics developments. It should be noted that
although we mainly focused on the septal MurJ, some evidence, like the D-
cycloserine experiments, MurJ depletion experiment, and MTSES experiments,

showed a global effect on *E. coli*, indicating that MurJ very likely works similarly in

collaboration with the elongasome, likely with the FtsW homologous protein RodA.

Material and methods

Strains and plasmids construction

Strains and plasmids used in this study are listed in Table. S1. To construct the MurJ

N-terminal fluorescent protein (FP) fusion plasmids, mCherry (mCh) (Cranfill et al.,

2016) or *E. coli* codon optimized mNeonGreen mNG (Meiresonne et al., 2017), were

firstly cloned into plasmids pTHV037 (Den Blaauwen et al., 2003) and pSAV057

(Alexeeva et al., 2010) by restriction and ligation (*Nco*I and *Eco*RI), to generate

plasmid pSAV047 and pSAV057-mNG, respectively. Subsequently, amplified MurJ

fragments with different linkers were cloned into these two plasmids by restriction

and ligation (*Eco*RI and *Hind*III), to generate the derived N-terminal fusion plasmids.

To construct the MurJ C-terminal FP fusion plasmids, MurJ gene was firstly cloned

into plasmids pTHV037 and pSAV057 with *Nco*I and *Bam*HI, to generate plasmids

pXL09 and pXL10. Subsequently, FP genes with different linkers were cloned into

these two plasmids with *Bam*HI and *Hind*III, to generate the derived C-terminal

fusion plasmids. GlpT fusion plasmid pXL28 was constructed in the same way. To

construct the MurJ mutant plasmids, QuickChange site directed mutagenesis (Agilent

technologies, Santa Clara, CA) and Gibson assembly (Gibson et al., 2009) approaches

were applied. Plasmid pXL74 was firstly generated from pXL05 by replacing the wild

type MurJ with the cysteine free mutant that was amplified from plasmid

pFLAGMurJΔCys (Butler et al., 2013). Single-cysteine-mutant plasmids were

constructed afterwards by replacing the original residues with cysteine. The double

mutation plasmid pXL93 was constructed by introducing the additional R52C

mutation into plasmid pXL82.

The MurJ deletion strain XL03 (LMC500 Δ*murJ*::FRT/pRC7MurJKan) was

constructed by λ-Red recombination (Datsenko & Wanner, 2000) in the presence of

the complementing plasmid pRC7MurJ (Butler et al., 2013). Primers priXL17 and

priXL18 were used to amplify the MurJ upstream homologous sequence region ahead

of MurJ start codon from the *E. coli* genome, primers priXL19 and priXL20 were

used to amplify the chloramphenicol (cat) cassette from plasmid pKD3(Datsenko &

Wanner, 2000), and primers priXL21 and priXL22 were used to amplify the
downstream homologous sequence that comes after MurJ stop codon from the *E. coli*
genome. An overlap PCR was performed using the obtained homologous sequences
and *cat* gene with primers priXL17 and priXL21 to generate the final recombination
fragment. This PCR product was used to generate the MurJ depletion strain sNM01
(LMC500 ΔmurJ::cat pRC7MurJKan). Subsequently, the chloramphenicol resistance
cassette was removed by plasmid pCP20 (Datsenko & Wanner, 2000), which resulted
in the final *murJ* deletion strain XL03/pRC7MurJΔKan (The kanamycin cassette on
this plasmid is also removable).

To prove the existence of the native *murJ* promoter, a chloramphenicol cassette was
inserted after the stop codon of *yceM* upstream of *murJ* with a similar strategy as
above. Primers priXL49 and priXL50 were used to amplify the upstream homologous
sequence from the *E. coli* genome, priXL54 and priXL52 were used to amplify the cat
cassette from plasmid pKD3, and priXL45 and priXL48 were used to amplify the
downstream homologous sequence from the *E. coli* genome. The amplified overlap
PCR product with primers priXL49 and priXL48 was used to construct the
recombinant strain XL04 that interrupted MurJ expression from the upstream gene
cluster. To construct MurJ chromosomal fusion strains that are expressed under its
native promoter, plasmids pXL36 and pXL37 were firstly constructed with Gibson
assembly: primers priXL51 and priXL52 were used to amplify the pKD3 backbone,
primers priXL45 and priXL46 were used to amplify the P_{murj} sequence from the *E.*
coli genome, primers priXL53 and priXL47 were used to amplify FPs-MurJ sequence
from plasmid pXL05 and pNM037. The final recombination fragments were
generated by overlap PCR of 2 fragments with primers priXL17 and priXL48 from
fragment *yceM-cat* amplified from the XL04 genome with primers priXL17 and
priXL50, and fragment *cat-P_{murj}-FPs-murJ* amplified from these assembled plasmids
with primers priXL54 and priXL48. To construct the MurJ chromosomal FP fusion
strains XL06 and XL07 expressing MurJ under control of the $P_{trcdown}$ promoter,
fragment *yceM-cat* was amplified from the XL04 genome with primers priXL17 and
priXL52, and fragment $P_{trcdown}$-FPs-MurJ was amplified from plasmids pXL06 or
pNM037 with primers priXL77 and priXL48. The two fragments were used for an
overlap PCR with primers priXL17 and priXL48 and the resulting sequence was used
to construct strains XL06 and XL07 from LMC500 by λ-Red recombination. To insert

the mNG-MurJ fusion in the chromosomal of temperature sensitive strains and deletion strains, PCR fragment *yceM-cat-P$_{trcdown}$-NG-(GGS)$_2$-MurJ* was firstly amplified from the XL06 genome with primer priXL17 and priXL48, and subsequently introduced into other strains by λ-Red recombination.

To construct an FtsN depletion strain, primers priXL140 and priXL141 were used to amplify the upstream sequence of FtsN from the *E. coli* genome, priXL54 and priXL52 were used to amplify the *cat* cassette from plasmid pKD3, and priXL138 and priXL139 were used to amplify the *araC-P$_{BAD}$* region from plasmid pJC83 (J. C. Chen & Beckwith, 2008). An overlap PCR with priXL140 and priXL138 was performed subsequently to generate the final product for recombination. The MurJ depletion strain was constructed in a similar way, primers priXL17 and priXL52 were used to amplify the *YceM-cat* region from the XL06 genome, priXL126 and priXL157 were used to amplify *araC-P$_{BAD}$* region from plasmid pKD46, priXL156 and priXL48 were used to amplify the MurJ region from the *E. coli* genome. The final PCR product for MurJ depletion was generated by an overlap PCR with primers priXL17 and priXL48.

All PCR amplifications were performed using the DNA polymerase pfuX7 prepared in our lab as described (Nørholm, 2010), and all restriction enzymes used were purchased from New England Biolabs Inc (NEB, Ipswich, MA).

Medium and growth conditions

LB medium (10 g of tryptone (Bacto laboratories, Australia), 10 g of NaCl (Merck, Kenilworth, NJ), 5 g of yeast extract (Duchefa, Amsterdam, The Netherlands) per liter) was used for growth in rich medium at 30 °C, 37 °C and 42 °C. The FtsW(*Ts*) strain was grown in LBΔNaCl medium (LB without NaCl and containing 0.1% glucose and 20 mg thymine per liter) as described previously (Pastoret et al., 2004). Gb1 minimal medium (6.33 g of K$_2$HPO$_4$ (Merck), 2.95 g of KH$_2$PO$_4$ (Riedel de Haen, Seelze, Germany), 1.05 g of (NH$_4$)$_2$SO$_4$ (Sigma, St. Louis, MO), 0.10 g of MgSO$_4$·7H$_2$O (Roth, Karlsruhe, Germany), 0.28 mg of FeSO$_4$·7H$_2$O (Sigma), 7.1 mg of Ca(NO$_3$)$_2$·4H$_2$O (Sigma), 4 mg of thiamine (Sigma), 2 mg of uracil (Sigma), 2 mg of lysine (Sigma), 2 mg of thymine (Sigma), and 0.5 % glucose (Merck) per liter, pH 7.0) was prepared for steady state growth at 28 °C as described (Van der Ploeg et al.,

2013). Protein expression was induced with isopropyl β-D-1-thiogalactopyranoside

(IPTG, Promega, Madison WI) or L-(+)arabinose (Sigma) at described

concentrations. All antibiotics were purchased from Sigma Aldric, Working

antibiotics concentrations were: 100 μg.mL^{-1} ampicillin (10 μg.mL^{-1} for Δ*tolA* and

Δ*pal* strains), 25 μg.mL^{-1} chloramphenicol, 50 μg.mL^{-1} kanamycin, 10 μg.mL^{-1}

tetracycline (for chromosomal recombinant strains, half of the concentrations were

used). Optical density (OD) was measured at 450 nm and 600 nm when grown in

minimal medium and rich medium, respectively.

Testing functionality of MurJ fusions

Functionality experiments were performed as described previously (Butler et al.,

2013, 2014) using the MurJ depletion strain XL03 that expresses a functional MurJ

copy from plasmid pRC7MurJΔKan. This plasmid has partitioning defects causing

some of the daughter cells to lose this plasmid and die after division, unless they are

complemented with a functional copy of MurJ since MurJ is essential for *E. coli*. In

addition, pRC7MurJΔKan contains the *lac*Z operon and thus cells that lost the

plasmid will also lose their ability to produce the blue color on 20 μg.L^{-1} X-gal

(Sigma) agar plates, as our pTHV and pSAV-based plasmids do not. The functional

FPs fusions to MurJ were obtained by selecting the white colonies on Ampicillin or

chloramphenicol X-gal agar dish after transformation. The growth and fluorescence

intensity of strains containing a fusion expressing plasmid were investigated with

Synergy Mx BioTek plate reader and microscopy.

Immunolocalization experiments

E. coli cells were fixed with FAGA (2.8% formaldehyde and 0.04% glutaraldehyde

final concentration) for 15-20 minutes in a shaking waterbath after steady state

growth in Gb1 minimal medium at 28 °C (Den Blaauwen et al., 2003). Subsequently,

cells were permeabilized and immunolabeled with antibodies as described

(Buddelmeijer et al., 2013). The antibodies against FtsZ, FtsN, MurG used in this

study were purified as described (Aarsman et al., 2005; Mohammadi et al., 2007).

Polyclonal antibodies against MraY were obtained from Rabbits that were injected

with an antigenic peptide that was designed with the Epiros program (Biosiris).

Based on the predicted topology by Bouhss et al (16,17) the peptide localizes in the

cytoplasmic loop between TMH 1 and 2. The sequence was H2N-

G54QVVRNDGPESHFS67C-COOH. The cysteine was added at the C-terminal to
perform the coupling with the KLH carrier protein.
709

Depletion or inactivation of of divisome proteins
Strain XL17, XL15, XL13 and XL14 that contain a MurJ FP fusion and the
temperature sensitive divisome mutant proteins FtsW, FtsI, FtsQ and FtsA,
respectively, were diluted 1:1000 from overnight cultures (grown at 30 °C) into fresh
LB medium (LB Δ NaCl medium for XL13) with 12.5 μ g.mL^{-1} chloramphenicol, and
grown at 30 °C to OD$_{600}$ of approximately 0.2. Cells were diluted 1:5 into the same
fresh medium that was pre-warmed at 30 °C and 42 °C, respectively, and kept
growing for 2 mass doublings. To deplete FtsN, strain XL23 was diluted 1:1000 from
an overnight culture (grown at 30 °C) into fresh LB medium with 12.5 μ g.mL^{-1}
chloramphenicol and 0.5% w/v glucose, and kept growing at 30 °C to OD$_{600}$ of
approximately 0.2. Induced with 0.2% w/v arabinose was set as control. After growth,
cells were immobilized on 1% agarose slides, and imaged live by phase contrast and
epifluorescence microscopy.
723

MTSES assay
Sodium (2-sulfonatoethyl) methanethiosulfonate (MTSES) was purchased from Santa
Cruz Biotechnology Company. A gradient assay on MTSES concentrations (0, 0.01,
0.05, 0.1 or 0.5 mg.mL^{-1}) was firstly applied on Gb1 steady state grown strain XL03
that expressed the MurJ A29C mutant. MurJ localization was determined by
microscopy at 10 min and 40 min after addition of MTSES. Upon induction of
expression with 20 μM IPTG for 2 MDs rapid cell lysis was observed after adding
MTSES for 10 min at concentrations 0.1mM or 0.5 mM. A pattern of MurJ midcell
localization was observed despite the increase of MTSES concentrations (Fig. S7A).
However, we noticed that lower concentration of MTSES (0.01 mM) was not
sufficient to influence the cell growth, and cells lysed too fast at higher MTSES
concentrations (above 0.05 mM) to determine MurJ localization at a longer time scale.
Since MTSES targets MurJ A29C by covalent binding, we suspected that the higher
expression of MurJ might help to increase the resistance to MTSES. Thus, 40 μM
IPTG was used for MurJ induction (2 MDs). Indeed, higher expression of A29C
slightly improved the survival of cells, with a yield of better fluorescent signal, and

MurJ midcell localization was able to be determined at both 10 min and 40 min after
MTSES addition. In addition, 0.5 mM MTSES was sufficient to show the influence
on cell growth. Thus, 0.5 mM MTSES and 40 μM IPTG inductions were used for the
investigation of MurJ localization in the presence of MTSES.

Microscope and image analysis

For localization imaging, cells were immobilized on 1.0% agarose (w/v in Gb1) pads
and imaged immediately. Fluorescence microscopy was carried out either with an
Olympus BX-60 fluorescence microscope equipped with a CoolSnap *fx*
(Photometrics) CCD camera, a 100×/N.A. 1.35 oil objective, and software 'ImageJ-
MicroManager', or with a Nikon Eclipse Ti microscope equipped with a C11440-
22CU Hamamatsu ORCA camera, a CFI Plan Apochromat DM 100× oil objective, an
Intensilight HG 130W lamp and the NIS elements software (version 4.20.01).

Images were analyzed with Coli-Inspector supported by the ObjectJ plugin for ImageJ
(version 1.49v) (Vischer et al., 2015). Briefly, the length and diameter of more than
1200 individual cells were marked and analysis in the phase contrast images.
Fluorescence and phase contrast images were aligned and fluorescence background
was subtracted as described (18). The fluorescence of each cell was collected in a one
pixel wide bar with the length of the cell. A map of the diameter or the fluorescence
localization and intensity was generated with the cells sorted according to increasing
cell from left to right. Because cells were grown to steady state, the length of the cells
can be directly correlated to the cell division cycle age. A collective profile is created
from all cell profiles in a map. They are first resampled to a normalized cell length of
100 data points, and then averaged to a single plot, in either 1 group or more age bins.
The FCplus (the extra fluorescence at mid cell in comparison to the fluorescence in
the rest of the cell) and Ringfraction (FCplus/Total cellular fluorescence) profiles
were generated with the "graph assistant" macro in ObjectJ (18).
.

ACKNOWLEDGEMENTS. We would like to thank Dr. N. Ruiz for critically
reading the manuscript and for the gift of plasmids pFLAGMurJΔCys and pRC7MurJ
for cloning and the functionality testing.
X.L. was supported by the Chinese Scholarship Council (File No.201406220123), and

References

Aarsman, M. E. G., Piette, A., Fraipont, C., Vinkenvleugel, T. M. F., Nguyen-
Distèche, M., & Den Blaauwen, T. (2005). Maturation of the *Escherichia coli*
divisome occurs in two steps. *Molecular Microbiology*, *55*(6), 1631–1645.
https://doi.org/10.1111/j.1365-2958.2005.04502.x

Al-Dabbagh B, Henry X, El Ghachi M, Auger G, Blanot D, Parquet C, Mengin-
Lecreulx D, B. A. (2008). Active Site Mapping of MraY , a Member of the
Polyprenyl-phosphate N- Acetylhexosamine 1-Phosphate Transferase
Superfamily , Catalyzing the First Membrane Step of Peptidoglycan
Biosynthesis †. *Biochemistry*, *47*(34), 8919–8928.
https://doi.org/10.1021/bi8006274

Alexeeva, S., Gadella, T. W. J., Verheul, J., Verhoeven, G. S., & Den Blaauwen, T.
(2010). Direct interactions of early and late assembling division proteins in
Escherichia coli cells resolved by FRET. *Molecular Microbiology*, *77*(2), 384–
398. https://doi.org/10.1111/j.1365-2958.2010.07211.x

Azam, M. A., & Jayaram, U. (2015). Inhibitors of alanine racemase enzyme: a
review. *Journal of Enzyme Inhibition and Medicinal Chemistry*, *6366*(August),
1–10. https://doi.org/10.3109/14756366.2015.1050010

Barreteau, H., Kovač, A., Boniface, A., Sova, M., Gobec, S., & Blanot, D. (2008).
Cytoplasmic steps of peptidoglycan biosynthesis. *FEMS Microbiology Reviews*,
32(2), 168–207. https://doi.org/10.1111/j.1574-6976.2008.00104.x

Bertsche, U., Kast, T., Wolf, B., Fraipont, C., Aarsman, M. E. G., Kannenberg, K., …
Vollmer, W. (2006). Interaction between two murein (peptidoglycan) synthases,
PBP3 and PBP1B, in *Escherichia coli*. *Molecular Microbiology*, *61*(3), 675–690.
https://doi.org/10.1111/j.1365-2958.2006.05280.x

Bisson Filho, A. W., Hsu, Y.-P., Squyres, G., Kuru, E., Wu, F., Jukes, C., … Garner,
E. (2017). Treadmilling by FtsZ filaments drives peptidoglycan synthesis and
bacterial cell division. *Science*, *355*, 739–743. https://doi.org/DOI:
10.1126/science.aak9973

Blaauwen, T. Den, Buddelmeijer, N., Aarsman, M. E. G., Hameete, C. M., &
Nanninga, N. (1999). Timing of FtsZ assembly in *Escherichia coli*. *Journal of
Bacteriology*, *181*(17), 5167–5175.

Bolla, J. R., Sauer, J. B., Wu, D., Mehmood, S., Allison, T. M., & Robinson, C. V.
(2018). Direct observation of the influence of cardiolipin and antibiotics on lipid
II binding to MurJ. *Nature Chemistry*, (January).
https://doi.org/10.1038/nchem.2919

Bouhss, A., Mengin-Lecreulx, D., Le Beller, D., & Van Heijenoort, J. (1999).
Topological analysis of the MraY protein catalysing the first membrane step of
peptidoglycan synthesis. *Molecular Microbiology*, *34*(3), 576–585.
https://doi.org/10.1046/j.1365-2958.1999.01623.x

Bouhss, A., Trunkfield, A. E., Bugg, T. D. H., & Mengin-Lecreulx, D. (2008). The

biosynthesis of peptidoglycan lipid-linked intermediates. *FEMS Microbiology*
Reviews. https://doi.org/10.1111/j.1574-6976.2007.00089.x

Buddelmeijer, N., Aarsman, M., & Blaauwen, T. den. (2013). Immunolabeling of
Proteins in situ in *Escherichia coli* K12 Strains. *Bio-Protocol*, *3*(15), 1–5.
https://doi.org/10.21769/BioProtoc.852

Butler, E. K., Davis, R. M., Bari, V., Nicholson, P. A., & Ruiz, N. (2013). Structure-
function analysis of MurJ reveals a solvent-exposed cavity containing residues
essential for peptidoglycan biogenesis in *Escherichia coli*. *Journal of*
Bacteriology, *195*(20), 4639–4649. https://doi.org/10.1128/JB.00731-13

Butler, E. K., Tan, W. B., Joseph, H., & Ruiz, N. (2014). Charge requirements of lipid
II flippase activity in *Escherichia coli*. *Journal of Bacteriology*, *196*(23), 4111–
4119. https://doi.org/10.1128/JB.02172-14

Chamakura, K. R., Sham, L.-T., Davis, R. M., Min, L., Cho, H., Ruiz, N., … Young,
R. (2017). A viral protein antibiotic inhibits lipid II flippase activity. *Nature*
Microbiology. https://doi.org/10.1038/s41564-017-0023-4

Chen, J. C., & Beckwith, J. (2008). FtsQ, FtsL and FtsI require FtsK, but not FtsN, for
co-localization with FtsZ during *Escherichia coli* cell division. *Molecular*
Microbiology. https://doi.org/https://doi.org/10.1046/j.1365-2958.2001.02640.x

Chen, X., Zaro, J. L., & Shen, W. C. (2013). Fusion protein linkers: Property, design
and functionality. *Advanced Drug Delivery Reviews*. Elsevier B.V.
https://doi.org/10.1016/j.addr.2012.09.039

Contreras-Martel, C., Martins, A., Ecobichon, C., Trindade, D. M., Matteï, P.-J.,
Hicham, S., … Dessen, A. (2017). Molecular architecture of the PBP2–MreC
core bacterial cell wall synthesis complex. *Nature Communications*, *8*(1), 776.
https://doi.org/10.1038/s41467-017-00783-2

Cranfill, P. J., Sell, B. R., Baird, M. A., Allen, J. R., Lavagnino, Z., de Gruiter, H. M.,
… Piston, D. W. (2016). Quantitative assessment of fluorescent proteins. *Nature*
Methods, *13*(7), 557–562. https://doi.org/10.1038/nmeth.3891

Dai, K., & Lutkenhaus, J. (1991). *ftsZ* is an essential cell division gene in *Escherichia*
coli. *J Bacteriol*, *173*(11), 3500–3506. https://doi.org/10.1046/j.1365-
2958.1997.4091773.x

Datsenko, K., & Wanner, B. (2000). One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proc. Nat. Acad. Sci. (USA)*, *971242*,
6640–6645. https://doi.org/https://doi.org/10.1073/pnas.120163297

de Roubin, M. R., Mengin-Lecreulx, D., & van Heijenoort, J. (1992). Peptidoglycan
biosynthesis in *Escherichia coli*: variations in the metabolism of alanine and D-
alanyl-D-alanine. *Journal of General Microbiology*, *138 Pt 8*(1 992), 1751–7.
https://doi.org/10.1099/00221287-138-8-1751

Den Blaauwen, T., Aarsman, M. E. G., Vischer, N. O. E., & Nanninga, N. (2003).
Penicillin-binding protein PBP2 of *Escherichia coli* localizes preferentially in the
lateral wall and at mid-cell in comparison with the old cell pole. *Molecular*
Microbiology, *47*(2), 539–547. https://doi.org/10.1046/j.1365-
2958.2003.03316.x

Den Blaauwen, T., De Pedro, M. A., Nguyen-Distèche, M., & Ayala, J. A. (2008).
Morphogenesis of rod-shaped sacculi. *FEMS Microbiology Reviews*, *32*(2), 321–
344. https://doi.org/10.1111/j.1574-6976.2007.00090.x

den Blaauwen, T., Hamoen, L. W., & Levin, P. A. (2017). The divisome at 25: the
road ahead. *Current Opinion in Microbiology*, *36*, 85–94.
https://doi.org/10.1016/j.mib.2017.01.007

Derouichei, R., Bénédetti, H., Lazzaroni, J. C., Lazdunski, C., & Lloubès, R. (1995).

Protein complex within *Escherichia coli* inner membrane: TolA N-terminal domain interacts with TolQ and TolR proteins. *Journal of Biological Chemistry*, *270*(19), 11078–11084. https://doi.org/10.1074/jbc.270.19.11078

Du, S., Pichoff, S., & Lutkenhaus, J. (2016). FtsEX acts on FtsA to regulate divisome assembly and activity. *Proceedings of the National Academy of Sciences*, 201606656. https://doi.org/10.1073/pnas.1606656113

Dubuisson, J. F., Vianney, A., & Lazzaroni, J. C. (2002). Mutational Analysis of the TolA C-Terminal Domain of *Escherichia coli* and Genetic Evidence for an Interaction between TolA and TolB. *Journal of Bacteriology*, *184*(16), 4620–4625. https://doi.org/10.1128/JB.184.16.4620-4625.2002

Egan, A. J. F., & Vollmer, W. (2015). The stoichiometric divisome: A hypothesis. *Frontiers in Microbiology*, *6*(MAY), 1–6. https://doi.org/10.3389/fmicb.2015.00455

Elhenawy, W., Davis, R. M., Fero, J., Salama, N. R., Felman, M. F., & Ruiz, N. (2016). The o-antigen flippase wzk can substitute for MurJ in peptidoglycan synthesis in *helicobacter pylori* and *Escherichia coli*. *PLoS ONE*, *11*(8), 1–16. https://doi.org/10.1371/journal.pone.0161587

Fay, A., & Dworkin, J. (2009). B*acillus subtilis* homologs of MviN (MurJ), the putative *Escherichia coli* lipid II flippase, are not essential for growth. *Journal of Bacteriology*, *191*(19), 6020–6028. https://doi.org/10.1128/JB.00605-09

Fenton, A. K., Mortaji, L. El, Lau, D. T. C., Rudner, D. Z., & Bernhardt, T. G. (2016). CozE is a member of the MreCD complex that directs cell elongation in *Streptococcus pneumoniae*. *Nature Microbiology*, *2*(December 2016), 16237. https://doi.org/10.1038/nmicrobiol.2016.237

Gerding, M. A., Liu, B., Bendezú, F. O., Hale, C. A., Bernhardt, T. G., & De Boer, P. A. J. (2009). Self-enhanced accumulation of FtsN at division sites and roles for other proteins with a SPOR domain (DamX, DedD, and RlpA) in *Escherichia coli* cell constriction. *Journal of Bacteriology*, *191*(24), 7383–7401. https://doi.org/10.1128/JB.00811-09

Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A., & Smith, H. O. (2009). Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nature Methods*, *6*(5), 343–345. https://doi.org/10.1038/nmeth.1318

Goehring, N. W., Gonzalez, M. D., & Beckwith, J. (2006). Premature targeting of cell division proteins to midcell reveals hierarchies of protein interactions involved in divisome assembly. *Molecular Microbiology*, *61*(1), 33–45. https://doi.org/10.1111/j.1365-2958.2006.05206.x

Goehring, N. W., Gueiros-filho, F., & Beckwith, J. (2005). Premature targeting of a cell division protein to midcell allows dissection of divisome assembly in *Escherichia coli*. *Genes & Development*, *19*(1), 127–137. https://doi.org/10.1101/gad.1253805

Gray, A. N., Egan, A. J. F., van't Veer, I. L., Verheul, J., Colavin, A., Koumoutsi, A., … Vollmer, W. (2015). Coordination of peptidoglycan synthesis and outer membrane constriction during *Escherichia coli* cell division. *eLife*, *4*(MAY), 1–29. https://doi.org/10.7554/eLife.07118

Hong, Y., Liu, M. A., & Reeves, P. R. (2018). Progress in our understanding of Wzx flippase for translocation of bacterial membrane lipid-linked oligosaccharide. *Journal of Bacteriology*, *200*(1), 1–14. https://doi.org/10.1128/JB.00154-17

Huang, Y. (2003). Structure and Mechanism of the Glycerol-3-Phosphate Transporter from *Escherichia coli*. *Science*, *301*(5633), 616–620. https://doi.org/10.1126/science.1087619

Ikeda, M., Sato, T., Wachi, M., Jung, H. K., Ishino, F., Kobayashi, Y., & Matsuhashi, M. (1989). Structural similarity among *Escherichia coli* FtsW and RodA proteins and *Bacillus subtilis* SpoVE protein, which function in cell division, cell elongation, and spore formation, respectively. *Journal of Bacteriology*, *171*(11), 6375–6378. https://doi.org/10.1128/jb.171.11.6375-6378.1989

Ishino, F., Hai Kwan Jung, Ikeda, M., Doi, M., Wachi, M., & Matsuhashi, M. (1989). New mutations *fts-36*, *lts-33*, and *ftsW* clustered in the mra region of the *Escherichia coli* chromosome induce thermosensitive cell growth and division. *Journal of Bacteriology*, *171*(10), 5523–5530. https://doi.org/10.1128/jb.171.10.5523-5530.1989

Khattar, M. M., Begg, K. J., & Donachie, W. D. (1994). Identification of FtsW and characterization of a new ftsW division mutant of *Escherichia coli*. *Journal of Bacteriology*, *176*(23), 7140–7147. https://doi.org/10.1128/jb.176.23.7140-7147.1994

Kuk, A. C. Y., Mashalidis, E. H., & Lee, S. Y. (2017). Crystal structure of the MOP flippase MurJ in an inward-facing conformation. *Nature Structural and Molecular Biology*, *24*(2), 171–176. https://doi.org/10.1038/nsmb.3346

Leclercq, S., Derouaux, A., Olatunji, S., Fraipont, C., Egan, A. J. F., Vollmer, W., … Terrak, M. (2017). Interplay between Penicillin-binding proteins and SEDS proteins promotes bacterial cell wall synthesis. *Scientific Reports*. https://doi.org/10.1038/srep43306

Li, G. W., Burkhardt, D., Gross, C., & Weissman, J. S. (2014). Quantifying absolute protein synthesis rates reveals principles underlying allocation of cellular resources. *Cell*, *157*(3), 624–635. https://doi.org/10.1016/j.cell.2014.02.033

Liu, B., Persons, L., Lee, L., & de Boer, P. A. J. (2015). Roles for both FtsA and the FtsBLQ subcomplex in FtsN-stimulated cell constriction in *Escherichia coli*. *Molecular Microbiology*, *95*(6), 945–970. https://doi.org/10.1111/mmi.12906

Lutkenhaus, J. (2009). FtsN - Trigger for septation. *Journal of Bacteriology*, *191*(24), 7381–7382. https://doi.org/10.1128/JB.01100-09

Macheboeuf, P., Contreras-Martel, C., Job, V., Dideberg, O., & Dessen, A. (2006). Penicillin binding proteins: Key players in bacterial cell cycle and drug resistance processes. *FEMS Microbiology Reviews*, *30*(5), 673–691. https://doi.org/10.1111/j.1574-6976.2006.00024.x

Manat, G., Roure, S., Auger, R., Bouhss, A., Barreteau, H., Mengin-Lecreulx, D., & Touzé, T. (2014). Deciphering the Metabolism of Undecaprenyl-Phosphate: The Bacterial Cell-Wall Unit Carrier at the Membrane Frontier. *Microbial Drug Resistance*, *20*(3), 199–214. https://doi.org/10.1089/mdr.2014.0035

Matsuzawa, H., Hayakawa, K., Sato, T., & Imahori, K. (1973). Characterization and genetic analysis of a mutant of *Escherichia coli* K-12 with rounded morphology. *J. Bacteriol.*, *115*(1), 436–442.

Meeske, A. J., Riley, E. P., Robins, W. P., Uehara, T., Mekalanos, J. J., Kahne, D., … Rudner, D. Z. (2016). SEDS proteins are a widespread family of bacterial cell wall polymerases. *Nature*, 1–15. https://doi.org/10.1038/nature19331

Meeske, A. J., Sham, L.-T., Kimsey, H., Koo, B.-M., Gross, C. a, Bernhardt, T. G., & Rudner, D. Z. (2015). MurJ and a novel lipid II flippase are required for cell wall biogenesis in *Bacillus subtilis*. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(20), 6437–42. https://doi.org/10.1073/pnas.1504967112

Meiresonne, N. Y., van der Ploeg, R., Hink, M. A., & den Blaauwen, T. (2017). Activity-Related Conformational Changes in d,d-Carboxypeptidases Revealed

by In Vivo Periplasmic Förster Resonance Energy Transfer Assay in *Escherichia*
coli. *mBio*. https://doi.org/10.1128/mBio.01089-17
Mengin-lecreulx, D., Flouret, B., Heijenoort, J. V. A. N., S, E. R. C. N. R., Biochimie,
I. De, & Paris-sud, U. (1983). Pool Levels of UDP N-Acetylglucosamine and
UDP N- Acetylglucosamine-Enolpyruvate in *Escherichia coli* and Correlation
with Peptidoglycan Synthesis. *Journal of Bacteriology*, *154*(3), 1284–1290.
https://doi.org/0021-9193/83/061284-07$02.00/0
Mengin-Lecreulx, D., Texier, L., Rousseau, M., & Van Heijenoort, J. (1991). The
murG gene of Escherichia coli codes for the UDP-N-acetylglucosamine:N-
acetylmuramyl-(pentapeptide) pyrophosphoryl-undecaprenol N-
acetylglucosamine transferase involved in the membrane steps of peptidoglycan
synthesis. *Journal of Bacteriology*, *173*(15), 4625–4636.
https://doi.org/10.1128/JB.173.15.4625-4636.1991
Mohammadi, T., Karczmarek, A., Crouvoisier, M., Bouhss, A., Mengin-Lecreulx, D.,
& Den Blaauwen, T. (2007). The essential peptidoglycan glycosyltransferase
MurG forms a complex with proteins involved in lateral envelope growth as well
as with proteins involved in cell division in *Escherichia coli*. *Molecular*
Microbiology, *65*(4), 1106–1121. https://doi.org/10.1111/j.1365-
2958.2007.05851.x
Mohammadi, T., Sijbrandi, R., Lutters, M., Verheul, J., Martin, N. I., Den Blaauwen,
T., … Breukink, E. (2014). Specificity of the transport of lipid II by FtsW in
Escherichia coli. *Journal of Biological Chemistry*, *289*(21), 14707–14718.
https://doi.org/10.1074/jbc.M114.557371
Mohammadi, T., van Dam, V., Sijbrandi, R., Vernet, T., Zapun, A., Bouhss, A., …
Breukink, E. (2011). Identification of FtsW as a transporter of lipid-linked cell
wall precursors across the membrane. *The EMBO Journal*, *30*(8), 1425–1432.
https://doi.org/10.1038/emboj.2011.61
Morgenstein, R. M., Bratton, B. P., Nguyen, J. P., Ouzounov, N., Shaevitz, J. W., &
Gitai, Z. (2015). RodZ links MreB to cell wall synthesis to mediate MreB
rotation and robust morphogenesis. *Proc Natl Acad Sci U S A*, *112*(40), 12510–
12515. https://doi.org/10.1073/pnas.1509610112
Neuhaus, F. C. (1962). The enzymatic synthesis of D-alanyl-D-alanine. I. Purification
and properties of D-alanyl-D-alanine synthetase. *Journal of Biological*
Chemistry, *237*(3).
Neuhaus, F. C., & Lynch, J. L. (1972). Studies on the inhibition of D-alanyl-D-
alanine synthetase by the antibiotic D-cycloserine. *BIOCHEMICAL AND*
BIOPHYSICAL RESEARCH COMMUNICATIONS, *8*(5), 377–382.
https://doi.org/https://doi.org/10.1016/0006-291X(62)90011-6
Nørholm, M. H. H. (2010). A mutant Pfu DNA polymerase designed for advanced
uracil-excision DNA engineering. *BMC Biotechnology*, *10*(1), 21.
https://doi.org/10.1186/1472-6750-10-21
Ortiz, C., Natale, P., Cueto, L., & Vicente, M. (2015). The keepers of the ring:
Regulators of FtsZ assembly. *FEMS Microbiology Reviews*, *40*(1), 57–67.
https://doi.org/10.1093/femsre/fuv040
Oswald, F., Varadarajan, A., Lill, H., Peterman, E. J. G., & Bollen, Y. J. M. (2016).
MreB-Dependent Organization of the E. coli Cytoplasmic Membrane Controls
Membrane Protein Diffusion. *Biophysical Journal*, *110*(5), 1139–1149.
https://doi.org/10.1016/j.bpj.2016.01.010
Pastoret, S., Fraipont, C., Blaauwen, T. Den, Aarsman, M. E. G., Thomas, A.,
Brasseur, R., & Nguyen-diste, M. (2004). Functional Analysis of the Cell

Division Protein FtsW of *Escherichia coli*. *Journal of Bacteriology*, *186*(24), 8370–8379. https://doi.org/10.1128/JB.186.24.8370

Pedro, M. A. De, Quintela, J. C., Höltje, J. V, & Schwarz, H. (1997). Murein segregation in *Escherichia coli*. *Microbiology*, *179*(9), 2823–2834. https://doi.org/doi: 10.1128/jb.179.9.2823-2834.1997

Pichoff, S., & Lutkenhaus, J. (2002). Unique and overlapping roles for ZipA and FtsA in septal ring assembly in *Escherichia coli*. *EMBO Journal*, *21*(4), 685–693. https://doi.org/10.1093/emboj/21.4.685

Pichoff, S., & Lutkenhaus, J. (2005). Tethering the Z ring to the membrane through a conserved membrane targeting sequence in FtsA. *Molecular Microbiology*, *55*(6), 1722–1734. https://doi.org/10.1111/j.1365-2958.2005.04522.x

Pomorski, T., & Menon, A. K. (2006). Lipid flippases and their biological functions. *Cellular and Molecular Life Sciences*, *63*(24), 2908–2921. https://doi.org/10.1007/s00018-006-6167-7

Priyadarshini, R., De Pedro, M. A., & Young, K. D. (2007). Role of peptidoglycan amidases in the development and morphology of the division septum in *Escherichia coli*. *Journal of Bacteriology*, *189*(14), 5334–5347. https://doi.org/10.1128/JB.00415-07

Prosser, G. a, & de Carvalho, L. P. S. (2013). Kinetic mechanism and inhibition of *Mycobacterium tuberculosis* D-alanine:D-alanine ligase by the antibiotic D-cycloserine. *The FEBS Journal*, *280*(4), 1150–66. https://doi.org/10.1111/febs.12108

Qiao, Y., Srisuknimit, V., Rubino, F., Schaefer, K., Ruiz, N., Walker, S., & Kahne, D. (2017). Lipid II overproduction allows direct assay of transpeptidase inhibition by [beta]-lactams. *Nature Chemical Biology*, *13*(7), 793–798. https://doi.org/10.1038/nchembio.2388

Rico, A. I., García-Ovalle, M., Mingorance, J., & Vicente, M. (2004). Role of two essential domains of *Escherichia coli* FtsA in localization and progression of the division ring. *Molecular Microbiology*, *53*(5), 1359–1371. https://doi.org/10.1111/j.1365-2958.2004.04245.x

Rowlett, V. W., & Margolin, W. (2015). The Min system and other nucleoid-independent regulators of Z ring positioning. *Frontiers in Microbiology*, *6*(MAY), 1–10. https://doi.org/10.3389/fmicb.2015.00478

Ruiz, N. (2008). Bioinformatics identification of MurJ (MviN) as the peptidoglycan lipid II flippase in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(40), 15553–7. https://doi.org/10.1073/pnas.0808352105

Ruiz, N. (2009). *Streptococcus pyogenes* YtgP (Spy-0390) complements *Escherichia coli* strains depleted of the putative peptidoglycan flippase MurJ. *Antimicrobial Agents and Chemotherapy*, *53*(8), 3604–3605. https://doi.org/10.1128/AAC.00578-09

Ruiz, N. (2015). Lipid Flippases for Bacterial Peptidoglycan Biosynthesis. *Lipid Insights*, *8*(Suppl 1), 21–31. https://doi.org/10.4137/LPI.S31783

Ruiz, N. (2016). Filling holes in peptidoglycan biogenesis of *Escherichia coli*. *Current Opinion in Microbiology*, *34*, 1–6. https://doi.org/10.1016/j.mib.2016.07.010

Sauvage, E., Kerff, F., Terrak, M., Ayala, J. A., & Charlier, P. (2008). The penicillin-binding proteins: Structure and role in peptidoglycan biosynthesis. *FEMS Microbiology Reviews*, *32*(2), 234–258. https://doi.org/10.1111/j.1574-6976.2008.00105.x

Scheffers, D. J., & Tol, M. B. (2015). LipidII: Just Another Brick in the Wall? *PLoS Pathogens*, *11*(12), 1–12. https://doi.org/10.1371/journal.ppat.1005213

Sham, L.-T., Butler, E. K., Lebar, M. D., Kahne, D., Bernhardt, T. G., & Ruiz, N. (2014). MurJ is the flippase of lipid-linked precursors for peptidoglycan biogenesis. *Science*, *345*(6193), 220–222. https://doi.org/10.1126/science.1254522

Shih, Y. L., Kawagishi, I., & Rothfield, L. (2005). The MreB and Min cytoskeletal-like systems play independent roles in prokaryotic polar differentiation. *Molecular Microbiology*, *58*(4), 917–928. https://doi.org/10.1111/j.1365-2958.2005.04841.x

Sieger, B., Schubert, K., Donovan, C., & Bramkamp, M. (2013). The lipid II flippase RodA determines morphology and growth in *Corynebacterium glutamicum*. *Molecular Microbiology*, *90*(5), 966–982. https://doi.org/10.1111/mmi.12411

Sun, Q., & Margolin, W. (1998). FtsZ Dynamics during the Division Cycle of Live *Escherichia coli* Cells, *180*(8), 2050–2056.

Suzuki, H., Nishimura, Y., & Hirota, Y. (1978). On the process of cellular division in *Escherichia coli*: a series of mutants of *E. coli* altered in the penicillin-binding proteins. *Proceedings of the National Academy of Sciences of the United States of America*, *75*(2), 664–668. https://doi.org/10.1073/pnas.75.2.664

Taschner, P. E., Huls, P. G., Pas, E., & Woldringh, C. L. (1988). Division behavior and shape changes in isogenic *ftsZ*, *ftsQ*, *ftsA*, *pbpB*, and *ftsE* cell division mutants of *Escherichia coli* during temperature shift experiments. *Journal of Bacteriology*, *170*(4), 1533–1540. https://doi.org/doi: 10.1128/jb.170.4.1533-1540.1988

Tsang, M. J., & Bernhardt, T. G. (2015). Guiding divisome assembly and controlling its activity. *Current Opinion in Microbiology*, *24*, 60–65. https://doi.org/10.1016/j.mib.2015.01.002

van der Ploeg, R., Goudelis, S. T., & den Blaauwen, T. (2015). Validation of FRET Assay for the Screening of Growth Inhibitors of *Escherichia coli* Reveals Elongasome Assembly Dynamics. *International Journal of Molecular Sciences*, *16*(8), 17637–54. https://doi.org/10.3390/ijms160817637

Van der Ploeg, R., Verheul, J., Vischer, N. O. E., Alexeeva, S., Hoogendoorn, E., Postma, M., … Den Blaauwen, T. (2013). Colocalization and interaction between elongasome and divisome during a preparative cell division phase in *Escherichia coli*. *Molecular Microbiology*, *87*(5), 1074–1087. https://doi.org/10.1111/mmi.12150

van Teeffelen, S., Wang, S., Furchtgott, L., Huang, K. C., Wingreen, N. S., Shaevitz, J. W., & Gitai, Z. (2011). The bacterial actin MreB rotates, and rotation depends on cell-wall assembly. *Proceedings of the National Academy of Sciences*, *108*(38), 15822–15827. https://doi.org/10.1073/pnas.1108999108

Vischer, N. O. E., Verheul, J., Postma, M., van den Berg van Saparoea, B., Galli, E., Natale, P., … den Blaauwen, T. (2015). Cell age dependent concentration of *Escherichia coli* divisome proteins analyzed with ImageJ and ObjectJ. *Frontiers in Microbiology*, *6*(JUN), 1–18. https://doi.org/10.3389/fmicb.2015.00586

Vollmer, W., Blanot, D., & De Pedro, M. A. (2008). Peptidoglycan structure and architecture. *FEMS Microbiology Reviews*, *32*(2), 149–167. https://doi.org/10.1111/j.1574-6976.2007.00094.x

Vollmer, W., Blanot, D., De Pedro, M. A., Scheffers, D., Pinho, M., Nikolaidis, I., … Dessen, A. (2005). Structural and mechanistic basis of penicillin-binding protein inhibition by lactivicins. *FEMS Microbiology Reviews*, *30*(2), 565–569.

https://doi.org/10.1128/MMBR.69.4.585

Wang, E., Walsh, C., & Walsh, C. (1978). Suicide substrates for the alanine racemase

of *Escherichia coli*. *Biochemistry*, *17*(7), 1313–1321.

https://doi.org/10.1021/bi00600a028

Weiss, D. S. (2015). Last but not least: New insights into how FtsN triggers

constriction during *Escherichia coli* cell division. *Molecular Microbiology*,

95(6), 903–909. https://doi.org/10.1111/mmi.12925

White, C. L., & Gober, J. W. (2012). MreB: Pilot or passenger of cell wall synthesis?

Trends in Microbiology, *20*(2), 74–79. https://doi.org/10.1016/j.tim.2011.11.004

Wood, B. W. A., & Gunsalusa, I. C. (1959). D-Alanine formation; a racemase in

Streptococcus faecalis. *Journal of Biological Chemistry*, *190*, 403–416.

Young, K. D. (2014). A flipping cell wall ferry. *Microbiology*, *345*(6193), 139–140.

https://doi.org/10.1126/science.1256585

Yousif SY, Broome-Smith JK, S. B. (1985). Lysis of *Escherichia coli* by beta-lactam

antibiotics: deletion analysis of the role of penicillin-binding proteins 1A and 1B.

J Gen Microbiol., *131*(10), 2839–45.

1137

1138

1139
Figure legends
Fig. 1. MurJ localizes both in the lateral wall and at midcell. Strain XL08 containing the
chromosomal mNG-MurJ fusion shows a cell-cycle-dependent localization in contrast to the
homogenously localizing integral membrane protein mNG-GlpT. Cells were grown in Gb1
minimal medium to steady state at 28°C. Left: MurJ profiles; Right: membrane control protein
GlpT profiles. (A) Collective profiles of localization of MurJ and GlpT. For each protein, the
diameter (black lines) and fluorescence (green lines) profiles along normalized cell length are
shown in 10 % age class bins. (B) Maps of diameter profiles (magenta) and fluorescence
profiles (green). Cells are plotted by length, ascending from left to right. (C) Phase contrast
image (left) and the corresponding fluorescence image (right) for each protein. Scale bar
equals 5 µm. More than 1200 cells were included for both analyses.

1159

1160

Fig. 2. Timing of MurJ midcell localization. mNG-(GGS)$_2$-MurJ cellular localization was
compared with the localization of divisomal and PG precursor synthesizing proteins. XL08
strain was grown to steady state in Gb1 medium. MurJ localization was determined in living
cells. After fixation with FAGA, the rest of cells were immunolabeled with antibodies against
FtsZ, FtsN, MraY and MurG. Diameter (black lines) and Fluorescence (colored lines) profiles
were plotted into 10 % age class bins along the normalized cell length in %. More than 1200
cells were included.

1168

1169

1170

1171

1172

1173

1174

1175

Fig. 3. MurJ requires the core divisome for its midcell localization. Chromosomal mNG-(GGS)$_2$-MurJ fusion was introduced into strains that harbor temperature sensitive, depletable, or deleted division proteins (except that the mNG-(GGS)$_2$-MurJ plasmid was expressed in *ΔtolA and Δpal strains, and the* mCherry-MurJ plasmid was expressed in *ΔamiABC* strain). (A-E) MurJ localization in XL08 (WT), *ftsA 1882* (Ts), *ftsQ1* (Ts), *ftsW* (Ts) and *ftsI2185* (Ts) backgrounds. Left: Localization at permissive temperature of 28 °C. Right: Localization after growth at the non-permissive temperature of 42 °C for 2 mass doublings (MD)s. (F) MurJ localization in FtsN depletion background at 30 °C. Left: localization in the presence of 0.2% w/v arabinose. Right: Localization of MurJ in the absence of arabinose for 2 MDs. (G-H) mCh-MurJ localization in *ΔmrcA* (PBP1A), *ΔmrcB* (PBP1B), *Δpal, ΔtolA* and *ΔamiABC* background at 30 °C, respectively. Scale bar equals 5 μm.

1192

1193

1194

Fig. 4. Localization of cell division proteins in the presence of the PBP3 inhibitor aztreonam

for 2 MDs. Cell division is inhibited but the localization and assemblage of the divisome

continues, whereas the localization of MurJ is lost. In Gb1 at 28 °C, steady state growing

XL08 was treated with 1 mg.L^{-1} aztreonam, MurJ localization was determined in living cells,

and FtsZ, FtsN, MraY and MurG were immunolabeled with antibodies after fixation. Diameter

(black lines) and fluorescence (colored lines) profiles were plotted into 10 % age class bins

along the normalized cell length. More than 1200 cells were included.

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

Fig. 5 Inhibition of lipid II biogenesis blocks MurJ midcell localization. In Gb1 at 28 °C, steady

state growing XL08 was treated with increasing concentrations of D-cycloserine (DCS), and

mNG-MurJ localization was determined at 30 minutes after addition of DCS. (A) Growth

curves were plotted as time against the natural logarithm of the optical density of the cells.

The black arrow indicates the addition of DCS. (B) Fluorescence maps profiles of MurJ

localization. Cells are plotted according to increasing cell length from top to bottom. From left

to right, the maps correspond to 0, 0.05, 0.1, 0.5, 1 and 2 µg.L^{-1} of DCS concentrations. More

than 1200 cells were included. (C) Phase contrast images (upper) and correspond

fluorescence images (down) of cells treated with DCS. Scale bar equals 5 µm.

1224

1225

1226

1227

1228

1229

Fig. 6. Expression of non-functional FtsW mutants abolishes MurJ midcell localization. XL08 expressing mutant FtsW R145A or FtsW K153N was grown in Gb1 minimal medium to steady state at 28 °C. IPTG (20 μM) was used to induce expression of the FtsW mutants from plasmid. MurJ localization was determined in living cells and the presence of the divisome was confirmed by immunolabeling of FtsN, after fixation. (A) Localization of MurJ and FtsN after expression of FtsW R145A. (B) Localization of MurJ and FtsN after expression of FtsW K153N. For each part: top, average localization profiles plotted along normalized cell length (n ≥1200 cells). Black lines indicate cell diameter. Green lines and red lines indicate MurJ and FtsN localization, respectively. Middle, map profiles of cell diameter and MurJ or FtsN localization sorted by ascending cell length. Bottom, phase contrast and fluorescence microscopy images. Scale bar equals 5 μm.

1248

Fig. 7. MurJ non-functional mutant R18A does not influence its localization. (A) The plasmid

expressing the mNG-MurJ R18A fusion was introduced into LMC500. This strain was grown

in Gb1 minimal medium to steady state at 28 °C, and expression was induced with 20 µM

IPTG for 2 MDs. The diameter (black lines) and fluorescence (green lines) profiles along

normalized cell length are shown in 10 % age class bins. More than 1200 cells were included.

Scale bar equals 5 µm. (B) The plasmid expressing the mNG-MurJ R18A fusion was

introduced into the MurJ depletion strain XL20. Mutant localization was determined in the

presence of arabinose (wild type MurJ expression) or glucose (wild type MurJ depletion).

Scale bar equals 1 µm.

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

Fig. 8. Localization of MurJ cysteine mutants in the presence of MTSES. Strains expressing
MurJ single-cysteine-mutants from plasmid were grown to steady state in Gb1 medium at 28
°C, and expression was induced with IPTG for 2 MDs. MurJ localization was determined at 10
min and 40 min after the addition of MTSES (for mutant S263C, localization was determined
at 10 min and 20 min because of its higher sensitivity to MTSES), Average of MurJ
localization profiles were plotted along normalized cell axis. Black lines indicate the MurJ
localization in the absence of MTSES, green and red lines indicate the MurJ localization at 10
and 40 min in the presence of MTSES, respectively. More than 1200 cells were included for
each experiment. (A-D) Only growth but not localization of functional mutants A29C, S263C,
N49C and E273C is affected by MTSES. (E) Neither growth nor localization of cysteine free
mutants is influenced by MTSES. (F) Neither growth nor localization of wild type MurJ is
influenced by MTSES. (G and H) Localization of non-functional MurJ mutants R18C and
R52C is not affected by MTSES.

1293
Fig. 9. Model of MurJ recruitment and function in septal peptidoglycan synthesis. (A)
Recruitment of MurJ to midcell when FtsW is activated. (B) MurJ binds substrate. (C) MurJ
flips lipid II across the IM by changing its structure from the inward-facing conformation into
the outward-facing state. (D) Unloaded MurJ changes back into the default inward-facing
state to participate in a new cycle. (i) Loss-of-function MurJ mutants that localize but are not
able to flip lipid II. (ii) Functional MurJ mutants that are inactivated by MTSES still localize but
are not able to flip lipid II.
1301
1302
1303

1304 Table 1. Functionality and localization of MurJ fluorescence proteins fusions with different

1305 linkers at N- or C- terminus

Fusions	Plasmids	Complementation[a]	Localization[b]	Function[c]
No fusion	pTHV037-MurJ	≈16h	ND	+
	pSAV057-MurJ	≈16h	ND	+
N-terminus	pTHV037-mCh-(GGS)$_n$-MurJ	≈16h	+	+
	pSAV057-mNG-(GGS)$_n$-MurJ	≈16h	+	+
	pSAV057-mKO-MurJ	≈16h	+	+
C-terminus	pTHV037-MurJ-(GGS)$_n$-mCh	≥3 days	−	−
	pSAV057-MurJ-(GGS)$_n$-mNG	≥3 days	−	−

1306 n: Number of GGS repeats (n = 0, 1, 2, 3)

1307 [a] after transformation, the time period needed for visible white colony appearance.

1308 [b] +, FP fused MurJ is fluorescent and localizes in the membrane; −, FP fused MurJ is

1309 fluorescent and localizes in the cytosol.

1310 [c] +, functional fusion; −, loss-of-function fusion.