Tuning Spatial Profiles of Selection Pressure to Modulate the Evolution of Resistance

Maxwell G. De Jong[1] and Kevin B. Wood[1, 2]

[1]*Department of Physics, University of Michigan, Ann Arbor, Michigan 48109, USA*
[2]*Department of Biophysics, University of Michigan, Ann Arbor, Michigan 48109, USA*[∗]

Spatial heterogeneity plays an important role in the evolution of drug resistance. While recent studies have indicated that spatial gradients of selection pressure can accelerate resistance evolution, much less is known about evolution in more complex spatial profiles. Here we use a stochastic toy model of drug resistance to investigate how different spatial profiles of selection pressure impact the time to fixation of a resistant allele. Using mean first passage time calculations, we show that spatial heterogeneity accelerates resistance evolution when the rate of spatial migration far exceeds that of mutation but slows fixation when mutation dominates. Interestingly, there exists an intermediate regime—characterized by comparable rates of migration and mutation—in which the rate of fixation can be either accelerated or decelerated depending on the spatial profile, even when spatially averaged selection pressure remains constant. Finally, we demonstrate that optimal tuning of the spatial profile can dramatically slow the spread and fixation of resistant subpopulations, which may lay the groundwork for optimized, spatially-resolved drug dosing strategies for mitigating the effects of drug resistance.

Drug resistance is a rapidly growing public health threat and a central impediment to the treatment of cancer, viruses, and microbial infections [1–4]. The battle against resistance has been largely fought at the molecular level, leading to an increasingly mature understanding of its underlying biochemical and genetic roots. At the same time, evolutionary biologists have long recognized resistance as a fundamentally stochastic process governed by the complex interplay between microbial ecology and evolutionary selection. The last decade, in particular, has seen a significant surge in efforts to develop and understand evolution-based treatment strategies to forestall resistance [5–16]. While the vast majority of this work focuses on spatially homogeneous environments, a number of recent studies, both theoretical and experimental, have demonstrated that spatial heterogeneity in drug concentration can dramatically alter the evolutionary dynamics leading to resistance [16–24]. On a practical level, the picture that emerges is somewhat bleak, as resistance evolution is dramatically accelerated in the presence of spatial gradients in drug concentration [18–20, 22–24] or heterogeneous drug penetration [17, 21]. Interestingly, however, recent work shows that this acceleration can be tempered or even reversed when the mutational pathway (i.e. the genotypic fitness landscape) leading to resistance contains fitness valleys [18], which are known to inhibit evolution [25–28]. Unfortunately, because the fitness landscape is a genetic property of the cells themselves, the potential for accelerated evolution appears to be "built in", making it difficult to combat in a treatment setting. However, these results raise the question of whether non-monotonic profiles of tunable properties of the system—for example, the spatial selection pressure— might also have the potential to slow evolution, even when the mutational pathway lacks the requisite fitness valleys.

Evolution in natural or clinical settings takes place in heterogeneous environments characterized by spatial fluctuations in multiple factors, including drug concentrations, nutrients, temperature, pH, and host immune responses, all of which potentially affect cellular growth. Understanding evolution and ecology in such spatially extended systems is a challenging and long-studied problem [29–31]. Recent studies have demonstrated rich dynamics when inter-cellular interactions are defined on heterogeneous complex networks [32–34], where spatial structure can (for example) promote invasive strategies in tumor models [33] or modulate fixation times on random landscapes [32]. Remarkably, in the weak selection limit, evolutionary dynamics can be solved for any population structure [34], providing extensive insight into game-theoretic outcomes on complex networks. In addition, theoretical tools from statistical physics have proven useful for understanding spatiotemporal dynamics in spatially structured populations in a wide range of contexts, including biologically-inspired Monte Carlo models [18], toy models of source-sink dynamics [19], stepping-stone models of spatial pattern formation [35], models of dispersion [36–38], and Moran meta-population models [39–41]. In a similar spirit, here we use stochastic models of evolution along with theoretical tools from statistical physics to investigate the effects of spatially heterogeneous fitness pressures on the evolution of resistance. In contrast to previous models defined on heterogeneous networks at the single-cell level, here we consider meta-populations connected via simple topologies and investigate the effects of spatial structure imposed by arbitrary distributions of selection pressure. While several elegant approaches exist for studying these models in particular limits (e.g. with a center manifold reduction) [39–41], here we instead use a classical mean first passage time approach based on adjoint equations to reduce the calculation of mean fixation times to a simple collection of linear equations that can be easily solved for arbitrary

spatial distributions of selection pressures. This method also allows us to find the fixation times from arbitrary initial states, which are often difficult to compute using other methods. Using this approach, we show that resistance evolution can be either accelerated or decelerated by spatial heterogeneities in selection pressure, even when the spatially averaged selection pressure remains constant. In addition, we demonstrate that tuning the spatial distribution of selection pressure can dramatically slow fixation when the subpopulations of resistant mutants are not uniformly distributed in space.

To investigate resistance evolution on a spatially heterogeneous landscape, we first consider a stochastic Moran-like model [42] of a finite population (N) consisting of ($N - n^*$) wild-type cells with fitness $r_0 \leq 1$ and n^* drug-resistant mutants with fitness r^*, which we set to unity without loss of generality. At each time step, cells are randomly selected for birth and death, with cells of higher fitness (in this case, resistant cells) chosen preferentially for division (see SI for full model with transition rates). Wild-type cells can mutate to become drug resistant at rate μ; we neglect reverse transitions to the drug-sensitive state. To incorporate spatial heterogeneity, we consider a simple spatially extended system with M distinct microhabitats, each containing N cells; cells are allowed to migrate at rate β between connected microhabitats (Fig. 1a). At each time t, the state of the system is characterized by the vector $n^*(x_i)$ whose components correspond to the number of mutants in each discrete microhabitat $x_i = 0, 1, ..., M - 1$. The system evolves according to a continuous time master equation

$$\frac{dP_m}{dt} = \sum_{m'} \Omega_{mm'} P_{m'}, \qquad (1)$$

where m and m' denote different states of the system and Ω is a $N^M \times N^M$ matrix whose entries depend on the wild-type fitness value $r_0(x_i)$ at each spatial location x_i (see SI). For tractability, we restrict our analysis to $M = 3$, which is the simplest model that allows for potentially non-monotonic fitness landscapes, such as fitness peaks and fitness valleys. In what follows, we refer to the vector $s(x_i) \equiv 1 - r_0(x_i)$ as the spatial profile of selection pressure, as it measures the difference in fitness between resistant and wild-type cells in each microhabitat (x_i). Intuitively, large values of $s(x_i)$ correspond to regions where the resistant mutant has a significant evolutionary advantage over the wild-type cells (e.g. regions of high drug concentration).

While Equation 1 is difficult to solve explicitly, it is straightforward to calculate quantities that describe the evolution of resistance in various spatial profiles. First, we note that the model consists of a single absorbing state—the fully resistant state ($n^*(x_i) = N$ for all x_i)—and the system will asymptotically approach this state. To characterize the speed of fixation in the presence of

different spatial profiles $s(x_i)$, we calculate the mean first passage times (MFTPs) between states, which obey [43, 44]

$$-1 = \sum_{m' \neq m_f} T(m_f | m') \Omega_{m', m_i} \qquad (2)$$

where $T(m_f | m_i)$ is the mean time required for a system initially in state m_i to first reach state m_f. We take m_f to be the fully resistant state ($n^*(x_i) = N$ for all x_i) and solve the coupled set of linear equations for $\tau_f^j \equiv T(m_f | j)$, where j is an index that runs over all initial states. In particular, when j is the fully wild-type population ($n^*(x_i) = 0$ for all x_i), we refer to the MFPT as the mean fixation time τ_f.

In the case of a single microhabitat, the mean fixation time τ_f would be expected to increase as selection pressure decreases (see SI for calculation). In the spatially extended case, τ_f would also be expected to increase when the selection pressure is globally decreased, though it should also depend on the spatial structure of the specific selection profile $s(x_i)$. To investigate the impact of spatial structure alone, we compared τ_f across different selection profiles $s(x_i)$, all of which were characterized by the same spatially averaged selection pressure, $\langle s \rangle = \sum_i s(x_i)/M$. For simplicity, we begin with a symmetric profile characterized by a background selection pressure s_0 in the edge habitats and a relative peak of height δs in the center habitat (Fig. 1a). This toy landscape has an average selection pressure of $\langle s \rangle = s_0 + \delta s/M$, and the parameters s_0 and δs are constrained by the fact that $0 \leq s(x_i) \leq 1$ at all spatial locations. We vary δs systematically to explore different selection profiles, which can include a single selection pressure valley ($\delta s < 0$), a homogeneous landscape ($\delta s = 0$), or a single selection pressure peak ($\delta s > 0$).

Interestingly, we find that modulating heterogeneity (δs) can increase or decrease τ_f for certain choices of migration and mutation rates, even when $\langle s \rangle$ is held constant (Fig. 1b). More generally, we find that the $\beta - \mu$ plane can be divided into three non-overlapping regions where the homogeneous landscape 1) leads to the smallest value of τ_f, 2) leads to the largest value of τ_f, or 3) does not correspond to an extremum τ_f (Fig. 2a). While we restrict ourselves primarily to $N = 25$, $\langle s \rangle = 0.167$, and to symmetric landscapes, we find qualitatively similar results (i.e. 3 distinct regions) for other values of $\langle s \rangle$ (Fig. S1), N (Fig. S2), as well as for permuted selection profiles (Fig. S3), globally coupled profiles (Fig. S3), and monotonic (gradient) selection profiles (Fig. S4).

Intuitively, the dynamics leading to fixation are dominated by three main timescales: τ_1: the time to have a stable, small population of mutants; τ_2: the time required for this population to grow to dominate the system; and τ_3: the time required for this large mutant population to become fixed in the population. The fixation time is

FIG. 1. (a) Stochastic model for emergence and spread of resistant cells (red) in a spatially extended population of sensitive cells (green). Each spatial habitat (x_i) contains N total cells. Cells migrate at a rate β between neighboring habitats, and sensitive cells mutate at a rate μ to resistant cells. The spatial distribution of selection pressure is characterized by a background value (s_0) and a peak height (δs). (b) Example plot of the mean fixation time for different landscapes with $\mu = 5 \times 10^{-3}$, $\beta = 0.08$, $N = 25$, and $\langle s \rangle = 0.167$. The time to fixation can be either faster (green) or slower (red) than the spatially homogeneous landscape with $\delta s = 0$. Inset: selection landscapes for $\delta s = -0.2$ and $\delta s = 0.5$.

thus the sum of these three different timescales, which are set by the selection pressure landscape as well as the parameters μ and β. In the limit $\beta << \mu$, heterogeneity always increases fixation time (Fig. 2b). In this regime, the three microhabitats are essentially independent, and τ_f becomes the slowest of three independent fixation times. Spatial heterogeneity necessarily results in a decreased selection pressure in at least one microhabitat, which in turn slows fixation (here dominated by τ_3).

In the limit $\mu << \beta$, any spatial heterogeneity leads to faster fixation (Fig. 2c). In this limit of small mutation rate, fixation is dominated by the time it takes to acquire resistant mutants—τ_f is dominated by τ_1. In this sense, the initial seed mutants are evolutionarily "expensive", and it is advantageous to have microhabitats with large selection pressures to prevent loss of the mutant by genetic drift. Spatial heterogeneity leads to increased selection pressure in at least one microhabitat and therefore decreases fixation time in this region (see also SI, including Fig. S6 and Fig. S7).

In general, spatial heterogeneity produces two conflict-

FIG. 2. Spatial heterogeneity can speed up or slow fixation depending on the rates of migration and mutation. (a) Phase diagram illustrates region of parameter space where the homogeneous landscape leads to a maximum (light blue), minimum (dark blue) or intermediate (medium blue) value of the in fixation time. Lower panels show sample fixation curves in the regions where any heterogeneity slows fixation (b, $\beta = 10^{-4}$, $\mu = 10^{-3}$) or accelerates fixation (c, $\beta = 10^{-2}$, $\mu = 10^{-6}$). $N = 25$ and $\langle s \rangle = 0.167$ in all panels.

ing influences on the fixation time. As spatial heterogeneity increases, the selection pressure of the maximum selection pressure microhabitat further increases, which allows mutants to safely emerge and decreases τ_1. But the selection pressure of the minimum selection pressure microhabitat necessarily decreases, which increases τ_3. In the two limiting cases of the parameters, one of these timescales dominates the fixation time, resulting in intuitive results. But outside of these limits, the fixation time is not dominated by a single timescale, resulting in non-trivial selection landscapes leading to the fastest fixation. To further investigate evolution in the intermediate regime, we calculated the minimum and maximum fixation times (τ_f^{min} and τ_f^{max}, respectively) as δs is modulated to create different selection pressure distributions (Fig. 3a). While in many cases the fixation time is decreased by only a few percent, we do find larger effects in the high and low migration limits (i.e. on the edges) of the intermediate regime. To understand how the spatial τ_f profiles change over this intermediate region, we fix μ and traverse across a trajectory in β as shown by the arrow in Fig. 3a. As we increase β, τ_f smoothly transitions from being minimized at $\delta s = 0$ to being maximized near $\delta s = 0$ (Fig. 3b).

Our results indicate that a judicious choice of selec-

FIG. 3. (a) Minimum fixation times (τ_f^{min}) over different se-
lection pressure distributions (relative to the maximum fixa-
tion times, τ_f^{max}) in the intermediate parameter region where
fixation can be both accelerated and decelerated. $N = 25$
and $\langle s \rangle = 0.167$. (b) Across a specific trajectory in the inter-
mediate region (arrow in panel (a)), the dependence of τ_f on
heterogeneity (δs) transitions smoothly from a state with a
minimum at $\delta s = 0$ (lightest curve) to one with a maximum
at $\delta s = 0$ (darkest curve). For ease of comparison, fixation
times are scaled to arbitrary units between 0 and 1.

tion pressure profile can potentially slow fixation of de
novo mutants. In addition, selection pressure profiles
can be optimized to mitigate the effects of resistance
once it has emerged. One advantage of the MFPT ap-
proach (i.e. solving Equation 2) is that it provides fixa-
tion times starting from all possible initial states, mak-
ing it straightforward to apply to cases where a resistant
subpopulation already exists. Specifically, consider a sit-
uation where a resistant subpopulation has arisen at a
particular spatial location. Is it possible to choose the
spatial distribution of selection pressure—for example,
by spatially dosing the drug—to minimize the time to
fixation from this state? As an illustrative example, we
consider a population consisting of $N/2$ mutants in the
center microhabitat and calculate the mean time to fix-
ation for different spatial profiles of selection pressure.
We then find the optimal value for δs—that is, the het-
erogeneity corresponding to the spatial landscape with
the slowest fixation time—in different regions of param-
eter space (Fig. 4a). The specific choice of spatial profile
significantly impacts the time to fixation from the initial
resistant subpopulation (Fig. 4b). We observe two dis-
tinct regions of parameter space that lead to two very
different dosing regimes (Fig. 4c). In the region $\mu >> \beta$,
slowest fixation occurs when we maximize the amount
of drug in the center microhabitat ($\delta s = 0.5$). On the
other hand, in the regime $\beta >> \mu$ fixation is optimally
slowed by maximizing the amount of drug in the two mi-
crohabitats without any initial mutants ($\delta s = -0.2$). In
contrast to the case with no initial mutants (e.g. Fig-
ure 2), fixation time is never maximized by choosing the
homogeneous profile.

We can understand this behavior by fixing a value of β
and changing μ across the two different regimes. When μ
is relatively large compared to β, fixation requires muta-
tions to emerge independently in each of the three micro-

FIG. 4. (a) Schematic: a subpopulation of resistant mutants
(red) arises at a particular spatial location. How can one
choose the spatial distribution of selection pressure (i.e. drug
concentration) to maximize the time to fixation? (b) Hetero-
geneity can significantly speed or slow fixation starting from
an initial resistant subpopulation consisting of $N/2$ cells in
the center habitat ($\mu = 10^{-5}$, $\beta = 8 \times 10^{-3}$). (c) The op-
timal spatial heterogeneity (δs) leading to the slowest mean
fixation time from an initial state of $(0, N/2, 0)$. Depending
on the specific parameter regime, the optimal selection pres-
sure profile is the one with the largest possible valley consis-
tent with $\langle s \rangle$ (black) or the one having the largest possible
peak (white). (d) Relative magnitude of $\tau_f^{\delta s_{max}}$ (mean fixa-
tion time at maximum value of δs) and $\tau_f^{\delta s_{min}}$ (mean fixation
time at minimum value of δs) as mutation rate decreases at
constant migration rate (green arrow, panel (c)). $N = 25$ and
$\langle s \rangle = 0.167$ in all panels.

habitats. As a result, the optimal strategy is to minimize
selection pressure in the microhabitats without mutants,
thereby increasing the fixation times in those spatial re-
gions. As we decrease μ, mutations become increasingly
"expensive" and fixation is achieved most slowly by mini-
mizing the selection pressure in the microhabitat with ini-
tial mutants, since losing this small initial population re-
quires waiting for another random mutation, which hap-
pens with a small probability in this parameter regime.
To further characterize these two regimes, we compare
the fixation times from a maximally peaked landscape
(δs is maximized) to that from a landscape with a large
valley (δs is minimized). The selection landscape that
leads to the slowest fixation rapidly becomes sub-optimal
as mutation rate is decreased at constant β (Fig. 4d).

Our results demonstrate that spatial heterogeneity in
selection pressure can modulate the mean fixation time of
resistant mutants, with homogeneous profiles leading to
maximum, minimum, or intermediate fixation times de-
pending on the rates of mutation and migration. Impor-
tantly, our results predict that spatial heterogeneity in

drug concentrations would impact populations of motile and non-motile cells in opposing ways, even when mutations rates are relatively similar. While heterogeneity is likely to accelerate evolution for populations of motile bacteria, similar to what is observed in experiments with *E. coli* [22, 24], our results predict slowed evolution for less motile cells (e.g. the nosocomial pathogen *E. faecalis* [45]) or cells with rapid mutation rates. Perhaps most interestingly, our results suggest counter-intuitive, spatially optimal profiles for slowing the spread of resistance sub-populations. In the long term, these results may lay the groundwork for optimized, spatially-resolved drug dosing strategies for mitigating the effects of drug resistance.

---

* kbwood@umich.edu

[1] S. B. Levy and B. Marshall, "Antibacterial resistance worldwide: causes, challenges and responses," *Nature medicine*, vol. 10, no. 12s, p. S122, 2004.

[2] C. Holohan, S. Van Schaeybroeck, D. B. Longley, and P. G. Johnston, "Cancer drug resistance: an evolving paradigm," *Nature reviews. Cancer*, vol. 13, no. 10, p. 714, 2013.

[3] M. M. Gottesman, "Mechanisms of cancer drug resistance," *Annual review of medicine*, vol. 53, no. 1, pp. 615–627, 2002.

[4] J. Davies and D. Davies, "Origins and evolution of antibiotic resistance," *Microbiology and Molecular Biology Reviews*, vol. 74, no. 3, pp. 417–433, 2010.

[5] M. Baym, L. K. Stone, and R. Kishony, "Multidrug evolutionary strategies to reverse antibiotic resistance," *Science*, vol. 351, no. 6268, p. aad3292, 2016.

[6] I. Bozic, J. G. Reiter, B. Allen, T. Antal, K. Chatterjee, P. Shah, Y. S. Moon, A. Yaqubie, N. Kelly, D. T. Le, *et al.*, "Evolutionary dynamics of cancer in response to targeted combination therapy," *Elife*, vol. 2, p. e00747, 2013.

[7] Y. Iwasa, M. A. Nowak, and F. Michor, "Evolution of resistance during clonal expansion," *Genetics*, vol. 172, no. 4, pp. 2557–2566, 2006.

[8] E. Toprak, A. Veres, J.-B. Michel, R. Chait, D. L. Hartl, and R. Kishony, "Evolutionary paths to antibiotic resistance under dynamically sustained drug selection," *Nature genetics*, vol. 44, no. 1, pp. 101–105, 2012.

[9] E. A. Yurtsev, H. X. Chao, M. S. Datta, T. Artemova, and J. Gore, "Bacterial cheating drives the population dynamics of cooperative antibiotic resistance plasmids," *Molecular systems biology*, vol. 9, no. 1, p. 683, 2013.

[10] R. Chait, A. Craney, and R. Kishony, "Antibiotic interactions that select against resistance," *Nature*, vol. 446, no. 7136, p. 668, 2007.

[11] J. P. Torella, R. Chait, and R. Kishony, "Optimal drug synergy in antimicrobial treatments," *PLoS computational biology*, vol. 6, no. 6, p. e1000796, 2010.

[12] E. Hansen, R. J. Woods, and A. F. Read, "How to use a chemotherapeutic agent when resistance to it threatens the patient," *PLoS biology*, vol. 15, no. 2, p. e2001110, 2017.

[13] A. Fischer, I. Vázquez-García, and V. Mustonen, "The value of monitoring to control evolving populations," *Proceedings of the National Academy of Sciences*, vol. 112, no. 4, pp. 1007–1012, 2015.

[14] H. Chung, T. D. Lieberman, S. O. Vargas, K. B. Flett, A. J. McAdam, G. P. Priebe, and R. Kishony, "Global and local selection acting on the pathogen stenotrophomonas maltophilia in the human lung," *Nature communications*, vol. 8, p. 14078, 2017.

[15] D. Nichol, P. Jeavons, A. G. Fletcher, R. A. Bonomo, P. K. Maini, J. L. Paul, R. A. Gatenby, A. R. Anderson, and J. G. Scott, "Steering evolution with sequential therapy to prevent the emergence of bacterial antibiotic resistance," *PLoS computational biology*, vol. 11, no. 9, p. e1004493, 2015.

[16] M. Gralka, D. Fusco, S. Martis, and O. Hallatschek, "Convection shapes the trade-off between antibiotic efficacy and the selection for resistance in spatial gradients," *Physical Biology*, vol. 14, no. 4, p. 045011, 2017.

[17] F. Fu, M. A. Nowak, and S. Bonhoeffer, "Spatial heterogeneity in drug concentrations can facilitate the emergence of resistance to cancer therapy," *PLoS Comput Biol*, vol. 11, no. 3, p. e1004142, 2015.

[18] P. Greulich, B. Waclaw, and R. J. Allen, "Mutational pathway determines whether drug gradients accelerate evolution of drug-resistant cells," *Physical Review Letters*, vol. 109, no. 8, p. 088101, 2012.

[19] R. Hermsen, J. B. Deris, and T. Hwa, "On the rapidity of antibiotic resistance evolution facilitated by a concentration gradient," *Proceedings of the National Academy of Sciences*, vol. 109, no. 27, pp. 10775–10780, 2012.

[20] T. B. Kepler and A. S. Perelson, "Drug concentration heterogeneity facilitates the evolution of drug resistance," *Proceedings of the National Academy of Sciences*, vol. 95, no. 20, pp. 11514–11519, 1998.

[21] S. Moreno-Gamez, A. L. Hill, D. I. Rosenbloom, D. A. Petrov, M. A. Nowak, and P. S. Pennings, "Imperfect drug penetration leads to spatial monotherapy and rapid evolution of multidrug resistance," *Proceedings of the National Academy of Sciences*, vol. 112, no. 22, pp. E2874–E2883, 2015.

[22] Q. Zhang, G. Lambert, D. Liao, H. Kim, K. Robin, C.-k. Tung, N. Pourmand, and R. H. Austin, "Acceleration of emergence of bacterial antibiotic resistance in connected microenvironments," *Science*, vol. 333, no. 6050, pp. 1764–1767, 2011.

[23] R. Hermsen and T. Hwa, "Sources and sinks: a stochastic model of evolution in heterogeneous environments," *Physical review letters*, vol. 105, no. 24, p. 248104, 2010.

[24] M. Baym, T. D. Lieberman, E. D. Kelsic, R. Chait, R. Gross, I. Yelin, and R. Kishony, "Spatiotemporal microbial evolution on antibiotic landscapes," *Science*, vol. 353, no. 6304, pp. 1147–1151, 2016.

[25] C. S. Gokhale, Y. Iwasa, M. A. Nowak, and A. Traulsen, "The pace of evolution across fitness valleys," *Journal of Theoretical Biology*, vol. 259, no. 3, pp. 613–620, 2009.

[26] Y. Iwasa, F. Michor, and M. A. Nowak, "Stochastic tunnels in evolutionary dynamics," *Genetics*, vol. 166, no. 3, pp. 1571–1579, 2004.

[27] D. M. Weinreich, L. Chao, and P. Phillips, "Rapid evolutionary escape by large populations from local fitness peaks is likely in nature," *Evolution*, vol. 59, no. 6, pp. 1175–1182, 2005.

[28] D. B. Weissman, M. M. Desai, D. S. Fisher, and M. W.

Feldman, "The rate at which asexual populations cross fitness valleys," *Theoretical population biology*, vol. 75, no. 4, pp. 286–300, 2009.

[29] S. T. Pickett and M. L. Cadenasso, "Landscape ecology: spatial heterogeneity in ecological systems," *Science*, vol. 269, no. 5222, p. 331, 1995.

[30] M. C. Whitlock and R. Gomulkiewicz, "Probability of fixation in a heterogeneous environment," *Genetics*, vol. 171, no. 3, pp. 1407–1417, 2005.

[31] L. A. Real and R. Biek, "Spatial dynamics and genetics of infectious diseases on heterogeneous landscapes," *Journal of the Royal Society Interface*, vol. 4, no. 16, pp. 935–948, 2007.

[32] S. Farhang-Sardroodi, A. Darooneh, M. Nikbakht, N. Komarova, and M. Kohandel, "The effect of spatial randomness on the average fixation time of mutants.," *PLoS computational biology*, vol. 13, no. 11, p. e1005864, 2017.

[33] A. Kaznatcheev, J. G. Scott, and D. Basanta, "Edge effects in game-theoretic dynamics of spatially structured tumours," *Journal of The Royal Society Interface*, vol. 12, no. 108, p. 20150154, 2015.

[34] B. Allen, G. Lippner, Y.-T. Chen, B. Fotouhi, N. Momeni, S.-T. Yau, and M. A. Nowak, "Evolutionary dynamics on any population structure," *Nature*, vol. 544, no. 7649, pp. 227–230, 2017.

[35] K. S. Korolev, M. Avlund, O. Hallatschek, and D. R. Nelson, "Genetic demixing and evolution in linear stepping stone models," *Reviews of modern physics*, vol. 82, no. 2, p. 1691, 2010.

[36] M. Khasin, E. Khain, and L. M. Sander, "Fast migration and emergent population dynamics," *Physical review letters*, vol. 109, no. 24, p. 248102, 2012.

[37] M. Khasin, B. Meerson, E. Khain, and L. M. Sander, "Minimizing the population extinction risk by migration," *Physical review letters*, vol. 109, no. 13, p. 138104, 2012.

[38] J. N. Waddell, L. M. Sander, and C. R. Doering, "Demographic stochasticity versus spatial variation in the competition between fast and slow dispersers," *Theoretical population biology*, vol. 77, no. 4, pp. 279–286, 2010.

[39] G. W. Constable and A. J. McKane, "Fast-mode elimination in stochastic metapopulation models," *Physical Review E*, vol. 89, no. 3, p. 032141, 2014.

[40] G. W. Constable and A. J. McKane, "Population genetics on islands connected by an arbitrary network: An analytic approach," *Journal of theoretical biology*, vol. 358, pp. 149–165, 2014.

[41] G. W. Constable and A. J. McKane, "Stationary solutions for metapopulation moran models with mutation and selection," *Physical Review E*, vol. 91, no. 3, p. 032711, 2015.

[42] C. Li, "The statistical processes of evolutionary theory," *American journal of human genetics*, vol. 14, no. 4, p. 438, 1962.

[43] C. W. Gardiner, *Handbook of stochastic methods for physics, chemistry and the natural sciences*, vol. 13 of *Springer Series in Synergetics*. Berlin: Springer-Verlag, third ed., 2004.

[44] N. V. Kampen, *Stochastic processes in physics and chemistry*. North Holland, 2007.

[45] D. B. Clewell, M. S. Gilmore, Y. Ike, and N. Shankar, *Enterococci: from commensals to leading causes of drug resistant infection*. Massachusetts Eye and Ear Infirmary, 2014.