TITLE:
Phenotypic variation in mitochondrial function across New Zealand snail populations

AUTHORS:
Emma S. Greimann[1], Samuel F. Ward[2], James D. Woodell[1], Samantha Hennessey[1], Michael R. Kline[1], Jorge A. Moreno[1], Madeline Peters[1], Jennifer L. Cruise[3], Kristi L. Montooth[4], Maurine Neiman[1], Joel Sharbrough*[1,5]

[1] University of Iowa, Iowa City, IA, USA
[2] Purdue University, West Lafayette, IN, USA
[3] University of St. Thomas, Saint Paul, MN, USA
[4] University of Nebraska, Lincoln, Lincoln, NE
[5] Current address: Colorado State University, Fort Collins, CO, USA

*CORRESPONDENCE TO:
Joel Sharbrough
Colorado State University
440 Biology Building
Fort Collins, CO 80523
jsharbro@rams.colostate.edu
(970) 495-2301

RUNNING HEADLINE:
Mitochondrial function in New Zealand snails

ABSTRACT

Mitochondrial function is critical for energy homeostasis and should shape how genetic variation

in metabolism is transmitted through levels of biological organization to generate stability in

organismal performance. Mitochondrial function is encoded by genes in two distinct and

separately inherited genomes - the mitochondrial genome and the nuclear genome - and selection

is expected to maintain functional mito-nuclear interactions. Nevertheless, high levels of

polymorphism in genes involved in these mito-nuclear interactions and variation for

mitochondrial function are nevertheless frequently observed, demanding an explanation for how

and why variability in such a fundamental trait is maintained. *Potamopyrgus antipodarum* is a

New Zealand freshwater snail with coexisting sexual and asexual individuals and, accordingly,

contrasting systems of separate vs. co-inheritance of nuclear and mitochondrial genomes. As

such, this snail provides a powerful means to dissect the evolutionary and functional

consequences of mito-nuclear variation. The lakes inhabited by *P. antipodarum* span wide

environmental gradients, with substantial across-lake genetic structure and mito-nuclear

discordance. This situation allows us to use comparisons across reproductive modes and lakes to

partition variation in cellular respiration across genetic and environmental axes. Here, we

integrated cellular, physiological, and behavioral approaches to quantify variation in

mitochondrial function across a diverse set of wild *P. antipodarum* lineages. We found

extensive across-lake variation in organismal oxygen consumption, mitochondrial membrane

potential, and behavioral response to heat stress, but few global effects of reproductive mode or

sex. Taken together, our data set the stage for applying this important model system for sexual

reproduction and polyploidy to dissecting the complex relationships between mito-nuclear

variation, performance, plasticity, and fitness in natural populations.

KEYWORDS

asexual reproduction, JC-1, mitochondria, oxygen consumption, *Potamopyrgus antipodarum*,

sexual reproduction

INTRODUCTION

The production of ATP *via* cellular respiration is a critical component of eukaryotic function and fitness[1-5], and the components of the oxidative phosphorylation (OXPHOS) pathway generally appear to be evolving under purifying selection[6-11]. Nevertheless OXPHOS genes are often polymorphic within species[3,12-15], with important implications for phenomena ranging from mito-nuclear incompatibilities[16] to mitochondrial introgression[17] to DNA barcoding[18]. How mitochondrial haplotype is connected to phenotype and how mitochondrial function is maintained over evolutionary time despite mutational pressure[19] remain unclear, despite extensive variation for metabolic and mitochondrial function in natural populations of a diverse array of species[3,20-27]. Although some variation in metabolic and mitochondrial traits has been linked to specific environmental correlates (*e.g.,* altitude[28,29], temperature[30], energy source[31]), we lack a systematic understanding of the genotype-phenotype-fitness map for traits that depend on mitochondrial function and how this map may vary across environmental gradients.

Teasing apart the sources of variation in mitochondrial function is further complicated by the genetic architecture underlying ATP production. The enzymes that carry out OXPHOS (excepting succinate dehydrogenase in animals, fungi, and most plants[32,33]) are encoded by genes that are partitioned across two separate cellular compartments (i.e., the nucleus and mitochondria) with different inheritance regimes (i.e., biparental vs. uniparental, respectively)[34]. The biology of these coevolving "mito-nuclear" interactions has been the subject of intense study[16,35-41], but the consequences of coevolution between genomes for function and fitness have only been evaluated in a handful of species and for a handful of mito-nuclear genotypes[16,42-45]. This research has nevertheless yielded critical insights into how mito-nuclear genotypic variation produces phenotypic variation, and demonstrating that mito-nuclear epistasis can be a powerful

force for shaping genetic variation in nature[46]. The separate inheritance of nuclear and

mitochondrial variation during sexual reproduction presents a challenge for natural selection in

shaping co-adapted mito-nuclear genotypes[47]. In particular, because nuclear genomes are

biparentally inherited and mitochondrial genomes are uniparentally inherited, mitochondrial

genomes will thus be swapped across diverse nuclear genomic backgrounds. This separate

inheritance of nuclear and mitochondrial genomes can result in the loss of particular high-fitness

mito-nuclear genotypes from populations, as selection favors the fixation of mitochondrial

genomes that have the highest fitness on average across the suite of nuclear variants in the

population[48]. Sex-linkage and sex-specific fitness effects comprise a particular set of conditions

that may enable the maintenance of mito-nuclear variation in populations[49]. The overwhelming

preponderance of sexual reproduction in nature would thus seem to imply that the reliable

transmission of particular mito-nuclear combinations is not all that important. What is perhaps

more likely is that we do not have a good understanding of how sexual reproduction affects the

connection between mito-nuclear genotype and mitochondrial performance.

Asexual reproduction, in which mitochondrial and nuclear genomes are co-inherited from

mother to daughter, offers a useful contrast for understanding the consequences of sexual

reproduction for mitochondrial function. In particular, mitochondrial genomes that are "trapped"

in lineages that have transitioned to asexuality are inextricably linked to their nuclear genomic

background for the remainder of their existence. Because asexual lineages are expected to

accumulate deleterious mutations in both the nuclear and mitochondrial genomes[50] as a result of

Muller's Ratchet[51] and the Hill-Robertson Effect[52], asexual lineages should exhibit reduced

mitochondrial performance relative to sexual lineages. However, co-inheritance of nuclear and

mitochondrial genomes also allows epistatic variation to become visible to natural selection,

such that asexual lineages may experience both relatively intense and efficient selection favoring

compensatory co-evolution between nuclear and mitochondrial loci. The extent to which asexual

lineages exhibit reduced vs. improved mitochondrial performance compared to close sexual

relatives will provide a useful guide for whether and how epistatic mito-nuclear variation

contributes to mitochondrial performance.

Potamopyrgus antipodarum is a New Zealand freshwater snail[53] featuring frequent co-

existence and competition among obligately sexual and obligately asexual lineages[54]. So similar

are these sexual and asexual lineages that the only reliably distinguishable character apart from

reproductive mode is that asexual lineages are polyploid and sexual lineages are diploid[55].

Because asexuality has arisen on multiple separate occasions within *P. antipodarum*[56,57], distinct

asexual lineages can be treated as repeated "natural experiments" into the consequences of

sexual reproduction for mitochondrial performance[8,58]. *Potamopyrgus antipodarum*

populations also feature extensive population structure in mitochondrial DNA sequences[8,56,59],

and asexual lineages harbor heritable variation for mitochondrial performance[27]. Notably, the

occasional production of male offspring by obligately asexual female *P. antipodarum*[60] makes it

possible to evaluate the phenotypic consequences of mitochondrial mutations that are neutral or

beneficial when present in females but deleterious when present in males. In particular, selection

against male-harming mutations is expected to be relaxed in asexual lineages, whose

mitochondrial genomes have been "trapped" in females for generations, such that asexual males

are predicted to exhibit especially poor mitochondrial performance. Together, these features

make *P. antipodarum* uniquely well suited for decoupling sexual reproduction from

mitochondrial genome inheritance, a necessary step if we hope to understand how mito-nuclear

epistasis contributes to organismal function and fitness in natural populations.

The observation that mitochondrial genomes of asexual *P. antipodarum* lineages

accumulate putatively harmful mutations more readily than do their sexual counterparts[8,58],

combined with the heritable variation for mitochondrial function present in asexual lineages[27]

leads to the expectation that asexual *P. antipodarum* will exhibit reduced mitochondrial

performance compared to sexual lineages. Here we tested that prediction as well as whether sex

(*i.e.*, male vs. female) and source population affect mitochondrial and physiological function at

organellar and organismal levels under laboratory conditions using wild-caught *P. antipodarum*.

MATERIALS AND METHODS

Field collections of *P. antipodarum*

The phenotypic and ecological similarity of sexual *vs.* asexual *P. antipodarum* enables direct

comparisons across reproductive modes, but also means that definitive determination of

reproductive mode (via flow cytometric genome size estimation – see below) requires snail

sacrifice. We therefore sampled field-collected snails from New Zealand lakes known to harbor

both sexual and asexual individuals[57,59,61]. Snails were collected from New Zealand lakes in

January 2015 and December 2016 and transported by hand in damp paper towels to the

University of Iowa. Upon arrival, snails were housed at 16ºC with a 18L:6D photoperiod and fed

Spirulina algae three times per week, as previously described[62]. We arbitrarily selected adult

snails from lake collections (each of which consisted of hundreds to thousands of individuals)

and isolated each snail in a 0.5 L glass container with 300ml carbon-filtered H_2O. Water was

changed weekly. All functional assays began immediately following isolation and were

completed within six months of arrival at the University of Iowa.

To maximize our ability to sample and compare sexual and asexual individuals from the

same lake, we assayed oxygen consumption under heat stress (details below) in 158 wild-caught

females from six lakes (Figure 1, Table 1) and used destructive sampling to determine

reproductive mode in snails that survived all three temperature trials. The recent discovery of

asexually produced males in *P. antipodarum*[60] and recognition of their unique potential to

provide insight into the capacity for mtDNA to harbor mutations that are deleterious in males but

neutral or beneficial in females motivated us to include both males and females in our next

experiment. Here, we assayed behavioral function in response to heat stress and mitochondrial

membrane potential in 46 wild-caught snails (Figure 1, Table 1).

Oxygen consumption under heat stress conditions

Because oxygen becomes limiting to ectotherms under elevated temperatures[63] and because *P.*

antipodarum demonstrates signs of stress at elevated (~30ºC) temperatures (*e.g.,* reduced

fecundity[64], elevated oxygen consumption[27], and decreased righting ability[27]), we measured

oxygen consumption using an aquatic respirometer as previously described[27] for 158 wild-caught

female *P. antipodarum* from each of six lakes. Oxygen consumption was assayed at three

different water temperatures: 16ºC (not stressful, and similar to New Zealand lake temperatures),

22ºC (moderately stressful), and 30ºC (stressful)[27]. Each snail was assayed at each temperature in

a randomly determined order, and only the 61 snails (38.6%) that survived across all three

temperature treatments were included in our analyses. We were unable to determine a single

source of mortality for those snails that did not survive all three temperature trials, although

exposure to high temperatures combined with hypoxic or near-hypoxic conditions inside the

respiration chamber may certainly have played a role. *Potamopyrgus antipodarum* are also

notoriously difficult to keep alive in laboratory conditions when isolated from other snails

(personal observation), and our temperature treatment procedure combined with recovery periods

required that snails spent many weeks in isolation. Wet mass for each individual was calculated

after each trial, and mean mass across all three trials was used in our final analyses.

Behavioral response to heat stress

Righting time and time to shell emergence following a startling stimulus increase with

temperature in *P. antipodarum*[27]*,* indicating that both assays can be used to assess heat-induced

stress. We quantified righting times and time to emergence under each of the same three

temperature treatments used for oxygen consumption in 46 wild-caught *P. antipodarum* and

compared behavior reaction norms across temperatures, lakes, reproductive modes, and sexes.

Snails were each assayed once at each temperature, and no snails died in the course of behavioral

analyses, which require ~50% less time spent (<1.05 hr) under the elevated temperature

treatment than oxygen consumption assays (2 hr) and no time spent under hypoxic or near-

hypoxic conditions.

Mitochondrial membrane potential

JC-1 is a small positively charged molecule that diffuses down the electrochemical gradient

across the inner mitochondrial membrane[65]. Under UV illumination, JC-1 fluoresces green if

dispersed and red if aggregated (e.g., when inside the mitochondrial matrix)[65]. As a result, the

median ratio of red: green fluorescence in freshly isolated live mitochondria stained with JC-1

can serve as a proxy for mitochondrial membrane potential, with a high ratio indicating high

membrane potential. We isolated mitochondria from 46 wild-caught *P. antipodarum* and

measured red: green ratios in JC-1-treated mitochondrial extracts as previously described[27] using

a Becton Dickenson LSR II flow cytometer. Using the set of 46 snails that completed behavioral

assays at all three temperatures, we compared median red: green JC-1 ratios across lakes,

reproductive modes, and sexes.

Determination of reproductive mode

Sexual *P. antipodarum* are diploid and asexual *P. antipodarum* are polyploid (triploid or

tetraploid[57,66]), allowing us to use flow cytometry to determine DNA content and infer

reproductive mode of individual snails. After completing all physiological and/or behavioral

assays blind with respect to snail reproductive mode, we dissected head tissue from each

individual snail, flash froze this tissue in liquid nitrogen, and stored the frozen tissue at -80ºC

until flow cytometry. We then homogenized head tissue in DAPI solution, passed this

homogenate through a 30 µm filter, and analyzed the suspended nuclei on a Becton-Dickinson

FACS Aria II following the protocol (including a chicken red blood cell standard) outlined in

previous work[67].

Statistical analyses

Using a mixed-effects model, we tested whether mass, temperature (16ºC, 22ºC, 30ºC), lake-of-

origin (n = 6), reproductive mode (sexual or asexual), and interactions of mass × temperature,

mass × lake, mass × reproductive mode, temperature × lake-of-origin, temperature ×

reproductive mode, mass × temperature × lake, and mass × temperature × reproductive mode

were good predictors of oxygen consumption per hour. A term for snail identity was fit as a

random intercept to account for repeated measures on individuals across temperatures. We

employed a similar framework for emergence time following a startling stimulus and righting

behavior after being flipped upside-down, except we also included terms for sex (male or

female) and a temperature × sex interaction, as we had both male and female snails in the sample

used in these analyses. Finally, we modeled mitochondrial membrane potential, measured as the

ratio of red: green fluorescence, as a function of lake, reproductive mode, and sex, using analysis

of variance (ANOVA) and a Levene's test to check for equal variances ($F_{5,40} = 2.044$, $p = 0.09$).

Because we lacked replication of reproductive mode and sex within all sampled lakes, and after

discovering a large lake effect in several phenotypic measures, we also fit sub-models predicting

each phenotypic measure within lakes. Sub-models were equivalent to those described above for

phenotypic measures but only included data from lakes that had replication for reproductive

mode, sex, or both.

We developed final models using backwards selection until only predictors with *p*-values

less than 0.05 remained. All main effects were kept in the model when part of a significant

interaction. To test assumptions of normality and heteroscedasticity of errors, we graphically

inspected residuals and log- or power-transformed response variables when necessary (Figure

S3). Power transformation parameters were estimated using the geoR package[68]. We performed

all statistical analyses in R[69], fitting fixed-effect models with the *lm* function, fitting mixed-

effects models using the lme4 package[70], estimating degrees of freedom for mixed-effect models

using Satterthwaite's approximation *via* the lmerTest package[71], and performing pairwise

comparisons of mixed model predictors using the emmeans package[72], which adjusts *p* values for

multiple comparison using the Tukey method. To test whether behavioral phenotypes were

correlated with mitochondrial membrane potential, we estimated pairwise Pearson correlation

coefficients for behavioral responses at each temperature and membrane potentials using the

Hmisc package[73] and corrected for multiple comparisons using the Holm procedure[74].

RESULTS

Prevalence of reproductive modes and sexes in two samples of wild-caught *P. antipodarum*

Our first sample was comprised of 158 female *P. antipodarum* collected from six New Zealand

lakes (Lakes Alexandrina, Clearwater, Kaniere, Paringa, Rotoroa, and Selfe [Figure 1]), among

which we were able to assay oxygen consumption at 16ºC, 22ºC, and 30ºC for 61 snails. We

determined ploidy (and therefore reproductive mode) for 57 out of these 61 snails. The sample

included 32 sexual snails, 25 asexual snails, and four snails of unknown reproductive mode

(Table 1). All 32 sexuals were diploid and all 25 asexual individuals were triploid. Only one of

our sampled lakes included multiple sexual and asexual snails (Lake Alexandrina), while the

other five were monomorphic for reproductive mode (Table 1).

Our second sample was comprised of 22 male and 24 female *P. antipodarum* collected

from six New Zealand freshwater lakes (Lakes Alexandrina, Ellery, Kaniere, Mapourika,

Rotoroa, and Selfe – Figure 1), among which we assayed emergence time and righting behavior

at 16ºC, 22ºC, and 30ºC, quantified mitochondrial membrane potential, and determined ploidy

for all 46 snails. This sample included 31 sexuals and 15 asexuals, and we sampled multiple

sexual and asexual individuals from within three different lakes (Lakes Ellery, Mapourika, and

Selfe). Lakes Mapourika and Selfe also had replication for sex, with both male and female

sexuals sampled from both lakes. However, we were only able to sample male and female snails

exhibiting both sexual and asexual reproduction in Lake Selfe (Table 1).

Oxygen consumption under stressful conditions

To partition variation in oxygen consumption across biological and environmental axes, we measured snail oxygen consumption in a closed-system aquatic respirometer under three different temperature treatments. Individual snail reaction norms from each of six lakes for oxygen consumption across temperature treatments are depicted in Figure S1a. We fit a mixed-effects model to our response variable of oxygen consumption per hour. After eliminating non-significant predictors using backwards selection, we found significant effects of snail wet mass ($\chi^2 = 10.82$, df = 1, $p = 0.0010$), temperature treatment ($\chi^2 = 45.91$, df = 2, $p < 0.0001$), and lake-of-origin ($\chi^2 = 12.74$, df = 5, $p = 0.0260$) (Table 2). Snail wet mass had a positive effect on oxygen consumption per hour (correlation slope estimate = 5,859.06), indicating that larger snails consume more oxygen per hour than do smaller snails (Figure S2). *Post hoc* pairwise t-tests revealed that snails consumed significantly more oxygen at 22ºC than at 16ºC (*t* ratio = -4.12, df = 118, $p = 0.0002$), at 30ºC than at 16ºC (*t* ratio = -6.72, df = 120, $p < 0.0001$), and at 30ºC than at 22ºC (*t* ratio = -2.663, df = 118, $p = 0.0238$) (Figure 2a). Similarly, *post hoc* pairwise t-tests of reaction norms across lakes revealed two statistical groups of lakes, with Lake Clearwater exhibiting especially low rates of oxygen consumption per hour. We did not find any difference in oxygen consumption per hour across reproductive modes (Figure 3a), even when we restricted our analyses to snails within Lake Alexandrina (Table S1, model e), the one lake where we had the snails required to perform such comparisons. Overall, these data indicate that lake-of-origin plays a major role in how snails respire in response to heat stress, and that asexual snails do not exhibit altered oxygen consumption compared to sexual snails overall. Still, low levels of within-lake sampling precluded a rigorous test of the effects of reproductive mode on oxygen consumption in response to heat stress.

Behavioral responses to heat stress

Using our second sample of 46 male and female *P. antipodarum* from six New Zealand lakes, we

assayed two behaviors that challenge snails, especially under heat stress conditions: emergence

time following a startling stimulus and righting ability after being flipped upside down.

Individual snail reaction norms from each of six lakes for emergence time and righting time

across temperature treatments are depicted in Figure S1b and c, respectively. We found that both

temperature ($\chi^2 = 48.53$, df = 2, $p < 0.0001$) and lake-of-origin ($\chi^2 = 11.34$, df = 5, $p = 0.0450$)

were significant predictors of emergence time (Table 2). In particular, *post hoc* t-tests of

temperature effects averaged across lakes revealed that snails emerged from their shells

significantly faster at 30ºC than at either 16ºC (*t* ratio = 6.42, df = 90, $p < 0.0001$) or 22ºC (*t* ratio

= 5.55, df = 90, $p < 0.0001$), but that there was no significant difference in emergence time at

16ºC vs. 22ºC (*t* ratio = 0.87, df = 90, $p = 0.66$) (Figure 2b). Despite the overall effect of lake-of-

origin on emergence time, *post hoc* pairwise comparisons of lakes averaged over temperature did

not reveal any obvious statistical groupings ($p > 0.05$ for all pairwise comparisons). We did not

detect overall effects of reproductive mode (Figure 3b) or sex (Figure 4a) on emergence time,

even when we limited our analyses to Lake Selfe (the only lake with replication for both sex and

reproductive mode) (Table S1, model f). Temperature was also a significant predictor of righting

behavior ($\chi^2 = 73.36$, df = 2, $p < 0.0001$), and *post hoc* pairwise analyses revealed that snails take

significantly longer to right themselves 30ºC than at 16ºC (*t* ratio = 6.42, df = 90, $p < 0.0001$)

and at 22ºC (*t* ratio = 6.42, df = 90, $p < 0.0001$), but not at 22ºC vs. 16ºC (*t* ratio = 6.42, df = 90,

$p < 0.0001$) (Table 2). In contrast to emergence time, reproductive mode ($\chi^2 = 4.48$, df = 1, $p =$

0.0343) and the reproductive mode × sex interaction ($\chi^2 = 12.67$, df = 2, $p = 0.0107$) were both

significant predictors of righting time (Table 2), with asexuals righting themselves faster at 16ºC

and 22ºC, but taking longer to right themselves at 30ºC than sexuals (Figure 3c). Lake-of-origin

was not a significant predictor of righting time overall (Figure 2c), but there were significant

differences among lakes (χ^2 = 12.67, df = 2, p = 0.0018) as well as a significant temperature ×

lake-of-origin interaction (χ^2 = 12.99, df = 4, p = 0.0113) when we limited our analyses to

females (Table S1, model s). We also tested whether behavioral responses were correlated with

one another and found that significant negative correlations exist between emergence time and

righting time at 16ºC (r^2 = 0.56, p < 0.0001), 22ºC (r^2 = 0.34, p < 0.0001), 30ºC (r^2 = 0.19, p <

0.0023), and between emergence time at 22ºC and righting time at 16ºC (r^2 = 0.049, p < 0.0044).

Together these data support a major role for lake-of-origin in determining behavioral response to

heat stress, and we see some evidence from this relatively small set of snails that reproductive

mode and sex have effects on righting behavior but not emergence.

Mitochondrial membrane potential

We next assayed mitochondrial membrane potential in freshly isolated mitochondrial fractions,

using the fluorescent dye JC-1, in this same set of 46 field-collected snails that were used for

behavioral trials. We found that sex was a significant predictor of mitochondrial membrane

potential (ANOVA $F_{1,44}$ = 6.84, p = 0.0122, Figure 4c), with males having higher mitochondrial

membrane potential than females. This difference between males and females was evident even

when we limited our analysis to asexuals (ANOVA $F_{1,6}$ = 12.09, p = 0.0132, Table S1, model k)

or to Lake Selfe, which featured full replication of reproductive mode and sex (ANOVA $F_{1,15}$ =

6.58, p = 0.0216, model h). While sexual males have higher mitochondrial membrane potentials

than sexual females, the difference is not statistically significant (Table S1, model u), indicating

that the higher membrane potential of asexual males are largely responsible for the effect of sex

in the overall dataset. Lake-of-origin (Figure 2d) and reproductive mode did not affect

mitochondrial membrane potential (Figure 3d), even when we limited our analyses to Lake Selfe

or to females within Lakes Mapourika and Selfe (Table S1). We did not find evidence of

correlations between mitochondrial membrane potential and either of the behavioral responses

from the same snails. In sum, we found a strong effect of sex, especially in the context of

asexuality, and no evidence of an effect of reproductive mode on mitochondrial membrane

potential, with the caveat that low-within lake sampling may prevent us from observing

relatively small effects.

DISCUSSION

Extensive work in the *P. antipodarum* system has documented lake-specific phenotypes and

local adaptation for resistance to infection by the trematode parasite *Atriophallophorus*

winterbourni [61,75-81], life history traits such as growth rate and size[82], and response to nutrient

limitation[67,83]. Here, we report the first evidence of lake-structured variation for metabolic and

behavioral function and sex-specific mitochondrial function in *P. antipodarum*. Combined with

marked population structure for mitochondrial genetic variation[8,56], this result suggests the

intriguing possibility that mitochondrial function is locally tuned in *P. antipodarum*.

While the elevated linkage disequilibrium expected to result from asexuality should

reduce the efficacy of natural selection in both nuclear[52,84-88] and mitochondrial genomes[19,50,89],

stable transmission of mito-nuclear genotypes may also facilitate rapid mito-nuclear coadaptation

and thereby local adaptation[47]. Surveys of mitochondrial genomes of asexual lineages[90-92],

including in *P. antipodarum*[8,56], have revealed elevated rates of putatively harmful mutations in

mitochondrial genomes compared to sexual lineages. Absent nuclear compensation for

mitochondrial function[37,93], we predict mitochondrial genomes carried by asexual lineages will

therefore be associated with reduced mitochondrial performance. However, we could not detect

any global decrease in mitochondrial, organismal, or behavioral performance in asexual vs.

sexual lineages in this geographically diverse sample. We must therefore conclude that either the

decline in mitochondrial performance in asexual lineages is only detectable among closely

related sexual and asexual pairs within the same lakes or that asexual lineages are compensating

for the increased mutation load in their mitochondrial genomes in some way. This compensation

could be through mechanisms of physiological homeostasis (see e.g.[94]) or through clonal

selection favoring lineages with mito-nuclear genotypes in which mutations in the nuclear

genome compensate for the deleterious mutations in the mitochondrial genome. One potentially

important source for genetic and/or physiological compensation might be polyploidy, as sexual

P. antipodarum are all diploid and asexual *P. antipodarum* are all polyploid[55]. Extra genome

copies in the nuclear genome may allow for mutational masking that facilitates broader traversal

of the adaptive landscape[95]. Physiologically, elevated genome size is positively correlated with

larger cell size[96], which likely alters the signaling, energetic, and stoichiometric landscape of the

cell[95]. Moreover, the expected larger cell sizes of polyploids vs. diploids might allow for more

and larger mitochondria, which in turn give rise to opportunities for compensatory regulation at

increasing levels of organization including the number of organelle genome copies per organelle,

the number of transcripts/proteins produced from each organelle genome copy, and even the

number of OXPHOS complexes formed per unit area of the mitochondrial inner membrane[97].

Across evolutionary time, if mitochondrial performance is indeed a critical component of fitness

in *P. antipodarum*, as it is for other animal taxa[1-5], asexual lineages with decreased mitochondrial

performance might be evolutionarily short lived and difficult to sample. Combined with the

strong lake effect observed here, it is clear that only extensive within-lake sampling combined

with integrative approaches that span levels of biological organization will be capable of

elucidating mechanisms of mitochondrial compensation in asexual lineages of *P. antipodarum.*

The nature of our sampling design means that the lake effects observed here could be the

result of either genetic or environmental forces. In particular, these effects are largely consistent

with our current understanding of *P. antipodarum* population genetic structure[56,59,61], in which

lakes and other geographic features of New Zealand (e.g., the Southern Alps[56]) act as major

barriers to gene flow among *P. antipodarum* subpopulations. Previous work investigating

mitochondrial function in lab-cultured iso-female asexual lineages of *P. antipodarum* (i.e., snails

that are asexual descendants of a single mother collected from the wild and isolated in the lab)

has identified heritable variation for mitochondrial, metabolic, and behavioral phenotypes[27].

However, because the snails used in this study were born and lived to reproductive maturity in

the wild and subsequently acclimated to laboratory conditions, it is possible that the lake effects

observed here might also be the result of phenotypic plasticity for metabolic rate. One tractable

angle for investigating the contributions of genetic vs. environmental variation in mito-nuclear

genotype to variation in mitochondrial performance can take advantage of extensive mito-

nuclear discordance observed in asexual lineages of *P. antipodarum*[59]. Because asexual lineages

stably transmit mito-nuclear genotypes across generations, asexual lineages that have trapped

reciprocal combinations of mitochondrial haplotypes with different nuclear genotypes can be

used to test whether mito-nuclear epistatic variation contributes to mitochondrial function and

ultimately, organismal fitness.

Maternal transmission of mitochondrial genomes precludes inheritance of all

mitochondrial mutations originating in males, with two primary consequences for the efficacy of

selection on the mitochondrial genome in sexual taxa. First, genes in the mitochondrial genome

of sexually reproducing lineages experience a reduction in the effective population size relative

to nuclear genes due to both the absence of recombination and uniparental inheritance. Second,

mutations in the mitochondrial genome that have sex-specific effects only experience effective

natural selection in females[98,99]. This latter phenomenon is predicted to result in the accumulation

of mutations that are neutral or beneficial in females but deleterious in males[100]. The lack of

widespread evidence for mitochondrial mutations with sex-specific fitness effects (but

see[35,101,102]) may point to mechanisms that prevent the spread of male-specific deleterious

mutations in mitochondrial genomes (*e.g.,* paternal leakage[103], inbreeding[104], kin selection[104],

and/or nuclear-encoded restorers of male function[42,105]). Asexually produced males, as are

occasionally produced in *P. antipodarum*[60], represent worst-case scenarios for male-specific

mitochondrial performance because their nuclear and mitochondrial genomes have been

"trapped" in females for generations but are now being expressed in a male context. Because

selection against male-harming mutations is therefore completely ineffective in asexual lineages,

asexual males are expected to exhibit particularly poor mitochondrial performance. We found

that male *P. antipodarum* have higher mitochondrial membrane potential than female

counterparts, leading us to speculate that male *P. antipodarum* do not suffer from sex-specific

mutations that decrease their ability to generate a proton motive force. Mitochondrial membrane

potential depends upon protons pumped by the OXPHOS complexes that are comprised of both

nuclear and mitochondrial gene products (*i.e.,* complexes I, III, and IV[32,33]). The implications are

that we should expect to detect reduced mitochondrial membrane potentials if male-harming

mutations have accumulated in *P. antipodarum* mitochondrial genomes. One of the most

intriguing findings presented here is that asexual males appear to be driving this pattern of

elevated mitochondrial membrane potential to a greater extent than sexual males. However,

because we do not know the relationship between mitochondrial membrane potential and fitness,

we cannot yet determine whether this sexual dimorphism in mitochondrial membrane potential

supports or refutes the presence of male-harming mitochondrial mutations in *P. antipodarum*.

Still, the marked increase in mitochondrial membrane potential among asexually produce males

offers a promising avenue for investigating the prevalence of evolutionary dynamics of male-

harming mutations.

CONCLUSIONS

We found strong effects of temperature on oxygen consumption, righting behavior, and

emergence time, lake effects on three out of four mitochondria-related phenotypes, a difference

among sexual and asexual individuals in one behavioral assay, and a sex-specific difference in

mitochondrial membrane potential. These results indicate that local conditions may play a major

role in modulating mitochondrial function in *P. antipodarum*. The strong lake effects observed

demand more extensive intra-lake sampling, in order to better test for the effects of reproductive

mode and sex on mitochondrial and organismal performance in *P. antipodarum*. Nevertheless,

with the current sampling, we were not able to detect a global effect of asexual reproduction to

decrease performance. This last finding points to physiological or genetic compensation for the

putative accumulation of deleterious mutations in mitochondrial genomes in asexual *P.*

antipodarum. Together, our results establish *P. antipodarum* as a model system for evaluating

the consequences of sexual reproduction for mitochondrial function and evolution and for

evaluating the strength and efficacy of selection against male-harming mitochondrial mutations

in sexual populations.

DATA ACCESSIBILITY

Oxygen consumption, behavioral, and mitochondrial membrane potential data have been posted

to https://github.com/jsharbrough/Potamo_mt_function and will be deposited in Dryad

(https://doi.org/10.5061/dryad.z08kprr8w) and FigShare

(https://doi.org/10.6084/m9.figshare.11782128).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

FUNDING

This work was funded by the National Science Foundation [MCB – 1122176, DEB – 1310825,

DEB – 1753851]; and the Iowa Academy of Sciences [ISF #13-10].

ACKNOWLEDGEMENTS

This work is dedicated in loving memory to J.J. Neiman-Brown, whose smile and curiosity give

inspiration to us all. This manuscript was written in part during the Biology Department Weekly

Writing Retreat at Colorado State University. Flow cytometry was performed at UI's Flow

Cytometry Facility. We thank two anonymous reviewers for comments on an earlier version of

this manuscript, and Laura Bankers, Kaitlin Hatcher, Katelyn Larkin, and Kyle McElroy for snail

collections. We thank the organizers, especially Karen Burnett, and all the participants of the

SICB 2020 Building Bridges from Genome to Phenome: Methods, Molecules, and Models

symposium for their participation in the symposium and helpful feedback on this work.

 TABLES

Table 1. Summary of source populations of *Potamopyrgus antipodarum* sampled from New Zealand lake collections.

Oxygen consumption assay

Lake	Latitude, Longitude	Sexual	Asexual	N/A[1]	Male	Female
Alexandrina	-43.900476, 170.453978	12	3	11	-	26
Clearwater	-43.602131, 171.043917	-	3	17	-	20
Kaniere	-42.832886, 171.14759	16	-	12	-	28
Paringa	-43.713068, 169.411348	4	-	20	-	24
Rotoroa	-41.855414, 172.637882	-	16	22	-	38
Selfe	-43.237765, 171.520449	-	3	19	-	22
Total	-	**32**	**25**	**101**	**0**	**158**

Behavior and mitochondrial membrane potential assays[2]

Lake	Latitude, Longitude	Sexual	Asexual	N/A	Male	Female
Alexandrina	-43.900476, 170.453978	3	-	-	3	-
Ellery	-44.046898, 168.654261	2	3	-	-	5
Kaniere	-42.832886, 171.14759	5	1	-	4	2
Mapourika	-43.315212, 170.204061	8	2	-	6	4
Rotoroa	-41.855414, 172.637882	4	1	-	-	5
Selfe	-43.237765, 171.520449	9	8	-	9	8
Total	-	**31**	**15**	**0**	**22**	**24**

[1] – N/A refers to snails that did not survive all three temperature trials

[2] – Same individual snails were used in behavioral and mitochondrial membrane potential assays

Table 2. Fixed-effect only and mixed-effects models of select predictors on oxygen consumption, righting time, emergence time, and mitochondrial membrane potential.

		Oxygen consumption[1]			
Model	**Factor**	**χ^2**	**df**	**p**	**Non-significant predictors[2]**
	Intercept[3]	5.32	1	0.0211	Lake-of-Origin × Mass × Temperature; Mass × Reproductive
a	Mass	10.82	1	0.0010	Mode × Temperature; Lake-of-Origin × Mass; Mass ×
	Temperature	45.91	2	< 0.0001	Temperature; Mass × Reproductive Mode; Reproductive Mode ×
	Lake-of-Origin	12.74	5	0.0260	Temperature; Reproductive Mode; Lake-of-Origin × Temperatu

		Emergence Time[4]			
Model	**Factor**	**χ^2**	**df**	**p**	**Non-significant predictors**
	Intercept[3]	138.14	1	< 0.0001	Reproductive Mode × Sex × Temperature; Sex × Temperature
b	Temperature	48.53	2	< 0.0001	Lake-of-Origin × Temperature; Reproductive Mode ×
	Lake-of-Origin	11.34	5	0.0450	Temperature; Reproductive Mode × Sex; Reproductive Mode; S

		Righting time[5]			
Model	**Factor**	**χ^2**	**df**	**p**	**Non-significant predictors**
	Intercept[3]	238.80	1	< 0.0001	
	Temperature	73.36	2	< 0.0001	Reproductive Mode × Sex × Temperature; Reproductiv
c	Reproductive Mode	4.48	1	0.0343	Mode × Temperature; Sex × Temperature; Lake-of-Orig
	Sex	2.49	1	0.12	× Temperature; Lake-of-Origin
	Reproductive Mode × Sex	6.52	1	0.0107	

		Mitochondrial membrane potential[6]			
Model	**Factor**	**Sum of Squares**	**df**	**F**	**p**	**Non-significant predictors**
	Intercept	3.911	1	64.53	< 0.0001	
d	Sex	0.42	1	6.84	0.0122	Reproductive Mode × Sex; Reproductive Mode; Lake-of-orig
	Residuals	2.67	44			

[1] – Type III Repeated-Measures Analysis of Deviance χ^2 Test of O$_2$ consumption per hour per gram

[2] – Semi-colon delimited list of non-significant predictors in order of elimination from the model (i.e., worst predictor listed first)

[3] – Snail ID was fit as a random intercept

[4] – Type III Repeated-Measures Analysis of Deviance χ^2 Test of Box-Cox-transformed emergence time (λ = -0.2845)

[5] – Type III Repeated-Measures Analysis of Deviance χ^2 Test of log-transformed righting times

[6] – Type III Analysis of Variance F Test of log-transformed ratios of red: green in mitochondrial extracts

Table S1. Fixed-effect only and mixed-effects models of select predictors on oxygen consumption, righting time, emergenc time, and mitochondrial membrane potential nested within lakes.

Oxygen consumption within Lake Alexandrina[1]

Model	Factor	χ^2	df	p	Non-significant predictors[2]
e	Intercept[3]	57.74	1	< 0.0001	Mass × Reproductive Mode × Temperature; Mass ×
	Mass	7.36	1	0.0067	Temperature; Reproductive Mode × Temperature; Ma
	Temperature	18.96	2	< 0.0001	× Reproductive Mode; Reproductive Mode

Emergence time within Lake Selfe[4]

Model	Factor	χ^2	df	p	Non-significant predictors
f	Intercept[3]	478.14	1	< 0.0001	Reproductive Mode × Sex × Temperature; Reproductive Mode × Sex; Reproductive Mode × Temperature; Sex × Temperature; Reproductive Mod Sex
	Temperature	26.85	2	< 0.0001	

Righting time within Lake Selfe[5]

Model	Factor	χ^2	df	p	Non-significant predictors
g	Intercept[3]	184.24	1	< 0.0001	Reproductive Mode × Sex × Temperature; Reproductive Mode × Temperatu Reproductive Mode × Sex; Reproductive Mode; Sex × Temperature; Sex
	Temperature	51.40	2	< 0.0001	

Mitochondrial membrane potential within Lake Selfe[6]

Model	Factor	Sum of Squares	df	F	p	Non-significant predictors
h	Intercept	0.95	1	15.04	0.0015	Reproductive Mode × Sex; Reproductive Mode
	Sex	0.42	1	6.58	0.0216	
	Residuals	0.95	15			

Emergence time within asexuals from Lake Selfe[4]

Model	Factor	χ^2	df	p	Non-significant predictors
i	Intercept[3]	233.31	1	< 0.0001	Sex × Temperature; Sex
	Temperature	11.64	2	0.0030	

Righting time within asexuals from Lake Selfe[5]

Model	Factor	χ^2	df	p	Non-significant predictors
j	Intercept[3]	89.48	1	< 0.0001	Sex × Temperature; Sex
	Temperature	27.75	2	< 0.0001	

Mitochondrial membrane potential within asexuals

Model	Factor	Sum of Squares	df	F	p	Non-significant predictors
k	Intercept	1.88	1	63.65	< 0.0001	Lake-of-Origin
	Sex	0.41	1	14.01	0.00246	
	Residuals	0.38	13			

Emergence time within sexuals from Lakes Mapourika and Selfe[4]

	Factor	χ^2	df	p	Non-significant predictors
l	Intercept[3]	494.44	1	< 0.0001	Lake-of-Origin × Sex × Temperature; Sex × Temperature; Lake-o Origin × Temperature; Lake-of-Origin × Sex; Lake-of-Origin; Se
	Temperature	16.19	2	0.00030	

Righting time within sexuals from Lakes Mapourika and Selfe[5]

	Factor	χ^2	df	p	Non-significant predictors
m	Intercept[3]	177.47	1	< 0.0001	Lake-of-Origin × Sex × Temperature; Lake-of-Origin × Temperatu Sex × Temperature; Lake-of-Origin × Sex; Lake-of-Origin; Sex
	Temperature	22.06	2	< 0.0001	

Mitochondrial membrane potential within sexuals from Lakes Mapourika and Selfe[6]

Model	Factor	Sum of Squares	df	*F*	*p*	Non-significant predictors
n	No significant predictors found					Lake-of-Origin × Sex; Sex; Lake-of-Origin

Emergence time within males from Lake Selfe[4]

Model	Factor	χ^2	df	*p*	Non-significant predictors
o	Intercept[3]	196.15	1	< 0.0001	Reproductive Mode × Temperature; Reproductive Mc
	Temperature	11.66	2	0.0029	

Righting time within males from Lake Selfe[5]

Model	Factor	χ^2	df	*p*	Non-significant predictors
p	Intercept[3]	104.39	1	< 0.0001	Reproductive Mode × Temperature; Reproductive Mc
	Temperature	20.79	2	< 0.0001	

Mitochondrial membrane potential within males from Lake Selfe[6]

Model	Factor	Sum of Squares	df	*F*	*p*	Non-significant predictors
q	No significant predictors found					Reproductive Mode

Emergence time within females from Lakes Ellery, Mapourika, and Selfe[4]

Model	Factor	χ^2	df	*p*	Non-significant predictors
r	Intercept[3]	601.98	1	< 0.0001	Lake-of-Origin × Reproductive Mode × Temperature; Lake-of-Ori × Reproductive Mode; Reproductive Mode × Temperature; Reproductive Mode; Lake-of-Origin × Temperature; Lake-of-Orig
	Temperature	19.84	2	< 0.0001	

Righting time within females from Lakes Ellery, Mapourika, and Selfe[5]

Model	Factor	χ^2	df	*p*	Non-significant predictors
s	Intercept[3]	87.62	1	< 0.0001	Lake-of-Origin × Reproductive Mode × Temperature Reproductive Mode × Temperature; Lake-of-Origin Reproductive Mode; Reproductive Mode
	Temperature	11.13	2	0.00383	
	Lake-of-Origin	12.67	2	0.00177	
	Temperature × Lake-of-Origin	12.99	4	0.01134	

Mitochondrial membrane potential within females from Lakes Ellery, Mapourika, and Selfe[6]

Model	Factor	Sum of Squares	df	*F*	*p*	Non-significant predictors
t	No significant predictors found					Reproductive Mode × Lake-of-Origin; Reproductive Mode; Lake-of-Origin

Mitochondrial membrane potential within sexuals

Model	Factor	Sum of Squares	df	*F*	*p*	Non-significant predictors
u	No significant predictors found					Lake-of-Origin; Sex

[1] – Type III Repeated-Measures Analysis of Deviance χ^2 Test of O_2 consumption per hour per gram

[2] – Non-significant predictors listed in order of elimination from the model

[3] – Snail ID was fit as a random intercept

[4] – Type III Repeated-Measures Analysis of Deviance χ^2 Test of Box-Cox-transformed emergence time (λ = -0.2845)

[5] – Type III Repeated-Measures Analysis of Deviance χ^2 Test of log-transformed righting times

[6] – Type III Analysis of Variance *F* Test of log-transformed ratios of red: green in mitochondrial extracts

FIGURES

Figure 1. Sampling *P. antipodarum* from New Zealand lakes. Samples from four lakes were

used in both experiments (grey circles – Lakes Alexandrina, Kaniere, Rotoroa, and Selfe),

samples from two lakes were used in the oxygen consumption experiment only (blue circles –

Lakes Clearwater and Paringa), and samples from a different two lakes were used for the

behavioral and mitochondrial membrane potential experiments only (orange circles – Lakes

Ellery and Mapourika).

Figure 2. Mitochondrial phenotypic variation across New Zealand lake populations of *P. antipodarum*. a) Oxygen consumption/hour/gram measured at 16ºC (gray), 22ºC (blue), and 30ºC (orange) in snails from six New Zealand lakes. There was a significant effect of mass, temperature, and lake-of-origin interaction on O_2 consumption (Table 2a). Red letters indicate *post hoc* statistical groupings. **b)** Time to emergence following a startling stimulus was measured at 16ºC (gray), 22ºC (blue), and 30ºC (orange) in snails from six New Zealand lakes. There was a significant effect of the temperature and lake-of-origin on emergence time (Table 2b). **c)** Time required for righting after being flipped was measured at 16ºC (gray), 22ºC (blue), and 30ºC

462   (orange) in snails from six New Zealand lakes. There was a significant effect of temperature,

463   reproductive mode, and reproductive mode × sex interaction on righting time (Table 2c). **d)** Log-

464   transformed ratios of red: green fluorescence following treatment with JC1 were measured using

465   flow cytometry. There was a significant effect of sex on red: green ratios (Table 2d).

Figure 3. Mitochondrial phenotypic variation across reproductive mode among field-collected *P. antipodarum*. Measurements collected from sexual individuals are depicted in blue and measurements collected from asexual individuals are depicted in red. **a)** Oxygen consumption per hour per gram assayed at 16ºC, 22ºC, and 30ºC. **b)** Time to emergence following a startling stimulus after incubation at 16ºC, 22ºC, and 30ºC. Two outliers are not shown for the sake of display. **c)** Righting time following being incubated at 16ºC, 22ºC, and

30ºC, and subsequently flipped over. **d)** Log-transformed ratios of red: green following treatment

with JC1 are depicted in sexual vs. asexual individuals.

Figure 4. Mitochondrial phenotypic variation across sexes among field-collected *P.*

***antipodarum*.** Measurements collected from females are depicted in blue and measurements

collected from males are depicted in yellow. **a)** Time to emergence following a startling

stimulus. **b)** Righting time after being flipped, **c)** Log-transformed ratios of red: green following

treatment with JC1.

Figure S1. Individual snail reaction norms across lakes in response to temperature stress.

Lines in each box (lake) represent individual reaction norms collected across three temperatures

(16ºC, 22ºC, and 30ºC) for each of three mitochondria-related phenotypes. Red lines represent

asexual individuals and blue lines represent sexual individuals. **a)** Oxygen consumption per hour

per gram. **b)** Time to emergence following a startling stimulus. **c)** Righting time after being

flipped.

Figure S2. Snail mass and oxygen consumption. a) Snail mass is positively correlated with O_2

consumption at 16ºC. **b)** Boxplot showing all datapoints (filled circles) depicting snail mass

across lakes. Snail mass varies across snails sampled from different lakes. Lowercase letters

above boxplots indicate statistical groupings at the α=0.05 level based on pairwise t-tests,

correcting for multiple comparisons using the Holm procedure.

Figure S3. QQ plots used to identify appropriate statistical transformations. Norm quantiles

plotted against: **a)** Oxygen consumption model residuals. **b)** Power-transformed emergence time

model residuals ($\lambda_E = -0.2845$). **c)** Log-transformed righting time model residuals. **d)** Log-

transformed ratio of red: green fluorescence following treatment with JC1.

REFERENCES

1. Chen JH, Hales CN, Ozanne SE. DNA damage, cellular senescence and organismal ageing: causal or correlative? Nucleic Acids Res 2007;35(22):7417-28.

2. Pike TW, Blount JD, Bjerkeng B, Lindstrom J, Metcalfe NB. Carotenoids, oxidative stress and female mating preference for longer lived males. Proc Biol Sci 2007;274(1618):1591-6.

3. Dowling DK, Friberg U, Lindell J. Evolutionary implications of non-neutral mitochondrial genetic variation. Trends Ecol Evol 2008;23(10):546-54.

4. Barreto FS, Burton RS. Elevated oxidative damage is correlated with reduced fitness in interpopulation hybrids of a marine copepod. Proc Biol Sci 2013;280(1767):20131521.

5. Dowling DK. Evolutionary perspectives on the links between mitochondrial genotype and disease phenotype. Biochim Biophys Acta 2014;1840(4):1393-403.

6. Blier PU, Dufresne F, Burton RS. Natural selection and the evolution of mtDNA-encoded peptides: evidence for intergenomic co-adaptation. Trends Genet 2001;17(7):400-6.

7. Montooth KL, Abt DN, Hofmann JW, Rand DM. Comparative genomics of *Drosophila* mtDNA: novel features of conservation and change across functional domains and lineages. J Mol 2009;69(1):94-114.

8. Neiman M, Hehman G, Miller JT, Logsdon JM, Jr., Taylor DR. Accelerated mutation accumulation in asexual lineages of a freshwater snail. Mol Biol Evol 2010;27(4):954-63.

9. Castellana S, Vicario S, Saccone C. Evolutionary patterns of the mitochondrial genome in Metazoa: exploring the role of mutation and selection in mitochondrial protein coding genes. Genome Biol Evol 2011;3:1067-1079.

10. Zhang F, Broughton RE. Mitochondrial-nuclear interactions: compensatory evolution or variable functional constraint among vertebrate oxidative phosphorylation genes? Genome Biol Evol 2013;5(10):1781-91.

11. Havird JC, Sloan DB. The roles of mutation, selection, and expression in determining relative rates of evolution in mitochondrial versus nuclear genomes. Mol Biol Evol 2016;33(12):3042-3053.

12. Ballard JW, Whitlock MC. The incomplete natural history of mitochondria. Mol Ecol 2004;13(4):729-44.

13. Galtier N, Nabholz B, Glemin S, Hurst GD. Mitochondrial DNA as a marker of molecular diversity: a reappraisal. Mol Ecol 2009;18(22):4541-50.

14. Bock DG, Andrew RL, Rieseberg LH. On the adaptive value of cytoplasmic genomes in plants. Mol Ecol 2014;23(20):4899-911.

15.   Dobler R, Rogell B, Budar F, Dowling DK. A meta-analysis of the strength and nature of cytoplasmic genetic effects. J Evol Biol 2014;27(10):2021-34.

16.   Ellison CK, Burton RS. Disruption of mitochondrial function in interpopulation hybrids of *Tigriopus californicus*. Evolution 2006;60(7):1382-91.

17.   Sloan DB, Havird JC, Sharbrough J. The on-again, off-again relationship between mitochondrial genomes and species boundaries. Mol Ecol 2017;26(8):2212-2236.

18.   Hebert PD, Ratnasingham S, deWaard JR. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. Proc Biol Sci 2003;270 Suppl 1:S96-9.

19.   Gabriel W, Lynch M, Burger R. Muller's ratchet and mutational meltdowns. Evolution 1993;47(6):1744-1757.

20.   Ursi S, Pedersen M, Plastino E, Snoeijs P. Intraspecific variation of photosynthesis, respiration and photoprotective carotenoids in *Gracilaria birdiae* (Gracilariales : Rhodophyta). Mar Biol 2003;142(5):997-1007.

21.   Sadowska ET, Labocha MK, Baliga K, Stanisz A, Wroblewska AK, Jagusiak W, Koteja P. Genetic correlations between basal and maximum metabolic rates in a wild rodent: Consequences for evolution of endothermy. Evolution 2005;59(3):672-681.

22.   Nilsson JA, Akesson M, Nilsson JF. Heritability of resting metabolic rate in a wild population of blue tits. J Evol Biol 2009;22(9):1867-1874.

23.   Arnqvist G, Dowling DK, Eady P, Gay L, Tregenza T, Tuda M, Hosken DJ. Genetic architecture of metabolic rate: environment specific epistasis between mitochondrial and nuclear genes in an insect. Evolution 2010;64(12):3354-63.

24.   Oellermann M, Portner HO, Mark FC. Mitochondrial dynamics underlying thermal plasticity of cuttlefish (*Sepia officinalis*) hearts. J Exp Biol 2012;215(Pt 17):2992-3000.

25.   Chung DJ, Bryant HJ, Schulte PM. Thermal acclimation and subspecies-specific effects on heart and brain mitochondrial performance in a eurythermal teleost (*Fundulus heteroclitus*). J Exp Biol 2017;220(Pt 8):1459-1471.

26.   Nespolo RF, Figueroa J, Solano-Iguaran JJ. Studying the evolutionary significance of thermal adaptation in ectotherms: The diversification of amphibians' energetics. J Therm Biol 2017;68(Pt A):5-13.

27.   Sharbrough J, Cruise JL, Beetch M, Enright NM, Neiman M. Genetic variation for mitochondrial function in the New Zealand freshwater snail *Potamopyrgus antipodarum*. J Hered 2017;108(7):759-768.

28.   Simonson TS, Yang YZ, Huff CD, Yun HX, Qin G, Witherspoon DJ, Bai ZZ, Lorenzo FR, Xing JC, Jorde LB and others. Genetic evidence for high-altitude adaptation in Tibet. Science 2010;329(5987):72-75.

29.   Storz JF, Scott GR, Cheviron ZA. Phenotypic plasticity and genetic adaptation to high-altitude hypoxia in vertebrates. J Exp Biol 2010;213(24):4125-4136.

30.   Clarke A, Johnston NM. Scaling of metabolic rate with body mass and temperature in teleost fish. J Anim Ecol 1999;68(5):893-905.

31.   Montooth KL, Marden JH, Clark AG. Mapping determinants of variation in energy metabolism, respiration and flight in *Drosophila*. Genetics 2003;165(2):623-35.

32.   Adams KL, Qiu YL, Stoutemyer M, Palmer JD. Punctuated evolution of mitochondrial gene content: high and variable rates of mitochondrial gene loss and transfer to the nucleus during angiosperm evolution. Proc Natl Acad Sci U S A 2002;99(15):9905-12.

33.   Rutter J, Winge DR, Schiffman JD. Succinate dehydrogenase - Assembly, regulation and role in human disease. Mitochondrion 2010;10(4):393-401.

34.   Rand DM, Haney RA, Fry AJ. Cytonuclear coevolution: the genomics of cooperation. Trends Ecol Evol 2004;19(12):645-653.

35.   Camus MF, Dowling DK. Mitochondrial genetic effects on reproductive success: signatures of positive intrasexual, but negative intersexual pleiotropy. Proc Biol Sci 2018;285(1879).

36.   Adrion JR, White PS, Montooth KL. The roles of compensatory evolution and constraint in aminoacyl tRNA synthetase evolution. Mol Biol Evol 2016;33(1):152-61.

37.   Osada N, Akashi H. Mitochondrial-nuclear interactions and accelerated compensatory evolution: evidence from the primate cytochrome C oxidase complex. Mol Biol Evol 2012;29(1):337-46.

38.   Barreto FS, Burton RS. Evidence for compensatory evolution of ribosomal proteins in response to rapid divergence of mitochondrial rRNA. Mol Biol Evol 2013;30(2):310-4.

39.   Sloan DB, Triant DA, Wu M, Taylor DR. Cytonuclear interactions and relaxed selection accelerate sequence evolution in organelle ribosomes. Mol Biol Evol 2014;31(3):673-82.

40.   van der Sluis EO, Bauerschmitt H, Becker T, Mielke T, Frauenfeld J, Berninghausen O, Neupert W, Herrmann JM, Beckmann R. Parallel structural evolution of mitochondrial ribosomes and OXPHOS complexes. Genome Biol Evol 2015;7(5):1235-51.

41.   Yan Z, Ye G, Werren JH. Evolutionary rate correlation between mitochondrial-encoded and mitochondria-associated nuclear-encoded proteins in insects. Mol Biol Evol 2019;36(5):1022-1036.

42.   Dowling DK, Friberg U, Arnqvist G. A comparison of nuclear and cytoplasmic genetic effects on sperm competitiveness and female remating in a seed beetle. J Evol Biol 2007;20(6):2113-2125.

43.   Hoekstra LA, Siddiq MA, Montooth KL. Pleiotropic effects of a mitochondrial-nuclear incompatibility depend upon the accelerating effect of temperature in *Drosophila*. Genetics 2013;195(3):1129-+.

44.   Paliwal S, Fiumera AC, Fiumera HL. Mitochondrial-nuclear epistasis contributes to phenotypic variation and coadaptation in natural isolates of *Saccharomyces cerevisiae*. Genetics 2014;198(3):1251-65.

45.   Touzet P, Meyer EH. Cytoplasmic male sterility and mitochondrial metabolism in plants. Mitochondrion 2014;19 Pt B:166-71.

46.   Hill GE, Havird JC, Sloan DB, Burton RS, Greening C, Dowling DK. Assessing the fitness consequences of mitonuclear interactions in natural populations. Biol Rev Camb Philos Soc 2019;94(3):1089-1104.

47.   Neiman M, Linksvayer TA. The conversion of variance and the evolutionary potential of restricted recombination. Heredity (Edinb) 2006;96(2):111-21.

48.   Clark AG. Natural selection with nuclear and cytoplasmic transmission. I. A deterministic model. Genetics 1984;107(4):679-701.

49.   Rand DM, Clark AG, Kann LM. Sexually antagonistic cytonuclear fitness interactions in *Drosophila melanogaster*. Genetics 2001;159(1):173-87.

50.   Neiman M, Taylor DR. The causes of mutation accumulation in mitochondrial genomes. Proc Biol Sci 2009;276(1660):1201-9.

51.   Muller HJ. Some genetic aspects of sex. Am Nat 1932;66(703):118-138.

52.   Hill WG, Robertson A. The effect of linkage on limits to artificial selection. Genet Res 1966;8(3):269-94.

53.   Winterbourn M. Population studies on the new zealand freshwater gastropod, *Potamoptrgus antipodarum* (Gray). J Molluscan Stud 1970;39(2-3):139-149.

54.   Lively CM. Evidence from a New Zealand snail for the maintenance of sex by parasitism. Nature 1987;328(6130):519-521.

55.   Jokela J, Lively CM, Dybdahl MF, Fox JA. Evidence for a cost of sex in the freshwater snail *Potamopyrgus antipodarum*. Ecology 1997;78(2):452-460.

56.   Neiman M, Lively CM. Pleistocene glaciation is implicated in the phylogeographical structure of *Potamopyrgus antipodarum*, a New Zealand snail. Mol Ecol 2004;13(10):3085-3098.

57. Neiman M, Paczesniak D, Soper DM, Baldwin AT, Hehman G. Wide variation in ploidy level and genome size in a New Zealand freshwater snail with coexisting sexual and asexual lineages. Evolution 2011;65(11):3202-16.

58. Sharbrough J, Luse M, Boore JL, Logsdon JM, Jr., Neiman M. Radical amino acid mutations persist longer in the absence of sex. Evolution 2018.

59. Paczesniak D, Jokela J, Larkin K, Neiman M. Discordance between nuclear and mitochondrial genomes in sexual and asexual lineages of the freshwater snail *Potamopyrgus antipodarum*. Mol Ecol 2013;22(18):4695-4710.

60. Neiman M, Larkin K, Thompson AR, Wilton P. Male offspring production by asexual *Potamopyrgus antipodarum*, a New Zealand snail. Heredity (Edinb) 2012;109(1):57-62.

61. Bankers L, Fields P, McElroy KE, Boore JL, Logsdon JM, Neiman M. Genomic evidence for population-specific responses to co-evolving parasites in a New Zealand freshwater snail. Mol Ecol 2017;26(14):3663-3675.

62. Zachar N, Neiman M. Profound effects of population density on fitness-related traits in an invasive freshwater snail. PLoS One 2013;8(11):e80067.

63. Abele E, Philip E, Gonzalez PM, Puntarulo S. Marine invertebrate mitochondria and oxidative stress. Front Biosci 2007;12:933-46.

64. Dybdahl MF, Kane SL. Adaptation vs. phenotypic plasticity in the success of a clonal invader. Ecology 2005;86(6):1592-1601.

65. Garner DL, Thomas CA. Organelle-specific probe JC-1 identifies membrane potential differences in the mitochondrial function of bovine sperm. Mol Reprod Dev 1999;53(2):222-9.

66. Wallace C. Parthenogenesis, sex and chromosomes in *Potamopyrgus*. J Molluscan Stud 1992;58:93-107.

67. Krist AC, Kay AD, Larkin K, Neiman M. Response to phosphorus limitation varies among lake populations of the freshwater snail *Potamopyrgus antipodarum*. PLoS One 2014;9(1):e85845.

68. Ribeiro Jr PJ, Diggle PJ, Ribeiro Jr MPJ, Suggests M. The geoR package. R news 2007;1(2):14-18.

69. Team RC. R: A language and environment for statistical computing [Internet]. Vienna, Austria; 2014. 2017.

70. Bates D, Sarkar D, Bates MD, Matrix L. The lme4 package. R package version 2007;2(1):74.

71.   Kuznetsova A, Brockhoff PB, Christensen RHB. Package 'lmerTest'. R package version 2015;2(0).

72.   Lenth R. emmeans: Estimated marginal means, aka least-squares means. R package version 1.4. 2019. 2019.

73.   Harrell Jr F. with contributions from CD and many others (2018) Hmisc: Harrell miscellaneous. R package version 4.1-1. 2019.

74.   Holm S. A simple sequentially rejective multiple test procedure. Scand J Stat 1979;6(2):65-70.

75.   Lively CM, Jokela J. Clinal variation for local adaptation in a host-parasite interaction.Proc Royal Soc B 1996;263(1372):891-897.

76.   Krist AC, Lively CM, Levri EP, Jokela J. Spatial variation in susceptibility to infection in a snail-trematode interaction. Parasitology 2000;121:395-401.

77.   Dybdahl MF, Krist AC. Genotypic vs. condition effects on parasite-driven rare advantage. J Evol Biol 2004;17(5):967-973.

78.   Jokela J, Dybdahl MF, Lively CM. The maintenance of sex, clonal dynamics, and host-parasite coevolution in a mixed population of sexual and asexual snails. Am Nat 2009;174:S43-S53.

79.   King KC, Delph LF, Jokela J, Lively CM. The geographic mosaic of sex and the Red Queen. Curr Biol 2009;19(17):1438-1441.

80.   Koskella B, Lively CM. Evidence for negative frequency-dependent selection during experimental coevolution of a freshwater snail and a sterilizing trematode. Evolution 2009;63(9):2213-21.

81.   Blasco-Costa I, Seppälä K, Feijen F, Zajac N, Klappert K, Jokela J. A new species of *Atriophallophorus* Deblock & Rosé, 1964 (Trematoda: Microphallidae) described from in vitro-grown adults and metacercariae from *Potamopyrgus antipodarum* (Gray, 1843) (Mollusca: Tateidae). J Helminthol 2019;94:e108.

82.   Larkin K, Tucci C, Neiman M. Effects of polyploidy and reproductive mode on life history trait expression. Ecol Evol 2016;6(3):765-78.

83.   Krist AC, Bankers L, Larkin K, Larson MD, Greenwood DJ, Dyck MA, Neiman M. Phosphorus availability in the source population influences response to dietary phosphorus quantity in a New Zealand freshwater snail. Oecologia 2017.

84.   Fisher RA. The genetical theory of natural selection. Oxford,: The Clarendon press; 1930. xiv, 272 p. p.

85.   Muller HJ. The relation of recombination to mutational advance. Mutat Res 1964;106:2-9.

86.   Birky CW, Jr., Walsh JB. Effects of linkage on rates of molecular evolution. Proc Natl Acad Sci U S A 1988;85(17):6414-8.

87.   Kondrashov AS. Classification of hypotheses on the advantage of amphimixis. J Hered 1993;84(5):372-87.

88.   Charlesworth B. The effects of deleterious mutations on evolution at linked sites. Genetics 2012;190(1):5-22.

89.   Normark BB, Moran NA. Opinion - Testing for the accumulation of deleterious mutations in asexual eukaryote genomes using molecular sequences. J Nat Hist 2000;34(9):1719-1729.

90.   Paland S, Lynch M. Transitions to asexuality result in excess amino acid substitutions. Science 2006;311(5763):990-992.

91.   Johnson SG, Howard RS. Contrasting patterns of synonymous and nonsynonymous sequence evolution in asexual and sexual freshwater snail lineages. Evolution 2007;61(11):2728-2735.

92.   Henry L, Schwander T, Crespi BJ. Deleterious mutation accumulation in asexual Timema stick insects. Mol Biol Evol 2012;29(1):401-8.

93.   Harrison JS, Burton RS. Tracing hybrid incompatibilities to single amino acid substitutions. Mol Biol Evol 2006;23(3):559-64.

94.   Matoo OB, Julick CR, Montooth KL. Genetic variation for ontogenetic shifts in metabolism underlies physiological homeostasis in *Drosophila*. Genetics 2019;212(2):537-552.

95.   Otto SP. The evolutionary consequences of polyploidy. Cell 2007;131(3):452-62.

96.   Beaulieu JM, Leitch IJ, Patel S, Pendharkar A, Knight CA. Genome size is a strong predictor of cell size and stomatal density in angiosperms. New Phytol 2008;179(4):975-86.

97.   Sharbrough J, Conover JL, Tate JA, Wendel JF, Sloan DB. Cytonuclear responses to genome doubling. Am J Bot 2017;104(9):1277-1280.

98.   Frank SA, Hurst LD. Mitochondria and male disease. Nature 1996;383(6597):224-224.

99.   Gemmell NJ, Metcalf VJ, Allendorf FW. Mother's curse: the effect of mtDNA on individual fitness and population viability. Trends Ecol Evol 2004;19(5):238-244.

100.   Camus MF, Clancy DJ, Dowling DK. Mitochondria, maternal inheritance, and male aging. Curr Biol 2012;22(18):1717-1721.

101.   Innocenti P, Morrow EH, Dowling DK. Experimental evidence supports a sex-specific selective sieve in mitochondrial genome evolution. Science 2011;332(6031):845-848.

102.   Patel MR, Miriyala GK, Littleton AJ, Yang HK, Trinh K, Young JM, Kennedy SR, Yamashita YM, Pallanck LJ, Malik HS. A mitochondrial DNA hypomorph of cytochrome oxidase specifically impairs male fertility in *Drosophila melanogaster*. Elife 2016;5.

103.   Kuijper B, Lane N, Pomiankowski A. Can paternal leakage maintain sexually antagonistic polymorphism in the cytoplasm? J Evol Biol 2015;28(2):468-480.

104.   Wade MJ, Brandvain Y. Reversing mother's curse: selection on male mitochondrial fitness effects. Evolution 2009;63(4):1084-1089.

105.   Delph LF, Touzet P, Bailey MF. Merging theory and mechanism in studies of gynodioecy. Trends Ecol Evol 2007;22(1):17-24.