1 **The fruitENCODE project sheds light on the genetic and epigenetic basis of convergent**
2 **evolution of climacteric fruit ripening**
3

4 Peitao Lü[a,*], Sheng Yu[a,*], Ning Zhu[a,*], Yun-Ru Chen[a,*], Biyan Zhou[b,*], Yu Pan[c], David
5 Tzeng[a], Joao Paulo Fabi[d], Jordi Garcia-Mas[e], Nengui Ye[f], Jianhua Zhang[f], Donald Grierson[g],
6 Zhangjun Fei[h], Jim Giovannoni[h], Silin Zhong[a]
7
a. State Key Laboratory of Agrobiotechnology, School of Life Sciences, The Chinese
University of Hong Kong, Hong Kong, China.
b. College of Horticulture, South China Agricultural University, Guangzhou, China.
c. College of Horticulture and Landscape Architecture, Southwest University, China
d. Department of Food Science and Experimental Nutrition, FCF, University of Sao Paulo,
Sao Paulo, Brazil.
e. IRTA, Centre for Research in Agricultural Genomics CSIC-IRTA-UAB-UB, Barcelona,
Spain.
f. Department of Biology, Hong Kong Baptist University. Hong Kong, China.
g. Zhejiang Provincial Key Laboratory of Horticultural Plant Integrative Biology, Zhejiang
University, Hangzhou, Zhejiang, China.
h. Boyce Thompson Institute for Plant Research, Cornell University, Ithaca, New York,
USA.
* These authors contributed equally to this work.

Abstract
Fleshy fruit evolved independently multiple times during angiosperm history. Many
climacteric fruits utilize the hormone ethylene to regulate ripening. The fruitENCODE
project shows there are multiple evolutionary origins of the regulatory circuits that govern
climacteric fruit ripening. Eudicot climacteric fruits with recent whole-genome duplications
(WGDs) evolved their ripening regulatory systems using the duplicated floral identity genes,
while others without WGD utilised carpel senescence genes. The monocot banana uses both
leaf senescence and duplicated floral-identity genes, forming two interconnected regulatory
circuits. H3K27me3 plays a conserved role in restricting the expression of key ripening
regulators and their direct orthologs in both the ancestral dry fruit and non-climacteric fleshy
fruit species. Our findings suggest that evolution of climacteric ripening was constrained by
limited availability of signalling molecules and genetic and epigenetic materials, and WGD
provided new resources for plants to circumvent this limit. Understanding these different
ripening mechanisms makes it possible to design tailor-made ripening traits to improve
quality, yield and minimize postharvest losses.

38
Introduction
Fleshy fruit ripening is a developmental process unique to flowering plants, in which the
chemical and physiological properties such as colour, aroma, flavour, texture and nutritional
contents of the seed-bearing organ are dramatically altered, in order to attract frugivores as
seed dispersers[1,2]. As they are now a vital part of human and animal diets, understanding fruit
ripening is increasingly important for global food security.
45

Many economically important fruits such as apple, banana and tomato can be harvested
mature but unripe, stored and treated with the plant hormone ethylene to complete the
ripening process. They show a concomitant increase in respiration and ethylene biosynthesis
upon initiation of ripening, and are referred to as climacteric fruits. Non-climacteric fruits
like grape and strawberry lack these features and do not require ethylene for ripening.

Fleshy fruit is a classic example of convergent evolution, and it has occurred multiple times
in the history of angiosperms. However, its molecular basis remained largely unknown, as
most of the studies have focused on a single model fruit tomato[3]. Tomato produces ethylene
at the onset of ripening, and disrupting ethylene biosynthesis or perception inhibits ripening[4].
Besides ethylene, a series of transcription factors such as COLORLESS NON-RIPENING
(CNR), NON-RIPENING (NOR) and RIPENING INHIBITOR (RIN) are also required for
tomato fruit ripening[3,5,6]. Among them, the MADS-box transcription factor RIN is most
extensive studied, and its mutant alleles are widely used in commercial tomato production.
The MADS gene family is present in all eukaryotes and has expanded considerably in plants.
RIN belongs to the MIKC-clade, members of which are involved in floral organ development.
RIN can bind directly to the promoter of key ripening genes, such as those involved in colour
change, fruit softening, aroma volatile production and sugar metabolism, and is also
responsible for regulating ethylene biosynthesis genes such as *ACO1* and *ACS2*[7].

In ripening tomato fruit, ethylene biosynthesis is autocatalytic, which is historically referred
to as system II ethylene, to be distinguished from the self-inhibitory system I ethylene
production in other tissues including immature fruit[8]. The autocatalytic nature suggests that
there is a positive feedback loop controlling ethylene production during fruit ripening.
However, ethylene is a stress hormone involved in many plant developmental processes such
as senescence and abscission[4]. Such an autocatalytic system involving a diffusible signal
molecule poses a major threat to the plant itself and its neighbours, as any leakiness could
cause growth disruption or even tissue death. It has been shown that DNA methylation plays
an important role in restricting the expression of tomato ripening genes[5,7]. Their promoter
regions bound by RIN are demethylated during fruit development, and silencing the fruit-
specific DNA demethylase can inhibit ripening[9].

While tomato is still the predominant model for ripening research, many fleshy fruit genomes
have now been sequenced. Tomato is also the only species reported so far that expresses a
DNA demethylase during fruit ripening[9], and questions arise regarding the tomato model
itself and whether or not it is universal. In addition, tomato has experienced whole-genome
duplication (WGD) ~71 Myr, and key ripening regulators including *RIN* and ethylene
biosynthesis genes *ACO1* and *ACS2* are paralog members of duplicated gene families[10].
Hence, other species without a ripening-specific demethylase and WGD might have evolved
a different mechanism.

Some fruits also have unique ripening traits that are absent in tomato. For example, both
climacteric and non-climacteric genotypes exist in melon and pear[11,12]. It has also been
suggested that banana operates an additional leaf-like self-inhibitory ethylene system I during

ripening, and unlike tomato, once its autocatalytic ethylene production starts, it becomes
ethylene-independent[13].
92
It is not possible to resolve complex convergent traits in diverse taxa such as fleshy fruit
ripening by sequencing and comparing genomes if the convergence occurred through the
evolution of different genes and pathways, or if the regulatory interactions are different. To
tackle this, we have used an ENCODE style functional genomic approach to systematically
characterize the molecular circuits controlling ripening in 13 plant species. We found three
major circuits adopted by different plants in the evolution of climacteric fruit. Part of the
genes and epigenetic marks are surprisingly conserved even in dry and non-climacteric
species, suggesting that these different ethylene-dependent fleshy fruit ripening mechanisms
are evolved from existing pathways in the ancestral angiosperm's leaf and carpel tissues.
102
103 **Results**
104
The fruitENCODE data
Our current fruitENCODE dataset comprises seven climacteric fruit species (apple, banana,
melon, papaya, peach, pear and tomato), four non-climacteric species (cucumber, grape,
strawberry and watermelon) and two dry fruit species (*Arabidopsis* and rice)
(**Supplementary Table 1-18**). We have generated 361 transcriptome profiles from their
leaves and fruit tissues at different developmental stages. We have also generated 66 tissue-
specific accessibility chromatin profiles, and identified between 36,584 to 45,116 DHSs that
could be associated to 43.28% to 71.93% of annotated genes in each species. The *cis*-
regulatory elements in the DHSs enabled us to predict and experimentally validate the
regulatory circuits controlling climacteric fruit ripening, and further recreate each circuit in
tobacco leave. In addition, we have generated 43 methylomes and 76 histone modification
profiles to investigate which epigenetic factors are responsible for regulating these ripening
circuits. All datasets can be accessed from the fruitENCODE database.
118
Tomato, apple and pear fruits use a MADS positive feedback loop
We first reconstructed the ripening regulatory circuit in the model fruit tomato that has three
key components: ethylene, transcription factor RIN and DNA methylation. The main features
of ethylene signalling are conserved in plants and signal perception by the receptors leads to
the stabilization of the EIN3-type transcription factors, which activate the downstream
genes[4]. From the open chromatin dataset, we found an EIN3 binding motif in the promoter of
RIN, the functional significance of which was confirmed by our EIN3 ChIP-Seq (**Fig. 1a**).
RIN is a MADS-box transcription factor, which functions in a multimeric complex with
TAGL1 that is constantly expressed during fruit development[14]. We have also performed
ChIP-Seq for RIN and TAGL1, and found that they can directly target the rate-limiting
ripening ethylene synthesis genes *ACS2* and *ACO1* (**Fig. 1a**). In addition, our DHS and
methylome data showed that the EIN3 binding site in the *RIN* promoter is demethylated and
become accessible in ripening fruit tissues (**Fig. 1a**).
132

Our findings suggest that EIN3 and RIN-TAGL1 form a positive feedback loop to synthesize
the autocatalytic system II ethylene during tomato fruit ripening (**Fig. 1b**). In addition,
ripening genes are also coupled to the loop through RIN-TAGL1. Our ChIP-Seq data showed
that the RIN-TAGL1 targets well-known ripening genes such as *EXPANSIN*, *PG* and *PL* that
are involved in fruit softening*, PSY*, *LOX* and *SuSy* that are involved in colour change, aroma
production and sugar metabolism, respectively (**Supplementary Table 28** and
Supplementary note 10.5). Given the central role of the MADS-box transcription factors in
this ripening model, we named it the MADS positive feedback loop.

To test this MADS positive feedback loop, we used the tobacco leaf transient expression
system that has a functional ethylene signalling pathway and endogenous EIN3 and ACO
activities, but lacks the RIN-TAGL1 and ACS activity. When we expressed the tomato *ACS2*
and *RIN* under their native promoters, and *TAGL1* was supplied under a constitutive
CaMV35S promoter, spontaneous ethylene synthesis was observed (**Fig. 1c**). Addition of 1-
MCP, a competitive inhibitor of ethylene, or deletion of the EIN3 binding motif in the *RIN*
promoter, interrupted the positive feedback loop and prevented ethylene synthesis,
confirming the autocatalytic nature of the regulation of ethylene synthesis (**Fig. 1d**).

We have examined transcriptome and open chromatin data from other species and found that
apple and pear have similar ripening-fruit-specific MADS genes with EIN3 binding motifs,
while their ethylene biosynthesis genes *ACS* and *ACO* have MADS transcription factor
binding motifs. These observations suggests that they could operate a tomato-like MADS
positive feedback loop (**Extended Data Figs 1 and 2**). We have used the tobacco leaf system
to test whether the apple and pear genes could generate autocatalytic ethylene. The results
showed that their MADS loops are indeed functional, and EIN3 motif deletion confirmed that
the EIN3 and these MADS genes could formed a positive feedback loop (**Extended Data**
Figs 1 and 2).

Since ethylene is a stress hormone, it is vital for the plant to repress this loop in non-ripening
tissues and confine autocatalytic ethylene action to specific tissues or organs, such as
ripening fruits, abscission zones and senescing leaves. It has been shown that whole-genome
demethylation is required for tomato fruit ripening[7,9]. However, our RNA-Seq data showed
that the expression of the apple and pear demethylases is not ripening-specific, neither are the
demethylase of other fleshy fruits species we examined (**Supplementary Table 16**).

Our methylome datasets confirmed that there is no tomato-like global demethylation event in
apple and pear (**Supplementary Fig 6**). Although we could identify hundreds of ripening
genes with promoter DMRs in all seven climacteric fruits we studied (**Supplementary Table**
10-12), only in tomato is the master ripening regulator in the positive feedback loop, MADS-
box *RIN* associated with DMR. For example, we have identified 11,911, 17,231 and 17,455
genes associated with promoter hypo-DMR in tomato, apple and pear, respectively, including
the apple *PSY* and *PG* genes, which are involved in colour change and fruit softening,
respectively (**Supplementary Table 12**). But the two MADS-box transcription factor genes
in the apple positive feedback loops are not (**Extended Data Fig. 1**). These results suggest

that although DNA methylation change in the promoter DHS is widespread, it might not be a
conserved mechanism utilized by climacteric fruits to regulate ripening genes.
179
However, from the histone modification datasets, we found that tomato, apple and pear have
the repressive histone mark H3K27me3 on their ethylene biosynthesis and MADS gene loci,
whereas they are removed in the ripe fruit tissues (**Fig. 1a, Extended Data Figs 1 and 2**).
The fruitENCODE data also includes multiple cultivars and mutants, and we found that the
ethylene-independent pear cultivar Dangshanshuli contains hyper-H3K27me3 in its *ACS*
locus compared to the ethylene-dependent cultivar Williams (**Extended Data Fig. 2**). From
the tomato ripening mutant collection, we also observed that the non-ripening mutant *nor*
contains hyper-H3K27me3 in the *ACS2* and *RIN* loci (**Supplementary Fig. 13**). These
suggests that H3K27me3, instead of DNA methylation, might play a conserved role in
regulating the MADS type ripening positive feedback loop.
190
Peach, papaya and melon operate a NAC positive feedback loop
Tomato has a WGD ~71 Mya, while apple and pear have a common WGD ~40 Mya. These
plants have a similar positive feedback loop using paralog member of the duplicated SEP-
clade MADS transcription factor family that controls floral organ development, suggesting
that neofunctionalization following WGD could have played a vital part in the evolution of
MADS-type climacteric fruit ripening (**Extended Data Fig. 3**). However, climacteric fruits
such as peach, papaya and some climacteric melon cultivars can also produce and require
autocatalytic ethylene for ripening, and did not undergo recent WGD. Their MADS direct
orthologs are constantly expressed during fruit development (**Supplementary Tables 23, 24**
and 25). If they have evolved a positive feedback loop to synthesize the ripening ethylene, it
will require different transcription factors.
202
We have examined the expression pattern of the papaya, peach and melon ethylene
biosynthesis genes, and identified their *ACS* and *ACO* genes that are specifically expressed
during fruit ripening (**Supplementary Figs 14 and 15**, **Supplementary Table 17**). Also
from the DHS datasets, we found NAC transcription factor binding motifs in those *ACS* and
ACO gene promoters. Intriguingly, all three species have a *NAC* gene that has ripening-
specific expression pattern similar to the *MADS-RIN* genes in tomato (**Extended Data Fig. 4**,
Supplementary Tables 23, 24 and 25). NAC is one of the largest plant-specific transcription
factor families with members involved in many developmental processes such as senescence,
stress, cell wall formation, and embryo development. All three ripening *NAC* genes we
identified are direct orthologs of the *Arabidopsis* carpel senescence-related *NAC* transcription
factors *NARS1/2*, and are distantly related to the leaf-senescence-related *AtNAP*[15,16]
(**Extended Data Fig. 4**). In addition, we have found EIN3 motif in the promoter DHSs of
these *NAC*, suggesting that instead of relying on the duplicated floral identity *MADS* gene,
plants without WGD might have evolved a positive feedback loop using the existing carpel
senescence *NAC* genes (**Fig. 2**).
218
To test this, we have ectopically expressed the peach *NAC* (ppa007577m) and *ACS*
(ppa004774m) genes under their native promoters in tobacco leaves, and found that they are

capable of generating ethylene (**Fig. 2b, c**). The ethylene production could be blocked by
treatment with inhibitor 1-MCP, suggesting that it is autocatalytic. We have also used motif
deletion assay to confirm that EIN3 binding to the *NAC* gene promoter is required for the
autocatalytic ethylene production. These suggest that peach operates a NAC-type positive
feedback loop.
226
In tomato, the downstream ripening genes are directly coupled to the positive feedback loop
through the MADS-box transcription factor RIN-TAGL1. To test whether it is the same for
plants operating this NAC-type loop, we use ChIP-Seq to identify the direct targets of the
peach NAC transcription factor (**Supplementary Table 25**). Our data showed that it indeed
binds to the promoters of the ethylene biosynthesis genes *ACS* and *ACO*, as well as key fruit
ripening genes such as those involved in carotenoid pigment accumulation, volatile
secondary metabolite production, cell wall softening and sugar accumulation (**Fig. 2,**
Supplementary note 7.5). The peach DNA methylation and histone modification data
showed that genes involved in the NAC positive feedback loop are also associated with the
repressive histone mark H3K27me3 in leaves and immature fruit tissues, but not in the ripe
fruit (**Fig. 2a**).
238
239 In papaya and climacteric melon cultivar Védrantais, we found the same NAC-type positive
240 feedback loops, with key genes associated with H3K27me3 in non-ripening tissues
241 (**Extended Data Figs 5 and 6**). We also tested their loops using the tobacco system, and
242 performed ethylene inhibitor treatment and EIN3 motif deletion to confirm that they are
243 capable of generating autocatalytic ethylene (**Extended Data Figs 5 and 6**).
244
Interestingly, our melon dataset includes the non-climacteric cultivar *Piel de Sapo*, which
does not require ethylene for ripening. The QTL (*ETHQV6.3*) responsible for the ethylene-
independent phenotype has been mapped to a region, where its *NAC* gene (*MELO3C016540*)
involved in the positive feedback loop is located[17]. This gene is down-regulated in *Piel de*
Sapo ripe fruit, when compared to other cultivars, while its promoter is hypermethylated and
the genebody is associated hyper-H3K27me3 (**Extended Data Fig. 6**). Similarly, the peach
NAC gene involved in the loop is also located in a previously identified QTL associated with
the late-ripening phenotype[18], suggesting that ripening traits resulting from mutations in the
NAC-type loop might have been selected during the breeding process, similar to the
widespread used of MADS gene *RIN* mutant allele in the modern tomato cultivars.
255
Monocot banana operates a dual-loop system
Banana is also a climacteric fruit that requires autocatalytic ethylene to ripen, and has
experienced three recent WGD[19]. The autocatalytic ethylene production in other climacteric
fruits such as tomato and peach can be interrupted by the ethylene-inhibitor, a scenario that
we could reproduce in tobacco by ectopic expression of their MADS or NAC-type positive
feedback loops. However, inhibitor treatment is unable to interrupt the ethylene production in
banana after ripening has been initiated, indicating a transition from autocatalytic to ethylene
independent ripening[13]. Our data suggests that banana has two positive feedback loops, and

the second one is able to maintain the ethylene synthesis when the first ethylene-dependent
loop is blocked (**Fig. 3**).

The first banana loop is similar to the NAC-type positive feedback loop in eudicots such as
peach, papaya and melon. We found that the banana NAC transcription factor
(Ma06_t33980.1) could activate the ripening ethylene biosynthesis genes *ACS*
(Ma04_t35640.1) and *ACO* (Ma07_t19730.1), while its own promoter contains an EIN3
binding motif like peach NAC gene (**Fig. 3b, Supplementary Fig 12**). To test the first loop,
we ectopically expressed the banana *NAC* and *ACS* genes under their native promoters in
tobacco and found that they are sufficient to generate ethylene in an autocatalytic manner
(**Fig. 3c**). Ethylene inhibitor 1-MCP, as well as EIN3 motif deletion, could block ethylene
production, suggesting this is a functional NAC positive feedback loop.

It should be noted that this banana ripening *NAC* gene is a direct ortholog of the rice leaf
senescence transcription factor OsNAP[20], and is distinct from the carpel senescence-related
NAC genes utilized by the eudicots climacteric fruits mentioned above (**Extended Data Fig.**
4). We have examined the gene expression and histone modifications in young, matured and
senescence banana leaves, and found that this *NAC* gene is also expressed during leaf
senescence (**Fig. 3a, Supplementary Table 20**). Our epigenome data showed that genes in
the first banana positive feedback loop are associated with similar tissue-specific H3K27me3
marks as those in the eudicots fruits, except that the banana *NAC* gene is not associated with
H3K27me3 in senescence leaf (**Fig. 3**).

Both the eudicot MADS- and NAC-type positive feedback loops are directly coupled to the
downstream ripening genes, which is confirmed by ChIP-Seq using tomato and peach as
example. Without a suitable banana NAC antibody, we have used the dual luciferase assay in
tobacco to confirm that the banana NAC in the positive feedback loop is capable of activating
downstream ripening genes such as *EXPANSIN*, *PG, MAN* and *INV* (**Supplementary Fig.**
12).

Interestingly, utilizing a leaf- instead of a carpel-senescence NAC in banana fruit ripening
poses a dilemma for the plant as some ripening genes, such as *EXPANSIN*, are not expressed
in leaf. We found that the *EXPANSIN* locus is associated with H3K27me3 marks in leaf,
suggesting that banana might rely on epigenetic factors to restrict its expression to ripening
fruit tissues (**Fig 3a**). In contrast, we did not observe H3K27me3 targeting the ripening
EXPANSIN gene in MADS or NAC-type eudicot fruits, as their *MADS* and *NAC* genes are
ripening-specific and are not expressed in leaf.

It has been previously shown that three MADS transcription factors (MADS1/2/5) are
involved in banana fruit ripening[21,22]. *MADS1/2* and tomato *RIN* are in the SEP-clade, while
MADS5 and the tomato *TAGL1* are in the AG-clade. It is important to note, however, that
MADS1/2 and *MADS5* are not direct orthologs of the tomato *RIN* and *TAGL1* (**Extended**
Data Fig. 3). The expression *MADS1/2/5* genes could be detected in both immature and
ripening fruits, and they are only associated with H3K27me3 in the leaf tissue (**Fig. 3a**). We

found a NAC motif in the banana *MADS1* gene promoter and a MADS motif in the *NAC* promoter, suggesting that they could form a second positive feedback loop to bypass the first loop that is ethylene dependent (**Fig. 3b**).

To test the second loop, we first used the tobacco system to show that the first NAC loop could be blocked by either inhibitor 1-MCP treatment or EIN3 motif deletion (**Fig. 3c**). Adding a second loop with *MADS1* driven by its native promoter and *MADS2/5* with the constitutive 35S promoter enabled the tobacco leaves to synthesize 98% more ethylene ($p=2.91 \times 10^{-5}$) (**Fig. 3d**). Most importantly, ethylene inhibitor 1-MCP was unable to block the ethylene production when the second loop was present, mimicking the behaviour of the ripening banana fruit. We have also confirmed that the second loop is dependent on the interaction of the *NAC* and *MADS1* genes by deleting the NAC motif in the promoter of *MADS1*, as well as deleting the MADS motif in the *NAC* promoter (**Fig. 3d**). Taken together, our results showed that banana fruit ripening is controlled by a dual positive feedback loop that consists of both *NAC* and *MADS* genes. These banana *NAC* and *MADS* genes are also associated with tissue-specific H3K27me3 marks. But they are of a different evolutionary origin compared to the NAC and MADS genes used by other eudicot climacteric fruits.

Direct orthologs of the ripening genes in non-climacteric and dry fruit species
To investigate the possible evolutionary origin of these three types of ripening circuits, we have examined four non-climacteric fruits (cucumber, grape, strawberry and watermelon). None of them have recent WGD. These species have direct orthologs of the carpel senescence *NAC* with H3K27me3 modification change and a ripening-specific gene expression pattern similar to those in the NAC-type climacteric fruit peach, papaya and melon (**Fig. 4a, Extended Data Figs 7 to 10**). However, their *NAC* genes either lack the EIN3 motif in their promoter DHS, or the NAC motif is absent from their ethylene biosynthesis genes, both factors which would preclude participation in a positive ethylene feedback loop. We have also examined their *MADS* genes that are direct orthologs of the tomato *RIN*, and found that they are expressed in fruits, and are associated with H3K27me3 marks in leaves.

Strawberry appears to be an exception. It possesses all the required *cis*-regulatory elements in its *NAC* (*mrna31150.1*) and ethylene biosynthesis gene *ACS* (*mrna11391.1*) to form a positive feedback loop (**Supplementary Fig. 16**). Its ethylene biosynthesis gene *ACS*, however, is constantly repressed by H3K27me3 during fruit development and ripening (**Extended Data Fig. 9**), again hinting the important roles of the *cis*-regulatory elements and epigenetic marks in evolving climacteric fruit ripening.

The dry fruit-bearing plant *Arabidopsis* also has direct orthologs of the *NAC* and *MADS* genes that are involved in climacteric fruit ripening. We have examined their H3K27me3 levels in *Arabidopsis* leaves, carpel and senescence silique, which is the equivalent tissue of a ripening fruit (**Fig. 4b**). We found its *NAC* gene (AT3G15510.1) showed reduced H3K27me3 level during silique senescence, while the MADS gene *SEP4* (AT2G03710.1) was also associated with tissue-specific H3K27me3.

The unique dual-loop system used by banana utilizes a direct ortholog of the rice leaf-
senescence *OsNAP* gene. We have profiled the gene expression and histone modification in
young and senescence rice leaves, as well as its carpel tissues. Our data confirmed that the
OsNAP gene is expressed in the senescing leaves, and is associated with H3K27me3 in the
young leaves and carpel, but not in the senescence leaves (**Fig. 4c**).

Taken together, our finding showed that other eudicot and monocot plants have direct
orthologs of the climacteric fruit ripening genes. These genes are involved in leaf senescence,
carpel senescence or floral development, and are associated with tissue-specific H3K27me3
marks. As fleshy fruits are believed to have evolved independently in multiple angiosperm
lineages, this could suggest that climacteric fruit ripening could have evolved from existing
mechanisms controlling leaf and floral tissue gene expression.

Discussion
The data from the fruitENCODE project shows that there are three major evolutionary paths
of the regulatory circuits that govern climacteric fruit ripening (**Fig. 5**). It is common that
different species often evolve similar features when exposed to the same selection pressure.
However, the probability of complex traits such as climacteric fruit ripening originating
multiple times through similar trajectories should be very small, unless there is strong
constraint. Hence, it might not be a coincidence that the four eudicot climacteric fruits
without recent WGD evolved a NAC-type ripening circuit that can be traced back to carpel
senescence, while those with WGD could utilize the duplicated floral identity genes. We
hypothesize that using the existing carpel senescence pathway, acquisition of *cis*-regulatory
elements in the ethylene biosynthesis genes to complete a positive feedback loop and finally
evolve *cis*-regulatory elements in the downstream ripening genes, which enabled them to be
coupled to the loop, could be one of the simplest paths to achieve climacteric fruit ripening.
Our findings also suggest that the evolution of climacteric fruit ripening was constrained by
the limited set of suitable signalling molecules, genetic or epigenetic building blocks
available in the carpel tissues, and WGD provided new resources for plants to circumvent this
limit.

The fruitENCODE data also enabled us to systematically compare the epigenetic changes
associated ripening in seven climacteric plants. Our data shows widespread promoter DNA
methylation and DHS changes, as well as genebody H3K27me3 changes (**Supplementary**
Tables 11 and 15). For example, 17231, 1759 and 630 apple genes are associated with hypo-
DMR, tissue-specific DHS and hypo-H3K27me3 in the ripe fruit tissue, respectively.
However, these DNA methylation and DHS changes are not always conserved across species.
For example, the tomato *CNR* gene promoter is associated with two well-characterized
DMRs and tissue-specific DHS[5], while they are absent in other species. In addition, for plants
that operate a NAC-type positive feedback loop, only the NAC gene in peach is associated
with promoter DMR and DHS changes (**Fig. 2**). However, the H3K27me3 changes in the
ripening ethylene biosynthesis genes, as well as the ripening NAC and MADS transcription
factor genes are highly conserved, and their direct orthologs in *Arabidopsis* and rice also

carry similar tissue-specific H3K27me3 marks, suggesting that the ripening epigenetic
changes could be originated from existing mechanisms in the ancestral dry fruit species.
398
Angiosperms became abundant toward end of Cretaceous (~65 Mya). There have been many
evolutionary transitions from an ancestral dry fruit to fleshy fruit during this period. It is
difficult to trace when these diverse fleshy fruit ripening mechanisms evolved. A common
triplication occurred ~71 Mya in the Solanaceae family that includes the climacteric fruit
tomato and non-climacteric fruit potato. This suggests that tomato is likely to have acquired
its ripening MADS loop after the tomato-potato split. Apple and pear shared a recent WGD
event ~40 Mya, and their ripening *MADS*, *ACS* and *ACO* genes are direct orthologs, which
differ from the tomato ones (**Extended Data Fig. 3, Supplementary Figs 14 and 15**),
suggesting they have a common ancestor that bore climacteric fruit. Future study of the
ripening regulatory circuits in wild species as well as the relatives of the domesticated crops
might shed light on the evolutionary origin of their ripening circuits, and how the non-
climacteric fruits make use of their *NAC* and duplicated *MADS* genes.
411
Banana is a monocot that has diverged from eudicots over 100 Mya. The ripening regulatory
system in banana is the most complex, but it is the only monocot fruit with high quality
genome that we have examined so far. It has experienced three WGDs, and utilizes a leaf-
senescence NAC transcription factor, as well as three MADS-box transcription factors to
complete the dual-loop system. However, further study in additional monocot species is
required to fully understand how they evolved fleshy fruit ripening.
418
Ethylene is a stress hormone often involved in tissue senescence and is a gaseous molecule
that can easily diffuse. Both attributes make it ideal as the signal controlling fruit ripening in
climacteric fruits. However, much less is known about the regulatory mechanisms in non-
climacteric fruits. Study of additional non-climacteric cultivars and mutants and comparison
with their climacteric counterparts could provide a new perspective to reassess how non-
climacteric ripening is regulated. Many plant hormones have been implicated in fleshy fruit
ripening[23]. We do observe ripening-associated expression of the ABA biosynthesis genes,
particularly in the non-climacteric melon cultivar PI 161375 (**Supplementary Fig. 23**),
which is perhaps not surprising as ABA is also associated with stress and involved in
senescence.
429
The fruitENCODE project has generated a comprehensive dataset for eleven fleshy fruit
species, which opens the door for addressing some important problems in agricultural
practices. For example, post-harvest loss is a major concern for horticulture produce
worldwide, especially in developing countries and persists even in modern developed food
supply chains. However, improvement in shelf life through manipulation of ethylene often
leads to undesirable reduction in fruit quality, which is expected as the downstream ripening
genes affecting quality are tightly coupled to the autocatalytic ethylene loop. With a
comprehensive annotation of the *cis*-regulatory elements, and much improved understanding
of their ripening regulators, it is possible to engineer specific genes to combat post-harvest
losses while retaining or improving quality.

Acknowledgments: This work is supported by Hong Kong UGC GRF-14119814/14104515, Area of Excellence Scheme AoE/M-403/16 and Shenzhen Peacock-KQTD201101. Sequencing data have been deposited in the NCBI Sequence Read Archive under the accession number PRJNA381300.

Author contributions: S.Z. designed the research, P.L., N.Z., Y.C., B.Z., Y.P., J.F., G.J. N.Y. and J.Z. performed the experiments, S.Y., D.T., S.Z. analysed the data, D.G., Z.F., J.G. and S.Z. wrote the paper.

Competing financial interests statement: The authors declare no conflict of interest.

References

1. Coombe, B. G. The development of fleshy fruits. *Annu. Rev. Plant Physiol.* **27,** 207–228 (1976).

2. *Biochemistry of Fruit Ripening*. (Springer Netherlands, 1993). doi:10.1007/978-94-011-1584-1

3. Giovannoni, J., Nguyen, C., Ampofo, B., Zhong, S. & Fei, Z. The epigenome and transcriptional dynamics of fruit ripening. *Annu. Rev. Plant Biol.* **68,** 61–84 (2017).

4. McManus, M. T. *The plant hormone ethylene*. (Wiley-Blackwell, 2012).

5. Manning, K. *et al.* A naturally occurring epigenetic mutation in a gene encoding an SBP-box transcription factor inhibits tomato fruit ripening. *Nat. Genet.* **38,** 948–952 (2006).

6. Vrebalov, J. A MADS-box gene necessary for fruit ripening at the tomato ripening-inhibitor (rin) Locus. *Science* **296,** 343–346 (2002).

7. Zhong, S. *et al.* Single-base resolution methylomes of tomato fruit development reveal epigenome modifications associated with ripening. *Nat. Biotechnol.* **31,** 154–159 (2013).

8. McMurchie, E. J., McGlasson, W. B. & Eaks, I. L. Treatment of fruit with propylene gives information about the biogenesis of ethylene. *Nature* **237,** 235–236 (1972).

9. Liu, R. *et al.* A DEMETER-like DNA demethylase governs tomato fruit ripening. *Proc. Natl Acad. Sci. USA* **112,** 10804–10809 (2015).

10. The Tomato Genome Consortium. The tomato genome sequence provides insights into fleshy fruit evolution. *Nature* **485,** 635–641 (2012).

11. Pech, J. C., Bouzayen, M. & Latché, A. Climacteric fruit ripening: Ethylene-dependent and independent regulation of ripening pathways in melon fruit. *Plant Sci.* **175,** 114–120 (2008).

12. Paul, V., Pandey, R. & Srivastava, G. C. The fading distinctions between classical

patterns of ripening in climacteric and non-climacteric fruit and the ubiquity of ethylene—An overview. *J. Food Sci. Technol.* **49,** 1–21 (2012).

13. Golding, J. B., Shearer, D., Wyllie, S. G. & McGlasson, W. B. Application of 1-MCP and propylene to identify ethylene-dependent ripening processes in mature banana fruit. *Postharvest Biol. Technol.* **14,** 87–98 (1998).

14. Martel, C., Vrebalov, J., Tafelmeyer, P. & Giovannoni, J. J. The tomato MADS-box transcription factor RIPENING INHIBITOR interacts with promoters involved in numerous ripening processes in a COLORLESS NONRIPENING-dependent manner. *Plant Physiol.* **157,** 1568–1579 (2011).

15. Kunieda, T. *et al.* NAC family proteins NARS1/NAC2 and NARS2/NAM in the outer integument regulate embryogenesis in *Arabidopsis. Plant Cell* **20,** 2631–2642 (2008).

16. Guo, Y. & Gan, S. AtNAP, a NAC family transcription factor, has an important role in leaf senescence. *Plant J.* **46,** 601–612 (2006).

17. Ríos, P. *et al. ETHQV6.3* is involved in melon climacteric fruit ripening and is encoded by a NAC domain transcription factor. *Plant J.* **91,** 671–683 (2017).

18. Pirona, R. *et al.* Fine mapping and identification of a candidate gene for a major locus controlling maturity date in peach. *BMC Plant Biol.* **13,** 166 (2013).

19. D'Hont, A. *et al.* The banana (*Musa acuminata*) genome and the evolution of monocotyledonous plants. *Nature* **488,** 213–217 (2012).

20. Liang, C. *et al.* OsNAP connects abscisic acid and leaf senescence by fine-tuning abscisic acid biosynthesis and directly targeting senescence-associated genes in rice. *Proc. Natl Acad. Sci. USA* **111,** 10013–10008 (2014).

21. Choudhury, S. R, Roy, S., Nag, A., Singh, S. K. & Sengupta, D. N. D.N. Characterization of an AGAMOUS-like MADS box protein, a probable constituent of flowering and fruit ripening regulatory system in banana. *PLoS One* **7,** e44361 (2012).

22. Elitzur, T. *et al.* Banana MaMADS transcription factors are necessary for fruit ripening and molecular tools to promote shelf-life and food security. *Plant Physiol.* **171,** 380–91 (2016).

23. Kumar, R., Khurana, A. & Sharma, A. K. Role of plant hormones and their interplay in development and ripening of fleshy fruits. *J. Exp. Bot.* **65,** 4561–4575 (2013).

Figure Legends

Figure 1 | MADS positive feedback loop controlling tomato fruit ripening. a, Gene loci involved in fruit ripening with browser tracks showing different epigenetic features, transcription factor binding and gene expression. **b**, Model for tomato fruit ripening regulation. In nonfruit tissues and immature fruits, key ripening regulators are repressed by promoter DNA hypermethylation and H3K27me3. When the epigenetic marks are removed in the matured fruits, ripening is initiated with a burst of autocatalytic ethylene production, followed by the activation of the downstream ripening genes. **c**, Ectopic expression of *RIN*, *TAGL1* and *ACS2* generated autocatalytic ethylene in tobacco. Ethylene inhibitor treatment or deleting the EIN3 binding motif in the *RIN* promoter disrupted the loop and blocked ethylene synthesis.

Figure 2 | NAC positive feedback loop controlling peach fruit ripening. a, Key gene loci involved in peach ripening with browser tracks showing different epigenetic features, transcription factor binding and gene expression. **b**, Model for peach fruit ripening regulation. Similar to the tomato model, the peach ripening genes are also repressed by promoter DNA hypermethylation and H3K27me3. **c**, Ectopic expression of the peach *NAC* and *ACS* genes generated autocatalytic ethylene in tobacco. Ethylene inhibitor treatment or deleting the EIN3 binding motif in the *NAC* promoter disrupted the loop and blocked ethylene synthesis.

Figure 3 | Banana fruit ripening controlled by a dual-loop system. Different from other climacteric fruits, once the banana autocatalytic ethylene production starts, it becomes ethylene independent, and we found that banana has two positive feedback loops to control ripening. **a,** Genes involved in banana ripening with browser tracks showing different epigenetic features and gene expression. **b**, Dual-loop model for banana fruit ripening regulation. **c**, Ectopic expression of the *NAC* and *ACS* genes from the first loop generated autocatalytic ethylene in tobacco. Ethylene inhibitor treatment or deleting the EIN3 binding motif in the *NAC* promoter disrupted the loop and blocked ethylene synthesis. **d**, Ectopic expression of the *NAC*, *ACS*, and the three *MADS* genes generated autocatalytic ethylene that could not be blocked by ethylene inhibitor treatment. Deletion of the NAC motif in *MADS1* promoter or the MADS motif in the NAC gene promoter disrupted the second loop.

Figure 4 | Direct orthologs of the ripening genes in non-climacteric and dry fruit species. The direct orthologs of the eudicot climacteric fruit ripening *NAC* and *MADS* genes in the non-climacteric fruit watermelon (**a**) and dry fruit *Arabidopsis* (**b**) are also associated with tissue-specific H3K27me3 mark. **c**, The direct orthologs of the monocot banana fruit ripening *NAC* and *MADS* genes.

Figure 5 | Speciation, fruit ripening types and polyploidization in different angiosperm lineages. Confirmed whole-genome duplications and triplications are shown with circles. Species bearing dry, fleshy climacteric and fleshy non-climacteric fruits are indicated by the color black, red and green, respectively. The three climacteric fruit types are shown in brackets.

 Extended Data Figure Legends
565

Extended Data Figure 1 | MADS positive feedback loop controlling apple fruit
ripening. a, Gene loci involved in fruit ripening with browser tracks showing different
epigenetic features and gene expression. **b**, Model for apple fruit ripening regulation.
The ethylene biosynthesis genes are associated with H3K27me3 in both leaf and
immature fruit, while the *MADS1* (MD14G1215600) is associated with H3K27me3
in leaf. **c**, Ectopic expression of *MADS1* and *ACS* (MD15G1302200) under their
native promoter and *MADS2* (MD06G1204300) with the CaMV35S promoter
generated autocatalytic ethylene in tobacco. Ethylene inhibitor treatment or deleting
the EIN3 binding motif in the *MADS1* promoter disrupted the loop and blocked
ethylene synthesis.
576

Extended Data Figure 2 | MADS positive feedback loop controlling pear fruit
ripening. a, Gene loci involved in fruit ripening with browser tracks showing different
epigenetic features and gene expression. **b**, Model for pear fruit ripening regulation.
The ethylene biosynthesis and the *MADS1* (PCP020055.1) gene are associated
with H3K27me3 in leaf leaf. In the non-climacteric pear cultivar that could no
synthesize ripening ethylene, the *ACS* gene became hyper-H3K27me3. **c**, Ectopic
expression of *MADS1* and *ACS* (PCP011500.1) under their native promoter and
MADS2 (PCP042878.1) with the CaMV35S promoter generated autocatalytic
ethylene in tobacco. Ethylene inhibitor treatment or deleting the EIN3 binding motif in
the *MADS1* promoter disrupted the loop and blocked ethylene synthesis.
587

Extended Data Figure 3 | MADS-box transcription factors involved in
climacteric fruit ripening. Phylogenetic tree showing the SEP-clade MADS-box
transcription factors associated with climacteric fruit ripening and their direct
orthologs in non-climacteric and dry fruit plants. Their gene expression and
H3K27me3 levels in leaves, immature and ripe fruit tissues are shown. For rice
genes, the data from young leaves, senescence leaves and ovary tissues are used.
For *Arabidopsis*, young leaves, young carpel and senescence silique data is shown.
Red: climacteric fruit ripening genes; green: non-climacteric fruit; purple: dry fruit.
596

Extended Data Figure 4 | NAC transcription factors involved in climacteric fruit
ripening. Phylogenetic tree showing NAC transcription factors associated with
climacteric fruit ripening and their direct orthologs in non-climacteric and dry fruit
plants. Their gene expression and H3K27me3 levels in leaves, immature and ripe
fruit tissues are shown. For rice genes, the data from young leaves, senescence
leaves and ovary tissues are used. For *Arabidopsis*, young leaves, young carpel and
senescence silique data is shown. Red: climacteric fruit ripening genes; green: non-
climacteric fruit; purple: dry fruit.
605

Extended Data Figure 5 | NAC positive feedback loop controlling papaya fruit
ripening. a, Key gene loci involved in fruit ripening with browser tracks showing
different epigenetic features and gene expression. **b**, Model for papaya fruit ripening
regulation. The papaya ripening genes are also repressed by promoter DNA
hypermethylation and H3K27me3. **c**, Ectopic expression of the *NAC*
(evm.model.supercontig_80.94) and *ACS* (evm.model.supercontig_1322.1) genes
generated autocatalytic ethylene in tobacco. Ethylene inhibitor treatment or deleting

the EIN3 binding motif in the *NAC* promoter disrupted the loop and blocked ethylene
synthesis.
615
Extended Data Figure 6 | NAC positive feedback loop controlling melon fruit
ripening. a, Key gene loci involved in fruit ripening with browser tracks showing
different epigenetic features and gene expression. **b**, Model for papaya fruit ripening
regulation. Key ripening genes are also repressed by promoter DNA
hypermethylation and H3K27me3. In non-climacteric variety PI, the *NAC*
(MELO3C016540T1) promoter is hypermethyated, while in non-climacteric variety
PS, the *NAC*, *ACS* (MELO3C021182T1) and *ACO* (MELO3C014437T1) loci are
associated with hyper-H3K27me3 **c**, Ectopic expression of the *NAC* and *ACS* genes
generated autocatalytic ethylene in tobacco. Ethylene inhibitor treatment or deleting
the EIN3 binding motif in the *NAC* promoter disrupted the loop and blocked ethylene
synthesis.
627
628 **Extended Data Figure 7 | Direct orthologs of the ripening genes in cucumber.**
629 In non-climacteric fruit cucumber, the carpel senescence related *NAC* gene is also
630 associated with tissue-specific H3K27me3. Additional members in the *NAC*, *ACS*
631 and *ACO* families are shown in the supplementary figures.
632
633 **Extended Data Figure 8 | Direct orthologs of the ripening genes in non-**
634 **climacteric fruit grape.** The *NAC* and *MADS* genes are associated with tissue-
635 specific H3K27me3.
636
637 **Extended Data Figure 9 | Direct orthologs of the ripening genes in strawberry.**
638 The *NAC* genes is associated with tissue-specific H3K27me3.
639
640 **Extended Data Figure 10 | Direct orthologs of the ripening genes in**
641 **watermelon.** Its *NAC* and *MADS* genes are associated with tissue-specific
642 H3K27me3.
643

Figure 1 | MADS positive feedback loop controlling tomato fruit ripening.

Figure 2 | NAC positive feedback loop controlling peach fruit ripening.

Figure 3 | Banana fruit ripening controlled by a dual-loop system.

Figure 4 | Direct orthologs of the ripening genes in non-climacteric and dry fruit species.

Figure 5 | Speciation, fruit ripening types and polyploidization in different angiosperm lineages

Extended Data Figure 1 | MADS positive feedback loop controlling apple fruit ripening.

Extended Data Figure 2 | MADS positive feedback loop controlling pear fruit ripening.

Extended Data Figure 3 | MADS-box transcription factors involved in climacteric fruit ripening.

Extended Data Figure 4 | NAC transcription factors involved in climacteric fruit ripening.

Extended Data Figure 5 | NAC positive feedback loop controlling papaya fruit ripening.

Extended Data Figure 6 | NAC positive feedback loop controlling melon fruit ripening.

Extended Data Figure 7 | Direct orthologs of the ripening genes in cucumber.

Extended Data Figure 8 | Direct orthologs of the ripening genes in non-climacteric fruit grape.

Extended Data Figure 9 | Direct orthologs of the ripening genes in strawberry.

Extended Data Figure 10 | Direct orthologs of the ripening genes in watermelon.