A eDNA-qPCR combination assay to non-invasively detect the presence of the

parasite *Schistocephalus solidus* inside its intermediate host, the threespine

stickleback.

Chloé Suzanne Berger[1] and Nadia Aubin-Horth[1,*]

[1]Département de Biologie et Institut de Biologie Intégrative et des Systèmes (IBIS),

Université Laval, Québec, Canada, G1V 0A6

*corresponding author: Nadia.Aubin-Horth@bio.ulaval.ca

ABSTRACT

To study parasite-host interactions, it is of major importance to detect the presence of the parasite during its development in the host. The *Schistocephalus solidus*-threespine stickleback pair has been used extensively to study host phenotypic alterations caused by a parasite with a complex life cycle. This cestode worm is localized inside the abdominal cavity of its intermediate fish host, which limits its detection to the time when it has grown substantially or to multiple infections. Here, we present a non-invasive species-specific quantitative real time PCR approach based on the detection of environmental DNA from the worm (eDNA), sampled directly in the abdominal cavity of the threespine stickleback. Using this approach on two fish populations (n=151), 98% of fish were correctly assigned to their *S. solidus* infection status. Among the 35 sticklebacks that harboured single or multiple infections, only 3 were not detected as infected. Furthermore, non-infected sticklebacks were never incorrectly identified as infected. There was a significant correlation between eDNA concentration and the total mass of parasites. We also assessed ventilation rates as a rapid and complimentary mean to detect infection (n=20). Ventilation rates were repeatable over 3 measurements, did not vary significantly between non-infected and infected fish, were not significantly correlated to the parasite index, and thus could not reliably be used to predict infection. Our eDNA approach could be used in this parasite-host system alone or in combination with morphological measurements to detect *S. solidus*.

Key Words: *Schistocephalus solidus*, threespine stickleback, parasite, detection, quantitative real-time PCR, environmental DNA

INTRODUCTION

Internal parasites with complex life cycles often have multiple effects on their hosts (Marcogliese and Cone 1997). Studying these parasite-hosts interactions requires to be able to track the parasite during its development in the hosts. One of the model systems in the study of phenotypic alterations of the intermediate host infected by a parasite with a complex life cycle is the interaction between *Schistocephalus solidus* and its obligatory intermediate host, the threespine stickleback (*Gasterosteus aculeatus*, (Barber 2013)). *S. solidus* is a freshwater flatworm with a complex life cycle that has to infect three distinct hosts in order to survive and reproduce: a copepod, the threespine stickleback, and a fish-eating bird. The threespine stickleback is the specific intermediate host of *S. solidus* and can be infected by one or several worms after consumption of a *S. solidus*-infected copepod. Within 24 hours after oral uptake, the worm penetrates the intestine wall of the fish and eventually migrates into the abdominal cavity where it establishes as a small plerocercoid (Hammerschmidt and Kurtz 2007). During the first twelve weeks, *S. solidus* shows a sigmoid growth curve in the stickleback abdominal cavity (Barber and Svensson 2003). When the worm has gained a sufficient mass, it becomes infective and can reproduce in the intestinal system of its final host, a fish-eating bird (Tierney and Crompton 1992). Previous studies demonstrated that *S. solidus* infection has major impacts on the stickleback energy demands, physiology and immunity. Furthermore, effects of the macro parasite on its fish host include behavioural perturbations that result in risky behaviours, such as loss of anti-predatory responses and of schooling tendency (reviewed in (Barber and Scharsack 2010)).

The whole life cycle of this parasite can be reproduced in the laboratory (Barber and

Scharsack 2010), which makes it highly amenable to manipulation. However, the

success rate of infection in the laboratory when experimental single infections are used

is around 15-20% ((Grécias et al. 2016); (Weber et al. 2017)). Thus, a high number of

individuals must be included and assayed in a study to obtain an acceptable final

sample size of infected ones. Furthermore, variation in infection rates for sticklebacks

caught in the wild is high (Weber et al. 2017), which means that wild individuals

considered as uninfected controls may turn out to be infected when studying wild

sticklebacks. A simple detection method would allow selecting, or avoiding, infected

individuals depending on the needs of a specific project. This would allow minimizing the

number of individuals used in an experimental procedure, an important aspect for the

refinement of animal use protocols. However, despite the fact that the *S. solidus*-

threespine stickleback system is a very useful model in behaviour, ecology, evolution,

immunology and parasitology (Barber 2013), there is currently a lack of a simple, reliable

and non-invasive method to systematically determine if a threespine stickleback is

infected by *S. solidus*.

Available methods to detect *S. solidus* are based on morphological criteria. Indeed,

sticklebacks parasitized by *S. solidus* display modifications in the shape of their head

and exhibit a swollen and distended belly. These morphological modifications are

exacerbated with an increase in the parasitic load (Dingemanse et al. 2009). The

principal morphometric technique that is thereby currently used to determine if a

stickleback is infected by *S. solidus* relies on a visual analysis of the level of the

abdominal distension of the fish. This allows to predict the infection status (infected or

not) and the parasite index (PI, the proportion of infected fish weight that is contributed

by parasite tissue) (Barber 1997). However, this visual approach is less reliable when

the stickleback is infected by one worm or by a few small worms, which do not induce

clear morphological changes. Furthermore, some confounding factors can impact the

level of the abdominal distension. During the breading season, gravid females exhibit a

distended belly that is very similar to the one of fish infected by *S. solidus,* and a large

stomach or liver can also influence the abdominal distension (Barber 1997). X-rays are

another morphometric technique that can be used in parasitology to determine the

infection status of a host, but several problems are associated with their application,

such as the difficulty to distinguish between the parasite and the surrounding fish tissue

(Shahidi et al. 1997) or the high financial cost that limits their application to some

laboratories. In this context, it is necessary to develop a method that will not be based

on morphological criteria.

Our objective was to design a low cost and non-lethal method to detect *S. solidus* in the

abdominal cavity of its threespine stickleback host. To do so, we used two

complementary approaches: detection of environmental DNA (eDNA) and measurement

of a physiological trait. Our first hypothesis was that *S. solidus* produces cell-free DNA in

the abdominal cavity of the fish, and that this eDNA could be identified by a PCR

approach to correctly detect *S. solidus* infection in sticklebacks (Bass et al. 2015). eDNA

produced by parasites is a common target in epidemiology to detect infection using a

PCR approach in organic samples like serum, feces ((Pontes et al. 2002); (Xu et al.

2017)) or plasma (Imwong et al. 2015). eDNA is also used to detect invasive species in

aquatic environments (Takahara et al. 2013). We thus tested a non-invasive quantitative

real-time PCR method (qPCR), based on eDNA extraction from fluids of the abdominal

cavity of the stickleback. We used qPCR for its sensitivity and higher capacity of

detecting false positive compared to PCR products detected on an electrophoresis gel

(Bass et al. 2015). We designed primers that amplify a gene sequence from the *S.*

solidus genome but that also do not amplify DNA from the specific host, the threespine

stickleback. To do so, we chose a nuclear gene (GEEE01010589.1 TSA:

Schistocephalus solidus ssol_TR119934_c5_g1_i2 transcribed RNA sequence) that

does not have known homologous sequences in other species (Hébert et al. 2016),

allowing to specifically detect *S. solidus*. We tested if we could assign a fish to a given

infection status. Since eDNA has been previously used to make quantitative predictions

about the biomass of organism detected (Lacoursiere-Roussel et al. 2016), we also

tested if the eDNA concentration we measured covaried with the total mass of the

parasite(s) present within the abdominal cavity of an individual.

Our second hypothesis was that infected individuals would show a significantly higher

ventilation rate, which could be used to distinguish between non-infected and infected

fish. As mentioned previously, the energy demands imposed by the presence of *S.*

solidus have been well documented and affect growth (Barber and Svensson 2003) and

reproductive output (Heins and Baker 2003). For example, oxygen consumption is

higher in infected sticklebacks compared to uninfected ones for a given swimming speed

(Lester 1971) and for fish of similar mass (Meakins and Walkey 1975). Infected fish shift

to surface respiration at a higher dissolved oxygen level than uninfected ones, being

only rivalled by females harbouring mature eggs, which are also energy demanding

(Meakins and Walkey 1975). We thus tested the use of the ventilation rate of healthy

and infected sticklebacks as a rapid and non-invasive indicator of *S. solidus* infection,

using a previously published protocol to measure ventilation in threespine stickleback (Di

Poi et al. 2016).

MATERIALS AND METHODS

Populations of interest

This study was conducted in two populations of threespine sticklebacks. The first

population was Lac Témiscouata, Québec (47°40'33"N 68°50'15"O, freshwater

environment), where sticklebacks are known to be infected by *S. solidus* and to show

typical behaviour alterations (Grecias et al. 2017). The second population was

composed of anadromous fish from a salt marsh at Isle-Verte, Québec (48°1'0"N

69°26'59"O; salinity 22-26 ppt (Poulin and FitzGerald 1989)), for which infection by *S.*

solidus is not expected as the worm does not tolerate high levels of salinity (Macat et al.

2015).

Fish sampling and rearing

144   In Lac Témiscouata, adults were caught in July 2016 and September 2017, and

145   juveniles in August 2016, using a seine and minnow traps. Juvenile sticklebacks from

146   the anadromous Isle-Verte salt marsh population were sampled in tide pools in July

2016 using hand nets. Sticklebacks were brought to the "Laboratoire Regional des Sciences Aquatiques" at Université Laval. Adults and juveniles from each population were kept in separate 80 L-tanks under a Light:Dark photoperiod and a temperature of 15°C reflecting the conditions in their natural environments (Québec, Canada). For anadromous fish, salinity was maintained at 28 ppt. Fish were fed daily with brine shrimps. All sticklebacks were adults at the start of the experiment. They were isolated in 2 L-tanks and were assigned an identification number. 151 sticklebacks were used: 96 from Lac Témiscouata and 55 from Isle-Verte.

Sampling eDNA from the host abdominal cavity

We sampled the internal fluids of the fish within the intra-peritoneal cavity to obtain eDNA. The fish of interest was taken out of its tank and placed on a sponge. The needle of a syringe filled with 100 µL of PBS (Phosphate Buffered Saline, pH 7.4 Life Technologies) was inserted into its abdominal cavity and PBS was injected. Then, without removing the needle from the abdominal cavity, the plunger of the needle was pulled back until the syringe filled with the PBS that was just injected (up to 100 µL). After the needle was removed, the fish was directly put back in its tank. Between 10 and 100 µL of PBS was obtained and directly added to a tube of 700 µL of Longmire lysis preservation buffer (Longmire et al. 1997). This protocol was done twice for each fish on two consecutive days. On the second day, the PBS used to wash the abdominal cavity was added to the same Longmire tube as the first sample (i.e. up to 200 µL in the 700 µL Longmire tube). Each day, the needle was inserted into a different face of the fish. Abdominal cavity fluid samples were placed at 4°C between the first and the second

day, and during the night following the second day. Samples were kept at -20°C before

DNA extraction. Fish were fed with brine shrimps after the sampling of the abdominal

cavity.

After these two samplings, fish were euthanized with an overdose of MS-222 (75 mg/L

mg/kg) followed by exsanguination and dissected to determine if they were infected by

S. solidus. Fish sex, size and mass, and *S. solidus* mass and number in each fish were

noted. The parasite index was calculated using the formula (total weight (mg) of

parasites / Total weight (mg) of host plus parasites) x 100. A value of 50 indicates a

situation where the total weight of parasites is equal to the net weight of the stickleback

(Arme and Owen 1967).

The protocol was approved by the Comité de Protection des Animaux de l'Université

Laval (CPAUL 2014069-3).

DNA extraction from samples

DNA extraction from the abdominal cavity samples was performed according to an

environmental DNA method developed by Lacoursière-Roussel et al. (Lacoursière-

Roussel et al. under revision), with small modifications to the protocol. To summarize, 30

µL of proteinase K (4 mg/mL) (VWR) was directly added to a Longmire buffer sample

and the tube was incubated at 55°C for the night. The sample was transferred into a

new 2 mL tube and 950 µL of phenol chloroform isoamyl alcohol (Invitrogen) was added.

After shaking and centrifugation, supernatant was transferred into a new tube and 950

µL of chloroform (VWR) was added. After shaking and centrifugation, a maximum of 750

µL of the supernatant was put into a new tube. 750 µL of ice cold isopropanol (Fisher

Scientific) and 375 µL of room temperature 5M sodium chloride NaCl were added to the

tube, which was placed at -20°C for the night. After centrifugation, all liquid was

removed from the tube and 1500 µL of 70% ethanol was added. After centrifugation,

ethanol was removed and the tube was air-dried. The DNA pellet was resuspended in

80 µL water and the tube was placed at 55°C for 10 minutes. The tube was placed at

4°C for the night and then stored at -20°C.

Design of primers specific to the *Schistocephalus solidus* genome

A pair of primers that amplifies a sequence from a nuclear gene (GEEE01010589.1

TSA: Schistocephalus solidus ssol_TR119934_c5_g1_i2 transcribed RNA sequence)

from the *S. solidus* genome was designed. The gene sequence was obtained from the

de novo transcriptome of *S. solidus* (sequence available on NCBI:

https://www.ncbi.nlm.nih.gov/nuccore/GEEE01010589.1?report=fasta) and does not

have known homologues in other species (Hébert et al. 2016). This nuclear gene, whose

function is unknown, is highly expressed during early developmental stages of the worm

when it is not able to reproduce (Hebert et al. 2017). We targeted the DNA sequence of

this gene with the forward for-CGGATTGTCTTCTCGTTGTA and reverse rev-

GGACAACCACTGTCCACTAA primers, designed with Primer3 (Untergasser et al.

2012). The primers generated a 200 bp amplicon. We verified with the Basic Local

Alignment Search Tool (nucleotide BLAST) on NCBI that the 200 bp amplicon did not

align with other sequences except for *S. solidus*.

PCR assays to determine specificity of amplification

Polymerase Chain Reaction (PCR) assays were conducted on genomic DNA samples to

confirm that the designed primers amplified only *S. solidus* DNA and not threespine

stickleback DNA. DNA was extracted from *S. solidus* and stickleback samples (Lac

Témiscouata) kept at -20°C in ethanol 90%. Tissue samples of 0.3 cm^2 (*S. solidus* body,

stickleback tail) were used for the extraction. Extraction was performed using a salt

method (Aljanabi and Martinez 1997). PCR reaction was performed in a total volume of

24.3 µL (13.75 µL of nuclease free water, 1 µL of 10mM dNTPs, 0.25 µL of TAQ DNA

polymerase (5U/µL Bio Basic), 2.5 µL of 10x TAQ buffer with MgCL$_2$, 5 µL of DNA (*S.*

solidus or stickleback) and 1.8 µL of primer mix (4.56 µM of each primer)). Two negative

controls were done (no primers, no DNA). The PCR program was realized with a

Mastercycler (Eppendorf) under these conditions: a denaturation step of 2 min at 95°C,

an annealing and elongation step of 40 cycles including 20 sec at 95°C, 40 sec at 56°C

and 60 sec at 72°C, and a final step of 10 min at 72°C. PCR products were visualized on

a 2% agarose gel using SYBR safe staining (Invitrogen).

qPCR assays

eDNA extracted from the abdominal cavity samples of fish was amplified with a real-time

quantitative polymerase chain reaction (qPCR) using the designed primers specific to *S.*

solidus. Each 96-well plate included eDNA samples extracted from the body cavity and a

standard curve obtained from a pool of *S. solidus* genomic DNA (3 individuals) diluted at

different concentrations. The concentrations of the 7-point standard curve ranged from

0.9 ng/µL to 0.01 ng/µL with a dilution factor of 2. qPCR reaction was performed in a

qPCR mix including 12.5 µL of PowerUp SYBR Green Master mix (Life Technologies)

and 1 µL of primer mix (4.56 µM of each primer). For the standard curve, genomic DNA

samples, 6.5 µL of RNAse free water and 5 µL of DNA were added to the qPCR mix. For

the eDNA samples, no water was used and 11.5 µL of solution containing the

resuspended eDNA was added to the qPCR mix (for a total qPCR mix volume of 25 µL

in each case). Two negative controls (no primers, no DNA) were done in each plate.

Each reaction was done in triplicate. qPCR was performed with a 7500 Real-Time PCR

system (Life Technologies). The amplification was realized according to the conditions

described by (Lacoursière-Roussel et al. 2016): 2 min at 50 °C, 10 min at 95 °C,

followed by 70 cycles of 15 sec at 95 °C and 60 sec at 56 °C. A melt curve protocol was

then performed with temperature going from 60 celsius to 95 celsius to detect primer

dimers, check specificity and obtain a Tm value for that sample, which can be compared

to the standard curve of Schistocephalus DNA.

Ventilation rate measurements

A subset of individuals used in the *S. solidus* eDNA detection experiment was used for

this analysis (captured in Lac Témiscouata in July 2016, n=20). Ventilation

measurements were repeated three days in a row with individuals sampled in the same

order each day. For each individual fish, a 250 mL beaker was filled with 50 mL of clean

water from the supply used to fill experimental 2 L-tanks. The individual was filmed from

the side (JVC Everio camera) to be able to detect opercular movement. At then end of

the three days of measurements, these fish were euthanized and their infection status

determined. Dissection showed that 13 individuals were infected and 7 were parasite-

free. All the infected fish were also detected as infected by the eDNA approach (see

below). Films were subsequently analyzed at 0.46 X speed by an observer (NAH) blind

to treatment using the open source VLC Media Player (Videolan). The first 30 seconds

of the movie where the fish was clearly visible were used to measure ventilation rates. If

the individual moved in a position where it was not visible during the first-30-second

sampling period, the ventilation measurement was made for the usable seconds within

that period and reported back to 30 seconds. The observer repeated all measurements

for the first day of sampling to assess accuracy. Measures done twice on the same fish

on the same video were highly correlated (Spearman r = 0.99) with most samples

having the exact same number of ventilations when measured twice, with a maximum

difference of 3 ventilation movements and an average difference of 0.7. A single

measurement per video was thus used. A ventilation rate was calculated by dividing the

number of ventilation measured in 30 seconds by 30. This measurement was used for

further statistical analyses. The total number of ventilation measured during a 30 second

interval was used to estimate the number of ventilation over 1 minute for graphical

purposes and for comparison to previous studies.

Analyses

Following qPCR amplification, the presence or the absence of *S. solidus* eDNA in the
body cavity samples was determined by comparing the Ct values (amplification curves)
and the Tm values (melt curves) of the samples with the ones of the standard curves. A
Ct ("cycle threshold") is the cycle at which a sample reaches a preset threshold of
fluorescence, which is the proxy for the quantity of eDNA. A sample with a low Ct means
that there were more templates in the original sample and that it thus reached the
threshold sooner in the 70 PCR cycles. A body cavity sample showing at least one of its
Ct values (among the triplicates) above 0 in combination with the associated Tm
value(s) close (+/- 2) to the mean Tm of the standard curve (79.64) was considered
positive for *S. solidus* eDNA. On the contrary, if a Ct value above 0 was associated with
a Tm value more distant to the mean Tm of the standard curve, then this value was
considered negative for *S. solidus* eDNA and the result of primer dimer formation.
Finally, if the body cavity sample had only "undetermined" Ct values, it was considered
negative for *S. solidus* eDNA, irrespectively of Tm values. For each fish, qPCR results
were analyzed with regard to its infection status (parasitized or not by *S. solidus* upon
dissection). A true positive was obtained when there was concordance between the
infection status and the qPCR results. A false negative was defined when the fish was
infected by *S. solidus*, but no traces of *S. solidus* eDNA were found using qPCR. A false
positive was associated with fish not infected by *S. solidus* but for which *S. solidus*
eDNA was inferred using qPCR results. All individual results are found in supplementary
data.

Statistical analysis was done in R version 3.3.3 (R Core Team 2017) using R Studio

(Version 1.0.136) and the ggplot2 package (Wickham 2009). To determine if our method

allows a quantitative estimate of parasite mass, we tested if the eDNA concentration

obtained for infected fish (using the Ct value) negatively correlated with their total

parasite mass, using a non-parametric Spearman correlation (a small Ct value should be

associated with a larger parasite mass and vice-versa if the eDNA method is

quantitative). We used the lowest Ct measured for a given fish if more than one triplicate

amplified DNA. Repeatability of ventilation rates over the three days was calculated

following (Lessells and Boag 1987). We compared ventilation rates between infected

and non-infected individuals using a t-test, as they followed a normal distribution (based

on a Shapiro-wilk test). We tested if ventilation rates covaried positively with the parasite

load (parasite index) using a non-parametric correlation (Spearman correlation).

RESULTS

Primers specificity

PCR tests performed with primers that target a sequence from the *S. solidus* nuclear gene NCBI ID GEEE01010589.1 (TSA: Schistocephalus solidus ssol_TR119934_c5_g1_i2 transcribed RNA sequence) confirm the amplification of *S. solidus* genomic DNA, and no amplification of threespine stickleback genomic DNA. The PCR product was sequenced using Sanger ABI3730xl (Plateforme de séquençage et de génotypage des génomes, Centre Hospital Universitaire du Québec) to confirm the amplification of the 200 pb amplicon in the *S. solidus* genome. Controls (no template, no primers) were negative as expected.

Detection of *S. solidus* DNA in the abdominal cavity of infected fish with qPCR

qPCR was performed in combination with a E-DNA extraction protocol to try to detect *S. solidus* presence in the abdominal cavity of sticklebacks. Standard curves realized with a pool of *S. solidus* genomic DNA samples diluted at different concentrations had mean Ct values ranging from 29.65 (highest concentration) to 36.86 (lowest concentration) and a mean Tm value of 79.64. All the controls were negative as expected. The earliest Ct value we obtained for abdominal cavity E-DNA samples with our protocol and amount of starting material varied between 34.4 and 49.6 for positively identified individuals. Using this approach, 148 fish out of 151 were correctly assigned to their infection status (infected or not by *S. solidus*, 98 % of true positive) (Table 1). The 3 remaining fish were false negative. We did not obtain any false positive with our approach.

In the Témiscouata population, 35 fish out of 96 were infected by *S. solidus.* Single and multiple infections (up to 6 worms) were reported upon dissection, with a median of 1.5 worm, illustrating the predominance of single or double infections. Note that one fish harboured 12 worms, but we did not obtain reliable mass for the worms and as such is included in the dataset only to calculate the infection status assignation rate. Among the infected sticklebacks, 18 exhibited a single worm upon dissection and no external visual clues suggested the presence of *S. solidus* in the fish body cavity (Figure 1). For all infected fish, the total parasite mass was comprised between 130 mg and 2356 mg and the parasite index was always below 50 (between 4.9 and 40.4), meaning that the total weight of parasites was always below the net weight of the host fish (Table 2). Thirty-two out of the 35 infected fish were successfully detected as infected using qPCR. For the 3 remaining infected fish, no trace of *S. solidus* eDNA was detected with our protocol (1.98% of samples tested, low false negative rate) (Table 1). The 3 fish had a total parasite mass that span almost the whole range found in the study (168 mg and 274 mg for the single infections and 1644 mg for the double infection, from the second smallest to the third largest). Their parasite index (respectively 12.5, 6.5 and 25.3) was above the minimal values from infected fish that were correctly detected (Table 2). They had no eDNA detected in any of the triplicates (Ct "undetermined"). As predicted, none of the fish from the Isle-Verte anadromous population were infected by *S. solidus* (n=55). We never detected *S. solidus* DNA in the samples of the uninfected fish from this population, nor from uninfected fish from Lac Temiscouata (0 false positive) (Table 1). There was a significant correlation between Ct values (which are inversely correlated to the amount

of *S. solidus* eDNA) and the total parasite mass found within a fish (one-sided Spearman

correlation, rho= -0.41, p = 0.01, figure 2).

Variations of ventilation rates between uninfected and infected sticklebacks

We measured the ventilation rates of 20 fish, 3 times over 3 consecutive days, and then

determined that 13 of these were infected. The infected individuals harboured from 1 to

5 worms with a total mass ranging between 618 mg and 2356 mg (parasite index

between 13.2 and 40.4). One infected individual died after the first ventilation

measurement and we could use the videos for only the 2 first days for one non-infected

fish. We had 3 ventilation measurements for all other individuals. Repeatability of

ventilation across 3 days for a given fish was 0.41 (Lessells and Boag 1987). As

ventilation rates were repeatable over this time scale, we used ventilation rates

measured on the first day in subsequent analysis, to reflect how this method would be

used to quickly detect infection. Ventilation rates estimated over 30 seconds and

reported in number of opercular movement per minute ranged between 99 and 146

opercular beats/min in un-infected fish and between 112 and 160 in *S. solidus* infected

ones (figure 3). The average number of ventilation per minute was 122 for uninfected

fish and 134 for infected fish. We found that ventilation rates did not vary significantly

between infected and non-infected individuals (two-sided t-test, t = -1.57, p = 0.15, n=20,

figure 3). Ventilation was positively correlated to the parasite index but this covariation

was not significant (one-sided Spearman correlation, rho = 0.35, p = 0.07, n=20). There

was a substantial overlap in ventilation rate between the two groups, which prevented

the use of this trait to discriminate between *S. solidus* infected and healthy individuals.

DISCUSSION

In parasite-host system studies, being able to detect the presence of a parasite during its development in the host is of major importance to better understand the interactions between the two organisms. Here, we demonstrate the power of a method based on detecting eDNA from the parasite *S. solidus* directly sampled from the abdominal cavity of its specific intermediate host, the threespine stickleback.

Using our eDNA-qPCR combination method in two populations of threespine sticklebacks, 148 fish out of 151 were correctly assigned to their infection status (infected or not), including 32 fish parasitized by *S. solidus* (with single and multiple infections). For 3 sticklebacks only (1.98% of samples tested), *S. solidus* eDNA was not detected despite the fact that these fish were infected (low false negative rate). We never assigned an "infected" status to an uninfected fish (0 false positive). The amount of S. solidus eDNA was correlated to the total mass of parasites found within a fish. The ventilation rate was repeatable when measured over 3 days, but had very low discriminatory power.

Environmental DNA has recently emerged as a promising tool to detect aquatic species that are sometimes at very low densities in past and present heterogeneous environments (Bass et al. 2015). Detecting eDNA therefore finds application in ecology and conservation ((Rees et al. 2014); (Thomsen and Willerslev 2015)). eDNA is also

combined with PCR approaches to detect pathogens in water (trematode *Ribeiroia ondatrae*, pathogen of amphibians: (Huver et al. 2015); liver fluke: (Hashizume et al. 2017)). The originality of our method to detect an internal parasite lies in the fact that the eDNA extraction was made from the internal fluids of the fish abdominal cavity instead of water samples, as *S. solidus* is free to move in the abdominal cavity of its host, potentially releasing cell-free DNA. We detected *S. solidus* eDNA in almost all infected fish (98% of true positive). We found only 3 false negatives out of 35 infected sticklebacks (1.98% of tested samples). In a preliminary test, we first tried to amplify *S. solidus* eDNA from body cavity samples or from water samples by standard PCR and to visualize the amplification products on an electrophoresis gel. For the body cavity samples, the PCR approach was less sensitive compared to the qPCR method, with a much higher false negative rate that made this simpler method unusable. For the water samples from tanks of infected fish, *S. solidus* DNA was never detected by PCR (although we could detect a stickleback positive control). We propose that concentrations of *S. solidus* DNA are too low or absent to be detected in the excretions of the fish.

All sticklebacks from the anadromous population (n=55) and 61 fish from the Témiscouata population were free of infection by *S. solidus*. Our method did not yield false positive, i.e. non-infected fish that would be detected as infected with our qPCR results. False positive could have resulted from a lack of specificity of the primers used. Thus, the absence of false positive for 116 samples confirms the specificity of the method. A false positive could also have resulted from a signal of an exposed

stickleback, i.e. a fish that was exposed to a *S. solidus*-infected copepod, but for which

infection did not succeed and the worm did not develop in its abdominal cavity. The

absence of false positive in our data suggests that our method does not allow to detect

exposed sticklebacks. However, all the fish used were of wild origin, which prevents us

to have a record of exposure over the life of an individual fish such that we do not know

if any of our non-infected fish were exposed to this parasite. Testing this possibility of

laboratory-exposed fish could add to the usefulness of the present detection method.

The 3 false negatives (infected sticklebacks from the Témiscouata population for which

S. solidus DNA was not detected using qPCR) had a total parasite mass and a parasite

index above the minimal values from infected fish that were correctly detected. This

observation indicates that these false negatives are not due to a lack of sensitivity of the

method at low parasitic weight. We rather propose that the critical step of our method is

during the sampling of the internal fluids of the fish intraperitoneal cavity. As mentioned

previously, *S. solidus* is free in the abdominal cavity of its host. When the PBS washing

is done, it is possible that the worm may be too far from the needle (e.g. between two

organs of the stickleback) to allow the capture of its DNA. To overcome this issue, we

performed body cavity sampling on two consecutive days, and each time the needle was

inserted into a different face of the fish. This approach was quite reliable and probably

resulted in our low false negative rate. However, we suggest that the eDNA sampling

protocol could be extended over three days in order to maximize *S. solidus* DNA

capture. The correlation we found between the levels of *S. solidus* amplified eDNA (Ct

values) and the parasitic load suggest that in the future this approach could be used in a

quantitative way to estimate not only the presence but the total mass of parasites within

a host. The correlation is non linear and there is large variation in total parasite mass for

a given Ct value and further refinement would be needed. Therefore, our method in its

current state gives a very reliable presence/absence response and its use for

quantitative assessment looks promising for the future.

Ventilation rates of un-infected fish were in the same range as found in previous studies

of freshwater sticklebacks (Bell et al. 2010, Di Poi et al. 2016). The ventilation rate was

repeatable when measured over 3 days, which suggest that a single measurement gives

sufficient information on ventilation requirements of each individual at this life stage and

that there are significant inter-individual differences in physiology. However, ventilation

rates did not differ between infected and non-infected sticklebacks, although the lowest

and highest rates were found in un-infected and infected fish respectively and there was

a trend for ventilation rates to covary positively with the parasite index. Therefore, the

discriminatory power of this simple measurement was low and could only be used as a

complementary indicator of infection by *S. solidus* in its fish host.

We expect the eDNA method presented here to find applications beyond the *S. solidus*-

threespine stickleback system. Indeed, a lot of fish species, including fish of economic

importance, have parasites found in their abdominal cavity (Hoffman 1999). To apply our

method to these systems, the main challenge would be to design highly specific primers

for the parasite of interest (Taylor et al. 2013). Our low cost method could be a simple

alternative to morphological-based methods already used in these systems.

Acknowledgments

The authors would like to thank the personnel of the Laboratoire Régional des Sciences

Aquatiques (LARSA) at Université Laval for their help with fish rearing. We thank Anaïs

Lacoursière-Roussel, Noémie Leduc, Alysse Perreault-Payette and Guillaume Côté for

their advice concerning eDNA extraction. We thank Christian Landry and Leonard Foster

for their comments during the development of this project. Anna Prudova from the Foster

lab first suggested to us to sample the stickleback abdominal cavity fluids to study the

Schistocephalus secretome, which led to this eDNA detection protocol.

Authors' contributions

C.S.B. designed the study with input from N.A.-H. C.S.B. performed the eDNA-qPCR

experiments and analyzed the data. N.A.-H. performed the ventilation rate

measurements and did the statistical analyses. C.S.B. and N.A-H. wrote the manuscript.

Ethics

This study was done in compliance with Animal Care and Use Program and was

approved by the Comité de Protection des Animaux de l'Université Laval (CPAUL

2014069-3).

Data availability

496   The complete dataset is available as a supplementary data file.

Competing interests

The authors declare no competing or financial interests.

REFERENCES

Aljanabi, S. M., and I. Martinez. 1997. Universal and rapid salt-extraction of high quality genomic DNA for
PCR-based techniques. Nucleic acids research **25**:4692-4693.
Arme, C., and R. W. Owen. 1967. Infections of the three-spined stickleback, Gasterosteus aculeatus L.,
with the plerocercoid larvae of Schistocephalus solidus (Müller, 1776), with special reference to
pathological effects. Parasitology **57**:301-314.
Barber, I. 1997. A non-invasive morphometric technique for estimating cestode plerocercoid burden in
small freshwater fish. JOURNAL OF FISH BIOLOGY **51**:654-658.
Barber, I. 2013. Sticklebacks as model hosts in ecological and evolutionary parasitology. Trends in
Parasitology **29**:556-566.
Barber, I., and J. P. Scharsack. 2010. The three-spined stickleback- Schistocephalus solidus system: an
experimental model for investigating host-parasite interactions in fish. Parasitology **137**:411-424.
Barber, I., and P. A. Svensson. 2003. Effects of experimental Schistocephalus solidus infections on
growth, morphology and sexual development of female three-spined sticklebacks, Gasterosteus
aculeatus. Parasitology **126**:359.
Bass, D., G. D. Stentiford, D. T. J. Littlewood, and H. Hartikainen. 2015. Diverse Applications of
Environmental DNA Methods in Parasitology. Trends in Parasitology **31**:499-513.
Bell, A. M., L. Henderson, and F. A. Huntingford. 2010. Behavioral and respiratory responses to stressors
in multiple populations of three-spined sticklebacks that differ in predation pressure. J Comp
Physiol B **180**:211-220.
Di Poi, C., J. Lacasse, S. M. Rogers, and N. Aubin-Horth. 2016. Evolution of stress reactivity in stickleback.
Evolutionary Ecology Research **17**:395-405.
Dingemanse, N. J., C. Oosterhof, F. Van Der Plas, and I. Barber. 2009. Variation in stickleback head
morphology associated with parasite infection. Biological Journal of the Linnean Society **96**:759-
768.
Grécias, L., F.-O. Hébert, C. S. Berger, I. Barber, and N. Aubin-Horth. 2016. Can the behaviour of
threespine stickleback parasitized with Schistocephalus solidus be replicated by manipulating
host physiology? The Journal of experimental biology **220**:237-246.
Grecias, L., J. Valentin, and N. Aubin-Horth. 2017. Testing the parasite mass burden effect on host
behaviour alteration in the Schistocephalus-stickleback system. bioRxiv.
Hammerschmidt, K., and J. Kurtz. 2007. Schistocephalus solidus: establishment of tapeworms in
sticklebacks--fast food or fast lane? Experimental Parasitology **116**:142-149.
Hashizume, H., M. Sato, M. O. Sato, S. Ikeda, T. Yoonuan, S. Sanguankiat, T. Pongvongsa, K. Moji, and T.
Minamoto. 2017. Application of environmental DNA analysis for the detection of Opisthorchis
viverrini DNA in water samples. Acta Tropica **169**:1-7.
Hebert, F. O., S. Grambauer, I. Barber, C. R. Landry, and N. Aubin-Horth. 2017. Major host transitions are
modulated through transcriptome-wide reprogramming events in Schistocephalus solidus, a
threespine stickleback parasite. Molecular Ecology **26**:1118-1130.
Hébert, F. O., S. Grambauer, I. Barber, C. R. Landry, and N. Aubin-Horth. 2016. Transcriptome sequences
spanning key developmental states as a resource for the study of the cestode Schistocephalus
solidus, a threespine stickleback parasite. GigaScience **5**:1-9.
Heins, D. C., and J. A. Baker. 2003. Reduction of Egg Size in Natural Populations of Threespine Stickleback
Infected with a Cestode Macroparasite. The Journal of Parasitology **89**:1-6.
Hoffman, G. L. 1999. Parasites of North American freshwater fishes. Cornell University Press.

Huver, J. R., J. Koprivnikar, P. T. J. Johnson, and S. Whyard. 2015. Development and application of an
eDNA method to detect and quantify a pathogenic parasite in aquatic ecosystems. Ecological
Applications **25**:991-1002.
Imwong, M., C. J. Woodrow, I. C. E. Hendriksen, J. Veenemans, H. Verhoef, M. A. Faiz, S. Mohanty, S.
Mishra, G. Mtove, S. Gesase, A. Seni, K. D. Chhaganlal, N. P. J. Day, A. M. Dondorp, and N. J.
White. 2015. Plasma concentration of parasite DNA as a measure of disease severity in
falciparum malaria. The Journal of infectious diseases **211**:1128.
Lacoursière-Roussel, A., K. Howland, E. Normandeau, E. Grey, P. Archambault, K. Deiner, D. M. Lodge, C.
Hernandez, N. Leduc, and L. Bernatchez. under revision. eDNA monitoring as a new surveillance
tool for coastal Arctic biodiversity. Molecular Ecology Resources.
Lacoursiere-Roussel, A., M. Rosabal, and L. Bernatchez. 2016. Estimating fish abundance and biomass
from eDNA concentrations: variability among capture methods and environmental conditions.
Mol Ecol Resour **16**:1401-1414.
Lacoursière-Roussel, A., M. Rosabal, and L. Bernatchez. 2016. Estimating fish abundance and biomass
from eDNA concentrations: variability among capture methods and environmental conditions.
Molecular Ecology Resources **16**:1401-1414.
Lessells, C. M., and P. T. Boag. 1987. Unrepeatable repeatabilities: a common mistake. The Auk **104**:116-
121.
Lester, R. J. G. 1971. The influence of Schistocephalus plerocercoids on the respiration of Gasterosteus
and a possible resulting effect on the behavior of the fish. Canadian Journal of Zoology **49**:361-
366.
Longmire, J. L., M. Maltbie, and R. J. Baker. 1997. Use of "Lysis Buffer" in DNA Isolation and Its
Implication for Museum Collections. OCCASIONAL PAPERS, MUSEUM OF TEXAS TECH
UNIVERSITY **163**:1-3.
Macat, Z., A. Bednarik, and M. Rulik. 2015. The three-spined stickleback (Gasterosteus aculeatus)
infection with Schistocephalus solidus in Hel marina (Puck Bay, Baltic Sea, Poland).
OCEANOLOGICAL AND HYDROBIOLOGICAL STUDIES **44**:11-17.
Marcogliese, D. J., and D. K. Cone. 1997. Food webs: a plea for parasites. Trends in Ecology & Evolution
12:320-325.
Meakins, R. H., and M. Walkey. 1975. The effects of parasitism by the plerocercoid of Schistocephalus
solidus Muller 1776 (Pseudophyllidea) on the respiration of the three-spined stickleback
Gasterosteus aculeatus L. JOURNAL OF FISH BIOLOGY **7**:817-824.
Pontes, L. A., E. Dias-Neto, and A. Rabello. 2002. Detection by polymerase chain reaction of Schistosoma
mansoni DNA in human serum and feces. The American journal of tropical medicine and hygiene
66:157.
Poulin, R., and G. J. FitzGerald. 1989. Risk of parasitism and microhabitat selection in juvenile
sticklebacks. Canadian Journal of Zoology **67**:14-18.
R Core Team. 2017. R: A language and environment for statistical computing. . R Foundation for
Statistical Computing, Vienna, Austria.
Rees, H. C., B. C. Maddison, D. J. Middleditch, J. R. Patmore, and K. C. Gough. 2014. The detection of
aquatic animal species using environmental DNA–a review of eDNA as a survey tool in ecology.
Journal of Applied Ecology **51**:1450-1459.
Shahidi, F., Y. M. Jones, and D. Kitts. 1997. Seafood safety, processing, and biotechnology. CRC Press.
Takahara, T., T. Minamoto, and H. Doi. 2013. Using environmental DNA to estimate the distribution of an
invasive fish species in ponds. PLoS ONE **8**:e56584.
Taylor, M. W., S. M. Kevin, K. Y. Michael, F. J. Stephen, H. L. Winsor, R. W. Andrew, and K. S. Michael.
2013. Robust detection of rare species using environmental DNA: the importance of primer
specificity. PLoS ONE **8**:e59520.

Thomsen, P. F., and E. Willerslev. 2015. Environmental DNA – An emerging tool in conservation for
monitoring past and present biodiversity. Biological Conservation **183**:4-18.
Tierney, J. F., and D. W. T. Crompton. 1992. Infectivity of Plerocercoids of Schistocephalus solidus
(Cestoda: Ligulidae) and Fecundity of the Adults in an Experimental Definitive Host, Gallus gallus.
The Journal of Parasitology **78**:1049-1054.
Untergasser, A., I. Cutcutache, T. Koressaar, J. Ye, B. C. Faircloth, M. Remm, and S. G. Rozen. 2012.
Primer3--new capabilities and interfaces. Nucleic acids research **40**:e115.
Weber, J. N., M. Kalbe, K. C. Shim, N. I. Erin, N. C. Steinel, L. Ma, and D. I. Bolnick. 2017. Resist Globally,
Infect Locally: A Transcontinental Test of Adaptation by Stickleback and Their Tapeworm
Parasite. The American naturalist **189**:43.
Wickham, H. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
Xu, J., Z. L. Duan, Z. X. Guan, Y. Y. Wang, C. Lin, T. T. Zhang, H. Q. Zhang, X. Qian, and C. M. Xia. 2017.
Early detection of circulating DNA of Schistosoma japonicum in sentinel mice models.
Experimental Parasitology **176**:82-88.

Table 1. Fish distribution according to their infection status at dissection (infected

or uninfected by *S. solidus*) and their qPCR results (*S. solidus* eDNA detected or

not). Numbers of fish from the Témiscouata population are indicated in normal font and

the ones from the anadromous Isle-Verte population are indicated in italic font.

	Status at dissection: infected	Status at dissection: uninfected
qPCR= detected as infected (*S. solidus* eDNA)	32; *0*	0; *0*
qPCR= detected as uninfected (no *S. solidus* eDNA)	3; *0*	61; *55*

615 **Table 2. Infection by *S. solidus* in the Témiscouata population.** Minimum, maximum

616 and median values for number of parasites, total parasite mass and parasite index.

617 Infected fish ID T12 was not included in calculations of min, max, median (no specific

618 data available other than known presence of worms).

number of infected fish	number of parasites per fish			total parasite mass (mg)			parasite index (PI, %)		
	min	max	median	min	max	median	min	max	median
35	1	6	1.5	130	2356	495	4.9	40.4	16.6

Figures

Figure 1. External morphology of threespine sticklebacks from Lac Témiscouata that were infected by *S. solidus* (single or multiple infections) or not. A. Non-infected fish. **B.** Infected fish with 1 *S. solidus* worm (parasite mass 618.2 mg; PI 13.2). **C.** Infected fish with 5 *S. solidus* worms (total parasite mass 2356 mg; PI 39.3). A and B do not exhibit abdominal distension while abdominal distension is clearly visible in C. Fish B and C were both detected as infected using the E-DNA method.

Figure 2. eDNA concentrations covary with total parasite mass. Low Ct values, which represent high levels of eDNA, are associated with a high parasite mass (Spearman correlation, rho = -0.41, p = 0.01, n= 31).

Figure 3. Ventilation rates do not differ significantly between un-infected and infected threespine sticklebacks (two-sided t-test, t = -1.57, p = 0.15, n=20). Ventilation per minute ranged between 99 and 146 opercular movement per minute in un-infected fish (n=7) and between 112 and 160 in *S. solidus* infected fish (n=13). The lower and upper hinges correspond to the 25th and 75th percentiles and the vertical bars extend to the largest value no further than 1.5 times the interquartile range.

Figure 1

Figure 2

Figure 3