1

Microhabitats shape diversity-productivity

relationships in freshwater bacterial communities

Marian L. Schmidt[1,2#], Bopaiah A. Biddanda[3], Anthony D. Weinke[3], Edna Chiang[1,4],

Fallon Januska[3], Ruben Props[1,5] & Vincent J. Denef[1]

#Corresponding author: marschmi@umich.edu

[1]Department of Ecology & Evolutionary Biology, University of Michigan, Ann Arbor, MI USA;

[2]Current Address: Department of Integrative Biology, University of Texas, Austin, TX USA;

[3]Annis Water Resources Institute, Grand Valley State University, Muskegon, MI, USA;

[4]Current Address: Department of Bacteriology, University of Wisconsin, Madison, WI, USA;

[5]Center for Microbial Ecology and Technology (CMET), Department of Biochemical and

Microbial Technology, Ghent University, Gent, Belgium

Abstract

Eukaryotic communities commonly display a positive relationship between biodiversity and ecosystem function (BEF) but the results have been mixed when assessed in bacterial communities. Habitat heterogeneity, a factor in eukaryotic BEFs, may explain these variable observations but it has not been thoroughly evaluated in bacterial communities. Here, we examined the impact of habitat on the relationship between diversity assessed based on richness, evenness, or phylogenetic diversity, and heterotrophic productivity. We sampled co-occurring free-living (more homogenous) and particle-associated (more heterogeneous) bacterial habitats in a freshwater, estuarine lake. Diversity measures, and not environmental variables, were the best predictors of particle-associated heterotrophic production. There was a strong, positive, linear relationship between particle-associated bacterial richness and heterotrophic productivity that strengthened with evenness. There were no observable BEF trends in free-living bacterial communities. Across both habitats, communities with more phylogenetically related taxa had higher per-capita heterotrophic production than communities of phylogenetically distantly related taxa. Our findings show that heterotrophic bacterial productivity is positively correlated with evenness and richness, negatively with phylogenetic diversity, and that BEF relationships are contingent on microhabitats. Our work adds to the understanding of the highly distinct contributions to community diversity and ecosystem functioning contributed by bacteria in free-living and particle-associated aquatic habitats.

Keywords: diversity-productivity, biodiversity-ecosystem function, bacterial communities, microhabitats, particle-associated, limnology, heterotrophic productivity

Introduction

Our planet is currently experiencing an extreme species extinction event (Thomas et al., 2004; Wake & Vredenburg, 2008, Ferrier et al., 2016). Concern about such declines in biodiversity has resulted in hundreds of studies evaluating the relationship between biodiversity and ecosystem function (BEF), with an even larger biodiversity effect in natural ecosystems compared to controlled experiments (Duffy et al., 2017). BEF relationships focusing on the number of species (*i.e.* species richness) as the biodiversity value are generally positive and asymptotic and thus biodiversity loss causes a small change in ecosystem function at first and then, at some tipping point, a dramatic decrease in function (Cardinale et al., 2012; Tilman et al., 2014). While the focus of local and global species loss is typically on eukaryotic organisms, bacterial species numbers have also been found to be decreasing at local scales within the human gut (Blaser, 2014) and terrestrial ecosystems (Singh et al., 2014). Of particular concern is the loss of the number of bacterial guilds responsible for key geochemical transformations, such as methane oxidation (Levine et al., 2011) that controls rates of methane emissions.

An extrapolation from eukaryotic relationships would predict there to be no richness-ecosystem function relationships for bacterial communities because they are generally composed of an order of magnitude more taxa than the communities in most eukaryotic BEF studies. Several studies have indicated no relationship between species richness with broad functional processes, such as heterotrophic respiration or biomass production, that are performed by many taxa (see figure 5 in Levine et al., 2011; Langenheder et al., 2006; Delgado-Baquerizo et al., 2016). Yet, other studies on broad processes such as denitrification (Philippot et al., 2013) and on narrow metabolic processes that are catalyzed by few bacterial taxa, such as methanotrophy (Levine et al., 2011), and the degradation of triclosan and microcystin (Delgado-Baquerizo et al.,

2016) did find evidence of bacterial richness and ecosystem function relationships. In addition to

the number of species, there are several other components of biodiversity including functional

trait or gene richness (*i.e.* abundance-unweighted) (Reich et al., 2004; Flynn et al., 2011; Evans

et al., 2017), species or functional dominance or evenness (*i.e.* abundance-weighted) (Wilsey &

Potvin, 2000; Wilsey & Polley, 2004; Kirwan et al., 2007; Wittebolle et al., 2009), and

phylogenetic diversity metrics (Flynn et al., 2011). Plant community evenness has been shown to

impact plant productivity more than richness (Wilsey & Potvin, 2000; Wilsey & Polley, 2004;

Kirwan et al., 2007) and the initial evenness of microbial microcosms has been found to be a key

factor in functional stability, even under the selective stressors of temperature and salt stress

(Wittebolle et al., 2009)

Phylogenetic relatedness, while having received less attention than richness and

evenness, may also influence BEF relationships. Indeed, some studies show relationships across

different ecosystems between phylogenetic diversity and ecosystem functions (Cadotte et al.,

2008; Jiang et al., 2010). However, research with freshwater green algae (Fritschie et al., 2014;

Venail et al., 2014) did not find this relationship. A more recent study found the opposite result

by showing that closely related green algal species had weaker competition and more facilitation

than distantly related species, thus resulting in higher productivity (Narwani et al., 2017). While

relationships between phylogenetic relatedness among community members and ecosystem

function have been assessed in bacterial systems (Tan et al., 2012; Galand et al., 2015; Roger et

al., 2016), much work has focused on low-diversity, experimentally-assembled communities

with bacteria that can be grown in culture. It is worthwhile to expand these findings to

communities with richness levels typically found in natural communities.

The nature of BEF relationships and the mechanism(s) that underpins them may depend

on habitat structure or heterogeneity. Increasing habitat heterogeneity has been found to enhance

the strength of BEF relationships (Tylianakis et al. 2008), presumably due to a greater role for

niche complementarity effects in heterogeneous environments (Cardinale 2011). While habitat

heterogeneity contributes to increased diversity within bacterial populations and communities

(Zhou et al., 2008; Shade et al., 2008), the influence of habitat heterogeneity on BEF

relationships remains unknown for bacterial systems.

In this study, we hypothesized that bacterial diversity would be positively correlated with

bacterial heterotrophic production, and that this relationship would be stronger in more

heterogeneous environments. We simultaneously surveyed free-living and particle-associated

surface water bacterial communities. These habitats have been extensively studied for their

ability to sustain distinct bacterial communities and ecosystem processes (Crump et al., 1999;

Bižić-Ionescu et al., 2014; Mohit et al., 2014; Schmidt et al., 2016). In addition, studies on model

aggregates had 3-fold higher bacterial protein production and two orders of magnitude higher

protease activity (Grossart et al., 2007), indicating particles can be an important place for

microbial activity in aquatic ecosystems. Particulate matter comprises a variety of types and sizes

of particles with some particles also harboring physicochemical gradients (Simon et al., 2002),

and hence represents a more heterogeneous habitat than the surrounding water. We tested BEF

relationships using a variety of diversity metrics including observed richness, evenness, and

phylogenetic diversity. We focused on heterotrophic bacterial production as our measure of

ecosystem function, as it is a key process affecting freshwater bacterial growth that in turn fuels

the eukaryotic food web (Cotner & Biddanda, 2002).

Methods

Lake sampling and sample processing

Surface water samples were collected at 1 meter depth from 4 long-term sampling stations

(Steinman et al., 2008) in mesotrophic Muskegon Lake (**Figure S1**), which is a freshwater

estuarine lake connecting the Muskegon River and Lake Michigan. These stations included the

mouth of the Muskegon River (43.250133,-86.2557), the channel to Bear Lake (43.238717,-

86.299283; a hypereutrophic lake), channel to Lake Michigan (43.2333,-86.3229; oligotrophic

lake), and the deepest basin of Muskegon Lake (43.223917,-86.2972; max depth = 24 m).

Samples were collected during the morning to early afternoon of 3 days in 2015 (May 12,

July 21, & September 30) aboard the R/V W.G. *Jackson*. All water samples were collected with

vertical Van Dorn samplers. Additionally, a vertical profile of temperature (T), pH, specific

conductivity (SPC), oxidation-reduction potential (ORP), chlorophyll (Chl*a*), total dissolved

solids (TDS), and dissolved oxygen (DO) was constructed at each station to characterize the

water column using a calibrated YSI 6600 V2-4 multiparameter water quality sonde (Yellow

Springs Instruments Inc.). Total Kjeldahl nitrogen (TKN), ammonia (NH_3), total phosphorus

(TP), and alkalinity (Alk) were processed from whole water while nitrate (NO_3), phosphate

(PO_4), and chloride (Cl^-) were hand filtered using a 60 mL syringe fitted with Sweeny filter

holder with a 13 mm diameter 0.45 µm pore size nitrocellulose filters (Millipore) and were

determined by standard wet chemistry methods in the laboratory (EPA, 1993).

Bacterial abundance by epifluorescence microscopy

Lake surface water samples were processed within 2-6 hours of their collection for determination

of heterotrophic bacterial abundance. Unfiltered lake water samples (5 mL) were preserved with

2% formalin and 1 mL subsamples were stained with acridine orange stain and filtered onto

black 25 mm 0.2 µm pore size polycarbonate filters (Millipore) at a maximum pressure of 0.1

Bar or 1.5 PSI. Prepared slides were stored frozen until manual enumeration by standard

epifluorescence microscopy at 1000x magnification under blue light excitation (Hobbie et al.

1977). Bacteria within the field of view (100 µm x 100 µm) that were not associated with any

particles were counted as free-living bacteria, whereas bacteria that were on particles were

counted as particle-associated. Sample filtration may bias counts due to free-living or particle-

associated cells being hidden on the underside of particles, free-living bacteria settling on top of

particles, or particle-associated cells dislodging. In the absence of any quantitative studies that

have rigorously addressed this issue, we have assumed the net effect of these opposing

methodological biases to be negligible in the present study.

Heterotrophic bacterial production measurements

Community-wide heterotrophic bacterial production was measured using [$_3$H] leucine

incorporation into bacterial protein in the dark (Kirchman et al. 1985; Simon and Azam, 1989).

At the end of the incubation with [$_3$H]-leucine, cold trichloroacetic acid-extracted samples were

filtered onto 3 µm filters that represented the leucine incorporation by particle-associated

bacteria (>3.0 µm). Each filtrate was collected and filtered onto 0.2 µm filters and the activity

therein represented incorporation of leucine by free-living bacteria (>0.2 µm-<3 µm). Measured

leucine incorporation during the incubation was converted to bacterial carbon production rate

using a standard theoretical conversion factor of 2.3 kg C per mole of leucine (Simon and Azam,

1989). Per-capita heterotrophic production was estimated by dividing heterotrophic production

by the cell counts measured in each fraction.

Preservation of bacterial filters in the field

Microbial biomass for the particle-associated (> 3 μm) and the free-living (3–0.22 μm) bacterial

fraction was collected by sequential in-line filtration on 3 μm isopore polycarbonate (TSTP, 47

mm diameter, Millipore, Billerica, MA, USA) and 0.22 μm Express Plus polyethersulfone

membrane filters (47 mm diameter, Millipore, MA, USA). We used 47 mm polycarbonate in-line

filter holders (Pall Corporation, Ann Arbor, MI, USA) and an E/S portable peristaltic pump with

an easy-load L/S pump head (Masterflex®, Cole Parmer Instrument Company, Vernon Hills, IL,

USA). The total volume filtered varied from 0.8–2.2 L with a maximum filtration time of 16

minutes per sample. Filters were submerged in RNAlater (Ambion) in 2 mL cryovials, frozen in

liquid nitrogen and transferred to a −80°C freezer until DNA extraction.

DNA extraction, sequencing and processing

DNA extractions were performed using an optimized method based on the AllPrep

DNA/RNA/miRNA Universal kit (Qiagen; McCarthy et al., 2015; details in supplementary

methods). Extracted DNA was sequenced using Illumina MiSeq V2 chemistry 2×250 (500

cycles) of dual index-labelled primers that targeted the V4 hypervariable region of the 16S rRNA

gene (515F/806R) (Caporaso et al., 2012; Kozich et al., 2013) at the Microbial Systems

Laboratories at the University of Michigan Medical School in July 2016. RTA V1.17.28 and

MCS V2.2.0 software were used to generate data. Fastq files were submitted to NCBI sequence

read archive under BioProject accession number PRJNA412984. We analyzed the sequence data

using MOTHUR V.1.38.0 (seed = 777; Schloss et al., 2009) based on the MiSeq standard

operating procedure accessed on 3 November 2015 and modified with time (see data

accessibility and supplemental methods). For classification of operational taxonomic units

(OTUs), a combination of the Silva Database (release 123; Quast et al., 2013) and the freshwater

TaxAss 16S rRNA database and pipeline (Rohwer et al., 2018, accessed August 18, 2016). All

non-bacterial and chloroplast sequences were pruned out of the dataset and replicate samples

were merged by summing sample sequencing read counts using the *merge_samples* function

(phyloseq). A batch script for our protocol can be found in this project's GitHub page in

https://github.com/DenefLab/Diversity_Productivity/blob/master/data/mothur/mothur.batch.taxa

ss.

Estimating Diversity

To get the best estimate of each diversity metric, each sample was subsampled to 6,664

sequences (the smallest library size) with replacement and were averaged over 100 trials.

Observed richness, Shannon entropy, and inverse Simpson's index were calculated using the

diversity function within the vegan (Oksanen et al., 2013) R package via the *estimate_richness*

function in the phyloseq (McMurdie and Holmes, 2013) R package. Simpson's Evenness was

calculated by dividing the inverse Simpson's index by the observed richness (Magurran, 2004).

To calculate phylogenetic diversity, we first removed OTUs that had a count of 2 sequences or

less throughout the entire dataset, as these are more prone to be artefacts originating from

sequencing errors or the OTU clustering algorithm. Representative sequences of each of the

1,891 remaining OTUs were collected from the aligned fasta file produced within mothur, and

header names in the mothur output fasta file were modified using bbmap (Bushnell, 2016) to

only include the OTU name. A phylogenetic tree was created with FastTree using the GTR+CAT

(general time reversible) model (Price et al., 2010). Mismatches between the species community

data matrix and the phylogenetic tree were checked with the *match.phylo.comm* command

(picante). Finally, both abundance-unweighted and -weighted phylogenetic diversity was

estimated using specifications described in the next paragraph with the picante R package.

The most common phylogenetic diversity (PD) measure is Faith's PD (Faith, 1992),

however, this metric is very strongly correlated with species richness (**Figure S2**). Instead, the

mean pairwise phylogenetic distance (or MPD) was calculated (*ses.mpd* function in the *Picante*

R package (Kembell et al., 2010), null.model = "independentswap"). The MPD measures the

average phylogenetic distance between all combinations of two taxa pulled from the observed

community and compares it with a null community of equal richness pulled from the gamma

diversity of all the samples (*see supplemental methods for more details*). Values higher than zero

indicate phylogenetic evenness or overdispersion (higher phylogenetic diversity) while values

less than zero indicate phylogenetic clustering (lower diversity) or that species are more closely

related than expected according to the null community (Kembel, 2009). Thus, this phylogenetic

metric is relative. From here, we will refer to the SES$_{MPD}$ as the "phylogenetic diversity" for

simplicity and clarity.

Statistical analysis

Further analysis of sequence data was performed in R version 3.4.2 (R Core Team 2017;

see supplemental methods for more details). To test which variable(s) were the best predictors of

community and per-capita heterotrophic production, we performed variable selection via a lasso

regression (using the *glmnet* R package, alpha = 1, and lambda.1se as the tuning parameter

(Friedman et al., 2010)) on all of the environmental, biodiversity, and principal component

variables. To further validate the lasso regression results, we performed ordinary least squares

(OLS) regressions on all variables, including the principal components (PCA) of the euclidean

distances of the environmental data. We used the Akaike information criterion (AIC) (accessed

with the *broom::glance()* command) to select the best performing OLS regression model.

Data and code availability

Original fastq files can be found on the NCBI sequence read archive under BioProject accession

number PRJNA412984. Processed data and code can be found on the GitHub page for this

project at https://deneflab.github.io/Diversity_Productivity/ with the main analysis at

https://deneflab.github.io/Diversity_Productivity/Final_Analysis.html

Results

Free-living communities had more cells but particle-associated communities had higher per-

capita heterotrophic production

We observed an order of magnitude more cells per milliliter ($p = 1 \times 10^{-6}$, Figure 1A) and

~2.5 times more community-wide heterotrophic production in the free-living fraction ($p = 0.024$,

Figure 1B). However, when calculated per-capita, particle-associated bacteria were on average

an order of magnitude more productive than free-living bacteria ($p = 7 \times 10^{-5}$, Figure 1C).

Particle-associated and free-living cell abundances in samples taken from the same water sample

did not correlate (Figure S3A). Heterotrophic production between corresponding free-living and

particle-associated fractions from the same water sample were positively correlated for both

community (Adjusted $R^2 = 0.40$, $p = 0.017$; Figure S3B) and per-capita production rates

(Adjusted $R^2 = 0.60$, $p = 0.003$; Figure S3C).

Particle-associated communities were more diverse in terms of observed richness and Shannon Entropy while free-living communities were more phylogenetically diverse

Across all samples, particle-associated bacterial communities were more diverse than free-living communities when considering richness and Shannon entropy (Figures 2A & S4A), but similar in the inverse Simpson's index and Simpson's evenness (Figure 2B & S4B).

Particle-associated bacterial community richness was always higher than in free-living communities and was maintained across the four sampling stations in the lake (Figure S5A). Particle-associated samples at the river and Bear Lake stations were on average more OTU-rich than the outlet to Lake Michigan and the Deep stations. Additionally, the river station had almost twice the inverse Simpson's value as compared with all other lake stations (Mean inverse Simpson Indices: Outlet = 23.6; Deep = 23.7; Bear = 35.3; River = 59.1; Figure S5A).

Particle-associated communities were more phylogenetically clustered than free-living communities based on unweighted phylogenetic diversity ($p = 0.01$, Figure 3A). Compared to other particle-associated samples, the outlet station that connects to oligotrophic Lake Michigan had a much larger unweighted phylogenetic diversity, indicating phylogenetic overdispersion (Figure S5A). Nevertheless, no sample across the entire dataset differed significantly from the null model with a significance threshold p-value of 0.05. There was no difference between weighted phylogenetic diversity in particle-associated versus free-living communities (Figure S5A).

Diversity-Productivity relationships were mostly observed in particle-associated communities

We analyzed BEF relationships for both community and per-capita production due to the distinct patterns of these two measures of heterotrophic production (Figure 1). There was a

strong, positive, linear BEF relationship between community-wide (Figures 2C-D & S4C-D) and

per-capita (Figures 2E-F & S4E-F) heterotrophic production and all richness and evenness

diversity metrics in the particle-associated communities, while no BEF relationships were

observed for the free-living communities. The inverse Simpson's index explained the most

amount of variation in community-wide heterotrophic production (Figure 2D; Adjusted R^2 =

0.69, p = 5 x 10^{-4}) and per-capita (Figure 2F; Adjusted R^2 = 0.69, p = 0.001). When the two data

points with the highest inverse Simpson's index and heterotrophic production were removed

from the regression (Figure 2D), the relationship was still significant and (Adjusted R^2 = 0.37; p

= 0.036), though not with richness (Adjusted R^2 = 0.12; p = 0.17). These results are also robust

across a range of minimum OTU abundance filtering thresholds (see *Sensitivity Analysis of Rare*

Taxa in the supplemental methods and Figure S6) and hold up for all threshold levels in inverse

Simpson and for richness until removal of OTUs observed 25 times (community-wide

heterotrophic production) and 15 times (per-capita heterotrophic production).

When the particle-associated and free-living samples were combined together into one

linear model to test an overall relationship between diversity and community-wide productivity,

there was no relationship (richness: p = 0.86; Shannon: p = 0.99; inverse Simpson: p = 0.36),

with the exception of a weak correlation for Simpson's Evenness (Adjusted R^2 = 0.12, p =

0.054). This further highlights the distinct BEF relationships across habitats. However, when

particle-associated and free-living samples were combined together into one linear model to test

an overall relationship between diversity and per-capita productivity, there was a strong

relationship with observed richness (Adjusted R^2 = 0.63, p = 3 x 10^{-6}), which broke down as

evenness was weighed more (Figure S7: Shannon: Adjusted R^2 = 0.52, 6 x 10^{-5}; Inverse

Simpson: Adjusted $R_2 = 0.48$, p = 2 x 10-4; Simpson's Evenness: p = 0.48). Thus, the relationship

between diversity and per-capita heterotrophic production was independent of habitat.

Phylogenetic diversity correlated with per-capita heterotrophic production but not with

community-wide production

Abundance-weighted phylogenetic diversity was not correlated with community or per-

capita heterotrophic production (Figure S8C - S8D) and therefore no further analyses were

performed with this diversity metric.

For unweighted phylogenetic diversity, we first determined the relationship between

community richness and phylogenetic diversity. There was a moderate, negative, linear

relationship when particle-associated and free-living samples were combined together into one

linear model to test an overall relationship between unweighted phylogenetic diversity and

observed richness (Figure 3B; Adjusted $R_2 = 0.35$, p = 0.001). To further validate this trend,

randomized communities were generated with an equal richness as the samples but with OTUs

randomly picked across the dataset. The unweighted phylogenetic diversity was then calculated

and regressed against each the randomized richness and there was no relationship (Figure S9;

Adjusted $R_2 = -0.02$, p = 0.44), verifying the negative relationship in the actual samples. When

particle-associated and free-living samples were individually run in separate linear models to test

for habitat-specific relationships between unweighted phylogenetic diversity and observed

richness, no trend was found in either particle-associated or free-living models (Figure 3B;

Particle: Adjusted $R_2 = 0.14$, p = 0.12; Free = Adjusted $R_2 = -0.10$, p = 0.97).

We then assessed the relationship between phylogenetic diversity and heterotrophic

productivity. Particle-associated and free-living phylogenetic diversities did not have individual

effects on community-wide or per-capita heterotrophic production. When data from both habitats

were combined into a single regression, there was no correlation between phylogenetic diversity

and community-wide heterotrophic production (Figure 3C). However, a negative correlation was

found when particle-associated and free-living samples were combined into one linear model to

test an overall relationship between unweighted phylogenetic diversity and per-capita

heterotrophic production (Figure 3D; Adjusted $R_2 = 0.42$, $p = 5 \times 10_{-4}$). Therefore, these two

results in combination indicated that communities composed of more phylogenetically similar

OTUs had a higher per-capita heterotrophic production rate.

Diversity, and not environmental variation, was the best predictor of particle-associated

heterotrophic production

311   To identify variables that best predicted community-wide and per-capita heterotrophic

production (*i.e.* remove variables that were correlated with each other and/or uninformative

variables), we performed lasso regression with all samples and individually with particle-

associated and free-living samples. For prediction of community-wide heterotrophic production,

only the inverse Simpson's index was selected for particle-associated samples whereas pH and

PC5 were selected for free-living samples, and no variables were selected when all samples were

included in the lasso regression. In contrast, for per-capita heterotrophic production, temperature

and the inverse Simpson's index were selected for particle-associated samples whereas pH was

the only predictor for free-living samples, and observed richness was the only predictor for all

samples (plotted in Figure S7A). Therefore, the best model for particle-associated microhabitats

always included inverse the inverse Simpson's index whereas free-living samples only included

environmental variables, such as pH.

323   To further verify that there were no confounding impacts of seasonal and environmental

variables on community-wide and per-capita heterotrophic production, we performed ordinary

least square (OLS) regressions and a dimension-reduction analysis of the environmental

variables through a principal components analysis (Table S1 & S2; Figure S10). Specifically, the

first 2 environmental axes explained ~70% of the environmental variation in the sampling sites

(Figure S10). Next, we predicted community-wide and per-capita heterotrophic production with

all environmental variables and the first six principal components as predictor variables with

individual particle-associated and free-living samples, and combined (*i.e.* all samples) models

(Table S1 & S2). The best single predictor of community-wide heterotrophic production was

inverse Simpson for particle-associated samples (AIC = 74.34; Adjusted R_2 = 0.69), pH for the

free-living samples (AIC =98.43; Adjusted R_2 = 0.49, p = 0.006), and pH for all samples (AIC =

192.16; Adjusted R_2 = 0.35) (Table S1). Whereas, the best single predictor of per-capita

heterotrophic production was inverse Simpson for particle-associated samples (AIC = 8.29;

Adjusted R_2 = 0.69), pH for the free-living samples (AIC = -2.39; Adjusted R_2 = 0.78), and

observed richness for all samples (AIC = 24.72; Adjusted R_2 = 0.63) (Table S2). Thus, the OLS

regressions are in agreement with the lasso regressions.

Discussion

We examined bacterial biodiversity-ecosystem function (BEF) relationships in relation to

two microhabitats within freshwater lakes: particulate matter and the surrounding water. First,

we found that community-wide and per-capita heterotrophic productivity of particle-associated

but not free-living bacterial communities showed a positive, linear BEF relationship with both

richness and evenness contributing. Second, particle-associated heterotrophic production was

better explained by diversity (*i.e.* inverse Simpson's index) than by environmental parameters.

Third, across both particle-associated and free-living communities, higher richness was

associated with lower phylogenetic diversity which, in turn, was associated with higher per-

capita heterotrophic bacterial production but not associated with community-wide heterotrophic

production.

Microbes have a large diversity of metabolisms and the choice of which to focus on may

inherently affect the BEF relationship. Indeed, "narrow" metabolic processes that are catalyzed

by a small subset of taxa within bacterial communities, such as some nitrogen and sulfur cycling,

have been found to display BEF relationships (Levine et al., 2011; Delgado-Baquerizo et al.,

2016). In contrast, for "broad" processes that are performed by the majority of taxa within a

bacterial community, such as heterotrophic production (*i.e.* focus of the present study) and

respiration, functional redundancy appears to weaken or remove the presence of BEF

relationships (Griffiths et al., 2000; Langenheder et al., 2006; Wertz et al., 2006; Levine et al.,

2011; Peter et al., 2011, Galand et al, 2015). These findings are in line with the absence of a BEF

relationship for free-living bacterial communities in our study.

However, the above results and hypotheses surrounding narrow and broad processes are

in conflict with the strong BEF relationship we observed in particle-associated bacterial

communities. As such, our study signifies that microhabitats or habitat heterogeneity can

influence bacterial BEF relationships, in agreement with previous research in eukaryotic systems

across a variety of ecosystems (Tylianakis et al., 2008; Cardinale 2011; Zeppilli et al., 2016). A

study using controlled stream mesocosms by Cardinale (2011) found that niche complementarity

effects are particularly important in more heterogeneous environments. In more heterogeneous

streams, algal populations used different nutrients and avoided direct competition for resources,

resulting in unique species occupying distinct and local microhabitats (Cardinale 2011). An

experimental study by Gravel et al. (2011) showed that BEFs depend on the legacy of previous

evolutionary events. Specifically, they found that after several hundred generations of evolution

on a variety of carbon substrates, generalist bacteria were more productive because of their more

efficient exploitation of the environmental heterogeneity. Finally, a recent study on freshwater

lake bacterial communities found a positive correlation between OTU evenness and the number

of dissolved organic matter (DOM) components (though the study did not evaluate ecosystem

function), suggesting that DOM resource heterogeneity may increase the diversity of bacterial

communities by creating equity among bacterial species (Muscarella et al., 2019).

Our observational study could not directly test the role of niche complementarity effects.

However, support for niche complementarity alone or in combination with species selection as

the mechanism underlying the BEF relationship in particle-associated habitats is provided by the

inverse Simpson's index being the strongest predictor of community-wide heterotrophic

production. As the inverse Simpson's index represents a measure of species dominance, it is

strongly affected by the evenness of abundant species. Communities that are more even have an

increased likelihood for complementary species to neighbor each other.

In our study, there are several reasons why heterogeneity of particulate matter may allow

for niche complementarity effects to occur and result in BEF relationships. First, particles have a

two-fold layer of heterogeneity as they (A) may be composed of different substrates such as

organic matter from terrestrial or aquatic environments and either heterotrophically or

photosynthetically derived (Grossart, 2010), and (B) each particle may comprise

physicochemical gradients as well (Simon et al., 2002). Second, microbial interactions are more

likely to occur between cells aggregated on particles as the interaction distances are usually

much shorter (Cordero & Datta, 2016) compared to free-living bacterial cells. In fact, genes

mediating social interactions, such as motility, adhesion, cell-to-cell transfer, antibiotic

resistance, mobile element activity, and transposases, have been found to be more abundant in

marine particles than compared to the surrounding water (Ganesh et al., 2014). Third, a recent

study by Ebrahimi et al. (bioRxiv) found that dense patches of bacterial cells on model marine

chitin particles promoted cross-feeding of oligosaccharides when particles were recalcitrant.

397   The importance of niche complementarity in microbial communities can also be deduced

from recent findings in the field of microbiology, which have shown widespread metabolic

interdependence among bacterial community members. First, a 2016 study that reconstructed

2,540 draft genomes of microbes found that most bacteria specialize in one particular step in

sulfur and nitrogen pathways and "hand-off" their metabolic byproducts to nearby organisms

(Anantharaman et al., 2016). It is likely that metabolic hand-offs, a specific form of bacterial

facilitation, will occur more in particle-associated compared to free-living communities. Indeed,

Datta et al.'s (2016) work on model marine chitin particles found that taxa that are incapable of

breaking down particles and instead rely on carbon produced by primary degraders thrive in later

phases of particle degradation, a repeatable result for three other polysaccharide substrates (Enke

et al., 2019). Second, Lilja and Johnson (2016) demonstrated that different microbial cell types

eliminate inter-enzyme competition by cross feeding, which increases substrate consumption by

allowing intracellular resources to go towards a single enzyme, rather than having two enzymes

that perform two separate reactions compete for nutrients within a cell. Third, some bacteria are

unable to grow in laboratory cultures unless they are in co-culture with other organisms, which

may be due to metabolic hand-offs or growth factors such as siderophores or catalases (Stewart,

2012).

414   Considering that (i) closely related taxa share more genes and metabolic pathways than

distantly related bacterial taxa (Konstantinidis & Tiedje, 2005; Kim et al., 2014) and (ii) bacteria

commonly have incomplete metabolic pathways, it may be possible that closely related bacteria

are most likely to exchange their metabolic byproducts. This may be why we found that new taxa

added to the community represented taxonomic clades similar to or already present in the

community, and that these communities with lower phylogenetic diversity (relative to expected)

had higher productivities. This result agrees with a recent study using freshwater algae and

vascular plants that found that co-cultures of more similar species were more productive

(Narwani et al., 2017). Similarly, a study of bacterial communities inhabiting Mediterranean

soils showed that plots containing more recently diverged lineages had higher ecosystem

function levels than when more distantly related lineages were present (Goberna & Verdú,

2018). However, other bacteria-focused studies found higher levels of antagonism with more

closely related taxa (Russel et al., 2017) and more bacterial productivity (measured through

colony forming units per mL) with more distantly related taxa (Venail and Vives, 2013).

Though, both of these studies were performed in the lab with r-selected (*i.e.* copiotrophic)

species grown in stable, warm, aerobic, agar plate conditions. Thus, Venail and Vives (2013) and

Russel et al. (2017) inherently break up potential interdependent relationship between bacteria

either by creating artificial communities or evaluating pairwise interactions and remove the

natural effect of spatial heterogeneity, environmental fluctuations, and the rest of the bacterial

community.

Previous studies on bacterial BEF relationships have used three approaches to manipulate

bacterial diversity (Krause et al., 2014): (1) removal of taxa (*e.g.* dilution to extinction) in which

complex communities are simplified (Franklin et al, 2001; Wertz et al., 2006; Peter et al., 2011;

Philippot et al., 2013; see Roger et al., 2016 for a review of this approach; fragmentation or

knockout in Bell, 2019), (2) addition of taxa (*e.g.* manually assembled communities) in culture

(Tan et al., 2012; Salles et al., 2009; invasion or coalescence in Bell, 2019), or (3) natural or

manipulated environmental communities (Griffiths et al., 2000; Levine et al., 2011; Galand et al.,

2015; Rivett & Bell, 2018). We took the third approach in this study. In contrast to the other two

approaches, this had the benefit of (a) maintaining high diversity with both abundant and rare

taxa, (b) including both r- and k-selected organisms, (c) allowing natural environmental and

ecological forcings to shape the community, and (d) evaluating BEF relationships in diversity

and productivity ranges that reflect natural communities. Admittedly, three inherent weaknesses

to our approach were that (a) we cannot measure all the potential variables that influence

heterotrophic productivity, which are especially and inherently difficult to measure within

particles, (b) we only have 24 samples for a 12 versus 12 study, and (c) our analysis is

correlational and we cannot manipulate the system to unequivocally separate causes and

consequences of bacterial production. For example, strong correlations with heterotrophic

production and pH in the free-living samples (Table S1 & S2) may point to pH being a

consequence of rather than a cause of varying production levels. This is because bacterial

production and bacterial respiration are positively correlated (del Giorgio & Cole, 1998) and

with increased respiration, pH may decrease due to CO_2 dissolution into the water.

Finally, we acknowledge that the typical sampling of bacterial communities and analysis

using DNA sequencing reflects *all* bacteria present in the community and not necessarily only

the *active* members of the community contributing to a given ecosystem function. In freshwater

systems, up to 40% of cells from the total community have been found to be inactive or dormant

(Jones and Lennon, 2010). In addition, leucine incorporation is not universal across all taxa

(Salcher et al, 2013). In this context, it is interesting to reflect on the richness in absence of

function (i.e. x-intercept) of the observed BEF relationship which is 295 (Figure 2C). This could

be interpreted as a baseline level of 295 particle-associated OTUs that are inactive (either dead or

dormant cells or environmental DNA) or incapable of incorporating leucine. This value

represents 35-85% of the total particle-associated communities and may obscure the *actual*

diversity (and BEF relationship) of the bacterial community (Carini et al., 2016).

In conclusion, we show that increased bacterial diversity leads to increased bacterial

heterotrophic production in particle-associated but not in free-living communities. As such, we

extend the validity of principles of the impact of microhabitat on BEF relationships from

Eukarya to Bacteria, contributing to current efforts to integrate ecological theories into the field

of microbiology (Barberán et al., 2014). Additionally, we show that communities with low

phylogenetic diversity have higher per-capita heterotrophic production rates, which we

hypothesize to be related to genome evolutionary patterns specific to bacteria that result in the

dependence on metabolic hand-offs. The unique nature of BEF relationships across particle-

associated and free-living habitats agrees with the distinct community assembly and functional

partitioning previously described between these two aquatic habitats (Bižić-Ionescu et al., 2014;

Ganesh et al., 2014; Mohit et al., 2014; Schmidt et al., 2016; Balmonte et al., 2018). Future

studies should focus on going beyond observations to predicting changes in ecosystem function

based on community changes (Bell, 2019).

Acknowledgements

This work was supported by the National Science Foundation Graduate Research Fellowship

Grant No. DGE 1256260 (MLS), the University of Michigan Office for Research MCubed

program (VJD), the American Society of Microbiology-Undergraduate Research Fellowship, the

University of Michigan Honors Summer Fellowship, and the Beckman Scholars Program (EC).

RP was supported by Ghent University (BOFDOC2015000601) and a Sofina Gustave-Boël grant

from the Belgian American Educational Foundation. We are grateful to the crew of the R/V

W.G. Jackson and the Grand Valley State University Robert B. Annis Water Resources Institute

science staff, and the generous help we received in the field from Amelia Waters and Daniel

S.W. Katz. Thank you to Kyle Buffin and Amadeus Twu for help with DNA extractions. Finally,

we thank Deborah Goldberg, George Kling, and members of the Denef, Dick, and Duhaime

laboratories for their comments on the manuscript.

References

Anantharaman, K. et al. 2016. Thousands of microbial genomes shed light on interconnected
biogeochemical processes in an aquifer system. Nature Communications 7:13219.
Balmonte, J.P., Teske, J.P., and C. Arnosti. 2018. Structure and function of high Arctic pelagic,
particle-associated and benthic bacterial communities. 20(8):2941–2954.
Barberán, A., E. O. Casamayor, and N. Fierer. 2014. The microbial contribution to
macroecology. Frontiers in Microbiology 5:1–8.
Bell, T. 2019. Next-generation experiments linking community structure and ecosystem
functioning. Environmental Microbiology Reports, 11(1):20–22.
Bižić-Ionescu, B., Zeder, M., Ionescu. D., Orlic, S., Fuchs, B.M., Grossart, H.P., and R. Amann.
2014. Comparison of bacterial communities on limnic versus coastal marine particles reveals
profound differences in colonization. Environmental Microbiology doi:10.1111/1462-
2920.12466.
Blaser, M. J. 2014. Missing Microbes: How the Overuse of Antibiotics Is Fueling Our Modern
Plagues. Henry Holt and Company LLC, New York 35:261.
Bushnell B. 2016. BBMap short read aligner. https://sourceforge.net/projects /bbmap/.
Cadotte, M. W., B. J. Cardinale, and T. H. Oakley. 2008. Evolutionary history and the effect of
biodiversity on plant productivity. Proceedings of the National Academy of Sciences
105:17012–17017.
Caporaso, J. G. et al. 2012. Ultra-high-throughput microbial community analysis on the Illumina
HiSeq and MiSeq platforms. The ISME Journal 6:1621–1624.
Cardinale, B. J. 2011. Biodiversity improves water quality through niche partitioning. Nature
472:86–89.
Cardinale, B. J. et al. 2012. Biodiversity loss and its impact on humanity. Nature 489:326–326.
Carini, P., P. J. Marsden, J. W. Leff, E. E. Morgan, M. S. Strickland, and N. Fierer. 2016. Relic
DNA is abundant in soil and obscures estimates of soil microbial diversity. Nature
Microbiology 2:16242.

Cordero, O. X., and M. S. Datta. 2016. ScienceDirect Microbial interactions and community assembly at microscales. Current Opinion in Microbiology 31:227–234.

Cotner, J. B., and B. A. Biddanda. 2002. Small players, large role: Microbial influence on biogeochemical processes in pelagic aquatic ecosystems. Ecosystems 5:105–121.

Crump, B.C., Armbrust, E.V., and J.A. Baross. 1999. Phylogenetic analysis of particle-attached and free-living bacterial communities in the Columbia River, its estuary, and the adjacent coastal ocean. Applied and Environmental Microbiology 65(7):3192–3204.

Datta, M. S., Sliwerska, E., Gore, J., Polz, M. F. and O. X. Cordero. 2016. Microbial interactions lead to rapid micro-scale successions on model marine particles. Nature Communications 7:11965.

Delgado-Baquerizo, M., L. et al. 2016. Lack of functional redundancy in the relationship between microbial diversity and ecosystem functioning. Journal of Ecology 104:936–946.

del Giorgio, P. A., and J. J. Cole. 1998. Bacterial Growth Efficiency in Natural Aquatic Systems. Annual Review of Ecology and Systematics 29:503–541.

Duffy, J.E., Godwin, C.M., and B.J. Cardinale. 2017. Biodiversity effects in the wild are common and as strong as key drivers of productivity. Nature 549:261–264.

Ebrahimi, A., Schwartzman, J., and O.X. Cordero. Cooperation and spatial self-organization determine ecosystem function for polysaccharide-degrading bacteria. bioRxiv:640961.

Enke, T.N., et al. 2019. Modular assembly of polysaccharide-degrading marine microbial communities. Current Biology 29:1–8.

EPA. 1993. Methods for the determination of inorganic substances in environmental samples. USEPA 600/R-93/100.

Evans, R., Alessi, A.M., Bird, S., McQueen-Mason, S.J., Bruce, N.C., and M.A. Brockhurst. 2017. Defining the functional traits that drive bacterial decomposer community productivity. The ISME Journal 11:1680–1687.

Faith, D. P. 1992. Conservation evaluation and phylogenetic diversity. Biological Conservation 61:1–10.

Ferrier S, K. N. Ninan, P. Leadley, R. Alkemade GK, M. Moraes R., & E. Y. Mohammed and Y. Trisurat. 2016. Overview and vision. In IPBES 2016: The methodological assessment report on scenarios and models of biodiversity and ecosystem services. S. Ferrier, K. N. Ninan, P. Leadley, R. Alkemade, L. A. Acosta, H. R. Akçakaya, L. Brotons, W. W. L. Cheung, V. Christensen, K. A. Harhash, J. KabuboMariara, C. Lundquist, M. Obersteiner, H. M. Pereira, G. Peterson, R. Pichs-Madruga, N. Ravindranath, C. Rondinini and B.A. Wintle (eds.), Bonn, Germany, Secretariat of the Intergovernmental Science-Policy Platform for Biodiversity and Ecosystem Services.

Flynn, D.F.B., Mirotchnick, N., Jain, M., Palmer, M.I., and S. Naeem. 2011. Functional and phylogenetic diversity as predictors of biodiversity–ecosystem-function relationships. Ecology 92(8): 1573–1581.

Franklin, R.B., Garland, J.L., Bolster, C.H., and A.L. Mills. 2001. Impact of dilution on microbial community structure and functional potential: Comparison of numerical

simulations and batch culture experiments. Applied and Environmental Microbiology 67(2):702–712.

Friedman, J., T. Hastie, and R. Tibshirani. 2010. Regularization Paths for Generalized Linear Models via Coordinate Descent. Journal of Statistical Software 33:1–22.

Fritschie, K. J., B. J. Cardinale, M. A. Alexandrou, and T. H. Oakley. 2014. Evolutionary history and the strength of species interactions: Testing the phylogenetic limiting similarity hypothesis. Ecology 95:1407–1417.

Galand, P. E., I. Salter, and D. Kalenitchenko. 2015. Ecosystem productivity is associated with bacterial phylogenetic distance in surface marine waters. Molecular Ecology 24:5785–5795.

Ganesh, S., D. J. Parris, E. F. DeLong, and F. J. Stewart. 2014. Metagenomic analysis of size-fractionated picoplankton in a marine oxygen minimum zone. The ISME journal 8:187–211.

Goberna, M. and M. Verdú. 2018. Phylogenetic-scale disparities in the soil microbial diversity–ecosystem functioning relationship. The ISME Journal 12:2152–2162.

Gravel, D., Bell, T., Barbera, C., Bouvier, T., Pommier, T., Venail, P., and N. Mouquet. 2011. Experimental niche evolution alters the strength of the diversity-productivity relationship. Nature 469:89–92.

Griffiths, B. S. et al. 2000. Ecosystem response of pasture soil communities to fumigation-induced microbial diversity reductions: an examination of the biodiversity-ecosystem function relationship. Oikos 90:279–294.

Grossart, H.P., Tang, K.W., Kiørboe, T., and H. Ploug. 2007. Comparison of cell-specific activities between free-living and attached bacteria using isolates and natural assemblages. FEMS Microbiology Letters 266:194–200.

Grossart, H. P. 2010. Ecological consequences of bacterioplankton lifestyles: Changes in concepts are needed. Environmental Microbiology Reports 2:706–714.

Hobbie, J. E., Daley, R. J., and S. Jasper. 1977. Use of nuclepore filter counting bacteria by fluorescence microscopy. Applied and Environmental Microbiology 33:1225–1228.

Jiang, L., J. Tan, and Z. Pu. 2010. An Experimental Test of Darwin's Naturalization Hypothesis. The American Naturalist 175:415–423.

Jones, S. E., and J. T. Lennon. 2010. Dormancy contributes to the maintenance of microbial diversity. Proceedings of the National Academy of Sciences 107:5881–5886.

Kembel, S. W. 2009. Disentangling niche and neutral influences on community assembly: Assessing the performance of community phylogenetic structure tests. Ecology Letters 12:949–960.

Kembel, S.W. et al.. 2010. Picante: R tools for integrating phylogenies and ecology. Bioinformatics 26:1463-1464.

Kim, M., Oh, H. S., Park, S. C., and J. Chun. 2014. Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. International Journal of Systematic and Evolutionary Microbiology 64:346–351.

Kirchman, D., E. K'nees, and R. Hodson. 1985. Leucine incorporation and its potential as a
measure of protein synthesis by bacteria in natural aquatic systems. Applied and
Environmental Microbiology 49:599–607.
Kirwan, L. et al. 2007. Evenness drives consistent diversity effects in intensive grassland
systems across 28 European sites. Journal of Ecology 95:530–539.
Konstantinidis, K. T. and J. M. Tiedje. 2005. Genomic insights that advance the species
definition for prokaryotes. Proceedings of the National Academy of Sciences 102:2567–
2572.
Kozich, J. J., S. L. Westcott, N. T. Baxter, S. K. Highlander, and P. D. Schloss. 2013.
Development of a dual-index sequencing strategy and curation pipeline for analyzing
amplicon sequence data on the miseq illumina sequencing platform. Applied and
Environmental Microbiology 79:5112–5120.
Krause, S. et al. 2014. Trait-based approaches for understanding microbial biodiversity and
ecosystem functioning. Frontiers in Microbiology 5:1–10.
Langenheder, S., E. S. Lindström, and L. J. Tranvik. 2006. Structure and Function of Bacterial
Communities Emerging from Different Sources under Identical Conditions. Applied and
Environmental Microbiology 72:212–220.
Levine, U. Y., T. K. Teal, G. P. Robertson, and T. M. Schmidt. 2011. Agriculture's impact on
microbial diversity and associated fluxes of carbon dioxide and methane. The ISME Journal
5:1683–1691.
Lilja, E. E. and D. R. Johnson. 2016. Segregating metabolic processes into different microbial
cells accelerates the consumption of inhibitory substrates. The ISME Journal 10:1–11.
Magurran, A. E. 2004. Chapter four: An index of diversity in Measuring Biological Diversity,
Wiley-Blackwell, Hoboken, NJ.
McMurdie, P. J., and S. Holmes. 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLoS ONE 8:e61217.
Mohit, V., Archambault, P., Toupoint, N., and C. Lovejoy. 2014. Phylogenetic differences in
attached and free-living bacterial communities in a temperate coastal lagoon during summer,
revealed via high-throughput 16S rRNA gene sequencing. Applied and Environmental
Microbiology 80: 2071–2083.
Muscarella, M.E., Boot, C. M., Broeckling, C. D., and J. T. Lennon. 2019. Resource
heterogeneity structures aquatic bacterial communities. The ISME Journal 13:2183–2195.
Narwani, A. et al. 2017. Ecological interactions and coexistence are predicted by gene
expression similarity in freshwater green algae. Journal of Ecology 105:580–591.
Oksanen, A. J. et al. 2015. vegan: Community Ecology Package. R package version 2.3-0.
Peter, H., S. Beier, S. Bertilsson, E. S. Lindström, S. Langenheder, and L. J. Tranvik. 2011.
Function-specific response to depletion of microbial diversity. The ISME Journal 5:351–361.
Philippot, L. et al. 2013. Loss in microbial diversity affects nitrogen cycling in soil. The ISME
Journal 7:1609–1619.

Price, M. N., P. S. Dehal, and A. P. Arkin. 2010. FastTree 2 – Approximately maximum-
likelihood trees for large alignments. PLoS ONE 5.

Quast, C. et al. 2013. The SILVA ribosomal RNA gene database project: Improved data
processing and web-based tools. Nucleic Acids Research 41:590–596.

R Core Team. 2017. R: A Language and Environment for Statistical computing. Vienna, Austria:
R Foundation for Statistical Computing. https:// www.R-project.org/.

Reich, P.B. et al. 2004. Species and functional group diversity independently influence biomass
accumulation and its response to CO_2 and N. Proceedings of the National Academy of
Sciences 101(27):10101–10106.

Rivett, D.W. and T. Bell. 2018. Abundance determines the functional role of bacterial
phylotypes in complex communities. Nature Microbiology:s41564-018-0180-0.

Roger, F., S. Bertilsson, S. Langenheder, O. A. Osman, and L. Gamfeldt. 2016. Effects of
multiple dimensions of bacterial diversity on functioning, stability and multifunctionality.
Ecology 97:2716–2728.

Rohwer, R. R., J. J. Hamilton, R. J. Newton, and K. D. McMahon. 2018. TaxAss: Leveraging a
custom freshwater database achieves fine-scale taxonomic resolution. mSphere 3(5):e00327-
18.

Russel, J., H. L. Røder, J. S. Madsen, M. Burmølle, and S. J. Sørensen. 2017. Antagonism
correlates with metabolic similarity in diverse bacteria. Proceedings of the National Academy
of Sciences:201706016.

Salcher, M.M., Posch, T., and J. Pernthaler. 2013. *In situ* substrate preferences of abundant
bacterioplankton populations in a prealpine freshwater lake. The ISME Journal 7:896–907.

Salles, J. F. et al. 2009. Community niche predicts the functioning of denitrifying bacterial
assemblages. Ecology, 90:3324–3332.

Schloss, P. D. et al. 2009. Introducing mothur: Open-source, platform-independent, community-
supported software for describing and comparing microbial communities. Applied and
Environmental Microbiology 75:7537–7541.

Schmidt, M.L., J.D. White, and V. J. Denef. 2016. Phylogenetic conservation of freshwater lake
habitat preference varies between abundant bacterioplankton phyla. Environmental
Microbiology 18(4):1212–1226.

Shade, A., S. E. Jones, and K. D. McMahon. 2008. The influence of habitat heterogeneity on
freshwater bacterial community composition and dynamics. Environmental Microbiology
10:1057–1067.

Simon, M., and F. Azam. 1989. Protein content and protein synthesis rates of planktonic marine
bacteria. Marine Ecology Progress Series 51:201–213.

Simon, M., H. P. Grossart, B. Schweitzer, and H. Ploug. 2002. Microbial ecology of organic
aggregates in aquatic ecosystems. Aquatic Microbial Ecology 28:175–211.

Singh, B. K. et al. 2014. Loss of microbial diversity in soils is coincident with reductions in some
specialized functions. Environmental Microbiology 16:2408–2420.

Steinman, A. D., Ogdahl, M., Rediske, R., Ruetz III, C.R., Biddanda, B. A., and L. Nemeth.
2008. Current status and trends in Muskegon Lake, Michigan. Journal of Great Lakes
Research 34:169–188.
Stewart, E. J. 2012. Growing unculturable bacteria. Journal of Bacteriology 194:4151–4160.
Tan, J., Z. Pu, W. A. Ryberg, and L. Jiang. 2012. Species phylogenetic relatedness, priority
effects, and ecosystem functioning. Ecology 93:1164–1172.
Thomas, C. D. et al. 2004. Extinction risk from climate change. Nature 427:145–8.
Tilman, D., F. Isbell, and J. M. Cowles. 2014. Biodiversity and Ecosystem Functioning. Annual
Review of Ecology, Evolution, and Systematics 45:471–493.
Tylianakis, J. M. et al. 2008. Resource heterogeneity moderates the biodiversity-function
relationship in real world ecosystems. PLoS Biology 6:0947–0956.
Venail, P. A., A. Narwani, K. Fritschie, M. A. Alexandrou, T. H. Oakley, and B. J. Cardinale.
2014. The influence of phylogenetic relatedness on species interactions among freshwater
green algae in a mesocosm experiment. Journal of Ecology 102:1288–1299.
Venail, P. A., and M. J. Vives. 2013. Phylogenetic distance and species richness interactively
affect the productivity of bacterial communities. Ecology 94:2529–2536.
Wake, D. B., and V. T. Vredenburg. 2008. Are we in the midst of the sixth mass extinction? A
view from the world of amphibians. Proceedings of the National Academy of Sciences
105:11466–11473.
Wertz, S., V. et al. 2006. Maintenance of soil functioning following erosion of microbial
diversity. Environmental Microbiology 8:2162–2169.
Wilsey, B.J. and H. W. Polley. 2004. Realistically low species evenness does not alter grassland
species richness productivity relationships. Ecology 85(10):2693–2700.
Wilsey, B.J. and C. Potvin. 2000. Biodiversity and ecosystem functioning: Importance of species
evenness in an old field. Ecology 81(4):887–892.
Wittebolle, L. et al. 2009. Initial community evenness favors functionality under selective stress.
Nature 458(2)623–626.
Zeppilli, D., A. Pusceddu, F. Trincardi, and R. Danovaro. 2016. Seafloor heterogeneity
influences the biodiversity–ecosystem functioning relationships in the deep sea. Scientific
Reports 6:26352.
Zhou, J., S. et al. 2008. Spatial scaling of functional gene diversity across various microbial
taxa. Proceedings of the National Academy of Sciences 105:7768–7773.
709

Figure 1: Bacterial counts, community-wide and per-capita heterotrophic production differ between microhabitats. Particle-associated and free-living samples were taken from four stations within Muskegon Lake during 2015 in May, July, and September. **(A)** Free-living bacteria were an order of magnitude (10_6 cells/mL) more abundant compared to particle-associated bacteria. **(B)** Free-living bacteria were more heterotrophically productive compared to particle-associated bacteria. **(C)** Particle-associated bacteria were disproportionately heterotrophically productive per cell compared to free-living bacteria.

Figure 2: Richness and inverse Simpson correlate with heterotrophic productivity. Top panel: Differences in **(A)** the observed richness and **(B)** the inverse Simpson diversity metrics between particle-associated (orange) and free-living (blue) habitats. **Middle panel:** Biodiversity and community-wide heterotrophic production (ugC/L/day) relationships. The y-axis between **(C)** and **(D)** is the same, however, the x-axis represents **(C)** richness and **(D)** inverse Simpson. **Bottom panel:** Biodiversity and $\log_{10}$(per-capita heterotrophic production) (ugC/cell/day) relationships. The y-axis between **(E)** and **(F)** is the same, however, the x-axis represents **(E)** richness and **(F)** inverse Simpson's index. Solid lines represent ordinary least squares models for the free-living (blue) and particle associated (orange) communities. All R_2 values represent the adjusted R_2 from an ordinary least squares model.

Figure 3: The relationship between heterotrophic productivity and unweighted phylogenetic diversity (SES$_{MPD}$; ses.mpd function in *picante* with null.model = "independentswap"). Positive phylogenetic diversity values represent communities that are phylogenetically diverse (*i.e.* overdispersed) while negative phylogenetic diversity values represent communities that are phylogenetically less diverse (*i.e.* clustered) compared to a null community with equal species richness. **(A)** Phylogenetic diversity was higher in free-living communities compared to particle-associated communities. **(B)** Negative relationship between observed richness and phylogenetic diversity. **(C)** Absence of phylogenetic diversity and community bulk heterotrophic production (µgC/L/day) relationships. **(D)** Negative phylogenetic diversity and per-capita heterotrophic production (µgC/cell/day) relationship. Linear models in figure **B** and **D** represent trends over all samples.

741   **Figure 1**

743 **Figure 2**

Figure 3