Predicting Response to Platin Chemotherapy Agents with Biochemically-inspired Machine Learning

Eliseos J. Mucaki[1], Jonathan Z.L. Zhao[1], Dan Lizotte[2,3], and §Peter K. Rogan[1,2,3,4,5]

Running Title:

Predicting Responses to Platin Drugs by Machine Learning

Author Affiliations

[1]Department of Biochemistry, Schulich School of Medicine and Dentistry, Western University,

London, Canada, N6A 2C1

[2]Department of Computer Science, Faculty of Science, Western University, London, Canada, N6A

2C1

[3]Department of Epidemiology & Biostatistics, Faculty of Science, Western University, London,

Canada, N6A 2C1

[4]Cytognomix Inc. London, Canada N5X 3X5

[5]Department of Oncology, Schulich School of Medicine and Dentistry, Western University,

London, Canada, N6A 2C1

Author Emails: emucaki@uwo.ca, jzhao293@uwo.ca, dlizotte@uwo.ca, progan@uwo.ca

§**Correspondence to:** Peter K. Rogan (progan@uwo.ca), Department of Biochemistry, Schulich
School of Medicine and Dentistry, Western University, London, Ontario, Canada, N6A 2C1. 1
(519) 661-4255.

ABSTRACT.

Selection of effective genes that accurately predict chemotherapy response could improve cancer outcomes. We compare optimized gene signatures for cisplatin, carboplatin, and oxaliplatin response, and respectively validate each with cancer patient data. Supervised support vector machine learning was used to derive gene sets whose expression was related to cell line GI_{50} values by backwards feature selection with cross-validation. Signatures at different GI_{50} thresholds distinguishing sensitivity from resistance contrast the contributions of genes at extreme vs. median thresholds. Ensembles of gene signatures at different thresholds are combined to reduce dependence on specific GI_{50} values for predicting drug response. The most accurate models for each platin are: cisplatin: *BARD1*, *BCL2*, *BCL2L1*, *CDKN2C*, *FAAP24*, *FEN1*, *MAP3K1*, *MAPK13*, *MAPK3*, *NFKB1*, *NFKB2*, *SLC22A5*, *SLC31A2*, *TLR4*, *TWIST1*; carboplatin: *AKT1*, *EIF3K*, *ERCC1*, *GNGT1*, *GSR*, *MTHFR*, *NEDD4L*, *NLRP1*, *NRAS*, *RAF1*, *SGK1*, *TIGD1*, *TP53*, *VEGFB*, *VEGFC;* oxaliplatin: *BRAF*, *FCGR2A*, *IGF1*, *MSH2*, *NAGK*, *NFE2L2*, *NQO1*, *PANK3*, *SLC47A1*, *SLCO1B1*, *UGT1A1*. TCGA bladder, ovarian and colorectal cancer patients were used to test cisplatin, carboplatin and oxaliplatin signatures (respectively), resulting in 71.0%, 60.2% and 54.5% accuracy in predicting disease recurrence and 59%, 61% and 72% accuracy in predicting remission. One cisplatin signature predicted 100% of recurrence in non-smoking bladder cancer patients (57% disease-free; N=19), and 79% recurrence in smokers (62% disease-free; N=35). This approach should be adaptable to other studies of chemotherapy response, independent of drug or cancer types.

INTRODUCTION

Chemotherapy regimens are selected based on overall outcomes for specific types and subtypes of cancer pathology, progression to metastasis, other high-risk indications, and prognosis[1,2], and variability in tumor resistance has led to tiered sequential strategies for selection of agents based on their overall efficacy[3]. We and others have developed machine learning (ML)-based gene signatures aimed at predicting response to specific chemotherapeutic agents and minimizing chemoresistance based on inhibition of growth or drug targets (GI_{50} or IC_{50})[4–6]. In this study, we present integrated ML models of platin responses (cis-, carbo- and oxaliplatin). Previous studies have reviewed the genes[7], gene products[8] and specific individual pathways[9], but not developed a comprehensive model of the global cell response to the drug. We use integrated models based on expression of multiple genes to predict responses to each of these platin agents, for the first time, at different resistance levels.

Cisplatin, carboplatin and oxaliplatin are each widely prescribed compounds for their antineoplastic effects. While each contains platinum to form adducts with tumour DNA, their effectiveness differs for specific types of cancers, such as bladder (cisplatin), ovarian (cisplatin and carboplatin) and colorectal cancer (oxaliplatin). Carboplatin differs in structure from cisplatin, exchanging the latter's dichloride ligands with a CBDCA (cyclobutane dicarboxylic acid) group, while oxaliplatin is paired with both a DACH (diaminocyclohexane) ligand and a bidentate oxalate group. These chelating ligands have greater stability and solubility to aqueous solutions, which lead to differences in drug toxicity compared to cisplatin[10]. Oxaliplatin can be up to two times as cytotoxic as cisplatin, but it forms fewer DNA adducts[11]. The large hydrophobic DACH ligand which overlaps the major groove is thought to prevent binding of certain DNA repair enzymes such as the POL polymerases, and may contribute to the low cross-resistance between

oxaliplatin and cisplatin and carboplatin[10]. While all three drugs can enter the cell via

copper transporters, organic cation transporters are oxaliplatin-specific and likely play a

role in its efficacy in colorectal cancer (CRC) cells where these transporters are

commonly overexpressed[7]. Oxaliplatin specifically plays a role in interfering with both

DNA and RNA synthesis, unlike cisplatin which only infers with DNA[12]. It is these

intrinsic properties between the platinum drugs which lead to differences in their activity

and resistance profiles, despite their similar mode of action.

We derived gene signatures to predict drug response at different sensitivity and

resistance levels for each of these agents. We and others have used supervised

learning algorithms, including random forest models[13]; support vector machine (SVM)

models[6]; neural networks[14]; and linear regression models[5] to make these predictions.

Pathway and network analysis of gene expression have been used to indicate hundreds

of genes potentially up- and down-regulated upon cisplatin treatment[15]. Cisplatin-specific

gene signatures have been developed with integrative approaches such as elastic net

regression using inferred pathway activity of bladder cancer cell line data[16]. These

methods have implicated genes that have not been described previously. Supervised ML

with biochemically-relevant genes has also been useful for predicting drug response[6]. A

concern with each of these ML approaches is that an insufficient number of samples

coupled to a large number of features, i.e. gene expression changes, in each sample

can result in overfitting of the model affecting its generalizability with other sources of

data[17]. We therefore reduce the number of dimensions by selecting genes biologically

relevant to the drugs under observation[6,17]. Additional selection criteria are necessary

when the number of genes implicated in peer-reviewed reports is still prohibitively large

compared to sample size.

Biochemically-inspired gene signatures have shown good performance in

predicting treatment response. A paclitaxel ML signature based on tumor gene

expression had a higher success predicting the pathological complete response rate

(pCR [18]) for sensitive patients (84% of patients with no / minimal residual disease) than

models based on differential gene expression (GE) analysis[6]. For gemcitabine, a

signature derived from both expression and copy number (CN) data from breast cancer

cell lines was derived, and subsequently applied to analysis of nucleic acids from patient

archival material. Multiple other outcome measures used to validate gene signatures

include prognosis[5], Miller-Payne response[19], and disease recurrence. Binary SVM

classifiers based on discrete time thresholds have been used to classify continuous

outcome measures such as prognosis and recurrence. By contrast, pCR is simpler to

interpret with binary SVM models. Nevertheless, differences in clinical recurrence have

been noted between patients demonstrated with pCR and those who do not exhibit

disease pathology[18]. This source of variability in defining patient response can confound

transferability of SVM models between different datasets.

We apply biochemically-inspired ML to predict and compare the cellular and

patient responses to cisplatin, carboplatin and oxaliplatin. We train models for

classification of platin resistance with cancer cell line data and validate with patient GE

and outcome data. Our previous gene signatures were based on median GI_{50} for each

drug[6]. This has been a necessary compromise, however in this study we consider

signatures that differ at the highest vs. the lowest levels of drug resistance. A series of

gene signatures are derived by shifting the GI_{50} thresholds that distinguish sensitivity

from resistance. The frequency of genes selected at median vs. extreme thresholds

highlights pathways that define these responses among different patient subsets.

RESULTS

Selection of Platin Drug Related Genes

We documented genes in the peer-reviewed literature associated with drug

effectiveness or response (Supplemental References). For cisplatin, carboplatin and

oxaliplatin, this implicated 178, 90, and 288 genes, respectively (Suppl. Table S1).

Multiple factor analyisis (MFA) was used to determine which genes were correlated to

GI_{50} in breast cancer cell lines through either GE and/or CN^{13}, significantly reducing the

sizes of the gene sets for cisplatin (N=39), carboplatin (N=28), and oxaliplatin (N=55).

Genes with significant relationships to GI_{50} and direction of correlation (positive or

inverse) are indicated in Figure 1. The diverse functions of these genes included

apoptosis, DNA repair, transcription, cell growth, metabolism, immune system, signal

transduction and membrane transport. Analysis of IC_{50} and gene expression levels for

cisplatin-treated bladder cancer cell lines confirmed these relationships evident from GI_{50}

values of different   breast cancer lines. IC_{50} values were related to GE for *CFLAR*,

FEN1, *MAPK3*, *MSH2*, *NFKB1*, *PNKP*, *PRKAA2*, and *PRKCA*[20]. Similarly, separate

bladder cell line IC_{50} values from the Genomics of Drug Sensitivity in Cancer project

(http://www.cancerrxgene.org; N=17) were correlated with GE for *CFLAR*, *FEN1*, and

NFKB1, in addition to *ATP7B*, *BARD1*, *MAP3K1*, *NFKB2*, *SLC31A2* and *SNAI1*.

We performed MFA on the GI_{50} values for cisplatin, carboplatin and oxaliplatin,

without consideration of either GE or CN. Responses to cis- and carboplatin were

directly correlated (a 6.2º separation between vectors), but neither was related to the

oxaliplatin response (Figure 2). Previous studies have shown that cisplatin-resistant cell

lines are generally sensitive to oxaliplatin[21–23].

SVM-based signatures were initially derived for each platin drug from breast

cancer cell line GE data. A 13-gene signature for cisplatin at the median GI_{50} threshold

(5.2% misclassification rate) consisted of *BARD1, BCL2L1, FAAP24, CFLAR, MAP3K1,*

MAPK3, NFKB1, POLQ, PRKAA2, SLC22A5, SLC31A2, TLR4, and *TWIST1*. A similarly

derived carboplatin signature included *AKT1*, *ATP7B*, *EGF*, *EIF3I*, *ERCC1*, *GNGT1*,

HRAS, *MTR*, *NRAS*, *OPRM1*, *RAD50*, *RAF1*, *SCN10A*, *SGK1*, *TIGD1*, *TP53*, and

VEGFB (10.4% misclassification). For oxaliplatin, the final SVM model consisted of

AGXT, *APOBEC2*, *BRAF*, *CLCN6*, *FCGR2A*, *IGF1*, *MPO*, *MSH2*, *NAGK*, *NAT2*,

NFE2L2, *NOTCH1*, *PANK3*, *PRSS1*, and *UGT1A1* (2.1% misclassification). A cisplatin

SVM generated from 17 bladder cancer cell lines in cancerRxgeneresulted in 2 equally

accurate signatures (with 11.8% misclassification) consisting of either *PNKP* and

PRKCA or *ATP7B*, *CFLAR*, *FEN1*, *MAPK3*, *NFKB1* and *SLC22A11*. These models were

not useful for predicting patient outcomes due to the limited size of the training set

GI_{50}-Threshold Independent Modeling

In our previous studies, we set median GI_{50} value as the threshold to

distinguished drug resistance and sensitivity[5,6]. An important question is whether the

genes contributing to drug response are consistent among different cell lines, each with

their own unique GI_{50} values. Different ML models were obtained by shifting the GI_{50}

threshold, which changed the labels of resistant vs. sensitive cell lines. After feature

selection, the compositions of the corresponding gene signatures for each threshold

were compared. Finally, ensemble averaging of all of these optimized Gaussian SVM

models derived for different GI_{50} thresholds was used to create a threshold-independent

ML-based signature.

Kinase (*MAPK3*, *MAP3K1*) genes and apoptotic family members (*BCL2*,

BCL2L1) were most the common in the cisplatin signatures at different GI_{50} thresholds,

with consistent representation of error-prone and base-excision DNA repair genes as

well (Figure 3A; Supplementary Table S2A). The kinases are more concentrated in

signatures with lower drug sensitivity thresholds, whereas *BCL2* and *BCL2L1* are more

ubiquitous at all levels. The error prone polymerases, *POLD1* and *POLQ,* are more

frequent in models with lower sensitivity thresholds, while the flap endonuclease *FEN1*

tends to be present at high levels of resistance. Thresholded models for carboplatin-

related genes commonly contained the apoptotic family member *AKT1*, transcription

regulation genes *ETS2* and *TP53*, as well as cell growth factors *VEGFB* and *VEGFC*,

although the latter was less common at lower sensitivity thresholds (Figure 3B).

Common oxaliplatin-related genes included transporters *SLCO1B1* and *GRTP1* (but not

SLC47A1), transcription genes *NFE2L2*, *PARP15* and *CLCN6*, as well as multiple

metabolism-related genes (Figure 3C).

SVM models were also derived using the cisplatin and/or carboplatin-treated

TCGA (The Cancer Genome Atlas) bladder urothelial carcinoma patients, using post-

treatment time to recurrence as a surrogate for different resistance thresholds (similarly

performed in Mucaki *et al.* [2017][24]; Supplementary Table S3). Similar trends to cell line

SVMs are apparent; *POLQ* is frequently selected when the resistance threshold is set at

a longer time period, while *FEN1* appears when it is shorter. However *BCL2*, which is

present in a majority of breast cancer cell line SVMs, is present in only one model

derived from TCGA data. Similarly, *MSH2* was rarely selected using cell lines, yet

appears in nearly all patient derived SVMs with > 1 year recurrence.

GI_{50}-thresholded models for each platin drug, generated with the breast cancer

cell line data, produced 70 cisplatin, 83 carboplatin, and 83 oxaliplatin SVM models,

respectively. Each model was validated using available platin-treated patient datasets[25–

29]. The chemotherapy response metadata differed between studies. Als *et al.*[28] reported

survival post-treatment, whereas Tsuji *et al.*[29] categorized patients as responders and

non-responders. TCGA provided two different measures which were used to assess

predictive accuracy in our models – chemotherapy response and disease-free survival.

Accuracy is similar using either measure (Supplementary Table S4A); however

recurrence and disease-free survival was used as the primary measure of response as it

was more often recorded in the TCGA data sets tested. Patients from Als *et al.* with a ≥

5 year survival post-treatment were labeled as sensitive to treatment. The differences

between these metadata may, in part, account for the differences in the prediction

accuracy of the thresholded SVM models.

At higher resistance thresholds for any platin drug (low GI_{50}), where more cell

lines are labeled sensitive, the positive class (disease-free survival) is correctly

classified, while the negative class (recurrence) is highly misclassified (Suppl. Figures 1

and 2). The reverse is true for models built using lower resistance thresholds (high GI_{50}).

We therefore state SVMs generated at these extreme thresholds are not very useful at

predicting patient data. When used to predict recurrence in the TCGA datasets,

sensitivity and specificity appears to be maximized in models where the GI_{50} threshold

for resistance was set near (but not necessarily at) the median (Suppl. Figure 1; Suppl.

Tables S4A to 4C). While this pattern holds true for Tsuji *et al.*[29], oxaliplatin models

where GI_{50} thresholds were set above the median could better separate primary and

metastatic CRC patients (best model predicting 92.6% metastatic and 60.7% primary

cancers; Suppl. Table S4C). While less consistent, cisplatin models generated with

thresholds above median GI_{50} performed better when evaluating the Als *et al.*[28] patient

dataset (Suppl. Figure 2).

Models were further evaluated for their accuracy in TCGA patients using various

recurrence times post-treatment to classify resistant and sensitive patients (0.5 - 5 years;

Supplemental Table S5A-C). The best performing cisplatin model (*BARD1, BCL2,*

BCL2L1, CDKN2C, FAAP24, FEN1, MAP3K1, MAPK13, MAPK3, NFKB1, NFKB2,

SLC22A5, SLC31A2, TLR4, TWIST1; GI_{50} of 5.11; hereby identified as **Cis1**) was able

to accurately predict 71.0% of bladder cancer patients who recurred after 18 mo. (N=31;

58.5% accurate for disease-free patients [N=41]). Response of TCGA bladder patients

treated with carboplatin (without cisplatin; N=19) were best predicted by **Cis12** two years

post-treatment (80% accurate for responding patients [N=5]; 93% for recurrent patients

[N=14]). The best performing carboplatin model (*AKT1, EIF3K, ERCC1, GNGT1, GSR,*

MTHFR, NEDD4L, NLRP1, NRAS, RAF1, SGK1, TIGD1, TP53, VEGFB, VEGFC;

designated **Car1** [Suppl. Table S5B]) was developed using a GI_{50} threshold of 4.42,

predicting recurrence of ovarian cancer after 4 years at an accuracy of 60.2% (N=302;

61.0% accurate for disease-free patients [N=108]). These models were also used to test

TCGA bladder cancer patients treated carboplatin but not cisplatin (N=19), of which the

best performing model (**Car73**) was 84% accurate for patients after 1 year of treatment

(100% for responding patients [N=11]; 62.5% accuracy for recurrent [N=8]). Two

additional carboplatin models are tied for overall accuracy (84%; **Car9** and **Car51**), but

more successfully predict non-responsive patients (87.5%; 82% accuracy for responding

patients). These three models share four genes: *AKT1, ETS2, GNGT1*, and *VEGFB.* For

oxaliplatin, the best performing model (*BRAF, FCGR2A, IGF1, MSH2, NAGK, NFE2L2,*

NQO1, PANK3, SLC47A1, SLCO1B1, UGT1A1; GI_{50} of 5.10; designated **Oxa1** [Suppl.

Table S5C]) accurately predicted 71.6% of the disease-free TCGA CRC patients after

one year (N=88; 54.5% accuracy predicting recurrence [N=11]). These models, as well

as the previously mentioned SVMs based on bladder cell line data, were added to the

online web-based SVM calculator (http://chemotherapy.cytognomix.com; introduced in

Dorman *et al.* [2016][6]) which can be used to predict platin response using normalized

gene expression values.

To evaluate the consistency in the prediction of TCGA bladder cancer patients

treated with cisplatin, the predicted distance from the hyperplane for all SVMs generated

were plotted for each patient with a short recurrence time (<6 mo., N=10; Supplementary

Figure 3). Despite showing similar levels of resistance to treatment, patterns differed

between patients. While these patients would be expected to be indicated as highly

cisplatin resistant (hyperplane distance < 0), two patients (TCGA-XF-A9SU and TCGA-

FJ-A871) were predicted sensitive across nearly all SVM models. Similar variation was

also seen in patients with either a long recurrence time (>4 years) or no recurrence at all

after 6 years (Suppl. Figure 4).

Threshold independent models were generated for each individual platin drug at

different GI_{50} thresholds through ensemble ML, which involves the averaging of

hyperplane distances for each model to generate a composite score for each TCGA

patient tested. Hyperplane distances across all 70 cisplatin models were similar, with a

mean score of -0.22 and a standard deviation of 3.5 hyperplane units (hu) across the set

of patient data. The ensemble model classified disease-free bladder cancer patients

with 59% accuracy and those with recurrent disease with 47% accuracy. Limiting

ensemble averaging to only cisplatin models generated at a moderate GI_{50} threshold

(ranging from 5.10 to 5.50) did not significantly improve accuracy (44% for disease-free

and 66% for recurrent patients; Suppl. Table S6A). For carboplatin, ensemble ML did not

produce significantly better predictions than random, regardless of the GI_{50} threshold

interval selected (Suppl. Table S6B) or the similar mean hyperplane distances (-0.11 +/-

3.9 hu). For oxaliplatin, the ensemble ML model (mean = -0.12 +/- 2.7 hu) was most

accurate after 1 year (60% accuracy for disease-free and 73% for recurrent patients;

Suppl. Table S6C). As in cisplatin, limiting this analysis to oxaliplatin SVM models with

moderate GI_{50} thresholds did not significantly increase accuracy.

To determine the impact of individual genes on overall model accuracy, each

gene within every SVM model was excluded, and model accuracy was reassessed

(Supplementary Table S2A; S2B and S2C contain carbo- and oxaliplatin models,

respectively). Genes which consistently significantly increase misclassification

(averaging > 16% increase) in moderate threshold SVMs (GI_{50} thresholds set from 5.1 to

5.5) include *ERCC2*, *POLD1*, *BARD1*, *BCL2*, *PRKCA* and *PRKCB*. *ERCC2* and *POLD1*

perform critical functions in nucleotide and base excision repair, respectively. *PRKCA*

and *PRKCB* are paralogs with significant roles in signal transduction. *BARD1* has been

shown to reduce apoptotic *BCL2* in the mitochondria[30], and has a key role in genomic

stability through its association with *BRCA1*. Genes with a high variance in increased

misclassification between different models include *NFKB1*, *NFKB2*, *TWIST1*, *TP63*,

PRKAA2, and *MSH2*. The variance of these genes may be due to epistatic interactions

with other biological components, including the other genes in the SVM. For example,

NFKB1 and *NFKB2* are jointly included in 7 SVMs generated at a moderate GI_{50}

threshold. There is evidence of possible epistasis in that the removal of either of these

genes, but not necessary both, will have a large impact in model misclassification rates

(≥ 18.0% increase). The misclassification variance of *NFKB1* with *NFKB2*, is significantly

lower than in SVM models lacking *NFKB2*.

Predicting cisplatin response in patients based on smoking history

Tobacco smoking is known as the highest risk factor for the development of

bladder cancer[31]. We therefore subdivided the patients based on their smoking history

and tested the thresholded models (Supplementary Tables S7 and S8). When testing

patients who were lifelong non-smokers, the prediction accuracy of **Cis1** predicted all

non-smoking patients who were recurrent after 18 months as cisplatin-resistant (N=5).

Prediction accuracy for disease-free patients was 57.1% (N=14). Another model (**Cis18**;

Suppl. Table S7) had performed equally as well for non-smokers, and these two models

share 7 genes: *BCL2*, *BCL2L1*, *FAAP24*, *MAP3K1*, *MAPK13*, *MAPK3*, and *SLC31A2*.

Threshold independent analysis predicted disease-free equally well, but recurrence was

less accurate (66.7%). Note that non-smokers make up a small subset of the patients

tested (N=19). Threshold-independent prediction of recurrence in patients with a

smoking history was 46% accurate (N=13), while disease-free patients were correctly

predicted at a rate of 58% (N=19). Recurrence in these patients was best predicted by a

model built at the median GI_{50} threshold (**Cis2**). Accuracy improved for both disease-free

(57.7% -> 61.9%) and recurrent patients (76.0% -> 78.6%) when excluding patients who

quit smoking more than 15 years before diagnosis. Genes in this SVM which are not

present in the two models which performed well for non-smokers include *CFLAR* and

PRKAA2.

Tobacco smoking has a significant impact on cytosine methylation levels in the

genome[32]. CpG island methylation has been associated with smoking pack years in a

subset of the TCGA bladder urothelial carcinoma patients[25]. We suspected that the level

of methylation measured in t SVMs which performed best for smoking and non-smoking

patients might differ, and with possible concomitant effects on GE. When ranking each

gene from **Cis1** by highest methylation and GE, 88 of 1080 patient: gene combinations

showed the expected inverse correlation between methylation levels and GE (i.e. high

methylation and low GE). Inverse correlation of methylation and GE was more common

than direct correlation (i.e. high methylation and high GE; N=17). However, direct

correlation was more common in patients with a recent smoking history (70.5%). This

pattern was also observed for **Cis2**, which best predicted recurrence in smokers. In

cases where methylation and GE are directly correlated, we propose that smoking may

alter expression by other effects, eg. mutagenic, rather solely than by epigenetic

inactivation through methylation.

To determine which genes in these models led to discordant predictions of

patient outcome, we conducted a bioinformatic experiment to gradually alter the

expression of each signature gene until the misclassification was corrected. If the GE

value required to cross this threshold exceeded ≥ 3-fold the highest/lowest expression of

that gene, it was interpreted as a minor contributor to the prediction. Genes which could

not correct a discordant prediction included *PRKAA2*, *NFKB1*, *NFKB2* and *TWIST1*.

Significant genes which, when altered, corrected discordant predictions included

MAP3K1, *MAPK3*, *SLC22A5* and *SLC31A2*. Altering *BCL2L1* expression was more

likely to correct the discordant predictions of **Cis1** (4 out of 5) than with **Cis2** (2 out of 4).

DISCUSSION

Using gene expression signatures, we derived both GI_{50} threshold-dependent

and -independent ML models which predict the chemotherapy responses for cisplatin,

carboplatin and oxaliplatin, respectively. The cisplatin model **Cis1** (Supplementary Table

S5A) most accurately predicted response in bladder cancer patients after 18 months,

and **Car1** (Suppl. Table S5B) best predicted response in ovarian cancer patients after 4

years. **Oxa1** (Suppl. Table S5C) more accurately predicted disease-free patients than

recurrent disease at the one year treatment threshold. The thresholds which best

represented time-to-recurrence differed between the platin drugs in each cancer type.

Cisplatin gene signatures had noticeably improved performance when smoking history

was taken into account.

The three platin drugs produce distinctly different gene signature models. Initial

gene sets exhibited some overlap between platin drugs (N=67 between any two platins),

but very few of these were correlated by MFA of GI_{50} with multiple platin drugs (*ATP7B*,

BCL2 and *MSH2*). Despite the close similarity between cisplatin and carboplatin GI_{50}

response (see Figure 2), only one gene (*ATP7B*) was related by MFA to GI_{50} levels of

both drugs. *BCL2* and *MSH2* correlated with both cisplatin and oxaliplatin GI_{50}. The

increase in misclassification caused by the elimination of *MSH2* from any SVM model in

which it was present was significant; for example, in oxaliplatin model **Oxa21** (Suppl.

Table S5C) and cisplatin model **Cis14** (Suppl. Table S5A) misclassification was

increased by 19.1% and 28.2%, respectively. These differences may reflect the

spectrum of activity, sensitivity, and toxicity of these signature genes[21–23,33,34].

Gene signature models derived from cell lines and tested on patients differ in

their outcome measures. The exact GI_{50} cell line threshold that is most predictive of

patient outcome is not known, and different groups use different methods to discretize

GI_{50} values[35,36]. Therefore, we developed ML models for platin drugs which predict drug

response without relying on arbitrary GI_{50} thresholds. For cisplatin, SVM ensemble

averaging generated on different resistance thresholds shows a small increase in

accuracy over most models, better representing the sensitive, disease-free class (59%

accuracy). Interestingly, ensemble averaging of only the models built using a moderate

GI_{50} thresholds yielded results which better represented the resistance class. This result

closer matches the accuracy of **Cis1**, and may be due to **Cis1** having a greater overall

impact on the ensemble prediction. When limiting ensemble averaging to only those

models with the highest area under the curve (AUC) at each resistance threshold,

differences in predictions were negligible. Ensemble ML can potentially avoid problems

with poor performance and overfitting by combining models that individually perform

slightly better than chance[37].

It is difficult to reconcile gene signatures without features known to be related to

chemoresistance with tumor biology. Our thresholding approach may reveal potentially

important genes and pathways associated with platin resistance. It would be preferable

to explore pathways related to signature genes to improve accuracy, identify potential

targets for further study of chemoresistance, and expand the model parameters to take

into account alternate states besides those captured in the original signature[38].

Signatures for resistance may be useful for developing targeted intervention to re-

sensitize tumours. For example, the mismatch repair (MMR) gene *MSH2* is commonly

present in gene signatures at high resistance levels for oxaliplatin, which is of interest,

as MMR deficiency has been shown to be predictive for oxaliplatin resistance[34]. Indeed,

MLH1, *MSH2* and *MSH6*-deficient cells are more susceptible to oxaliplatin, despite

MMR-deficiency being associated with cisplatin resistance[33]. The autoimmune disease-

associated gene *SIAE*, which has been previously shown to have a strong negative

correlation to oxaliplatin response in advanced CRC patients[39], was selected in the

majority of thresholded oxaliplatin models (Supplementary Table S2C). The gene *BCL2*,

which was commonly selected for cisplatin (Figure 3A), was rarely selected for

oxaliplatin (Figure 3C). At the highest levels of resistance to cisplatin, models were

enriched for genes belonging to DNA repair, anti-oxidative response, apoptotic pathways

and drug transporters (Figure 3A). These gene pathways are known to be involved in

cisplatin resistance[40,41] and these specific genes may be explored in subsequent work to

identify the contribution to chemotherapy response in a biochemical context.

It may be feasible to predict responses to combination chemotherapy with the

models described here. Not included in the present analysis were signatures for

methotrexate, vinblastine, and doxorubicin, which comprise the MVAC cocktail used to

treat bladder cancer. This was due primarily to a lack of patients treated with this drug

combination in the TCGA bladder dataset (N=11). Individual signatures for several of

these drugs have been derived and analyzed using the patient data from METABRIC

(Molecular Taxonomy of Breast Cancer International Consortium)[24]. A reasonable

approach to predicting combination chemotherapy would first determine the probability

of sensitivity or resistance to individual drugs, accounting for the misclassification rate by

each (defined as d_1, …, d_k). The ML classifiers output these probabilities, analogous to

their misclassification rates in a set of patients treated identically. If the model predicts

that the patient is sensitive to drug d_1 with 90% probability, and sensitive to drug d_2 with

5% probability, the probability of sensitivity to the combination is 1 - (1 - 0.9)*(1 - 0.05) =

90.5%, and the probability of resistance is 9.5%. The correlated responses could be

estimated for drug pairs, d_1 and d_2, and then adjusted for the combined probability of the

pair to d_{12}, based on the features that are shared by the signatures of both drugs. The

probability of sensitivity would then be given by 1 - (1 - d_{12})*(1 - d_3)*…*(1 - d_k).

The predictive accuracy for the same model could differentiate highly between

the two datasets. **Cis3** (Supplemental Table S5A) had a high predictive accuracy and

AUC for TCGA bladder cancer patients (AUC=0.64). However, the AUC was lower when

applied to the Als *et al.*[28] dataset (AUC=0.18). Patient metadata in the latter study only

indicated patient survival times, while we base the expected TCGA patient outcome on

time to disease recurrence. As the basis of our expected outcome differs between

datasets, these differences may be acting as a confounding factor to determine accuracy

of gene signatures. The datasets also differ in how expression was measured

(microarray vs. RNA-seq). The relevance of models based on training and testing data

from different platforms can affect the accuracy of validation, which might not be

improved by data normalization. In this study, datasets were subjected to z-score

normalization. In subsequent studies, other techniques to correct for some of these

effects have been described and could be applied[42].

In summary, we describe GI_{50}- or IC_{50}-threshold-independent ML models to

predict chemotherapy response to platin agents in cancer patients. Ensemble machine

learning produced combined signatures that were more accurate than most individual

models generated with different thresholds. Genes associated cisplatin response

included those which exacerbate resistance in patients with a history of smoking. The

methodology described here should be adaptable to other drugs and cancer types. With

a range of models for multiple drugs, it may be possible to improve the efficacy of

treatment by tailoring treatment to a patient's specific tumour biology, and reduce

treatment duration by limiting the number of different therapeutic regimens prescribed

before achieving a successful response[43].

MATERIALS AND METHODS

Data and preprocessing

Microarray GE and data were from breast cancer cell lines were used to train

ML-based gene signatures of drug response based on respective growth or target

inhibition data (GI_{50} or IC_{50}). Cell lines were treated with either cisplatin (N=39),

carboplatin (N=46), or oxaliplatin (N=47)[13]. Bladder cancer cell line GE and IC_{50}

measurements for cisplatin were obtained from cancerRxgene (N=17). However, all

testing was performed on breast cancer cell line data as the number of bladder cancer

cell lines was insufficient to produce accurate signatures. RNA-seq GE and survival

measurements were downloaded from TCGA for bladder urothelial carcinoma (N=72

patients treated with cisplatin)[25], ovarian epithelial tumor (N=410 treated with

carboplatin)[26] and colorectal adenocarcinoma (N=99 treated with oxaliplatin)[27]. GE of

cisplatin-treated patients of cell carcinoma of the urothelium (N=30)[28] and for oxaliplatin-

treated CRC patients (N=83)[29] were obtained from the Gene Expression Omnibus.

Clinical metadata and GE for TCGA patients were obtained from Genomic Data

Commons (https://gdc.cancer.gov/), while methylation HM450 (Illumina) data for these

patients was downloaded from cBioPortal[44].

Initial gene sets for developing signatures for each drug were identified from

previously published literature (see Supplemental References) and databases, such as

PharmGKB and DrugBank[45,46]. Genes were then eliminated that showed no or little

expression in TCGA bladder cancer patients (where RNA-seq by Expectation

Maximization [RSEM] is < 5.0 for majority of individuals). The final gene sets were

chosen using MFA to analyze interactions between GE, CN, and GI_{50} data for the drug

of interest[47]. Genes whose GE and/or CN showed a direct or inverse correlation with

GI_{50} were selected for SVM training. This reduces the overall number of genes for SVM

analysis, and thus helps to avoid a data to size sample imbalance. For cisplatin, MFA

was repeated using IC_{50} values for 17 bladder cancer cell lines; however, the available

CN data generally showed a lack of variation in the cell lines for these genes. Instead,

the available IC_{50} values for three other cancer drugs (doxorubicin, methotrexate and

vinblastine) were compared with the IC_{50} of cisplatin by MFA.

Applying an SVM model directly to patient data without a normalization approach

is imprecise when training and testing data are not obtained using similar methodology

(i.e. different microarray platforms). To compare the cell line GE microarray data and the

patient RNA-seq GE datasets, expression values were normalized by conversion to z-

scores using MATLAB[48]. Although Log2 intensity values from microarray data were not

available for TCGA samples, RNA-seq based GE and log_2 intensities from microarray

data are highly correlated[49].

Machine Learning

SVM were trained with breast cancer cell line GE datasets[13] with the Statistics

Toolbox in MATLAB[48] using Gaussian kernel functions (fitcsvm), and then tested with a

leave-one-out cross-validation (using '*crossval'* and '*leaveout'* options). A greedy

backwards feature selection algorithm was used to improve classification accuracy[50].

The gene subset with the lowest misclassification with cross-validation is selected as the

model for subsequent testing with patient GE and clinical data.

Derivation of gene signatures for different drug resistance thresholds

We have previously set a conventional GI_{50} threshold distinguishing sensitivity

from resistance at the *median* of the range of drug concentrations that inhibited cell

growth by 50%[6]. We hypothesized that different gene signatures could be derived for

different levels of drug resistance by varying this threshold. ML experiments for

classifying resistance or sensitivity at GI_{50} values generated a series of optimized

Gaussian SVM models whose performance were assessed with patient expression data

for each signature. A heat map which illustrates the frequencies of genes appearing in

these models was created with the R language *hist2d* function.

A composite gene signature was created by ensemble averaging of all models

generated at each resistance threshold. Ensemble averaging combines signatures

through averaging the weighted accuracy of a set of related models[37]. The decision

function for the ensemble classifier is the mean of the decision function scores of the

component classifiers, weighted by the AUC.

Randomization

The potential for models to overfit data during training and/or feature selection

was assessed by permutation analysis with randomized cell line labels and with random

sets of genes, as described previously[6]. Using the median cisplatin GI_{50} as the

resistance threshold, 10,000 models based on random gene selection (15 genes) had

higher rates of misclassification than the best median SVM models (2 signatures with

7.7% misclassification). Cisplatin, carboplatin and oxaliplatin GE data for random cell

line label combinations (n=10,000) generated only 8, 1 and 1 signatures, respectively,

with lower error rates than the best biochemically-inspired signatures.

ACKNOWLEDGEMENTS

Katherina Baranova contributed to the cisplatin gene signatures and Dimo Angelov

developed automated feature selection. We thank Murray Junop for commenting on the

manuscript. Compute Canada and Shared Hierarchical Academic Research Computing

Network (SHARCNET) provided high performance computing and storage facilities.

CONFLICTS OF INTEREST

PKR cofounded CytoGnomix Inc., which hosts the interactive resource described in this

study for prediction of responses to chemotherapy agents. The other authors have no

conflicts of interest.

AUTHOR CONTRIBUTIONS

PKR and DL designed the methodology. EJM and JZ performed analyses. EJM and

PKR wrote the manuscript.

FUNDING

PKR is supported by NSERC (RGPIN-2015-06290), Canadian Foundation for

Innovation, Canada Research Chairs, and CytoGnomix.

REFERENCES

1. Cardoso, F. *et al.* Locally recurrent or metastatic breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann. Oncol.* **23,** vii11– vii19 (2012).

2. Oostendorp, L. J., Stalmeier, P. F., Donders, A. R. T., van der Graaf, W. T. & Ottevanger, P. B. Efficacy and safety of palliative chemotherapy for patients with advanced breast cancer pretreated with anthracyclines and taxanes: a systematic review. *Lancet Oncol.* **12,** 1053–1061 (2011).

3. Alfarouk, K. O. *et al.* Resistance to cancer chemotherapy: failure in drug response from ADME to P-gp. *Cancer Cell Int.* **15,** 71 (2015).

4. Gąsowska-Bodnar, A. *et al.* Survivin Expression as a Prognostic Factor in Patients With Epithelial Ovarian Cancer or Primary Peritoneal Cancer Treated With Neoadjuvant Chemotherapy: *Int. J. Gynecol. Cancer* **24,** 687–696 (2014).

5. Hatzis, C. *et al.* A genomic predictor of response and survival following taxane-anthracycline chemotherapy for invasive breast cancer. *JAMA* **305,** 1873–1881 (2011).

6. Dorman, S. N. *et al.* Genomic signatures for paclitaxel and gemcitabine resistance in breast cancer derived by machine learning. *Mol. Oncol.* **10,** 85–100 (2016).

7. Zhang, S. *et al.* Organic Cation Transporters Are Determinants of Oxaliplatin Cytotoxicity. *Cancer Res.* **66,** 8847–8857 (2006).

8. Poisson, L. M. *et al.* A metabolomic approach to identifying platinum resistance in ovarian cancer. *J. Ovarian Res.* **8,** (2015).

9. Cadoná, F. C. *et al.* Guaraná a Caffeine-Rich Food Increases Oxaliplatin Sensitivity of Colorectal HT-29 Cells by Apoptosis Pathway Modulation. *Anticancer Agents Med. Chem.* **16,** 1055–1065 (2016).

10. Kasparkova, J., Vojtiskova, M., Natile, G. & Brabec, V. Unique Properties of DNA

Interstrand Cross-Links of Antitumor Oxaliplatin and the Effect of Chirality of the

Carrier Ligand. *Chem. – Eur. J.* **14,** 1330–1341 (2008).

11. Woynarowski, J. M. *et al.* Oxaliplatin-Induced Damage of Cellular DNA. *Mol.*

Pharmacol. **58,** 920–927 (2000).

12. Tashiro, T., Kawada, Y., Sakurai, Y. & Kidani, Y. Antitumor activity of a new platinum

complex, oxalato (trans-l-1,2-diaminocyclohexane)platinum (II): new experimental

data. *Biomed. Pharmacother.* **43,** 251–260 (1989).

13. Daemen, A. *et al.* Modeling precision treatment of breast cancer. *Genome Biol.* **14,**

R110 (2013).

14. Yuan, Y. *et al.* Identification of the biomarkers for the prediction of efficacy in first-line

chemotherapy of metastatic colorectal cancer patients using SELDI-TOF-MS and

artificial neural networks. *Hepatogastroenterology.* **59,** 2461–2465 (2012).

15. L'Espérance, S., Bachvarova, M., Tetu, B., Mes-Masson, A.-M. & Bachvarov, D.

Global gene expression analysis of early response to chemotherapy treatment in

ovarian cancer spheroids. *BMC Genomics* **9,** 99 (2008).

16. Nickerson, M. L. *et al.* Molecular analysis of urothelial cancer cell lines for modeling

tumor biology and drug response. *Oncogene* (2016). doi:10.1038/onc.2016.172

17. Yuryev, A. Gene expression profiling for targeted cancer treatment. *Expert Opin.*

Drug Discov. **10,** 91–99 (2015).

18. Sataloff, D. M. *et al.* Pathologic response to induction chemotherapy in locally

advanced carcinoma of the breast: a determinant of outcome. *J. Am. Coll. Surg.* **180,**

297–306 (1995).

19. Ogston, K. N. *et al.* A new histological grading system to assess response of breast

cancers to primary chemotherapy: prognostic significance and survival. *Breast*

Edinb. Scotl. **12,** 320–327 (2003).

20. Earl, J. *et al.* The UBC-40 Urothelial Bladder Cancer cell line index: a genomic

resource for functional studies. *BMC Genomics* **16,** 403 (2015).

21. Rixe, O. *et al.* Oxaliplatin, tetraplatin, cisplatin, and carboplatin: Spectrum of activity

in drug-resistant cell lines and in the cell lines of the national cancer institute's

anticancer drug screen panel. *Biochem. Pharmacol.* **52,** 1855–1865 (1996).

22. Mehmood, R. K. Review of Cisplatin and oxaliplatin in current immunogenic and

monoclonal antibody treatments. *Oncol. Rev.* **8,** 256 (2014).

23. Kweekel, D. M., Gelderblom, H. & Guchelaar, H.-J. Pharmacology of oxaliplatin and

the use of pharmacogenomics to individualize therapy. *Cancer Treat. Rev.* **31,** 90–

105 (2005).

24. Mucaki, E. J. *et al.* Predicting Outcomes of Hormone and Chemotherapy in the

Molecular Taxonomy of Breast Cancer International Consortium (METABRIC) Study

by Biochemically-inspired Machine Learning. *F1000Research* **5,** 2124 (2017).

25. Robertson, A. G. *et al.* Comprehensive Molecular Characterization of Muscle-

Invasive Bladder Cancer. *Cell* **171,** 540-556.e25 (2017).

26. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian

carcinoma. *Nature* **474,** 609–615 (2011).

27. Cancer Genome Atlas Network. Comprehensive molecular characterization of

human colon and rectal cancer. *Nature* **487,** 330–337 (2012).

28. Als, A. B. *et al.* Emmprin and survivin predict response and survival following

cisplatin-containing chemotherapy in patients with advanced bladder cancer. *Clin.*

Cancer Res. Off. J. Am. Assoc. Cancer Res. **13,** 4407–4414 (2007).

29. Tsuji, S. *et al.* Potential responders to FOLFOX therapy for colorectal cancer by

Random Forests analysis. *Br. J. Cancer* **106,** 126–132 (2012).

30. Tembe, V. *et al.* The BARD1 BRCT domain contributes to p53 binding, cytoplasmic

and mitochondrial localization, and apoptotic function. *Cell. Signal.* **27,** 1763–1771

(2015).

31. Freedman, N. D., Silverman, D. T., Hollenbeck, A. R., Schatzkin, A. & Abnet, C. C.

Association between smoking and risk of bladder cancer among men and women.

JAMA **306,** 737–745 (2011).

32. Joehanes, R. *et al.* Epigenetic Signatures of Cigarette Smoking. *Circ. Cardiovasc.*

Genet. (2016). doi:10.1161/CIRCGENETICS.116.001506

33. Raymond, E., Faivre, S., Chaney, S., Woynarowski, J. & Cvitkovic, E. Cellular and

Molecular Pharmacology of Oxaliplatin1. *Mol. Cancer Ther.* **1,** 227–235 (2002).

34. Alex, A. K. *et al.* Response to Chemotherapy and Prognosis in Metastatic Colorectal

Cancer With DNA Deficient Mismatch Repair. *Clin. Colorectal Cancer* (2016).

doi:10.1016/j.clcc.2016.11.001

35. Sos, M. L. *et al.* Predicting drug susceptibility of non-small cell lung cancers based

on genetic lesions. *J. Clin. Invest.* **119,** 1727–1740 (2009).

36. Laderas, T. G., Heiser, L. M. & Sönmez, K. A Network-Based Model of Oncogenic

Collaboration for Prediction of Drug Sensitivity. *Front. Genet.* **6,** (2015).

37. Clemen, R. T. Combining forecasts: A review and annotated bibliography. *Int. J.*

Forecast. **5,** 559–583 (1989).

38. Airley, R. *Cancer chemotherapy*. (Wiley-Blackwell, 2009).

39. Li, X.-X. *et al.* RNA-seq identifies determinants of oxaliplatin sensitivity in colorectal

cancer cell lines. *Int. J. Clin. Exp. Pathol.* **7,** 3763–3770 (2014).

40. Borst, P., Rottenberg, S. & Jonkers, J. How do real tumors become resistant to

cisplatin? *Cell Cycle Georget. Tex* **7,** 1353–1359 (2008).

41. Wernyj, R. & Morin, P. Molecular mechanisms of platinum resistance: still searching

for the Achilles heel. *Drug Resist. Updat.* **7,** 227–232 (2004).

42. Johnson, W. E., Li, C. & Rabinovic, A. Adjusting batch effects in microarray

expression data using empirical Bayes methods. *Biostat. Oxf. Engl.* **8,** 118–127

(2007).

43. Akamatsu, N., Nakajima, H., Ono, M. & Miura, Y. Increase in acetyl CoA synthetase

activity after phenobarbital treatment. *Biochem. Pharmacol.* **24,** 1725–1727 (1975).

44. Gao, J. *et al.* Integrative analysis of complex cancer genomics and clinical profiles

using the cBioPortal. *Sci. Signal.* **6,** pl1 (2013).

45. Whirl-Carrillo, M. *et al.* Pharmacogenomics Knowledge for Personalized Medicine.

Clin. Pharmacol. Ther. **92,** 414–417 (2012).

46. Law, V. *et al.* DrugBank 4.0: shedding new light on drug metabolism. *Nucleic Acids*

Res. **42,** D1091–D1097 (2014).

47. Abdi, H. & Williams, L. J. Principal component analysis. *Wiley Interdiscip. Rev.*

Comput. Stat. **2,** 433–459 (2010).

48. MATLAB and Statistics Toolbox Release 2012b, The MathWorks, Inc., Natick,

Massachusetts, United States.

49. Marioni, J. C., Mason, C. E., Mane, S. M., Stephens, M. & Gilad, Y. RNA-seq: an

assessment of technical reproducibility and comparison with gene expression arrays.

Genome Res. **18,** 1509–1517 (2008).

50. Bermingham, M. L. *et al.* Application of high-dimensional feature selection:

evaluation for genomic prediction in man. *Sci. Rep.* **5,** 10312 (2015).

 FIGURE LEGENDS

 Figure 1. Schematic of platinum drug sensitivity and resistance genes which showed

 MFA correlation for GI_{50} of A) cisplatin, B) carboplatin, and C) oxaliplatin. The genes

 used to derive the SVM are shown in context of their effect in the cell and role in cisplatin

 mechanisms of action. GE and CN correlation with inhibitory drug concentration by MFA

 of breast (GI_{50}) and bladder (IC_{50}) cancer cell line data.

 Figure 2: GI_{50} values for cell lines treated with the three platin drugs were plotted in

 order of ascending oxaliplatin GI_{50}. For most cell lines, there is a visible trend between

 the GI_{50} for cisplatin and carboplatin, reflecting the correlation between the two drugs

 seen by MFA. Despite this correlation, carboplatin shows a much smaller variance (0.22)

 compared to cisplatin (0.37; oxaliplatin variance is 0.34).

 Figure 3. The variation in gene composition of SVMs at different GI_{50} thresholds for A)

 cisplatin, B) carboplatin, and C) oxaliplatin. GI_{50} intervals are indicated on the left, with

 the number of cell lines with GI_{50} values within said intervals in brackets. Each box

 represents the density of genes appearing in optimized Gaussian SVM models in those

 functional categories, with darker grey indicating frequent genes in indicated GI_{50}

 threshold intervals, while lighter grey indicates less commonly selected genes. The

 number of thresholded models used to derive the density plot within each interval is

 equal (or greater, in the case of multiple equally performing models) to the number of

 cell lines within that GI_{50} interval.

 Supplementary Figure 1. Classification accuracy of models on TCGA bladder cancer

 patients treated with cisplatin and/or carboplatin as the resistance threshold is varied.

 Recurrence and disease-free survival are used as a binary measure to assess

 performance. The x-axis indicates movement of the resistance threshold, with more cell

lines labeled sensitive on the left and more labeled resistant on the right. Maximal AUC

is indicated by the downward arrows.

Supplementary Figure 2. Classification accuracy of SVM models for cisplatin, at a

range of response thresholds, were assessed using gene expression data for cisplatin-

treated bladder cancer patients from Als *et al.* [28]. Patients with a ≥ 5 year survival post-

treatment were labeled sensitive. Red arrows indicate the SVM models with the highest

positive predictive value (PPV) in the accuracy of classification of patient outcome.

Supplementary Figure 3. Hyperplane distance calculated by all thresholded SVMs for

recurrent (<6 months) TCGA patients. Each diagram represents the predictions of all

SVMs for all patients who had recurrence less than 6 months after treatment (N=10).

Each point represents an SVM, where the x-axis represents the number of cell lines set

to resistant (in order of lowest to highest GI_{50}), and the y-axis represents the calculated

hyperplane distance. A negative hyperplane distance would represent a prediction of

resistance to cisplatin. Despite this, some patients show a strong preference towards

predictions of sensitivity (i.e. TCGA-XF-A9SU).

Supplementary Figure 4. Hyperplane distance calculated by all thresholded SVMs for

sensitive TCGA patients. Each diagram represents the predictions of all SVMs for all

patients who had recurrence > 4 years after treatment (top; N=3), or patients who

showed no recurrence after 6 years (bottom; N=6). Each point represents an SVM,

where the x-axis represents the number of cell lines set to resistant (in order of lowest to

highest GI_{50}), and the y-axis represents the calculated hyperplane distance. A positive

hyperplane distance would represent a prediction of sensitivity to cisplatin.

a.

b.

C.

Resistant

GI$_{50}$
(# cell lines in interval)

Apoptosis • Immune System • Metabolism • Signal Transduction • Transcription / DNA Repair • Membrane Transport • Uncharacterized

3.50 – 4.50 (4)
4.51 – 4.80 (7)
4.81 – 5.10 (7)
5.11 – 5.40 (7)
5.41 – 5.70 (14)
5.71 – 6.00 (4)

BCL2 • CSMD1 • ICAM5 • FCGR2A • HLA-B • LTA • MPO • SIAE • PRSS1 • AGXT • NAT2 • NQO1 • PANK3 • SGPP2 • UGT1A1 • NAGK • PROC • APOBEC2 • KIT • KLC3 • BRAF • IGF1R • KISS1 • NOTCH1 • IGF1 • KLF3 • HIF1A • NFE2L2 • PARP15 • MSH2 • SLC47A1 • CLCN6 • ABOG2 • SLCO1B1 • GRTP1

Sensitive

0% 20% 40% 60% 80% 100%

Frequency of Genes Selected in SVM for GI$_{50}$ interval