Trait paranoia shapes inter-subject synchrony in brain activity during an ambiguous

social narrative

Emily S. Finn*[1], Philip R. Corlett[2], Gang Chen[3], Peter A. Bandettini[1], R. Todd Constable[4]

[1]Section on Functional Imaging Methods, Laboratory of Brain and Cognition, National Institute

of Mental Health; Bethesda, Md.

[2]Department of Psychiatry, Yale School of Medicine; New Haven, Conn.

[3]Scientific and Statistical Computing Core, National Institute of Mental Health; Bethesda, Md.

[4]Department of Radiology and Biomedical Imaging, Yale School of Medicine, New Haven, Conn.

*Corresponding author: emily.finn@nih.gov

ABSTRACT

Individuals often interpret the same event in different ways. How do personality traits modulate

brain activity evoked by a complex stimulus? Here we report results from a naturalistic paradigm

designed to draw out both neural and behavioral variation along a specific dimension of interest,

namely paranoia. Participants listen to a narrative during functional MRI describing an ambiguous

social scenario, written such that some individuals would find it highly suspicious, while others

less so. Using inter-subject correlation analysis, we identify several brain areas that are

differentially synchronized during listening between participants with high- and low trait-level

paranoia, including theory-of-mind regions. Follow-up analyses indicate that these regions are

more active to mentalizing events in high-paranoia individuals. Analyzing participants' speech as

they freely recall the narrative reveals semantic and syntactic features that also scale with paranoia.

Results indicate that a personality trait can act as an intrinsic 'prime', yielding different neural and

behavioral responses to the same stimulus across individuals.

That different individuals may see the same event in different ways is a truism of human

nature. Examples are found at many scales, from low-level perceptual judgments to interpretations

of complex, extended scenarios. This latter phenomenon is known as the "Rashomon effect"[1] after

a 1950 Japanese film in which four eyewitnesses give contradictory accounts of a crime and its

aftermath, raising the point that for multifaceted, emotionally charged events, there may be no

single version of the truth.

What accounts for these individual differences in interpretation? Assuming everyone has

access to the same perceptual information, personality traits may bias different individuals toward

one interpretation or another. Paranoia is one such trait, in that individuals with strong paranoid

tendencies may be more likely to assign a nefarious interpretation to otherwise neutral events[2].

While paranoia in its extreme is a hallmark symptom of schizophrenia and other psychoses, trait-

level paranoia exists as a continuum rather than a dichotomy[3,4]: on a behavioral level, up to 30

percent of people report experiencing certain types of paranoid thoughts (e.g., 'I need to be on my

guard against others') on a regular basis[5] and trait paranoia in the population follows an

exponential, rather than bimodal, distribution[6].

Few neuroimaging studies have investigated paranoia as a continuum; the majority simply

contrast healthy controls and patients suffering from clinical delusions. However, a handful of

reports from subclinical populations describe patterns of brain activity that scale parametrically

with tendency toward paranoid or delusional ideation. For example, it has been reported that

higher-paranoia individuals show less activity in the medial temporal lobe during memory retrieval

and less activity in the cerebellum during sentence completion[7], less activity in temporal regions

during social reflection[8] and auditory oddball detection[9], but higher activity in the insula and

medial prefrontal cortex during self-referential processing[10] and differential patterns of activity in

these regions as well as the amygdala while viewing emotional pictures[11].

Such highly controlled paradigms enable precise inferences about evoked brain activity,

but potentially at the expense of real-world validity. For example, brain response to social threat

is often assessed with decontextualized static photographs of unfamiliar faces presented rapidly in

series[12]. Compare this to threat detection in the real world, which involves perceiving and

interacting with both familiar and unfamiliar faces in a rich, dynamic social context. Paranoid

thoughts that eventually reach clinical significance usually have a slow, insidious onset, involving

complex interplay between a person's intrinsic tendencies and his or her experiences in the world.

In studying paranoia and other trait-level individual differences, then, is important to complement

highly controlled paradigms with more naturalistic stimuli.

Narrative is an attractive paradigm for several reasons. First, narrative is an ecologically

valid way to study belief formation in action. Theories of fiction posit that readers model narratives

in a Bayesian framework in much the same way as real-world information[13], and story

comprehension and theory-of-mind processes share overlapping neural resources[14]. Second, a

standardized narrative stimulus provides identical input, so any variation in interpretation reflects

individuals' intrinsic biases in how they assign salience, learn and form beliefs. Third, from a

neuroimaging perspective, narrative listening is a continuous, engaging task that involves much of

the brain[15] and yields data lending itself to innovative, data-driven analyses such as inter-subject

correlation[16,17].

Previous work has shown that experimenters can manipulate patterns of brain activity

during naturalistic stimuli by explicitly instructing participants to focus on different aspects of the

stimulus. For example, Cooper et al. reported that activity patterns in temporal and frontal regions

varied according to whether listeners were told to pay attention to action-, space- or time-related

features of short stories[18]. Lahnakoski et al. showed participants the same movie twice, asking

them to adopt different perspectives each time, and found differences in neural synchrony

depending on which perspective had been taken[19]. Most recently, Yeshurun et al. presented

participants with a highly ambiguous story with at least two plausible—but very different—

interpretations, and used explicit primes to bias each participant toward one interpretation or the

other. Responses in higher-order brain areas, including default mode, were more similar among

participants who had received the same prime, indicating that shared beliefs have a powerful effect

on how individuals perceive an identical stimulus[20]. However, while informative, these studies

have all relied on an explicit prime or instruction; they cannot explain why individuals often

spontaneously arrive at different interpretations of the same stimulus.

In this work, we use participants' intrinsic personality traits as an implicit prime, relating

individual differences in trait paranoia to brain activity during a naturalistic task in which

participants are faced with complex, ambiguous social circumstances. Using an original narrative,

we show that while much of the brain is synchronized across all participants during story listening,

stratifying participants based on trait paranoia reveals an additional set of regions with stereotyped

activity only among high-paranoia individuals; many of these are regions involved in theory-of-

mind and mentalizing. An encoding model of the task suggests that these regions, including the

temporal pole and medial prefrontal cortex, are particularly sensitive to "mentalizing events" when

the main character is experiencing an ambiguous social interaction or explicitly reasoning about

other characters' intentions. Finally, we measure participants' behavioral reactions to the narrative

by analyzing their speech as they freely recall the story, and identify semantic and syntactic

features that vary dimensionally with trait paranoia. Together, results indicate that a personality

trait, in this case paranoia, can modulate both neural and behavioral responses to a single stimulus

across individuals.

RESULTS

Behavioral data and task performance

We created a fictional narrative to serve as the stimulus for this study. The narrative

described a main character faced with a complex social scenario that was deliberately ambiguous

with respect to the intentions of certain characters; it was designed such that different individuals

would interpret the events as more nefarious and others as less so. A synopsis of the story is given

in Supplementary Note 1.

Twenty-two healthy participants listened to a pre-recorded audio version of the narrative

(total duration = 21:50 min:sec, divided into three parts) during fMRI scanning. Following each

of the three parts, participants answered three challenging multiple-choice comprehension

questions to ensure they had been paying attention. Performance was very accurate (15 of the 22

subjects answered 9/9 [100%] questions correctly, while five answered 8/9 [89%] correctly and

two answered 7/9 [78%] correctly). Self-report data indicated that subjects generally found the

narrative engaging and easy to pay attention to (engagement rating on a scale of 1 to 5: mean =

3.8, s.d. = 0.96, median = 4, median absolute deviation [m.a.d.] = 0.72; attention rating: mean =

4.1, s.d. = 0.87, median = 4, m.a.d. = 0.66).

During a separate behavioral visit one week prior to the scan, participants completed

several self-report questionnaires and behavioral tasks to assess personality traits and cognitive

abilities (see Fig. 1a for a schematic of the experimental protocol). Our primary measure of interest

was subscale A from the Green et al. Paranoid Thoughts Scale[21] (GPTS-A), henceforth referred to

as trait paranoia score. We administered this scale on a different day, and placed it amongst other

tasks unrelated to paranoia, to minimize any priming effects or demand characteristics that might

influence participants' eventual reactions to the narrative. Possible scores on the GPTS-A range

from 16 to 80; higher scores are generally observed only in clinical populations[21]. In our healthy

sample, we observed a right-skewed distribution that nonetheless had some variance (range = 16-

40, mean = 20.6, s.d. = 6.3; median = 18.5, m.a.d. = 4.0; see Fig. 1b for a histogram of the

distribution). This is consistent with observations from much larger sample sizes that trait paranoia

follows an exponential, rather than normal, distribution in the healthy population[5,6,21].

Story listening evokes widespread neural synchrony

Our primary approach for analyzing the fMRI data was inter-subject correlation (ISC),

which is a model-free way to identify brain regions responding reliably to a naturalistic stimulus

across subjects[16,17]. In this approach, the timecourse from each voxel in one subject's brain across

the duration of the stimulus is correlated with the timecourse of the same voxel in a second

subject's brain. Voxels that show high correlations in their timecourses across subjects are

considered to have a stereotyped functional role in processing the stimulus. The advantage of this

approach is that it does not require the investigator to have an *a priori* model of the task, nor to

assume any fixed hemodynamic response function.

In a first-pass analysis, we calculated ISC at each voxel across the whole sample of n = 22

participants, using a recently developed statistical approach that relies on a linear mixed-effects

model with crossed random effects to appropriately account for the correlation structure of the

data[22]. Results are shown in Fig. 2. As expected given the audio-linguistic nature of the stimulus,

ISC was highest in primary auditory cortex and language regions along the superior temporal lobe,

but we also observed widespread ISC in other parts of association cortex, including frontal,

parietal, midline and temporal areas, as well as the posterior cerebellum. These results replicate

previous reports that complex naturalistic stimuli induce stereotyped responses across participants

in not only the relevant primary cortex, but also higher-order brain regions[15,16,23].

Also as expected, ISC was generally lower or absent in primary motor and somatosensory

cortex, although we did observe significant ISC in parts of primary visual cortex, despite the fact

that there was no timecourse of visual input during the story. (To encourage engagement, we had

participants fixate on a static photograph that was thematically relevant to the story during

listening, so the observed ISC in visual cortex may reflect similarities in the timecourse of

internally generated imagery across participants.)

Paranoia modulates neural response to the narrative

Having established that story listening evokes widespread neural synchrony across all

participants, we next sought to determine if there were brain regions whose degree of ISC was

modulated by trait paranoia. Using a median split of GPTS-A scores, we stratified our sample into

a low-paranoia (GPTS-A $\leq$ 18, n = 11) and high-paranoia (GPTS-A $\geq$ 19, n = 11) group (Fig. 1b).

We then used the same linear mixed-effects model described above formulated as a two-group

contrast to reveal areas that are differentially synchronized across paranoia levels.

We opted for a median split rather than using raw paranoia score as a continuous covariate

because of the unique challenge of an ISC-based analysis, which, to take advantage of all the

information contained in the cross-subject correlation matrix (Fig. 1c), requires any covariates to

be at the subject *pair* level, rather than the level of individual subjects. Because trait paranoia is a

single scalar value per participant, it is difficult to calculate a meaningful pairwise metric. (Median

splits can also mitigate the influence of extreme values, such as the two participants with GPTS-

A $\geq$ 38 [cf. Fig. 1b], ensuring these do not have an outsize effect on the results.) Still, we conducted

post-hoc tests to investigate continuous relationships with raw GPTS-A score whenever possible

to respect the inherently continuous nature of this trait, and to facilitate interpretation.

We were primarily interested in three contrasts. First, which voxels show greater ISC

among pairs of high-paranoia participants versus low-paranoia participants, or vice versa? Second

and third, which voxels show greater ISC among pairs of low- or high-paranoia participants,

respectively (i.e., low-low or high-high), than pairs of participants mismatched for group (i.e.,

high-low)? All three contrasts reveals regions whose response timecourses are modulated by trait

paranoia in some way. These contrasts are schematized in Fig. 1c.

Results are shown in Fig. 3. In the first contrast, several regions emerged as being more

synchronized in the high-paranoia group relative to the low-paranoia group. Significant clusters

were found in the left temporal pole (Talairach coordinates for center of mass: [+46.7, -10.0, -

26.2]), left precuneus ([+10.8, +71.0, +35.9]), and two regions of the right medial prefrontal cortex

(mPFC; one anterior [-8.1, -46.9, +16.3] and one dorsal [+2.9, -14.8, +45.1]; Fig. 3a). Searches for

these coordinates on Neurosynth, an automated fMRI results synthesizer for mapping between

neural and cognitive states[24], indicated that for the left temporal pole and right anterior mPFC

clusters, top meta-analysis terms included "mentalizing", "mental states", "intentions", and

"theory mind". There were no regions showing a statistically significant difference in the reverse

direction (low-paranoia > high-paranoia).

In the second contrast (Fig. 3b, cool colors), pairs of low-paranoia participants were more

synchronized than pairs of inter-group participants in the left lateral occipital gyrus (center of mass:

[+31.3, +86.1, +14.0], Neurosynth: "objects", "scene", "encoding"), and in the third contrast (Fig.

3b, warm colors), pairs of high-paranoia participants were more synchronized than pairs of inter-

group participants in the right angular gyrus ([-44.8, +57.9, +37.9], Neurosynth: "beliefs").

Interestingly, there were no voxels of statistically significant overlap between the second and third

contrasts, indicating that no single region had a timecourse that was equally synchronized within

groups but qualitatively different between groups. Instead, for most of the regions that emerged

from the three contrasts, the relationship between trait paranoia and timecourse synchrony is best

expressed by the Anna Karenina principle: all paranoid participants are alike; all not-paranoid

participants are not-paranoid in their own way (except in the lateral occipital gyrus, where it is the

opposite).

As these regions were obtained via dichotomization into groups, we also conducted post-

hoc tests to determine if ISC remained sensitive to finer-grained differences in trait paranoia. We

were primarily interested in two regions that emerged from the first contrast, the left temporal pole

and right medial PFC, since these are known from prior literature to be involved in theory of mind

and mentalizing. To determine whether ISC in these regions scales monotonically with trait

paranoia, we visualized the participant-by-participant ISC matrices with participants ordered by

trait paranoia score (Fig. 4a. 4c). Visual inspection suggests a relatively continuous increase in ISC

values as one moves down and to the right along the diagonal, which represents pairs of

increasingly high-paranoia participants. To quantify this, we plotted each participant's median ISC

with all other participants (i.e., the median of each row of the ISC matrix) against their paranoia

rank within the sample (i.e., 1–22; Fig. 4b and d). For both ROIs, participants with higher paranoia

rank tended to have higher median ISC ($r_s = 0.71$ and $r_s = 0.63$ for the left temporal pole and right

medial PFC, respectively; both $p < 0.002$). We used paranoia rank rather than raw score to mitigate

the influence of the two participants with extreme paranoia scores (≥ 38; cf. Fig. 1b).

Effects are specific to paranoia

We conducted several control analyses to rule out the possibility that the observed group differences were driven by a factor other than trait paranoia. (For all analyses in this section, we checked for both categorical and continuous relationships with paranoia; full results are reported in Table 1.)

For example, if the high-paranoia participants have better overall attentional and cognitive abilities, they might simply be paying closer attention to the story, inflating ISC values but not necessarily because of selective attention to ambiguous or suspicious details. However, there were no differences between high- and low-paranoia participants on any of the cognitive tasks we administered (verbal IQ, vocabulary, fluid intelligence or working memory), making it unlikely that observed differences are due to trait-level differences in attention or cognition. As for state-level attention during the story, there was no relationship between paranoia and number of comprehension questions answered correctly, total word count during the recall task, or self-report measures of engagement and attention. We also explored potential imaging-based confounds, and found that paranoia was not related to amount of head motion during the scan (as measured by mean framewise displacement), number of censored frames, or temporal signal-to-noise ratio (tSNR). Paranoia groups did not differ in age or sex breakdown. Thus we are reasonably confident that the observed effects are driven by true trait-level differences in paranoia between individuals.

Activity to mentalizing events scales with paranoia

Results of the first contrast from the two-group ISC analysis indicated that certain brain regions showed a more stereotyped response in high-paranoia versus low-paranoia individuals. What features of the narrative were driving activity in these regions? In theory, ISC allows for reverse correlation, in which peaks of activation in a given region's timecourse are used to recover the stimulus events that evoked them[16]. In practice, this is often difficult. Especially with narrative stimuli, in which structure is built up over relatively long timescales[15], it is challenging to pinpoint exactly which event—word, phrase, sentence—triggered an increase in BOLD activity.

Rather than rely on reverse correlation, a data-driven *decoding* approach, we took an *encoding* approach: we modeled events in the task that we hypothesized would stimulate differing interpretations across individuals, and evaluated the degree to which certain regions of interest (ROIs) responded to such events, using a general linear model (GLM) analysis. Specifically, we

labeled sentences in the story when the main character was experiencing an ambiguous (i.e., possibly suspicious) social interaction, and/or sentences when she was explicitly reasoning about the intentions of other characters. For brevity, we refer to these timepoints as "mentalizing events." In creating the regressor, all events were time-locked to the end of the last word of the labeled sentences, when participants are presumably evaluating information they just heard and integrating it into their situation model of the story.

We hypothesized that the two ROIs from the previous analysis known to be involved in theory-of-mind and mentalizing, the left temporal pole and right medial PFC, would be more active to mentalizing events in individuals with higher trait paranoia. We included two additional ROIs, the left temporo-parietal junction (TPJ) and left Heschl's gyrus, as a positive and negative control, respectively. We selected the left TPJ as a positive control because of its well-established role in theory-of-mind and mentalizing processes, and the fact that it emerged as highly synchronized across all participants (cf. Fig. 2) but did not show a group difference (cf. Fig. 3); thus we hypothesized that this region should respond to mentalizing events in all participants, regardless of trait paranoia. Conversely, left Heschl's gyrus (primary auditory cortex) should only respond to low-level acoustic properties of the stimulus and not show preferential activation to mentalizing events in either group or the sample as a whole. See Fig. 5a for ROI locations.

For each participant, we regressed the timecourse of each of these four ROIs against the mentalizing-events regressor and compared the resulting regression coefficients between groups (Fig. 5b). Compared to low-paranoia individuals, high-paranoia individuals showed stronger responses in both the left temporal pole (two-sample $t(20) = 2.71$, $p_{adj} = 0.014$) and right medial PFC ($t(20) = 3.36$, $p_{adj} = 0.007$). As hypothesized, responses in the left TPJ were strong across the whole sample (one-sample $t(21) = 8.73$, $p < 0.0001$), but there was no significant difference between groups in this ROI ($t(20) = 0.67$, $p_{adj} = 0.34$). Also as hypothesized, the sample as a whole did not show a significant response to these events in primary auditory cortex (one-sample $t(21) = 0.44$, $p = 0.66$), and there was no group difference ($t(20) = 0.47$, $p_{adj} = 0.34$).

To confirm that these results hold if paranoia is treated as a continuous variable, we conducted additional post-hoc tests in which we correlated participants' paranoia ranks and regression coefficients for all four ROIs (Fig. 4c). As expected, response to suspicious events was significantly related to paranoia rank in the left temporal pole ($r_s = 0.57$, $p = 0.005$) and right medial

PFC (r_s = 0.64, p = 0.001), but not in the left TPJ (r_s = -0.04, p = 0.86) or left Heschl's gyrus (r_s =

0.02, p = 0.95).

As an additional control, to check that this effect was specific to mentalizing events and

not just any sentence offset, we crated an inverse regressor comprising all *non*-mentalizing events

(i.e., by flipping the binary labels from the mentalizing-events regressor, such that all sentences

were labeled *except* those containing an ambiguous social interaction or explicit mentalizing as

described above). There were no differences between paranoia groups in any of the four ROIs in

response to non-mentalizing sentences (Fig. 4d), and no continuous relationships between

regression coefficient and paranoia rank (Fig. 4e). This indicates that trait paranoia is associated

with differential sensitivity of the left temporal pole and right medial PFC to not just any

information, but specifically to socially ambiguous information that presumably triggers theory-

of-mind processes.

Paranoia modulates behavioral response to the narrative

Having established that trait paranoia modulates individuals' brain responses to an

ambiguous narrative, we next sought to determine if this trait also modulates their behavioral

responses to the narrative. In other words, does trait-related (intrinsic) paranoia bear upon state-

related (stimulus-evoked) paranoia? If the observed differences in neural activity propagate up to

conscious perception and interpretation of the stimulus, then participants' subjective experiences

of the narrative should also bear a signature of trait paranoia.

Immediately following the scan, participants completed a post-narrative battery that

consisted of free-speech prompts followed by multiple-choice items to characterize their beliefs

and feelings about the story. For the first item, participants were asked to retell the story in as much

detail as they could remember, and their speech was recorded. Participants were allowed to speak

for as long as they wished on whatever aspects of the story they chose. Without guidance from the

experimenter, participants recalled the story in rich detail, speaking an average of 1,081 words

(range = 399-3,185, s.d. = 610).

Audio recordings of participants' speech were transcribed and submitted to the language

analysis software Linguistic Inquiry and Word Count[25] (LIWC). The output of LIWC is one vector

per participant describing the percentage of speech falling into various semantic and syntactic

categories. Example semantic categories are positive emotion ('love', 'nice'), money ('cash',

'owe'), and body ('hands', 'face'), while syntactic categories correspond to parts of speech such
as pronouns, adjectives and prepositions; there are 67 categories in total.

Using partial least-squares regression, we searched for relationships between speech
features and trait paranoia score. More than 72 percent of the variance in paranoia score could be
accounted for by the first component of speech features; the loadings of semantic and syntactic
categories for this component are visualized in Fig. 6a. The feature with the highest positive
loading—indicating a positive relationship with paranoia—was affiliation, a category of words
describing social and familial relationships (e.g., 'ally', 'friend', 'social'). Also associated with
high trait paranoia was frequent use of adjectives as well as anxiety- and risk-related words (e.g.,
'bad', 'crisis'); drives, a meta-category that includes words concerning affiliation, achievement,
power, reward and risk; and health-related words (e.g., 'clinic', 'fever', 'infected'; recall that the
story featured a doctor treating patients in a remote village; cf. Supplementary Note 1). Features
with strongly negative loadings—indicating an inverse relationship with paranoia—included male
references (e.g., 'him', 'his', 'man', 'father'); anger-related words ('yell', 'annoyed'); function
words ('it', 'from', 'so', 'with'); and conjunctions ('and', 'but', 'until'). Fig. 6b contains specific
examples for selected categories from participants' speech transcripts.

After the free-speech prompts, participants answered a series of multiple-choice questions
(see Supplementary Table 1 for the full questionnaire). First, they were asked to rate the degree to
which they were experiencing various emotions (suspicion, paranoia, sadness, happiness,
confusion, anxiety, etc; 16 in total) on a scale from 1 to 5. Most of ratings skewed low—for
example, the highest paranoia rating was 3, and only six subjects rated their paranoia level higher
than 1. Interestingly, there was no significant correlation between trait paranoia score and self-
reported paranoia ($r_s = -0.02$, $p = 0.91$) or suspicion ($r_s = 0.11$, $p = 0.62$) following the story. Neither
were any of the other emotion ratings significantly correlated with trait-level paranoia (all
uncorrected $p > 0.12$; see Fig. 6c).

Second, participants were asked to rate the three central characters on six personality
dimensions (trustworthy, impulsive, considerate, intelligent, likeable, naïve; see Supplementary
Fig. 1a). Third, they were asked to rate the likelihood of each of six scenarios (see Supplementary
Fig. 1b), and finally, to indicate (via forced-choice options) what they believed the main character
would do next, as well as what they themselves would do in her situation.

None of the individual questionnaire items significantly correlated with trait paranoia. However, to facilitate comparison with the speech data, we submitted the questionnaire data to a second partial least-squares regression to search for multidimensional relationships. This analysis revealed a first component of questionnaire responses that accounted for 62 percent of the variance in trait paranoia (Supplementary Fig. 1c). Features with the highest positive loadings, indicating a positive relationship with paranoia, included certain answers about what individuals thought the main character might do next as well as what they would do in her place (e.g., escape from the situation), as well as feeling more uncomfortable and suspicious following the story. Features with the highest negative loadings, indicating an inverse relationship with paranoia, included feeling more amused, inspired and hopeful following the story, as well a tendency to agree with one of the scenarios ("Juan and the other villagers had not known anything about the disease before Carmen arrived").

Overall, then, we found signatures of paranoia in story-evoked behavior using both free speech and self-report measures. Participants' free speech was slightly more sensitive than their answers on the multiple-choice questionnaire. Self-report is a coarse measure that may suffer from response bias; behavior provides a richer feature set that allows for the discovery of more subtle associations. In studying nuanced individual differences, then, these results highlight the desirability of capturing behavior in both traditional and naturalistic ways.

DISCUSSION

Here we have shown that a personality trait can act as a lens, or "implicit prime", through which individuals perceive ambiguous events, shaping both their neural and behavioral responses to an identical stimulus. Previous work using naturalistic tasks has shown that brain activity and behavioral responses are sensitive to experimenter instructions, i.e., an explicit prime[19,20], or to the nature of the stimulus itself, i.e., whether it is more or less compelling or entertaining[26-28]. The present study extends these results in an important new direction, suggesting that there is substantial implicit variation in the brain's response to a naturalistic stimulus that stems from trait-level individual differences.

Our results have implications for the neural correlates of both trait- and state-related paranoia. Those with higher trait paranoia may have more stereotyped brain responses because

suspicious and/or paranoid schemas come to mind more readily for these individuals; the idea that
certain individuals tend to engage certain constructs more frequently across time and situations
has been termed "chronic accessibility"[29]. The relative hyperactivity of theory-of-mind regions to
mentalizing events in high-paranoia individuals fits with the conception of paranoia as "over-
mentalizing", or the tendency to excessively attribute (malevolent) intentions to other people's
actions[30]. Both regions of differential response, the temporal pole and medial PFC, are sometimes,
but not always, reported in theory-of-mind tasks broadly construed; individual differences may at
least partially explain the inconsistencies in the literature[31].

While the present study included only healthy controls with subclinical paranoia, it may
provide a useful starting point for the study of paranoid or persecutory delusions in schizophrenia
and related illnesses. Delusions with a persecutory theme account for roughly 70-80 percent of all
delusions. This high prevalence is stable across time[32] and geo-cultural factors[33-36], suggesting a
strong biological component. Persecutory delusions are also the type most strongly associated with
anger and most likely to be acted upon, especially in a violent manner[37]. Thus, understanding the
neurobiological basis of paranoid delusions is a critical problem in psychiatry.

But because delusions typically have a slow, insidious onset, it is nearly impossible to
retrospectively recover triggering events in individual patients. A related challenge is that while
thematically similar, each patient's delusion is unique in its details. Thus it is difficult to devise
material that will evoke comparable responses across patients. One solution is to craft a model
context using a stimulus that is ambiguous yet controlled—i.e., identical across participants,
permitting meaningful comparisons of time-locked evoked activity— such as the one used in this
work. Paradigms such as this one may shed light on mechanisms of delusion formation and/or
provide eventual diagnostic or prognostic value.

While there is little work investigating brain activity during naturalistic stimuli in
psychiatric populations, a handful of studies have used such paradigms in autism, finding that
autistic individuals are less synchronized with one another and with typically developing controls
while watching movies of social interactions[38-40]. Notably, the degree of asynchrony scales with
autism-spectrum phenotype severity in both the patient and control groups[39]. It is interesting to
juxtapose these reports with the present results, in which individuals with a stronger paranoia
phenotype were *more* synchronized during exposure to socially relevant material; ultimately, this
fits with the notion of autism and psychosis as opposite ends of the same spectrum, involving

hypo- and hyper-mentalization, respectively[41,42]. Future studies should combine naturalistic

stimuli with ISC-based analyses that cut across diagnostic labels to examine how neural responses

vary across the full range of human phenotypes.

From a methodological perspective, much of the fMRI research on individual differences

has shifted in recent years from measuring activation in task-based conditions to measuring

functional connectivity, predominantly at rest[43-47]. Both paradigms suffer from limitations:

traditional tasks are so tightly controlled that they often lack ecological validity; resting-state

scans, on the other hand, are entirely unconstrained, making it difficult to separate signal from

noise. Naturalistic tasks may be a happy medium for studying both group-level functional brain

organization as well as individual differences[48,49]. We and others argue that such tasks could serve

as a "stress test" to draw out individual variation in brain and behaviors of interest[50-54], enhancing

signal in the search for neuroimaging-based biomarkers and permitting more precise inferences

about the sources of individual differences in neural activity.

METHODS

Participants

A total of 23 healthy volunteers participated in this study. Data from one participant was excluded

due to excessive head motion and self-reported falling asleep during the last third of the narrative.

Thus, the final data set used for analysis contained 22 participants (11 females; age range = 19-35

years, mean = 27, s.d. = 4.4). All participants were right-handed, native speakers of English, with

no history of neurological disease or injury, and were not on psychoactive medication at the time

of scanning. All participants provided written informed consent in accordance with the

Institutional Review Board of Yale University. The experiment took place over two visits to the

laboratory. Participants were paid $25 upon completion of the first visit (behavioral assessments)

and $75 upon completion of the second visit (MRI scan); all participants completed both visits.

Stimulus

An original narrative was written by author E.S.F. to serve as the stimulus for this experiment. For

a synopsis of the story, see Supplementary Note 1. The full audio recording, as well as a complete

transcript, are available in the "stimuli" directory at the following URL:

https://openneuro.org/datasets/ds001338/. To mitigate confounds associated with education level

or verbal IQ, we wrote the narrative text to be easy to comprehend, with a readability level of

78.1/100 and a grade 5.5 reading level as calculated by the Flesch-Kinkaid Formula.

Audio recording. A male native speaker of English read the story aloud and his speech was

recorded using high-quality equipment at Haskins Laboratories (New Haven, Conn.). The speaker

was instructed to read in a natural, conversational tone, but without excess emotion. The final

length of the audio recording was 21:50.

Experimental protocol

Session 1: Behavior. Approximately one week prior to the scan visit, participants came to the

laboratory to complete a battery of self-report and behavioral tasks. While our primary measure of

interest was the Green et al. Paranoid Thoughts Scale (GPTS)[21], we also administered several other

psychological scales and cognitive assessments, in part to help reduce any demand characteristics

that would allow participants to intuit the purpose of the study. We chose the GPTS because it

provides a meaningful assessment of trait-level paranoia in clinical, but crucially, also subclinical

and healthy populations. In a previous study, score on this scale best predicted feelings of

persecution following immersion in a virtual-reality environment[55]. The full GPTS contains two

subscales, A and B, which pertain to ideas of social reference and ideas of persecution,

respectively. We focused on subscale A, as it produces a wider range of scores in subclinical

populations[21].

The following cognitive tests were administered via the web interface of the University of

Pennsylvania Computerized Neuropsychological Test Battery (PennCNP;

penncnp.med.upenn.edu)[56]: SRAVEN (short Raven's progressive matrices, a measure of

abstraction and mental flexibility, or fluid intelligence); SPVRT (short Penn logical reasoning test,

a measure of verbal intelligence); and LNB2 (letter n-back, a measure of working memory). We

also administered the word reading test from the Wide Range Achievement Test 3 (WRAT)[57], a

measure of reading and vocabulary.

Session 2: MRI scan. The full audio recording was divided into three segments of length 8:46,

7:32, and 5:32, respectively; each of these segments was delivered in a continuous functional run

while participants were in the scanner. To ensure attention, after each run, subjects answered three

challenging multiple-choice comprehension questions regarding the content of the part they had

just heard, for a total of nine questions. Immediately upon exiting the scanner, participants

completed a post-narrative questionnaire that consisted of open-ended prompts to elicit free

speech, followed by multiple-choice items. These are described further below.

MRI data acquisition and preprocessing

Scans were performed on a 3T Siemens TimTrio system at the Yale Magnetic Resonance

Research Center. After an initial localizing scan, a high-resolution 3D volume was collected using

a magnetization prepared rapid gradient echo (MPRAGE) sequence (208 contiguous sagittal slices,

slice thickness = 1 mm, matrix size 256 × 256, field of view = 256 mm, TR = 2400 ms, TE = 1.9

ms, flip angle = 8°). Functional images were acquired using a multiband T2*-sensitive gradient-

recalled single shot echo-planar imaging pulse sequence (TR = 1000 ms, TE = 30 ms, voxel size

= 2.0mm^3, flip angle = 60°, bandwidth = 1976 Hz/pixel, matrix size = 110 × 110, field of view =

220 mm × 220 mm, multiband factor = 4).

We acquired the following functional scans: 1) an initial eyes-open resting-state run

(6:00/360 TRs in duration) during which subjects were instructed to relax and think of nothing in

particular; 2) a movie-watching run using Inscapes [58] (7:00/420 TRs); 3) three narrative-listening

runs corresponding to parts 1, 2 and 3 of the story (21:50/1310 TRs in total); and 4) a post-

narrative, eyes-open resting-state run (6:00/360 TRs) during which subjects were instructed to

reflect on the story they had just heard. The present work focuses exclusively on data acquired

during narrative listening. The narrative stimulus was delivered through MRI-compatible audio

headphones and a short "volume check" scan was conducted just prior to the first narrative run to

ensure that participants could adequately hear the stimulus above the scanner noise. To promote

engagement, during the three narrative runs, participants were asked to fixate on a static image of

a jungle settlement and to actively imagine the story events as they unfolded.

Following conversion of the original DICOM images to NIFTI format, AFNI (Cox 1996)

was used to preprocess MRI data. The functional time series went through the following

preprocessing steps: despiking, head motion correction, affine alignment with anatomy, nonlinear

alignment to a Talairach template (TT_N27), and smoothing with an isotropic FWHM of 5 mm.

A ventricle mask was defined on the template and intersected with the subject's cerebrospinal fluid

(CSF) mask to make a subject-specific ventricle mask. Regressors were created from the first three

principal components of the ventricles, and fast ANATICOR (Jo et al 2010) was implemented to

provide local white matter regressors. Additionally, the subject's 6 motion time series, their

derivatives and linear polynomial baselines for each of the functional runs were included as

regressors. Censoring of time points was performed whenever the per-time motion (Euclidean

norm of the motion derivatives) was ≥0.3 or when ≥10% of the brain voxels were outliers.

Censored time points were set to zero rather than removed altogether (this is the conventional way

to do censoring, but especially important for inter-subject correlation analyses, to preserve the

temporal structure across participants). The final output of this preprocessing pipeline was a single

functional run concatenating data from the three story runs (total duration = 21:50, 1310 TRs). All

analyses were conducted in volume space and projected to the surface for visualization purposes.

We used mean framewise displacement (MFD), a per-participant summary metric, to

assess the amount of head motion in the sample. MFD was overall relatively low (after censoring:

mean = 0.075 mm, s.d. = 0.026, range = 0.035-0.14). Number of censored time points during the

story was overall low but followed a right-skewed distribution (range = 0-135, median = 4, median

absolute deviation = 25). All 22 participants in the final analysis retained at least 89 percent of the

total time points in the story, so missing data was not a substantial concern. Still, we performed

additional control analyses to ensure that number of censored timepoints and amount of head

motion were not associated with paranoia score in any way that would confound interpretation of

the results (see Table 1).

Inter-subject correlation

Following preprocessing, inter-subject correlation (ISC) during the story was computed

across all possible pairs of subjects (*i,j*) using AFNI's 3dTcorrelate function, resulting in 231

(n*(n-1)/2, where n = 22) unique ISC maps, where the value at each voxel represents the Pearson's

correlation between that voxel's timecourse in subject *i* and its timecourse in subject *j*.

To identify voxels demonstrating statistically significant ISC across all 231 subject pairs,

we performed inference at the single-group level using a recently developed linear mixed-effects

(LME) model with a crossed random-effects formulation to accurately account for the correlation

structure embedded in the ISC data [22]. This approach has been characterized extensively, including

a comparison to non-parametric approaches, and found to demonstrate proper control for false

positives and good power attainment [22]. The resulting map was corrected for multiple comparisons

and thresholded for visualization using a voxelwise false discovery rate threshold of q < 0.001

(Fig. 2).

In a second analysis, we stratified participants according to a median split of scores on the

GPTS-A subscale. We used these groups to identify voxels that had higher ISC values within one

paranoia group or the other, or higher ISC values within rather than across paranoia groups. To

this end, we used a two-group formulation of the LME model. This model gives the following

outputs: voxelwise population ISC values within group 1 (G_{11}); voxelwise population ISC values

within group 2 (G_{22}); voxelwise population ISC values between the two groups that reflect the ISC

effect between any pair of subjects with each belonging to different groups (G_{12}). These outputs

can be compared to obtain several possible contrasts. Here, we were primarily interested in three

of these contrasts: 1) G_{11} versus G_{22}, 2) G_{11} versus G_{12}, and 3) G_{22} versus G_{12}. The maps resulting

from each of these contrasts were thresholded using an initial voxelwise threshold of p < 0.002

and controlled for family-wise error (FWE) using a cluster size threshold of 50 voxels,

corresponding to a corrected *p*-value of 0.05. We opted for a particularly stringent initial p-

threshold in light of recent concerns about false positives arising from performing cluster

correction on maps with more lenient initial thresholds [59].

Event-related analysis

Creating the regressor. A forced-aligner (Gentle; https://lowerquality.com/gentle/) was

used to obtain precise timing information for each word in the narrative, by aligning the audio file

with its transcript. One of the authors (E.S.F.) manually labeled sentences containing either an

ambiguous social interaction or an instance of the main character mentalizing about other

characters' intentions using a binary scoring system (1 = ambiguous social interaction or

mentalizing present in sentence, 0 = neither ambiguous social interaction nor mentalizing present).

Four additional, independent raters previously naïve to the narrative listened to the same version

that was played to participants in the scanner. They were then given a written version of the

narrative broken down by sentence and asked to label each sentence as described above. Sentences

that were labeled by at least three of the five raters were included in the final set of events. There

were 48 sentences that met this criteria, with 17, 13 and 18 occurring in parts 1, 2 and 3 of the

narrative, respectively.

Events were timestamped based on the TR corresponding to the offset of the last word of

each labeled sentence. These timestamps were convolved with a canonical hemodynamic response

function (HRF) to create the mentalizing-events regressor. Our assumption that evaluation and

integration would happen primarily at the end of the sentence was based on theories of text

comprehension, which hold that readers/listeners segment continuous linguistic information online

into larger units of meaning, or "macropropositions"; the mental models that listeners use to

represent narratives are thus updated primarily at event boundaries [60-62]. Empirical neurobiological

support for this comes from Whitney et al.[63], who showed, using a 23-minute continuous narrative

stimulus, that sentence boundaries coinciding with narrative shifts—defined as shifts in character,

time, location, or action— evoked more brain activity than sentence boundaries not coincident

with such shifts. Additional neuroimaging evidence comes from Zacks et al.[64], who demonstrated

transient changes in brain activity that were time-locked to event boundaries during movie

viewing.

However, some degree of evaluation and integration could also be happening online as

participants listen to the event, and ideally the results from the regression would not depend on

methodological choices about which parts of the sentence to model. To test this, we created a

second version of the regressor, this time treating the entire sentence as a mini-block by modeling

all TRs in each of the labeled sentences. Results were unchanged (see Supplementary Fig. 2). Thus

we are confident that the results are robust to this methodological choice.

As a control analysis, we also created a regressor that was the inverse of the above

regressor, by flipping the binary labels (0 or 1) for all sentences and convolving the corresponding

sentences offset timestamps with the HRF; we refer to this as the non-mentalizing-events

regressor.

ROI definition and GLM. For the left temporal pole and right medial PFC, ROIs were

defined based on the cluster-corrected group-comparison map for the contrast $ISC_{high} > ISC_{low}$ (cf.

Fig. 3a). For the left temporo-parietal junction (TPJ) and left Heschl's gyrus, spherical ROIs were

created by placing a sphere with radius 4 mm around a central coordinate. In the case of the TPJ

this was the peak voxel in this region identified by the whole-sample ISC analysis (cf. Fig. 2;

Talairach *xyz*, [+53, +55, +18]). In the case of Heschl's gyrus, this was selected anatomically

(Talairach *xyz*, [-41, -24, +9]; as in Schönwiesner et al.[65]).

Timecourses for each ROI were extracted from each participant's preprocessed functional

data using AFNI's 3dmaskave function and regressed against both the mentalizing- and non-

mentalizing-events regressors to obtain a regression coefficient for each participant for each ROI.

These regression coefficients were then compared across groups using two-sample t-tests corrected

for four multiple comparisons. In the case of the two control ROIs (TPJ and Heschl's gyrus) for

the mentalizing-events regressor, these coefficients were also pooled across both groups and

submitted to a one-sample t-test to test for a significant deviation from zero.

Free speech capture

Immediately following their exit from the scanner, we gave participants the following

prompts and recorded their speech: (1) "Please retell the story in as much detail as you can

remember"; and (2) "What did you think of the story as a whole? In particular, did anything strike

you as strange or confusing? How do you feel after listening to the story?" Here we focus on data

acquired from the first prompt, as participants consistently talked for much longer to this one than

to the second one (since they tended to preempt answers to second prompt in their answer to the
first).
598
599 **Multiple-choice questionnaire**
600   Following the free-speech prompts, we had participants complete a computerized multiple-
601 choice questionnaire to assess their feelings toward and beliefs about the story. A full list of items
602 is provided in Supplementary Table 1; there were 47 in total.
603
Analysis of speech features
Audio recordings of participants' retelling of the story were professionally transcribed by
a third-party company. We submitted the resulting transcripts to Linguistic Inquiry and Word
Count (LIWC; www.liwc.net)[25], a software program that takes as input a given text and counts the
percentage of words falling into different syntactic and semantic categories. Because LIWC was
developed by researchers with interests in social, clinical, health, and cognitive psychology, the
language categories were created to capture people's social and psychological states.
We restricted LIWC output to the 67 linguistic (syntactic and semantic) categories,
excluding categories relating to metadata (e.g., percentage of words found in the LIWC
dictionary), as well as categories irrelevant to spoken language (e.g., punctuation). Thus, our final
LIWC output was a 22x67 matrix where each row corresponds to a participant and each column
to a category.
These categories can be scaled very differently from one another. For example, words in
the syntactic category "pronoun" accounted for between 10.3-20.5 percent of speech transcripts,
while words in the semantic category "leisure" accounted for only 0-1.09 percent. To give
approximately equal weight to all categories, we standardized each category (to have zero mean
unit variance) across participants before performing partial least squares regression (PLSR) as
described in the next section. This ensures that the resulting PLS components are not simply
dominated by variance in categories that are represented heavily in all human speech.
623
Relating story-evoked behavior to paranoia
To determine which speech features were most related to trait paranoia, we submitted the
data to a partial least squares regression (PLSR) with the z-scored speech features as X (predictors)

and trait paranoia score as Y (response), implemented in Matlab as *plsregress*. PLSR is a latent

variable approach to modeling the covariance structure between two matrices, which seeks to find

the direction in X space that explains the maximum variance in Y space. It is well suited to the

current problem, because it can handle a predictor matrix with more variables than observations,

as well as multi-collinearity among the predictors.

In a first-pass analysis, we ran a model with 10 components to determine the number of

components needed to explain most of the variance in trait paranoia. Results of this analysis

indicated that the first component was sufficient to explain 72.3 percent of the total variance in

paranoia score, so we selected just this component for visualization and interpretation. Feature

loadings for this component are visualized in Fig. 6a.

In a parallel analysis, we submitted participants' answers to the multiple-choice

questionnaire to a PLSR as the X (predictor) matrix, again with paranoia score as the Y (response)

variable. Results of this analysis indicated that the first component was sufficient to explain 61.5

percent of the variance in paranoia score. Feature loadings for this component are visualized in

Supplementary Fig. 1c.

Data availability

Source data generated during this study, including raw MRI data and the full narrative

stimulus (audio and text), are available at: [https://openneuro.org/datasets/ds001338/].

Code availability

More information about this project, including links to code and other supporting material,

can be found at: [https://esfinn.github.io/projects/ParanoiaStory.html].

REFERENCES

651
652
1   Davis, B., Anderson, R. & Walls, J. *Rashomon Effects: Kurosawa, Rashomon and Their*
Legacies. (Taylor & Francis, 2015).
2   Savulich, G., Freeman, D., Shergill, S. & Yiend, J. Interpretation Biases in Paranoia.
Behav. Ther. **46**, 110-124, (2015).
3   Johns, L. C. & Van Os, J. The Continuity of Psychotic Experiences in the General
Population. *Clin. Psychol. Rev.* **21**, 1125-1141, (2001).
4   Insel, T. *et al.* Research Domain Criteria (Rdoc): Toward a New Classification Framework
for Research on Mental Disorders. *Am. J. Psychiatry.* **167**, 748-751, (2010).
5   Freeman, D. *et al.* Psychological Investigation of the Structure of Paranoia in a Non-
Clinical Population. *The British Journal of Psychiatry* **186**, 427-435, (2005).
6   Bebbington, P. E. *et al. The Structure of Paranoia in the General Population.* Vol. 202
(2013).
7   Whalley, H. C. *et al.* Correlations between Fmri Activation and Individual Psychotic
Symptoms in Un-Medicated Subjects at High Genetic Risk of Schizophrenia. *BMC*
Psychiatry **7**, 61, (2007).
8   Brent, B. K. *et al.* Subclinical Delusional Thinking Predicts Lateral Temporal Cortex
Responses During Social Reflection. *Soc. Cogn. Affect. Neurosci.*, nss129, (2012).
9   Sumich, A., Castro, A. & Kumari, V. N100 and N200, but Not P300, Amplitudes Predict
Paranoia/Suspiciousness in the General Population. *Pers. Individ. Dif.* **61**, 74-79, (2014).
10   Modinos, G., Renken, R., Ormel, J. & Aleman, A. Self-Reflection and the Psychosis-Prone
Brain: An Fmri Study. *Neuropsychology* **25**, 295-305, (2011).
11   Modinos, G. *et al.* Multivariate Pattern Classification Reveals Differential Brain Activation
During Emotional Processing in Individuals with Psychosis Proneness. *Neuroimage* **59**,
3033-3041, (2012).
12   Bishop, S. J. Neural Mechanisms Underlying Selective Attention to Threat. *Ann. N. Y.*
Acad. Sci. **1129**, 141-152, (2008).
13   Kukkonen, K. Bayesian Narrative: Probability, Plot and the Shape of the Fictional World.
Anglia **132**, 720-739, (2014).
14   Mar, R. A. The Neural Bases of Social Cognition and Story Comprehension. *Annu. Rev.*
Psychol. **62**, 103-134, (2011).
15   Lerner, Y., Honey, C. J., Silbert, L. J. & Hasson, U. Topographic Mapping of a Hierarchy
of Temporal Receptive Windows Using a Narrated Story. *J. Neurosci.* **31**, 2906-2915,
(2011).
16   Hasson, U., Nir, Y., Levy, I., Fuhrmann, G. & Malach, R. Intersubject Synchronization of
Cortical Activity During Natural Vision. *Science* **303**, 1634-1640, (2004).
17   Hasson, U., Malach, R. & Heeger, D. J. Reliability of Cortical Activity During Natural
Stimulation. *Trends in Cognitive Sciences* **14**, 40-48, (2010).
18   Cooper, E. A., Hasson, U. & Small, S. L. Interpretation-Mediated Changes in Neural
Activity During Language Comprehension. *Neuroimage* **55**, 1314-1323, (2011).
19   Lahnakoski, J. M. *et al.* Synchronous Brain Activity across Individuals Underlies Shared
Psychological Perspectives. *Neuroimage* **100**, 316-324, (2014).
20   Yeshurun, Y. *et al.* Same Story, Different Story: The Neural Representation of Interpretive
Frameworks. *Psychol. Sci.* **28**, 307-319, (2017).

21   Green, C. *et al.* Measuring Ideas of Persecution and Social Reference: The Green Et Al.
Paranoid Thought Scales (Gpts). *Psychol. Med.* **38**, 101-111, (2008).
22   Chen, G., Taylor, P. A., Shin, Y.-W., Reynolds, R. C. & Cox, R. W. Untangling the
Relatedness among Correlations, Part Ii: Inter-Subject Correlation Group Analysis through
Linear Mixed-Effects Modeling. *NeuroImage* **147**, 825-840, (2017).
23   Yarkoni, T., Speer, N. K. & Zacks, J. M. Neural Substrates of Narrative Comprehension
and Memory. *Neuroimage* **41**, 1408-1425, (2008).
24   Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C. & Wager, T. D. Large-Scale
Automated Synthesis of Human Functional Neuroimaging Data. *Nature Methods* **8**, 665,
(2011).
25   Linguistic Inquiry and Word Count: Liwc 2015 (Pennebaker Conglomerates, Austin, TX,
2015).
26   Schmälzle, R., Häcker, F. E. K., Honey, C. J. & Hasson, U. Engaged Listeners: Shared
Neural Processing of Powerful Political Speeches. *Soc. Cogn. Affect. Neurosci.* **10**, 1137-
1143, (2015).
27   Dmochowski, J. P. *et al.* Audience Preferences Are Predicted by Temporal Reliability of
Neural Processing. *Nature communications* **5**, 4567, (2014).
28   Ki, J. J., Kelly, S. P. & Parra, L. C. Attention Strongly Modulates Reliability of Neural
Responses to Naturalistic Narrative Stimuli. *J. Neurosci.* **36**, 3092-3101, (2016).
29   Higgins, E. T., King, G. A. & Mavin, G. H. Individual Construct Accessibility and
Subjective Impressions and Recall. *J. Pers. Soc. Psychol.* **43**, 35, (1982).
30   Brüne, M. "Theory of Mind" in Schizophrenia: A Review of the Literature. *Schizophr.*
Bull. **31**, 21-42, (2005).
31   Carrington, S. J. & Bailey, A. J. Are There Theory of Mind Regions in the Brain? A Review
of the Neuroimaging Literature. *Hum. Brain Mapp.* **30**, 2313-2335, (2009).
32   Cannon, B. J. & Kramer, L. M. Delusion Content across the 20th Century in an American
Psychiatric Hospital. *Int. J. Soc. Psychiatry* **58**, 323-327, (2012).
33   Tateyama, M., Asai, M., Hashimoto, M., Bartels, M. & Kasper, S. Transcultural Study of
Schizophrenic Delusions. Tokyo Versus Vienna and Tubingen (Germany).
Psychopathology **31**, 59-68, (1997).
34   Stompe, T. *et al.* Comparison of Delusions among Schizophrenics in Austria and in
Pakistan. *Psychopathology* **32**, 225-234, (1998).
35   Gutiérrez-Lobos, K., Schmid-Siegel, B., Bankier, B. & Walter, H. Delusions in First-
Admitted Patients: Gender, Themes and Diagnoses. *Psychopathology*, (2001).
36   Brakoulias, V. & Starcevic, V. A Cross-Sectional Survey of the Frequency and
Characteristics of Delusions in Acute Psychiatric Wards. *Australasian Psychiatry* **16**, 87-
91, (2008).
37   Coid, J. W., Ullrich, S., Kallis, C. & et al. The Relationship between Delusions and
Violence: Findings from the East London First Episode Psychosis Study. *JAMA Psychiatry*
70, 465-471, (2013).
38   Hasson, U. *et al.* Shared and Idiosyncratic Cortical Activation Patterns in Autism Revealed
under Continuous Real-Life Viewing Conditions. *Autism Research* **2**, 220-231, (2009).
39   Salmi, J. *et al.* The Brains of High Functioning Autistic Individuals Do Not Synchronize
with Those of Others. *NeuroImage: Clinical* **3**, 489-497, (2013).

40 Byrge, L., Dubois, J., Tyszka, J. M., Adolphs, R. & Kennedy, D. P. Idiosyncratic Brain
Activation Patterns Are Associated with Poor Social Comprehension in Autism. *J.*
Neurosci. **35**, 5837-5850, (2015).

41 Crespi, B. & Badcock, C. Psychosis and Autism as Diametrical Disorders of the Social
Brain. *Behav. Brain Sci.* **31**, 241-261, (2008).

42 Ciaramidaro, A. *et al.* Schizophrenia and Autism as Contrasting Minds: Neural Evidence
for the Hypo-Hyper-Intentionality Hypothesis. *Schizophr. Bull.* **41**, 171-179, (2015).

43 Kelly, C., Biswal, B. B., Craddock, R. C., Castellanos, F. X. & Milham, M. P.
Characterizing Variation in the Functional Connectome: Promise and Pitfalls. *Trends in*
Cognitive Sciences **16**, 181-188, (2012).

44 Castellanos, F. X., Di Martino, A., Craddock, R. C., Mehta, A. D. & Milham, M. P. Clinical
Applications of the Functional Connectome. *Neuroimage* **80**, 527-540, (2013).

45 Finn, E. S. *et al.* Functional Connectome Fingerprinting: Identifying Individuals Using
Patterns of Brain Connectivity. *Nat. Neurosci.* **18**, 1664-1671, (2015).

46 Smith, S. M. *et al.* A Positive-Negative Mode of Population Covariation Links Brain
Connectivity, Demographics and Behavior. *Nat. Neurosci.* **18**, 1565-1567, (2015).

47 Rosenberg, M. D. *et al.* A Neuromarker of Sustained Attention from Whole-Brain
Functional Connectivity. *Nat. Neurosci.* **advance online publication**, (2015).

48 Ren, Y., Nguyen, V. T., Guo, L. & Guo, C. C. Inter-Subject Functional Correlation Reveal
a Hierarchical Organization of Extrinsic and Intrinsic Systems in the Brain. *Sci. Rep.* **7**,
10876, (2017).

49 Vanderwal, T. *et al.* Individual Differences in Functional Connectivity During Naturalistic
Viewing Conditions. *bioRxiv*, (2016).

50 Finn, E. S. & Constable, R. T. Individual Variation in Functional Brain Connectivity:
Implications for Personalized Approaches to Psychiatric Disease. *Dialogues Clin.*
Neurosci. **18**, 277-287, (2016).

51 Dubois, J. & Adolphs, R. Building a Science of Individual Differences from Fmri. *Trends*
in Cognitive Sciences **20**, 425-443, (2016).

52 Koyama, M. S. *et al.* Imaging the at-Risk Brain: Future Directions. *J. Int. Neuropsychol.*
Soc. **22**, 164-179, (2016).

53 Finn, E. S. *et al.* Can Brain State Be Manipulated to Emphasize Individual Differences in
Functional Connectivity? *Neuroimage*, (2017).

54 Guo, C. C., Hyett, M. P., Nguyen, V. T., Parker, G. B. & Breakspear, M. J. Distinct
Neurobiological Signatures of Brain Connectivity in Depression Subtypes During Natural
Viewing of Emotionally Salient Films. *Psychol. Med.* **46**, 1535-1545, (2016).

55 VALMAGGIA, L. R. *et al.* Virtual Reality and Paranoid Ideations in People with an 'at-
Risk Mental State' for Psychosis. *The British Journal of Psychiatry* **191**, s63-s68, (2007).

56 Gur, R. C. *et al.* Computerized Neurocognitive Scanning:: I. Methodology and Validation
in Healthy People. *Neuropsychopharmacology* **25**, 766-776, (2001).

57 Wilkinson, G. S. *Wrat-3: Wide Range Achievement Test Administration Manual.* (Wide
Range, Incorporated, 1993).

58 Vanderwal, T., Kelly, C., Eilbott, J., Mayes, L. C. & Castellanos, F. X. Inscapes: A Movie
Paradigm to Improve Compliance in Functional Magnetic Resonance Imaging.
Neuroimage **122**, 222-232, (2015).

59 Eklund, A., Nichols, T. E. & Knutsson, H. Cluster Failure: Why Fmri Inferences for Spatial
Extent Have Inflated False-Positive Rates. *Proc. Natl. Acad. Sci. USA*, 201602413, (2016).

60   Zwaan, R. A. & Radvansky, G. A. Situation Models in Language Comprehension and
787   Memory. *Psychol. Bull.* **123**, 162-185, (1998).
788   61   Johnson-Laird, P.   (Cambridge, MA: Harvard University Press, 1983).
789   62   Van Dijk, T. A., Kintsch, W. & Van Dijk, T. A. *Strategies of Discourse Comprehension*.
790   (Academic Press New York, 1983).
791   63   Whitney, C. *et al.* Neural Correlates of Narrative Shifts During Auditory Story
792   Comprehension. *Neuroimage* **47**, 360-366, (2009).
793   64   Zacks, J. M. *et al.* Human Brain Activity Time-Locked to Perceptual Event Boundaries.
794   *Nat. Neurosci.* **4**, 651, (2001).
795   65   Schönwiesner, M. *et al.* Heschl's Gyrus, Posterior Superior Temporal Gyrus, and Mid-
796   Ventrolateral Prefrontal Cortex Have Different Roles in the Detection of Acoustic
797   Changes. *J. Neurophysiol.* **97**, 2075-2082, (2007).
798
799

Acknowledgements

This research was supported by a National Science Foundation Graduate Research Fellowship to E.S.F., and by the Intramural Research Program of the NIMH (annual report ZIAMH002783). Portions of this study used the computational capabilities of the NIH HPC Biowulf cluster (*biowulf.nih.gov*). The authors thank Paul Taylor for advice on preprocessing pipelines for naturalistic task data, Richard Betzel for sharing code to visualize component feature loadings (Fig. 6a and Supplementary Fig. 2), and Javier Gonzalez-Castillo, Daniel Handwerker, David Jangraw, Peter Molfese, Monica Rosenberg, and Tamara Vanderwal for helpful discussion.

Author contributions

E.S.F, P.R.C and R.T.C conceived the study. E.S.F. developed experimental materials and performed data collection. E.S.F. and G.C. analyzed the data. E.S.F., P.R.C., P.A.B. and R.T.C interpreted results. E.S.F. wrote the manuscript with comments from all other authors.

Competing financial interests

The authors have no competing financial interests to declare.

817
Figure 1. Experimental protocol, distribution of trait-level paranoia and inter-subject
correlation analysis. a) Schematic of experimental protocol. Participants came to the laboratory
for an initial behavioral visit, during which they completed several computerized cognitive tasks
as well as self-report psychological scales, one of which was the Green et al. Paranoid Thoughts
Scale (GPTS)[21]. To minimize demand characteristics and/or priming effects, the fMRI scan visit
took place approximately one week later. During this visit, subjects listened to an ambiguous social
narrative in the scanner and then completed an extensive post-narrative battery consisting of both
free-speech prompts and multiple-choice items. b) Distribution of scores on the GPTS-A subscale
across n = 22 participants, and median split used to stratify participants into low ($\leq$ 18, blue) and
high ($\geq$ 19, orange) trait-level paranoia. c) Schematic of inter-subject correlation (ISC) analysis.
Following normalization to a standard template, the inter-subject correlation of activation
timecourse during narrative listening was computed for each voxel (v, yellow square; enlarged
relative to true voxel size for visualization purposes) for each pair of subjects (i,j), resulting in a
matrix of pairwise correlation coefficients (r values). These values were then compared across
paranoia groups using voxelwise linear mixed-effects models with crossed random effects to
account for the non-independent structure of the correlation matrix[22].
834

835
836
Figure 2. Narrative listening evokes widespread inter-subject correlation across the whole
sample. Voxels showing significant inter-subject correlation (ISC) across the timecourse of
narrative listening in all participants (n = 22). As expected, the highest ISC values were observed
in auditory cortex, but several regions of association cortex in the temporal, parietal, frontal and
cingulate lobes as well as the cerebellum also showed high synchrony. Also included are three
representative axial slices from the cerebellum. Results are displayed at a voxelwise false-
discovery rate (FDR) threshold of q < 0.001).
844

845
846
Figure 3. Trait-level paranoia modulates patterns of inter-subject correlation during
narrative listening. a) Results frma whole-brain, voxelwise contrast revealing brain regions that
are more synchronized between pairs of high-paranoia participants than pairs of low-paranoia
participants (contrast schematized in top panel, cf. Fig. 1C). Significant clusters were detected in
the left temporal pole, two regions in the right medial prefrontal cortex (one anterior and one dorsal
and posterior), and the left precuneus. No clusters were detected in the opposite direction (low >
high). b) Results from two whole-brain, voxelwise contrasts revealing brain regions that are more
synchronized within a paranoia group than across paranoia groups. The first contrast revealed that
left lateral occipital cortex was more synchronized within the low-paranoia group (i.e., low-low
pairs) than across groups (i.e., high-low pairs; contrast schematized in top panel, cf. Fig 1C). The
second contrast revealed that right angular gyrus was more synchronized within the high-paranoia
group (i.e., high-high pairs) than across groups. For all three contrasts, results are shown at an
initial threshold of p < 0.002 with cluster correction corresponding to p < 0.05.
860

Figure 4. Inter-subject correlation scales continuously with trait paranoia. Post-hoc analyses for two regions of interest (ROIs) that emerged from the dichotomized contrast between high- and low-paranoia groups (cf. Fig. 3a): left temporal pole (top row) and right medial prefrontal cortex (PFC, bottom row). a) Location of ROI (left) and participant-by-participant inter-subject correlation (ISC) matrix (right) for the left temporal pole. Participants are ordered by increasing trait paranoia score. Each matrix element reflects the correlation between two participants' activation timecourses in the left temporal pole during narrative listening. Higher correlations are visible as one moves to the right and down along the diagonal, representing pairs of increasingly high-paranoia individuals. b) Scatter plot of paranoia rank versus median ISC value—i.e., the median of each row of the ISC matrix in (a). Each dot represents a participant. Rank correlation indicates a significant monotonic relationship between trait paranoia and median ISC in left temporal pole ($r_s = 0.71$, $p = 0.0002$). c) Location of ROI and participant-by-participant ISC matrix for the right medial PFC. Participants are ordered as in (a). d) Scatter plot of each participant's paranoia rank versus their median ISC value in the right medial PFC. As in (b), rank correlation indicates a significant monotonic relationship between paranoia rank and median ISC ($r_s = 0.63$, $p = 0.0016$).

880
881
Figure 5. Response to mentalizing events is stronger in high- compared to low-paranoia
individuals. a) Regions of interest (ROIs) for the event-related analysis. LTmpPole, left temporal
pole; RmPFC, right medial prefrontal cortex; LTPJ, left temporo-parietal junction; LHeschl, left
Heschl's gyrus. b) Comparison of beta coefficients for each ROI for the mentalizing-events
regressor between paranoia groups (low, blue; high, orange). Each dot represents a subject. Boxes
represent the median and $25^{th}/75^{th}$ percentiles, and whiskers represent the minimum and maximum.
*p = 0.01; **p < 0.007; n.s., not significant (p-values adjusted to control the false discovery rate).
c) Comparison of beta coefficients for each ROI for the non-mentalizing-events regressor (the
inverse of the mentalizing-events regressor shown in (b)). Each dot represents a subject. Boxes
represent the median and $25^{th}/75^{th}$ percentiles, and whiskers represent the minimum and maximum.
d) Beta coefficients for the mentalizing-events regressor plotted against paranoia rank (coefficients
are the same as in (b)). Left panel: the two ROIs in which beta coefficient was hypothesized to
scale with trait paranoia (LTmpPole and RmPFC). Right panel: the two control ROIs (LTPJ and
LHeschl). Correlations between paranoia rank and beta coefficient: LTmpPole, $r_s = 0.57$, p =
0.005; RmPFC, $r_s = 0.64$, p = 0.001; LTPJ, $r_s = -0.04$, p = 0.86, LHeschl, $r_s = 0.02$, p = 0.95. e)
Beta coefficients for the non-mentalizing-events regressor plotted against paranoia rank
(coefficients are the same as in (c)). Left and right panels as in (d). Correlations between paranoia
rank and beta coefficients (all n.s.): LTmpPole, $r_s = -0.28$, p = 0.21; RmPFC, $r_s = -0.22$, p = 0.33;
LTPJ, $r_s = 0.085$, p = 0.71; LHeschl, $r_s = 0.17$, p = 0.44.

Figure 6. Speech analysis reveals a signature of trait-level paranoia in behavioral response to the narrative. a) Loadings of all semantic and syntactic categories for the first component from a partial least squares regression relating features of speech during narrative recall to trait-level paranoia score, sorted by strength and direction of association with paranoia (those positively related to paranoia at top in orange; those inversely related at bottom in blue). b) Example sentences from participant speech transcripts containing words falling into the three of the top positive categories (affiliation, health and anxiety) and one of the top negative categories (anger). c) Rank correlations between participants' trait-level paranoia and their self-report measures of 16 emotions following the narrative (self-report was based on a Likert scale from 1 to 5). Dotted lines represent approximate threshold for a significant correlation a p < 0.05 (uncorrected). Gray shaded area indicates non-significance.

		Categorical (low vs. high)		Continuous	
		t	*p*	Spearman *r*	*p*
Demographics	Age	0.81	0.43	-0.11	0.62
	Sex*	1.64	0.20	--	--
	Education (yrs)	-0.24	0.81	-0.15	0.49
Cognitive ability	Working memory: Letter n-back (precision)	-0.45	0.66	0.16	0.47
	Fluid intelligence: Raven's matrices (total correct of 9 items)	0.00	1.00	-0.03	0.89
	Vocabulary: WRAT Word Reading (total correct of 42 items)	-1.42	0.17	0.31	0.16
	Verbal IQ: Penn logical reasoning test (total correct of 8 items)	0.23	0.82	-0.01	0.96
	Words of 6+ letters (free recall)	-1.03	0.32	0.04	0.85
	Words per sentence (free recall)	0.31	0.76	-0.18	0.43
fMRI data quality	Head motion (mean FD; mm)	0.94	0.36	0.01	0.96
	No. frames censored	-0.70	0.49	-0.08	0.74
	Average tSNR	-1.12	0.28	0.23	0.30
Attention to stimulus	No. comprehension questions correct	-0.31	0.76	0.08	0.72
	Total word count, free recall	1.00	0.33	-0.26	0.24
	Self-reported attention	0.48	0.63	-0.02	0.95
	Self-reported engagement	0.89	0.39	-0.10	0.65

Table 1. Trait paranoia was unrelated to potential confounding variables. There were no
significant differences between high- and low-paranoia participants in terms of demographics,
cognitive abilities, fMRI data quality or attention to the stimulus. Categorical comparisons were
carried out using Student's t-tests between the low and high paranoia groups as determined by
median split (degrees of freedom for all t-tests = 20). Continuous comparisons were carried out
using Spearman (rank) correlation between raw paranoia score and the variable of interest. All p-
values are raw (uncorrected). *Measured with a chi-squared test. FD, framewise displacement;
tSNR, temporal signal-to-noise ratio; WRAT, Wide Range Achievement Test.

a Visit 1

Behavior
- psychological scales
- cognitive tasks

approx. 1 week

Visit 2

fMRI scan
- story listening

Post-narrative battery
- free recall (speech)
- multiple-choice items

"One day, a doctor named Carmen gets an email..."

b

Low
n = 11

High
n = 11

median

GPTS-A score

c

Subj 1 2 ... n

ISC_{[low, low]}

ISC_{[high, low]}

ISC_{[high, high]}

$r^v_{i,j}$

Signal

Time

r value

0.16

a

Contrast:
ISC{high, high} > ISC{low, low}

b

Contrasts:
ISC{low, low} > ISC{low, high}
ISC{high, high} > ISC{low, high}

Occipital view

L   R

r value

0.08

−0.08

0.04

−0.04

a L temporal pole

b

trait paranoia

Subj

trait paranoia

ISC (*r* value)

median ISC (*r* value)

trait paranoia (rank)

$r_s = 0.71$

c R medial PFC

d

trait paranoia

Subj

trait paranoia

ISC (*r* value)

median ISC (*r* value)

trait paranoia (rank)

$r_s = 0.63$

a

LIWC loadings onto first PLSR component

affiliation
adjectives
health
anxiety
drives
pers. pron: 1st plural
female references
family
see
prepositions
insight
achievement
articles
perceptual processes
comparisons
biological processes
positive emotion
reward
verbs
risk
affective processes
interrogatives
pers. pron: 1st singular
present focus
leisure
negative emotion
discrepancy
social processes
space
work
differentiation
past focus
death
hear
friends
motion
money
pers. pron: 3rd singular
time
tentative
relativity
feel
auxiliary verbs
positive processes
future focus
adverbs
body
power
sadness
sexual
relig
negations
home
ingestion
pronouns
impersonal pron.
seasonal
certainty
causation
quantifiers
pers. pron: 3rd plural
conjunctions
function words
anger
male references

b

Affiliation

"...but they become **friends** once she gets to know her better."

Health

"...there were a series of **patients** that had a **fever** accompanied by a **rash**..."

Anxiety

"...she almost **panicked**..."

Anger

"...and she **yells** at him."

c

excited
amused
hopeful
happy
inspired
confused
paranoid
disturbed
threatened
suspicious
surprised
calm
anxious
sad
frustrated
uncomfortable

$p < 0.05$ $p < 0.05$

Correlation with trait paranoia (r_s)