1   **The ERK MAPK pathway modulates Gq-dependent locomotion in**

2   ***Caenorhabditis elegans***

5   Brantley Coleman, Irini Topalidou, and Michael Ailion

Department of Biochemistry, University of Washington, Seattle, WA, 98195

Running Title: The ERK pathway modulates Gq signaling

Keywords: Gq signaling, ERK MAPK pathway, Rho small GTPase, *C. elegans*, G

protein

Corresponding author:

Michael Ailion

Department of Biochemistry

University of Washington

Box 357350

1705 NE Pacific St

Seattle, WA 98195

Phone: 206-685-0111

email: mailion@uw.edu

Abstract

The heterotrimeric G protein Gq regulates neuronal activity through distinct downstream effector pathways. In addition to the canonical Gq effector phospholipase Cβ, the small GTPase Rho was recently identified as a conserved effector of Gq. To identify additional molecules important for Gq signaling in neurons, we performed a forward genetic screen in the nematode *Caenorhabditis elegans* for suppressors of the hyperactivity and exaggerated waveform of an activated Gq mutant. We isolated two mutations affecting the MAP kinase scaffold protein KSR-1 and found that KSR-1 modulates locomotion downstream of or in parallel to the Gq-Rho pathway. Through epistasis experiments, we found that the core ERK MAPK cascade is required for Gq-Rho regulation of locomotion, but that the canonical ERK activator LET-60/Ras may not be required. Through neuron-specific rescue experiments, we found that the ERK pathway functions in head acetylcholine neurons to control Gq-dependent locomotion. Additionally, expression of activated LIN-45/Raf in head acetylcholine neurons is sufficient to cause an exaggerated waveform phenotype and hypersensitivity to the acetylcholinesterase inhibitor aldicarb, similar to an activated Gq mutant. Taken together, our results suggest that the ERK MAPK pathway modulates the output of Gq-Rho signaling to control locomotion behavior in *C. elegans.*

Introduction

Introduction

The heterotrimeric G protein Gq is a conserved regulator of neurotransmission in

metazoans. Gq is highly expressed in neurons in mammals and in the nematode *C.*

elegans (Wilkie *et al.* 1991; Lackner *et al.* 1999). In its canonical signaling pathway, Gq

activates phospholipase Cβ (PLCβ) to cleave the membrane lipid phosphatidylinositol

4,5-bisphosphate (PIP2) into inositol trisphosphate (IP3) and diacylglycerol (DAG)

(Rhee 2001). An increased DAG concentration at the synapse helps trigger synaptic

vesicle release (Miller *et al.* 1999; Lackner *et al.* 1999).

In addition to activating PLCβ, Gq directly binds and activates the Rho guanine

nucleotide exchange factor (GEF) Trio, which in turn activates the small GTPase Rho

(Lutz *et al.* 2005, 2007; Williams *et al.* 2007). In mature *C. elegans* neurons, the Rho

ortholog RHO-1 regulates synaptic activity through multiple G protein-dependent

mechanisms. First, RHO-1 acts downstream of the G_{12}-class G protein GPA-12 by

binding to and inhibiting the diacylglycerol kinase DGK-1. Inhibition of DGK-1 allows

DAG to accumulate at the synapse, thereby increasing synaptic vesicle release

(McMullan *et al.* 2006; Hiley *et al.* 2006). Second, Gq-Rho signaling promotes

neurotransmitter release by recruiting the sphingosine kinase SPHK-1 to presynaptic

terminals (Chan *et al.* 2012). Third, Gq-Rho signaling positively regulates the NCA-

1/NALCN cation channel to regulate locomotion (Topalidou *et al.* 2017a). Here we

identify the extracellular signal-related kinase mitogen-activated protein kinase (ERK

MAPK) pathway as a positive regulator of neuronal activity acting downstream of or in

parallel to Gq and Rho in acetylcholine neurons.

ERK MAPK signaling acts extensively in animal development, cellular

proliferation, and cancer signaling (Yoon and Seger 2006; Karnoub and Weinberg 2008;

Sun *et al.* 2015). ERKs are highly expressed in mammalian neurons (Boulton *et al.*

1991; Ortiz *et al.* 1995) and act in both the nucleus and at the synapse to regulate

synaptic activity and plasticity (Thomas and Huganir 2004; Sweatt 2004; Mao and Wang

2016b). In *C. elegans*, the ERK pathway is required for multiple developmental events

including specification of the vulva (Sternberg 2005; Sundaram 2013), and also acts in

several types of neurons to control behavior. ERK signaling is activated in response to

odorants in the AWC sensory neuron to regulate chemotaxis to volatile odorants and in

AIY interneurons to mediate odor adaptation (Hirotsu *et al.* 2000; Hirotsu and Iino 2005;

Chen *et al.* 2011; Uozumi *et al.* 2012). ERK is also activated in the ASER sensory

neuron to regulate chemotaxis to salt (Tomioka *et al.* 2006; Tomida *et al.* 2012). ERK

signaling regulates foraging behavior by acting in the IL1, OLQ, and RMD neurons

(Hamakawa *et al.* 2015). Finally, the ERK pathway has been shown to act in

interneurons to regulate the nose touch response, a mechanosensory behavior (Hyde *et*

al. 2011).

In the canonical ERK MAPK pathway, extracellular ligand binding activates

transmembrane receptor tyrosine kinases (RTKs), and adaptor proteins recruit a GEF to

activate the small GTPase Ras (LET-60 in *C. elegans*). Upon Ras activation, LIN-45/Raf

translocates to the plasma membrane where it interacts with Ras and the scaffold

protein KSR-1. KSR-1 facilitates the activation of LIN-45/Raf and the subsequent

phosphorylation of the MAPK cascade consisting of LIN-45/Raf, MEK-2/MEK, and MPK-

1/ERK (Sundaram 2013). In this study, we found that the ERK MAPK pathway

consisting of KSR-1, LIN-45/Raf, MEK-2/MEK and MPK-1/ERK modulates Gq-Rho

signaling in acetylcholine neurons, but that surprisingly LET-60/Ras may not be

required.

Materials and Methods

***C. elegans* strains**

All strains were cultured using standard methods and were maintained at 20°C.

Table S1 contains all the strains used in this study.

Isolation and mapping of the ksr-1(ox314) and ksr-1(yak10) mutations

The *ox314* and *yak10* mutants were isolated from an ENU mutagenesis

suppressor screen of the activated Gq mutant *egl-30(tg26)* (Ailion *et al.* 2014). We

mapped the *ox314* mutation by its activated Gq suppression phenotype using single

nucleotide polymorphisms (SNPs) in the Hawaiian strain CB4856 as described (Davis *et*

al. 2005).The *ox314* mutation was mapped to an approximately 709 kb region in the

middle of the X chromosome between SNPs on cosmids F45E1 and F53A9 (SNPs

F45E1[1] and pkP6158). This region included 159 predicted protein-coding genes. A

complementation test of *ox314* and *yak10* in the *egl-30(tg26)* background showed these

to be alleles of the same gene. Whole genome sequencing (see below) identified these

as mutations in *ksr-1*, and we confirmed this by performing a complementation test with

the deletion allele *ksr-1(ok786).*

Whole genome sequencing

Strains EG4198 *egl-30(tg26); ox314* and XZ1340 *egl-30(tg26); yak10* were

sequenced to identify candidate mutations. DNA was purified according to the Hobert

Lab protocol (http://hobertlab.org/whole-genome-sequencing/). Ion Torrent sequencing

was performed at the University of Utah DNA Sequencing Core Facility. Each data set

contained roughly 18,400,000 reads of a mean read length of 160 bases, resulting in

about 30X average coverage of the *C. elegans* genome. The sequencing data were

processed on the Galaxy server at usegalaxy.org (Afgan *et al.* 2016). SNPs and indels

were identified and annotated using the Unified Genotyper and SnpEff tools (DePristo *et*

al. 2011; Cingolani *et al.* 2012). After filtering for mutations in open reading frames, we

found each strain to have unique stop mutations in *ksr-1*, in the middle of the interval

where we mapped *ox314*. *ox314* is a G to A transition that causes a stop codon at

amino acid K463, and *yak10* is an A to T transversion that causes a stop codon at

W254.

Locomotion assays

Track waveform and radial locomotion assays were performed on 10 cm

nematode growth medium (NGM) plates seeded with 400 µl of OP50 *E. coli* culture and

spread with sterile glass beads. Bacterial lawns were grown at 37°C for 16 hrs and the

plates were stored at 4°C until needed. For track waveform measurements, five first day

adult animals were placed on a plate and allowed to roam for 2-5 min. We then

recorded each animal's tracks following forward locomotion. Track pictures were taken

at 40X on a Nikon SMZ18 microscope with the DS-L3 camera control system. Pictures

of worm tracks were processed using ImageJ. Period and 2X amplitude were measured

freehand using the line tool. For each worm, we calculated the average

period/amplitude ratio of five individual track bends (Figure 1C). For assays with the

temperature sensitive allele *sos-1(cs41)*, all strains were grown at 20°C and shifted to

the non-permissive temperature of 25°C for 24 hours before being assayed. For radial

locomotion assays, ten to fifteen first day adult animals were picked to the center of a

plate and were then allowed to move freely for 40 minutes. The positions of worms were

marked and the distances of the worms from the starting point were measured. For all

waveform and radial locomotion assays, the experimenter was blind to the genotypes of

the strains assayed.

Microscopy

Photographs of moving worms were taken at 60X on a Nikon SMZ18 microscope

with the DS-L3 camera control system. The worms were age-matched as first day

adults grown at 20°C.

Constructs and transgenes

Plasmids were constructed using the three-slot multisite Gateway cloning system

(Invitrogen). Plasmids and primers used are found in Tables S2 and S3. The *ksr-1*, *lin-*

45, *let-60*, and *egl-30* cDNAs were amplified by PCR from a worm cDNA library and

cloned into [1-2] Gateway entry vectors. Activating Raf mutations T626E/T629D were

introduced into the *lin-45* cDNA vector by two sequential site-directed mutagenesis

reactions (Q5 kit, NEB) with primers oBC094/095 and oBC096/097, respectively, and

then confirmed by sequencing. The dominant Ras mutation S17N was introduced into

the *let-60* cDNA by site-directed mutagenesis with primers oBC109/110. The *tg26*

activating Gq mutation R243Q was introduced into the *egl-30* cDNA by site-directed

mutagenesis with primers oET254/255. cDNAs were cloned into expression constructs

under different neuronal promoters using the multisite Gateway system. Specifics of

promoter fragments used are given in Table S2. For expression in different subclasses

of acetylcholine neurons, we used the *Punc-17H* promoter that is expressed in head

acetylcholine neurons (Hammarlund *et al.* 2007; Topalidou *et al.* 2017a) and the *Punc-*

17β promoter that is expressed in ventral cord acetylcholine motor neurons (Charlie *et*

al. 2006; Topalidou *et al.* 2017a). Proper expression of *ksr-1*, *lin-45* and *let-60* was

confirmed by including an operon GFP::H2B in the [2-3] slot of the expression

constructs. The operon GFP template *tbb-2 3'utr::gpd-2 operon::GFP::H2B:cye-1 3'utr*

(Frøkjær-Jensen *et al.* 2012) results in untagged proteins whose expression can be

monitored by GFP expression.

Injections and chromosomal integrations

Worms carrying the activated *lin-45* transgenes *Punc-17::lin-45** and *Punc-*

17H::lin-45* as extrachromosomal arrays were generated by injecting pBC37 or pBC44

at 20 ng/µL or 10 ng/µL respectively along with co-injection markers pCFJ104 (*Pmyo-*

3::mCherry) at 5 ng/µL, pCFJ90 (*Pmyo-2::mCherry*) at 2.5 ng/µL, and the carrier DNA

Litmus 38i to a final concentration of 100 ng/µL DNA (Mello *et al.* 1991). Worms carrying

the dominant negative *let-60* transgene *Prab-3::let-60(S17N)* were generated by

injecting pBC50 at 20 ng/ul with the same co-injection markers. Worms carrying the

activated Gq *egl-30(tg26)* transgene *Punc-17H::egl-30(tg26)* were generated by

injecting pET102 at 10 ng/ul with the co-injection marker pCFJ90 (*Pmyo-2::mCherry*) at

1 ng/µL. MosSCI lines were generated as described (Frøkjær-Jensen *et al.* 2012) using

an injection mix containing 10-15 ng/µL targeting vector, 50 ng/µL pCFJ601 (*Peft-*

3::Mos1 transposase), negative selection markers pGH8 (*Prab-3::mCherry*) at 10 ng/μL,

pCFJ104 (*Pmyo-3::mCherry*) at 5 ng/μL, pCFJ90 (*Pmyo-2::mCherry*) at 2.5 ng/μL,

pMA122 (*Phsp16.41::peel-1*) at 10 ng/μL, and carrier DNA Litmus 38i to a final

concentration of 100 ng/μL DNA.

Extrachromosomal arrays were integrated into the genome by exposure to 4000

rads of gamma irradiation. Irradiated young adult hermaphrodites were transferred to 10

cm OP50 plates (5 worms/plate) and grown to starvation. The plates were chunked and

grown to starvation twice more to enrich for stably expressing lines. When nearly

starved, 8 animals per plate were picked to individual plates. The progeny were then

screened for 100% stable transmission, indicating integration into the genome.

Integration was confirmed by mapping the transgene to a chromosome.

Aldicarb assays

35 mm aldicarb assay plates were poured with NGM supplemented with 1 mM

aldicarb. The plates were seeded with 5 uL OP50 and dried at room temperature

overnight. Animals were picked onto the OP50 lawn to begin the assay (time 0) and

then kept at room temperature. Every 15 minutes, animals were scored for paralysis by

lightly touching the nose of the animal with an eyebrow hair. Animals were scored as

paralyzed if the worm displayed no locomotor response to three nose touches and had

no pharyngeal pumping. Animals that left the OP50 lawn were picked back onto the

food.

Statistical analysis

P values were determined using GraphPad Prism 5. Normally distributed data

sets were analyzed with a one-way ANOVA and Bonferroni's *post hoc* test when group

size was unequal, or with Tukey's *post hoc* test when group size was equal. Data sets

with non-normal distribution (using the Shapiro-Wilk normality test) were analyzed with

a Kruskal-Wallis test and Dunn's *post hoc* test. Data sets with multiple independent

variables were analyzed by two-way ANOVA and Bonferroni's *post hoc* test.

Reagent and data availability

Strains and plasmids are listed in Tables S1 and S2 and are available upon

request. Primers are listed in Table S3. The authors state that all data necessary for

confirming the conclusions presented in the article are represented fully within the

article and Supplemental Material.

Results

KSR-1 regulates locomotion downstream of or in parallel to Gq

In *C. elegans,* the heterotrimeric G protein Gq regulates synaptic vesicle release

(Hu *et al.* 2015). Gq is a key regulator of neuromuscular activity, as loss-of-function

mutants in *egl-30* are nearly paralyzed (Brundage *et al.* 1996) whereas the gain-of-

function mutant *egl-30(tg26)* has hyperactive locomotion with an exaggerated loopy

waveform (Doi and Iwasaki 2002; Bastiani *et al.* 2003) (Figure 1, A, C, and D). To

identify pathways required for Gq signaling, we performed a forward genetic screen in

C. elegans for suppressors of the activated Gq mutant *egl-30(tg26)*. This screen has

been used to identify new components of the Gq signal transduction pathway and

proteins important for dense-core vesicle function (Williams *et al.* 2007; Ailion *et al.*

2014; Topalidou *et al.* 2016; Topalidou *et al.* 2017a; b). Two previously uncharacterized

suppressors identified in this screen, *ox314* and *yak10*, showed similar suppression of

the loopy waveform and hyperactivity of *egl-30(tg26)* animals (Figure 1D). When

crossed away from the *egl-30(tg26)* background, both mutants moved with wild-type

waveform (Figure 1D), but at a slightly slower rate. We mapped the *ox314* allele near

the center of the X chromosome (see Materials and Methods), and a complementation

test showed that *ox314* and *yak10* are mutations in the same gene since they fail to

complement in an *egl-30(tg26)* background.

We used whole genome sequencing to identify *ox314* and *yak10* as nonsense

mutations in *ksr-1* (Figure 1B, see Materials and Methods). KSR-1 is a scaffold protein

that facilitates the localization and interactions required for the Ras-mitogen activated

protein kinase (MAPK) cascade consisting of Raf, MEK, and ERK (Kornfeld *et al.*

1995b; Sundaram and Han 1995; Nguyen *et al.* 2002; Zhang *et al.* 2013). The deletion

allele *ksr-1(ok786)* also suppressed the loopy waveform of the activated Gq mutant

identically to *ox314* and *yak10.* These results suggest that KSR-1 activity is required for

regulation of locomotion rate and waveform by Gq.

The ERK MAPK cascade acts to promote Gq signaling

256   Because the loss of the MAPK scaffold *ksr-1* suppresses the activated Gq

mutant *egl-30(tg26)*, we asked whether other components of the Ras-ERK pathway

would also suppress. Since the core components of the Ras-ERK pathway are required

for viability, we built double mutants of activated Gq with reduction-of-function mutations

in genes at each step of the ERK cascade. Mutations in Raf (*lin-45(sy96)*)*,* MEK (*mek-*

2(n1989), *mek-2(ku114)*), and ERK (*mpk-1(ga117)*, *mpk-1(oz140)*) all suppressed the

loopy waveform of activated Gq animals similarly to *ksr-1(ok786)* (Figure 2, A and B;

Figure S1, A and B). However, mutations in Ras (*let-60(n2021)*) and the upstream

pathway activators EGF (*lin-3(e1417)*) and the EGF receptor (*let-23(sy12)*) did not

suppress activated Gq (Figure 2C). Because *let-60* is required for viability, most *let-60*

alleles including *n2021* are partial loss-of-function (Beitel *et al.* 1990). We also analyzed

the dominant negative D119N allele *let-60(sy93)* that disrupts Ras binding to guanine

nucleotides and thus prevents Ras activation (Han and Sternberg 1991). We found that

let-60(sy93) also did not suppress the loopy waveform of activated Gq (Figure 2D).

Additionally, we expressed the dominant negative *let-60* mutation S17N specifically in

neurons and did not observe suppression of the loopy waveform of activated Gq (Figure

S2D). These results support the possibility that ERK activation in this pathway occurs

through a Ras-independent mechanism.

Because partial loss-of-function mutations in the ERK MAPK pathway genes

downstream of Ras showed clear suppression of activated Gq, we were surprised to

find that partial loss-of-function mutations in Ras did not suppress. If LET-60/Ras is

indeed not required, Gq might instead activate the ERK pathway via other Ras-

subfamily proteins. To test this possibility, we made double mutants of activated Gq with

putative null alleles of R-Ras/*ras-1,* M-Ras/*ras-2,* and Rap1/*rap-1* and found that they

also did not suppress activated Gq (Figure S2A). To further investigate whether this

pathway acts independently of Ras, we tested mutations in GEFs that activate Ras. The

temperature-sensitive RasGEF mutant *sos-1(cs41)* did not suppress activated Gq when

shifted to the non-permissive temperature (Figure S2B). Additionally, a null mutation in

the neuronal RasGEF *rgef-1* also did not suppress activated Gq (Figure S2C). In

genetic screens for vulval induction mutants, additional factors such as the PP2A

subunit *sur-6* (Sieburth *et al.* 1999) and ion transporter *sur-7* (Yoder *et al.* 2004) were

identified as positive regulators of Ras-ERK activity. However, the *sur-6(sv30)* and *sur-*

7(ku119) mutations did not suppress activated Gq locomotion (data not shown). These

data suggest either that only a low level of Ras activity is needed to properly activate

ERK signaling downstream of Gq, or that ERK signaling acts independently of LET-

60/Ras and other known *C. elegans* Ras family proteins to regulate locomotion

downstream of Gq.

KSR-1 and the ERK MAPK cascade modulate Rho signaling

Three classes of suppressor mutations were isolated in our forward genetic

screen of activated Gq, as characterized by their molecular roles and unique

suppression phenotypes (Topalidou *et al.* 2017a; b). We grouped together a class of

suppressor mutations including *ox314, yak10*, and the RhoGEF Trio (*unc-73* in *C.*

elegans) by their characteristic strong suppression of the loopy waveform of activated

Gq (Topalidou *et al.* 2017a; b), suggesting that *ksr-1* might act in the same pathway as

unc-73.

We have shown that Gq regulates locomotion via the small GTPase Rho (RHO-1

in *C. elegans*) (Topalidou *et al.* 2017a). Transgenic expression of an activated RHO-1

mutant (G14V) in acetylcholine neurons (here called "*rho-1(gf)*") causes worms to have

a loopy waveform and impaired locomotion (McMullan *et al.* 2006) (Figure 3A). To

examine whether *ksr-1* acts in the Gq-Rho pathway we tested whether mutations in *ksr-*

1 suppress the phenotypes of *rho-1(gf)* worms. We found that the *ksr-1* alleles *ok786,*

ox314, and *yak10* all suppressed the loopy waveform of *rho-1(gf)* worms (Figure 3, A,

B, and D). Because *rho-1(gf)* worms have a slow locomotion rate and loopy waveform,

these mutants do not efficiently travel long distances. We used radial locomotion assays

(see Materials and Methods) to quantify the locomotion phenotype of *rho-1(gf)* worms.

rho-1(gf) ksr-1 double mutants had an increased radial distance traveled compared to

rho-1(gf) alone (Figure 3B). These data suggest that KSR-1 acts downstream of or in

parallel to the Gq-Rho pathway to regulate locomotion.

Since *ksr-1* mutants suppress the exaggerated waveform of both activated Gq

and activated Rho animals, we expected that loss of other ERK pathway components

would also suppress activated Rho. We made double mutants of activated Rho (*rho-*

1(gf)) with reduction-of-function alleles of the Ras-ERK pathway and found that

mutations in Raf, MEK, and ERK suppressed the *rho-1(gf)* loopy waveform phenotype

(Figure 3, A, C, and D). However, the *let-60(n2021)* and *let-60(sy93)* Ras mutations did

not suppress the loopy waveform of *rho-1(gf)* worms (Figure 3, A, C, and D). These

data suggest that the ERK pathway acts downstream of or in parallel to the Gq-Rho

pathway to regulate locomotion, possibly in a Ras-independent manner.

The ERK MAPK cascade acts in acetylcholine neurons to control locomotion

Members of the ERK pathway are expressed in neurons in *C. elegans* (Dent and

Han 1998; Hunt-Newbury *et al.* 2007), and Gq and Rho act in acetylcholine neurons to

promote synaptic release and regulate locomotion (Lackner *et al.* 1999; McMullan *et al.*

2006). To determine whether the ERK pathway also acts in neurons to modulate Gq

signaling, we expressed the *ksr-1* cDNA under promoters driving expression in specific

types of neurons. Single-copy transgenic expression of *ksr-1* under an acetylcholine

neuron promoter (*Punc-17*) or under a head acetylcholine neuron promoter (*Punc-17H*)

fully reversed the *ksr-1* suppression of the loopy waveform of activated Gq worms

(Figure 4). *ksr-1* expression in ventral cord acetylcholine motor neurons (*Punc-17β*) or

GABA neurons (*Punc-47)* did not significantly reverse the *ksr-1* suppression of activated

Gq (Figure 4). This suggests that ERK signaling primarily functions in the acetylcholine

interneurons or motor neurons of the head to modulate Gq-dependent locomotion.

We have shown that the ERK pathway is necessary for Gq-dependent effects on

locomotion. To determine whether ERK signaling is sufficient to modulate locomotion,

we expressed an activated form of *lin-45*/Raf specifically in acetylcholine neurons. Raf

kinase activity is regulated via conserved phosphorylation events, and phosphomimetic

mutations T626E/T629D in the kinase activation loop of *lin-45* are sufficient to confer

constitutive Raf activity (Chong *et al.* 2001). We found that expression of activated Raf

in acetylcholine neurons (*Punc-17*) causes a loopy waveform similar to activated Gq

and Rho mutants and similar limited dispersal in radial locomotion assays (Figure 5, A

and B; Figure S3, A and B). Additionally, expression of activated Raf or activated Gq in

head acetylcholine neurons also caused a loopy waveform (Figure S3, A-C).

Gq and Rho promote acetylcholine release at the worm neuromuscular junction

(Lackner *et al.* 1999; Miller *et al.* 1999; McMullan *et al.* 2006). To determine if Raf

activation affects acetylcholine release at the neuromuscular junction, we assayed for

sensitivity to the acetylcholinesterase inhibitor aldicarb. Mutants with reduced

acetylcholine secretion are resistant to aldicarb, whereas mutants with increased

acetylcholine secretion are hypersensitive to aldicarb (Mahoney *et al.* 2006). Activated

Gq and Rho mutants have increased rates of paralysis when exposed to aldicarb

(Lackner *et al.* 1999; McMullan *et al.* 2006). We found that expression of activated Raf

in all acetylcholine neurons or in head acetylcholine neurons also led to aldicarb

hypersensitivity, similar to activated Gq and Rho mutants (Figure 5, C and D). However,

we found that a *ksr-1* mutation does not suppress the aldicarb hypersensitivity of

activated Gq or Rho mutants, and the *ksr-1* mutant on its own has similar aldicarb

sensitivity to wild type (Figure 5, E and F). These results suggest that the ERK pathway

is not required for synaptic transmission, but is sufficient to stimulate synaptic

transmission when constitutively activated. Furthermore, these data suggest that ERK

pathway activation in head acetylcholine neurons is sufficient to promote acetylcholine

release by motor neurons at the neuromuscular junction.

Discussion

In this study we identified KSR-1 and the ERK MAPK cascade as modulators of

Gq-Rho signaling. We found that ERK signaling acts in head acetylcholine neurons

(interneurons or motor neurons of the head) to modulate Gq-dependent locomotion

behavior, especially the waveform of the animal. Our data support the model that Gq-

Rho activation of the ERK pathway may be independent of its canonical regulator, the

small GTPase Ras/LET-60 (Figure 6).

Gq regulates multiple aspects of *C. elegans* locomotion behavior, including the

rate of locomotion and waveform. Previous work has shown that these behaviors are at

least partially genetically separable, with different pathways downstream of Gq

contributing differentially to these behaviors and with Gq likely acting in several classes

of neurons including both head acetylcholine neurons and ventral cord acetylcholine

motor neurons (Lackner *et al.* 1999; Topalidou *et al.* 2017a). Here we focused on Gq

regulation of the waveform, which is strongly dependent on Rho and ERK. The most

parsimonious model supported by our findings is that Gq, Rho, and the ERK pathway

function cell autonomously in the head acetylcholine neurons to regulate the waveform

(Figure 6). Other components of the Gq-Rho pathway have been previously shown to

act in head acetylcholine neurons to regulate the waveform (Topalidou *et al.* 2017a; b),

and here we show that KSR-1 is required in head acetylcholine neurons to modulate

Gq-dependent effects on the waveform. Additionally, expression of activated LIN-45/Raf

in head acetylcholine neurons is also sufficient to regulate the waveform and cause

loopy locomotion. These data suggest that ERK pathway activity in head acetylcholine

neurons is both necessary and sufficient to regulate worm waveform.

Though the ERK pathway is required for Gq and Rho-dependent regulation of the

waveform, it is not required for the aldicarb hypersensitivity of activated Gq and Rho

mutants. This indicates that Gq and Rho regulation of acetylcholine release at the

neuromuscular junction is at least partially independent of Gq and Rho regulation of the

locomotion waveform. The simplest model is that Gq and Rho signaling regulates the

waveform by acting in head acetylcholine neurons through an ERK-dependent pathway,

but Gq and Rho signaling regulates synaptic transmission at the neuromuscular junction

through an ERK-independent pathway. Overexpression of an activated Gq mutant in

ventral cord acetylcholine motor neurons is sufficient to cause an aldicarb

hypersensitive phenotype (Lackner *et al.* 1999), suggesting that Gq may act cell

autonomously in the ventral cord motor neurons to promote acetylcholine release by

these neurons. But as with activated Raf, it is also possible that Gq activity in head

acetylcholine neurons may contribute to acetylcholine release by downstream ventral

cord motor neurons.

The ERK signaling cascade has been well-studied for its regulation of cellular

proliferation and differentiation (Sun *et al.* 2015), but also plays important roles in

mature neurons and has been associated with synaptic plasticity and memory (Impey *et*

al. 1999). In addition to activating transcription, ERKs regulate synaptic plasticity both

presynaptically and postsynaptically by phosphorylating relevant substrates. ERKs

phosphorylate the presynaptic proteins synapsin I and Munc18-1, and postsynaptic

proteins such as scaffolds, $K_v4.2$ potassium channels, and Group I metabotropic

glutamate receptors (Jovanovic *et al.* 2000; Thomas and Huganir 2004; Sweatt 2004;

Kushner *et al.* 2005; Boggio *et al.* 2007; Vara *et al.* 2009; Mao and Wang 2016a;

Schmitz *et al.* 2016). Our findings suggest that the ERK pathway controls Gq-dependent

locomotion in *C. elegans* and that activated ERK promotes synaptic transmission.

In contrast to many developmental and neuronal roles of Ras-dependent ERK

signaling in *C. elegans,* our data suggest that Raf-MEK-ERK signaling may modulate

Gq-Rho output independently of Ras/LET-60. One caveat to the conclusion that the

ERK pathway acts independently of Ras to control Gq-dependent locomotion is that the

alleles of *let-60*/Ras tested here are not null (Han and Sternberg 1990, 1991; Han *et al.*

1990; Beitel *et al.* 1990). However, other than *mpk-1(ga117)*, the alleles we used of *lin-*

45, *mek-2* and *mpk-1* are also not null (Han *et al.* 1993; Kornfeld *et al.* 1995a; Wu *et al.*

1995; Lackner and Kim 1998; Hsu *et al.* 2002)**,** yet were able to suppress an activated

Gq mutant. Furthermore, the *let-*60/Ras alleles used here have stronger phenotypes in

vulval development than the two weak *mek-2*/MEK alleles we used (*n1989* and *ku114*)

(Beitel *et al.* 1990; Kornfeld *et al.* 1995a; Wu *et al.* 1995) and the *let-60(n2021)*/Ras

mutant has comparable phenotypes to the *lin-45(sy96)*/Raf and *mpk-1(ga117)*/ERK

alleles for chemotaxis to odorants and the regulation of foraging behavior (Hirotsu *et al.*

2000; Hamakawa *et al.* 2015). Thus, either LET-60/Ras is not required to modulate Gq

signaling, or ERK signaling in the locomotor circuit requires a lower threshold of Ras

activity and the *let-60* mutants we used have sufficient levels of active Ras to modulate

Gq signaling, but not enough for vulval induction or other behaviors.

If Ras is not required, how instead might KSR-Raf-MEK-ERK signaling be

activated? Normally, active Ras recruits Raf to the plasma membrane but it has been

shown that artificial recruitment of Raf to the plasma membrane in the absence of Ras

is sufficient to activate Raf signaling (Stokoe *et al.* 1994; Marais *et al.* 1995). Thus,

another possible way to activate the ERK pathway would be for Raf to be recruited to

the membrane by proteins other than Ras. The most obvious candidates for recruiting

Raf are other members of the Ras family such as Rap1 or R-Ras that have a conserved

effector-binding domain (Reiner and Lundquist 2016); it has been reported that Rap1

can mediate Ras-independent activation of Raf downstream of a Gq-coupled receptor

(Guo *et al.* 2001). However, mutations in the worm orthologs of Rap1/RAP-1 and R-

Ras/RAS-1 did not suppress the activated Gq mutant, indicating that these Ras-like

proteins are not required, at least individually, to activate Raf in this pathway. It is

possible that more than one of these Ras family members function redundantly to

activate Raf, or that Raf is activated independently of Ras family proteins. There have

been other reported cases of Ras-independent activation of the Raf-MEK-ERK pathway

(Robbins *et al.* 1992; Honda *et al.* 1994; van Biesen *et al.* 1996; Ueda *et al.* 1996;

Drosten *et al.* 2014), some of which involve G protein signaling and protein kinase C,

though the precise mechanisms involved are unclear.

Gq, Rho, and ERK are also required for the *C. elegans* behavioral and immune

response to infection by the bacterium *M. nematophilum* (Nicholas and Hodgkin 2004;

McMullan *et al.* 2012). Interestingly, LET-60/Ras was reported to not be required for the

immune response (Nicholas and Hodgkin 2004) or only partially required (McMullan *et*

al. 2012). Additionally, LET-60/Ras was not required for the increased sensitivity to

aldicarb caused by bacterial infection (McMullan *et al.* 2012), at least as assayed using

the *let-60(n2021)* allele. Thus, the same neuronal Ras-independent ERK pathway we

describe here may also modulate Gq signaling in the neuronal response to bacterial

infection and possibly the innate immune response as well.

Our epistasis analysis suggests that ERK signaling acts genetically downstream

of or in parallel to Gq-Rho signaling. How might Gq-Rho signaling lead to ERK

activation? ERK activation could occur via a linear pathway downstream of Gq and Rho,

or ERK could signal in parallel and converge downstream of Rho to affect neuronal

activity (Figure 6). There is precedence for Gq activating ERK via a linear pathway. In

the pharyngeal muscle of *C. elegans*, ERK activity is increased by Gq-dependent

signaling through protein kinase C (You *et al.* 2006). In the AWC olfactory neuron, ERK

activity is increased downstream of Gq signaling via the RasGEF RGEF-1 (Chen *et al.*

2011; Uozumi *et al.* 2012). Protein kinase C and RGEF-1 are both activated by DAG,

probably produced by the canonical Gq-PLCβ pathway. By contrast, we found that ERK

signaling regulates locomotion by modulating the output of the Gq-Rho pathway that

acts in parallel to Gq-PLCβ, and does not depend on RGEF-1.

How might the ERK pathway modulate neuronal activity downstream of Gq-Rho

signaling? Rho regulates neuronal activity and synaptic release through several

mechanisms in *C. elegans* neurons, any of which could be targets of ERK signaling.

One possible ERK effector is the NCA/NALCN cation channel that acts genetically

downstream of Gq-Rho to regulate locomotion rate and waveform (Topalidou *et al.*

2017a; b). Though ERK has not been directly connected to NCA/NALCN, mammalian

ERK regulates neuronal excitability by directly phosphorylating voltage-gated sodium

and potassium channels (Schrader *et al.* 2006; Stamboulian *et al.* 2010) and by

regulating channel expression (Yang *et al.* 2015). Given the many possible ways that

ERK may regulate excitability or synaptic transmission, *C. elegans* genetics is well

suited to determine relevant effectors of Gq-Rho-ERK signaling.

Acknowledgements

We thank Brooke Jarvie for the isolation of *ksr-1(yak10)*; Stephen Nurrish for the

activated Rho worm strain; Jordan and Jill Hoyt for help with whole-genome sequence

analysis; Laura Taylor for help with irradiation of extrachromosomal arrays; Dana Miller

for use of her Nikon SMZ18 microscope and camera. Some strains were provided by

the CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40

OD010440). This work was supported by an Ellison Medical Foundation New Scholar

Award and by NIH grants R00 MH082109 and R56 NS100843 to M.A.

Figure Legends

Figure 1. *ksr-1* mutations suppress activated Gq.

(A) A *ksr-1* mutation suppresses the coiled posture of activated Gq. The activated Gq mutant *egl-30(tg26)* has small size and deep body bends. *ksr-1(ok786)* suppresses the exaggerated body bends and small size of *egl-30(tg26)* worms.

(B) Gene structure of *ksr-1* locus. Locations of the *egl-30(tg26)* suppressor alleles *ox314* and *yak10* are indicated, as well as the position of the *ok786* deletion. The gene structure was drawn using Exon-Intron Graphic Maker (www.wormweb.org/exonintron)

(C) A *ksr-1* mutation suppresses the exaggerated waveform of *egl-30(tg26)* mutants. Straightened images of tracks left in bacterial lawns show similar waveform for wild type and *ksr-1(ok786)* worms. *egl-30(tg26)* mutants have an exaggerated waveform, creating tracks with a large amplitude relative to the period. *ksr-1*(*ok786)* suppresses the exaggerated waveform of the *egl-30(tg26)* mutant.

(D) *ksr-1* mutations suppress the *egl-30(tg26)* exaggerated waveform. *ksr-1* nonsense alleles *ox314* and *yak10* and the deletion allele *ok786* strongly but incompletely suppress the *egl-30(tg26)* exaggerated waveform. $N \geq 12$, *** $P < 0.001$, one-way ANOVA with Bonferroni's *post hoc* test. All *egl-30(tg26); ksr-1* double mutants were significantly different than *ksr-1* single mutants ($P<0.05$), indicating that suppression is incomplete.

Figure 2. Mutations in the ERK MAPK pathway suppress activated Gq.

(A) Mutations in the MAPKKK *lin-45*/Raf and the MAPK *mpk-1*/ERK suppress the coiled

posture of *egl-30*(*tg26*) worms. Partial loss of Ras *let-60* activity does not suppress

activated Gq posture.

(B) Waveform quantification of ERK pathway mutants. Mutations in *lin-45*, *mek-2*, and

mpk-1 suppress activated Gq *egl-30*(*tg26*) similar to *ksr-1* mutations. $N \geq 12$, *** $P <$

0.001, one-way ANOVA with Bonferroni's *post hoc* test.

(C) Signaling components upstream of Raf do not suppress activated Gq. Mutations in

the EGF ligand (*lin-3*) or the EGFR (*let-23*) do not affect the exaggerated waveform of

egl-30(*tg26*) animals. The Ras partial loss-of-function mutation *let-60(n2021)* does not

suppress activated Gq waveform. $N \geq 12$, *** $P < 0.001$, one-way ANOVA with

Bonferroni's *post hoc* test.

(D) The *let-60(sy93)* dominant negative mutation in Ras does not suppress activated Gq

waveform. $N \geq 13$, n.s., not significant, one-way ANOVA with Bonferroni's *post hoc* test.

Figure 3. Mutations in *ksr-1* and the ERK pathway suppress activated Rho.

(A) Mutations in ERK pathway genes *ksr-1*, *lin-45*, *mek-2*, and *mpk-1* visibly suppress

the coiled posture of animals expressing an activated Rho mutant (G14V) in

acetylcholine neurons (*nzIs29[Punc-17::rho-1(gf)]*). Reduction of LET-60/Ras activity

does not suppress activated Rho.

(B) The *ksr-1(ok786)*, *ksr-1(ox314)*, and *ksr-1(yak10)* mutations suppress the

locomotion of activated Rho animals as shown by radial locomotion assays. $N \geq 38$, ***

$P < 0.001$, ** $P < 0.01$, Kruskal-Wallis test with Dunn's *post hoc* test.

(C) Mutations in *lin-45* and *mek-2* suppress the locomotion defect of activated Rho

animals as shown by radial locomotion assays. The *let-60*(*n2021*) mutation does not

significantly suppress activated Rho locomotion. *N* ≥ 50, *** *P* < 0.001, Kruskal-Wallis

test with Dunn's *post hoc* test.

(D) Waveform quantification of ERK pathway mutants. Mutations in *ksr-1*, *lin-45*, *mek-2*,

and *mpk-1* show similar suppression of activated Rho, but the Ras reduction-of-function

mutant *let-60(n2021)* or dominant negative *let-60(sy93)* do not suppress activated Rho.

N ≥ 14, * P < 0.05, n.s., not significant, one-way ANOVA with Bonferroni's *post hoc* test,

compared to *rho-1(gf)*.

Figure 4. KSR-1 acts in head acetylcholine neurons to modulate Gq signaling.

Single-copy expression of the *ksr-1* cDNA exclusively in acetylcholine neurons (*Punc-*

17, *yakSi26* transgene) or head acetylcholine neurons (*Punc-17H, yakSi27* transgene)

is sufficient to reverse the *ksr-1* suppression of the loopy waveform of activated Gq

animals. *ksr-1* expression in ventral cord acetylcholine neurons (*Punc-17β, yakSi28*

transgene) or GABA neurons (*Punc-47*, *yakSi29* transgene) does not reverse the *ksr-1*

suppression of the activated Gq exaggerated waveform. *N* ≥ 12, *** *P* < 0.001, ** *P* <

0.01, n.s., not significant, Kruskal-Wallis test with Dunn's *post hoc* test.

Figure 5. Increased Raf activity in acetylcholine neurons is sufficient to regulate

locomotion and increase acetylcholine release.

(A) Transgenic lines expressing an activated form of *lin-45*/Raf (T626E/T629D) in

acetylcholine neurons (*yakIs34[Punc-17::lin-45(gf)]*) have exaggerated body bends and

coiled posture similar to the activated Gq mutant *egl-30*(*tg26*) and to animals expressing

activated Rho in acetylcholine neurons (*nzIs29[Punc-17::rho-1(gf)]*). The wild type and

egl-30(tg26) photos are the same as shown in Figure 1A, and the *rho-1(gf)* photo is the

same as the one in Figure 3A.

(B) Expression of activated Rho (*nzIs29[Punc-17::rho-1(gf)]*) or Raf (*yakIs34[Punc-*

17::lin-45(gf)]) in acetylcholine neurons impairs coordinated locomotion similarly to

activated Gq (*egl-30(tg26)*). *N* ≥ 19, * *P* < 0.05, *** *P* < 0.001, Kruskal-Wallis test with

Dunn's *post hoc* test.

(C) Animals expressing activated Gq, Rho, or Raf are hypersensitive to the

acetylcholinesterase inhibitor aldicarb. Activated Gq (*egl-30*(*tg26*)), activated Rho

expressed in acetylcholine neurons (*nzIs29[Punc-17::rho-1(gf)]*), and activated Raf

expressed in acetylcholine neurons *yakIs34[Punc-17::lin-45(gf)]*) become paralyzed

significantly faster than wild type animals when exposed to 1 mM aldicarb. All strains

are significantly different from wild type at t = 60, 75, 90, and 105 minutes. *N* ≥ 61, *** *P*

< 0.001, two-way ANOVA with Bonferroni's *post hoc* test.

(D) Animals expressing activated Raf in head acetylcholine neurons are hypersensitive

to aldicarb. The *yakIs34[Punc-17::lin-45(gf)]* integrated array expressing activated Raf in

all acetylcholine neurons and the *yakEx154[Punc-17H::lin-45(gf)]* (#1) and

yakEx168[Punc-17H::lin-45(gf)] (#2) extrachromosomal arrays expressing activated Raf

in head acetylcholine neurons show similar hypersensitivity to aldicarb. All strains are

significantly different from wild type at t = 60, 75, 90, and 105 minutes. N ≥ 47, ** P <

0.01, two-way ANOVA with Bonferroni's *post hoc* test.

(E) *ksr-1* is not required for the aldicarb hypersensitivity of activated Gq. Paralysis of

ksr-1(*ok786*) animals on 1 mM aldicarb is not significantly different from wild type. The

ksr-1 deletion *ok786* does not suppress the aldicarb hypersensitivity of activated Gq

(*egl-30*(*tg26*)). *N* ≥ 53.

(F) *ksr-1* is not required for the aldicarb hypersensitivity of activated Rho. Paralysis

of *ksr-1(ok786)* animals on 1 mM aldicarb is not significantly different from wild type.

The *ksr-1* deletion *ok786* does not suppress the aldicarb hypersensitivity of the

activated Rho mutant *nzIs29[Punc-17::rho-1(gf)]*. N ≥ 41.

Figure 6. Model for Gq-Rho-ERK signaling. Gq directly activates Trio and Rho (solid

arrow) (Lutz et al. 2007). The core ERK cascade acts either downstream of Rho or in

parallel (dashed arrows) to modulate locomotion behavior. ERK activation occurs

independently of the extracellular growth factor LIN-3, its receptor LET-23, or Ras/LET-

60. It is possible that Raf/LIN-45 is activated by a Ras family member other than the

canonical worm Ras/LET-60. Our model suggests that Gq-Rho signaling regulates the

worm waveform by acting cell autonomously in head acetylcholine neurons in an ERK-

dependent pathway. By contrast, we propose that Gq-Rho signaling regulates synaptic

transmission at the neuromuscular junction by acting in an ERK-independent pathway,

possibly in the ventral cord motorneurons themselves.

Figure S1. Additional mutations in *mek-2* and *mpk-1* suppress activated Gq.

(A) The *mek-2(ku114)* mutation suppresses the exaggerated waveform of *egl-30*(*tg26*).

N ≥ 14, *** *P* < 0.001, one-way ANOVA with Bonferroni's *post hoc* test.

(B) The *mpk-1(oz140)* mutation suppresses the exaggerated waveform of *egl-30(tg26)*.

$N \geq 12$, *** $P < 0.001$, one-way ANOVA with Bonferroni's *post hoc* test.

Figure S2. The exaggerated waveform of activated Gq does not depend on the

RasGEFs *sos-1* and *rgef-1*, or the Ras family members *ras-1*, *ras-2*, and *rap-1*.

(A) Deletion mutations *ras-1(gk243)* and *ras-2(ok682),* or the nonsense mutation *rap-*

1(pk2082) do not suppress the waveform of the activated Gq mutant *egl-30(tg26)*. $N \geq$

18, n.s., not significant, one-way ANOVA with Bonferroni's *post hoc* test.

(B) Loss of the RasGEF *sos-1* does not suppress activated Gq waveform. Animals

carrying the temperature sensitive *sos-1(cs41)* mutation were incubated for 24 hours at

the non-permissive temperature of 25° and assayed for their waveform. $N \geq 15$, n.s., not

significant, one-way ANOVA with Tukey's *post hoc* test.

(C) The RasGEF deletion mutation *rgef-1(ok675)* does not suppress the exaggerated

waveform of activated Gq. $N = 16$, n.s., not significant, one-way ANOVA with Tukey's

post hoc test.

(D) Neuronal expression of the dominant negative Ras mutant S17N (*yakEx158[Prab-*

3::let-60(S17N)]) does not suppress the coiled posture of the activated Gq mutant *egl-*

30(tg26).

Figure S3. Expression of activated LIN-45/Raf or EGL-30/Gq in head acetylcholine

neurons is sufficient to cause locomotion with an exaggerated waveform.

(A, B) The *yakIs34[Punc-17::lin-45(gf)]* integrated array expressing activated *lin-*

45(T626E/T629D) in all acetylcholine neurons and the *yakEx154[Punc-17H::lin-45(gf)]*

extrachromosomal array expressed in head acetylcholine neurons both cause a coiled

posture (A) and exaggerated waveform (B). *N* ≥ 12, *** *P* < 0.001, * *P* < 0.05, Kruskal-

Wallis test with Dunn's *post hoc* test.

(C) Expression of the activated Gq mutant *egl-30(tg26)* R243Q in head acetylcholine

neurons (*Punc-17H::egl-30(tg26)*) causes an exaggerated waveform. N ≥ 7, ** P < 0.01,

unpaired T test, two-tailed. Overexpression of activated Gq is toxic and these animals

were slow growing and had reduced viability and fertility.

Table S1. List of strains

N2   Bristol wild strain

XZ1151   *egl-30(tg26) I*

EG6699   *ttTi5605 II; unc-119(ed3) III*

EG6207   *unc-119(ed3) III*

RB915   *ksr-1(ok786) X*

EG4782   *nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II*

MT4866   *let-60(n2021) IV*

PS436   *let-60(sy93) IV*

PS427   *lin-45(sy96) IV*

MT8666   *mek-2(n1989) I*

SD378   *mpk-1(ga117) / dpy-17(e164) unc-79(e1068) III*

CB1417   *lin-3(e1417) IV*

PS5131   *let-23(sy12) / mIn1[dpy-10(e128) mIs14] II*

UP604   *sos-1(cs41) V*

VC450   *ras-1(gk243) II*

RB852   *ras-2(ok682) III*

TZ181   *rap-1(pk2082) IV*

RB848   *rgef-1(ok675) V*

MH538   *mek-2(ku114) I; let-60(n1046) IV*

MT8186   *mpk-1(oz140)/unc-79(e1068) dpy-17(e164) III*

The following strains were produced during this study:

EG314   *ksr-1(ox314) X*

XZ10   *ksr-1(yak10) X*

EG4198   *egl-30(tg26) I ; ksr-1(ox314) X*

XZ1340   *egl-30(tg26) I ; ksr-1(yak10) X*

XZ1511   *egl-30(tg26) I; ksr-1(ok786) X*

660	XZ1465	*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ksr-1(ox314) X*
661	XZ1547	*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ksr-1(yak10) X*
662	XZ2042	*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ksr-1(ok786) X*
663	XZ1615	*egl-30(tg26) I; let-60(n2021) / nT1[qIs51(Pmyo-2::GFP; Ppes-10::GFP; F22B7.9::GFP)] IV*
664	XZ1626	*egl-30(tg26) I; lin-3(e1417) IV*
665	XZ1630	*egl-30(tg26) I; let-23(sy12) / mIn1[dpy-10(e128) mIs14] II*
666	XZ1677	*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; let-60(n2021) / nT1[qIs51(Pmyo-2::GFP;*
667		*Ppes-10::GFP; F22B7.9::GFP)] IV*
668	XZ1548	*egl-30(tg26) I; lin-45(sy96) IV*
669	XZ1556	*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; lin-45(sy96) IV*
670	XZ1850	*egl-30(tg26) mek-2(n1989) / hT2[bli-4(e937) let(q782) qIs48] I*
671	XZ1851	*mek-2(n1989) / hT2[bli-4(e937) let(q782) qIs48] I; nzIs29[Punc-17::rho-1(G14V), unc-*
672		*122::gfp] II*
673	XZ1700	*mpk-1(ga117) / qC1[dpy-19(e1259) glp-1(q339) qIs26(Plag-2::GFP, rol-6(su1006))] III*
674	XZ1668	*egl-30(tg26) I; mpk-1(ga117) / qC1[dpy-19(e1259) glp-1(q339) qIs26(Plag-2::GFP, rol-*
675		*6(su1006))] III*
676	XZ2043	*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; mpk-1(ga117) / qC1[dpy-19(e1259) glp-*
677		*1(q339) qIs26(Plag-2::GFP, rol-6(su1006))] III*
678	XZ2077	*egl-30(tg26) I; let-60(sy93) IV*
679	XZ2106	*egl-30(tg26) I; let-60(sy93) / nT1[qIs51(Pmyo-2::GFP; Ppes-10::GFP; F22B7.9::GFP)] IV*
680	XZ2150	*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; let-60(sy93) / nT1[qIs51(Pmyo-2::GFP;*
681		*Ppes-10::GFP; F22B7.9::GFP)] IV*
682	XZ2158	*egl-30(tg26) I; yakEx158[Prab-3::let-60(S17N)::tbb-2 3'UTR::OPERON::GFP, Pmyo-*
683		*2::mCherry, Pmyo-3::mCherry]*
684	XZ1855	*egl-30(tg26) I; ras-1(gk243) II*
685	XZ1856	*egl-30(tg26) I; ras-2(ok682) III*
686	XZ2045	*egl-30(tg26) I; rap-1(pk2082) IV*
687	XZ2046	*egl-30(tg26) I; sos-1(cs41) V*

XZ2101   *egl-30(tg26) I; rgef-1(ok675) V*

XZ1921   *egl-30(tg26) sur-6(sv30) / hT2[bli-4(e937) let(q782) qIs48] I*

XZ1857   *egl-30(tg26) I ; sur-7(ku119) X*

XZ1854   *nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ras-2(ok682) III*

XZ1852   *nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; rap-1(pk2082) IV*

XZ1880   *yakSi26[Punc-17::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1881   *yakSi27[Punc-17H::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1882   *yakSi28[Punc-17β::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1883   *yakSi29[Punc-47::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1884   *egl-30(tg26) I; yakSi26[Punc-17::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-1(ok786)*

X

XZ1885   *egl-30(tg26) I; yakSi27[Punc-17H::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-*

1(ok786) X

XZ1946   *egl-30(tg26) I; yakSi28[Punc-17β::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-*

1(ok786) X

XZ1947   *egl-30(tg26) I; yakSi29[Punc-47::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-1(ok786)*

X

XZ2015   *unc-119(ed3) III; yakEx122[Punc-17::lin-45* cDNA::tbb-2 3'UTR::OPERON::GFP, Pmyo-*

3::mCherry]

XZ2050   *unc-119(ed3) III; yakIs34[Punc-17::lin-45* cDNA::tbb-2 3'UTR::OPERON::GFP, Pmyo-*

3::mCherry] X

XZ2130   *egl-30(tg26) I; mpk-1(oz140) / oxTi619[Peft-3::TdTomato::H2B cb-unc-119(+)] III*

XZ2131   *egl-30(tg26) mek-2(ku114) I / hT2[bli-4(e937) let(q782) qIs48] I;III*

XZ2119   *ttTi5605 II; unc-119(ed3) III; yakEx154[Punc-17H::lin-45*::tbb-2 3'UTR::OPERON::GFP,*

Pmyo-2::mCherry, Pmyo-3::mCherry]

XZ2154   *yakEx168[Punc-17H::lin-45*::tbb-2 3'UTR::OPERON::GFP, Pmyo-2::mCherry, Pmyo-*

3::mCherry]

XZxxxx   *yakEx[Punc-17H::egl-30(tg26), Pmyo-2::mCherry]* (no strain or allele # given because strain

was not viable enough to be frozen)

Table S2. List of plasmids

<u>Gateway destination vectors</u>

pCFJ150   Gateway destination vector for insertion at chr II Mos site ttTi5605

<u>Gateway entry clones</u>

pEGB05   *Prab-3* [4-1] (pan-neuronal promoter. 1207 bp of the *rab-3* promoter upstream of the

ATG)

pGH1   *Punc-17* [4-1] (acetylcholine neuron promoter. 3229 bp of the *unc-17* promoter upstream

of and including the ATG)

pADA180   *Punc-17H* [4-1] (head acetylcholine neuron promoter. derived from the *unc-17* promoter

of pGH1, carrying a 127 bp deletion that removes the nerve cord β enhancer)

pMA23   *Punc-17β* [4-1] (body acetylcholine neuron promoter. a fusion of the *unc-17* nerve cord β

enhancer to the minimal *unc-17* promoter consisting of the 382 bp upstream of and including the ATG)

pMH522   *Punc-47* [4-1] (GABA neuron promoter. 1254 bp of the *unc-47* promoter upstream of and

including the ATG)

pBC11   *ksr-1* cDNA [1-2]

pBC26   *lin-45* cDNA [1-2]

pBC35   *lin-45** (T626E/T629D) cDNA [1-2]

pBC48   *let-60(S17N)* cDNA

pET97   *egl-30(tg26)* cDNA [1-2]

pADA126   *let-858* 3'UTR [2-3]

pCFJ326   *tbb-2* 3'UTR::OPERON::*GFP* [2-3]

<u>Gateway expression constructs</u>

pBC13   *Prab-3::ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

743	pBC31	*Punc-17::ksr-1* cDNA*::tbb-2* 3'UTR::OPERON::GFP_pCFJ150
744	pBC32	*Punc-17H::ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150
745	pBC33	*Punc-17β::ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150
746	pBC34	*Punc-47::ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150
747	pBC37	*Punc-17::lin-45** cDNA*::tbb-2* 3'UTR::OPERON::GFP_pCFJ150
748	pBC44	*Punc-17H::lin-45** cDNA*::tbb-2* 3'UTR::OPERON::GFP_pCFJ150
749	pBC50	*Prab-3::let-60(S17N)* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150
750	pET102	*Punc-17H::egl-30(tg26)* cDNA::*let-858* 3'UTR_pCFJ150
751		

Table S3. List of primers

753	oBC011	Forward *ksr-1* cDNA for Gateway [1-2] ENTR vector
754		GGGGACAAGTTTGTACAAAAAAGCAGGCTCAatgatgcaaacccaagttgc
755	oBC012	Reverse *ksr-1* cDNA for Gateway [1-2] ENTR vector
756		GGGGACCACTTTGTACAAGAAAGCTGGGTGaaatgtcgactcgtaactttttcatc
757	oBC085	Forward *lin-45* cDNA for Gateway [1-2] ENTR vector
758		GGGGACAAGTTTGTACAAAAAAGCAGGCTcaATGAGTCGGATTAATTTCAAAAAG
759	oBC086	Reverse *lin-45* cDNA for Gateway [1-2] ENTR vector
760		GGGGACCACTTTGTACAAGAAAGCTGGGTgCTAAATGAGACCATAGACATTGTAGTATG
761	oBC094	Reverse *lin-45* T626E mutagenesis
762		gttcactgtccatttcgttttgacctctgccaagccgaaatctccaatttt
763	oBC095	Forward *lin-45* T626E mutagenesis
764		aaaattggagatttcggcttggcagaggtcaaaacgaaatggacagtgaac
765	oBC096	Reverse *lin-45* T629D mutagenesis
766		cctccgttcactgtccatttatctttgacctctgccaagccga
767	oBC097	Forward *lin-45* T629D mutagenesis
768		tcggcttggcagaggtcaaagataaatggacagtgaacggagg
769	oBC105	Forward *let-60* cDNA for Gateway [1-2] ENTR vector
770		GGGGACAAGTTTGTACAAAAAAGCAGGCTCAatgacggagtacaagcttgtgg

771 oBC106   Reverse *let-60* cDNA for Gateway [1-2] ENTR vector

772   GGGGACCACTTTGTACAAGAAAGCTGGGTGtcacattatttgacacttcttcttcttttg

773 oBC109   Reverse *let-60* S17N mutagenesis

774   ttgaatggtgagtgcatttttaccaactcctccatctccaactacca

775 oBC110   Forward *let-60* S17N mutagenesis

776   tggtagttggagatggaggagttggtaaaaatgcactcaccattcaa

777 oET252   Forward *egl-30* cDNA for Gateway [1-2] ENTR vector

778   GGGGACAAGTTTGTACAAAAAAGCAGGCTcaATGGCCTGCTGTTTATCCG

779 oET253   Reverse *egl-30* cDNA for Gateway [1-2] ENTR vector

780   GGGGACCACTTTGTACAAGAAAGCTGGGTgTTACACCAAGTTGTACTCCTTCAG

781 oET254   Forward *egl-30(tg26)* R243Q mutagenesis

782   TGTGACAACGAGAACCGAATGGAAGAATCGAAAGC

783 oET255   Reverse *egl-30(tg26)* R243Q mutagenesis

784   GCTTTCGATTCTTCCATTCGGTTCTCGTTGTCACA

Literature Cited

Afgan E., Baker D., Beek M. van den, Blankenberg D., Bouvier D., *et al.*, 2016 The

Galaxy platform for accessible, reproducible and collaborative biomedical

analyses: 2016 update. Nucleic Acids Res. 44: W3–W10.

Ailion M., Hannemann M., Dalton S., Pappas A., Watanabe S., *et al.*, 2014 Two Rab2

interactors regulate dense-core vesicle maturation. Neuron 82: 167–180.

Bastiani C. A., Gharib S., Simon M. I., Sternberg P. W., 2003 Caenorhabditis elegans

Gαq Regulates Egg-Laying Behavior via a PLCβ-Independent and Serotonin-

Dependent Signaling Pathway and Likely Functions Both in the Nervous System

and in Muscle. Genetics 165: 1805–1822.

Beitel G. J., Clark S. G., Horvitz H. R., 1990 Caenorhabditis elegans ras gene let-60

acts as a switch in the pathway of vulval induction. Nature 348: 503–509.

Boggio E. M., Putignano E., Sassoè-Pognetto M., Pizzorusso T., Giustetto M., 2007

Visual Stimulation Activates ERK in Synaptic and Somatic Compartments of Rat

Cortical Neurons with Parallel Kinetics. PLoS ONE 2: e604.

Boulton T. G., Nye S. H., Robbins D. J., Ip N. Y., Radzlejewska E., *et al.*, 1991 ERKs: A

family of protein-serine/threonine kinases that are activated and tyrosine

phosphorylated in response to insulin and NGF. Cell 65: 663–675.

Brundage L., Avery L., Katz A., Kim U. J., Mendel J. E., *et al.*, 1996 Mutations in a C.

elegans Gqalpha gene disrupt movement, egg laying, and viability. Neuron 16:

999–1009.

Chan J. P., Hu Z., Sieburth D., 2012 Recruitment of sphingosine kinase to presynaptic

terminals by a conserved muscarinic signaling pathway promotes

neurotransmitter release. Genes Dev. 26: 1070–1085.

Charlie N. K., Schade M. A., Thomure A. M., Miller K. G., 2006 Presynaptic UNC-31

(CAPS) is required to activate the G alpha(s) pathway of the Caenorhabditis

elegans synaptic signaling network. Genetics **172**: 943–961.

Chen L., Fu Y., Ren M., Xiao B., Rubin C. S., 2011 A RasGRP, C. elegans RGEF-1b,

Couples External Stimuli to Behavior by Activating LET-60 (Ras) in Sensory

Neurons. Neuron 70: 51–65.

Chong H., Lee J., Guan K. L., 2001 Positive and negative regulation of Raf kinase

activity and function by phosphorylation. EMBO J. 20: 3716–3727.

Cingolani P., Platts A., Wang L. L., Coon M., Nguyen T., *et al.*, 2012 A program for

annotating and predicting the effects of single nucleotide polymorphisms, SnpEff:

SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly

6: 80–92.

Davis M. W., Hammarlund M., Harrach T., Hullett P., Olsen S., *et al.*, 2005 Rapid single

nucleotide polymorphism mapping in C. elegans. BMC Genomics 6: 118.

Dent J. A., Han M., 1998 Post-embryonic expression pattern of C. elegans let-60 ras

reporter constructs. Mech. Dev. 72: 179–182.

DePristo M. A., Banks E., Poplin R., Garimella K. V., Maguire J. R., *et al.*, 2011 A

framework for variation discovery and genotyping using next-generation DNA

sequencing data. Nat. Genet. 43: 491–498.

Doi M., Iwasaki K., 2002 Regulation of retrograde signaling at neuromuscular junctions

by the novel C2 domain protein AEX-1. Neuron 33: 249–259.

Drosten M., Sum E. Y. M., Lechuga C. G., Simón-Carrasco L., Jacob H. K. C., *et al.*,

2014 Loss of p53 induces cell proliferation via Ras-independent activation of the

Raf/Mek/Erk signaling pathway. Proc. Natl. Acad. Sci. U. S. A. 111: 15155–

15160.

Frøkjær-Jensen C., Davis M. W., Ailion M., Jorgensen E. M., 2012 Improved Mos1-

mediated transgenesis in C. elegans. Nat. Methods 9: 117–118.

Guo F. F., Kumahara E., Saffen D., 2001 A CalDAG-GEFI/Rap1/B-Raf cassette couples

M(1) muscarinic acetylcholine receptors to the activation of ERK1/2. J. Biol.

Chem. 276: 25568–25581.

Hamakawa M., Uozumi T., Ueda N., Iino Y., Hirotsu T., 2015 A role for Ras in inhibiting

circular foraging behavior as revealed by a new method for time and cell-specific

RNAi. BMC Biol. 13: 6.

Hammarlund M., Palfreyman M. T., Watanabe S., Olsen S., Jorgensen E. M., 2007

Open syntaxin docks synaptic vesicles. PLoS Biol. **5**: e198.

Han M., Sternberg P. W., 1990 let-60, a gene that specifies cell fates during C. elegans

vulval induction, encodes a ras protein. Cell 63: 921–931.

Han M., Aroian R. V., Sternberg P. W., 1990 The let-60 locus controls the switch

between vulval and nonvulval cell fates in Caenorhabditis elegans. Genetics 126:

899–913.

Han M., Sternberg P. W., 1991 Analysis of dominant-negative mutations of the

Caenorhabditis elegans let-60 ras gene. Genes Dev. 5: 2188–2198.

Han M., Golden A., Han Y., Sternberg P. W., 1993 C. elegans lin-45 raf gene

participates in let-60 ras-stimulated vulval differentiation. Nature 363: 133–140.

Hiley E., McMullan R., Nurrish S. J., 2006 The Galpha12-RGS RhoGEF-RhoA signalling

pathway regulates neurotransmitter release in C. elegans. EMBO J. 25: 5884–

5895.

Hirotsu T., Saeki S., Yamamoto M., Iino Y., 2000 The Ras-MAPK pathway is important

for olfaction in Caenorhabditis elegans. Nature 404: 289–293.

Hirotsu T., Iino Y., 2005 Neural circuit-dependent odor adaptation in C. elegans is

regulated by the Ras-MAPK pathway. Genes Cells 10: 517–530.

Honda Z., Takano T., Gotoh Y., Nishida E., Ito K., *et al.*, 1994 Transfected platelet-

activating factor receptor activates mitogen-activated protein (MAP) kinase and

MAP kinase kinase in Chinese hamster ovary cells. J. Biol. Chem. 269: 2307–

2315.

Hsu V., Zobel C. L., Lambie E. J., Schedl T., Kornfeld K., 2002 Caenorhabditis elegans

lin-45 raf is essential for larval viability, fertility and the induction of vulval cell

fates. Genetics 160: 481–492.

Hu Z., Vashlishan-Murray A. B., Kaplan J. M., 2015 NLP-12 engages different UNC-13

proteins to potentiate tonic and evoked release. J. Neurosci. 35: 1038–1042.

Hunt-Newbury R., Viveiros R., Johnsen R., Mah A., Anastas D., *et al.*, 2007 High-

throughput in vivo analysis of gene expression in Caenorhabditis elegans. PLoS

Biol. 5: e237.

Hyde R., Corkins M. E., Somers G. A., Hart A. C., 2011 PKC-1 acts with the ERK MAPK

signaling pathway to regulate C. elegans mechanosensory response. Genes

Brain Behav. 10: 286–298.

Impey S., Obrietan K., Storm D. R., 1999 Making New Connections. Neuron 23: 11–14.

Jovanovic J. N., Czernik A. J., Fienberg A. A., Greengard P., Sihra T. S., 2000

Synapsins as mediators of BDNF-enhanced neurotransmitter release. Nat.

Neurosci. 3: 323.

Karnoub A. E., Weinberg R. A., 2008 Ras oncogenes: split personalities. Nat. Rev. Mol.

Cell Biol. 9: 517–531.

Kornfeld K., Guan K. L., Horvitz H. R., 1995a The Caenorhabditis elegans gene mek-2

is required for vulval induction and encodes a protein similar to the protein kinase

MEK. Genes Dev. 9: 756–768.

Kornfeld K., Hom D. B., Horvitz H. R., 1995b The ksr-1 gene encodes a novel protein

kinase involved in Ras-mediated signaling in C. elegans. Cell 83: 903–913.

Kushner S. A., Elgersma Y., Murphy G. G., Jaarsma D., Woerden G. M. van, *et al.*,

2005 Modulation of presynaptic plasticity and learning by the H-ras/extracellular

signal-regulated kinase/synapsin I signaling pathway. J. Neurosci. 25: 9721–

9734.

Lackner M. R., Kim S. K., 1998 Genetic analysis of the Caenorhabditis elegans MAP

kinase gene mpk-1. Genetics 150: 103–117.

Lackner M. R., Nurrish S. J., Kaplan J. M., 1999 Facilitation of Synaptic Transmission

by EGL-30 Gqα and EGL-8 PLCβ: DAG Binding to UNC-13 Is Required to

Stimulate Acetylcholine Release. Neuron 24: 335–346.

Lutz S., Freichel-Blomquist A., Yang Y., Rümenapp U., Jakobs K. H., *et al.*, 2005 The

guanine nucleotide exchange factor p63RhoGEF, a specific link between Gq/11-

coupled receptor signaling and RhoA. J. Biol. Chem. 280: 11134–11139.

Lutz S., Shankaranarayanan A., Coco C., Ridilla M., Nance M. R., *et al.*, 2007 Structure

of Galphaq-p63RhoGEF-RhoA complex reveals a pathway for the activation of

RhoA by GPCRs. Science 318: 1923–1927.

Mahoney T. R., Luo S., Nonet M. L., 2006 Analysis of synaptic transmission in

Caenorhabditis elegans using an aldicarb-sensitivity assay. Nat. Protoc. 1: 1772–

1777.

Mao L.-M., Wang J. Q., 2016a Regulation of Group I Metabotropic Glutamate Receptors
by MAPK/ERK in Neurons. J. Nat. Sci. 2: e268.

Mao L.-M., Wang J. Q., 2016b Synaptically Localized Mitogen-Activated Protein
Kinases: Local Substrates and Regulation. Mol. Neurobiol. 53: 6309–6315.

Marais R., Light Y., Paterson H. F., Marshall C. J., 1995 Ras recruits Raf-1 to the
plasma membrane for activation by tyrosine phosphorylation. EMBO J. 14: 3136–
3145.

McMullan R., Hiley E., Morrison P., Nurrish S. J., 2006 Rho is a presynaptic activator of
neurotransmitter release at pre-existing synapses in C. elegans. Genes Dev. 20:
65–76.

McMullan R., Anderson A., Nurrish S., 2012 Behavioral and Immune Responses to
Infection Require Gαq- RhoA Signaling in C. elegans. PLOS Pathog. 8:
e1002530.

Mello C. C., Kramer J. M., Stinchcomb D., Ambros V., 1991 Efficient gene transfer in
C.elegans: extrachromosomal maintenance and integration of transforming
sequences. EMBO J. 10: 3959–3970.

Miller K. G., Emerson M. D., Rand J. B., 1999 Goalpha and diacylglycerol kinase
negatively regulate the Gqalpha pathway in C. elegans. Neuron 24: 323–333.

Nguyen A., Burack W. R., Stock J. L., Kortum R., Chaika O. V., *et al.*, 2002 Kinase

Suppressor of Ras (KSR) Is a Scaffold Which Facilitates Mitogen-Activated

Protein Kinase Activation In Vivo. Mol. Cell. Biol. 22: 3035–3045.

Nicholas H. R., Hodgkin J., 2004 The ERK MAP kinase cascade mediates tail swelling

and a protective response to rectal infection in C. elegans. Curr. Biol. 14: 1256–

1261.

Ortiz J., Harris H. W., Guitart X., Terwilliger R. Z., Haycock J. W., *et al.*, 1995

Extracellular signal-regulated protein kinases (ERKs) and ERK kinase (MEK) in

brain: regional distribution and regulation by chronic morphine. J. Neurosci. 15:

1285–1297.

Reiner D., Lundquist E. A., 2016 Small GTPases. WormBook Online Rev. C Elegans

Biol. doi/10.1895/wormbook.1.67.2

Rhee S. G., 2001 Regulation of phosphoinositide-specific phospholipase C. Annu. Rev.

Biochem. 70: 281–312.

Robbins D. J., Cheng M., Zhen E., Vanderbilt C. A., Feig L. A., *et al.*, 1992 Evidence for

a Ras-dependent extracellular signal-regulated protein kinase (ERK) cascade.

Proc. Natl. Acad. Sci. U. S. A. 89: 6924–6928.

Schmitz S. K., King C., Kortleven C., Huson V., Kroon T., *et al.*, 2016 Presynaptic

inhibition upon CB1 or mGlu2/3 receptor activation requires ERK/MAPK

phosphorylation of Munc18-1. EMBO J. 35: 1236–1250.

Schrader L. A., Birnbaum S. G., Nadin B. M., Ren Y., Bui D., *et al.*, 2006 ERK/MAPK
regulates the Kv4.2 potassium channel by direct phosphorylation of the pore-
forming subunit. Am. J. Physiol. Cell Physiol. 290: C852-861.

Sieburth D. S., Sundaram M., Howard R. M., Han M., 1999 A PP2A regulatory subunit
positively regulates Ras-mediated signaling during Caenorhabditis elegans vulval
induction. Genes Dev. 13: 2562–2569.

Stamboulian S., Choi J.-S., Ahn H.-S., Chang Y.-W., Tyrrell L., *et al.*, 2010 ERK1/2
mitogen-activated protein kinase phosphorylates sodium channel Na(v)1.7 and
alters its gating properties. J. Neurosci. 30: 1637–1647.

Sternberg P. W., 2005 Vulval development. WormBook Online Rev. C Elegans Biol.
doi/10.1895/wormbook.1.6.1

Stokoe D., Macdonald S. G., Cadwallader K., Symons M., Hancock J. F., 1994
Activation of Raf as a result of recruitment to the plasma membrane. Science
264: 1463–1467.

Sun Y., Liu W.-Z., Liu T., Feng X., Yang N., *et al.*, 2015 Signaling pathway of
MAPK/ERK in cell proliferation, differentiation, migration, senescence and
apoptosis. J. Recept. Signal Transduct. Res. 35: 600–604.

Sundaram M., Han M., 1995 The C. elegans ksr-1 gene encodes a novel Raf-related
kinase involved in Ras-mediated signal transduction. Cell 83: 889–901.

Sundaram M. V., 2013 Canonical RTK-Ras-ERK signaling and related alternative

pathways. WormBook Online Rev. C Elegans Biol. doi/10.1895/wormbook.1.80.2

Sweatt J. D., 2004 Mitogen-activated protein kinases in synaptic plasticity and memory.

Curr. Opin. Neurobiol. 14: 311–317.

Thomas G. M., Huganir R. L., 2004 MAPK cascade signalling and synaptic plasticity.

Nat. Rev. Neurosci. 5: 173–183.

Tomida T., Oda S., Takekawa M., Iino Y., Saito H., 2012 The Temporal Pattern of

Stimulation Determines the Extent and Duration of MAPK Activation in a

Caenorhabditis elegans Sensory Neuron. Sci Signal 5: ra76.

Tomioka M., Adachi T., Suzuki H., Kunitomo H., Schafer W. R., *et al.*, 2006 The

insulin/PI 3-kinase pathway regulates salt chemotaxis learning in Caenorhabditis

elegans. Neuron 51: 613–625.

Topalidou I., Cattin-Ortolá J., Pappas A. L., Cooper K., Merrihew G. E. *et al.*, 2016 The

EARP Complex and Its Interactor EIPR-1 Are Required for Cargo Sorting to

Dense-Core Vesicles. PLoS Genet. **12**: e1006074.

Topalidou I., Chen P.-A., Cooper K., Watanabe S., Jorgensen E. M., *et al.*, 2017a The

NCA-1 and NCA-2 Ion Channels Function Downstream of Gq and Rho To

Regulate Locomotion in Caenorhabditis elegans. Genetics 206: 265-282.

Topalidou I., Cooper K., Pereira L., Ailion M., 2017b Dopamine negatively modulates

the NCA ion channels in C. elegans. PLoS Genet. 13: e1007032.

Ueda Y., Hirai S. i, Osada S. i, Suzuki A., Mizuno K., *et al.*, 1996 Protein kinase C

activates the MEK-ERK pathway in a manner independent of Ras and dependent

on Raf. J. Biol. Chem. 271: 23512–23519.

Uozumi T., Hirotsu T., Yoshida K., Yamada R., Suzuki A., *et al.*, 2012 Temporally-

regulated quick activation and inactivation of Ras is important for olfactory

behaviour. Sci. Rep. 2: 500.

van Biesen T., Hawes B. E., Raymond J. R., Luttrell L. M., Koch W. J., *et al.*, 1996 G(o)-

protein alpha-subunits activate mitogen-activated protein kinase via a novel

protein kinase C-dependent mechanism. J. Biol. Chem. 271: 1266–1269.

Vara H., Onofri F., Benfenati F., Sassoè-Pognetto M., Giustetto M., 2009 ERK activation

in axonal varicosities modulates presynaptic plasticity in the CA3 region of the

hippocampus through synapsin I. Proc. Natl. Acad. Sci. U. S. A. 106: 9872–9877.

Wilkie T. M., Scherle P. A., Strathmann M. P., Slepak V. Z., Simon M. I., 1991

Characterization of G-protein alpha subunits in the Gq class: expression in

murine tissues and in stromal and hematopoietic cell lines. Proc. Natl. Acad. Sci.

U. S. A. 88: 10049–10053.

Williams S. L., Lutz S., Charlie N. K., Vettel C., Ailion M., *et al.*, 2007 Trio's Rho-specific

GEF domain is the missing Gαq effector in C. elegans. Genes Dev. 21: 2731–

2746.

Wu Y., Han M., Guan K. L., 1995 MEK-2, a Caenorhabditis elegans MAP kinase kinase,

functions in Ras-mediated vulval induction and other developmental events.

Genes Dev. 9: 742–755.

Yang F., Sun W., Yang Y., Wang Y., Li C.-L., *et al.*, 2015 SDF1–CXCR4 signaling

contributes to persistent pain and hypersensitivity via regulating excitability of

primary nociceptive neurons: involvement of ERK-dependent Nav1.8 up-

regulation. J. Neuroinflammation 12: 219–233.

Yoder J. H., Chong H., Guan K.-L., Han M., 2004 Modulation of KSR activity in

Caenorhabditis elegans by Zn ions, PAR-1 kinase and PP2A phosphatase.

EMBO J. 23: 111–119.

Yoon S., Seger R., 2006 The extracellular signal-regulated kinase: multiple substrates

regulate diverse cellular functions. Growth Factors 24: 21–44.

You Y., Kim J., Cobb M., Avery L., 2006 Starvation activates MAP kinase through the

muscarinic acetylcholine pathway in Caenorhabditis elegans pharynx. Cell

Metab. 3: 237–245.

Zhang H., Koo C. Y., Stebbing J., Giamas G., 2013 The dual function of KSR1: a

pseudokinase and beyond. Biochem. Soc. Trans. 41: 1078–1082.

1018

1019

1020

Figure 1

A

wild type
300 μm
egl-30(tg26)
egl-30(tg26); ksr-1(ok786)

B

ksr-1

yak10
W254>stop

ox314
K463>stop

1000 bp

ok786 deletion

C

wild type
2x amplitude
period

ksr-1(ok786)

egl-30(tg26)

egl-30(tg26); ksr-1(ok786)

D

period/amplitude

wild type
ksr-1(ox314)
ksr-1(yak10)
ksr-1(ok786)
egl-30(tg26)
ksr-1(ox314)
ksr-1(yak10)
ksr-1(ok786)

egl-30(tg26)

***
***
***

Figure 2

Figure 3

A

B

C

D

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

Figure S3

A

B

C