1   **The ERK MAPK pathway modulates Gq-dependent locomotion in**

2   ***Caenorhabditis elegans***

5   Brantley Coleman, Irini Topalidou, and Michael Ailion

Department of Biochemistry, University of Washington, Seattle, WA, 98195

Running Title: The ERK pathway modulates Gq signaling

Keywords: Gq signaling, ERK MAPK pathway, Rho small GTPase, *C. elegans*, G

protein

Corresponding author:

Michael Ailion

Department of Biochemistry

University of Washington

Box 357350

1705 NE Pacific St

Seattle, WA 98195

Phone: 206-685-0111

email: mailion@uw.edu

Abstract

The heterotrimeric G protein Gq regulates neuronal activity through distinct

downstream effector pathways. In addition to the canonical Gq effector phospholipase

Cβ, the small GTPase Rho was recently identified as a conserved effector of Gq. To

identify additional molecules important for Gq signaling in neurons, we performed a

forward genetic screen in the nematode *Caenorhabditis elegans* for suppressors of the

hyperactivity and exaggerated waveform of an activated Gq mutant. We isolated two

mutations affecting the MAP kinase scaffold protein KSR-1 and found that KSR-1

modulates locomotion downstream of or in parallel to the Gq-Rho pathway. Through

epistasis experiments, we found that the core ERK MAPK cascade is required for Gq-

Rho regulation of locomotion, but that the canonical ERK activator LET-60/Ras may not

be required. Through neuron-specific rescue experiments, we found that the ERK

pathway functions in acetylcholine neurons to control Gq-dependent locomotion.

Additionally, expression of activated LIN-45/Raf in acetylcholine neurons is sufficient to

cause an exaggerated waveform phenotype and hypersensitivity to the

acetylcholinesterase inhibitor aldicarb, similar to an activated Gq mutant. Taken

together, our results suggest that the ERK MAPK pathway modulates the output of Gq-

Rho signaling to control locomotion behavior in *C. elegans.*

Introduction

The heterotrimeric G protein Gq is a conserved regulator of neurotransmission in metazoans. Gq is highly expressed in neurons in mammals and in the nematode *C. elegans* (Wilkie *et al.* 1991; Lackner *et al.* 1999). In its canonical signaling pathway, Gq activates phospholipase Cβ (PLCβ) to cleave the membrane lipid phosphatidylinositol 4,5-bisphosphate (PIP2) into inositol trisphosphate (IP3) and diacylglycerol (DAG) (Rhee 2001). An increased DAG concentration at the synapse helps trigger synaptic vesicle release (Miller *et al.* 1999; Lackner *et al.* 1999).

In addition to activating PLCβ, Gq directly binds and activates the Rho guanine nucleotide exchange factor (GEF) Trio, which in turn activates the small GTPase Rho (Lutz *et al.* 2005, 2007; Williams *et al.* 2007). In mature *C. elegans* neurons, the Rho ortholog RHO-1 regulates synaptic activity through multiple G protein-dependent mechanisms. First, RHO-1 acts downstream of the G_{12}-class G protein GPA-12 by binding to and inhibiting the diacylglycerol kinase DGK-1. Inhibition of DGK-1 allows DAG to accumulate at the synapse, thereby increasing synaptic vesicle release (McMullan *et al.* 2006; Hiley *et al.* 2006). Second, Gq-Rho signaling promotes neurotransmitter release by recruiting the sphingosine kinase SPHK-1 to presynaptic terminals (Chan *et al.* 2012). Third, Gq-Rho signaling positively regulates the NCA-1/NALCN cation channel to regulate locomotion (Topalidou *et al.* 2017a). Here we identify the extracellular signal-related kinase mitogen-activated protein kinase (ERK MAPK) pathway as a positive regulator of neuronal activity acting downstream of or in parallel to Gq and Rho in acetylcholine neurons.

ERK MAPK signaling acts extensively in animal development, cellular

proliferation, and cancer signaling (Yoon and Seger 2006; Karnoub and Weinberg 2008;

Sun *et al.* 2015). ERKs are highly expressed in mammalian neurons (Boulton *et al.*

1991; Ortiz *et al.* 1995) and act in both the nucleus and at the synapse to regulate

synaptic activity and plasticity (Thomas and Huganir 2004; Sweatt 2004; Mao and Wang

2016b). In *C. elegans*, the ERK pathway is required for multiple developmental events

including specification of the vulva (Sternberg 2005; Sundaram 2013), and also acts in

several types of neurons to control behavior. ERK signaling is activated in response to

odorants in the AWC sensory neuron to regulate chemotaxis to volatile odorants and in

AIY interneurons to mediate odor adaptation (Hirotsu *et al.* 2000; Hirotsu and Iino 2005;

Chen *et al.* 2011; Uozumi *et al.* 2012). ERK is also activated in the ASER sensory

neuron to regulate chemotaxis to salt (Tomioka *et al.* 2006; Tomida *et al.* 2012). ERK

signaling regulates foraging behavior by acting in the IL1, OLQ, and RMD neurons

(Hamakawa *et al.* 2015). Finally, the ERK pathway has been shown to act in

interneurons to regulate the nose touch response, a mechanosensory behavior (Hyde *et*

al. 2011).

In the canonical ERK MAPK pathway, extracellular ligand binding activates

transmembrane receptor tyrosine kinases (RTKs), and adaptor proteins recruit a GEF to

activate the small GTPase Ras (LET-60 in *C. elegans*). Upon Ras activation, LIN-45/Raf

translocates to the plasma membrane where it interacts with Ras and the scaffold

protein KSR-1. KSR-1 facilitates the activation of LIN-45/Raf and the subsequent

phosphorylation of the MAPK cascade consisting of LIN-45/Raf, MEK-2/MEK, and MPK-

1/ERK (Sundaram 2013). In this study, we found that the ERK MAPK pathway

consisting of KSR-1, LIN-45/Raf, MEK-2/MEK and MPK-1/ERK modulates Gq-Rho

signaling in acetylcholine neurons, but that surprisingly LET-60/Ras may not be

required.

Materials and Methods

C. elegans strains

All strains were cultured using standard methods and were maintained at 20°C.

Table S1 contains all the strains used in this study.

Isolation and mapping of the ksr-1(ox314) and ksr-1(yak10) mutations

The *ox314* and *yak10* mutants were isolated from an ENU mutagenesis

suppressor screen of the activated Gq mutant *egl-30(tg26)* (Ailion *et al.* 2014). We

mapped the *ox314* mutation by its activated Gq suppression phenotype using single

nucleotide polymorphisms (SNPs) in the Hawaiian strain CB4856 as described (Davis *et*

al. 2005).The *ox314* mutation was mapped to an approximately 709 kb region in the

middle of the X chromosome between SNPs on cosmids F45E1 and F53A9 (SNPs

F45E1[1] and pkP6158). This region included 159 predicted protein-coding genes. A

complementation test of *ox314* and *yak10* in the *egl-30(tg26)* background showed these

to be alleles of the same gene. Whole genome sequencing (see below) identified these

as mutations in *ksr-1*, and we confirmed this by performing a complementation test with

the deletion allele *ksr-1(ok786).*

Whole genome sequencing

Strains EG4198 *egl-30(tg26); ox314* and XZ1340 *egl-30(tg26); yak10* were

sequenced to identify candidate mutations. DNA was purified according to the Hobert

Lab protocol (http://hobertlab.org/whole-genome-sequencing/). Ion Torrent sequencing

was performed at the University of Utah DNA Sequencing Core Facility. Each data set

contained roughly 18,400,000 reads of a mean read length of 160 bases, resulting in

about 30X average coverage of the *C. elegans* genome. The sequencing data were

processed on the Galaxy server at usegalaxy.org (Afgan *et al.* 2016). SNPs and indels

were identified and annotated using the Unified Genotyper and SnpEff tools (DePristo *et*

al. 2011; Cingolani *et al.* 2012). After filtering for mutations in open reading frames, we

found each strain to have unique stop mutations in *ksr-1*, in the middle of the interval

where we mapped *ox314*. *ox314* is a G to A transition that causes a stop codon at

amino acid K463, and *yak10* is an A to T transversion that causes a stop codon at

W254.

Locomotion assays

Track waveform and radial locomotion assays were performed on 10 cm

nematode growth medium (NGM) plates seeded with 400 µl of OP50 *E. coli* culture and

spread with sterile glass beads. Bacterial lawns were grown at 37°C for 16 hrs and the

plates were stored at 4°C until needed. For track waveform measurements, five first day

adult animals were placed on a plate and allowed to roam for 2-5 min. We then

recorded each animal's tracks following forward locomotion. Track pictures were taken

at 40X on a Nikon SMZ18 microscope with the DS-L3 camera control system. Pictures

of worm tracks were processed using ImageJ. Period and 2X amplitude were measured

freehand using the line tool. For each worm, we calculated the average

period/amplitude ratio of five individual track bends (Figure 1C). For assays with the

temperature sensitive allele *sos-1(cs41)*, all strains were grown at 20°C and shifted to

the non-permissive temperature of 25°C for 24 hours before being assayed. For radial

locomotion assays, ten to fifteen first day adult animals were picked to the center of a

plate and were then allowed to move freely for 40 minutes. The positions of worms were

marked and the distances of the worms from the starting point were measured. For all

waveform and radial locomotion assays, the experimenter was blind to the genotypes of

the strains assayed.

Microscopy

Photographs of moving worms were taken at 60X on a Nikon SMZ18 microscope

with the DS-L3 camera control system. The worms were age-matched as first day

adults grown at 20°C.

Constructs and transgenes

Plasmids were constructed using the three-slot multisite Gateway cloning system

(Invitrogen). Plasmids and primers used are found in Tables S2 and S3. The *ksr-1* and

lin-45 cDNAs were amplified by PCR from worm cDNA library and cloned into [1-2]

Gateway entry vectors. Activating Raf mutations T626E/T629D were introduced into the

lin-45 cDNA vector by two sequential site-directed mutagenesis reactions (Q5 kit, NEB)

with primers oBC094/095 and oBC096/097, respectively, and then confirmed by

sequencing. The *ksr-1* and activated *lin-45* cDNAs were cloned into expression

constructs under different neuronal promoters using the multisite Gateway system.

Proper expression of *ksr-1* and activated *lin-45* was confirmed by including an operon

GFP::H2B in the [2-3] slot of the expression constructs. The operon GFP template *tbb-2*

3'utr::gpd-2 operon::GFP::H2B:cye-1 3'utr (Frøkjær-Jensen *et al.* 2012) results in

untagged proteins whose expression can be monitored by GFP expression.

Injections and chromosomal integrations

Worms carrying the activated *lin-45* transgenes *Punc-17::lin-45** and *Punc-*

17H::lin-45* as extrachromosomal arrays were generated by injecting pBC37 or pBC44

at 20 ng/µL or 10 ng/ µL respectively along with co-injection markers pCFJ104 (*Pmyo-*

3::mCherry) at 5 ng/µL, pCFJ90 (*Pmyo-2::mCherry*) at 2.5 ng/µL, and the carrier DNA

Litmus 38i to a final concentration of 100 ng/µL DNA (Mello *et al.* 1991). MosSCI lines

were generated as described (Frøkjær-Jensen *et al.* 2012) using an injection mix

containing 10-15 ng/µL targeting vector, 50 ng/µL pCFJ601 (*Peft-3::Mos1* transposase),

negative selection markers pGH8 (*Prab-3::mCherry*) at 10 ng/µL, pCFJ104 (*Pmyo-*

3::mCherry) at 5 ng/µL, pCFJ90 (*Pmyo-2::mCherry*) at 2.5 ng/µL, pMA122

(*Phsp16.41::peel-1*) at 10 ng/µL, and carrier DNA Litmus 38i to a final concentration of

100 ng/µL DNA.

Extrachromosomal arrays were integrated into the genome by exposure to 4000

rads of gamma irradiation. Irradiated young adult hermaphrodites were transferred to 10

cm OP50 plates (5 worms/plate) and grown to starvation. The plates were chunked and

grown to starvation twice more to enrich for stably expressing lines. When nearly

starved, 8 animals per plate were picked to individual plates. The progeny were then

screened for 100% stable transmission, indicating integration into the genome.

Integration was confirmed by mapping the transgene to a chromosome.

Aldicarb assays

35 mm aldicarb assay plates were poured with NGM supplemented with 1 mM

aldicarb. The plates were seeded with 5 uL OP50 and dried at room temperature

overnight. Animals were picked onto the OP50 lawn to begin the assay (time 0) and

then kept at room temperature. Every 15 minutes, animals were scored for paralysis by

lightly touching the nose of the animal with an eyebrow hair. Animals were scored as

paralyzed if the worm displayed no locomotor response to three nose touches and had

no pharyngeal pumping. Animals that left the OP50 lawn were picked back onto the

food.

Statistical analysis

P values were determined using GraphPad Prism 5. Normally distributed data

sets were analyzed with a one-way ANOVA and Bonferroni's *post hoc* test when group

size was unequal, or with Tukey's *post hoc* test when group size was equal. Data sets

with non-normal distribution (using the Shapiro-Wilk normality test) were analyzed with

a Kruskal-Wallis test and Dunn's *post hoc* test. Data sets with multiple independent

variables were analyzed by two-way ANOVA and Bonferroni's *post hoc* test.

Reagent and data availability

Strains and plasmids are listed in Tables S1 and S2 and are available upon

request. Primers are listed in Table S3. The authors state that all data necessary for

confirming the conclusions presented in the article are represented fully within the

article and Supplemental Material.

Results

KSR-1 regulates locomotion downstream of Gq

In *C. elegans,* the heterotrimeric G protein Gq regulates synaptic vesicle release

(Hu *et al.* 2015). Gq is a key regulator of neuromuscular activity, as loss-of-function

mutants in *egl-30* are nearly paralyzed (Brundage *et al.* 1996) whereas the gain-of-

function mutant *egl-30(tg26)* has hyperactive locomotion with an exaggerated loopy

waveform (Doi and Iwasaki 2002; Bastiani *et al.* 2003) (Figure 1A, C, D). To identify

pathways required for Gq signaling, we performed a forward genetic screen in *C.*

elegans for suppressors of the activated Gq mutant *egl-30(tg26)*. Two suppressors

identified in this screen, *ox314* and *yak10*, showed similar suppression of the loopy

waveform and hyperactivity of *egl-30(tg26)* animals (Figure 1D). When crossed away

from the *egl-30(tg26)* background, both mutants moved with wild-type waveform (Figure

1D), but at a slightly slower rate. We mapped the *ox314* allele near the center of the X

chromosome (see Materials and Methods), and a complementation test showed that

ox314 and *yak10* are mutations in the same gene since they fail to complement in an

egl-30(tg26) background.

We used whole genome sequencing to identify *ox314* and *yak10* as nonsense

mutations in *ksr-1* (Figure 1B, see Materials and Methods). KSR-1 is a scaffold protein

that facilitates the localization and interactions required for the Ras-mitogen activated

protein kinase (MAPK) cascade consisting of Raf, MEK and ERK (Kornfeld *et al.* 1995b;

Sundaram and Han 1995; Nguyen *et al.* 2002; Zhang *et al.* 2013). The deletion allele

ksr-1(ok786) also suppressed the loopy waveform of the activated Gq mutant identically

to *ox314* and *yak10.* These results suggest that KSR-1 activity is required for regulation

of locomotion rate and waveform by Gq.

The ERK MAPK cascade acts to promote Gq signaling

Because the loss of the MAPK scaffold *ksr-1* suppresses the activated Gq

mutant *egl-30(tg26)*, we asked whether other components of the Ras-ERK pathway

would also suppress. Since the core components of the Ras-ERK pathway are required

for viability, we built double mutants of activated Gq with reduction-of-function mutations

in genes at each step of the ERK cascade. Mutations in Raf (*lin-45(sy96)*)*,* MEK (*mek-*

2(n1989), *mek-2(ku114)*), and ERK (*mpk-1(ga117)*, *mpk-1(oz140)*) all suppressed the

loopy waveform of activated Gq animals similarly to *ksr-1(ok786)* (Figure 2A,B; Figure

S1A,B)*.* However, mutations in Ras (*let-60(n2021)*) and the upstream pathway

activators EGF (*lin-3(e1417)*) and the EGFR (*let-23(sy12)*) did not suppress activated

Gq (Figure 2C). Because *let-60* is required for viability, most *let-60* alleles including

n2021 are partial loss-of-function (Beitel *et al.* 1990). We also analyzed the dominant

negative D119N allele *let-60(sy93)* that disrupts Ras binding to guanine nucleotides and

thus prevents Ras activation (Han and Sternberg 1991). We found that *let-60(sy93)* also

did not suppress the loopy waveform of activated Gq (Figure 2D), supporting the

possibility that ERK activation occurs through a Ras-independent mechanism.

Because partial loss-of-function mutations in the ERK MAPK pathway genes

downstream of Ras showed clear suppression of activated Gq, we were surprised to

find that partial loss-of-function mutations in Ras did not suppress. If LET-60/Ras is

indeed not required, Gq might instead activate the ERK pathway via other Ras-

subfamily proteins. To test this possibility, we made double mutants of activated Gq with

putative null alleles of R-Ras/*ras-1,* M-Ras/*ras-2,* and Rap1/*rap-1* and found that they

also did not suppress activated Gq (Figure S2A). To further investigate whether this

pathway acts independently of Ras, we tested mutations in GEFs that activate Ras. The

temperature-sensitive RasGEF mutant *sos-1(cs41)* did not suppress activated Gq when

shifted to the non-permissive temperature (Figure S2B). Additionally, a null mutation in

the neuronal RasGEF *rgef-1* also did not suppress activated Gq locomotion (Figure

S2C), supporting the possibility that this pathway is Ras-independent. In genetic

screens for vulval induction mutants, additional factors such as the PP2A subunit *sur-6*

(Sieburth *et al.* 1999) and ion transporter *sur-7* (Yoder *et al.* 2004) were identified as

positive regulators of Ras-ERK activity. However, the *sur-6(sv30)* and *sur-7(ku119)*

mutations did not suppress activated Gq locomotion (data not shown). These data

suggest either that only a low level of Ras activity is needed to properly activate ERK

signaling downstream of Gq, or that ERK signaling acts independently of LET-60/Ras

and other known *C. elegans* Ras family proteins to regulate locomotion downstream of

Gq.

KSR-1 and the ERK MAPK cascade modulate Rho signaling

Three classes of suppressor mutations were isolated in our forward genetic

screen of activated Gq, as characterized by their molecular role and unique suppression

phenotypes (Topalidou *et al.* 2017a; b). We grouped together a class of suppressor

mutations including *ox314, yak10*, and the RhoGEF Trio (*unc-73* in *C. elegans*) by their

characteristic strong suppression of the loopy waveform of activated Gq (Topalidou *et*

al. 2017a; b), suggesting that *ksr-1* might act in the same pathway as *unc-73*.

We have shown that Gq regulates locomotion via the small GTPase Rho (RHO-1

in *C. elegans*) (Topalidou *et al.* 2017a). Transgenic expression of an activated RHO-1

mutant (G14V) in acetylcholine neurons (here called "*rho-1(gf)*") causes worms to have

a loopy waveform and impaired locomotion (McMullan *et al.* 2006) (Figure 3A). To

examine whether *ksr-1* acts in the Gq-Rho pathway we tested whether mutations in *ksr-*

1 suppress the phenotypes of *rho-1(gf)* worms. We found that the *ksr-1* alleles *ok786,*

ox314, and *yak10* all suppressed the loopy waveform of *rho-1(gf)* worms (Figure 3A).

Because *rho-1(gf)* worms have a slow locomotion rate and loopy waveform, these

mutants do not efficiently travel long distances. We used radial locomotion assays (see

Materials and Methods) to quantify the locomotion phenotype of *rho-1(gf)* worms. *rho-*

1(gf) ksr-1 double mutants had an increased radial distance traveled compared to *rho-*

1(gf) alone (Figure 3B). These data suggest that KSR-1 acts downstream of or in

parallel to the Gq-Rho pathway to regulate locomotion.

Since *ksr-1* mutants suppress the exaggerated waveform of both activated Gq

and activated Rho animals, we expected that loss of other ERK pathway components

would also suppress activated Rho. We made double mutants of activated Rho (*rho-*

1(gf)) with reduction-of-function alleles of the Ras-ERK pathway and found that

mutations in Raf, MEK, and ERK suppressed the *rho-1(gf)* loopy waveform and

decreased locomotion phenotypes (Figure 3A,C). However, the *let-60(n2021)* Ras

mutation did not suppress the loopy waveform or radial locomotion defect of *rho-1(gf)*

worms (Figure 3A,C). These data suggest that the ERK pathway acts downstream of or

in parallel to the Gq-Rho pathway to regulate locomotion, possibly in a Ras-independent

manner.

308   The ERK MAPK cascade acts in acetylcholine neurons to control locomotion

Members of the ERK pathway are expressed in neurons in *C. elegans* (Dent and

Han 1998; Hunt-Newbury *et al.* 2007), and Gq and Rho act in acetylcholine neurons to

promote synaptic release and regulate locomotion(Lackner *et al.* 1999; McMullan *et al.*

2006) To determine whether the ERK pathway also acts in neurons to modulate Gq

signaling, we expressed the *ksr-1* cDNA under promoters driving expression in specific

types of neurons. Single-copy transgenic expression of *ksr-1* under an acetylcholine

neuron promoter (*Punc-17*) or under a head acetylcholine neuron promoter (*Punc-17H)*

fully reversed the *ksr-1* suppression of the loopy waveform of activated Gq worms

(Figure 4). *ksr-1* expression in ventral cord acetylcholine motor neurons (*Punc-17β*) or

GABA neurons (*Punc-47)* did not significantly reverse the *ksr-1* suppression of activated

Gq (Figure 4). This suggests that ERK signaling primarily functions in the acetylcholine

interneurons or motor neurons of the head to modulate Gq-dependent locomotion.

We have shown that the ERK pathway is necessary for Gq-dependent effects on

locomotion. To determine whether ERK signaling is sufficient to modulate locomotion,

we expressed an activated form of *lin-45*/Raf specifically in acetylcholine neurons. Raf

kinase activity is regulated via conserved phosphorylation events, and phosphomimetic

mutations T626E/T629D in the kinase activation loop of *lin-45* are sufficient to confer

constitutive Raf activity (Chong *et al.* 2001). We found that expression of activated Raf

in acetylcholine neurons (*Punc-17*) causes a loopy waveform similar to activated Gq

and Rho mutants and similar limited dispersal in radial locomotion assays (Figure 5A,

B).

To determine if Raf activation affects acetylcholine release, we assayed for

sensitivity to the acetylcholinesterase inhibitor aldicarb. Mutants with reduced

acetylcholine secretion are resistant to aldicarb, whereas mutants with increased

acetylcholine secretion are hypersensitive to aldicarb (Mahoney *et al.* 2006). Activated

Gq and Rho mutants have increased rates of paralysis when exposed to aldicarb

(Lackner *et al.* 1999; McMullan *et al.* 2006) . We found that expression of activated Raf

in acetylcholine neurons also led to aldicarb hypersensitivity, similar to activated Gq and

Rho mutants (Figure 5C). However, we found that a *ksr-1* mutation does not suppress

the aldicarb hypersensitivity of an activated Gq mutant, and the *ksr-1* mutant on its own

has similar aldicarb sensitivity to wild type (Figure 5D). These results suggest that the

ERK pathway is not necessary for synaptic transmission, but is sufficient to stimulate

synaptic transmission when constitutively activated.

Discussion

In this study we identified KSR-1 and the ERK MAPK cascade as downstream

modulators of Gq-Rho signaling. We found that ERK signaling acts in the acetylcholine

interneurons or motor neurons of the head to modulate Gq-dependent locomotion. We

also found that activation of the MAPKKK Raf/LIN-45 in acetylcholine neurons causes

increased sensitivity to the acetylcholinesterase inhibitor aldicarb, probably reflecting

increased release of acetylcholine at neuromuscular synapses. Our data support the

model that Gq-Rho activation of the ERK pathway may be independent of its canonical

regulator, the small GTPase Ras/LET-60 (Figure 6).

The ERK signaling cascade has been well-studied for its regulation of cellular

proliferation and differentiation (Sun *et al.* 2015), but also plays important roles in

mature neurons and has been associated with synaptic plasticity and memory (Impey *et*

al. 1999). In addition to activating transcription, ERKs regulate synaptic plasticity both

presynaptically and postsynaptically by phosphorylating relevant substrates. ERKs

phosphorylate the presynaptic proteins synapsin I and Munc18-1, and postsynaptic

proteins such as scaffolds, $K_v4.2$ potassium channels, and Group I metabotropic

glutamate receptors (Jovanovic *et al.* 2000; Thomas and Huganir 2004; Sweatt 2004;

Kushner *et al.* 2005; Boggio *et al.* 2007; Vara *et al.* 2009; Mao and Wang 2016a;

Schmitz *et al.* 2016). Our findings suggest that the ERK pathway controls Gq-dependent

locomotion in *C. elegans* and that activated ERK promotes synaptic transmission.

In contrast to many developmental and neuronal roles of Ras-dependent ERK

signaling in *C. elegans,* our data suggest that Raf-MEK-ERK signaling may modulate

Gq-Rho output independently of Ras/LET-60. One caveat to the conclusion that the

ERK pathway acts independently of Ras to control Gq-dependent locomotion is that the

alleles of *let-60*/Ras tested here are not null (Han and Sternberg 1990, 1991; Han *et al.*

1990; Beitel *et al.* 1990). However, other than *mpk-1(ga117)*, the alleles we used of *lin-*

45, *mek-2* and *mpk-1* are also not null (Han *et al.* 1993; Kornfeld *et al.* 1995a; Wu *et al.*

1995; Lackner and Kim 1998; Hsu *et al.* 2002)**,** yet were able to suppress an activated

Gq mutant. Furthermore, the *let*-60/Ras alleles used here have stronger phenotypes in

vulval development than the two weak *mek-2*/MEK alleles we used (*n1989* and *ku114*)

(Beitel *et al.* 1990; Kornfeld *et al.* 1995a; Wu *et al.* 1995) and the *let-60(n2021)*/Ras

mutant has comparable phenotypes to the *lin-45(sy96)*/Raf and *mpk-1(ga117)*/ERK

alleles for chemotaxis to odorants and the regulation of foraging behavior (Hirotsu *et al.*

2000; Hamakawa *et al.* 2015). Thus, either LET-60/Ras is not required to modulate Gq

signaling, or ERK signaling in the locomotor circuit requires a lower threshold of Ras

activity and the *let-60* mutants we used have sufficient levels of active Ras to modulate

Gq signaling, but not enough for vulval induction or other behaviors.

If Ras is not required, how instead might KSR-Raf-MEK-ERK signaling be

activated? Normally, active Ras recruits Raf to the plasma membrane but it has been

shown that artificial recruitment of Raf to the plasma membrane in the absence of Ras

is sufficient to activate Raf signaling (Stokoe *et al.* 1994; Marais *et al.* 1995). Thus,

another possible way to activate the ERK pathway would be for Raf to be recruited to

the membrane by proteins other than Ras. The most obvious candidates for recruiting

Raf are other members of the Ras family such as Rap1 or R-Ras that have a conserved

effector-binding domain (Reiner and Lundquist 2016); it has been reported that Rap1

can mediate Ras-independent activation of Raf downstream of a Gq-coupled receptor

(Guo *et al.* 2001). However, mutations in the worm orthologs of Rap1/RAP-1 and R-

Ras/RAS-1 did not suppress the activated Gq mutant, indicating that these Ras-like

proteins are not required, at least individually, to activate Raf in this pathway. It is

possible that more than one of these Ras family members function redundantly to

activate Raf, or that Raf is activated completely independently of Ras family proteins.

There have been other reported cases of Ras-independent activation of the Raf-MEK-

ERK pathway (Robbins *et al.* 1992; Honda *et al.* 1994; van Biesen *et al.* 1996; Ueda *et*

al. 1996; Drosten *et al.* 2014), some of which involve G protein signaling and protein

kinase C, though the precise mechanisms involved are unclear.

Gq, Rho, and ERK are also required for the *C. elegans* behavioral and immune

response to infection by the bacterium *M. nematophilum* (Nicholas and Hodgkin 2004;

McMullan *et al.* 2012). Interestingly, LET-60/Ras was reported to not be required for the

immune response (Nicholas and Hodgkin 2004) or only partially required (McMullan *et*

al. 2012). Additionally, LET-60/Ras was not required for the increased sensitivity to

aldicarb caused by infection (McMullan *et al.* 2012), at least as assayed using the *let-*

60(n2021) allele. Thus, the same neuronal Ras-independent ERK pathway we describe

here appears to also modulate Gq signaling in the neuronal response to bacterial

infection and possibly the innate immune response as well.

Our epistasis analysis suggests that ERK signaling acts genetically downstream

of or in parallel to Gq-Rho signaling. How might Gq-Rho signaling lead to ERK

activation? ERK activation could occur via a linear pathway downstream of Gq and Rho,

or ERK could signal in parallel and converge downstream of Rho to affect neuronal

activity (Figure 6). There is precedence for Gq activating ERK via a linear pathway. In

the pharyngeal muscle of *C. elegans*, ERK activity is increased by Gq-dependent

signaling through protein kinase C (You *et al.* 2006). In the AWC olfactory neuron, ERK

activity is increased downstream of Gq signaling via the RasGEF RGEF-1 (Chen *et al.*

2011; Uozumi *et al.* 2012). Protein kinase C and RGEF-1 are both activated by DAG,

probably produced by the canonical Gq-PLCβ pathway. By contrast, we found that ERK

signaling regulates locomotion by modulating the output of the Gq-Rho pathway that

acts in parallel to Gq-PLCβ, and does not depend on RGEF-1.

How might the ERK pathway modulate neuronal activity downstream of Gq-Rho

signaling? Rho regulates neuronal activity and synaptic release through several

mechanisms in *C. elegans* neurons, any of which could be targets of ERK signaling.

One possible ERK effector is the NCA/NALCN cation channel that acts genetically

downstream of Gq-Rho to regulate locomotion rate and waveform (Topalidou *et al.*

2017a; b). Though ERK has not been directly connected to NCA/NALCN, mammalian

ERK regulates neuronal excitability by directly phosphorylating voltage-gated sodium

and potassium channels (Schrader *et al.* 2006; Stamboulian *et al.* 2010) and by

regulating channel expression (Yang *et al.* 2015). Given the many possible ways that

ERK may regulate excitability or synaptic transmission, *C. elegans* genetics is well

suited to determine relevant effectors of Gq-Rho-ERK signaling.

Acknowledgements

We thank Stephen Nurrish for the activated Rho worm strain; Jordan and Jill Hoyt for help with Galaxy analysis; Laura Taylor for help with irradiation of extrachromosomal arrays; Dana Miller for use of her Nikon SMZ18 microscope and camera. Some strains were provided by the CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). This work was supported by an Ellison Medical Foundation New Scholar Award and by NIH grants R00 MH082109 and R56 NS100843 to M.A.

Figure Legends

Figure 1. *ksr-1* mutations suppress activated Gq.

(A) A *ksr-1* mutation suppresses the exaggerated body bends of activated Gq. The activated Gq mutant *egl-30(tg26)* has small size and deep body bends. *ksr-1(ok786)* suppresses the exaggerated body bends and small size of *egl-30(tg26)* worms.

(B) Gene structure of *ksr-1* locus. Locations of the *egl-30(tg26)* suppressor alleles *ox314* and *yak10* are indicated, as well as the position of the *ok786* deletion. The gene structure was drawn using Exon-Intron Graphic Maker (www.wormweb.org/exonintron)

(C) A *ksr-1* mutation suppresses the exaggerated body bend waveform of *egl-30(tg26)* mutants. Straightened images of tracks left in bacterial lawns show similar waveform for wild type and *ksr-1(ok786)* worms. *egl-30(tg26)* mutants have an exaggerated waveform, creating tracks with a large amplitude relative to the period. *ksr-1*(*ok786*) suppresses the exaggerated body bends of *egl-30(tg26)* mutants.

(D) A *ksr-1* mutation suppresses the *egl-30(tg26)* waveform. *ksr-1* nonsense alleles *ox314* and *yak10* and the deletion allele *ok786* strongly suppress the *egl-30(tg26)* exaggerated waveform. $N \geq 12$ *** $P < 0.001$, one-way ANOVA with Bonferroni's *post hoc* test.

Figure 2. Mutations in the ERK MAPK pathway suppress activated Gq.

(A) Mutations in the MAPKKK *lin-45*/Raf, the MAPKK *mek-2*/MEK, and the MAPK *mpk-1*/ERK suppress the exaggerated waveform of *egl-30*(*tg26*) worms. Partial loss of Ras *let-60* activity does not suppress activated Gq waveform.

(B) Waveform quantification of ERK pathway mutants show levels of *egl-30(tg26)*

suppression similar to *ksr-1* alleles. $N \geq 12$ *** $P < 0.001$, one-way ANOVA with

Bonferroni's *post hoc* test.

(C) Signaling pathways upstream of the ERK pathway do not suppress activated Gq.

Mutations in the EGFR (*let-23*) or the EGF ligand (*lin-3*) do not affect the exaggerated

waveform of *egl-30(tg26)* animals. The Ras partial loss-of-function mutation *let-*

60(n2021) does not suppress activated Gq waveform. $N \geq 12$ *** $P < 0.001$, one-way

ANOVA with Bonferroni's *post hoc* test.

(D) The *let-60(sy93)* dominant negative mutation in Ras does not suppress activated Gq

waveform. $N \geq 13$, n.s., not significant, one-way ANOVA with Bonferroni's *post hoc* test.

Figure 3. Mutations in *ksr-1* and the ERK pathway suppress activated Rho.

(A) Mutations in the ERK pathway visibly suppress the exaggerated body bends of

animals expressing an activated Rho mutant (G14V) in acetylcholine neurons

(*nzIs29[Punc-17::rho-1(gf)]*). Reduction of LET-60/Ras activity does not suppress the

activated Rho waveform.

(B) The *ksr-1(ok786), ksr-1(ox314),* and *ksr-1(yak10)* mutations suppress the

locomotion of activated Rho animals as shown by radial locomotion assay. $N \geq 38$ *** P

< 0.001, ** $P < 0.01$, Kruskal-Wallis test with Dunn's *post hoc* test.

(C) Mutations in *lin-45* and *mek-2* suppresses the locomotion defect of activated Rho

animals as shown by radial locomotion assays. The *let-60(n2021)* and *mpk-1(ga117)*

mutations do not significantly suppress activated Rho locomotion. $N \geq 50$ *** $P < 0.001$,

Kruskal-Wallis test with Dunn's *post hoc* test.

Figure 4. KSR-1 acts in acetylcholine neurons to modulate Gq signaling.

Single-copy expression of the *ksr-1* cDNA exclusively in acetylcholine neurons (*Punc-17*, *yakSi26* transgene) or head acetylcholine neurons (*Punc-17H, yakSi27* transgene) is sufficient to reverse the *ksr-1* suppression of the loopy waveform of activated Gq animals. *ksr-1* expression in ventral cord acetylcholine neurons (*Punc-17β*, *yakSi28* transgene) or GABA neurons (*Punc-47*, *yakSi29* transgene) does not reverse the *ksr-1* suppression of the activated Gq exaggerated waveform. $N \geq 12$ *** $P < 0.001$, ** $P < 0.01$, n.s., not significant, Kruskal-Wallis test with Dunn's *post hoc* test.

Figure 5. Increased ERK signaling in acetylcholine neurons is sufficient to regulate locomotion and increase acetylcholine release.

(A) Transgenic lines expressing an activated form of *lin-45* (T626E/T629D) in acetylcholine neurons (*yakIs34[Punc-17::lin-45(gf)]*) have exaggerated body bends and coiling behavior similar to the activated Gq mutant *egl-30*(*tg26*) and to animals expressing activated Rho in acetylcholine neurons (*nzIs29[Punc-17::rho-1(gf)]*). The wild type and *egl-30(tg26)* photos are the same as shown in Figure 1A, while the *rho-1(gf)* photos is the same as the one in Figure 3A.

(B) Expression of activated Rho (*nzIs29[Punc-17::rho-1(gf)]*) or Raf (*yakIs34[Punc-17::lin-45(gf)]*) in acetylcholine neurons impairs coordinated locomotion similarly to activated Gq (*egl-30(tg26)*). $N \geq 19$ * $P < 0.05$, *** $P < 0.001$, Kruskal-Wallis test with Dunn's *post hoc* test.

(C) Animals expressing activated Gq, Rho, or Raf are hypersensitive to the

acetylcholinesterase inhibitor aldicarb. Activated Gq (*egl-30*(*tg26*)), activated Rho

expressed in acetylcholine neurons (*nzIs29[Punc-17::rho-1(gf)]*), and activated Raf

expressed in acetylcholine neurons *yakIs34[Punc-17::lin-45(gf)]*) become paralyzed

significantly faster than wild type animals when exposed to 1 mM aldicarb. All strains

are significantly different from wild type at t = 60, 75, 90, and 105 minutes. $N \geq 61$ *** *P*

< 0.001, two-way ANOVA with Bonferroni's *post hoc* test.

(D) *ksr-1* is not necessary for the aldicarb hypersensitivity of activated Gq. Paralysis of

ksr-1(*ok786*) animals on 1 mM aldicarb is not significantly different from wild type. The

ksr-1 deletion *ok786* does not suppress the aldicarb hypersensitivity of activated Gq

(*egl-30*(*tg26*)). $N \geq 53$.

Figure 6. Model for Gq-Rho-ERK signaling. Gq directly activates Trio and Rho (solid

arrow) (Lutz et al. 2007). The core ERK cascade acts either downstream of Rho or in

parallel (dashed arrows) to modulate locomotion. ERK activation occurs independently

of the extracellular growth factor LIN-3, its receptor LET-23, or Ras/LET-60. It is

possible that Raf/LIN-45 is activated by a Ras family member other than the canonical

worm Ras/LET-60.

Figure S1. Additional mutations in *mek-2* and *mpk-1* suppress activated Gq.

(A) The *mek-2(ku114)* mutation suppresses the exaggerated waveform of *egl-30*(*tg26*)

worms. $N \geq 14$ *** *P* < 0.001, one-way ANOVA with Bonferroni's *post hoc* test.

(B) The *mpk-1(oz140)* mutation suppresses the exaggerated waveform of *egl-30*(*tg26*)

worms. *N* ≥ 12 *** *P* < 0.001, one-way ANOVA with Bonferroni's *post hoc* test.

Figure S2. Gq-dependent locomotion does not depend on the RasGEFs *sos-1* and

rgef-1, or the Ras family members *ras-1*, *ras-2*, and *rap-1*.

(A) Deletion mutations *ras-1(gk243)* and *ras-2(ok682),* or the nonsense mutation *rap-*

1(pk2082) do not suppress the waveform of the activated Gq mutant *egl-30(tg26)*. *N* ≥

18, n.s., not significant, one-way ANOVA with Bonferroni's *post hoc* test.

(B) Loss of the RasGEF *sos-1* does not suppress activated Gq waveform. Temperature

sensitive *egl-30(tg26); sos-1(cs41)* mutant animals were incubated for 24 hours at the

non-permissive temperature of 25° and assayed for their waveform. *N* ≥ 15, n.s., not

significant, one-way ANOVA with Tukey's *post hoc* test.

(C) The RasGEF deletion mutation *rgef-1(ok675)* does not suppress the exaggerated

waveform of activated Gq. *N* = 16, n.s., not significant, one-way ANOVA with Tukey's

post hoc test.

Figure S3. Expression of activated LIN-45/Raf in head acetylcholine neurons is

sufficient to cause locomotion with exaggerated body bends.

The *yakIs34[Punc-17::lin-45(gf)]* integrated array expressing activated *lin-*

45(T626E/T629D) in all acetylcholine neurons and the *yakEx154[Punc-17H::lin-45(gf)]*

extrachromosomal array expressed in head acetylcholine neurons both cause

exaggerated body bends and coiled locomotion. *N* ≥ 12 *** *P* < 0.001, * *P* < 0.05,

Kruskal-Wallis test with Dunn's *post hoc* test.

Table S1. List of strains

Table S1. List of strains

N2   Bristol wild strain

XZ1151   *egl-30(tg26) I*

EG6699   *ttTi5605 II; unc-119(ed3) III*

EG6207   *unc-119(ed3) III*

RB915   *ksr-1(ok786) X*

EG4782   *nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II*

MT4866   *let-60(n2021) IV*

PS436   *let-60(sy93) IV*

PS427   *lin-45(sy96) IV*

MT8666   *mek-2(n1989) I*

SD378   *mpk-1(ga117) / dpy-17(e164) unc-79(e1068) III*

CB1417   *lin-3(e1417) IV*

PS5131   *let-23(sy12) / mIn1[dpy-10(e128) mIs14] II*

UP604   *sos-1(cs41) V*

VC450   *ras-1(gk243) II*

RB852   *ras-2(ok682) III*

TZ181   *rap-1(pk2082) IV*

RB848   *rgef-1(ok675) V*

MH538   *mek-2(ku114) I; let-60(n1046) IV*

MT8186   *mpk-1(oz140)/unc-79(e1068) dpy-17(e164) III*

The following strains were produced during this study:

EG314   *ksr-1(ox314) X*

XZ10   *ksr-1(yak10) X*

EG4198   *egl-30(tg26) I ; ksr-1(ox314) X*

XZ1340   *egl-30(tg26) I ; ksr-1(yak10) X*

XZ1511   *egl-30(tg26) I; ksr-1(ok786) X*

XZ1465　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ksr-1(ox314) X*

XZ1547　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ksr-1(yak10) X*

XZ2042　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ksr-1(ok786) X*

XZ1615　*egl-30(tg26) I; let-60(n2021) / nT1[qIs51(Pmyo-2::GFP; Ppes-10::GFP; F22B7.9::GFP)] IV*

XZ1626　*egl-30(tg26) I; lin-3(e1417) IV*

XZ1630　*egl-30(tg26) I; let-23(sy12) / mIn1[dpy-10(e128) mIs14] II*

XZ1677　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; let-60(n2021) / nT1[qIs51(Pmyo-2::GFP;*

Ppes-10::GFP; F22B7.9::GFP)] IV

XZ1548　*egl-30(tg26) I; lin-45(sy96) IV*

XZ1556　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; lin-45(sy96) IV*

XZ1850　*egl-30(tg26) mek-2(n1989) / hT2[bli-4(e937) let(q782) qIs48] I*

XZ1851　*mek-2(n1989) / hT2[bli-4(e937) let(q782) qIs48] I; nzIs29[Punc-17::rho-1(G14V), unc-*

122::gfp] II

XZ1700　*mpk-1(ga117) / qC1[dpy-19(e1259) glp-1(q339) qIs26(Plag-2::GFP, rol-6(su1006))] III*

XZ1668　*egl-30(tg26) I; mpk-1(ga117) / qC1[dpy-19(e1259) glp-1(q339) qIs26(Plag-2::GFP, rol-*

6(su1006))] III

XZ2043　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; mpk-1(ga117) / qC1[dpy-19(e1259) glp-*

1(q339) qIs26(Plag-2::GFP, rol-6(su1006))] III

XZ2046　*egl-30(tg26) I; sos-1(cs41) V*

XZ1855　*egl-30(tg26) I; ras-1(gk243) II*

XZ1856　*egl-30(tg26) I; ras-2(ok682) III*

XZ2045　*egl-30(tg26) I; rap-1(pk2082) IV*

XZ2101　*egl-30(tg26) I; rgef-1(ok675) V*

XZ1921　*egl-30(tg26) sur-6(sv30) / hT2[bli-4(e937) let(q782) qIs48] I*

XZ1857　*egl-30(tg26) I ; sur-7(ku119) X*

XZ1854　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; ras-2(ok682) III*

XZ1852　*nzIs29[Punc-17::rho-1(G14V), unc-122::gfp] II; rap-1(pk2082) IV*

XZ1880　*yakSi26[Punc-17::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1881   *yakSi27[Punc-17H::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1882   *yakSi28[Punc-17β::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1883   *yakSi29[Punc-47::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II*

XZ1884   *egl-30(tg26) I; yakSi26[Punc-17::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-1(ok786)*

X

XZ1885   *egl-30(tg26) I; yakSi27[Punc-17H::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-*

1(ok786) X

XZ1946   *egl-30(tg26) I; yakSi28[Punc-17β::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-*

1(ok786) X

XZ1947   *egl-30(tg26) I; yakSi29[Punc-47::ksr-1 cDNA::tbb-2 3'UTR::OPERON::GFP] II; ksr-1(ok786)*

X

XZ2015   *unc-119(ed3) III; yakEx122[Punc-17::lin-45* cDNA::tbb-2 3'UTR::OPERON::GFP, Pmyo-*

3::mCherry]

XZ2077   *egl-30(tg26) I; let-60(sy93) IV*

XZ2106   *egl-30(tg26) I; let-60(sy93) / nT1[qIs51(Pmyo-2::GFP; Ppes-10::GFP; F22B7.9::GFP)] IV*

XZ2050   *unc-119(ed3) III; yakIs34[Punc-17::lin-45* cDNA::tbb-2 3'UTR::OPERON::GFP, Pmyo-*

3::mCherry] X

XZ2130   *egl-30(tg26) I; mpk-1(oz140) / oxTi619[Peft-3::TdTomato::H2B cb-unc-119(+)] III*

XZ2131   *egl-30(tg26) mek-2(ku114) I / hT2[bli-4(e937) let(q782) qIs48] I;III*

XZ2119   *ttTi5605 II; unc-119(ed3) III; yakEx154[Punc-17H::lin-45*::tbb-2 3'UTR::OPERON::GFP,*

Pmyo-2::mCherry, Pmyo-3::mCherry]

Table S2. List of plasmids

Gateway destination vectors

pCFJ150   Gateway destination vector for insertion at chr II Mos site ttTi5605

Gateway entry clones

pEGB05   *Prab-3* [4-1]

pGH1   *Punc-17* [4-1]

pADA180   *Punc-17H* [4-1] (head acetylcholine neurons)

pMA23   *Punc-17β* [4-1] (body acetylcholine neurons)

pMH522   *Punc-47* [4-1]

pBC11   *ksr-1* cDNA [1-2]

pBC26   *lin-45* cDNA [1-2]

pBC35   *lin-45** (T626E/T629D) cDNA [1-2]

pCFJ326   *tbb-2* 3'UTR::OPERON::*GFP* [2-3]

Gateway expression constructs

pBC13   *Prab-3::ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

pBC31   *Punc-17::ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

pBC32   *Punc-17H*::*ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

pBC33   *Punc-17β*::*ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

pBC34   *Punc-47*::*ksr-1* cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

pBC37   *Punc-17::lin-45** cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

pBC44   *Punc-17H::lin-45** cDNA::*tbb-2* 3'UTR::OPERON::GFP_pCFJ150

Table S3. List of primers

oBC011   Forward *ksr-1* cDNA for Gateway [1-2] ENTR vector

656   GGGGACAAGTTTGTACAAAAAAGCAGGCTCAatgatgcaaacccaagttgc

657   oBC012   Reverse *ksr-1* cDNA for Gateway [1-2] ENTR vector

658   GGGGACCACTTTGTACAAGAAAGCTGGGTGaaatgtcgactcgtaacttttcatc

659   oBC085   Forward *lin-45* cDNA for Gateway [1-2] ENTR vector

660   GGGGACAAGTTTGTACAAAAAAGCAGGCTcaATGAGTCGGATTAATTTCAAAAAG

661   oBC086   Reverse *lin-45* cDNA for Gateway [1-2] ENTR vector

662   GGGGACCACTTTGTACAAGAAAGCTGGGTgCTAAATGAGACCATAGACATTGTAGTATG

664 oBC094   Reverse *lin-45* T626E mutagenesis

665   gttcactgtccatttcgttttgacctctgccaagccgaaatctccaatttt

666 oBC095   Forward *lin-45* T626E mutagenesis

667   aaaattggagatttcggcttggcagaggtcaaaacgaaatggacagtgaac

668 oBC096   Reverse *lin-45* T629D mutagenesis

669   cctccgttcactgtccatttatctttgacctctgccaagccga

670 oBC097   Forward *lin-45* T629D mutagenesis

671   tcggcttggcagaggtcaaagataaatggacagtgaacggagg

Literature Cited

Afgan E., Baker D., Beek M. van den, Blankenberg D., Bouvier D., *et al.*, 2016 The Galaxy

platform for accessible, reproducible and collaborative biomedical analyses: 2016

update. Nucleic Acids Res. 44: W3–W10.

Ailion M., Hannemann M., Dalton S., Pappas A., Watanabe S., *et al.*, 2014 Two Rab2

interactors regulate dense-core vesicle maturation. Neuron 82: 167–180.

Bastiani C. A., Gharib S., Simon M. I., Sternberg P. W., 2003 Caenorhabditis elegans Gαq

Regulates Egg-Laying Behavior via a PLCβ-Independent and Serotonin-Dependent

Signaling Pathway and Likely Functions Both in the Nervous System and in Muscle.

Genetics 165: 1805–1822.

Beitel G. J., Clark S. G., Horvitz H. R., 1990 Caenorhabditis elegans ras gene let-60 acts as a

switch in the pathway of vulval induction. Nature 348: 503–509.

Boggio E. M., Putignano E., Sassoè-Pognetto M., Pizzorusso T., Giustetto M., 2007 Visual

Stimulation Activates ERK in Synaptic and Somatic Compartments of Rat Cortical

Neurons with Parallel Kinetics. PLoS ONE 2: e604.

Boulton T. G., Nye S. H., Robbins D. J., Ip N. Y., Radzlejewska E., *et al.*, 1991 ERKs: A family of

protein-serine/threonine kinases that are activated and tyrosine phosphorylated in

response to insulin and NGF. Cell 65: 663–675.

Brundage L., Avery L., Katz A., Kim U. J., Mendel J. E., *et al.*, 1996 Mutations in a C. elegans

Gqalpha gene disrupt movement, egg laying, and viability. Neuron 16: 999–1009.

Chan J. P., Hu Z., Sieburth D., 2012 Recruitment of sphingosine kinase to presynaptic

terminals by a conserved muscarinic signaling pathway promotes neurotransmitter

release. Genes Dev. 26: 1070–1085.

Chen L., Fu Y., Ren M., Xiao B., Rubin C. S., 2011 A RasGRP, C. elegans RGEF-1b, Couples

External Stimuli to Behavior by Activating LET-60 (Ras) in Sensory Neurons. Neuron

70: 51–65.

Chong H., Lee J., Guan K. L., 2001 Positive and negative regulation of Raf kinase activity and

function by phosphorylation. EMBO J. 20: 3716–3727.

Cingolani P., Platts A., Wang L. L., Coon M., Nguyen T., *et al.*, 2012 A program for annotating

and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the

genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly 6: 80–92.

Davis M. W., Hammarlund M., Harrach T., Hullett P., Olsen S., *et al.*, 2005 Rapid single

nucleotide polymorphism mapping in C. elegans. BMC Genomics 6: 118.

Dent J. A., Han M., 1998 Post-embryonic expression pattern of C. elegans let-60 ras reporter

constructs. Mech. Dev. 72: 179–182.

DePristo M. A., Banks E., Poplin R., Garimella K. V., Maguire J. R., *et al.*, 2011 A framework

for variation discovery and genotyping using next-generation DNA sequencing data.

Nat. Genet. 43: 491–498.

Doi M., Iwasaki K., 2002 Regulation of retrograde signaling at neuromuscular junctions by

the novel C2 domain protein AEX-1. Neuron 33: 249–259.

Drosten M., Sum E. Y. M., Lechuga C. G., Simón-Carrasco L., Jacob H. K. C., *et al.*, 2014 Loss of

p53 induces cell proliferation via Ras-independent activation of the Raf/Mek/Erk

signaling pathway. Proc. Natl. Acad. Sci. U. S. A. 111: 15155–15160.

Frøkjær-Jensen C., Davis M. W., Ailion M., Jorgensen E. M., 2012 Improved Mos1-mediated

transgenesis in C. elegans. Nat. Methods 9: 117–118.

Guo F. F., Kumahara E., Saffen D., 2001 A CalDAG-GEFI/Rap1/B-Raf cassette couples M(1)

muscarinic acetylcholine receptors to the activation of ERK1/2. J. Biol. Chem. 276:

25568–25581.

Hamakawa M., Uozumi T., Ueda N., Iino Y., Hirotsu T., 2015 A role for Ras in inhibiting

circular foraging behavior as revealed by a new method for time and cell-specific

RNAi. BMC Biol. 13: 6.

Han M., Sternberg P. W., 1990 let-60, a gene that specifies cell fates during C. elegans vulval

induction, encodes a ras protein. Cell 63: 921–931.

Han M., Aroian R. V., Sternberg P. W., 1990 The let-60 locus controls the switch between

vulval and nonvulval cell fates in Caenorhabditis elegans. Genetics 126: 899–913.

Han M., Sternberg P. W., 1991 Analysis of dominant-negative mutations of the

Caenorhabditis elegans let-60 ras gene. Genes Dev. 5: 2188–2198.

Han M., Golden A., Han Y., Sternberg P. W., 1993 C. elegans lin-45 raf gene participates in

let-60 ras-stimulated vulval differentiation. Nature 363: 133–140.

Hiley E., McMullan R., Nurrish S. J., 2006 The Galpha12-RGS RhoGEF-RhoA signalling

pathway regulates neurotransmitter release in C. elegans. EMBO J. 25: 5884–5895.

Hirotsu T., Saeki S., Yamamoto M., Iino Y., 2000 The Ras-MAPK pathway is important for

olfaction in Caenorhabditis elegans. Nature 404: 289–293.

Hirotsu T., Iino Y., 2005 Neural circuit-dependent odor adaptation in C. elegans is regulated

by the Ras-MAPK pathway. Genes Cells 10: 517–530.

Honda Z., Takano T., Gotoh Y., Nishida E., Ito K., *et al.*, 1994 Transfected platelet-activating

factor receptor activates mitogen-activated protein (MAP) kinase and MAP kinase

kinase in Chinese hamster ovary cells. J. Biol. Chem. 269: 2307–2315.

Hsu V., Zobel C. L., Lambie E. J., Schedl T., Kornfeld K., 2002 Caenorhabditis elegans lin-45

raf is essential for larval viability, fertility and the induction of vulval cell fates.

Genetics 160: 481–492.

Hu Z., Vashlishan-Murray A. B., Kaplan J. M., 2015 NLP-12 engages different UNC-13

proteins to potentiate tonic and evoked release. J. Neurosci. 35: 1038–1042.

Hunt-Newbury R., Viveiros R., Johnsen R., Mah A., Anastas D., *et al.*, 2007 High-throughput

in vivo analysis of gene expression in Caenorhabditis elegans. PLoS Biol. 5: e237.

Hyde R., Corkins M. E., Somers G. A., Hart A. C., 2011 PKC-1 acts with the ERK MAPK

signaling pathway to regulate C. elegans mechanosensory response. Genes Brain

Behav. 10: 286–298.

Impey S., Obrietan K., Storm D. R., 1999 Making New Connections. Neuron 23: 11–14.

Jovanovic J. N., Czernik A. J., Fienberg A. A., Greengard P., Sihra T. S., 2000 Synapsins as

mediators of BDNF-enhanced neurotransmitter release. Nat. Neurosci. 3: 323.

Karnoub A. E., Weinberg R. A., 2008 Ras oncogenes: split personalities. Nat. Rev. Mol. Cell

Biol. 9: 517–531.

Kornfeld K., Guan K. L., Horvitz H. R., 1995a The Caenorhabditis elegans gene mek-2 is

required for vulval induction and encodes a protein similar to the protein kinase

MEK. Genes Dev. 9: 756–768.

Kornfeld K., Hom D. B., Horvitz H. R., 1995b The ksr-1 gene encodes a novel protein kinase

involved in Ras-mediated signaling in C. elegans. Cell 83: 903–913.

Kushner S. A., Elgersma Y., Murphy G. G., Jaarsma D., Woerden G. M. van, *et al.*, 2005

Modulation of presynaptic plasticity and learning by the H-ras/extracellular signal-

regulated kinase/synapsin I signaling pathway. J. Neurosci. 25: 9721–9734.

Lackner M. R., Kim S. K., 1998 Genetic analysis of the Caenorhabditis elegans MAP kinase

gene mpk-1. Genetics 150: 103–117.

Lackner M. R., Nurrish S. J., Kaplan J. M., 1999 Facilitation of Synaptic Transmission by EGL-

30 Gqα and EGL-8 PLCβ: DAG Binding to UNC-13 Is Required to Stimulate

Acetylcholine Release. Neuron 24: 335–346.

Lutz S., Freichel-Blomquist A., Yang Y., Rümenapp U., Jakobs K. H., *et al.*, 2005 The guanine

nucleotide exchange factor p63RhoGEF, a specific link between Gq/11-coupled

receptor signaling and RhoA. J. Biol. Chem. 280: 11134–11139.

Lutz S., Shankaranarayanan A., Coco C., Ridilla M., Nance M. R., *et al.*, 2007 Structure of

Galphaq-p63RhoGEF-RhoA complex reveals a pathway for the activation of RhoA by

GPCRs. Science 318: 1923–1927.

Mahoney T. R., Luo S., Nonet M. L., 2006 Analysis of synaptic transmission in

Caenorhabditis elegans using an aldicarb-sensitivity assay. Nat. Protoc. 1: 1772–

1777.

Mao L.-M., Wang J. Q., 2016a Regulation of Group I Metabotropic Glutamate Receptors by

MAPK/ERK in Neurons. J. Nat. Sci. 2: e268.

Mao L.-M., Wang J. Q., 2016b Synaptically Localized Mitogen-Activated Protein Kinases:

Local Substrates and Regulation. Mol. Neurobiol. 53: 6309–6315.

Marais R., Light Y., Paterson H. F., Marshall C. J., 1995 Ras recruits Raf-1 to the plasma

membrane for activation by tyrosine phosphorylation. EMBO J. 14: 3136–3145.

McMullan R., Hiley E., Morrison P., Nurrish S. J., 2006 Rho is a presynaptic activator of

neurotransmitter release at pre-existing synapses in C. elegans. Genes Dev. 20: 65–

76.

McMullan R., Anderson A., Nurrish S., 2012 Behavioral and Immune Responses to Infection

Require Gαq- RhoA Signaling in C. elegans. PLOS Pathog. 8: e1002530.

Mello C. C., Kramer J. M., Stinchcomb D., Ambros V., 1991 Efficient gene transfer in

C.elegans: extrachromosomal maintenance and integration of transforming

sequences. EMBO J. 10: 3959–3970.

Miller K. G., Emerson M. D., Rand J. B., 1999 Goalpha and diacylglycerol kinase negatively

regulate the Gqalpha pathway in C. elegans. Neuron 24: 323–333.

Nguyen A., Burack W. R., Stock J. L., Kortum R., Chaika O. V., *et al.*, 2002 Kinase Suppressor

of Ras (KSR) Is a Scaffold Which Facilitates Mitogen-Activated Protein Kinase

Activation In Vivo. Mol. Cell. Biol. 22: 3035–3045.

Nicholas H. R., Hodgkin J., 2004 The ERK MAP kinase cascade mediates tail swelling and a

protective response to rectal infection in C. elegans. Curr. Biol. 14: 1256–1261.

Ortiz J., Harris H. W., Guitart X., Terwilliger R. Z., Haycock J. W., *et al.*, 1995 Extracellular

signal-regulated protein kinases (ERKs) and ERK kinase (MEK) in brain: regional

distribution and regulation by chronic morphine. J. Neurosci. 15: 1285–1297.

Reiner D., Lundquist E. A., 2016 Small GTPases. WormBook Online Rev. C Elegans Biol.

doi/10.1895/wormbook.1.67.2

Rhee S. G., 2001 Regulation of phosphoinositide-specific phospholipase C. Annu. Rev.

Biochem. 70: 281–312.

Robbins D. J., Cheng M., Zhen E., Vanderbilt C. A., Feig L. A., *et al.*, 1992 Evidence for a Ras-

dependent extracellular signal-regulated protein kinase (ERK) cascade. Proc. Natl.

Acad. Sci. U. S. A. 89: 6924–6928.

Schmitz S. K., King C., Kortleven C., Huson V., Kroon T., *et al.*, 2016 Presynaptic inhibition

upon CB1 or mGlu2/3 receptor activation requires ERK/MAPK phosphorylation of

Munc18-1. EMBO J. 35: 1236–1250.

Schrader L. A., Birnbaum S. G., Nadin B. M., Ren Y., Bui D., *et al.*, 2006 ERK/MAPK regulates

the Kv4.2 potassium channel by direct phosphorylation of the pore-forming subunit.

Am. J. Physiol. Cell Physiol. 290: C852-861.

Sieburth D. S., Sundaram M., Howard R. M., Han M., 1999 A PP2A regulatory subunit

positively regulates Ras-mediated signaling during Caenorhabditis elegans vulval

induction. Genes Dev. 13: 2562–2569.

Stamboulian S., Choi J.-S., Ahn H.-S., Chang Y.-W., Tyrrell L., *et al.*, 2010 ERK1/2 mitogen-

activated protein kinase phosphorylates sodium channel Na(v)1.7 and alters its

gating properties. J. Neurosci. 30: 1637–1647.

Sternberg P. W., 2005 Vulval development. WormBook Online Rev. C Elegans Biol.

doi/10.1895/wormbook.1.6.1

Stokoe D., Macdonald S. G., Cadwallader K., Symons M., Hancock J. F., 1994 Activation of Raf

as a result of recruitment to the plasma membrane. Science 264: 1463–1467.

Sun Y., Liu W.-Z., Liu T., Feng X., Yang N., *et al.*, 2015 Signaling pathway of MAPK/ERK in cell

proliferation, differentiation, migration, senescence and apoptosis. J. Recept. Signal

Transduct. Res. 35: 600–604.

Sundaram M., Han M., 1995 The C. elegans ksr-1 gene encodes a novel Raf-related kinase

involved in Ras-mediated signal transduction. Cell 83: 889–901.

Sundaram M. V., 2013 Canonical RTK-Ras-ERK signaling and related alternative pathways.

WormBook Online Rev. C Elegans Biol. doi/10.1895/wormbook.1.80.2

Sweatt J. D., 2004 Mitogen-activated protein kinases in synaptic plasticity and memory.

Curr. Opin. Neurobiol. 14: 311–317.

Thomas G. M., Huganir R. L., 2004 MAPK cascade signalling and synaptic plasticity. Nat. Rev.

Neurosci. 5: 173–183.

Tomida T., Oda S., Takekawa M., Iino Y., Saito H., 2012 The Temporal Pattern of Stimulation

Determines the Extent and Duration of MAPK Activation in a Caenorhabditis elegans

Sensory Neuron. Sci Signal 5: ra76.

Tomioka M., Adachi T., Suzuki H., Kunitomo H., Schafer W. R., *et al.*, 2006 The insulin/PI 3-

kinase pathway regulates salt chemotaxis learning in Caenorhabditis elegans.

Neuron 51: 613–625.

Topalidou I., Chen P.-A., Cooper K., Watanabe S., Jorgensen E. M., *et al.*, 2017a The NCA-1

and NCA-2 Ion Channels Function Downstream of Gq and Rho To Regulate

Locomotion in Caenorhabditis elegans. Genetics 206: 265-282.

Topalidou I., Cooper K., Pereira L., Ailion M., 2017b Dopamine negatively modulates the

NCA ion channels in C. elegans. PLoS Genet. 13: e1007032.

Ueda Y., Hirai S. i, Osada S. i, Suzuki A., Mizuno K., *et al.*, 1996 Protein kinase C activates the

MEK-ERK pathway in a manner independent of Ras and dependent on Raf. J. Biol.

Chem. 271: 23512–23519.

Uozumi T., Hirotsu T., Yoshida K., Yamada R., Suzuki A., *et al.*, 2012 Temporally-regulated

quick activation and inactivation of Ras is important for olfactory behaviour. Sci.

Rep. 2: 500.

van Biesen T., Hawes B. E., Raymond J. R., Luttrell L. M., Koch W. J., *et al.*, 1996 G(o)-protein

alpha-subunits activate mitogen-activated protein kinase via a novel protein kinase

C-dependent mechanism. J. Biol. Chem. 271: 1266–1269.

Vara H., Onofri F., Benfenati F., Sassoè-Pognetto M., Giustetto M., 2009 ERK activation in

axonal varicosities modulates presynaptic plasticity in the CA3 region of the

hippocampus through synapsin I. Proc. Natl. Acad. Sci. U. S. A. 106: 9872–9877.

Wilkie T. M., Scherle P. A., Strathmann M. P., Slepak V. Z., Simon M. I., 1991 Characterization

of G-protein alpha subunits in the Gq class: expression in murine tissues and in

stromal and hematopoietic cell lines. Proc. Natl. Acad. Sci. U. S. A. 88: 10049–10053.

Williams S. L., Lutz S., Charlie N. K., Vettel C., Ailion M., *et al.*, 2007 Trio's Rho-specific GEF

domain is the missing Gαq effector in C. elegans. Genes Dev. 21: 2731–2746.

Wu Y., Han M., Guan K. L., 1995 MEK-2, a Caenorhabditis elegans MAP kinase kinase,

functions in Ras-mediated vulval induction and other developmental events. Genes

Dev. 9: 742–755.

Yang F., Sun W., Yang Y., Wang Y., Li C.-L., *et al.*, 2015 SDF1–CXCR4 signaling contributes to

persistent pain and hypersensitivity via regulating excitability of primary

nociceptive neurons: involvement of ERK-dependent Nav1.8 up-regulation. J.

Neuroinflammation 12: 219–233.

Yoder J. H., Chong H., Guan K.-L., Han M., 2004 Modulation of KSR activity in Caenorhabditis

elegans by Zn ions, PAR-1 kinase and PP2A phosphatase. EMBO J. 23: 111–119.

Yoon S., Seger R., 2006 The extracellular signal-regulated kinase: multiple substrates

regulate diverse cellular functions. Growth Factors 24: 21–44.

You Y., Kim J., Cobb M., Avery L., 2006 Starvation activates MAP kinase through the

muscarinic acetylcholine pathway in Caenorhabditis elegans pharynx. Cell Metab. 3:

237–245.

Zhang H., Koo C. Y., Stebbing J., Giamas G., 2013 The dual function of KSR1: a pseudokinase

and beyond. Biochem. Soc. Trans. 41: 1078–1082.

Figure 1

A

wild type | 300 μm | *egl-30(tg26)* | *egl-30(tg26); ksr-1(ok786)*

B

ksr-1

yak10 W254>stop | *ox314* K463>stop | 1000 bp

ok786 deletion

C

wild type | 2x amplitude | period

ksr-1(ok786)

egl-30(tg26)

egl-30(tg26); ksr-1(ok786)

D

period/amplitude

wild type, ksr-1(ox314), ksr-1(yak10), ksr-1(ok786), egl-30(tg26), ksr-1(ox314), ksr-1(yak10), ksr-1(ok786)

*** *** ***

egl-30(tg26)

Figure 2

Figure 3

A

rho-1(gf)   150 μm

rho-1(gf);
ksr-1(ok786)

rho-1(gf);
let-60(n2021)

rho-1(gf);
lin-45(sy96)

rho-1(gf);
mek-2(n1989)

rho-1(gf);
mpk-1(ga117)

B

distance from center (mm)

*** *** **

wild type, ksr-1(ox314), ksr-1(yak10), ksr-1(ok786), rho-1(gf), ksr-1(ox314), ksr-1(yak10), ksr-1(ok786)

rho-1(gf)

C

distance from center (mm)

n.s. n.s. *** ***

wild type, let-60(n2021), lin-45(sy96), mek-2(n1989), mpk-1(ga117), rho-1(gf), let-60(n2021), lin-45(sy96), mek-2(n1989), mpk-1(ga117)

rho-1(gf)

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

Figure S3

wild type   300 μm   *Punc-17::lin-45(gf)*   *Punc-17H::lin-45(gf)*