Molecular map of GNAO1-related disease phenotypes and reactions to therapy

Ivana Mihalek[a,1], Jeff L. Waugh[b,c], Meredith Park[d], Saima Kayani[b], Annapurna Poduri[c, d, e], and Olaf Bodamer[f,g,1]

[a]Department of Molecular Medicine and Biotechnology, Faculty of Medicine, University of Rijeka, Rijeka, Croatia
[b]Department of Pediatrics, University of Texas Southwestern, Dallas, TX
[c]Department of Neurology, Boston Children's Hospital, Harvard Medical School, Boston, MA
[d]Epilepsy Genetics Program, Department of Neurology, Boston Children's Hospital, Boston, MA
[e]F.M. Kirby Neurobiology Center, Boston Children's Hospital, Boston, MA
[f]Division of Genetics and Genomics, Boston Children's Hospital, Harvard Medical School, Boston, MA
[g]Broad Institute of MIT and Harvard University, Cambridge, MA
[1]corresponding author

September 21, 2020

The GNAO1 gene codes for the most commonly expressed $G\alpha$ protein in the central nervous system. Pathogenic GNAO1 variants result in early-onset neurological phenotypes, sometimes with distinct epilepsy or movement disorder manifestations, and sometimes with both manifestations in the same patient.

The existing extensive knowledge about G-protein coupled receptor (GPCR) signaling provides the input needed to quantitatively evaluate how mutations modify the GPCR signal. This in turn allows rational interpretation of distinct phenotypes arising from mutations in GNAO1. In this work we outline a model that enables understanding of clinical phenotypes at a molecular level, and anticipating symptoms and treatment response for future patients.

We show that the epileptic phenotype arises necessarily in mutations affecting the catalytic pocket of GNAO1, which results in the improper withdrawal of the signal. Mutations close to the interface with GNAO1's downstream effector cut the signal propagation in that

direction, and manifest as a movement disorder phenotype without epilepsy. Quantifying the reported reaction to therapy highlights the tendency of the latter group to be unresponsive to the therapies in use to date. We argue, however, that the largest number of mutations can impact several aspects of GNAO1 function at once, resulting in the continuum of phenotypes observed in patients. In addition to the practical implications for decision making process in choosing therapy for GNAO1 patients, analyzing this system provides a precision medicine paradigm in which individual genomic information and understanding the implications of variants at the protein level can inform and guide the treatment process.

Introduction

From the reports [1, 2, 3, 4, 5], and the early compilations [6, 7, 8, 9, 10], of disease causing variants in GNAO1 gene, a picture emerges of a source of complex phenotypes, with equally complex and sometimes contradictory responses to treatment. GNAO1 (guanine nucleotide binding protein, subunit α, type "o") is the most common type of Gα in the central nervous system (CNS). It functions at the core of multiple neurotransmitter pathways, and is expressed in multiple types of neurons [11]. In particular, GNAO1 expresses profusely in striatum, [12, 13], Table S1, the region of the brain responsible for the smooth transition between movement sequences [14]. Unsurprisingly then, GNAO1 mutations cause a spectrum of neurological phenotypes, roughly falling into two categories: movement disorders (MD) or epilepsy-related (E) phenotypes. While most patients manifest symptoms from both categories, some patients have exclusively MD or E manifestations. The most frequent MD phenotype is chorea, though the spectrum can include or be limited to myoclonus or dystonia. Epilepsy-related disorders are typically early-onset, often with status epilepticus and/or seizures, refractory to a multi anti-epileptic drug regimen. The epilepsy-dominant patients universally have global developmental delay, often accompanied by severe hypotonia, and typically manifest stereotypies and difficulty with motor coordination. The fact that the two patterns may completely segregate suggests that distinct classes of molecular abnormality underlie GNAO1-related symptoms. At the same time, the movement and epilepsy phenotypes do overlap in the majority of cases, suggesting that a single mutation can impact multiple aspects of GNAO1 function, eluding the straightforward gain- and loss-of-function classification.

The reports on GNAO1 therapy and its effectiveness are not systematic. However, if we consider drugs in terms of their main targets, roughly two classes of disease variants emerge: one responsive to modulation of GABA and dopamine receptors, and the other refractory to all reported therapy. In large part, the first class overlaps with the variants close to the catalytic pocket of GNAO1 [10], while the second overlaps with those close to the interface with its effector. The overlap is incomplete, and the therapy response much more complex than this schematic classification. The challenge lies precisely in characterizing the molecular lesions in a way that reconciles the seemingly contradictory treatment requirements and predicts medication efficacy, thus reducing side effects and time to benefit.

In this work we show that even though the concrete aspects of phenotype in this system may depend on the individual genome background and development history, the genotype-phenotype correlation can be rationalized by considering the core loop of the G protein signaling pathway, and the way in which disease-related mutations modify the signal.

Molecular players in GNAO1 signaling pathway

The model we describe below centers on the GPCR signaling as a membrane bound GTP-ase signaling pathway [15], Fig. S4 and Movie S1, with its three obligate components: a GTPase, a guanine nucleotide exchange factor (GEF), and a GTPase-activating protein (GAP). It is a particular feature of GPCR signaling, in contrast to the signaling through small GTPases, that the GTPase, Gα, forms a complex with G$\beta\gamma$ [16]. This complex dissociates upon the interaction with activated GPCR, creating

a branching, or bipartite, signal [17], with Gα and G$\beta\gamma$ independently influencing their downstream effectors.

GTPase: GNAO1

GNAO1 is severalfold more abundant than any other Gα protein in the CNS [11]. Neurons and neuroendocrine cells are the primary site of expression of the gene [18, 13]. Known G$\beta\gamma$ partners of GNAO1 are encoded by GNB1 (as Gβ) and GNG2 (as Gγ) [19], though the existence of other partners is likely. As is the case with GNAO1, GNB1 and GNG2 are preferentially expressed in brain [20]. Of the five genes coding for Gβ [21], GNB1 is the only one that has been related to autosomal dominant neurodevelopmental disease phenotypes, typically global developmental delays and hypotonia, sometimes accompanied by epilepsy and/or dystonia [22, 23]. Known disease-related GNB1 mutations are located almost exclusively at the interface with Gα [23], Movie S1.

GEF: G-protein coupled receptor (GPCR)

GNAO1 proteins are known to interact with GPCRs from the GABA, adrenergic, dopamine, and muscarinic group, and more [18], underscoring the diversity and extent of systemic damage incurred with GNAO1 mutations. When activated by their respective agonist, GPCRs act as GEFs for GNAO1.

GAP: RGS protein

Multiple RGS proteins accelerate the GTPase activity of GNAO1, including RGS1, RGS16, and GAIP, and almost certainly others. This interaction is possibly competing with the effector binding [11].

Effectors of Gα and G$\beta\gamma$

The prototypical downstream effectors of GNAO1 signaling are adenylate cyclase for Gα, and ion channels, phospholipase C-β, or phosphoinositide-3-kinase for G$\beta\gamma$ [18]. GNAO1 functions as inhibitor of adenylate cyclase (ADCY) *in vitro* [24]. Of all ADCY genes, ADCY5 most convincingly coexpresses with GNAO1 in striatum (Table S1). This interaction is further supported by the analogous co-expression in mouse brain [25], Table S2, and by the recently established connection between mutations in ADCY5 and movement disorders [26, 27].

In parallel, GNAO1's partner G$\beta\gamma$ regulates calcium, potassium and sodium ion channels [28, 11]. The potassium channels KCNQ2 and GIRK, in particular have been implicated, respectively, in the early onset epileptic encephalopathy (EOEE) [29] and general epileptic phenotypes in human and in animal models [30]. GIRK, in addition, has been reported to be directly regulated by G$\beta\gamma$ in cardiac myocytes, though the subunit through which G$\beta\gamma$ regulates GIRK is the heart specific GIRK4, homologous to the neuronal counterparts GIRK2 and GIRK3 [31].

With the mediators of signaling downstream of GNAO1 only partly understood at the time of this writing, we investigated what can be learned from the core GTPase cycle, without knowing the precise mechanism through which the effectors create the phenotype, or even without knowing the effectors' exact identity.

Functional regions of GNAO1 and the location of disease mutations

GNAO1 processes almost all of the (inter)actions involved in GPCR signaling cycle: its central functional part is the catalytic pocket that is normally occupied by either GDP or GTP at any time, and it also interfaces Gβ, GPCR, RGS, and the effector(s), Movie S1. To these functional regions we should add the folding core - buried residues responsible for the proper folding of the protein. GNAO1 must handle all of these through its relatively small structure; even more restrictively, through its catalytic domain. This has three strong implications: the functional interfaces overlap, all residues in

the catalytic domain belong to at least one functional group, and all residues have very little freedom to vary (Movie S1). In particular, the same interface is responsible for the interaction with Gβ and all of the known effectors of Gα proteins. This further implies that a single mutation can have impact on multiple signalling pathways, and on multiple points in a given pathway.

Response to therapy, from the drug target perspective

Our ultimate goal is to be able to understand which therapy, and why, might be applicable for each genotype. As the first step, then, we systematize the therapy response recorded and reported so far (Dataset S1), according to the location of the resulting lesion on GNAO1 protein.

For each drug we extract the activity values toward its main protein targets, which are typically transmembrane receptors (not GNAO1 itself). Drugs commonly target several members of a protein family, and we do not know for certain which members elicit the effect. To make the information more compact and tractable, we group all members of one target family into a single representative target, additionally labeled by the regulation direction (up- or down-regulation; Fig. 1).

A representative target is assigned a high positive score if it has been targeted with strong affinity, if its effect was documented in multiple patients, and if it has repeatedly been noted as effective in reducing the GNAO1-related symptoms. In the opposite case, when the attempted regulation of the target has repeatedly been documented as ineffective, we assign a high negative score. For details see Supplementary Information (SI) "Building the target-response profiles." In Fig. 1 the score is represented by the color gradient, blue for negative, red for positive, and the results shown as a heatmap.

The most striking feature in Fig. 1 is the clustering of mutation positions into responsive and non-responsive, in particular to GABA and dopamine-receptor manipulation. In the following we will attempt to rationalize that distinction using the existing knowledge about GNAO1 function.

Structural map of disease variants and responses to therapy

Next, we collate the information about the therapy response with the structural location of the disease mutations (Dataset S1). As we noted previously [10], the epilepsy-related (E) phenotypes usually stem from from lesions near the catalytic pocket Fig. 2A, though there is an additional cluster of positions resulting in E phenotypes which cannot be interpreted based on our current understanding of Gα proteins, Movie S1. Movement disorder (MD) only phenotypes are quite strikingly clustered at the effector interface, Fig. 2B.

Rationalizing the phenotype through modeling the signal transmitted by GNAO1 mutants

So why do E or E+MD generating mutations cluster around the catalytic pocket and respond to the reported therapy while effector interface, MD-only mutations do not? A qualitative answer is that the mutations in the catalytic pocket, counterintuitively, lead to hyperexcitable system. This affects both branches of the signal. Treatment for G$\beta\gamma$ entails reducing general neuronal excitability, and, for Gα, enhancing GABA receptors (since GNAO1, at least, *inhibits* its effector, the hyperactive GNAO1 means hyper-downregulated GABA signaling; adding to the complexity of the pathway, GABA-ergic neurons expressing GNAO1 may themselves have an inhibitory role [13]).

At the other end of the problem, and the protein, none of the attempted MD treatments reported so far addresses the issue of broken effector (down-) regulation, as is the case when a variant degrades the GNAO1-effector interaction. In the case of ADCY5, if that is indeed the preferred effector of GNAO1, that would entail blocking or downregulation of adenylyl cyclase [33], for which purpose no drugs have been approved so far [34].

We can however develop a better intuition about the systemic effects of GNAO1 mutations by modeling the interrelated reactions within its signalling cycle.

Modeling the GPCR-related system of biomolecular reactions

We model the GPCR signaling system as a set of reaction rate equations [35], in which the relative concentrations of molecular species (complexes and individual molecules) change in a mutually interdependent way - products of one reaction being the input into another - and all details of the interactions are hidden in the rate constants (Fig. S5). The concentrations [36, 37, 38, 39, 40, 41, 42] and rate constants [43, 44, 45, 46, 47] are chosen to follow as closely as possible the known experimental values (SI, Table S3 and "Modeling agonist dose response"). It is important to note that at this point the model is an explanatory device, rather than a predictive tool, see Discussion below.

Signal transduction in the wild-type GPCR system

Let us first consider, assisted by the simulation, how the system behaves in its wild-type form. Before the arrival of the agonist signal, most of the Gα is inactivated by the GDP in its catalytic pocket, and tied in G trimer with G$\beta\gamma$. For all practical purposes, the exchange of GDP for GTP never happens spontaneously, and the system is thus paused in the equilibrium state (Fig. 3A).

The signal consists in taking the system out of this equilibrium, and letting it relax back (SI, Movie S1). The arrival of the signal, from the perspective of the G-trimer, consists of an active guanine-nucleotide exchange factor (GEF) appearing in the system, with the main effect of enabling the replacement of GDP with GTP. In the GPCR-signaling system, the GEF is the GPCR itself. The 'appearing' - in the functional sense - of the GPCR is the allosteric response to extracellular agonist binding. The ensuing GDP for GTP exchange is accompanied by the disassembly of the trimer and its release from the receptor. From this point on, the signal branches in two directions. In the context of this work, G$\beta\gamma$ is traditionally related to epileptic phenotype (if deregulated), through its interaction with potassium channels, and Gα to regulating cAMP signaling through ADCY, implicated in movement disorders. The cycle of interactions (not to be confused with the signal) is closed with Gα, assisted by GAP, catalyzing the GTP hydrolysis, followed by the reassembly of the G-trimer.

Each of these interactions is an ongoing process that repeats as long as the participating entities are available. Thus, the overall effect of an active GEF is the repartitioning of the available Gα and G$\beta\gamma$ into an active population, consisting of Gα and G$\beta\gamma$ bound to their respective effectors, and the populations of free and GPCR bound G-trimers (Fig. 3B). The relative size of each molecular population depends on the ratio of interaction rates and relative abundance of molecular species (SI, Fig. S5).

Once the signal is terminated (by removal of the GPCR agonist) the system begins its return to the original, equilibrium state. Thus a brief salvo of agonist molecules results in an intracellular signal extended in time, of the characteristic shape [48], shown in Fig. 3C. The GPCR-G signal decays on the timescale of seconds to tens of seconds [49, 50], depending on the mechanism through which the agonist appears and is removed from the system [14]. In our model we use an acetylcholine-like mechanism, in which the agonist molecule is removed by the action of acetylcholinesterase (AChE). AChE cleaves the agonist and thus removes it from the system. This should be understood as a stand-in for any agonist removal mechanism that may be used by a GPCR system (such as dopamine reuptake). The particular choice may modify the timescale but not the signaling cycle itself.

As a measure of the strength of the signal traveling along each of the two branches we use the fraction, or percentage, of Gα and G$\beta\gamma$ effectors engaged in interaction with Gα and G$\beta\gamma$, respectively, Fig. 3C. These interactions are assumed to result, for example, in the regulation of ion current by G$\beta\gamma$ and ADCY inhibition by Gα, which are not explicitly modeled.

Signal modification by changing reaction rates

The height and the shape of the signal depend on the catalytic rate of GNAO1, as well as its affinity for the other molecules in the core signaling cycle. These properties – in our model reflected in the reaction rates (Fig. S5) – can be modified by mutations in the peptide, and depend on the physical properties of the replacement. In principle, each mutation will have some effect on each of the interactions in the system. However, the relative size of the impact on various interaction interfaces is expected to depend on the distance/proximity to each of them. When designing a decision platform for precision medicine, the change in reaction rates should be either evaluated in molecular simulation, or preferably determined experimentally. We choose to limit our assessment of mutations to those that *weaken* an interaction, and observe signal propagation through the system given the reduced interaction rates. In all of the clinical cases reported so far, the mutations were *de novo* and dominant; therefore, in our simulations only half of the Gα population is mutated.

Null mutants

First, let us note that as long as the GNAO1 outnumber their GPCR and effector partners, null mutants – the early stop codons (nonsense and frameshift mutations), and mutations causing the protein misfolding – the only effect of which is to half the GNAO1 population, have no effect on the GCPR signal, Fig. S6. Consistently, no early stop codon mutations were reported in GNAO1 patients, while gnomAD v.2.1.1 [51] reports one healthy person heterozygotic in L269 to nonsense mutation, a position sufficiently early in the sequence for the elimination of the transcript by nonsense mediated decay.

Mutations affecting GTP to GDP catalysis

The most commonly seen dysfunction of GNAO1 mutants can be understood in terms of their reduced catalytic capability. This may be the result of degraded interaction with the activating protein (RGS; Fig. 4A), or the alteration of the catalytic pocket (directly or remotely; Fig. 4B).

Many positions mutated in disease are in the close proximity to the catalytic pocket (G45, S47, R177, N270, D273) all closer than $4\overset{\circ}{A}$ to the substrate; Dataset S1), and given the precision required for the process, it is reasonable to assume that as a result the catalysis is degraded, rather than being enhanced by chance. Additionally, keeping in mind that the catalytic domain is relatively small, it is conceivable that even a mutation not directly within the catalytic pocket can distort its geometry, and exert a similar effect (I56 and D174, tentatively).

Slow catalytic process means that Gα is stuck with the GTP bound, and as long as GTP is present, Gα is primed to interact with the effector, resulting in the extended signal. The increased Gα activity means increased *downregulation* of the effector, which seems to result in MD phenotypes that do not include chorea or ballism. Rather, the phenotypes include severe hypotonia, myoclonic seizures, spasticity, sometimes dystonia.

G$\beta\gamma$ branch follows Gα closely in this case, Fig. 4A. We would expect, thus, that both sets of symptoms react well to a single therapy consisting of reduction of the availability of the agonist, damping down the signal. This is admittedly difficult to discern from the available case descriptions in the literature. The patients *were* reported (Dataset S1) to respond to reduction of monoamine neurotransmitter availability (tetrabenazine) and GABA replacement (vigabatrin). In the cases of extreme dysfunction of the catalytic pocket, no therapy may be effective ("Exotic effects" below; Dataset S1).

Mutations affecting the interaction with the Gα effector

The number of ways in which the Gα signal can be modified exclusively is small. Either direct degradation of the effector interface, or, sometimes, a chance superposition (see below) can generate the effect. Our best candidate for the mutation positions that result in the degraded effector interface

is S207. If the interface with the effector is degraded, we expect the signal of reduced amplitude, though not necessarily of significantly shortened duration Fig. 4C. The problem, from the therapeutic perspective, is that the result is an intensified effector signal, presumably cAMP, for which no approved downregulator exists. The reported phenotypes include severe choreoathetosis, dystonia, and ballism, difficult to treat even in the ICU, Dataset S1 and the references therein. Mostly poor or no response to dopamine and GABA manipulation has been reported in these cases, Fig. 1.

Double impact mutations

Because the catalytic domain of GNAO1 is relatively small, it is feasible for a mutation to have impact on multiple aspects of its function. In particular, some of the experimentally available data [36], can be modelled by assuming that some mutations have double impact, simultaneously affecting the catalytic pocket and the effector binding interface (SI, "Modeling agonist dose response"). Note that this mechanism is different than mutants exerting impact on both Gα and G$\beta\gamma$ signalling branch. Both branches are always affected, with very little exception. Independently of that effect, the shape of the distorted signal depends on which Gα interactions are degraded. For example, while a mutation that slows down the catalysis intensifies and prolongs both branches of the signal, Fig. 4A, a double impact mutation may suppress the Gα branch, while enhancing the G$\beta\gamma$, Fig. 5A. A straightforward example of the latter is provided by mutations in G203 and G204, less than $6\mathring{A}$ from GDP and $8\mathring{A}$ from effector, that result in both EOEE, and severe choreoathetotic symptoms, but without the severe hypotonia that we see in the context of the (exclusive) degradation of the catalysis efficiency.

A particular consequence of double impact mutations is that they may result in superposition of two effects moving the signal in two opposite directions (enhancement vs. suppression). In extreme cases and in the right environment (relative concentration of agonist, GPCR and the effector) this may lead to their mutual compensation, with the resulting signal for one branch close to its wildtype shape, Fig.5B.

A prototypical example is provided by G40, a residue buried deep inside the structure, some $6\mathring{A}$ away from the substrate and $9\mathring{A}$ from the effector. Several cases have been reported, all with bulky amino acid replacements (G40W, G40R, G40E). The behavior of G40R mutant under conditions measured by Feng *et al* [36] can be modelled well under the assumption that it produces sizable effect on both interfaces (SI, "Modeling agonist dose response"). The distance from G40 to the catalytic pocket and the effector is comparable to the G203 and G204 cases, yet the only movement disorders recorded were described as limb twitching. We suggest that this is the result of the effector interface being degraded just to the extent that can be compensated by the concomitant slowing down of the catalytic process, which extends the time in which Gα is available for the interaction.

Another extreme example of this effect id provided by the mutations in E246 and R209, E246K and R209H in particular. In the experiments by Feng *et al* they are indistinguishable from the wild-type. Even more strikingly, gnomAD database records four heterozygote individuals with E246K mutation that are apparently healthy [51]. This time, in the model at least, the ratio of rate constants is moved in the other direction, Fig.5C, masking the impact on the catalytic pocket brought about by disruption of the salt bridge between E246 and R209, exposing only the severe movement disorder phenotype.

Mutations affecting the interaction with the GPCR

The disease mutations involving the Gα C-terminal helix through which it interacts with GPCR are rare (I344del, R349-G352delinsQGCA), and the phenotype relatively mild (seizures limited to early childhood, staring spells; mild spasticity, oromotor apraxia). A single mutated allele with decreased affinity for GPCR should not be able to produce a noticeable effect Fig. 6A if the ratio of GPCRs to Gα is small (see SI, "Parametrizing the GPCR signaling system"). Even when that is not the case, when the ratio is 1:1, the effect is small, Fig. S8, and the reported cases responded to trihexyphenidyl, presumably enhancing the availability of dopamine and thus overcoming the weak baseline signal.

It is possible that the effect of the R349-G352delinsQGCA is exerted through a more subtle change

in the function of the protein (see SI, "Forme fruste of GNAO1 phenotype") leading to the enhanced GDP for GTP exchange. This, in our model, leads to increase in the signal peak intensity Fig. 6B. The patient responded well to levetiracetam, believed to reduce the availability of the agonist.

Epilepsy cluster

When discussing the effector-only impact mutations, an important note should be made of the cluster of positions away from all known $G\alpha$ interacting interfaces, Movie S1, mostly causing epilepsy-only phenotype (peptide positions 231, 275, 279, 284, 291, and possibly 228), Movie S1, suggesting a function that is specific to GNAO1, which means that it is probably *not* related to folding and structure stability (though it should be noted here that Y231C, F275S, and I279N did not express well when transfected into human embryonic kidney cells in the experiments of Feng *et al* [36]). Another possibility is that the effect of these mutations is related to an interaction interface specific to GNAO1. Indeed, that region is conserved in all vertebrate GNAO1s. Taken together, these observations suggest the possibility of so-far unrecognized GNAO1 effector, possibly in different cell types, leading to epilepsy phenotypes not mediated by $G\beta\gamma$.

Exotic effects: empty catalytic pocket

Some mutations may lead to special behavior, completely abolishing some or multiple interactions in the GPCR signaling cycle. The case in question here is G42R mutation, presenting a unique combination of severe hypotonia and choreoathetosis, the first usually associated with extended $G\alpha$ signal, and the latter with the lack thereof. In Fig. S7 we show the behavior of the "empty pocket" mutant - a $Go\alpha$ mutant that does not bind its substrate - described in the series of works by Yu *et al* [45, 46, 47] It has two peculiar properties of blocking its native GPCR, even in the absence of the agonist, while not binding its $G\beta\gamma$. Depending on the ratio of concentrations of $G\alpha$, GPCR and the effector in the system these properties can lead to permanently elevated baseline signal for the $G\beta\gamma$ and almost completely suppressed $G\alpha$, the latter corresponding well with the severe choreoathetosis. G42R is a good candidate for this model because the bulky substitution should completely displace the last phosphate group in the nucleoside making both the binding and the catalysis very unlikely. To move this argument forward, we need to learn how the hypotonia arises in this context in the first place.

Discussion

The continuum of GNAO1 phenotypes, as we have argued, is the consequence of the continuum of changes that a mutation can produce in this protein. Naively, one might expect that a mutation having an impact on a single aspect of GNAO1 function should manifest as a single well-defined phenotype. There are at least two problems with that assumption. One, because of the small, compact, geometry of the protein, it is almost impossible to have a mutation that will have an isolated effect on a single function; and two, because the GPCR signalling cycle branches and closes back onto itself, it is almost impossible to change one branch ($G\alpha$, MD-related, or $G\beta\gamma$, E-related) without affecting the other.

The closest we can come to having a clear-cut impact are the mutations that affect only the catalytic pocket. Even this results both in EEOE and milder, non-choreoathetotic MD symptoms, owing to the branched signal that does not withdraw properly. Accordingly, the patients with these variants seem to react to lowering neuronal excitability or neurotransmitter availability. Increasing GABA-sensitivity of the downstream neurons also makes sense in the context of striatum, since we are reasonably sure that GNAO1 there functions as downregulator of ADCY5 in GABAergic neurons.

If a distant mutation,that is a mutation not in the immediate vicinity of any of the functional sites, weakens the reaction with the effector, the change can be compensated for by increasing the availabilty of the neurotransmitter. We can begin to see, though, the conflicting requirement that it poses - the

distant mutation is likely to have a simultaneous impact on the catalytic pocket, making the G$\beta\gamma$ branch hyperactive and in the need of neurotransmitter *reduction*.

When the mutation completely abolishes the interaction with the effector, as might be the case with some mutations exactly at the interface, an MD-phenotype arises refractory to all reported therapies. We suspect the reason is that the available drugs do not address the underlying problem - the broken downregulation of the effector. It cannot be addressed by enhancing the GPCR cycle, because of its disconnect from the effector, thus the downregulation should be tackled directly. Yet, even for the prototypical effector, ADCY5, the needed downregulator is not available for the purpose, as of this writing (see [34], though).

How do we, then, approach therapeutically a novel GNAO1 variant? Since the phenotypes of various molecular lesions overlap greatly, in particular in our limited ability to describe them, it is unlikely that we will ever be able to discern the impact on the molecular level just by observing the phenotype. Even the mapping of a genetic variant onto protein structure proves to be insufficient, because of the indirect impact it may have through distorting the structure. A dedicated experimental pipeline for quantitative characterization of GNAO1 mutants, in terms of their level of expression and reaction rates with its partners, might be of help to the growing number of patients diagnosed with GNAO1 related disease. Coupled with better understanding of relative abundance of molecular members of the GPCR signalling cycle in the relevant types of neurons, a computational model, such as the one used in our discussion here, would become a predictive tool useful in choosing the first line of drug therapy.

But even in a hypothetical case where we can perfectly quantify the impact of a mutation on each of GNAO1's functions, we are still left with the problem of specifically targeting the system that as its input and output has such generic entities as dopamine and GABA. Looking ahead, it might prove cost efficient in the near future to move to a gene therapeutic approach.

GNAO1-related diseases are caused by autosomal dominant mutations. Our model suggests why: ultimately, the mutants exert their effect by competing nonproductively for the resources in the the signaling cycle, It may seem counterintuitive, then, that in the healthy population there are many nonsynonymous mutations in the highly conserved catalytic domain of GNAO1 [51], Movie S1. An important lesson can be drawn from null mutants, the mutations that completely remove a GNAO1 allele from the pool of expressed proteins. While there are several such mutations reported in the healthy population, none are reported in the context of the GNAO1 disorders. This points to a straightforward path to gene therapy: silencing of the faulty copy of the gene.

Materials and methods

GNAO1 splice

All mutations discussed in this work refer to the GNAO1 splice most commonly expressed in the brain, NCBI accession number NM_020988.2.

Molecular structures - modeling and visualization

The model for active GNAO1 protein structure was build on the template from PDB [52] identifier 3c7k [53], using Swissmodel [54]. The same PDB entry was used to model the interaction between Gα and and RGS domain. The interaction with GPCR modeled after PDB entry 3sn6 [55], and interaction with acetylcholinesterase after 6r3q [56]. Mapping between various complex structures was done using deconStruct [57]. The illustrations were generated using PyMol [58].

Modeling GPCR-G signal

The set of equations describing the core GPCR signaling cycle, Fig. S4, was solved using BioNetGen 2.5 [59]. The input for BioNetGen was initially prepared using V-Cell [60]. The parametrization was taken from the published biochemical experiments [45, 46, 44, 43], details: Fig. S5.

Code availabilty

The full analysis pipeline used in this work can be found in GitHub repository (github.com/dogmaticcentral/gnao1), and the code needed to reproduce figures in the text in the compute capsule on CodeOcean (codeocean.com/capsule/8747824).

References

[1] Nakamura K, Kodera H, Akita T, Shiina M, Kato M, et al. (2013) De novo mutations in gnao1, encoding a gα o subunit of heterotrimeric g proteins, cause epileptic encephalopathy. The American Journal of Human Genetics 93: 496–505.

[2] Gawlinski P, Posmyk R, Gambin T, Sielicka D, Chorazy M, et al. (2016) Peho syndrome may represent phenotypic expansion at the severe end of the early-onset encephalopathies. Pediatric neurology 60: 83–87.

[3] Law CY, Chang STL, Cho SY, Yau EKC, Ng GSF, et al. (2015) Clinical whole-exome sequencing reveals a novel missense pathogenic variant of gnao1 in a patient with infantile-onset epilepsy. Clinica Chimica Acta 451: 292–296.

[4] Marcé-Grau A, Dalton J, López-Pisón J, García-Jiménez MC, Monge-Galindo L, et al. (2016) Gnao1 encephalopathy: further delineation of a severe neurodevelopmental syndrome affecting females. Orphanet journal of rare diseases 11: 38.

[5] Yilmaz S, Turhan T, Ceylaner S, Gökben S, Tekgul H, et al. (2016) Excellent response to deep brain stimulation in a young girl with gnao1-related progressive choreoathetosis. Child's Nervous System 32: 1567–1568.

[6] Talvik I, Møller RS, Vaher M, Vaher U, Larsen LH, et al. (2015) Clinical phenotype of de novo gnao1 mutation: case report and review of literature. Child Neurology Open 2.

[7] Ananth AL, Robichaux-Viehoever A, Kim YM, Hanson-Kahn A, Cox R, et al. (2016) Clinical course of six children with gnao1 mutations causing a severe and distinctive movement disorder. Pediatric neurology 59: 81–84.

[8] Waak M, Mohammad SS, Coman D, Sinclair K, Copeland L, et al. (2017) Gnao1-related movement disorder with life-threatening exacerbations: movement phenomenology and response to dbs. J Neurol Neurosurg Psychiatry : jnnp–2017.

[9] Danti FR, Galosi S, Romani M, Montomoli M, Carss KJ, et al. (2017) Gnao1 encephalopathy broadening the phenotype and evaluating treatment and outcome. Neurology Genetics 3: e143.

[10] Kelly M, Park M, Mihalek I, Rochtus A, Gramm M, et al. (2019) Spectrum of neurodevelopmental disease associated with the gnao1 guanosine triphosphate–binding region. Epilepsia 60: 406–418.

[11] Jiang M, Bajpayee NS (2009) Molecular mechanisms of go signaling. Neurosignals 17: 23–41.

[12] Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, Shen EH, Ng L, et al. (2012) An anatomically comprehensive atlas of the adult human brain transcriptome. Nature 489: 391.

[13] Abela L, Kurian MA (2018) Postsynaptic movement disorders: clinical phenotypes, genotypes, and disease mechanisms. Journal of inherited metabolic disease 41: 1077–1091.

[14] Purves D, Augustine G, Fitzpatrick D, Katz L, Lamantia AS, et al. (2001) Neurotransmitter Release and Removal, Sinauer Associates.

[15] Scheffzek K, Ahmadian MR (2005) Gtpase activating proteins: structural and functional insights 18 years after discovery. Cellular and Molecular Life Sciences CMLS 62: 3014–3038.

[16] Higashijima T, Ferguson K, Sternweis P, Smigel MD, Gilman AG (1987) Effects of mg2+ and the beta gamma-subunit complex on the interactions of guanine nucleotides with g proteins. Journal of Biological Chemistry 262: 762–766.

[17] Clapham DE, Neer EJ (1997) G protein $\beta\gamma$ subunits. Annual review of pharmacology and toxicology 37: 167–203.

[18] Wettschureck N, Offermanns S (2005) Mammalian g proteins and their cell type specific functions. Physiological reviews 85: 1159–1204.

[19] Garcia-Higuera I, Thomas TC, Yi F, Neer EJ (1996) Intersubunit surfaces in g protein heterotrimers analysis by cross-linking and mutagenesis of β. Journal of Biological Chemistry 271: 528–535.

[20] Bastian F, Parmentier G, Roux J, Moretti S, Laudet V, et al. (2008) Bgee: integrating and comparing heterogeneous transcriptome data among species. In: Data Integration in the Life Sciences. Springer, pp. 124–131.

[21] Gray KA, Yates B, Seal RL, Wright MW, Bruford EA (2014) Genenames. org: the hgnc resources in 2015. Nucleic acids research 43: D1079–D1085.

[22] Petrovski S, Küry S, Myers CT, Anyane-Yeboa K, Cogné B, et al. (2016) Germline de novo mutations in gnb1 cause severe neurodevelopmental disability, hypotonia, and seizures. The American Journal of Human Genetics 98: 1001–1010.

[23] Hemati P, Revah-Politi A, Bassan H, Petrovski S, Bilancia CG, et al. (2018) Refining the phenotype associated with gnb1 mutations: Clinical data on 18 newly identified patients and review of the literature. American Journal of Medical Genetics Part A 176: 2259–2275.

[24] Ghahremani MH, Cheng P, Lembo PM, Albert PR (1999) Distinct roles for gαi2, gαi3, and g$\beta\gamma$ in modulation of forskolin-or gs-mediated camp accumulation and calcium mobilization by dopamine d2s receptors. Journal of Biological Chemistry 274: 9238–9245.

[25] Lein ES, Hawrylycz MJ, Ao N, Ayres M, Bensinger A, et al. (2007) Genome-wide atlas of gene expression in the adult mouse brain. Nature 445: 168–176.

[26] Raskind WH, Friedman JR, Roze E, Méneret A, Chen DH, et al. (2017) Adcy5-related dyskinesia: Comments on characteristic manifestations and variant-associated severity. Movement Disorders 32: 305–306.

[27] Carecchio M, Mencacci NE, Iodice A, Pons R, Panteghini C, et al. (2017) Adcy5-related movement disorders: Frequency, disease course and phenotypic variability in a cohort of paediatric patients. Parkinsonism & Related Disorders .

[28] Sprang SR, Chen Z, Du X (2007) Structural basis of effector regulation and signal termination in heterotrimeric gα proteins. Advances in protein chemistry 74: 1–65.

[29] Macdonald R, Gallagher M (2014) The genetic epilepsies. In: Rosenberg's Molecular and Genetic Basis of Neurological and Psychiatric Disease, Elsevier, chapter 84.

[30] Villa C, Combi R (2016) Potassium channels and human epileptic phenotypes: An updated overview. Frontiers in cellular neuroscience 10.

[31] Krapivinsky G, Kennedy ME, Nemec J, Medina I, Krapivinsky L, et al. (1998) G$\beta\gamma$ binding to girk4 subunit is critical for g protein-gated k+ channel activation. Journal of Biological Chemistry 273: 16946–16952.

[32] Waskom M (2020). Seaborn: statistical data visualization¶. URL `seaborn.pydata.org`.

[33] Seifert R, Lushington GH, Mou TC, Gille A, Sprang SR (2012) Inhibitors of membranous adenylyl cyclases. Trends in pharmacological sciences 33: 64–78.

[34] Bravo CA, Vatner DE, Pachon R, Zhang J, Vatner SF (2016) A food and drug administration–approved antiviral agent that inhibits adenylyl cyclase type 5 protects the ischemic heart even when administered after reperfusion. Journal of Pharmacology and Experimental Therapeutics 357: 331–336.

[35] Resat H, Petzold L, Pettigrew MF (2009) Kinetic modeling of biological systems. Computational Systems Biology : 311–335.

[36] Feng H, Sjögren B, Karaj B, Shaw V, Gezer A, et al. (2017) Movement disorder in gnao1 encephalopathy associated with gain-of-function mutations. Neurology 89: 762–770.

[37] Guo W, Shi L, Filizola M, Weinstein H, Javitch JA (2005) Crosstalk in g protein-coupled receptors: changes at the transmembrane homodimer interface determine activation. Proceedings of the National Academy of Sciences of the United States of America 102: 17495–17500.

[38] Milligan G (2007) G protein-coupled receptor dimerisation: molecular basis and relevance to function. Biochimica et Biophysica Acta (BBA)-Biomembranes 1768: 825–835.

[39] Navarro G, Cordomí A, Zelman-Femiak M, Brugarolas M, Moreno E, et al. (2016) Quaternary structure of a g-protein-coupled receptor heterotetramer in complex with g i and g s. BMC biology 14: 26.

[40] Javitch JA (2004) The ants go marching two by two: oligomeric structure of g-protein-coupled receptors. Molecular pharmacology 66: 1077–1082.

[41] Nobles M, Benians A, Tinker A (2005) Heterotrimeric g proteins precouple with g protein-coupled receptors in living cells. Proceedings of the National Academy of Sciences of the United States of America 102: 18706–18711.

[42] Traut TW (1994) Physiological concentrations of purines and pyrimidines. Molecular and cellular biochemistry 140: 1–22.

[43] Calebiro D, Nikolaev VO, Gagliani MC, de Filippis T, Dees C, et al. (2009) Persistent cAMP-signals triggered by internalized G-protein–coupled receptors. PLoS biology 7: e1000172.

[44] Zhong H, Wade SM, Woolf PJ, Linderman JJ, Traynor JR, et al. (2003) A spatial focusing model for g protein signals. Journal of Biological Chemistry 278: 7278–7284.

[45] Yu B, Slepak VZ, Simon MI (1997) Characterization of a goα mutant that binds xanthine nucleotides. Journal of Biological Chemistry 272: 18015–18019.

[46] Yu B, Simon MI (1998) Interaction of the xanthine nucleotide binding goα mutant with g protein-coupled receptors. Journal of Biological Chemistry 273: 30183–30188.

[47] Yu B, Gu L, Simon MI (2000) Inhibition of subsets of g protein-coupled receptors by empty mutants of g protein α subunits in go, g11, and g16. Journal of Biological Chemistry 275: 71–76.

[48] Lodish H, Berk A, Zipursky SL, Matsudaira P, Baltimore D, et al. (2000) Neurotransmitters, synapses, and impulse transmission, WH Freeman.

[49] Yamada M, Inanobe A, Kurachi Y (1998) G protein regulation of potassium ion channels. Pharmacological reviews 50: 723–757.

[50] Holt JC, Jordan PM, Lysakowski A, Shah A, Barsz K, et al. (2017) Muscarinic acetylcholine receptors and m-currents underlie efferent-mediated slow excitation in calyx-bearing vestibular afferents. Journal of Neuroscience 37: 1873–1887.

[51] Karczewski KJ, Francioli LC, Tiao G, Cummings BB, Alföldi J, et al. (2020) The mutational constraint spectrum quantified from variation in 141,456 humans. bioRxiv : 531210.

[52] Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, et al. (2000) The protein data bank. Nucleic acids research 28: 235–242.

[53] Slep KC, Kercher MA, Wieland T, Chen CK, Simon MI, et al. (2008) Molecular architecture of gαo and the structural basis for rgs16-mediated deactivation. Proceedings of the National Academy of Sciences 105: 6243–6248.

[54] Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, et al. (2014) Swiss-model: modelling protein tertiary and quaternary structure using evolutionary information. Nucleic acids research 42: W252–W258.

[55] Rasmussen SG, DeVree BT, Zou Y, Kruse AC, Chung KY, et al. (2011) Crystal structure of the β2adrenergic receptor-gs protein complex. Nature 477: 549.

[56] Qi C, Sorrentino S, Medalia O, Korkhov VM (2019) The structure of a membrane adenylyl cyclase bound to an activated stimulatory g protein. Science 364: 389–394.

[57] Zhang ZH, Lee HK, Mihalek I (2010) Reduced representation of protein structure: implications on efficiency and scope of detection of structural similarity. BMC bioinformatics 11: 155.

[58] DeLano W (2002). The pymol molecular graphics system.

[59] Harris LA, Hogg JS, Tapia JJ, Sekar JA, Gupta S, et al. (2016) Bionetgen 2.2: advances in rule-based modeling. Bioinformatics 32: 3366–3368.

[60] Schaff JC, Vasilescu D, Moraru II, Loew LM, Blinov ML (2016) Rule-based modeling with virtual cell. Bioinformatics 32: 2880–2882.

MUTATION EFFECT ON THE PROTEIN LEVEL

Figure 1: **The response profiles for the protein positions of GNAO1 disease variants with reported therapy.** Each reported drug is characterized in terms of its main target(s). A drug is termed "effective" if it elicits a response. For most of the patients this does not mean resolution of their symptoms. Drug target family names are standard Human Genome Nomenclature Committee names. See Legend in Dataset S1 for the full list. Most drugs target several members of a gene family, and/or multimeric receptors transcribed from several genes. Particularly discriminating are responses to modulation of GABA, dopmamine and muscarinic actylcholine receptors, as well as synaptic vesicle glycoprotein 2A, and transporters from solute carrier group. Clustering of drug response profiles shows variants falling into two categories: drug responsive and drug unresponsive. The latter category overlaps mostly but not completely with mutations at the interface with the effector (see Fig. 2). Heatmap and clustering by Seaborn [32].

Figure 2: **Structural map of disease-related variants in GNAO1, and the main features of the effective therapy reported for their treatment.** Orange, surface representation: GPCR. Deep red surface representation: effector. Cartoon representation, white: GNAO1 protein backbone. Sphere representation: position of missense variants (Dataset S1, Legend). Red: movement disorder (MD), blue: epilepsy (E), green: both phenotypes. Cases when different phenotype combinations were reported in different patients are indicated by the combination of the basic RGB colors, Fig. S9. Gene family names same as in Fig. 1. Arrow up(down): up(down)-regulation or increased (reduced) availability. Strikethrough: the therapy shown to be ineffective. This is a low resolution map, in the sense that it ignores subdivision in response possibly introduced by different amino acid replacements at the same mutation position. **A)** View toward the catalytic pocket. Pink: Mg ion; magenta sticks: nucleotide. **B)** View toward the back side of the catalytic pocket, the putative region of interaction with the effector.

Figure 3: **GPCR signal transiently takes the system out of its equilibrium state. A)** Without agonist present, all G units are bound in trimers. **B)** With agonist, majority the units cycle between trimer, activation by GPCR, interaction with effectors and deactivation with the help of RGS. **C)** Signal: agonist appears, it is removed, and the system returns to its original state. Blue lines: Gα. Yellow lines: G$\beta\gamma$. The interacting partner is indicated by dot notation.

Figure 4: **Signal modification.** By mutation that **A)** changes the interaction with RGS, **B)** alters the efficiency of GTP to GDP catalysis, and **C)** changes the interaction with the effector. Blue lines: Gα signal. Yellow lines: G$\beta\gamma$ signal. Dashed: wild type. The interacting partner is indicated by dot notation. In all cases the mutation is assumed to reduce the forward rate of interaction. Note that the only way for a mutation to downregulate the interaction with the effector is to exclusively modify the interaction with its interface, which is much less likely than having a mutation with multiple effects.

Figure 5: **Compensating effect of double-impact mutations.** A simulated effect of a mutation having a simultaneous impact on the catalysis and effector binding. **A)** In this hypothetical case the Gα signal (blue) is suppressed, and G$\beta\gamma$ (yellow) is increased. However, **B)** a small increase in the catalysis rate can make Gα branch behave almost like the wild type, while **C)** a small decrease can produce analogous effect in the G$\beta\gamma$ branch. Dashed: wild type.

Figure 6: **Impact of mutations at the GPCR interface.** **A)** Effect of weakening the interaction between Gα and the GPCR is nonexistent. In the system in which two alleles compete for a much smaller number of GPCRs, the wild-type allele outcompetes the mutant. **B)** If, however, the mutation enhances the GDP for GTP exchange, the effect is a notable increase in the signal amplitude. Dashed: wild type.