A fungal pathogen that robustly manipulates the behavior of *Drosophila*

***melanogaster* in the laboratory**

Carolyn Elya[1*], Tin Ching Lok[1#], Quinn E. Spencer[1#], Hayley McCausland[1], Ciera C. Martinez[1], Michael

B. Eisen[1,2,3*]

[1] Department of Molecular and Cell Biology, University of California, Berkeley, CA

[2] Department of Integrative Biology, University of California, Berkeley, CA

[3] Howard Hughes Medical Institute, University of California, Berkeley, CA

[*] Correspondence: cnelya@gmail.com, mbeisen@berkeley.edu

[#] These authors contributed equally to this publication

Abstract

Many microbes induce striking behavioral changes in their animal hosts, but how they achieve these effects

is poorly understood, especially at the molecular level. This is due in large part to the lack of a robust system

amenable to modern molecular manipulation. We recently discovered a strain of the behavior-manipulating

fungal fly pathogen *Entomophthora muscae* infecting wild adult *Drosophila* in Northern California, and

developed methods to reliably propagate the infection in lab.-reared *Drosophila melanogaster*. Our lab.-

infected flies manifest the moribund behaviors characteristic of *E. muscae* infections: on their final day of

life they climb to a high location, extend their proboscides and become affixed to the substrate, then finally

raise their wings to strike a characteristic death pose that clears a path for spores that are forcibly ejected

from their abdomen to land on and infect other flies. Using a combination of descriptive, histological,

molecular and genomic techniques, we have carefully characterized the progress of infection in lab.-reared

flies in both the fungus and host. Enticingly, we reveal that *E. muscae* invades the fly nervous system early

in infection, suggesting a direct means by which the fungus could induce behavioral changes. Given the

vast toolkit of molecular and neurobiological tools available for *D. melanogaster*, we believe this newly

established *E. muscae* system will permit rapid progress in understanding how microbes manipulate animal

behavior.

Introduction

Among the most extraordinary products of evolution are microorganisms that are able to manipulate animal

behavior to their advantage. Some have achieved fame in the popular press, like the fungus *Ophiocordyceps*

unilateralis that makes ants wander away from their nests and climb to an optimal height before sprouting

through their heads to rain down infectious spores [1], or the protozoan *Toxoplasma gondii* that suppresses

rodents' innate fear of cat odors to aid the return of the parasite to a cat's stomach where it can sexually

reproduce [2]. However, the mechanisms by which these and other microbes hijack the animal nervous

system have remained elusive.

Entomophthora, from the Greek meaning "insect destroyer", is an aptly-named genus within the basal

fungal lineage Zoopagomycota consisting of species that infect, alter the behavior of, then kill their insect

hosts [3]. *Entomophthora muscae*, first described in 1855 [4] in house flies (*Musca domestica*), is a fungal

species complex that exclusively targets dipterans [5,6]. A fly infected with *E. muscae* exhibits a striking

set of behaviors: shortly before sunset on its final day of life, the fly climbs to a high location (a behavior

known as "summiting"), extends its proboscis and becomes affixed to the substrate on which it stands via

fungal holdfasts [7–9]. The fly's wings then lift up and away from its dorsal abdomen, striking a final death

pose that is thought to be ideal for fungal dispersal [9].

Over the course of the next few hours, the fungus within the dead fly differentiates into structures called

conidiophores that emerge through the weakest points in the fly's cuticle, usually the intersegmental

membranes of the dorsal abdomen, giving the cadavers a distinct banding pattern [10]. A primary conidium

(also referred to as a "spore") forms at the tip of each conidiophore; once mature, these conidia are forcibly

ejected into the surrounding environment in order to land on the cuticle of a susceptible fly host [11,12].

Launched primary conidia are polynucleated, campanulate (bell-shaped) and are surrounded by a sticky

"halo" that serves to adhere the conidium where it lands. If successful in landing on the cuticle of a new

host, the conidium germinates, using both mechanical and enzymatic force to bore through the cuticle and

into the fly's hemolymph [10,13]. If the primary conidium misses its target or fails to germinate upon

landing on the host [14], it can sporulate anew to generate a smaller secondary conidium [11]. Off-target

conidia can continue to re-sporulate and give rise to smaller, higher order conidia until a host is reached or

resources are exhausted [11]. Once inside the fly, the fungus is initially sustained by nutrients in the

hemolymph then later consumes the fat body as an energy source [10]. When available resources are

depleted, the fungus elicits the end of life behaviors and the fungal life cycle begins again.

A range of fly species and even non-dipterans can be infected and killed by *E. muscae* in the laboratory,

though not all infected hosts manifest the stereotypical end-of-life behaviors, and susceptibility has not

been found to track with host phylogeny [15,16]. *E. muscae* has almost exclusively been observed and

studied in muscoid flies (especially the house fly, *Musca domestica*), organisms for which we have very

few experimental tools [17]. Thus, despite inspiring curiosity and intrigue for over a century, how *E. muscae*

achieves control of its host remains poorly understood, with essentially no information as to what is

occurring at the molecular level in either fungus or host.

In 2015, we observed several wild *Drosophila* in separate sites in Northern California with the

characteristic death pose and fungal banding of *E. muscae* infections, and saw an unprecedented opportunity

to study a behavior-changing fungus in the laboratory species, *Drosophila melanogaster*. Here, we describe

the isolation and subsequent characterization of this *E. muscae* strain and its impact on *D. melanogaster*

behavior in the laboratory, and present the *E. muscae-D. melanogaster* as a model for developing a

mechanistic understanding of parasitic manipulation of host behavior.

Results

Discovery and isolation of *E. muscae* from wild *Drosophila*

In June 2015, we established a stable food source (organic fruits in a clean dish pan, referred to henceforth

as the "fendel") at a field site in Berkeley, CA to collect wild *Drosophila* for a separate study (see [18]). In

late July 2015, we noticed that several flies had died with raised wings at the bottom of the fendel and, upon

closer inspection, observed remnants of fungal growth and sporulation on these dead flies (Fig S1A). We

suspected that these animals had been killed by the fungal pathogen *Entomophthora muscae*, though there

have been only a few reports of *E. muscae* infection in wild *Drosophila [19–21]*.

We first confirmed that these flies had been killed by *E. muscae* by genotyping a dozen representative

cadavers at the ITS and 28S (LSU) rDNA loci and searching for similar sequences with BLAST (Fig

S1B,C). PCR genotyping of the host at the cytochrome oxidase II (COII) locus [22] demonstrated that

susceptible host species included *D. melanogaster*, *D. immigrans, D. simulans* and *D. hydei*, which are all

commonly observed in Berkeley, CA. The fungal sequences for all of the typed cadavers were identical at

these two loci, consistent with one *E. muscae* strain being responsible for this epizootic event. Species

identification within the *E. muscae* species complex (which will hereafter be referred to as *E. muscae*) has

historically relied on conidial morphology (and, to a lesser extent, host species), but is expanding to include

molecular data [23]. Still, the taxonomic boundaries between strains and species within this group are still

unclear. To distinguish our strain (or possibly species) from others reported, we will henceforth refer to our

isolate as *E. muscae Berkeley*.

We were intrigued by the possibility that the presence of *E. muscae* in *Drosophila* would allow us to

establish an infection in lab.-reared flies. However, our initial observations were all of dead flies that had

already ejected their spores (Fig S1A). Studies in *M. domestica* have shown that, at room temperature, the

majority of *E. muscae'*s infectious spores are ejected within the first approximately twelve hours of an

infected host's death, and lose infectivity within 48 hours of landing on a non-host substrate [24]. Thus, to

culture *E. muscae Berkeley* we needed to procure freshly-killed flies to ensure access to viable conidia.

The repeated observation of *E. muscae Berkeley*-killed *Drosophila* demonstrated that the infection was

circulating in the population of flies at our field site. We therefore reasoned that some of the flies that were

visiting our fendel should be infected. Previous *E. muscae* research had demonstrated that the fungus only

kills hosts once a day, around sunset [9]. Thus, we collected flies once every morning (1-2 hours after

sunrise) from the fendel and monitored them nightly (1-3 hours after sunset), looking for animals that had

recently died in the stereotyped death pose.

Using a single, wild cadaver, we first established a culture of *E. muscae Berkeley in vitro,* by

inoculating liquid media previously reported to support *E. muscae* growth [23]. Genotyping the resultant

culture at both the ITS and 28S loci verified that we had isolated the same strain as the one that had killed

the previously observed cadavers (Fig S2).

To establish an *in vivo* infection, wild cadavers were co-housed overnight in a confined space with

healthy, lab.-reared CantonS *D. melanogaster*, and exposed flies were monitored nightly for two weeks to

identify *E. muscae Berkeley* cadavers. We repeated this process daily for several weeks before we were

able to passage the infection. We were aware that our standard fly diet contained a small amount of the

preservative tegosept (0.09%), but did not anticipate that this would be problematic since infected wild flies

still died of infection after being housed on this diet for up to eight days (Fig S3). However, it was only

when we began housing flies on food devoid of the preservative tegosept that we were able to successfully

passage the infection.

Once we had transferred *E. muscae Berkeley* to lab.-reared flies, we assessed the impacts of several

variables on infection efficacy, ultimately arriving at an optimized propagation protocol (Fig S4). Briefly,

we expose flies to *E. muscae* by embedding six freshly-killed, infected cadavers headfirst in sucrose agar

and confining 50 young (eclosed within the past 24 hours) CantonS adults of mixed sex with these cadavers

for 24 hours in a cool, humid environment on an inverted 12:12 light:dark cycle. After 24 hours,

confinement is relieved and flies are transferred to a medium free of tegosept. Exposed flies are housed at

room temperature with moderate humidity and monitored daily for death by fungus.

Figure 1. Wild drosophilids killed by *Entomophthora muscae* Berkeley. A) Cadavers found among sampled flies 65 minutes (above) and 40 minutes (below) after sunset. *E. muscae Berkeley* has not grown through the host cuticle. B) Cadavers found among sampled flies 120 minutes (above) and 160 minutes (below) after sunset. *E. muscae Berkeley* has grown through the host cuticle and will soon start to eject conidia. C) Cadavers as discovered *in situ* in fendel at least 12 hours after sunset. *E. muscae Berkeley* has grown through the host cuticle and ejected conidia, some of which have landed on the cadavers' wings.

<u>Description of *E. muscae Berkeley* infection in CantonS flies</u>

With *E. muscae Berkeley* stably propagating *in vivo*, we next focused on carefully observing the process of infection in CantonS flies. By eye, infected flies are hard to distinguish from their healthy counterparts both morphologically and behaviorally until they begin to exhibit end-of-life behaviors (Fig 2A). Exposed flies bear melanized scars that form following spore entry through the cuticle, which are most apparent when the point of entry is the pale ventral abdomen. However, not all flies that are penetrated by the fungus are successfully infected and killed, as we have observed animals with scarring that survive beyond seven days after exposure, and have found that housing exposed flies on diet with anti-fungal significantly improves survival (Fig S4). At 72 hours after exposure and beyond, infected flies generally have more opaque abdomens than uninfected flies due to abundant fungal growth. Under our conditions, ~80% of

CantonS flies are killed four to seven days after exposure to *E. muscae Berkeley*, with the majority of deaths

occurring at 96 and 120 hours (Fig 2B). While by eye infected animals behave normally until the onset of

end-of-life behaviors, analysis of infected fly activity revealed that infected flies exhibit a marked decrease

in total activity compared to healthy counterparts beginning about 36 hours before time of death, which

presently is the best indication of imminent mortality for a given fly (Fig 2C).

On the last day of life, *E. muscae Berkeley* infected flies stop moving 0-5 hours before sunset (Fig 2D).

Taking time of last movement as a proxy for time of death, this observation agrees with reports of *E. muscae*

in house flies [9]. Also consistent with previous reports, flies exposed to *E. muscae Berkeley* and housed

under complete darkness die sporadically throughout the day rather than in a gated fashion (Fig S5, [9]).

As healthy flies housed for 168 hours in complete darkness maintain circadian rhythm, this suggests that

environmental cues and/or a fungal clock are required to coordinate the timing of death, as has been

previously suggested [9]. Of note, flies housed in complete darkness are still observed to die in elevated

positions. This suggests that summiting behavior relies predominantly on gravitaxis rather than phototaxis.

Figure 2. Characteristics of *E. muscae Berkeley* infected CantonS. A) Typical female fly over the course

of infection. Arrows denote conidia that have landed on the cuticle but not yet bored into the hemolymph.

Arrowheads indicate melanization of the fly cuticle that has occurred in response to conidia boring into

hemolymph. B) Time of death for flies infected as per standardized protocol (Fig S4). C) Activity profile

of control flies or *E. muscae Berkeley*-infected flies measured using the *Drosophila* activity monitor

(DAM). D) Time of last movement as measured using the DAM. Each blue circle represents the time of

last movement observed for one cadaver. Flies were exposed to *E. muscae Berkeley* as per standardized

protocol. Dotted line indicates the light-dark transition (L:D transition). E) *E. muscae Berkeley*-infected fly

exhibiting proboscis extension tens of minutes before death. Arrow indicates extended proboscis adhered

to the surface. Real time footage of an *E. muscae Berkeley*-infected fly undergoing proboscis extension is

available as Movie S1. F) *E. muscae Berkeley*-infected fly exhibiting wing raising immediately prior to

death. Arrow indicates original positioning of wings. Time elapsed is given in minutes:seconds. Real time

footage of an *E. muscae Berkeley*-infected fly undergoing wing raising is available as Movies S2, S3. G)

E. muscae Berkeley-killed CantonS summited and adhered to a wooden dowel. Graph to the right indicates

position of death for flies housed in vials without (Dowel -) or with (Dowel +) a wooden dowel. H) Most

commonly observed wing positions of *E. muscae Berkeley*-killed CantonS. Complete wing raising is

observed in most cadavers; wing lowering is consistently observed in a small fraction of cadavers.

On the last day of life, flies infected with *E. muscae Berkeley* show a precipitous decline. The first

portent of imminent death is that flies cease to fly. Though they can still walk and are responsive to

perturbations (i.e. poking with a paintbrush or jostling their container), they will not take flight. After they

have lost the ability (or desire) to fly, moribund flies will begin to exhibit a shaky and slowed gait which is

usually coincident with an upward climbing or movement towards a vertical surface. Many flies reach

elevated positions before they lose the ability (or desire) to continue moving (even when perturbed by the

experimenter), but some succumb to immobility before they leave the ground. When provided a thin,

wooden dowel as a summiting substrate, more flies are observed to die in elevated positions, mostly on the

dowel itself (Fig 2G). Interestingly, we have noticed that when drips of medium are present on the side of

a vial, flies that die on the side of the vial are preferentially found on these drips. It is unclear if this indicates

a preference for the medium as a climbing substrate (versus the smooth plastic of a fly vial) or if the flies

are attempting to eat until their very last.

Once the fly stops walking, it extends its proboscis until it makes contact with the surface on which

it is standing (Fig 2E). The extension of the proboscis is shaky and can occur slowly relative to extension

in response to a nutritive stimulus, and we have observed in multiple instances that the labella of infected

flies do not spread as is typically observed when uninfected flies eat (see Movie S1). Typically, once the

proboscis has made contact with the surface, the fly may move its legs in what appears to be an apparent

attempt to escape, but the material that emanates from the proboscis is sufficient to keep it anchored in

place. After the proboscis has adhered, the fly then begins to raise its wings up and away from the dorsal

abdomen (Fig 2F). This process has been observed to take on the order of ~10 minutes, with wing raising

occurring in small bursts, reminiscent of the inflation of a balloon (see Movies S2 and S3). Curiously, a

persistent minority of infected flies die with their wings lowered down onto their abdomen rather than with

wings elevated (Fig 2H). By applying pressure to the thorax of these flies, the wings are observed to

"toggle" into the upright position, suggesting that the same muscles are involved in raising and lowering.

The fly may continue to twitch its legs and antenna for several minutes after the wings have reached their

final position but will shortly cease moving.

After death, the fungus inside of the fly continues to differentiate into conidiophores, conidia-

launching structures, that grow out into the environment through weak points in the fly's cuticle. Over the

course of several minutes, each conidiophore forms a single primary conidium (Movie S7) which, upon

maturation, is forcibly ejected into the environment. Using time lapse imaging, we observe that conidia

begin to launch approximately five hours after sunset and continue doing so for several hours at ambient

temperature and humidity (Fig 3A). We observed that conidia form and launch asynchronously within a

given cadaver, and not all conidiophores are guaranteed to launch what appear to be mature conidia. Using

high speed videography, we were able to capture the motion of conidial ejection (Fig 3B), and determine

that conidia leave the conidiophore at an initial velocity of ~21 miles per hour (~9.4 meters/second). These

speeds are comparable to those observed in coprophilous fungi, which are among the fastest observed

velocities of organisms relative to their size known in the natural world [25]. In addition, we obtained high

speed footage of primary conidia landing (Fig 3C), which shows conclusively that conidia and halo land

concurrently, an observation that supports the fungal canon mechanism of spore discharge [26].

Figure 3. Fungal transmission from *E. muscae Berkeley* killed cadavers. A) Sporulation time lapse in

E. muscae Berkeley killed cadavers. Time listed in each frame is the time that has elapsed since the light-

dark transition. One image was taken every minute for ~24 hours with three cadavers situated on a cotton

flug at ambient temperature and humidity. The arrow in the second frame indicates the first primary

conidium observed to land on the camera's lens, indicating the start of conidial ejection (i.e. sporulation).

Animated time lapse available as Movie S4. B) Time lapse of the ejection of a primary conidium from a

sporulating cadaver as captured at 54,000 frames per second (fps). Arrowheads indicates conidium that
launches and the vacant conidiophore that remains after launch. Animated time lapse available as Movie
S5. C) Time lapse of a primary conidium landing on the lid of a coverslip as captured at 18,000 fps. The
conidium lands as one complete unit, supporting the fungal cannon mechanism of primary conidium
ejection in *E. muscae*. Arrowheads indicates the position where the primary conidium lands. Animated time
lapse available as Movie S6. D) Primary conidium adhered to glass coverslip and stained with fluorescent
nuclear dye (Hoechst 33342). The conidium is surrounded by a halo of co-ejected material (h).

To compare *E. muscae Berkeley* with other reported isolates, we collected primary conidia and
measured their key morphological traits (e.g. Fig 3D). Our measurements are most similar to primary
conidia from *E. muscae sensu strictu* rather than other members of the *E. muscae* species complex (Table
1). As expected, secondary conidia were observed to form from primary conidia that had landed on non-
productive surfaces (host wing or agar substrate) (Movie S8).

Table 1. Morphology of primary conidia of *E. muscae Berkeley* compared to other reported *E. muscae*
strains.

Report	Isolated from	Infected host	# of nuclei	Diameter of nuclei (μm)	Conidium length (μm)	Conidium width (μm)
Present study[1]	*Drosophila spp.*	*D. melanogaster*	13.9-14.9 (8-22)	3.7-4.1 (2.8-5.8)	23.5-26.1 (19.2-31.8)	18.4-20.5 (14.6-26.7)
[21][2]	*Drosophila spp.*	*Drosophila spp.*	15.3-15.9	4.4	26.0-27.6 (23-30)	21.7-22.9 (17-27)
[27][3]	*M. domestica*	*D. suzukii*	12.8-13.8 (11–16)	NA	27.9-29.1 (25.2-36.8)	22.4-23.2 (19.7-27.6)
[28][4]	*M. domestica*	*M. domestica*	15.2-20.2 (10-27)	3.9-4.4 (3.5-5.5)	26.9-31.1 (21-35)	20.4-24.2 (16-29)

[1] Measurements are given as range in means of three series of 50 objects per host.
[2] Measurements given as reported, unknown number of conidia measured
[3] Measurements are given as reported, nuclei of 12 conidia from one animal were counted, 20 conidia were
measured for length and width

[4] Redescribed type species of *E. muscae* in type host, *M. domestica*, for reference. Measurements were taken

of 8-27 series of 50 objects per host.

<u>Transcriptional profiles of *E. muscae Berkeley* and *D. melanogaster* over the course of infection</u>

To gain a first comprehensive look into how *E. muscae Berkeley* infection progresses in *D.*

melanogaster at the molecular level, we next measured how transcription changes in both host and fungus

at 24 hour time points. We knew that in any given exposure vial there are a mix of infected and uninfected

animals and faced the complication that, early on, infected flies are phenotypically indistinguishable from

uninfected animals. However, we felt confident that we would be able to distinguish infected from

uninfected exposed animals after sequencing mRNA since a) only a few *E. muscae Berkeley* reads would

be needed to confirm that *E. muscae Berkeley* was present and b) it would be unlikely that infected and

uninfected exposed animals would demonstrate identical transcriptional profiles. To favor our chances of

collecting three infected flies for each time point, we collected six *E. muscae Berkeley* exposed CantonS

females at each 24, 48 and 72 hours, three exposed flies at 96 hours, and six fresh cadavers, three at each

96 hours and 120 hours. In parallel, we collected three CantonS females at each time point that were

subjected to a "mock" exposure (housed under identical conditions but in the absence of cadavers). We

prepared and sequenced mRNA libraries from each individual sampled, then aligned the reads to the *D.*

melanogaster transcriptome reference and a bespoke *E. muscae Berkeley* transcriptome reference, which

was assembled from reads that failed to align to the *D. melanogaster* reference. This initial *E. muscae*

Berkeley transcriptome was contaminated by some fly RNAs. After removal of the bulk of contaminating

transcripts based on BLAST alignment and expression levels, our final *E. muscae Berkeley* reference

contained 86,509 transcripts, including a small number of contaminating *Drosophila* transcripts to which

an average of 2.6% of reads from uninfected, control flies aligned (see Methods for details).

We first examined the percentage of reads that aligned to host or fungus in each of our time course

samples (Figure 4A). We observe that *E. muscae Berkeley* reads are low abundance until 72 hours after

exposure at which point a significant portion of the total reads align to the *E. muscae Berkeley*

transcriptome. This likely reflects that the fungus does not begin to actively divide until between 48 and 72

hours and is consistent with our previous observation that *E. muscae Berkeley* rRNA is not reliably

detectable by endpoint reverse transcription PCR until at least 72 hours after exposure (Fig S6). Notably,

the majority of reads from cadavers align to *E. muscae Berkeley* rather than fly. Strikingly, two of our

cadavers show only trace amounts of *D. melanogaster* RNA at the point of sampling.

Figure 4. Gene expression time course of *E. muscae Berkeley*-infected CantonS flies. A) Percentage of

reads aligned to *D. melanogaster* reference versus *E. muscae Berkeley* reference using Kallisto. Samples

are separated into controls (healthy animals who were mock exposed), exposed (animals who exposed to

E. muscae Berkeley and were alive at the time of sampling) and cadavers (animals who were killed by *E.*

muscae Berkeley, dead at the time of sampling) and are color-coded according to the time point at which

they were collected (i.e. 24, 48, 72, 96 or 120 hours). B) *E. muscae Berkeley* expression data from *E. muscae*

Berkeley-exposed and cadaver samples. Complete linkage hierarchical gene clustering by gene was

performed in Gene Cluster 3.0 after filtering out across all genes that are expressed at least at ten TPM in

at least three out of 27 samples (10,809 transcripts total), then log transforming and centering on the mean

value for each transcript. Samples are ordered by percentage of *E. muscae Berkeley* reads as a fraction of

the total reads aligned (above). The scale bar for the heatmap is given to the right of the plot. Two 96 hour

exposed samples that show an aberrant immune response compared to all other exposed samples are

indicated by asterisks. C) *D. melanogaster* expression data from control, *E. muscae Berkeley*-exposed and

E. muscae Berkeley-killed cadavers. Complete linkage hierarchical gene clustering by gene was performed

in Gene Cluster 3.0 after filtering out across all genes that are expressed at least at two TPM in at least three

out of 42 samples (10,875 transcripts total), then log transforming and centering on the mean value for each

transcript. Samples are ordered by percentage of *E. muscae Berkeley* reads as a fraction of the total reads

aligned (above). The scale bar for the heatmap is given to the right of the plot. Two 96 hour exposed samples

that show an aberrant immune response compared to all other exposed samples are indicated by asterisks.

D) Genes that are consistently over or under-expressed compared to controls over the first 72 hours after

exposure to *E. muscae Berkeley*. Top: Volcano plot for all genes over the first 72 hours after exposure. P-

value is determined by ANOVA grouping 24-72 hour control vs. 24-72 hour exposed samples. Genes with

p-value below 0.001 are shown in color. Bottom: Panther GO-term analysis (complete biological process)

of genes overexpressed (red) or under-expressed (blue) in exposed animals compared to controls.

We next surveyed gene expression in *E. muscae Berkeley* across our exposed samples. As different

exposed individuals vary in their rate of infection by *E. muscae Berkeley*, we reasoned that it would be most

informative to order our samples based on *E. muscae Berkeley* titer, which we approximated using the

proportion of reads that aligned to *E. muscae Berkeley* of total reads aligned to either the *E. muscae Berkeley*

or *D. melanogaster* references (Figure 4B). The bulk of transcripts are not expressed until three days after

exposure, which could simply be a consequence of the fungus being low abundance until this time point.

Interestingly, there are three groupings of genes (Groups i-iii) that demonstrate patterns that cannot be

explained merely by fungal abundance in the samples. Group i consists of genes that are expressed early

and depressed later on in infection, Group ii contains genes that turn on during the later phases of growth

in the living host but are turned off after the fly is killed and Group iii consists of genes that do not turn on

until after the fungus has killed the host. At present, there is almost nothing known what gene products

these transcripts encode, as there is little homology between entries in protein databases and the translated

open reading frames in these transcripts.

Next, we examined host gene expression patterns across all of our samples, again ordering samples

based on the proportion of *E. muscae Berkeley* aligned reads among all total and clustering genes by

expression pattern (Figure 4C). Host gene expression segregates into six major groupings (Table S1). Group

i and Group iii contain genes with low expression in controls and early infection, but increased expression

as infection continues, and are enriched for genes involved in epithelial integrity and sensory processes,

respectively. Group ii shows induction of expression all exposed samples except for two taken at 96 hours

(asterisks) and is highly enriched for genes involved in the innate immune response to fungi. Group iv

contains genes that, broadly speaking, are expressed in controls and early infection but not during later

infection and is enriched for genes involved in the metabolism of carbohydrates, steroids and lipids, as well

as cellular respiration. Group v and Group vi largely trend in the same direction, with genes in both groups

tending to be expressed in control samples 72 hours and later and a handful of exposed samples, including

three samples (one 48 hour sample and two 96 hour samples) that behave aberrantly compared to other

biological replicates. These groups are enriched for genes with functions in broad and basic cell functions

including DNA synthesis and repair, transcription, translation and cell cycle control (Group v) as well as

protein localization, mitochondrial translation, autophagy and homeostasis (Group vi).

Following our initial overview of host transcription, we next looked at genes that were consistently

different between control and exposed samples from 24-72 hours (Figure 4D). We excluded all cadaver

samples (both 96 and 120 hours) from this pooled analysis because the animals are dead, and variations in

gene expression would be confounded by mRNA degradation. We also opted to exclude animals at 96 hours

because two of these three samples do not show immune induction (Figure 4C, Group ii). One-way

ANOVA analysis between exposed and control animals from 24-72 hours demonstrated that genes that are

under-expressed in exposed animals are enriched for a handful of metabolic processes, including arginine

and gluatamine synthesis. Interestingly, both arginine and glutamine are amino acids synthesized from the

Kreb's cycle intermediate alpha-ketoglutarate. In times of starvation, the cell would be expected to

prioritize generating ATP via the Kreb's cycle over synthesizing these amino acids. The idea that the fly is

starving is consistent with these enrichments and also with the observation that basic cell metabolism

(macromolecule synthesis) is substantially decreased at 72 hours (Fig S7).

The same analysis shows that genes that are over-expressed in exposed animals are enriched for

immune function, including the melanization defense response and Toll-dependent pathways. *E. muscae*

Berkeley infection relies on boring through the host cuticle which should elicit an initial melanization

response, consistent with our observation. However, it is generally thought that *E. muscae Berkeley* evades

the host immune response once inside the fly because it grows without a cell wall (i.e. protoplastically) and

therefore does not present antigens that can alert the fly immune system to infection [29,30]. Examining

expression patterns of all genes annotated as having immune function, we see a large induction of immune

gene expression at 24-48 hours which includes genes both involved in the melanization response and genes

that specifically respond to fungal infection. In addition, we see overexpression of several groups of

immune genes compared to uninfected controls that persists into late infection (72 and 96 hours) and even

into death (96 and 120 hours) (Fig S8) These data suggest that the initial immune response may not be

strictly limited to wound repair and show that the host immune system remains engaged throughout

infection.

E. muscae Berkeley is present in the fly nervous system 48 hours after exposure

To better understand the process of *E. muscae Berkeley* infection in *D. melanogaster*, we next used

a histological approach to examine the interior anatomy of exposed flies. Analogous to the transcriptomic

time course, we collected adult flies (a mix of 50 males and females) every 24 hours for the first 168 hours

after *E. muscae Berkeley* or mock exposure. Flies were fixed before embedding and sectioning in paraffin

then stained with Safranin O/Fast Green (SFG), a contrast staining method that facilitates the differentiation

of fungal versus host cells (Richard Humber, personal communication), though is more commonly used for

plant histology. We identified *E. muscae Berkeley* morphology by examining *E. muscae Berkeley*-killed

hosts. While there is slide-to-slide variability in the exact hue for a given tissue stained with SFG, generally,

we observed that SFG-stained *E. muscae Berkeley* hyphal bodies have nuclei that stain red (or dark purple)

and cytoplasm that stains purple (Figure 5). *E. muscae Berkeley* nuclei are consistently sized throughout

the host which helps in distinguishing them from host *D. melanogaster* cells.

Figure 5. *E. muscae Berkeley* is consistently present in the nervous system starting 48 hours after exposure. CantonS flies were exposed or mock-exposed (control) to *E. muscae Berkeley* starting 3-5 hours after the light-dark transition and were subsequently sampled at 24, 48, 72 or 96 hours for histological analysis. For each time point, 4-6 individual, paraffin-embedded flies were sectioned at 8 μm, stained using

Safranin O/Fast Green to identify fungal morphology and location and imaged at 20x magnification (Zeiss

Axio Scan.Z1 slide scanner). Only male flies are shown here for ease of comparison. No differences in the

progression of the infection were observed between males and females. An inset of the brain and the

abdomen are shown for each sample. A) Uninfected fly with major anatomical features labeled as follows:

e - eye, b - brain, g - gut, t - testes, f - fat body. B) At 24 hours after exposure there is significant

immunological activity in the abdomen; the nervous system is devoid of fungal cells. C) At 48 hours after

exposure *E. muscae Berkeley* cells are present in the brain (white arrowheads) and/or ventral nerve cord

(VNC) of all but one sample where *E. muscae Berkeley* cells abut but have not yet entered brain. A handful

of *E. muscae Berkeley* cells are observed in the abdominal and/or thoracic hemolymph at 48 hours. The gut

and testes are not invaded by fungus. D) At 72 hours after exposure, *E. muscae Berkeley* can be found

throughout the body cavity and the amount of visible fat body has decreased. *E. muscae Berkeley* titers

have increased in the nervous system. E) In a living fly at 96 hours after exposure (the first point at which

a fly may be killed by *E. muscae Berkeley* infection), fungus occupies virtually all available volume in the

hemolymph. *E. muscae Berkeley* titers have increased in the nervous system, gut and gonads remain

uninvaded. F) In an *E. muscae Berkeley*-killed fly (cadaver) at 96 hours after exposure, only traces of host

organs remain in the abdomen and the nervous system has been considerably degraded. No fat body cells

are observed. *E. muscae Berkeley* cells differentiate into conidiophores, cell-walled structures that will

pierce through weak points of the cuticle to produce and launch infectious conidia. Black scale bars are 100

μm. All living animals shown are males; cadaver's sex is undetermined (the gonads have been consumed

by the fungus.)

We then carefully examined SFG-stained sections from exposed and control flies to determine

where the fungus resides and how fly tissue is impacted over the course of infection (Figure 5). As we

observed no difference in fungal localization between males in females, all samples from a given time point

are described regardless of sex. In control animals, sagittal sections consistently show abundant fat body

cells in the abdomen surrounding the gut and gonads. Fat body is also apparent, though less abundant, in

the head and thorax. The thorax is predominantly occupied by muscles (generally staining red), which are

also observed in the legs. At 24 hours after exposure, we observed hemocyte activity in the abdomen, with

all other tissues indistinguishable from controls. Though hemocyte activity indicates that the immune

system is responding to the fungus, we could not unambiguously identify the fungal cells anywhere in the

body cavity at this time point. We must therefore conclude that the fungus adopts a morphology that is

different from that of cadavers.

At 48 hours after exposure, fungal cells are consistently observed in the brain and/or ventral nerve

cord (VNC; 4 out of 5 samples). In the one case where fungus had not invaded the nervous system, hyphal

bodies were apparent immediately adjacent to the brain, abutting the blood brain barrier. A handful of

fungal cells are also observed in the abdomen or thorax, with some samples showing hemocyte activity as

in 24 hour samples. At 72 hours after exposure, fungal growth is apparent throughout the body cavity and

some hemocyte activity can still be observed. The fat body is depleted compared to earlier time points and

fungus is apparent between muscle fibers, but the gut and gonads all appear indistinguishable from controls.

In addition, fungal titers increase in the brain and VNC. In infected animals that survived 96 hours, fungal

growth is rampant throughout the entire body cavity (head, thorax and abdomen), with the fat body

substantially depleted and fungus residing between muscle fibers. There is no apparent damage to the gut

or gonads. Occasional hemocyte activity can still be observed; fungal titers continue to increase in the brain

and VNC. In *E. muscae Berkeley* killed cadavers, fungus is apparent throughout the body cavity, especially

in the abdomen. The gut and gonads have been completely degraded by the fungus, the brain has begun to

be degraded and the muscles are largely intact.

To confirm that the morphologies observed in the nervous system at 48 hours after exposure and

beyond were *E. muscae Berkeley*, we used fluorescence *in situ* hybridization (FISH) to specifically label

E muscae Berkeley cells within the context of an infected fly. By performing FISH with a fluorescently-

labeled DNA probe targeting the most abundant repeated 18mer in the *E. muscae Berkeley* genome

(~11,000 copies, Bronski, Elya and Eisen, unpublished), we verified that *E. muscae Berkeley* is present in

the brain and VNC in infected animals (Figure 6).

Figure 6. Fluorescence in situ hybridization confirms that *E. muscae Berkeley* resides in the nervous system during infection. CantonS flies were exposed or mock-exposed (control) to *E. muscae Berkeley* starting 3-5 hours after the light:dark transition and were subsequently sampled at 24, 48, 72 or 96 hours for histological analysis. For time points 48 hours and later, at least 3 individual, paraffin-embedded flies were sectioned at 8 μm and subjected to FISH with an *E. muscae Berkeley*-specific 18mer DNA probe labeled at the 5' end with AlexaFluor633. Sections were imaged at 40x magnification on a confocal fluorescent microscope (Zeiss 800 LSM). A) Psuedo-coronal section of a female sampled 96 hours after infection stained with an *E. muscae Berkeley*-specific probe and DAPI. Regions shown at higher detail in B and C located are denoted by white boxes. Scale bar is 200 μm. B) Enlargement of top region in A

showing each DAPI, AlexaFluor633 and merged images of this area. *E. muscae* Berkeley nuclei are

strongly labeled and can be observed inside the host neuropil. Fungal nuclei are also observed in the head

outside of the brain. Scale bar is 50 µm. C) Enlargement of bottom region in A showing each DAPI,

AlexaFluor633 and merged images of this area. *E. muscae* nuclei are observed in abundance outside of gut

and ovaries. Scale bar is 50 µm.

Discussion

A remarkably diverse array of microbes have independently evolved the ability to alter animal behavior.

The prospect of understanding how they do this is intrinsically fascinating, and potentially of great practical

value as a means to understand how animal behaviors are generated and how we might manipulate them in

disease therapy, pest control and other contexts. While an increasing number of researchers have been

drawn to these systems, sustained progress has been limited by the lack of anything approaching an ideal

model system. We believe that the combination presented here of a strain of *E. muscae* that infects wild

Drosophila, protocols for propagating this strain in lab.-reared flies, and the robust induction of behaviors

in the laboratory has the potential as to serve the model system that finally allows us to successfully wield

the tools of modern molecular genetics and neuroscience to describe the molecular mechanisms that

underlie at least one example of microbial manipulation of animal behavior.

E. muscae in wild drosophilids

Though to our knowledge we are the first to study a naturally *Drosophila*-infecting strain of *E. muscae* in

the laboratory, we are not the first to encounter *E. muscae* circulating in wild *Drosophila*. In 1927, Goldstein

reported finding *Empusa muscae* (now *E. muscae*)-infected cadavers of *Drosophila repleta* as well as

Musca domestica at Columbia University in New York state, stating that an epidemic of *E. muscae* had

been observed for the previous four years in this location [20]. In 1969, Turian and Wuest reported

observing *E. muscae*-infected cadavers of wild *Drosophila hydei* in a rotting fruit bait in Geneva,

Switzerland [19]. In 2002, Keller et. al. reported morphological parameters for an *E. muscae* strain

(putatively identified as *E. ferdinandii*, a member of the *E. muscae* species complex) infecting *Drosophila*

spp in Switzerland [21].

Notably, the discovery of *E. muscae Berkeley* has not been our only observation of *E. muscae*

infecting wild fruit flies. In fall of 2014, members of our group caught two individuals from the southern

bay area that were some days later found killed by *E. muscae* (Quan and Schiabor, unpublished). During a

return to the same site in fall 2015, we recovered several additional *E. muscae* cadavers. In the fall of 2016,

a wild drosophilid collected from a site in the north bay was also killed by *E. muscae*. Interestingly, all of

these samples from Northern California, including those found in the fendel, are identical at the LSU and

ITS loci. Samples recovered from a colleague at two distinct locations in Southern California show distinct

sequences at these loci, suggesting that multiple strains (or species) are infecting wild *Drosophila*. Drawing

from these observations as well as other unpublished reports of *E. muscae* infections in fruit flies across the

continental United States (D. Tighe, S. Dara, B. de Bivort), we propose that *E. muscae* infections in wild

Drosophila populations are more common than heretofore recognized. Based on our experience, the

infections seem to be positively correlated with high, local densities of fruit flies in temperate habitats,

which is consistent with how *E. muscae* infection is propagated.

It remains unclear both 1) if the strain(s) or species infecting *Drosophila spp* are distinct from those

that infect other fly species and 2) the degree of specificity for *Drosophila spp* over other dipterans. We

have observed that *E. muscae Berkeley*-infected *D. melanogaster* cadavers are capable of infecting *M.*

domestica in the laboratory, but it is unclear if this infection occurs frequently in the wild. Whether infection

can occur naturally would depend on the ecology of the two different host species (i.e. if they interact

frequently enough to expect the exchange of *E. muscae* infection) Our understanding of strain diversity and

host specificity would greatly benefit from the collection of more molecular and ecological data.

<u>The progression of *E. muscae Berkeley* infection in lab.-reared *D. melanogaster*</u>

Taken together, our RNAseq and histology time course data describe the typical progression of *E. muscae*

infection in lab.-reared *D. melanogaster*. At 24 hours after exposure, flies show a robust antifungal immune

response, though the fungus is nearly undetectable within the fly by histology, indicating that it is at low

titer. At 48 hours, the fungus has begun to adopt the morphology which it will assume until killing the host.

Fungus is observable within the host brain and VNC, though overall fungal titer is still quite low. As the

abdomen is the most likely point of initial entry for the fungus (it is the biggest target for the fungus to hit),

we suspect that the fungus has travelled from the point of entry to the CNS, indicating tropism for neural

tissue early in the infection. Elements of the host's immune response are still activated. At 72 hours, fungus

was apparent throughout the body cavity, in the thorax (between muscle fibers), abdomen (surrounding but

not invading the gut and gonads) and also in the limbs and halteres. The fat body is significantly depleted

by this point; the host's dampened metabolism suggests an internal starvation state. At 96 hours, if the fly

has not succumbed to infection (i.e. there are still energy reserves available to the fungus), fungal titer will

continue to increase and the fat body will continue to be depleted. Two-three hours after death, flies that

have been killed by the fungus show no intact abdominal organs and nervous systems that are being broken

down.

While the trajectory of infection is consistent, it is important to recognize that just because two

animals have been exposed for the same duration of time that these two animals will not progress through

infection identically. In our RNAseq data, we noticed that the host gene expression in exposed animals at

24 and 48 hours tended to be more variable than those for 72 hours. We imagine that this was due to chance,

that we picked animals that were at similar points in infection at 72 hours, whereas we picked animals that

were more variable at other time points. This may at least partly explain why we observed so much

differential expression at the 72 hour time point in exposed versus controls whereas less was observed at

earlier time points, especially at 48 hours. It is likely that several factors play into whether or not an infection

succeeds and how quickly it progresses (e.g. initial exposure titer, size of host, nutritional status of host

etc.). Thus, future work should consider how to determine a metric to gauge progress of infection so that

similarly-progressed samples can be compared.

E. muscae Berkeley infection and host immune response

The entomopathogen community has believed that *E. muscae* evades the host response by growing

protoplastically (i.e. without a cell wall, components of which would be recognized and targeted by the host

immune system). In both the gypsy moth and the greater wax moth, it has been shown that the host immune

cells recognize walled *Entomophaga* fungal cells, but there is little cellular response to protoplasts [31,32].

Based on these findings, it has been posited that the host does not detect the ever-increasing fungal mass

within until the end of infection when the fungus puts on a cell wall that contains epitopes that the host can

recognize [29,30]. As a result of ostensibly evading the immune system, it has also been hypothesized that

*E. musca*e does not generate toxins, as it would have no incentive to do so in the absence of attack by the

host [29].

Our data show that there is a robust initial response to *E. muscae Berkeley* exposure. Many of the

immune genes that are induced with *E. muscae Berkeley* have also been observed to be induced by exposure

to other, more generalist fungal pathogens (e.g. *Beauveria bassiana*, *Metarhizium anisopliae* [33,34]).

These data clearly indicate that the host detects an invader early on in infection. Furthermore, there is a

detectable immune response through the length of infection (Fig S8), though at this point we cannot say if

this response is a slow disengagement of the initial response or stimulated *de novo* by the growing fungus.

Interestingly, the living animals sampled at 96 hours for RNAseq are inconsistent in their host

transcriptional immune response: two of the three animals more closely resemble control animals than

infected animals in host transcription (Figure 4C). At least two scenarios could explain this observation. It

is possible that both of these animals were in the process of recovering from infection (i.e. the immune

system was effectively combatting the fungus) or there was a delay in the course of infection compared to

contemporaneous samples. The proportion of fungal reads present in these samples is lower than what

would be expected for late time points, which is consistent with either scenario. At this point we simply do

not know if every instance of a spore hitting a fly leads to a productive fungal infection. There is some

evidence to the contrary: we have consistently observed that some highly-exposed flies die prematurely.
These animals are generally smaller than others in the vial and are often covered in spores. This could
indicate that getting hit by too many spores (an unlikely outcome in the natural world) leads to an
overwhelmed fly (e.g. overactive immune system or accelerated fungal growth) that dies before being
manipulated. These flies do not sporulate, though it is possible that they do produce resting spores. On the
other hand, we have observed that survival of exposed flies is substantially increased when flies are exposed
to small quantities of anti-fungal. This indicates that there are ways of either halting or slowing an infection,
though whether the fly's immune system is generally capable of doing this is unknown.

Though five of our six cadavers sampled for transcriptomics have similar levels of immune gene
transcripts compared to *E. muscae Berkeley*-exposed animals sampled at 72 hours, the sixth cadaver
exhibits higher expression of anti-fungal peptides Drosomycin and Metchkinowin, the beta-glucan receptor
GNBP3-like and several IM family genes. It is possible that this fly is demonstrating the proposed immune-
system overload and resultant spike in immune system gene expression [29]. It might be the case that this
immune spike occurs in all animals; we could have sampled too late to observe it in the other five samples
but were able to see it in one sample that was late to respond. However, one can imagine that in the presence
of copious fungal epitopes the fly immune system would continue to go be highly engaged until death, not
drop back down to levels comparable to earlier time points (i.e. 72 hours), as seen in the majority of sampled
cadavers. At present, we are inclined to interpret this odd-sample-out as a fluke rather than an indication of
a moribund immune spike.

<u>Why is *E. muscae Berkeley* in the brain?</u>
Our work demonstrated that *E. muscae Berkeley* is present in the nervous system system relatively early in
infection, just 48 hours following exposure. *E. muscae Berkeley*'s invasion of the nervous system grants
the fungus direct access to host neurons and may be mechanistically important for achieving behavioral
manipulation of the host fly. However, we should be careful to consider any and all possible ways *E. muscae*
Berkeley could alter host behavior before jumping to this conclusion.

We can imagine four general mechanisms by which *E. muscae* is able to achieve behavioral
manipulation. The fungus could invade the nervous system in order to localize adjacent to and impinge on
the activity of particular neurons through chemical or physical means. However, we are skeptical that this
is the case as our observations do not support specific localization of the fungus in the CNS.

A second possibility is that the fungus invades the nervous system in order to gain access to either
a particular group or groups of neurons or all neurons generally, but does not localize within the CNS in a
stereotyped manner. Rather it is sufficient that it has crossed the blood brain barrier, which insulates the
nervous system from the activities in the hemolymph and allows for the selective transport of compounds

to and from the hemolymph, allowing the fungus to modulate the activity of neurons by secreting

compounds that diffuse throughout the CNS. The secreted compounds could be specific, only altering the

activity of a subset of susceptible neurons, or could be more general, changing activity over many or all

neurons.

A third possibility is that the fungus does not need to invade the nervous system in order to change

the host's behavior. The fungus could be secreting a compound into a hemolymph that is capable of crossing

the blood brain barrier and altering neuronal activity. Alternatively, the fungus could be secreting a

compound into the hemolymph that changes the host's internal state (either directly or by leading the host

to respond in a way that causes the internal state to change) which leads the animal to respond by executing

one or more of the end-of-life behaviors.

Lastly, it's possible that the fungus does not secrete compounds to induce these behaviors, but by

destroying fly tissues elicits the series of observed behaviors. While we believe this last scenario to be

highly unlikely, it cannot yet be ruled out.

For these last two proposed mechanisms, the fungus would not need to invade the CNS in order to

affect behavior. In these cases, the fungus could be invading the CNS as a means of escaping immune

surveillance. By establishing a reservoir in the CNS, the fungus could replenish dying cells in the

hemolymph in order to ensure that the infection took hold. Alternatively, the fungus could invade the CNS

because it provides a rich, nutritive substrate to sustain the fungus. This scenario is inconsistent with our

histological data both from flies that are not executing end-of-life behaviors (Figure 5) and flies that are

executing end-of-life behaviors show that the brain is largely intact, indicating that the fungus abstains from

consuming these tissues until host death.

Our observation that *E. muscae Berkeley* invades the host nervous system contrasts with the recent

report that *Ophiocordyceps unilateralis,* a fungal pathogen that induces very similar end-of-life phenotypes

in the ant host *Camponotus castaneus* is absent from the brain at the point of behavioral manipulation [35].

Interestingly, another Entomophthoralean fungus, *Strongwellsea magna,* is also known to invade the

nervous system of its lesser house fly host (*Fannia canicularis*) during infection [36]. In this case, the

author proposed that this did not have consequences for behavior.

Surprisingly, transcriptomic analysis of dissected brains from exposed females at 24, 48 and 72

hours with confirmed *E. muscae Berkeley* infections failed to show differential gene expression compared

to uninfected controls but did show an increase in *E. muscae Berkeley* titer (taking *E. muscae Berkeley*

reads as a proxy) (Fig S9). Though these samples were not collected at the point of behavioral manipulation

by the fungus, it is surprising that there are no major transcriptional changes within the brain at these time

points, and suggests that behavioral modification may be largely independent of transcriptional changes in

the brain.

<u>Behavior and beyond: the utility of the *E. muscae Berkeley-D. melanogaster* system</u>

The past decade has seen an explosion of tools for characterizing and manipulating the nervous system of

D. melanogaster, including a catalog of the types and corresponding expression patterns of its

approximately 100,000 neurons, a complete map of connections in the brain, reagents for conditionally

activating or disabling specific sets of neurons as well as purifying these cells, and methods for

automatically tracking and classifying behaviors in populations. Our development of a robust system of

microbially induced behavior manipulation in *D. melanogaster* will allow us, and we hope many others, to

leverage the powerful molecular and neurobiological toolkit of *D. melanogaster* to explore the molecular

basis of this fascinating but still mysterious biological phenomenon.

Acknowledgements

This research was supported by the MBE's Investigator Award from the Howard Hughes Medical Institute

and CE's Graduate Research Fellowship from the National Science Foundation. This work used the Vincent

J. Coates Genomics Sequencing Laboratory at UC Berkeley, supported by NIH S10 OD018174

Instrumentation Grant. The authors are grateful to Richard Humber whose expertise in all things

Entomophthora, and his eagerness to educate us, was invaluable throughout, Michael Vahey and Brian

Belardi in Dan Fletcher's group for assistance with high-speed videography, Ciera Martinez for developing

a protocol to embed and section flies in paraffin, Steve Ruzin and Denise Schneides at the Berkeley Imaging

Facility (College of Natural Resources, UC Berkeley) for their expertise and patience in microtomy and all

things microscopy, Jen-Yi Lee for her guidance in using equipment in the Molecular Imaging Center

(College of Natural Resources, UC Berkeley), Kristin Scott, Russell Vance, Damian Elias and Richard

Calendar for advice, feedback and support during this project, William Ludington for his reminder that

"fruit flies like a banana", and Michael Bronski for introducing the Eisen Lab to the wonderful world of

entomopathogens. CE also acknowledges the support of Nora, Bruce and Kevin.

Author Contributions

CE discovered *Drosophila* infected with *Entomophthora* in her backyard, figured out how to reliably

passage the infection to lab-reared flies, carried out all of the experiments described in this paper, made all

the figures and wrote the manuscript. TCL and QES helped maintain populations of infected flies for

experiments and provided experimental assistance, and HM assisted with the early characterization of the

Entomophthora infection. MBE encouraged the pursuit of this project by CE, obtained funding for the

research, provided feedback and ideas throughout, and edited and advised on figures and the manuscript.

Funding

All of the research described in this paper was funded by an HHMI Investigator award to MBE. CE was

supported by a National Science Foundation Graduate Research Fellowship. CCM was supported by a

National Science Foundation Postdoctoral Fellowship.

Materials and Methods

Fly husbandry

Healthy wild-type, CantonS Wolbachia-free (WF) D. melanogaster were reared on Koshland diet (0.68%

agar, 6.68% cornmeal, 2.7% yeast, 1.6% sucrose, 0.75% sodium tartrate tetrahydrate, 5.6 mM CaCl2, 8.2%

molasses, 0.09% tegosept, 0.77% ethanol, 0.46% propionic acid) supplemented with activated dry yeast

pellets (Red Star) at 21C on a 12:12 light:dark photoperiod. Any time the photoperiod needed to be adjusted,

flies were reared from third instar or earlier on the desired photoperiod to ensure that emerging adults were

properly entrained.

Fendel tending

Wild fruit flies were caught by directly aspirating from an uncovered plastic dishwashing pan (aka the

"fendel") that was baited with quartered organic watermelon and an assortment of other organic fruits.

Aspirated flies were transferred onto Koshland diet and housed at ambient temperature and ambient

humidity. Baiting and capture was performed in the spring through early fall of 2015 at a personal residence

in Berkeley, CA.

PCR genotyping

DNA was extracted from individual cadavers or 1.5 mL of *in vitro* culture using the QIAamp Micro Kit

(QIAGEN) following the tissue protocol. These DNA preparations were used to amplify the desired

sequences.   Entomophthora-specific   ITS   primers   (emITS: emITS-1   5'-

TGGTAGAGAATGATGGCTGTTG-3', emITS-4 5'- GCCTCTATGCCTAATTGCCTTT-3') or fungal-

specific large subunit primers (LSU: LR3-1 5'- GGTCCGTGTTTCAAGAC-3', LR0R-4 5'-

GTACCCGCTGAACTTAAGC-3') were used to genotype Entomophthora (James et al 2006); cytochrome

oxidase   II   primers   (tLEU: 5'   ATGGCAGATTAGTGCAATGG   3'   and   tLYS: 5'

GTTTAAGAGACCAGTACTTG 3') were used to genotype infected Drosophila hosts (Liu and

Beckenbach (1992). Each reaction was performed using GoTaq 2x colorless mastermix (Promega) with

800 nM of each forward and reverse primer with the following thermocycling conditions: 95C for 5 min

followed by 35 iterations of 95C for 30 seconds, 51C for 30 seconds then 72C for 1 min/kb then 72C for

an additional 10 minutes. Reactions were checked by gel electrophoresis in 1% agarose. Successful

reactions were prepared for sequencing using ExoSap-IT (Affymetrix) per manufacturer's instructions and

submitted with each amplification primer for Sanger sequencing. Assembly of forward and reverse reads

was attempted to generate a consensus sequence using Seqman Pro (DNA Lasergene v.10). Sequences were

searched against the BLAST NT database using blastn.

Isolating *E. muscae Berkeley in vitro*

To grow E. muscae Berkeley in vitro, first spores were collected using the ascending conidia collection

method (i.e. by placing a fresh cadaver in the bottom of a sterile petri dish and allowing the cadaver to

sporulate overnight) (144). The following morning, the lid of the dish was rinsed with 10 mL of Grace's

insect medium (1x) supplemented with L-glutamine, 3.33g/L lactalbumin hydrolysate, 3.33g/L yeastolate

(ThermoFisher Scientitfic # 11605- 094) containing 5% fetal bovine serum (Invitrogen) and transferred to

a vented, treated T25 tissue culture flask (Corning # 10-126-10) using sterile technique. The culture was

then incubated at room temperature in the dark until growth was apparent (non- homogenous white spheres

floating in the medium). The culture was genotyped with emITS and LSU primers to confirm that it was E.

muscae and was an identical match to the cadaver that produced the spores which started the culture

("Fendel mama") at these loci. The culture was periodically examined at 100-400x on a compound

microscope to confirm proper morphology and absence of contamination.

Isolating and optimizing *in vivo E. muscae* Berkeley infection

Wild fruit flies sampled from the fendel and housed on Koshland food were monitored nightly for two

weeks after capture for deaths due to *E. muscae* Berkeley. Freshly killed cadavers were separated from their

living conspecifics by briefly anesthetizing live flies via cold anesthesia (incubating 2-3 minutes in a

residential freezer). Fresh cadavers (anywhere from 1-10, depending on availability) were placed on chunks

of organic banana in a wide Drosovial with kimwipes to sop up excess moisture. Approximately 50 healthy

flies were then transferred onto the prepared vial by flipping (no anesthesia was used). The plug of the vial

was pushed down to confine the flies within a few centimeters to improve the likelihood that they would

encounter flying spores. Leaving the exposed flies with the spent cadavers was initially problematic as we

were working without access to anesthesia or a microscope and had to identify new cadavers by naked eye.

Additionally, the raw banana began to ferment and break down, leading to excess moisture which was
prematurely killing some of our exposed flies. To avoid these issues, the exposed flies were transferred to
a new banana/kimwipe vial after the first 48 hours. This was done by allowing the living flies to climb into
an empty "holding" vial then flipping them onto the fresh vial. The flies were monitored daily for deaths;
cadavers were removed after allowing healthy flies to climb into a "holding" vial and used to start new
exposure vials.

In vivo propagation of *E. muscae Berkeley* infection

Cadavers are collected daily 2-5 hours after the end of the 12 hour light cycle from exposure vials that are
between 96 and 168 hours (4 and 7 days) old. All flies that will die on this day because of *E. muscae*
Berkeley infection will be dead by this time and will be extremely swollen with fungal growth, making
them obvious among the living flies. CO_2 is used to anesthetize the living flies in exposure vials and collect
cadavers, which are placed in a petri dish with a piece of Whatman paper wetted with DI water to mitigate
static. Molten cadaver embedding medium is preparing by microwaving solidified AS solution (1.5% agar,
10% sucrose) and poured into a clean 100 x 15 mm petri dish just enough to cover the bottom of the dish.
As soon as the agar has set, six cadavers are embedded head first in a circle of diameter <2 cm with their
wings on the outside of the circle. The wings are pressed into the agar to ensure they do not intercept any
launched conidia. The agar is allowed to completely set before continuing. The cadaver circle is cut out
from the set agar by using an empty, wide-mouth Drosophila vial (FlyStuff). The agar disc containing the
fly circle is then transferred, cadaver-side up, into an empty, wide-mouth Drosophila vial. A ruler is used
to mark 2 cm above the surface of the agar. CantonS flies reared on the same 12 hour light cycle on
Koshland medium are collected from eclosing vials using CO_2 anesthesia. Fifty healthy flies are added to
the vial container the agar disc with cadavers and tapped down until all flies are under the 2 cm mark. A
Droso-plug (FlyStuff) is pushed into the vial such that its bottom is flush with the 2 cm mark. The vials is
incubated for the first 24 hours at 18C in a humid chamber (~95% humidity, 2L plastic beaker lined at the
bottom with wetted paper towels or kimwipes and covered with foil), to encourage sporulation. After 24
hours, the Droso-flug is lifted to relieve the confinement of the flies and the vial is moved to a 21C incubator
(~60% humidity). After 48 hours, the exposed flies are transferred onto GB+ medium (40% organic banana
[w/v], 2% agar, 0.3% propionic acid) without anesthesia; incubation continues at 21C. Cadavers are
collected daily 2-45 hours after the end of the 12 hour light cycle from exposure vials that are between 96
and 168 hours (4 and 7 days) old. The process is repeated daily to supply cadavers for experiments and
ensure the infection is maintained.

Photography and videography

Pictures and videos of dead and dying flies (extending probosces and raising wings) were taken with a

Nexus 5x (Google) or iPod Touch (Apple) aided by attaching a macrolens (Luxsure) to the device camera

lens or by aligning the device camera lens with the eyepiece of a dissecting scope. Images were taken under

ambient light, humidity and temperature.

Time lapse microscopy was taken via a USB microscope (DinoLite Digital Microscope Pro) using

DinoLite software v1.12 (Figure 3A) or on a Nikon 80i compound microscope equipped with a Hamamatsu

black and white camera (C11440) using MetaMorph software (v. 7.8.00, Molecular Devices) (Figure 3B,C).

Each time lapse consists of images collected once a minute for the indicated duration. Images were taken

under ambient temperature and humidity.

High-speed videos (18,000-54,000 fps) were filmed with a 5x objective on a Axiovert 200

microscope (Zeiss) equipped with an Photron Fastcam 1024PCI. Cadavers were mounted in 1.5% agar on

a glass slide and arranged such that forming conidiophores and conidia were visible (for spore ejection) or

such that cadavers sporulated onto a #1 glass coverslip in the plane of the camera (for spore landing). Video

was captured via Photron Fastcam Viewer software, running at the indicated frames per second in end

trigger mode (new frames were continually captured and old frames dumped until the user triggered the

filming to stop). Spores or landing sites were manually watched until a spore disappeared or appeared, then

video was stopped and last ~10 seconds of footage were manually searched for the spore launching or

landing event.

<u>Circadian timing of death</u>

CantonS WF flies were reared on a 12:12 light cycle (photophase 1 am – 1 pm or 7 pm - 7 am PDT, as

indicated). For experiments run in complete darkness, 25 flies were exposed to E. muscae Berkeley via the

standard anesthesia-dependent protocol. All exposure vials were set up before the utilized cadavers

sporulated (within 2 hours of the light-dark transition). Exposure vials were housed in a humid chamber in

a dark 21C incubator wrapped in blackout cloth for approximately 24 hours before loading into Drosophila

activity monitors (DAMs, Trikinetics). Before loading flies, DAM tubes (5 mm x 65 mm, polycarbonate)

were prepared containing such that one end of the tube held approximately one cm of 1.5% agar, 10%

sucrose and was wrapped in parafilm to prevent drying out.

To load the flies, all accessible light sources were disabled before removing the humid chamber

from 21C and placing on the bench at RT. Vials of flies were kept under a foil-lined box as they waited to

be processed. One vial at a time was retrieved from the box, knocked out with CO_2 under a dissecting scope

whose LED light ring was covered with a red gel filter (Neewer), sorted by sex and loaded into individual

DAM tubes with the aid of a red LED flashlight (KMD Aero) before capping each tube with an autoclaved

cotton plug. For each vial, eight females and eight males were chosen for loading into DAM vials.

Unexposed flies (i.e. controls) were always processed before proceeding to exposed flies. CO2 pad was

wiped down with 70% ethanol between vial types to present cross-contamination. DAMs were loaded from

bottom to top row, filling a row and securing each tube with rubber bands before proceeding to the next.

Loaded tubes were kept under a foil-lined box to prevent light exposure. When all loading was finished,

DAMs were covered with blackout cloth and transported back to the 21C incubator. There they were

attached to the DAM interface unit and recording began, binning counts in 30 second intervals. Recording

continued for ~170 hours until the experiment was stopped.

Like with loading, experiments were stopped by first disabling all light sources, then carefully

disconnecting and removing DAMs from the 21C incubator to not disturb adjacent experiments, and sealing

incubator before turning on the overhead lights for manual inspection. Each DAM tube was inspected to

see if the fly was dead or alive. If dead, the tube was inspected for evidence of sporulation to determine if

the cause of death was patent *E. muscae* infection. For experiment run on a 12:12 light cycle, flies were

exposed as above but without concern for light contamination; resultant DAMs were housed on a 12:12

light cycle for the duration of the experiment. Each channel was checked daily following sundown to see

which flies had died within the previous 24 hours.

DAM data were processed using Python to determine time of last movement (accurate to 30

seconds) and to plot movements over time. For each channel, the reported time of last movement was

manually cross-checked using the plot of activity data. In cases where there was an erroneous movement

(i.e. a signal occurring more than 24 hours after the fly's last movement), the time of last death was manually

re-assigned. For data visualization, data were binned into 15-30 minute intervals and the average

movements of unexposed animals (controls), exposed or entrained light cycle of exposed flies and cadavers

and the time of last movement for each observed cadaver were plotted in Prism (GraphPad). There were

generally no obvious differences between male and female activity for the unexposed animals so sexes were

combined for data analysis.

<u>Collection and staining of primary conidia</u>

Three to six fresh cadavers (i.e. those who had not yet sporulated) were collected from exposure vials using

the anesthesia dependent methods detailed above. Sporulation chambers were prepared as follows: a small

piece of Whatman paper was placed in the base of a small petri dish (60 mm x 15 mm) and wetted with DI

water. A bloated cadaver was chosen for each chamber and its wings were removed. The cadaver was

placed in the middle of the Whatman paper and the chamber was topped with a custom, 3D-printed top that

included a square opening slightly smaller than a standard 22x22 mm coverslip. The top and bottom were

sealed using parafilm and a new coverslip was placed over the opening. Cadavers were left in the chambers

at room temperature to sporulate. Coverslips were changed every 30 minutes to 1 hour, as needed, and

promptly stained for microscopy by applying Hoechst (1 ug/mL). Spores were imaged on a compound

microscope at 40x for measuring conidia attributes; exact distances were determined by calibration with a

0.01 mm micrometer (OMAX). For each attribute (number of conidia, length and width of conidia, diameter

of nuclei), at least 50 different primary conidia were counted from three different cadavers.

RNA experiments

RNA was prepared from each thawed sample by homogenizing with an RNase-free pestle (Kimble Chase),

washing the pestle with 750 uL Trizol, then proceeding using the manufacturer's protocol. For reverse-

transcription PCR (RT-PCR) and transcriptomic experiments, three mock vials and three Three mock vials

and three exposure vials were started with 25 CantonS WF flies 0-1 days old (RT-PCR, whole flies) or 1-2

days old (dissected brains) with either 0 (mock) or 6 (exposure) cadavers embedded in AS. Flies were

incubated for the first 24 hours at 18C confined to 2 cm with cadavers, then moved to 21C where the

confinement was relieved. Flies were transferred to GB+ at 48 hours where they continued to be housed at

21C. (Vials were sampled every 24 hours for 72 (dissected brains), 96 (RT-PCR) or 120 (whole flies) hours

by anesthetizing the entire vial with CO2. Exposed animals were preferentially selected based on evidence

of contact with fungus (e.g. spores or melanization responses thereto visible on cuticle). Sampling for each

time point consistently occurring between 2-3 hours following the light-dark transition. Before sampling,

all equipment used to manipulate flies (e.g. CO2 pad, forceps etc.) were treated with 10% bleach, wiped

with DI water then sprayed with 70% ethanol. All materials that handled flies (CO2 gun, pad, forceps) were

treated with 10% bleach and rinsed with DI water between sampling exposure vials. Control vials were

always sampled first. Sampled material (either whole fly or dissect brain) was immersed in 250 uL Trizol

then immediately flash frozen with liquid nitrogen. Samples were stored at -80C until extraction.

RT-PCR

One control female (24-96 hours) and two infected females (24-72 hours) or one fresh cadaver (96 hours)

from each vial were collected as described above. RNA was prepared from each thawed sample by

homogenizing with an RNase-free pestle (Kimble Chase), washing the pestle with 750 uL Trizol, then

proceeding using the manufacturer's protocol. RNA was then treated with Turbo DNase (ThermoScientific)

per the manufacturer's protocol and quantified using a Qubit Fluorometer (Qubit RNA HS assay kit,

ThermoFisher Scientific). For each sample, 1 μL or 160 ng of DNase-treated RNA, whichever was more,

was added to a new, nonstick tube and mixed with two pmol primer emITS1, 770 nM dNTPs in a final

volume of 13 μL. The reaction was incubated at 65C for 5 minutes then incubated on ice for at least 1

minute before proceeding. To the mixture was added 5x First Strand Buffer (1x final, ThermoFisher

Scientific), 100 mM DTT (5 mM final, ThermoFisher Scientific), 1 uL RNaseOUT (ThermoFisher

Scientific) then 200 units of SuperScript III RT (ThermoFisher Scientific). After thorough mixing, each

tube was incubated at 55C for 60 minutes to reverse transcribe then 70C for 15 minutes to heat kill the

transcriptase. To amplify E.muscae-specific cDNA, 2 uL of the reverse transcription reaction was mixed

with GoTaq 2x colorless mastermix (1x final, Promega) and 500 nM each primers emITS1 and emITS4

(5'- GCCTCTATGCCTAATTGCCTTT-3') then run on a thermocycler with the following settings: 95C

for 5 min followed by 35 iterations of 95C for 30 seconds, 61C for 30 seconds then 72C for 30 seconds

then 72C for an additional 10 minutes. Four µL of each reaction was analyzed by gel electrophoresis in 1%

agarose.

<u>Whole fly *in vivo* RNAseq time course</u>

One control female (24-120 hours) and two infected females (24-72 hours), one infected female and one

cadaver (96 hours) or one cadaver (120 hours) from each vial were collected as described above. RNA was

prepared from each thawed sample by homogenizing with an RNase-free pestle (Kimble Chase), washing

the pestle with 750 uL Trizol, then proceeding using the manufacturer's protocol. RNA was quantified

using a Qubit Fluorometer (Qubit RNA HS assay kit, ThermoFisher Scientific) and quality was checked by

running on a RNA 6000 Pico chip on a Bioanalyzer 2100 (Agilent Technologies). High quality RNA was

then treated with Turbo DNase (ThermoScientific) per the manufacturer's protocol. RNAseq libraries were

prepared with the TruSeq RNA v2 kit (Illumina) using 500 ng of input RNA per sample. Samples were

multiplexed 21 samples to a lane and sequenced using 100 bp paired-end reads on a HiSeq 4000 at the QB3

Vincent J. Coates Genomic Sequencing Facility at UC Berkeley.

<u>Dissected brain RNAseq</u>

Brains were individually dissected and sampled from first three control and then three exposed females.

Each animal was dissected in sterile 1x PBS in its own disposable dissection chamber (35mm petri dish

lined with 2-3% agar) and dissecting forceps were treated with 3.5% hydrogen peroxide then rinsed with

sterile water between samples to prevent nucleic acid carryover. The body of each animal was saved and

subjected to a DNA extraction using the manufacturer's provided protocol for the isolation of genomic

DNA from tissues (QIAamp DNA Micro kit, QIAGEN) eluting in 20 µL of buffer AE. For each fly body,

1 µL was used to template a PCR reaction consisting of 12.5 µL GoTaq, 2 µL of each primer emITS1 and

emITS4 (10 µM stocks), and 7.5 µL water for a final volume of 25 µL. Reactions were cycled with the

following conditions: 95C for 5 minutes followed by 35 cycles of 95C for 30 seconds, 51C for 30 seconds

and 72C for 1 minute, then a final 10 minute extension at 72C. Reactions were analyzed via gel

electrophoresis to confirm that all exposed animals had come into contact with *E. muscae* Berkeley and that

control animals were uninfected.

RNA was prepared from each thawed sample by homogenizing with an RNase-free pestle (Kimble

Chase), washing the pestle with 750 uL Trizol, then proceeding using the manufacturer's protocol. RNA

was quantified using a Qubit Fluorometer (Qubit RNA HS assay kit, ThermoFisher Scientific) and quality

was checked by running on a RNA 6000 Pico chip on a Bioanalyzer 2100 (Agilent Technologies). One

replicate control RNA sample for the 48 hour time point was lost prior to library preparation so was omitted.

High quality RNA was then treated with Turbo DNase (ThermoScientific) per the manufacturer's protocol.

RNAseq libraries were prepared with the TruSeq RNA v2 kit (Illumina) using all of the extracted RNA for

each brain, 17-75 ng of input RNA per sample. Samples were multiplexed 17 samples to a lane in equimolar

amounts and sequenced using 100 bp paired-end reads on a HiSeq 4000 at the QB3 Vincent J. Coates

Genomic Sequencing Facility at UC Berkeley.

E. muscae reference transcriptome assembly

An initial reference (Emus-Ref1) was assembled from reads from exposed *in vivo* time course samples that

had first failed to align as pairs to the D. melanogaster transcriptome (r6.11, HiSat2) then failed to align as

singletons to the D. melanogaster genome (r.611, bowtie2) using TRINITY with the developer's

recommended settings [37]. After assembly, all *in vivo* time course reads were aligned to Emus-Ref1 to

assess contamination of non-*E. muscae* sequences. All Emus-Ref1 transcripts were searched using blastn

for homology (evalue 1e-50 or smaller) to organisms not annotated as fungi or virus. These transcripts were

removed to generate Emus-Ref2. All *in vivo* time course reads were aligned to Emus-Ref2 to assess

contamination of non-E. muscae sequences. Transcripts that were not expressed by any sample (TPM = 0)

or where TPM of uninfected samples accounted for more than 10% of TPM summed across all samples

were removed to generate Emus-Ref3. Transcriptome completeness was estimated by BUSCO v1.1

analysis using the fungal reference set (1438 BUSCOs).

RNAseq data analysis

To calculate gene expression, reads were pseudo-aligned to the appropriate reference (dmelDBGP6.rel85

for D. melanogaster or Emus-Ref3 for E. muscae) and transcript abundance was estimated using Kallisto

[38]. Data were analyzed using hierarchical clustering by gene (Cluster 3.0), ANOVA between grouped

treatments (scipy.stats) and GO term analysis (Panther [39]). Hierarchical clustering heatmaps were

generated in Java TreeView; other data were plotted in matplotlib (Python), Prism (GraphPad) or Excel

2013 (Microsoft).

Paraffin embedding and microtomy of whole flies

Two mock and two exposure vials were started daily for seven days each with 50 CantonS WF flies 0-1

days old with either 0 (mock) or 6 (exposure) cadavers embedded in AS. Flies were incubated for the first

24 hours at 18C confined to 2 cm with cadavers, then moved to 21C where the confinement was relieved.

Flies were transferred to GB+ at 48 hours where they continued to be housed at 21C. Vials were sampled

every 24 hours via CO2 anesthesia then infiltrated and embedded in paraffin. For detailed protocol, see

dx.doi.org/10.17504/protocols.io.k5ecy3e. Briefly, flies were fixed 24-36 in ice-cold Carnoy's (6:3:1

ethanol:chloroform:glacial acetic acid) at 4C. Samples were next dehydrated by stepping through a series

of increasing ethanol concentrations Samples were then transitioned into Histoclear (National Diagnostic)

before slowly introducing Paraplast (Sigma). Samples were infiltrated with Paraplast for at least 84 hours

at 60C with gentle shaking before embedding in base molds with embedding rings (Thermo Scientific) and

drying overnight. Samples were stored at room temperature until they were sectioned at 8 μm with an

RM2255 microtome (Leica), applied to Polysine slides (ThermoFisher Scientific) and dried overnight at

42C. Sections were stored at room temperature for up to three weeks before Safranin O/Fast Green FCF

staining or up to one week before fluorescence in situ hybridization (FISH).

<u>Safranin O/Fast Green FCF staining of paraffin sections</u>

Slide-mounted sections were dewaxed with two, 10 minute changes of Histoclear then rehydrated to 70%

ethanol with a decreasing ethanol series. Sections were then stained one-at-a-time following Johansen's

Safranin and Fast Green protocol [40] then checked under a dissecting scope before mounting in DEPEX

mounting medium (Electron Microscope Sciences) and drying overnight. Slides were imaged using a 20x

objective with the Axio Scan.Z1 (Zeiss).

<u>Fluorescent *in situ* hybridization (FISH) of paraffin sections</u>

Slide-mounted sections were dewaxed with two, 10 minute changes of Histoclear then rehydrated to 70%

ethanol with a decreasing ethanol series. Slides were incubated in 0.2 M HCl at 37C for 45-60 minutes and

rinsed in DI water before applying 80 μL of hybridization solution (20 mM Tris-HCl pH 8.0, 0.9 M NaCl,

0.01% sodium dodecyl sulphate, 30% formamide) containing 100 pmol/μL of an *E. muscae* Berkeley-

specific DNA probe (AlexaFluor633-5'-TGCTAAAACAGCACAGTT-3', ThermoFisher Scientific).

Slides were incubated overnight in a humid chamber at room temperature. The following day, slides were

briefly washed in 1x PBS with 0.3% Triton-X100, rinsed in 1x PBS and mounted in ProLong Gold with

DAPI (ThermoFisher Scientific). Slides were cured for 24 hours before imaging on a LSM 800 confocal

microscope (Zeiss) with 5x-40x air objectives.

Supporting Information

Figure S1. Discovery of *E. muscae Berkeley*. A) *E. muscae Berkeley*-killed cadavers discovered in bait on

July 25, 2015. Note remnants of fungal growth through the intersegmental membrane of dorsal abdomen

(above and below) and spores deposited on wings (above and below) and legs (below). B,C) BLAST

(blastn) results as trees (fast minimum evolution) for B) consensus *E. muscae Berkeley* ITS (13 sequences)

and C) *E. muscae Berkeley* 28S (11 sequences). Gray legend bars show difference in percent identity

between sequences.

Figure S2. Confirmation that *E. muscae* Berkeley is growing *in vitro*. A) Alignment of ITS region from

in vitro culture (iv_culture) and consensus ITS sequence from fendel cadavers (fendel). B) Same as A, but

for LSU region.

Figure S3. Time between capture of wild *Drosophila* and death by *E. muscae* infection when housed on Koshland diet. Flies were housed at ambient temperature and humidity in a second-story apartment during August and September 2015 in Berkeley, CA. A total of 63 cadavers of wild *Drosophila* were observed out of approximately 900 flies aspirated and monitored during this time.

Figure S4. Optimization of *in vivo E. muscae* infection of CantonS WF *D. melanogaster* under

laboratory conditions. A-E) Percentage of infected cadavers at 96 hours (4 days), 120 hours (5 days) or

144 hours (6 days) after exposure to *E. muscae* Berkeley upon varying A) age; B) sex; C) temperature and

humidity; D) CO2 anesthesia for 20 minutes at the indicated post-exposure time or E) housing on 0.09%

tegosept for 120 hours beginning 48 hours after exposure. 18C > 21C indicates that vials were incubated

24 hours at 18C, 100% humidity then transferred to 21C, ~60% humidity. If no percent is indicated, then

humidity was ~60%. All vials used 4-6 cadavers to establish infection and were set up using the anesthesia-

independent protocol. Replicate vials for each condition are shown above the graph. For all panels each vial

contained 50 flies. F) Summary of method for *in vivo* propagation of *E. muscae* in CantonS WF *D.*

melanogaster. Briefly, 50 healthy, young (eclosed within the last 24 hours) CantonS flies of mixed sex are

confined within 2 cm of a circle of 6 cadavers embedded head-first in sucrose agar. Vials are incubated 24

hours at 18C and ~100% humidity. On Day 1 (24 hours since exposure) the vial plugs are raised to the top

of the vial and incubation continues for the next 24 hours at 21C with ~60% humidity. On Day 2 (48 hours

since exposure), flies are moved away from cadavers and onto GB+ diet. From Days 3-7 (96 to 168 hours

since exposure), vials are monitored 2-5 hours following "sundown" to collect fresh cadavers. These

cadavers are then used to begin new vials (Day 0).

https://youtu.be/111Wks5wYCs

Movie S1. *E. muscae* **Berkeley-infected CantonS fly undergoing end-of-life proboscis extension.** Video

recorded through the eyepiece of a dissecting microscope on a Nexus 5x phone (Google). Capture and

playback are in real time.

https://youtu.be/IiqJcpfBAGQ

Movie S2. *E. muscae* **Berkeley-infected CantonS fly undergoing end-of-life wing raising, viewed from**

the side. Video was captured with a Nexus 5x phone (Google) and macro lens (Luxsure). Capture and

playback are in real time.

https://youtu.be/DdR-iSdYG6A

Movie S3. *E. muscae* **Berkeley-infected CantonS fly undergoing end-of-life wing raising, viewed head-**

on. Video was captured with a Nexus 5x phone (Google) and macro lens (Luxsure). Capture and playback

are in real time.

https://youtu.be/qDCZJmhWkbU

Movie S4. Animated time lapse of *E. muscae* **Berkeley-infected CantonS cadavers undergoing spore**

production and ejection. Sporulation time lapse in *E. muscae Berkeley* killed cadavers. Time listed at the

top right corner of each frame is the time that has elapsed since the light-dark transition. One image was

taken every minute for ~24 hours with three cadavers situated on a cotton flug at ambient temperature and

humidity. Images are played back at 10 fps.

1000

1001   https://youtu.be/T5Pi0JoEm0I

1002   **Movie S5. A primary conidium is ejected from a conidiophore formed in an** *E. muscae* **Berkeley-**

1003   **killed cadaver.** Arrowheads indicates conidium that launches and the vacant conidiophore that remains

1004   after launch. Video was captured at 54,000 frames per second (fps) at 5x magnification; frames are played

1005   back at 5 fps. Scale bar is 25 μm.

1006

1007   https://youtu.be/B3BHDwvagzg

1008   **Movie S6. A primary conidium lands on the lid of a polystryene petri dish.** The conidium lands as one

1009   complete unit, supporting the fungal cannon mechanism of primary conidium ejection in *E. muscae*. Video

1010   was captured at 18,000 fps at 5x magnification; frames are played back at 5 fps. Scale bar is 25 μm.

1011

Figure S5. *E. muscae Berkeley*-infected CantonS flies housed in constant darkness do not consistently

die in a gated fashion. Each circle represents the time of last movement observed for one cadaver as

detected by the DAM. Flies were raised on the indicated 12:12 light:dark cycle then exposed to *E. muscae*

Berkeley as per standardized protocol. Dotted line indicates the expected light-dark transition based on the

entrained light cycle. A) Flies entrained on a 12:12 light:dark cycle with photophase beginning at 1 am and

scotophase beginning at 1 pm. B) Flies entrained on a 12:12 light:dark cycle with photophase beginning at

7 pm and scotophase beginning at 7 am.

Figure S6. Reverse-transcription PCR of *E. muscae* Berkeley ITS sequence in exposed and control flies. All samples were run on the same gel with equal loading volumes; samples to the right of the dashed line were run on the lower half of the same gel containing samples to the left of dotted line (see methods for sample generation). DNA ladder (5 µL Hyperladder 1 kb, Bioline) was run in the first lane of each gel half. Samples are shown in chronological order, with the time point indicated above the left-most sample. Black lines indicated exposed flies; white lines indicate unexposed flies. Exposed flies collected at 96 hours were cadavers that had died of *E. muscae* Berkeley infection. Plus (+) indicates positive control (*E. muscae* Berkeley *in vitro* RNA template); minus (-) indicates additional negative control (*D. melanogaster* RNA from earlier experiment, before the discovery and introduction of *E. muscae* Berkeley to the laboratory).

https://youtu.be/GeTRUiBIW8s

Movie S7. Formation of primary conidia atop conidiophores. The intersegmental membranes of a fresh *E. muscae Berkeley*-killed cadaver were imaged at 20x magnification every minute for 222 minutes beginning 4 hours and 10 minutes after sunset. The time lapse is played back at 10 fps. Scale bar is 50 µm.

https://youtu.be/1vrXbDWHQCw

Movie S8. Formation of secondary conidia from off-target primaries. The wing of an *E. muscae* Berkeley-killed cadaver was imaged at 20x magnification every minute for 361 minutes beginning 10 hours and 30 minutes after the light-dark transition. The time lapse is played back at 10 fps. Scale bar is 100 µm.

Table S1. GO term enrichments of host gene groups as given in Figure 4C. Panther GO-term analysis (complete biological process) for genes in Groups i-vi (Figure 4C).

➢ See file Table_S1.pdf

1043

Figure S7. Genes exhibiting differential expression between flies after mock exposure or exposure to

E. muscae **at 24, 48 or 72 hours.** A) Left: Volcano plot for all genes at 24 hours. P-value is determined by

ANOVA grouping control vs. exposed samples. Genes with p-value < 0.001 are shown in color. Right:

Panther GO-term analysis (complete biological process) of genes overexpressed in exposed animals (red)

or control animals (blue). B) Left: Volcano plot for all genes at 48 hours. P-value is determined by ANOVA

grouping control vs. exposed samples. Genes with p-value under 0.001 are shown in color. Right: No

significant GO term enrichments were found for differentially-expressed genes. C) Left: Volcano plot for

all genes at 72 hours. P-value is determined by ANOVA grouping control vs. exposed samples. Genes with

p-value under 0.001 are shown in color. Right: Panther GO-term analysis (complete biological process) of

genes overexpressed in control animals (blue). There are no significant GO term enrichments for set of

genes overexpressed in exposed samples.

Group A	Group B	
PGRP-LB	AttC	Tep2
CG10433	Mtk	MP1
Sp7	Dro	NimB3
PGRP-SA	DptB	Hayan
Spn28Dc	PGRP-SD	CG42259
grass	Drs13	
Spn27A	Drs12	
spirit	CG15065	
AttA	SPH93	
AttB	SPE	
TotM	GNBP-like3	
CecC	NimB1	
CecB	IM3	
Listericin	IM2	
IM18	IM14	
DptA	IM23	
edin	IM1	
CG30098	nec	
CecA2	IM4	
CecA1	Cpr49Ac	

1055

Figure S8. Expression of immune genes over the course of infection of *D. melanogaster* by *E. muscae*

Berkeley. Complete linkage hierarchical gene clustering by gene was performed in Gene Cluster 3.0 on all

genes annotated with defense response (GO 0006952) after filtering out across all genes that are expressed

at least at two TPM in at least three out of 42 samples (10,875 transcripts total), then log transforming and

centering on the mean value for each transcript. Samples are ordered by percentage of *E. muscae* Berkeley

reads as a fraction of the total reads aligned (above). The scale bar for the heatmap is given to the right of

the plot. Two 96 hour exposed samples that show an aberrant immune response compared to all other

exposed samples are indicated by asterisks.

1064

Figure S9. Host gene expression in the brain is stable over the first 72 hours of *E. muscae Berkeley*

infection. Right) All pairwise linear correlations between samples from *E. muscae Berkeley*-infected whole

fly RNAseq time course. Left) All pairwise linear correlations between samples from *E. muscae Berkeley*-

infected dissected brain pilot RNAseq time course. Samples are named in the following format:

HourTypeReplicate, with "C" indicating controls, "E" indicating exposed flies living at the time sampled

and "cad" indicates that the fly had been killed by *E. muscae Berkeley* before sampling. For example,

"24C1" indicates a the first replicate control sample (uninfected fly) taken at 24 hours after mock exposure.

Black rectangles outline rows and columns containing correlation values for control samples. Control

samples are denoted on each axis with a black bar.

References

1. Hughes DP, Andersen SB, Hywel-Jones NL, Himaman W, Billen J, Boomsma JJ. Behavioral
mechanisms and morphological symptoms of zombie ants dying from fungal infection. BMC Ecol.
2011;11: 13.

2. Vyas A, Kim S-K, Giacomini N, Boothroyd JC, Sapolsky RM. Behavioral changes induced by
Toxoplasma infection of rodents are highly specific to aversion of cat odors. Proc Natl Acad Sci.
2007;104: 6442–6447.

3. Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, et al. A phylum-level
phylogenetic classification of zygomycete fungi based on genome-scale data. Mycologia. 2016;108:
1028–1046.

4. Cohn F. Empusa muscae und die krankheit der Stubenfliegen. Ein Beitrag zur Lehre von den durch
parasitische Pilze charakterisierten Epidemien. Nova acta Academiae Caesareae Leopoldino-Carolinae
Germanicae Naturae Curiosorum. 1855;25: 299–360.

5. Keller S. Entomophthora muscae als Artenkomplex. Mitteilungen der Schweizerischen Entomologischen
Gesellschaft. agris.fao.org; 1984;57: 131–132.

6. MacLeod DM, Müller-Kögler E, Wilding N. Entomophthora species with E. muscae-like conidia.
Mycologia. JSTOR; 1976;68: 1–29.

7. Roy HE, Steinkraus DC, Eilenberg J, Hajek AE, Pell JK. Bizarre Interactions and Endgames:
Entomopathogenic Fungi and Their Arthropod Hosts. Annu Rev Entomol. 2006;51: 331–357.

8. Balazy S. On Rhizoids of Entomophthora-muscae (Cohn) Fresenius (Entomophthorales,
Entomophthoraceae). Mycotaxon. 1984;19: 397–407.

9. Krasnoff SB, Watson DW, Gibson DM, Kwan EC. Behavioral effects of the entomopathogenic fungus,
Entomophthora muscae on its host Musca domestica: Postural changes in dying hosts and gated pattern of
mortality. J Insect Physiol. Elsevier; 1995;41: 895–903.

10. Brobyn PJ, Wilding N. Invasive and developmental processes of Entomophthora muscae infecting
houseflies (Musca domestica). Trans Br Mycol Soc. Elsevier; 1983;80: 1–8.

1. Mullens BA, Rodriguez JL. Dynamics of Entomophthora muscae (Entomophthorales:
Entomophthoraceae) Conidial Discharge from Musca domestica (Diptera: Muscidae) Cadavers. Environ
Entomol. ee.oxfordjournals.org; 1985;14: 317–322.

2. Mullens, B A Rodriguez, J L Meyer, J A. An Epizootiological Study of Entomophthora muscae in
Muscoid Fly Populations on Southern California Poultry Facilities, with Emphasis on Musca domestica.
Hilgardia. 1987;55.

3. Charnley AK. Physiological aspects of destructive pathogenesis in insects by fungi: a speculative review.
In: J M Anderson Adm Rayner, editor. Invertebrate-microbial interactions: Joint symposium of the British
Mycological Society and the British Ecological Society held at the University of Exeter 1982. 1984. pp.
229–270.

4. Watson DW, Petersen JJ. Sexual Activity of Male Musca domestica (Diptera: Muscidae) Infected with
Entomophthora muscae (Entomophthoraceae: Entomophthorales). Biol Control. Elsevier; 1993;3: 22–26.

5. Steinkraus DC, Kramer JP. Susceptibility of sixteen species of Diptera to the fungal pathogen
Entomophthora muscae (Zygomycetes: Entomophthoraceae). Mycopathologia. 1987;100: 55–63.

6. Kramer JP, Steinkraus DC. Culture of Entomophthora muscae in vivo and its infectivity for six species of
muscoid flies. Mycopathologia. Springer Netherlands; 1981;76: 139–143.

7. Keller S. Arthropod-pathogenic Entomophthorales: Biology, Ecology, Identification. 2007.

8. Elya C, Zhang V, Ludington WB, Eisen MB. Stable Host Gene Expression in the Gut of Adult
Drosophila melanogaster with Different Bacterial Mono-Associations. PLoS One. Public Library of
Science; 2016;11: e0167357.

9. Turian G, Wuest J. MYCOSES CAUSED BY ENTOMOPHTHORACEAE AFFECTING
POPULATIONS OF ANTS AND DROSOPHILA. Mitteilungen der Schweizerischen Entomologischen
Gesellschaft. 1969;42: 197–201.

10. Goldstein B. An Empusa Disease of Drosophila. Mycologia. Mycological Society of America; 1927;19:
97–109.

11. Keller S. The genus Entomophthora (Zygomycetes, Entomophthorales) with a description of five new
species. SYDOWIA-HORN-. VERLAG FERDINAND BERGER; 2002;54: 157–197.

22. Liu H, Beckenbach AT. Evolution of the mitochondrial cytochrome oxidase II gene among 10 orders of
insects. Mol Phylogenet Evol. 1992;1: 41–52.

23. Hajek AE, Papierok B, Eilenberg J. Chapter IX - Methods for study of the Entomophthorales. In: Lacey
LA, editor. Manual of Techniques in Invertebrate Pathology (Second Edition). San Diego: Academic
Press; 2012. pp. 285–316.

24. Kalsbeek V, Pell JK, Steenberg T. Sporulation by Entomophthora schizophorae (Zygomycetes:
Entomophthorales) from housefly cadavers and the persistence of primary conidia at constant
temperatures and relative humidities. J Invertebr Pathol. Elsevier; 2001;77: 149–157.

25. Yafetto L, Carroll L, Cui Y, Davis DJ, Fischer MWF, Henterly AC, et al. The fastest flights in nature:
high-speed spore discharge mechanisms among fungi. PLoS One. 2008;3: e3237.

26. Humber RA. An alternative view of certain taxonomic criteria used in the Entomophthorales
(Zygomycetes) [Fungi, insect pathogens]. Mycotaxon. apr-jun1981;13. Available:
http://agris.fao.org/agris-search/search.do?recordID=US8112089

27. Becher PG, Jensen RE, Natsopoulou ME, Verschut V, De Fine Licht HH. Infection of Drosophila suzukii
with the obligate insect-pathogenic fungus Entomophthora muscae. J Pest Sci. Springer Berlin
Heidelberg; 2017; 1–7.

28. Keller S, Kalsbeek V, Eilenberg J. Redescription of Entomophthora muscae (Cohn) Fresenius. Sydowia.
1999;51: 197–209.

29. Boomsma JJ, Jensen AB, Meyling NV, Eilenberg J. Evolutionary interaction networks of insect
pathogenic fungi. Annu Rev Entomol. 2014;59: 467–485.

30. De Fine Licht HH, Jensen AB, Eilenberg J. Comparative transcriptomics reveal host-specific nucleotide
variation in entomophthoralean fungi. Mol Ecol. 2017;26: 2092–2110.

31. Butt TM, Hajek AE, Humber RA. Gypsy Moth Immune Defenses in Response to Hyphal Bodies and
Natural Protoplasts of Entomophthoralean Fungi. J Invertebr Pathol. 1996;68: 278–285.

32. Beauvais A, Latge J-P, Vey A. The Role of Surface Components of the Entomopathogenic Fungus
Entomophaga aulicae in the Cellular Immune Response of Galleria mellonella (Lepidoptera). J Gen
Microbiol. 1989;135: 489–498.

33. Valanne S, Wang J-H, Rämet M. The Drosophila Toll signaling pathway. J Immunol. 2011;186: 649–656.

34. Levitin A, Whiteway M. Drosophila innate immunity and response to fungal infections. Cell Microbiol. 2008;10: 1021–1026.

35. Fredericksen MA, Zhang Y, Hazen ML, Loreto RG, Mangold CA, Chen DZ, et al. Three-dimensional visualization and a deep-learning model reveal complex fungal parasite networks in behaviorally manipulated ants. Proc Natl Acad Sci U S A. 2017; doi:10.1073/pnas.1711673114

36. Humber RA. The systematics of the genus Strongwellsea (Zygomycetes: Entomophthorales). Mycologia. 1976;68: 1042–1060.

37. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol. 2011;29: 644–652.

38. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol. 2016;34: 525–527.

39. Mi H, Poudel S, Muruganujan A, Casagrande JT, Thomas PD. PANTHER version 10: expanded protein families and functions, and analysis tools. Nucleic Acids Res. 2016;44: D336–42.

40. Ruzin SE. Plant Microtechnique and Microscopy. Oxford University Press; 1999.